1

Enrichment of halogenated organic compounds and their degrading

microorganisms in the deepest ocean

Rulong Liu[1*], Hui Wei[1#], Zhiao Xu[1#], Yuheng Liu[1], Zhixuan Wang[1], Jiani He[1], Min

Luo[1], Federico Baltar[1, 2], Yunping Xu[1], Qirui Liang[1], Liting Huang[1], Li Wang[1*],

Jiasong Fang[1, 3, 4*]

[1] College of Oceanography and Ecological Science, Shanghai Ocean University,

Shanghai, China

[2] Department of Functional and Evolutionary Ecology, University of Vienna, Vienna,

Austria

[3] Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for

Marine Science and Technology, Qingdao, China

[4] Department of Natural Sciences, Hawaii Pacific University, Honolulu, HI, USA

*Correspondence:

Rulong Liu, rlliu@shou.edu.cn

Li Wang, wang-l@shou.edu.cn

Jiasong Fang, jsfang@shou.edu.cn

[#], these authors contribute equally.

Abstract

The hadal trenches, the deepest regions of the ocean, serve as the final resting place for marine particles and as "tunnels" for material exchange between the ocean and Earth's interior. Despite their extreme conditions, the trenches contain high content of organic carbon (OC) and active microbial carbon turnover, are hotspots for deep-sea OC degradation and unique microbial processes. However, little is known about the OC components and microbial metabolisms driving the OC degradation in the trenches. This study reveals unexpectedly high concentrations of halogenated organic compounds (HOCs) in the Mariana Trench sediments, with about 1 halogen per 58 carbon atoms in the OC pool. Systematic analysis of deep-sequenced metagenomes and metagenome-assembled-genomes from the global ocean shows significantly higher abundance of the genes for biodegradation of HOCs (dehalogenation) in trench microbes, with microorganisms capable of dehalogenation belonging to 16 phyla and 52 orders, 75% of which identified as HOCs degraders for the first time. Microcosms simulating trench conditions showed rapid degradation of typical HOCs and transcription of genes related with HOCs metabolisms, demonstrating the active HOCs degradation by the trench microorganisms. The findings suggest the HOCs metabolism as an important process in the OC remineralization of deep-sea trenches, advancing understanding of deep-sea carbon cycling and microbial survival strategies.

Introduction

The deep ocean, covering 65% of the Earth's surface and housing 75% of the ocean's prokaryotic biomass, plays a crucial role in the remineralization and long-term storage of marine organic carbon [1]. However, our understanding of the metabolism of deep-sea microorganisms and the carbon turnover processes they drive remains limited, especially concerning carbon cycling in the ocean's deepest regions, known as hadal trenches. The hadal trenches, which are found at depths exceeding 6,000 meters, are among the most inaccessible and least understood marine environments on the planet [2, 3]. Located in subduction zones, the hadal trenches serve as both the endpoint for sinking organic matter from the upper ocean and as a "channel" for material exchange between the ocean and the Earth's deep interior [3]. Recent studies have shown that the hadal zone is rich in organic matter and exhibits unexpectedly active microbial organic carbon turnover, emerging as "hotspots" for organic carbon remineralization in the deep ocean [2, 4]. Furthermore, the extreme conditions of the hadal zone, such as high pressure and geographic isolation, have led to the development of unique microbial communities and special metabolic processes, forming a distinct "trench biosphere" [5, 6]. Understanding the processes and mechanisms by which hadal microbes drive the degradation of sedimentary organic carbon is essential for advancing knowledge of carbon cycling and life processes in deep-sea environments [2, 3].

While studies have demonstrated that hadal trench microorganisms exhibit potential for chemoautotrophic metabolism such as ammonia oxidation [7], metagenomic and genomic analyses suggest that microbial taxa present in the trench sediment predominantly adopt a heterotrophic lifestyle, possessing metabolic genes and

pathways for degradation of a wide range of organic matter, including simple sugars, peptides, fatty acids, proteins and other organic detritus [6, 8]. Particularly, most of the studies demonstrated the metabolic potential of hadal trench microbes to utilize multiple complex and recalcitrant organic compounds, such as hydrocarbons, polysaccharides and aromatic compounds [6, 8]. These findings are consistent with the higher humification of organic carbon in the bottom of hadal sediments [9], suggesting that the utilization of recalcitrant organic carbon might be a crucial metabolic strategy for hadal microbes [8, 10]. However, only a limited number of these metabolic processes, such as hydrocarbon degradation, have been experimentally validated combined evidences of biogeochemical detection of substrates and the assessment of microbial degradation activities [11, 12]. Due to the diverse and variable sources of organic matter[3], along with the unique biological, geological, and physicochemical factors regulating the dynamics of materials in hadal trenches [13, 14], the composition of organic matter in hadal sediments is highly complex. Consequently, our understanding of the components of organic carbon (OC) in hadal trenches and the microbial processes driving OC degradation in these sediments remains limited.

Halogenated organic compounds (HOCs) are widely recognized as environmentally hazardous substances, raising significant global concerns. As persistent organic pollutants, most anthropogenic HOCs exhibit high toxicity to various organisms and a strong tendency to bioaccumulate, causing considerable harm to the marine ecosystems [15]. In addition to human sources, large quantities of natural HOCs are also produced through various biological and abiotic processes [16, 17]. The ocean, being rich in halogen atoms, serves as the largest source of natural HOCs on Earth [18], with high level of HOCs detected in sinking particles [19]. Due to their recalcitrant nature, HOCs

are thought to be transported to the deep ocean via mechanisms such as biological pump and ultimately sequestered in deep-sea sediment [15, 19, 20]. However, little is known about the content of HOCs and their fate in the deep ocean.

Hadal trenches serve as major depositories of different types of pollutants [21, 22]. Recent studies have identified a variety of HOCs, such as organochlorine pesticides dichlorodiphenyltrichloroethane (DDT) and chlordane, industrial pollutants PCB and polybrominated diphenyl ethers (PBDE), in the macro-organisms and sediments of the Mariana Trench [22, 23, 24, 25]. In addition, our previous work revealed that genes encoding complete metabolic pathways for HOCs degradation are present in genomes of hadal Chloroflexi, one of the dominant taxa in in trench sediments [10, 26]. The finding suggests the possibility of microbial HOCs degradation in the hadal trench sediment. Nevertheless, the metabolic processes, degradation activities as well as the significance of the HOCs degradation in hadal trench carbon cycling is not explored. In this study, we applied an optimized combustion-IC based approach to determine the content and composition of total HOCs in sediments of the Mariana Trench. Surprisingly, the results revealed that the total HOCs represent a large proportion of OC pool in the trench sediments. Systematic metagenomic and genomic analysis further showed that genes encoding enzymes and pathways for HOCs degradation are enriched and prevalent in hadal trenches sediments, and are widely distributed across more than half of the microbial taxa. Microcosm experiments simulating trench conditions further demonstrated active degradation of HOCs by trench microbial communities. These findings provide novel insights into the significance of HOCs and their microbial degradation processes in carbon cycling of the hadal zone.

Results and Discussion

High content of HOCs in hadal trench sediments

A method based on combustion-ion chromoatography were developed and optimized to show high accuracy, precision and reproducibility in determining the content of adsorbable organic halogens (AOX) including fluorine, chlorine and bromine (AOF, AOCl, AOBr), and the corresponding insoluble organic halogens (IOF, IOCl, IOBr, IOX), as well as total organic halogen contents (TOBr, TOCl, TOF, TOX) in marine sediments [27]. In this study, the content and composition of HOCs were determined in 38 sediment samples of 4 sediment cores taken from 5842-10901 meters of the Mariana Trench, which resulted in a total number of 430 data points of different components of HOCs (Fig. 1; Fig. S1; Table S1). The results showed that HOCs presented in all of the trench sediments tested (Fig. 1B). The values of AOX, which indicates dissolved HOCs adsorbable by activated carbon, ranged from 4.90±0.70 to 383.14±64.17 mg/kg, while the insoluble HOCs (IOX) ranged from 71.82±0.50 to 459.96±21.90 mg/kg (Fig. 1B), were significantly higher than the content of AOX ($p < 0.01$, t-text). The dissolved fraction of HOCs were mainly dominated by organic chlorine compounds (AOCl), which were significantly more abundant than AOBr and AOF. In contrast, organic fluorine compounds (IOF) showed the highest abundance in the insoluble HOCs, followed by IOCl (Fig. 1B). Organic bromine compounds showed the lowest abundance in both the dissolved and insoluble HOCs (Fig. 1B). The concentration of total organic halogen (TOX) in sediments of the Mariana Trench ranged from 110.33 ± 2.71 to 594.15 ± 15.28 mg/kg, with an average value of 292.47 mg/kg (Fig. 1B). Although hydrocarbons and lipid biomarkers, e.g. alkanes, carboxylic acids, alcohols and GDGTs, have been reported to be potential components in OCs of hadal trench sediments, these compounds generally showed low abundance

(up to 15 mg/kg) [11, 12, 28]. HOCs determined in this study were at least one magnitude

higher than the reported OC compounds, making HOCs the largest known component

of organic matter in hadal trench sediments. The contents of total organic carbon were

relatively stable in the tested samples, ranged from 0.48 – 0.69% (0.57 ± 0.06 %)

(Fig. 1B). As a result, the calculated molar ratios of carbon to halogen atoms (C:X)

were ranged from 17.8 to 121.8, with an average value of 49.2. This indicates an

average of one halogen atom for every 49.2 organic carbon atoms, suggesting a high

proportion of HOCs within the organic carbon pool in the tested trench sediments.

Currently, understanding on components of deep-sea organic matter is scarce, and to

the best of our knowledge, only one existing study reported the TOX values in the

deep ocean and from only two stations [27]. The values of TOX in sediments of the MT

were significantly higher than those reported in coastal and other deep-sea regions,

despite the lower TOC content in the trench sediments (Fig. S2) [27]. Consequently, the

proportion of HOCs were significantly higher in organic carbon of the trench

sediments than those in the coastal or other deep-sea sediments (evidenced by the

lower C:X ratio) (Fig. S2).

HOCs in marine environments might be natural or anthropogenic sourced [17, 18]. The

hadal trenches have been reported to accumulate many types of manmade pollutants

such as black carbon [29], mercury [30], and plastics [21]. It was therefore not surprising to

have the anthropogenic HOCs in the trenches, and indeed, various types of

anthropogenic HOCs, such as PCBs, PBDEs, and halogenated organic pesticides have

been reported in animals of several trenches, including the MT [22, 23, 24, 31]. However,

the reported anthropogenic HOCs generally showed low concentration in the trench

sediment, with average values ranged from 0 to 2 µg/kg [22, 23, 24]. Our results showed

that the concentration of TOX were magnitudes higher than all of the reported

anthropogenic HOCs, suggesting that large amount of HOCs might be come from

other sources. Carbon isotopic compositions (δ13C) of TOC ranged between -20.0 to -

22.6 ‰, with an average value of -21.4 ‰ in the sediments analyzed (Fig. 1B), which

indicating that the organic matter in the sediments of the MT was mainly marine

sourced [32]. Such finding was consistent with conclusions driven by previous studies

carried out in the same region using the carbon isotopic or biomarker evidences [32, 33].

Marine sourced HOCs might therefore be an important contributor for the measured

high content of total HOCs in the hadal sediment of the MT. In fact, previous studies

have shown that HOCs can be formed biologically via synthesis of marine organisms

or abiotically via physical/chemical halogenation of natural organic matter in seawater

[17, 18, 19], and that the marine particles can be a natural sink for HOCs [19]. The TOX

values determined in the trench sediments here were comparable with those in deep-

sea particles (180-694 mg/kg) [19], supporting the potentials of contribution from

marine particles.

Enrichment of dehalogenase in sediments of the Mariana Trench

HOCs are typically considered to be resistant to microbial degradation [16]. The

potential for microorganisms residing in hadal trench sediments to metabolize and

degrade HOCs was investigated through a comprehensive metagenomic analysis

focusing on the occurrence, relative abundance, and distribution of dehalogenase,

which is the key enzyme for the biodegradation of HOCs [34, 35, 36]. A total of 54

metagenomes from global ocean sediments were examined, comprising 27 deeply

sequenced metagenomes from sediment cores collected from depths of 5,437 to

10,954 meters in the Mariana Trench, alongside 27 metagenomes from other deep-sea

and shallow coastal sediments (Fig. 1A, Table S2). Following sequence preprocessing

and functional annotation against the KEGG database, various types of dehalogenase

genes were identified and validated through BLASTP search against the UniProt

protein database. The relative abundance of these confirmed dehalogenase genes was

quantified and expressed as genes per million (GPM). The results indicated the

presence of dehalogenases in all metagenomes derived from Mariana Trench

sediments (Fig. 2A). Specifically, four distinct types of dehalogenases were identified:

haloalkane dehalogenase, S-2-haloacid dehalogenase, haloacetate dehalogenase, and

reductive dehalogenase (Fig. 2A). These enzymes facilitate the biodegradation of

various HOCs through either hydrolytic or reductive mechanisms [18], suggesting a

significant metabolic versatility regarding HOCs within trench sediments. Notably,

the relative abundance of haloalkane dehalogenase and S-2-haloacid dehalogenase

was significantly higher in Mariana Trench sediments compared to other deep-sea and

shallow sediments ($p < 0.001$, t-test) (Fig. 2A), indicating an enrichment of

dehalogenation metabolism within the microbiome of the Mariana Trench. These

results align with the high levels of HOCs found in the sediment of the trench (Fig.

1B), implying that trench microorganisms may utilize HOCs as substrates for living.

Furthermore, phylogenetic analysis of 2,300 haloalkane dehalogenase sequences from

the Mariana Trench metagenomes, in conjunction with all manually annotated and

non-redundant haloalkane dehalogenase sequences available in the UniProt-Swiss

database, revealed that the microbial communities in the Mariana Trench encompass

all three types of haloalkane dehalogenases (HDL-I, II, and III) (Fig. 2B). This

suggests functional complementation among these enzymes, as different types of the

enzyme exhibit varying substrate specificities [37]. Interestingly, the sequences derived

from the trench metagenomes formed unique clusters that were distinct from known

haloalkane dehalogenases (Fig. 2B), indicating the presence of novel phylotypes that

may possess unique structural and functional characteristics, which were potentially

resulting from long-term adaptation to the extreme environmental conditions of the

hadal trench, such as low temperatures and high pressures [2, 3].

Prevalence of dehalogenation metabolism among hadal trench microorganisms

In this study, a total of 2544 metagenome-assembled-genomes (MAGs) were

reconstructed from the 27 metagenomes derived from the Mariana Trench sediments.

After dereplication at 95% average nucleotide identity (ANI), 285 representative

genomes were obtained. Functional annotation and homology confirmation revealed

that over half of the trench microbial taxa, specifically 50.2% (143 MAGs), harbored

genes encoding dehalogenases (Fig. 2A; Table S3). These MAGs showed

contamination < 4.99%, the genome size ranged from 0.58 to 5.31 Mbp, the GC

content ranged from 59.4 to 65.4%, and 79 of them had a genome completeness

greater than 80% (Fig. 2A; Table S3). Among the 143 MAGs, 78 MAGs contained

genes for S-2-haloacid dehalogenase, 11 for haloacetate dehalogenase, 79 for

haloalkane dehalogenase, and 23 for reductive dehalogenase. While the majority of

the MAGs harbored a single type of dehalogenase, 40 MAGs contained genes for

multiple dehalogenase types, with one MAG from the Chloroflexota phylum

(MT_10840m_10-14cm.31) possessing genes for all four dehalogenases (Fig. 2A).

Phylogenomic analysis and taxonomic classification against the GTDB database

revealed that the 143 MAGs with dehalogenation potentials were distributed across 16

bacterial phyla and 20 classes, including 60 novel species and 15 novel genera (Fig.

2A; Table S3). By comparing against the mibPOPdb database, which is a manually

curated integrative resource for microbial bioremediation of persistent organic

pollutants (POP) [38, 39], 11 of the 16 phyla were reported to have the capability of

dehalogenation for the first time (Fig. 2A), thereby greatly enhancing the current

understanding of the diversity of dehalogenation microorganisms. The MAGs

encoding dehalogenases were found to be widely distributed across all analyzed MT

sediment samples, with the total relative abundance of dehalogenation

microorganisms constituting between 8% and 21% of the metagenome reads (Fig.

2B). Majority of the MAGs (96 MAGs) showed high occurrence frequency, being

present in over 80 % of the MT sediment samples (Fig. 2B, Table S4). These findings

suggest that the dehalogenation microorganisms were both prevalent and dominant in

the microbial communities of the trench sediments.

Functional annotation utilizing the KEGG, COG, and PROKKA databases revealed

that the dehalogenation population, represented by the 143 dehalogenation MAGs,

contained gene sets that encode pathways for the degradation of various HOC

compounds with distinct molecular structures. Specifically, the population was

associated with the degradation of 18 compounds categorized into groups such as

chlorocyclohexane and chlorobenzene, chloroalkane and chloroalkene,

fluorobenzoate, as well as polychlorobiphenyl (PCB) (Fig. 3, Fig. S3, Table S5).

These represent nearly all documented degradation pathways for HOCs in the KEGG

database, suggesting the wide range of substrate spectra for the dehalogenation

population identified in hadal trench environments. Majority of the HOCs compounds

(15 of 18), e.g. γ- hexachlorocyclohexane, 2,6-dichloropenol, 2,4-

dichlorophenoxyaceate, 2,4-dichloroaniline, cis-dichloropropene, trans-

dichloropropene, 1,2-dichloroethane, dichloromethane, 2-fluorobenzoate, 3-

fluorobenzoate, 4-chlorobenzoate, and 4-chlorobiphenyl, could be completely

degraded via entering the TCA cycle (Fig. 3, Table S5). Furthermore, coding genes for

each of the functional enzymes involved in these degradation pathways was identified

across various genomes from multiple phyla (Fig. 3, Table S5), indicating a functional

redundancy that facilitates the efficient degradation of substrates [40]. The exact

molecular composition of HOCs present in trench sediment remains undetermined.

Nevertheless, as previously discussed, they are hypothesized to comprise a mixture of

both anthropogenic and natural HOCs. The anthropogenic HOCs are characterized by

low structural diversity [16], with majority of their degradation pathways being

annotated in our analysis (Fig. 3, Table S5). In contrast, natural HOCs exhibit a

considerably higher diversity in their molecular structures [16, 17]. However, as the

primary source of organic matter in deep-sea and hadal trenches, phytoplankton and

sinking particles have been identified as containing HOCs that were predominant by

aliphatic forms [19]. Considering the diverse degradation pathways identified in the

MAGs, the dehalogenation population possess the potential to efficiently utilize HOC

substrates present in hadal trenches.

In addition to direct degradation of HOCs, the dehalogenation population showed

potentials to contribute to other ecological functions. Firstly, the degradation

pathways of HOCs annotated involved were generally long, with many steps and

produce series of intermediates (Fig. 3). These intermediate compounds, with lowered

recalcitrancy compared to their original compounds, can also be substrates for other

heterotrophic taxa, supporting the potential role of the dehalogenation

microorganisms as mediators of microbial interactions within hadal trenches [16, 41].

Furthermore, the dehalogenation microbial populations also showed the capability to

drive the biogeochemical cycling of multiple key elements other than halogen and

carbon. Particularly, the sulfur oxidation seemed to be common in the dehalogenation

population, with 49-87% of the MAGs harbored gene sets for different steps of sulfur

oxidation (Fig. 3, Fig. S3, Table S5). Moreover, 43% and 45% of the MAGs also

showed potentials of iron reduction and iron oxidation, respectively (Fig. 3, Fig. S3,

Table S5). Overall, the findings demonstrate a significant role of dehalogenation

microorganisms in the functions and structures of hadal trench ecosystems.

Active HOCs degradation driven by hadal trench microorganisms

The degradation activity of HOCs by hadal trench microorganisms was demonstrated

by inoculating typical HOCs compounds—PCB and γ-HCH—into sediments from the

Mariana Trench (MBR06, 6477 m depth) and incubating them for 270 days under

high pressure (60 MPa) and low temperature (4 °C) to simulate *in-situ* conditions.

Time-series sampling and monitoring revealed a rapid decrease of substrate

abundance in all treatments containing trench microorganisms (non-sterile systems),

whereas no significant changes occurred in the control systems (sterile sediments)

(Fig. 4A). For both substrates, the most substantial decrease took place within the first

30 days, leaving less than 40% of the original material. PCB levels continued to

decline throughout the 270-day period, whereas γ-HCH remained relatively stable

after day 30. Reads recruited to metagenomes and metatranscriptomes indicated that

the dehalogenation population, i.e., the 143 MAGs with dehalogenation potential,

were present and actively expressed in the microcosms (Fig. 4B). Dominant

dehalogenation MAGs included those from the orders UBA1845

(p_Planctomycetota), Hydrogenedentiales (p_Hydrogenedentota), SAR324 (p_SAR

324), Longimicrobiales (p_Gemmatimonadota), UBA5794 (p_Actinomycetota),

UBA9160 (p_Myxococcota), Woeseiales, Pseudomonadales, Kiloniellales

(p_Proteobacteria), and SM23-28-2, UBA1151, SAR202, UBA3496, DSTF01

(p_Chloroflexota) (Fig. 4B). Among these, MAGs from the order Pseudomonadales

were the most abundant in the microcosms (Fig. 4B). The majority of functional

genes associated with HOC degradation pathways were found in high relative

abundance in the metagenomes of the microcosms (Fig. 4C, Table S6), with over half

of these genes being actively expressed in the transcriptome (Fig. 4D, Table S6).

Notably, genes such as *dhaA*, *EC:1.1.1.1*, *clcD*, *catB*, *EC1.14.13.-*, and *ALDH*, which

encode haloalkane dehalogenase, alcohol dehydrogenase,

carboxymethylenebutenolidase, muconate cycloisomerase, 2-octaprenylphenol

hydroxylase, and aldehyde dehydrogenase, respectively, exhibited high transcriptional

abundance (Fig. 4D). Overall, the results from the chemical and microbial analyses in

the microcosm experiments indicated rapid degradation of typical HOCs, alongside

active expression of microorganisms and functional genes related with dehalogenation

metabolism. This suggests that hadal trench microorganisms actively metabolize HOC

under the high pressure and low temperature conditions. However, this cultivation

experiment was limited to two specific human-made halogenated organic compounds.

Future research is necessary to explore the metabolic activity of deep-sea

microorganisms on other types of HOCs, particularly natural ones. Additionally, since

this experiment utilized enriched sediment microorganisms in a laboratory setting, the

degradation rates observed may differ from those in natural environments. Future in-

situ cultivation studies are needed for a more accurate evaluation of the degradation

rates and flux of HOCs.

Implications for deep-sea biogeochemical cycling and environmental

bioremediation

The deep-sea environments constitute a significant portion of the Earth's microbial biomass and production, thereby playing a crucial role in regulation of global carbon cycling and climate change[1]. However, there remains a limited understanding of the metabolic processes of deep-sea microorganisms and the carbon turnover mechanisms they facilitate. On the other hand, over the past few decades, dehalogenation processes and the microorganisms involved have attracted a wide range of research interests, primarily due to their potential applications in the bioremediation of dioxin pollutants [42]. Nevertheless, their ecological roles in carbon cycling of the natural environments have largely been overlooked [16, 41]. This study presents novel evidence showing a surprisingly high concentration of HOCs within the TOC pool of deep-sea trenches. It also highlights the enrichment and prevalence of HOCs metabolic potentials across a diverse array of microbial taxa, as well as the active degradation of typical HOCs by microorganisms in the Mariana Trench under simulated *in-situ* conditions. These results indicate that HOCs may serve as important substrates for microorganisms inhabiting trench environments, and that the metabolism of HOCs may represent a common strategy for the survival of microorganisms in harsh conditions of the hadal trenches, such as the presence of recalcitrant organic matter. The findings not only enhance the current understanding of microbial processes and survival strategies in extreme marine environments, but also underscore the significance of HOCs degradation in the microbial carbon turnover in the Mariana Trench. Given that trenches are "hotspots" for microbial carbon turnover and early diagenesis in the deep ocean [4], the observed active degradation process of HOCs in the MT sediments is believed to have great contribution to the deep-sea organic carbon remineralization. Furthermore, our results also showed that dehalogenases present in all of the microbial metagenomes from various habitats across the global

ocean (Fig. 2), with potential dehalogenation microorganisms, such as Chloroflexi, being widespread and dominant in the sediments of the hadal trenches and other deep ocean habitats [43, 44]. Dehalogenation activities have even been recorded in deep subseafloor sediments down to 358 m below the seafloor [45, 46]. Although research on this topic remains limited, existing evidence suggests that HOC metabolism is likely widespread in deep-sea environments. This implies that the microbial degradation of HOCs may play an important yet overlooked role in the global deep-sea carbon cycle, warranting further investigation in future studies.

Conversely, HOCs have been regarded as hazardous pollutants pose substantial environmental risks, and they are believed to ultimately become sequestered in deep-sea sediments due to their persistent nature [15, 20]. The findings of this study demonstrate that HOCs can serve as substrates for a diverse array of microbial taxa in deep-sea environments, where they can be effectively degraded. This provides valuable new insights into the fate and environmental implications of HOCs within deep-sea ecosystems. Additionally, the high diversity of dehalogenation microorganisms along with the unique phylogenetic traits of dehalogenase identified in hadal trenches, represent promising resources for bioremediation efforts.

Material and Methods

Sampling sites and sample processing

Sediment samples from MBR02 (11.327°N,142.188°E, water depth 10901 m),
MBR04 (10.761°N,142.274°E, water depth 5842 m), MBR05 (10.979°N, 141.950°E,
water depth 6957), and MBR06 (10.813°N,141.180°E, water depth 6477 m) of the
Mariana Trench were taken during the R/V *Tansuoyihao* TS15 cruise on December

2019 (Fig. 1). Samples were taken by a box corer attached to a Hadal Lander. The procedures of the sample processing were the same as in [10, 26]: after recovering onboard, the sediment samples were immediately subsampled using sterile polypropylene corers, and most of the cores were stored under -80 °C until analysis. The pore water was collected at 1–2 cm intervals using Rhizon samplers. Aliquots for DOM analysis were stored in a pre-combusted (525 °C for 4 h) brown glass bottle, sealed with an acid-cleaned Teflon-lined cap, and frozen at −20 °C.

Measurement of halogenated organic matter and geochemical parameters

The method for sample pretreatment and HOCs measurement followed our previous study Wei et al.,[27]. Briefly, sediment cores for HOCs measurement were thawed and divided at 2 cm interval along the vertical depth profile. The most outer layer of each depth fraction was removed to avoid halogen contamination, and the remaining sample was freeze-dried and ground. Twenty milligram sediment was then weighted out from each sample and mixed with 50 mL sodium nitrate washing solution, shaking for 2 h at 140 rpm and then stay overnight to remove inorganic halogens [27]. The sediment slurry was then filtered through a polycarbonate (PC) membrane to collect insoluble fraction of the sediment. Eighty milligram of activated carbon was added to the filtrate and shake for 2 h at 140 rpm. The activated carbon with dissolved organic matter adsorbed was collected on a PC membrane by filtration. The two PC membranes were then subject to combustion for measurement of content of insoluble and absorbable HOCs, respectively. The protocols for combustion and detection of different halogen ions by IC were the same as Wei et al., [27]. All labware utilized were carefully selected and pretreated according to Wei et al. (2024) to secure that they were free of halogen contamination. Three replicates were taken for each sample.

Triplicate sample processing blanks and analytical blanks were involved in each batch

of experiment to indicate the presence and level of the background contamination. All

measured values were calibrated by subducting the background values in blank

controls from the same batch of experiment.

Content of TOC (wt.%) was determined with a TOC analyzer (Multi N/C 3100

Analytik Jena) according to the procedure of Xu et al. [47]. Carbon isotopic composition

of TOC was determined according to Luo et al., [32], using high temperature

combustion on a Vario Pyro Cube Elemental Analyzer connected to an Isoprime 100

Continuous Flow Isotope Ratio Mass Spectrometer. Stab le isotope results were

reported using the per mil notation (δ, ‰) relative to the V-PDB standard. Pore-water

DOC concentrations were determined by a high-temperature catalytic combustion

method using a Shimadzu TOC-L total carbon analyzer with a precision of $\pm 3\%$ [9].

Inorganic nutrients (NO_3^-, NO_2^-, NH_4^+, and PO_4^{3-}) were determined using a

QuAAtro autoanalyzer (Seal Analytical, United Kingdom) [48].

Metagenomic sequencing and public data collection

Sediment cores were thawed on ice and were depth fractioned at 5 cm intervals (2019

cruise). Total genomic DNA was extracted from 10-20 g of sediments from each

depth fraction using the PowerMax® Soil Kit (MoBio Laboratories, USA), and then

purified by DNeasy® PowerClean Pro Cleanup Kit (Qiagen, Germany). DNA

fragment libraries were prepared by shearing genomic DNA from each sample and

were then subjected to metagenomic sequencing on the illumine MiSeq platform in

Majorbio Bio-Pharm Technology Co. Ltd (Shanghai, China). In total, 8 metagenomes

were successfully sequenced and generated 335.4 Gb of raw reads. The data size for

individual metagenomes ranged from 27.2 - 50.1 Gb, with an average value of 41.9

Gb. The metagenomes were combined with 46 additional deep-sequenced

metagenomes from NCBI database (including 19 metagenomes from sediments of the

Mariana Trench and 27 from other habitats of the global ocean) (Table S1), and were

co-analyzed for distribution of dehalogenase and related microorganisms.

Assembly, functional annotation, and dehalogenase genes identification

Metagenomic reads from each sample were quality filtered using Trimmomatic v.

0.38 [49] with parameters specified as "LEADING:30 TRAILING:30 CROP:90

HEADCROP:10 SLIDINGWINDOW:4:25 MINLEN:50," and were separately

assembled with metaSPAdes v. 3.15.5 (-m 700) [50]. Gene predictions were performed

against the assembled contigs using Prodigal v. 2.6.3 (-p meta) [51], and genes were then

dereplicated to construct the non-redundant gene database using CD-HIT v. 4.8.1 with

default parameters [52]. Functional annotation was performed using BlastKOALA

against the KEGG database with default parameters [53]. The gene sequences annotated

as any types of dehalogenases were extracted and confirmed using BLASTp search

against UniProt protein database [54].Only the sequences showed $\geq$ 60% identity, $\geq$ 90%

coverage and E values < 10^{-5} with existing dehalogenase sequences in the UniProt

database were selected for downstream analysis.

Relative abundance determination and phylogenetic analysis of dehalogenase

Clean reads from each metagenome were mapped back to confirmed dehalogenase

genes sequences using BWA v. 0.7.17 [55]. BWA output was then converted to a sorted

and indexed bam file using SAMtools v. 1.9 [56].The number of mapped reads and

length of each gene sequence from the bam file was utilized for calculation of the

length normalized gene abundance, which was expressed as GPM (genes per million).

The relative abundance of each type of dehalogenase was calculated by summing the

GPM values of the confirmed gene sequences.

Phylogenetic analysis was conducted to reveal the differences between the hadal

trench dehalogenase and the existing counterparts. A total of 2300 BLASTP

confirmed haloalkane dehalogenase gens sequences were extracted from the 28

Mariana Trench metagenomes, translated in to protein sequences, and co-analyzed

with 37 previously characterized haloalkane dehalogenase sequences from the

Uniport database. The sequences were aligned with MAFFT v. 7.467 [57] and trimmed

with trimAl software [58], with default parameters. A maximum-likelihood phylogenetic

tree was constructed with IQ-Tree [59] with 100 bootstrap replicates.

Genome binning and dereplication

The quality-filtered reads were mapped back to the assembled contigs using Bowtie2

(v. 2.3.4.1) [60], and coverage was determined according to the mapping results with the

jgi_summa- rize_bam_contig_depths script [60]. Metagenome binning was conducted

for assemblies longer than 2500 bp using MetaBAT v. 2.12.1 [61] with default

parameters. Quality of the MAGs was assessed by CheckM v. 1.1.2 using the line-

age_wf workflow [62], and only MAGs with completeness > 50% and contamination <

5% were kept for further analysis. Redundant bins were subsequently dereplicated

using dRep v. 2.3.2 [63] at 95% average nucleotide identity (ANI) (all other parameters

were set to the default), and MAG with the highest quality was selected from each

cluster for downstream analysis. The genome size was estimated by dividing the size

of the MAG by its estimated completeness.

Taxonomic assignment, identification of dehalogenation taxa and calculation of their relative abundance

Taxonomic classification of the MAGs was determined using GTDB-Tk, which is based on the phylogenetically calibrated Genome Taxonomy Database (GTDB) [64]. Coding sequences in the MAGs were predicted using Prodigal v. 2.6.3 with default setting [51]. Functional annotation was performed by using BlastKOALA against the KEGG database with default parameters [53]. The annotation results were manually examined to screen for the MAGs with dehalogenase annotated. The sequences of dehalogenase genes were then extracted from each candidate MAG and confirmed using BLASTp search against UniProt protein database. Only the MAGs containing gene sequences showing $\geq 60\%$ identity and $\geq 90\%$ coverage with existing dehalogenase in the UniProt database were selected as the potential dehalogenation microorganisms.

Phylogenomic tree was constructed for the confirmed dehalogenation MAGs, using the 43 universal single-copy genes (SCGs) used by CheckM [62]. Protein sequences of the SCGs were identified using HMMER v. 3.1b2 [65] with default parameters, individually aligned with MAFFT v. 7.467 [57] and then concatenated. Phylogenomic tree was constructed based on the alignment using FastTree2 v. 2.1.11 [66], with a JTT model, a gamma approximation and 100 bootstrap replicates. Archaeal genome (GCF_004799605.1) was utilized as the root. The phylogenomic tree was visualized with iTOL [67].

Relative abundances of the dehalogenation MAGs were calculated by mapping the

trimmed reads to the manually curated and refined MAGs with CoverM (v.0.6.1;

https://github.com/wwood/CoverM) in genome mode including the dereplication flag

using the default aligner Minimap2 (v.2.21; https://docs.csc.fi/apps/minimap2/) in

short-read mode, discarding unmapped reads. The final relative abundance of each

MAG is the percentage of the MAG in the mapped fraction of each sample.

Reconstruction of metabolic network for dehalogenation microorganisms

Systematic metabolic capabilities of the dehalogenation microbial taxa were explored

by systematic annotation of MAGs containing dehalogenase genes. Coding sequences

in the MAGs were annotated against the KEGG [53], Cluster of Orthologous Groups

(COG) [68], as well as PROKKA databases [69], following the steps in Liu et al. [10]. The

pathways related with energy metabolism and biogeochemical cycling were also

annotated with METABOLIC [70]. Metabolic network was reconstructed for each of the

MAGs by summarizing the annotated pathways.

Coding sequences in the MAGs were annotated against the KEGG[53], Cluster of

Orthologous Groups (COG)[68], as well as PROKKA databases[69], following the steps in

Liu et al.[10]. The pathways related with energy metabolism and biogeochemical

cycling were further annotated with METABOLIC[70]. Metabolic network was

reconstructed for each of the MAGs by summarizing the annotated pathways.

High-pressure microcosms

Microcosm set up and sampling

Sediments were taken from site MBR06 (water depth 6477 m) of the Mariana Trench

using a box corer during RV Tansuoyihao TS15 cruise on December 2019. After

removing the outer layers, around 2.0 kg sediment were taken by pre-combusted

(450 °C, 4 h) stainless steel scoops. Sediment samples were transferred into sterile

incubation bags (polypropylene) and stored under 4 °C for 20 months, which leads to

the enrichment of dehalogenation microbial taxa (relative abundance of

dehalogenation microbial taxa increased from < 20% to 53% of total mapped

metagenomic reads).

After thorough mixing with a pre-combusted stainless steel scoop, the sediments were

divided into polypropylene incubation bags and mixed with 0.2 μm filtered seawater

taken from 6000m depth of the Mariana Trench. Each of the 50 ml incubation bag

contained 15 g sediment and 30 ml filtration sterilized seawater, and then either 1.5

mg of 4-Chlorobiphenyl (4-PCB) or 3 mg of γ-Hexachlorocyclohexane (γ-HCH) was

amended as the representatives of anthropogenic HOCs with low or high numbers of

halogen atoms, respectively. No-substrate controls and no-microorganism controls

were set up simultaneously with the experimental groups. No-substrate controls are

incubation bags contained only sediment and sterilized seawater, and without

halogenated organic matter substrates. No microorganism controls are incubation bags

containing autoclaving sterilized sediments and filtration sterilized seawater, and

either type of halogenated organic substrates with the same ratios as the experiment

groups. All of the incubation bags for experimental groups and the controls were put

into high pressure incubation vessels, and were incubated under 60 MPa and 4 °C to

simulate the *in-situ* pressure and temperature. Triplicate incubation bags were taken

for each treatment at day 0, day 30, day 60, day 90, and day 270 to monitoring the

changes of organic substrates and microbial metabolisms.

4-PCB and γ-HCH measurement

Around 15 mL of sediment slurry from each incubation bags were freeze dried and

ground. After mixing with 100 µL surrogate standard solution (2,4,5,6-Tetrachlorom-

xylene, 500 mg/L), the sample was transferred into Accelerated Solvent Extraction

(ASE) cells for extraction with (1:1, v/v) (72 °C, 20 h for 4-PCB, and 65 °C, 22h for

HCH). The yielded extract from each cell was then evaporated to a final volume of 1

mL and then cleaned up in CNWBOND Florisil PR SPE cartridge (60-100 mesh, 1

g/6 mL). First, 1 g of Na_2SO_4 was added to the cartridge and 10 mL of n-hexane was

added to pre-condition the column. Then the extract was transferred into the column

and eluted by 10 mL of acetone:n-hexane (1:1). The collected effluent was

concentrated to 5 mL. Before instrumental analysis, 100 µL of the effluent was

transferred into amber GC vials, and 50 µL of injection standard (50 mg /L, 1-Bromo-

2-nitrobenzene for 4-PCB analysis, Pentachloronitrobenzene for HCH analysis) was

added to calculate recoveries.

The 4-PCB and HCH were analyzed by using gas chromatograph (Agilent 7890B;

DB-5 column, 15m× 0.32mm×0.25 µm) equipped with 63Ni-ECD detector (for

analysis of HCH) and MSD detector (for analysis of PCB). Samples (1 µL) were

injected in the splitless injection mode with high purity nitrogen as carrier gas (1

mL/min for 4-PCB analysis, and 2 mL/min for HCH analysis). For analysis of 4PCB,

the temperature of the column was set at 40°C for 2 min, then raised to 280°C

$(20°C·min^{-1})$, and finally retained at 280°C for 5 min. The injection port and detector

temperatures were kept at 270°C and 280°C, respectively. For analysis of HCH, the

temperature of the column was set at 100°C for 2 min, raised to 220°C $(15°C·min^{-1})$

and retained for 5 min, then raised to 260 °C (15°C·min^{-1}) and retained for 20 min.

The injection port and detector temperatures were kept at 220°C and 300°C,

respectively.

The quality control steps included sample processing blanks, limit of detection,

recovery tests, replicates, and standard curves. Sample processing blanks were

involved for each batch of experiment to indicate possible contamination. All blanks

were below the limits of detection (LOD). The LODs were set to be triple the standard

deviation of the blanks, which were averagely 0.1 µg/g for PCB and 0.01µg/g for

HCH. Spiked recovery tests of the standards showed acceptable recoveries ranging

from 81.80% to 85.40% for 4-PCB, with the relative standard deviation (RSD) less

than ±2.57%, and from 91.59% to 94.63 % for HCH, with the RSD less than ±

4.74% . The standard curves had high degree of linearity ($R^2 > 0.999$) for all the

compounds.

Metagenomic and metatranscriptomic analysis

Total DNA and RNA were coextracted from each of the microcosm samples using

PowerSoil Total RNA Isolation Kit and DNA Elution Accessory Kit (MoBio Lab,

United States). The concentration and quality of DNA and RNA were determined

using both Qubit 4.0 (Thermo Fisher Scientific, Waltham, USA) and Nanodrop One

(Thermo Fisher Scientific, Waltham, USA). Quality of RNAs were also determined

through the Agilent 4200 system (Agilent Technologies, Waldbronn,

Germany). All of the DNA samples were directly subject to library preparation using

the ALFA-SEQ DNA library Prep Kit (Findrop Biosafety Tech., China). The RNA

samples went through whole transcriptome amplification using RNA REPLI-g Cell

WGA & WTA Kit (Qiagen, Germany), and subject to library construction using

ALFA-SEQ RNA library Prep Kit II (Findrop Biosafety Tech., China). The DNA

libraries and RNA libraries were then subject to metagenome and metatranscriptome

sequencing, respectively, in Magigene Biotechnology Co. Ltd (Guangzhou, China)

using Illumina novaseq6000 (PE150 sequencing).

The sequencing was successful for all of the metagenomes. Metagenomic reads from

each sample were quality filtered, assembled, and genes were predicted and

dereplicated following the same procedure as mentioned earlier. Functional annotation

was performed against the KEGG, COG and PROKKA databases. The gene

sequences involved in any of the HOCs degradation pathways were extracted and

confirmed via BLASTp. Clean reads from each metagenome were mapped back to

predicted genes using BWA v. 0.7.17 and the gene length normalized relative

abundance of each gene were expressed as GPM. The relative abundances of the

dehalogenation genomes were estimated by mapping the metagenomic reads to

sequences of the corresponding MAGs and expressed as GPM.

Duplicated metatranscriptomes were successfully sequenced for both PCB and HCH

microcosm samples taken from 90[th] day. Metatranscriptomic raw reads were quality

filtered using Trimmomatic v. 0.38 [49], and non-coding RNAs were removed using

sortmerna v. 4.2.0 [71]. Clean reads of each metatranscriptome were then mapped to the

predicted protein coding genes of the metagenome from the same sample. The

transcription level of each gene was expressed as transcripts per million (TPM).

Similarly, the transcription level of the dehalogenation genomes was estimated by

mapping the metatranscriptome clean reads to sequences of the corresponding MAGs

and expressed as TPM.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (grant

numbers 42276149, 92251303) and the Shanghai Frontiers Research Fund of the

Hadal Biosphere. We also acknowledge organizers, technological teams and crews in

the cruises of M.V. Zhangjian and R.V. Tansuoyihao.

References:

1.   Herndl, G. J., Bayer, B., Baltar, F., Reinthaler, T. Prokaryotic life in the deep

ocean's water column. Annu. Rev. Mar. Sci. **15**, 461–83 (2023).

2.   Du M, et al. Geology, environment, and life in the deepest part of the world's

oceans. Innovation (New York, NY) 2, 100109 (2021).

3.   Jamieson A. The Hadal Zone: Life in the Deepest Oceans (2015).

4.   Glud RN, et al. Hadal trenches are dynamic hotspots for early diagenesis in the

deep sea. Communications Earth & Environment 2, 21 (2021).

5.   Nunoura T, et al. Hadal biosphere: Insight into the microbial ecosystem in the

deepest ocean on Earth. Proceedings of the National Academy of Sciences of the

United States of America 112, E1230-E1236 (2015).

6.   Zhou YL, Mara P, Cui GJ, Edgcomb VP, Wang Y. Microbiomes in the Challenger

Deep slope and bottom-axis sediments. Nature Communications 13, 1515 (2022).

7.   Thamdrup B, et al. Anammox bacteria drive fixed nitrogen loss in hadal trench

sediments. Proceedings of the National Academy of Sciences of the United States of

America 118, 46 (2021).

8.   Chen P, et al. Revealing the full biosphere structure and versatile metabolic

functions in the deepest ocean sediment of the Challenger Deep. Genome Biology 22,

207 (2021).

9.   Hu TC, et al. Probing sedimentary DOM in the deepest sector of Earth's surface.

Marine Chemistry 237,104033 (2021).

10. Liu RL, et al. Novel Chloroflexi genomes from the deepest ocean reveal

metabolic strategies for the adaptation to deep-sea habitats. Microbiome 10, 1-17

(2022).

11. Guan H, et al. Composition and origin of lipid biomarkers in the surface

sediments from the southern Challenger Deep, Mariana Trench. Geoscience Frontiers

10, 351-360 (2019).

12. Liu JW, et al. Proliferation of hydrocarbon-degrading microbes at the bottom of

the Mariana Trench. Microbiome 7, 47 (2019).

13. Chu MF, et al. Earthquake-enhanced dissolved carbon cycles in ultra-deep ocean

sediments. Nature Communications 14, 5427 (2023).

14. Jiang HC, et al. Three-layer circulation in the world deepest hadal trench. Nature

Communications 15, 8949 (2024).

15. Sanganyado E, Chingono KE, Gwenzi W, Chaukura N, Liu WH. Organic

pollutants in deep sea: Occurrence, fate, and ecological implications. Water Research

205, 117658 (2021).

16. Atashgahi S, Häggblom MM, Smidt H. Organohalide respiration in pristine

environments: implications for the natural halogen cycle. Environmental

Microbiology 20, 934-948 (2018).

17. Gribble GW. Naturally Occurring Organohalogen Compounds-A Comprehensive

Review. Progress in the chemistry of organic natural products 121, 1-546 (2023).

18. Liu YH, Wang L, Liu RL, Fang JS. Biogeochemical cycling of halogenated

organic compounds in the ocean: Current progress and future directions. Deep-Sea

Research Part I-Oceanographic Research Papers 205, 104237 (2024).

19. Leri AC, et al. A marine sink for chlorine in natural organic matter. Nature

Geoscience 8, 620-624 (2015).

20. Wagner CC, et al. A Global 3‐D Ocean Model for PCBs: Benchmark Compounds

724 for Understanding the Impacts of Global Change on Neutral Persistent Organic

725 Pollutants. 33, 469 - 481 (2019).

727 21. Peng GY, Bellerby R, Zhang F, Sun XR, Li DJ. The ocean's ultimate trashcan:

728 Hadal trenches as major depositories for plastic pollution. Water Research 168,

729 115121 (2020).

731 22. Sobek A, et al. Organic matter degradation causes enrichment of organic

732 pollutants in hadal sediments. Nature Communications 14, 2012 (2023).

734 23. Cui JT, et al. Occurrence of Halogenated Organic Pollutants in Hadal Trenches of

735 the Western Pacific Ocean. Environmental Science & Technology 54, 15821-15828

736 (2020).

738 24. Dasgupta S, et al. Toxic anthropogenic pollutants reach the deepest ocean on

739 Earth. Geochemical Perspectives Letters 7, 22-26 (2018).

741 25. Jamieson AJ, Malkocs T, Piertney SB, Fujii T, Zhang ZL. Bioaccumulation of

742 persistent organic pollutants in the deepest ocean fauna. Nature Ecology & Evolution

743 1, 0051 (2017).

745 26. Liu RL, et al. Bulk and Active Sediment Prokaryotic Communities in the Mariana

746 and Mussau Trenches. Frontiers in Microbiology 11, 1521 (2020).

748 27. Wei H, Liu YH, Huang LT, Wang L, Fang JS, Liu RL. Determining the

abundance, composition and spatial distribution of organohalogens in marine

sediments using combustion-ion chromatography. Marine Environmental Research

199,106626 (2024).

28. Xiao WJ, Xu YP, Canfield DE, Wenzhöfer F, Zhang CL, Glud RN. Strong linkage

between benthic oxygen uptake and bacterial tetraether lipids in deep-sea trench

regions. Nature Communications 15,3439 (2024).

29. Zhang X, et al. The hadal zone is an important and heterogeneous sink of black

carbon in the ocean. Communications Earth & Environment 3, 25 (2022).

30. Blum JD, Drazen JC, Johnson MW, Popp BN, Motta LC, Jamieson AJ. Mercury

isotopes identify near-surface marine mercury in deep-sea trench biota. Proceedings

of the National Academy of Sciences of the United States of America 117, 29292-

29298 (2020).

31. Xie JQ, et al. First evidence and potential sources of novel brominated flame

retardants and BDE 209 in the deepest ocean. Journal of Hazardous Materials

448,130974 (2023).

32. Luo M, et al. Benthic Carbon Mineralization in Hadal Trenches: Insights From In

Situ Determination of Benthic Oxygen Consumption. Geophysical Research Letters

45, 2752-2760 (2018).

33. Xiao WJ, Wang YS, Liu YS, Zhang X, Shi LL, Xu YP. Predominance of

hexamethylated 6-methyl branched glycerol dialkyl glycerol tetraethers in the Mariana Trench: source and environmental implication. Biogeosciences 17, 2135-2148 (2020).

34. Fang H, et al. Metagenomic analysis reveals potential biodegradation pathways of persistent pesticides in freshwater and marine sediments. Science of the Total Environment 470, 983-992 (2014).

35. Thomas AW, Lewington J, Hope S, Topping AW, Weightman AJ, Slater JH. Environmentally directed mutations in the dehalogenase system of Pseudomonas putida strain PP3. Archives of Microbiology 158, 176-182 (1992).

36. Zhao Y, Ma J, Zhang L, Tian K, Yao D, Yun Liu. Biodegradation mechanism of organochlorine pesticides: a review. Environ. Prot. Chem. Ind. 41, 551-558 (2021).

37. Koudelakova T, et al. Substrate specificity of haloalkane dehalogenases. The Biochemical journal 435, 345-354 (2011).

38. Ngara TR, Zeng PJ, Zhang HJ. mibPOPdb: An online database for microbial biodegradation of persistent organic pollutants. Imeta 1, e45 (2022).

39. Ruan Z, Xu X, Chen K, Qiao W, Jiang J. Recent advances in microbial catabolism of persistent organic pollutants. 60, 2763-2784 (2020).

40. Li DD, et al. Quantifying functional redundancy in polysaccharide-degrading

prokaryotic communities. Microbiome 12, 120 (2024).

41. Yang Y, et al. Roles of Organohalide-Respiring Dehalococcoidia in Carbon

Cycling. Msystems 5, e00757-e00719 (2020).

42. Adrian L, Löffler F. Organohalide-Respiring Bacteria (2016).

43. Hoshino T, et al. Global diversity of microbial communities in marine sediment.

117, 27587-27597 (2020).

44. Wu JX, et al. Biogeographic distribution, ecotype partitioning and controlling

factors of Chloroflexi in the sediments of six hadal trenches of the Paciflc Ocean.

Science of the Total Environment 880, 163323 (2023).

45. Futagami T, Morono Y, Terada T, Kaksonen AH, Inagaki F. Dehalogenation

Activities and Distribution of Reductive Dehalogenase Homologous Genes in Marine

Subsurface Sediments. Applied and Environmental Microbiology 75, 6905-6909

(2009).

46. Futagami T, Morono Y, Terada T, Kaksonen AH, Inagaki F. Distribution of

dehalogenation activity in subseafloor sediments of the Nankai Trough subduction

zone. Philosophical Transactions of the Royal Society B-Biological Sciences 368, 368

(2013).

47. Xu YP, et al. Distribution, Source, and Burial of Sedimentary Organic Carbon in

Kermadec and Atacama Trenches. Journal of Geophysical Research-Biogeosciences

126, e2020JG006189 (2021).

48. Liu R, et al. Depth-Resolved Distribution of Particle-Attached and Free-Living

Bacterial Communities in the Water Column of the New Britain Trench. Front

Microbiol 9, 625 (2018).

49. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30, 2114-2120 (2014).

50. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. Genome Research 27, 824-834 (2017).

51. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:

prokaryotic gene recognition and translation initiation site identification. Bmc

Bioinformatics 11, 119 (2010).

52. Li WZ, Godzik A. Cd-hit: a fast program for clustering and comparing large sets

of protein or nucleotide sequences. Bioinformatics 22, 1658-1659 (2006).

53. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG

Tools for Functional Characterization of Genome and Metagenome Sequences.

Journal of Molecular Biology 428, 726-731 (2016).

54. Liu WG, Schmidt B, Müller-Wittig W. CUDA-BLASTP: Accelerating BLASTP

on CUDA-Enabled Graphics Hardware. Ieee-Acm Transactions on Computational

Biology and Bioinformatics 8, 1678-1684 (2011).

55. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics 25, 1754-1760 (2009).

56. Etherington GJ, Ramirez-Gonzalez RH, MacLean D. bio-samtools 2: a package

for analysis and visualization of sequence and alignment data with SAMtools in Ruby.

Bioinformatics 31, 2565-2567 (2015).

57. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version

7: Improvements in Performance and Usability. Molecular Biology and Evolution 30,

772-780 (2013).

58. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25, 1972-

1973 (2009).

59. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.

Molecular Biology and Evolution 32, 268-274 (2015).

60. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature

Methods 9, 357-U354 (2012).

61. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. PeerJ 3, e1165

(2015).

62. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Research 25, 1043-1055 (2015).

63. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes

through de-replication. Isme Journal 11, 2864-2868 (2017).

64. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to

classify genomes with the Genome Taxonomy Database. Bioinformatics 36, 1925-

1927 (2020).

65. Eddy SR. A new generation of homology search tools based on probabilistic

inference. Genome informatics International Conference on Genome Informatics 23,

205-211 (2009).

66. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood

trees for large alignments. PLoS One 5, e9490 (2010).

67. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display

and annotation of phylogenetic and other trees. Nucleic Acids Research 44, W242-

W245 (2016).

68. Tatusov RL, et al. The COG database: an updated version includes eukaryotes.

Bmc Bioinformatics 4, 41 (2003).

69. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30,

2068-2069 (2014).

70. Zhou ZC, et al. METABOLIC: high-throughput profiling of microbial genomes

for functional traits, metabolism, biogeochemistry, and community-scale functional

networks. Microbiome 10, 33 (2022).

71. Kopylova E, Noé L, Touzet H. SortMeRNA: fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. Bioinformatics 28, 3211-3217 (2012).

 Figure legends

Figure 1. (A) Locations of sediment cores involved in the analysis of this study. The

red stars with squares indicate the stations and samples with detailed geochemical

data analyzed. (B) The geochemical parameters of the sediment cores from the

Mariana Trench, including the concentrations of adsorbable organic halogens (AOF,

AOCl, AOBr and AOX), and the corresponding insoluble organic halogens (IOF,

IOCl, IOBr, IOX and IOX), as well as total organic halogen contents (TOBr, TOCl,

TOF, and TOX). The values of total organic carbon (TOC) and their carbon isotopic

compositions ($\delta 13C$), as well the molar ratio of calculated molar ratios of organic

carbon to halogen atoms (C:X) were also shown.

Figure 2. (A) Relative abundance of different types of dehalogenases in the

metagenomes from the global ocean. The values were expressed as genes per million

(GPM). (B) Maximum-likelihood phylogenetic tree of haloalkane dehalogenase

identified from the sediment of the Mariana Trench. Red arrows indicate the location

of previously characterized haloalkane dehalogenase sequences from the Uniport

database. A sequence of epoxide hydrolase (accession no. P07099) was used as the

root.

Figure 3. (A) Maximum-likelihood phylogenomic tree of the 143 MAGs containing

genes encoding dehalogenase. An archaeal genome (accession no.

GCF_004799605.1) was utilized as the root. The colored backgrounds indicate the

phylum level classification, and different symbols indicate the types dehalogenase

genes each MAGs contained. (B) The relative abundance of each MAG in the

metagenomes from different sediment samples of the Mariana Trench. The numbers at

the lefthand side indicate the total relative abundance of the 143 MAGs.

Figure 4. The detailed pathways for the HOCs degradation and related

biogeochemical cycling by the dehalogenation microbial taxa (the 143 MAGs)

identified from the sediment of the Mariana Trench. The name of the encoding gene

involved in each step of degradation was shown, and the value in the brackets shows

the number of MAGs containing the gene. The pie charts beside the gene names

indicate the phylum level classification and proportion of the MAGs containing the

gene.

Figure 5. Degradation of typical HOCs compounds by microbial community from the

sediment of the Mariana Trench, tested using the simulated high-pressure microcosms

(A) Decreasing of both substrates with incubation time in the treatments with intact

trench sediments. (B) Relative abundance (GPM) and transcription activities (TPM)

of the MAGs containing dehalogenase during different times of the incubation. (C)

and (D), relative abundance and transcription activities of functional genes related

with pathways for degradation of HOCs.

Fig. 1

Fig. 2

970 **Fig. 3**

Fig. 4

Fig. 5