Leveraging long-read assemblies and machine learning to enhance short-read transposable element detection and genotyping

Austin Daigle[1,2]*, Logan S. Whitehouse[1,2], Roy Zhao[3], JJ Emerson[3], and Daniel R. Schrider[1]*

[1]Department of Genetics, University of North Carolina, Chapel Hill, NC 27599

[2]Curriculum in Bioinformatics and Computational Biology, University of North Carolina, Chapel Hill, NC 27599

[3]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697

*Correspondence to: drs@unc.edu and adaigle@unc.edu

Running title: Detecting TE insertions with machine learning

Keywords: transposable elements, variant calling, short-read sequencing, *Drosophila melanogaster*, genomics

ABSTRACT

Transposable elements (TEs) are parasitic genomic elements that are ubiquitous across the tree of life and play a crucial role in genome evolution. Advances in long-read sequencing have allowed highly accurate TE detection, though at a higher cost than short-read sequencing. Recent studies using long reads have shown that existing short-read TE detection methods perform inadequately when applied to real data. In this study, we use a machine learning approach (called TEforest) to discover and genotype TE insertions and deletions with short-read data by using TEs detected from long-read genome assemblies as training data. Our method first uses a highly sensitive algorithm to discover potential TE insertion or deletion sites in the genome, extracting relevant features from short-read alignments. To discriminate between true and false TE insertions, we train a random forest model with a labeled ground-truth dataset for which we have calculated the same set of short-read features. We conduct a comprehensive benchmark of TEforest and traditional TE detection methods using real data, finding that TEforest identifies more true positives and fewer false positives across datasets with different read lengths and coverages, while also accurately inferring genotypes and the precise breakpoints of insertions. By learning short-read signatures of TEs previously only discoverable using long reads, our approach bridges the gap between large-scale population genetic studies and the accuracy of long-read assemblies. This work provides a user-friendly tool to study the prevalence and phenotypic effects of TE insertions across the genome.

INTRODUCTION

Transposable elements (TEs) are genetic sequences capable of replicating themselves throughout the genome and that shape genome structure, gene expression, and evolutionary dynamics (Bourque et al. 2018; Drongitis et al. 2019; Makałowski et al. 2019; Wells and Feschotte 2020). TE insertions can impact phenotypes through the direct disruption of DNA sequences (Finnegan 1992), TE-induced chromosomal rearrangements (Montgomery et al. 1987; Montgomery et al. 1991), and changes in gene expression (Feschotte 2008; Lee 2015). Though most insertions leading to phenotypic changes are thought to be deleterious (Charlesworth and Langley 1989; Lynch 2007), numerous examples of adaptations driven by TE insertions have accumulated (reviewed in Casacuberta and González 2013; Schrader and Schmitz 2019).

To comprehensively quantify the impact of TEs on evolutionary processes, studies usually rely on tools that can detect and genotype TE insertions using genome re-sequencing data. Although recent breakthroughs in long-read sequencing have dramatically improved our ability to pinpoint and characterize TE insertions with unprecedented detail (Kirov et al. 2021; Han et al. 2022; Hoyt et al. 2022; Rech et al. 2022; Mohamed et al. 2023), these technologies remain more costly and less scalable than their short-read counterparts for the time being. Consequently, most large-scale population genetic studies rely heavily on short-read data, requiring robust computational methods for detecting and genotyping TEs from these more accessible, but inherently limited, resources.

Though numerous TE detection methods have been developed since the advent of whole-genome sequencing, most TE callers that use short-read sequence data (e.g. Illumina) follow a similar general strategy: identifying discordant read pairs, where one read maps to the reference genome and the other to a TE sequence, to identify locations of TE insertions (see Figure 1A). Additionally, some methods also use split reads, where part of a single read maps to the reference genome and another portion of that read shares homology with a TE, to precisely identify TE insertion breakpoints. This process is followed by a series of filtering steps to limit the prediction of false positives (reviewed in Makałowski et al. 2019). These approaches can be complicated by the difficulty of designing filters that apply to all read lengths and insert sizes, work for diverse TE family sequences, and do not filter out TE insertions in repetitive regions or nearby other structural variations relative to the reference genome. While these approaches have proven useful, their accuracy often varies substantially according to the properties of the input data, including read length, sequencing coverage, or the degree of repetitiveness in the genome of the species being examined (Vendrell-Mir et al. 2019; Yu et al. 2021; Verneret et al. 2024). Furthermore, because most previous studies comparing the performance of TE detectors relied on benchmark sets of simulated TE insertions (Nelson et al. 2017; Chen et al. 2023; Verneret et al. 2024) or a small number of genomes assembled with long reads (Rishishwar et al. 2017; Vendrell-Mir et al. 2019; Yu et al. 2021) to perform these assessments, our understanding of the shortfalls and relative efficacy of TE detection methods is incomplete. Simulated data in particular may not adequately represent the complexity and messiness of short-read mapping patterns around true TE insertions. On the other hand, large high-quality datasets of known TE insertions such as the long-read based population-scale *Drosophila melanogaster* assemblies and

TE annotations generated by Rech *et al.* (2022), open the possibility for more rigorous testing of

TE detection algorithms. Crucially, when both long- and short-read data are available for the

same genomes, short-read-based detection methods can also be benchmarked using a set of high-

confidence TE insertions. Moreover, the availability of such data allows us to reframe short-read

TE detection as a machine learning problem—rather than designing a method to detect what we

think the patterns of short-read mapping around a TE insertion *ought* to look like, we can train a

machine learning classifier to detect the actual signatures of known TE insertions in real

empirical datasets.

Here, we present TEforest, a machine learning method that enhances short-read TE

detection and genotyping by learning predictive features directly from high-confidence TE

insertions previously detected from long-read assemblies. First, TEforest uses a sensitive initial

scanning algorithm to identify a large set of potential TE insertions. TEforest then employs a

random forest classifier to simultaneously discriminate between true and false TE candidates and

genotype the insertions as heterozygous or homozygous. By training TEforest to examine a rich

suite of features—drawn from multiple read mapping signatures and tested across variable read

lengths and coverages—we are able to not only achieve higher performance than existing tools,

but also provide more reliable genotype predictions, precise breakpoint predictions, and accurate

allele frequency estimates. TEforest is freely available at

https://github.com/SchriderLab/TEforest.git.

METHODS

Algorithm overview

TEforest accepts as input (1) paired-end short-read fastq files, (2) a reference genome in fasta

format, (3) a TE consensus library in fasta format, and (4) a BED file detailing reference TE

locations (Figure 1B). The algorithm first identifies genomic regions that may contain non-

reference TE insertions by finding reads that map to TE consensus sequences and TEs annotated

in the reference genome. After a small number of filters are applied to the candidate insertion

sites, a comprehensive set of features summarizing read alignments within each candidate region

are computed and transformed into feature vectors. These vectors are then classified by a random

forest model as either a homozygous TE insertion, a heterozygous TE insertion, or no insertion.

These feature vectors can also be used for training a model if the true genotypes are available.

Finally, the algorithm attempts to pinpoint precise breakpoint locations using split-read evidence.

For TEs annotated in the reference genome, we use an additional random forest model trained to

detect presence/absence using the same feature vectors of the non-reference model.

Figure 1: (**A**) Diagram depicting read-mapping information used to detect non-reference TE
insertions with paired-end short reads. Short reads are aligned to the reference genome, and reads
where one read in the pair maps to the reference genome and the other to a TE sequence
(discordantly mapping reads) or reads where one pair is split between the reference and TE
sequence (split reads) are quantified. (**B**) Overview of the TEforest pipeline. Input and output
files are shown in ovals, important branching points in the pipeline are shown in diamonds, and
computational steps of the pipeline are shown in rectangles. (**C**) An example of short-read
alignment patterns around a TE insertion, displayed in IGV. Reads mapping to a TE sequence
elsewhere in the genome are shown in colors.

Algorithm to detect non-reference insertions

Discovery of regions with candidate TE insertions

To discover regions of the genome with potential TE insertions, the fastq reads are preprocessed

with *fastp* to trim adaptors and low-quality sequences (Chen 2023) and then mapped to the TE

consensus sequences as well as sequences annotated as TEs in the reference genome using the

BWA-MEM 2 algorithm, with default settings (Li and Durbin 2009; Vasimuddin et al. 2019).

Reads that map to TE sequences are then mapped back to the reference genome using the same

software/settings. Nested TE sequences, where an annotation of one TE overlaps with another

TE, are omitted to prevent errors where sequences aligning to one TE are misattributed to

another nested TE. For each TE family, the locations of read pairs where one of the two reads

maps to the TE sequence (whether it be the canonical sequence or an annotated copy of that TE

family found in the reference genome) are used to identify candidate regions for TE insertions;

we refer to these pairs as "TE-mapping read pairs". Initially, the candidate region consists of any

contiguous stretch of sites with coverage by a TE mapping read. To avoid confounding reads

mapped to reference TEs with non-reference insertions, candidate regions overlapping with a

reference TE of the same family as well as those contained entirely within a reference TE of a

different family are filtered out.

Because we observed that many false positive regions are very short in length, regions

<154 bp are also filtered out to improve computational efficiency. This length is slightly larger

than the largest read length used in our dataset and was chosen because many of these false

positive regions are the length of a single read. Shortening this filtering length did not improve

performance for shorter read lengths (not shown). To ensure that our candidate region fully

encompassed all TE-mapping read pairs around a putative TE insertion, all remaining regions are

expanded a further 200 bp in either direction. Because many TE insertions contain gaps in

coverage of TE-mapping read pairs (Figure 1C), we next merge any overlapping candidate

regions for the same TE family. The implementation of this algorithm benefitted from open

source bioinformatics packages for working with short-read alignments and genomic coordinate

ranges including SAMtools (Danecek et al. 2021), SeqKit2 (Shen et al. 2024), and

GenomicRanges (Lawrence et al. 2013).

Calculation of feature vectors

Here we expand on concepts described by Hill and Unckless (2019), who proposed the use of a feature vector as input for machine learning models to detect structural variants. This feature vector representation summarizes many aspects of the alignment of short reads to the genome calculated on a per-base level (all features are defined in Table 1). Features are calculated based on the alignment of all reads to the reference genome as well as to a TE family-specific BAM file, which contains the alignments of TE-mapping read pairs discovered during the candidate region discovery step. For each site in the candidate region a per-base pair sum is calculated for all features listed in Table 1 by extracting information from the CIGAR string, bitwise flag, coverage, and other alignment information reported by the BAM file containing the alignment of all reads to the reference genome. For each site, the value of each of these features is divided by the by the total number of reads mapping to that site. A concise vector summary of these features for a candidate region is created using the mean, standard deviation, median, and interquartile range (IQR) for each statistic across all sites in the candidate region, resulting in 21×2×4=168 feature summaries in total (21 features measured for both TE- and reference-genome mapping read pairs, and each summarized by four values) in the final vector.

Feature	Description (T=1, F=0)
Cigar1	Base matches reference.
Cigar2	Deletion has occurred in the read.
Cigar3	Insertion in one or more bases in the read, preceding this base.
Cigar4	Soft clip in the read at this position.
Cigar5	Hard clip in the read at this position.
Paired	Read has a pair.
Proper Pair	Both reads in the pair are aligned with correct orientation and distance.
Is Read1 Unmapped	First read in the pair is unmapped.
Is Read2 Unmapped	Second read in the pair is unmapped.
Is Read1 Rev Comp	First read in the pair is reverse complemented.
Is Read2 Rev Comp	Second read in the pair is reverse complemented.
Is First Read	Is the first read in the pair.
Is Second Read	Is the second read in the pair.
Split	Read is part of a split alignment.
Long Insert	Template length of this read is in the top 1% of template lengths in this region.
Short Insert	Template length of this read is in the bottom 2% of template lengths in this region.
Parallel Read	Both read and its mate align in the same direction.
Everted Read	Both read and its mate align in opposite directions.
Discordant Read	Mate of the read is not mapped to the same chromosome.
Template Length	Distance between the two ends in the read pair on the reference genome.
Quality	Quality score of the alignment at this position.

Table 1: Description of features used to describe the alignment of short reads to the reference genome. Each feature is first calculated for each basepair of each read and summed for each basepair before summary statistics (mean, median, sd, and IQR) are calculated for the feature across the region. Features are calculated using the full BAM alignment as well as a BAM file containing only reads aligning to the TE of interest.

Random forest model

When the locations of true TE insertions are known, labeled feature vectors can be used for

training a machine learning model. We used a random forest classifier, implemented in scikit-

learn (Pedregosa et al. 2011), and trained with 500 estimators (i.e. decision trees), using the Gini

impurity criterion to measure split quality. To address potential class imbalance, we

set class_weight="balanced", which adjusts weights inversely proportional to class frequencies.

The number of features to consider when looking for the best split was set to the square root of

the total number of features. All other hyperparameters were set to their default values in scikit-

learn. The possible output classes for the classifier are homozygous insertion, heterozygous

insertion, or no insertion.

Breakpoint detection

After the random forest model described above is used for TE detection, all candidate regions

classified as homozygous and heterozygous insertions are further processed to identify precise

breakpoints of TE insertions. TE-mapping read pairs are checked for split reads, and if none are

identified, then all read pairs mapping to the candidate region are checked for split reads. For

each split read, we locate the site in the candidate region where the 3'-most site in the read maps

to before the split, calling this a split-read-end. We find the top two sites with the largest number

of split-read-ends. We use these two sites as the final TE breakpoints, as most TEs make

staggered double strand breaks when inserting into the genome, creating a target site duplication

at the insertion site (Craig 2007; Linheiro and Bergman 2012). If only one site with split reads is

identified, that site is used as the insertion site. If no split reads are identified, then the center of

the candidate region is used as the breakpoint.

Algorithm to infer the presence/absence of TE insertions annotated in the reference

genome

When examining short-read data from a given individual, TEforest classifies TEs annotated in

the reference genome as present or absent in a similar manner to non-reference insertions, though

without a genotyping step. For this task, no detection of putative candidate regions or

breakpoints is needed. User-provided coordinates of reference TEs are expanded 500 basepairs

in either direction to capture information about reads surrounding the TE breakpoints. Feature

vectors are calculated as previously described, and a random forest classifier is trained and used

to label a reference TE as present or absent using the same hyperparameters used for classifying

non-reference TEs.

Testing TEforest on high-quality *Drosophila melanogaster* data

Creation of synthetic heterozygous genomes for training/testing TEforest's genotyping of non-

reference TEs

A subset of the *Drosophila melanogaster* genomes annotated with the manually curated TE

library created by Rech *et al.* (2022) were used for training and testing the TEforest model. As

this annotation only contains homozygous insertions found in the long-read assembly, we

selected six genomes with low heterozygosity (as reported in Supplemental Table 2 of Rech *et*

al. (2022) to minimize the risk of our model identifying false positive insertions that are present
in only a subset of flies from the inbred line used for DNA extraction and sequencing.
To include known heterozygous insertions in model training, we created synthetic
heterozygous genomes by combining reads from two genomes sequenced with the same read
length. To do this, reads from both genomes were first preprocessed by *fastp* then mapped to the
reference genome. A final heterozygous fastq file was created such that half the reads from
contigs 2R and 3R were taken from each genome, while all other chromosome arms were taken
from one of the genomes in the pair, with reads in each chromosome arm downsampled to the
same mean coverage. Thus, most differences between the two lines on chr2R and chr3R would
be heterozygous, while all variants on the remaining chromosome arms would be homozygous.
To train on a diverse set of read lengths, we created synthetic heterozygous genomes from
genomes sequenced with 54, 125, and 151 bp reads (one synthetic genome per read length).
230
Training and validation dataset for non-reference insertions
TEforest was trained using non-reference insertions from contigs 3L, 3R, and X and tested on
contigs 2L and 2R to ensure that no insertions used in model training were used in model testing.
Additionally, all candidate regions found inside the heterochromatic regions were filtered out as
these regions were not annotated by Rech *et al.* (2022). A full description of the numbers of
labeled candidate regions used for training and testing can be found in Table 2. Separate models
were trained and tested at target coverages of 5X, 10X, 20X, 30X, 40X, and 50X.
238

Training/testing	Read length (bp)	Insert size (bp)	# homozygotes	# heterozygotes	# candidate regions without true positives
Training	54	287	204	252	778
Training	125	208	200	272	25863
Training	151	436	202	312	4080
Testing	54	287	129	219	255
Testing	125	208	110	192	10390
Testing	151	436	108	218	1586

239
240 **Table 2:** Numbers of homozygous, heterozygous, and false positive candidate regions used as
241 input for training and testing the non-reference insertion model (50X coverage).
242
243
244
245

 Training and validation data for reference insertions

247 A model to detect reference TE presence and absence was trained using highly homozygous

248 genomes only because reference insertions tend to be homozygous present or absent. Reads were

249 downsampled to the same target coverages and training and testing was conducted on the same

250 contigs as the non-reference model. The numbers of presence and absence calls found in training

251 and testing can be found in Table 3.

Training/testing	Read length (bp)	Insert size (bp)	# present	# absent
Training	54	287	1668	1213
Training	125	208	1681	1219
Training	151	436	1554	1328
Testing	54	287	374	320
Testing	125	208	389	319
Testing	151	436	333	362

253
254 **Table 3:** Numbers of candidate regions with a TE present or absent used as input for training and
255 testing the reference insertion model (50X coverage).
256
Benchmarking other TE detectors

To compare TEforest to other TE detectors, we utilized McClintock, a meta-pipeline that

implements popular TE detection methods in a controlled workflow (Nelson et al. 2017; Chen et

al. 2023). We ran McClintock with default settings, using PoPoolationTE (Kofler et al. 2012),

PoPoolationTE2 (Kofler et al. 2016), RetroSeq (Keane et al. 2013), TEFLoN (Adrion et al.

2017), TEMP (Zhuang et al. 2014), and TEMP2 (Yu et al. 2021). These methods were chosen

based on their ability to produce more than zero calls for all read lengths tested and finish

running within five days using 8 CPU cores.

Methods were benchmarked for accuracy of TE detection, genotyping, and breakpoint

localization. To assess the overall ability to detect TEs, calls made by a TE caller within 500 bp

of a TE of the same family in the truth dataset were counted as true positives, with the rest being

counted as false positives. Overlapping TEs of the same type in the truth dataset were condensed

to avoid the double-counting of nested TEs; this was not required for nested TEs of different

types. The truth dataset provided by Rech *et al.* (2022) does not provide perfectly precise

breakpoints because it does not report the target site duplications (TSDs) produced by the

staggered cuts of transposase upon TE integration (Craig 2007; Linheiro and Bergman 2012);

rather, it provides an approximate breakpoint near the two breakpoints of the TSD. For this

reason, the center of two breakpoints predicted by each TE caller was used to calculate distances

between the truth dataset breakpoint and the prediction of the TE caller. If one breakpoint is

predicted, we simply found the distance between this breakpoint and the truth dataset breakpoint.

To assess genotyping accuracy, we calculated the F_1 scores separately for homozygous and

heterozygous genotypes and reported the macro-averaged F_1 score as the overall measure of

performance. For example, for the homozygous F_1 score, the true positives are correctly

genotyped homozygotes, false positives are heterozygotes classified as homozygotes, and false

negatives are homozygotes classified as heterozygotes, and vice-versa for the heterozygous F_1

score. We additionally performed a separate benchmarking of breakpoint accuracy and

genotyping performance for the intersection of calls made by TEforest, TEMP2, and RetroSeq at

a given coverage.

Benchmarking allele frequency accuracy

To assess the ability of TE callers to accurately predict the allele frequencies of non-reference

TEs in a population of individuals, we benchmarked the performance of all previously used TE

callers in 13 genomes from the *Drosophila* Synthetic Population Resource, sequenced with 54 bp

reads, that were included in our truth dataset. The median coverage of these genomes was 46X,

so the TEforest model trained with 50X coverage was used for this task. Only calls within the

euchromatic regions of contigs 2L, 2R, 3L, 3R, and X were used for benchmarking. As these

genomes are highly homozygous, each true positive call (a prediction of a TE insertion of the

same family within 500 bp of an annotated insertion site) was counted as a frequency of one,

with a max frequency of 13. For each insertion in the population, the true allele frequency was

subtracted from the predicted allele frequency.

RESULTS

TEforest outperforms other TE detection methods

We developed TEforest, a machine-learning tool that detects the positions of transposon

insertions in a reference genome using short reads (Figure 1 and Methods). This is accomplished

by identifying read pairs where one read maps to an annotated set of TE sequences and the other

to the reference genome (TE-mapping read pairs), extracting a comprehensive set of features

describing read-mapping patterns in those genomic regions containing TE-mapping read pairs,

then using a random forest algorithm to predict if the region contains a true TE insertion. By

comprehensively incorporating all available read mapping information from both true TE

insertions and non-insertion sites, our approach attempts to capture a comprehensive spectrum of

read-mapping patterns, thereby enabling the model to more effectively distinguish genuine

insertions from false positives. We trained and tested TEforest using the annotated TE insertions

in six *D. melanogaster* genomes sequenced with both long and short reads by Chakraborty *et al.*

(Chakraborty et al. 2019) and Rech *et al.* (2022). Because these genomes were sequenced from

inbred lines and we wished to train TEforest to be able to detect and genotype both homozygous

and heterozygous TE insertions, we created synthetic heterozygous non-reference TE insertions

by combining the reads of highly homozygous genomes for some chromosome arms, while

others remained homozygous. These data were partitioned into training and testing sets as

described in the Methods and in Table 2. In addition to benchmarking TEforest on this hold-out

test set, we benchmarked six other TE detectors available in the McClintock meta-pipeline.

Overall, TEforest demonstrated better accuracy than competing methods across our three

tested read lengths and varying coverages (Figure 2A-C; SFigure 1&2A-C). TEforest's superior

performance is largely due to an increase in recall, as most TE detectors including TEforest

called very few false positives. In general, the F_1 score (the harmonic mean of recall and

precision) increased with coverage for all callers due to an increased number of informative

reads. Interestingly, TEforest tended to approach its highest F_1 score (~0.84) at low coverages

(10-30X) before leveling off, while other callers continued to show slight increases up to 50X

coverage. However, no method achieved recall above 75% for any coverage, indicating that a

subset of TEs was undetectable with any amount of short-read data (Figure 2D; SFigure 1&2D).

In the case of TEforest, the majority of these false negatives (69%) were lost in the candidate

region detection stage instead of being mislabeled by the random forest model, which could be

due to a lack of supporting reads or proximity to an annotated reference TE resulting in the

region being filtered out (Methods).

The other TE callers tested exhibited more variable performance than TEforest across

different read lengths and coverages. When tested with 151 bp reads, TEMP2 had the next-best

F_1 score compared to TEforest from 5X-30X coverage (e.g. 0.68 at 5X coverage and 0.77 at 30X

coverage, vs. 0.71 at 5X coverage and 0.84 at 30X coverage for TEforest), before seeing a rapid

decline in recall at 40X (leading to a F_1 score of 0.62 for TEMP2 vs. 0.84 for TEforest) and 50X

coverage (F_1 of 0.58 for TEMP2 vs. 0.84 for TEforest). RetroSeq continued to steadily increase

in recall, eventually approaching but not exceeding the performance of TEforest at 50X coverage

(F_1=0.81).

Figure 2: The performance of TEforest compared to other short-read TE callers at detecting non-reference TE insertions annotated by long read assemblies of *D. melanogaster* strains. Short reads used for detection of TEs were 151 bp with ~436 bp insert sizes. True positives were defined as predicting a TE insertion of the correct TE family with any prevalence within 500 bp of the true insertion site. Performance was quantified with (**A**) F_1 scores, (**B**) precision, and (**C**) recall. The proportions of TEs that were successfully detected by TEforest, lost in candidate region detection stage, or misclassified by the random forest model for each read length are shown in Panel **D**. The mean breakpoint accuracy of TE callers for all (**E**) true positive calls or (**F**) true positive calls shared by TEforest, RetroSeq and TEMP2 was quantified by finding the distance between the center of true positive breakpoint ranges predicted by the TE callers and the breakpoint in the truth dataset.

When tested with 125 bp reads, TEMP2 was consistently the next-best caller behind TEforest (TEforest's F_1=0.80 at 50X), though it still exhibited a decline in performance when increasing coverage to 50X (F_1=0.76 at 40X and 0.70 at 50X; SFigure 1), although this was not as severe as in the 151 bp data set. In contrast to the 151 bp dataset, RetroSeq's performance lagged behind the other TE callers until 50X coverage (F_1=0.49 at 40X and 0.59 at 50X), and did not approach the F_1 score of TEforest. Interestingly, the median insert size for this dataset was 208, meaning that the paired end reads typically had no gap between them when aligned to the genome. As RetroSeq only utilizes information from discordant paired-end reads rather than information about split reads for TE detection, it may struggle to detect TEs in this context. In contrast to the 151 and 125 bp datasets, when tested on the 54 bp dataset, TEforest exhibited lower recall and F_1 scores at 5X and 10X coverage (F_1=0.58 at 5X) compared to TEMP and TEMP2, which maintained their high F_1 scores at even 5X coverage (F_1=0.80 and 0.83, respectively; SFigure 2). From 20X-50X coverage, F_1 scores were approximately equal among these callers (~0.85). We also note that in the 54 bp dataset TEMP2 did not exhibit any of the decline in performance at higher coverages that it did in the 151 and 125 bp data sets.

TEforest non-reference calls are close to the true insertion breakpoints

We additionally approximated the breakpoint accuracy of TEforest and other TE detectors by measuring the mean absolute difference between the true and predicted insertion sites for each method. For the 151 bp dataset, TEforest's mean breakpoint estimation error was <25 bp for all coverages, gradually improving to ~10 bp at 50X coverage (Figure 2E). TEforest's predictions were closer on average to the true breakpoint than all callers except for TEFLoN at lower coverages (<30X), and comparable to TEFLoN and TEMP at higher coverages. The full distribution of breakpoint distances from the truth shows that TEforest, TEMP2, and TEFLoN predict ~50% of calls 0 or 1 bp away from the annotated breakpoint, while an excess of calls >=25 bp away from the annotated breakpoint causes TEMP2 to be 2X further than TEforest from the true breakpoint on average (SFigure 3). RetroSeq had an excess of predictions at intermediate (~10 bp) and long (>25 bp) distances, which is not unexpected because it does not use split reads to narrow down its breakpoint locations. Callers also improved as the coverage increased for the 125 bp dataset, though nearly all callers performed similarly to each other in this context (SFigure 1E). TEforest produced more distant calls relative to other algorithms in the 54 bp

context, likely due to its reliance on split short reads whose paired read maps to a TE, which are

less common for shorter reads (SFigure 2E).

As coverage decreased, each method's recall decreased (although this trend was less

severe for TEforest). We hypothesized this may lead to inaccurate comparisons of callers

because more difficult to detect TEs may also be more difficult to accurately detect breakpoints

for. We compared the intersection of TEs recovered by TEforest, TEMP2, and RetroSeq, and

found that TEforest more accurately predicted breakpoint locations than TEMP2 and RetroSeq

with 151 and 125 bp reads, but for 50 bp reads was ~2X further from the true breakpoint relative

to TEMP2 and RetroSeq at 30X coverage and above (Figure 2F; SFigure 1&2F). Overall,

TEforest can be relied upon to produce calls reasonably close to the breakpoint of a TE insertion.

TEforest accurately genotypes non-reference TE insertions

We also assessed the ability of TEforest and other callers to accurately genotype known TE

insertions. Other callers estimate the fraction of genomes in the sample from which DNA was

collected that contain an insertion, yielding a prediction that can range anywhere from 0 to 1.

While some tools refer to this as the TE insertion's "frequency", to avoid conflation with

population frequency we refer to this as the insertion's "prevalence". In contrast, since we

designed TEforest to classify insertions as heterozygotes or homozygotes, it makes discrete

predictions of 0 (homozygous absent) 0.5 (heterozygous) or 1 (homozygous present). Thus, to

allow for a comparison of genotyping accuracy among methods, we transformed the predictions

made by the other methods into distinct genotypes by labeling estimates below 0.25 as

homozygous absent, between 0.25 and 0.75 as heterozygotes, and above 0.75 were called

homozygous present.

In the dataset with 151 bp reads at 30X coverage, the predicted prevalence of true

positive predictions were located near the true value for TEforest (predicted mean prevalence of

0.98 for homozygous insertions and 0.53 for heterozygous insertions) and RetroSeq

(homozygous mean prevalence=0.94, heterozygous mean=0.50), while TEMP2 systematically

underestimated the prevalence (homozygous mean=0.46, heterozygous mean=0.18), despite

having a high F_1 score of 0.77 for detecting insertions at this coverage (Figure 3A-F). The

average prevalence predicted by TEforest for homozygous true positive insertions (that is, the

average of its predicted genotypes) remained close to one across all coverages, while it tended to

misclassify heterozygotes as homozygotes at lower coverages and make more accurate

predictions at higher coverages (Figure 3I). Other callers such as RetroSeq, TEFLoN, and

PopoolationTE also had reasonable average prevalences, while TEMP and TEMP2

underpredicted the true prevalence of insertions. We also calculated a genotyping F_1 score,

where we compute F_1 scores separately for homozygous and heterozygous genotypes and

reported the macro-averaged F_1 score as the overall measure of performance. TEforest displayed

a competitive genotyping F_1 score relative to other across callers across coverages when

calculated for all of its calls (Figure 3G) or when comparing to the intersection of TE insertions

detected by TEforest, RetroSeq, and TEMP2 (Figure 3H). Despite recovering fewer known TEs,

PopoolationTE and TEFLoN performed well at the genotyping task for those TEs they were able

to detect (Figure 3G).

Figure 3: The performance of TEforest compared to other short-read TE callers at genotyping non-reference TE insertions. Short reads used for detection of TEs were 151 bp with ~436 bp insert sizes. Frequency predictions for insertions that were homozygous, heterozygous, or absent were calculated for (**A**) TEforest, (**B**) RetroSeq, and (**C**) TEMP2, where true positive absences refer to candidate regions identified by TEforest with no TE insertion. The median prevalence is shown as a black line. (**D**-**F**) The accuracy of these frequency predictions is quantified in confusion matrices, where heterozygous predictions are defined as frequency predictions between 0.25 and 0.75 and homozygous predictions are above 0.75. The genotyping F_1 score for (**G**) all true positive calls or (**H**) true positive calls shared by TEforest, RetroSeq and TEMP2 was quantified using true positive predictions of each caller. The mean for the predicted prevalence of true positive homozygous (circles) and heterozygous (triangles) insertions are shown in panel **I**, calculated using only the prevalences of true positive predictions of each caller (false positives are excluded).

For 125 bp reads with short insert sizes, patterns of genotyping were largely the same as

for the 151 bp dataset, with TEFLoN, PopoolationTE, and TEforest performing well (SFigure 4).

For example, at 30X coverage TEforest's mean prevalence was 0.96 for homozygotes and 0.55

for heterozygotes. TEMP and TEMP2 also underpredicted the prevalence of insertions at this

read length (e.g. for TEMP2 at 30X coverage, homozygous mean prevalence=0.50, heterozygous

mean=0.18). RetroSeq, which also performed poorly at detecting known TE insertions,

overpredicted the prevalence of heterozygous insertions for this dataset across all coverages (at

30X coverage, homozygous mean=1.00, heterozygous mean=0.77; SFigure 4I). Genotyping

performance for all callers was generally higher for the 54 bp reads than other read lengths

(SFigure 5), which is to be expected as a more reads are sequenced per genome to reach a given

coverage, and the space between the two reads on a sequenced fragment is larger. TEMP and

TEMP2 still underpredicted the prevalence of TE insertions, though this bias was less severe

than for the longer read lengths (SFigure 5C&I). Specifically, at 30X coverage, the mean

prevalence for homozygotes was 0.99, 0.98, and 0.94 for TEforest, RetroSeq, and TEMP2,

respectively while the mean prevalence for heterozygotes was 0.52, 0.54, 0.39.

TEforest false negatives include truncated and nested TEs

Despite its high recall relative to other methods, TEforest still failed to detect a considerable

fraction of true positive non-reference insertions (22.5% of the true positive calls in genome with

151 bp reads at 50X coverage). We hypothesized that insertions of TEs that were shorter or more

fragmented relative to the consensus TE sequence, nested inside of other TE insertions, and/or

located near other structural variants would be more difficult to detect than full-length TE

insertions in non-repetitive regions. Focusing on TEforest false negatives for the genome

sequenced with 151 bp reads at 50X coverage, we found that insertions that were not detected in

the candidate region detection stage of the algorithm were significantly shorter in total length

(Figure 4A) compared to their consensus TE sequence (SFigure 6). Lengths of TE insertions

were taken from Supplementary Table 10 from Rech *et al.* (2022). Additionally, we found that

~20% of these false negatives consisted of TEs from the families labeled as INE-1 and Gypsy-2-

Dsim in Rech *et al.* (2022), and that most of these insertions from these families were lost in the

candidate region stage (STable 1). INE-1 is a TE family that has been thought to be inactive for

millions of years (Kapitonov and Jurka 2003; Singh and Petrov 2004), and thus most copies are

expected to be old and fragmented, so low performance on this family is not surprising. In

addition, TEs nested with copies from different TE families were much more difficult to detect,

with 28/37 TEs nested with different TE families not identified by TEforest, with 20 of these lost

in the candidate region detection stage (STable 2). TEs nested with copies of the same TE family

were easier to detect (though not as easy as unnested TEs), with ~50% undetected, and all lost in

the candidate region detection stage.

Figure 4: (A) Lengths of non-reference TEs insertions that were in the genome with 151 bp with ~436 bp insert sizes, divided into classes depending on whether they were unnested and not detected in the candidate region identification stage (Candidate FN), unnested and mislabeled as absences by the random forest classifier (Classifier FN), nested and lost in either the candidate regions or classifier step (Nested FN), or successfully detected by TEforest (True Positives). Overhead bars represent the results of pairwise Wilcox rank-sum tests. **(B)** Upset plot representing the results of combining different TE callers on the F_1 Score, Precision, and Recall for detecting non-reference TEs.

Combining TE insertion detectors improves performance

To determine whether TEforest false negatives were identified by other methods and whether its improved performance stemmed from recovering insertions missed by all other methods, we evaluated the combined performance of various TE detectors in the genome with 151 bp reads at 50X coverage. We found that while TEforest was able to identify twice as many unique true positive insertions as the caller with the next highest number of unique true positives (RetroSeq), ~96% of true positive insertions found by any caller, including TEforest, were identified by at least one additional caller (SFigure 7A). Notably, of those TE insertions identified by at least one caller, only 6% were not recovered by TEforest.

While all callers produced false positives that were not found by any other callers, 23% of false positives were shared with at least one other caller (SFigure 7B). Since recall was a bigger limiting factor than precision in the F_1 score, this resulted in prediction sets consisting of the union of predictions made by a combination of callers generally outperforming a single caller in terms of F_1 as well as recall (Figure 4B). However, TEforest performed better (F_1 score=0.852, recall=0.771, precision=0.954) than many more combinations of callers than other methods. While most combinations of callers that outperformed TEforest included TEforest, including the best combination of TEforest, TEMP, and PopoolationTE (F_1 score=0.870, recall=0.813, precision=0.937), there were several combinations that did not, such as the third-best combination of TEMP, RetroSeq, and PopoolationTE (F_1 score=0.867, recall=0.802, precision=0.942). While these results are informative as to how the union of different TE call sets may be used to improve performance, in light of the variation in accuracy that some methods show across read lengths and coverages, we caution readers that the relative performance of these different combinations may change across datasets.

TEforest uses many features to make its predictions

We analyzed feature importances in TEforest's non-reference model to identify the key features driving accurate TE detection and their relative contributions (Table 4), as measured by both permutation importance and impurity-based performance. Permutation importance measures the decrease in model performance when a feature's values are randomly shuffled, reflecting the feature's predictive power (Breiman 2001). Impurity-based importance, on the other hand, quantifies the average reduction in Gini impurity achieved by a feature during decision tree

splits, highlighting its role in decision-making within the Random Forest model (Breiman 2002).

The top ten features, ranked by permutation importance, were primarily related to discordantly

mapping reads, split reads, and proper pairing of reads. In many cases the standard deviation or

interquartile range (IQR) of a feature was more important for the model than central tendency

metrics such as the mean or median. In general, features with high permutation importance had

high impurity-importance, though the relative ordering of certain features differed for each

method.

Feature	Permutation Importance	Impurity-Based Importance
TE specific Discordant Read IQR	0.002919 ± 0.000105	0.049117
Paired mean	0.001568 ± 0.000058	0.020140
Paired sd	0.001398 ± 0.000118	0.014301
TE specific Discordant Read mean	0.000952 ± 0.000058	0.037814
Proper Pair mean	0.000579 ± 0.000036	0.034105
Cigar4 sd	0.000223 ± 0.000058	0.020924
Cigar1 sd	0.000157 ± 0.000079	0.019859
Quality IQR	0.000111 ± 0.000033	0.004595
Proper Pair sd	0.000108 ± 0.000021	0.047337
Discordant Read median	0.000105 ± 0.000024	0.019143

523
Table 4: Feature importance metrics for the top non-reference model features. The table lists the
top ten features ranked by permutation importance, which measures the decrease in model
performance when feature values are randomly shuffled. Values are reported as the mean
importance ± standard deviation across permutations. Impurity-based importance reflects the
average reduction in Gini impurity achieved by the feature during decision tree splits in the
Random Forest model.
530
One particularly informative feature was the TE-specific Discordant Read IQR,

visualized in Figure 5. This feature exhibited clear stratification across model predictions:

homozygotes had the highest values, heterozygotes showed intermediate values, and absences

had the lowest values. However, overlap near the edges of these distributions highlighted the

model's reliance on additional features to maintain its high predictive performance. False

positives had higher values for this feature compared to correctly predicted negatives, but they

remained at the lower end of the range seen in true heterozygotes or homozygotes. Conversely,

false negatives exhibited lower values than true positives but still exceeded those of true

negatives. These patterns suggest that while TEforest excels at leveraging key features for

prediction, the inclusion of complementary features is crucial for resolving edge cases and

maintaining accuracy across complex scenarios.

Figure 5: The distribution of the TE-specific discordant read interquartile range (IQR), the most important feature in the non-reference TE model, across various prediction categories, representing the relationship between true and predicted zygosity classifications. Categories include true positives (TP_homo, TP_het); true negatives (TN); candidate regions without true positives (FP_homo, FP_het); false negatives (FN_homo, FN_het); and misclassifications (misclass_homo, misclass_het), which involve errors such as misclassifying homozygotes or heterozygotes. Boxplots display the distribution of the values reported for each candidate region, with outliers shown as points. 11 outlier points >1 in the TN category were cropped out of this plot to improve the visualization.

553 **TEforest is robust to read lengths not in its training set, while TE callers show unexpected**
554 **performance gains with shorter reads**

555 To evaluate whether TEforest's performance would decline when tested on read lengths not

556 included in its training set, we conducted an analysis using trimmed reads from our synthetic

557 genomes. Reads originally 151 and 125 bp in length were trimmed at the 3' end down to 100 bp

558 using *fastp* (Chen 2023), and the trimmed reads were downsampled to 30X genome coverage to

559 enable a direct comparison with the previously tested 30X model.

560 For the reads that were trimmed from 151 bp down to 100 bp, TEforest encouragingly

561 maintained an F_1 score of 0.84 despite a 50% reduction in read length (SFigure 8). The recall for

562 TEforest improved slightly from 0.75 to 0.76, while precision decreased to from 0.95 to 0.94.

563 These results suggest that TEforest's feature vectors are resilient to variations in read length,

564 preserving its ability to accurately identify TE insertions. Surprisingly, the performance of all

565 other TE callers improved with these trimmed reads, with F_1 scores increasing across the board.

566 PopoolationTE2 exhibited the largest improvement, with its F_1 score increasing from 0.32 to

567 0.46, while TEFLoN had the smallest improvement, from 0.55 to 0.57. Despite these gains, none

568 of the callers outperformed TEforest. The improvements in other callers' performance were

569 primarily driven by higher recall, though this came at the cost of slightly reduced precision in

570 most cases, with the exception of PopoolationTE2.

571 For the 125 bp reads, TEforest exhibited a slight decline in F_1 score after trimming,

572 dropping from 0.80 to 0.77 (SFigure 9). Recall decreased from 0.67 to 0.65, while precision

573 declined from 0.97 to 0.96. In contrast, the performance of other TE callers improved even more

574 dramatically with trimming than the 151 bp reads. RetroSeq showed the largest increase, with its

575 F_1 score rising from 0.33 to 0.68. Notably, TEMP2 surpassed TEforest's performance, achieving

576 an F_1 score of 0.80, up from 0.76.

577 These unexpected increases in the performance of other TE callers may be attributed to

578 their reliance on discordant and split reads that can be mapped to TE sequences. While split

579 reads are a valuable resource for precisely detecting the breakpoints of TE insertions, it is more

580 challenging to correctly map each piece of the split read compared to mapping a discordant read.

581 When reads spanning a TE insertion are trimmed, the length of the effectively unsequenced

582 portion of the fragment is increased, making it more likely that the two reads will be on either

583 side of the breakpoint. This results in an increased number of discordant reads mapping to TE

sequences (that is, non-TE mapping split reads become TE mapping discordant reads), likely

enhancing the confidence of these tools in identifying TE insertions. In contrast, TEforest

incorporates split-read information directly into its feature vectors, regardless of whether the read

mapper correctly aligns the split piece back to the TE consensus sequence. This capability of

TEforest is particularly desirable, as it allows researchers to utilize the full range of information

present in their dataset without needing to manipulate or preprocess the data extensively to

achieve optimal performance.

Allele frequency prediction: Can we use TEforest for population genetics?

To characterize the impact of any class of mutation, it is essential not only to identify these

mutations, but also to genotype them accurately enough to measure their frequency in a

population. For example, the distribution of allele frequencies in a population, also called the site

frequency spectrum (SFS) is a useful tool for understanding the population genetic forces

governing a class of mutations (Williamson et al. 2005; Eyre-Walker et al. 2006; Keightley and

Eyre-Walker 2007; Emerson et al. 2008). To accurately measure allele frequencies in a

population, a TE caller will need to successfully detect and genotype the TE in all individuals

carrying the allele (unless there are multiple detection/genotyping errors that happen to offset

one another). For example, false positive predictions could skew the SFS towards rare mutations,

potentially inflating estimates of the strength of natural selection acting against new mutations

(Emerson et al. 2008; Johri et al. 2022).

To assess whether the genotyping accuracy of TE detection methods examined here is

sufficient to produce accurate estimates of the SFS of TE insertions, we ran all previously tested

TE detection methods in 13 genomes, sequenced with 54 bp reads, from the *Drosophila* synthetic

population resource included in our truth dataset (Chakraborty et al. 2019; Rech et al. 2022). We

assessed the allele frequency prediction accuracy for each non-reference insertion in the

population by subtracting the true TE frequency from the number of true positive predictions

(Figure 6A). Due to the tendency towards false negatives instead of false positives, all TE callers

underpredicted the true TE frequency on average. TEMP and TEforest predicted frequencies

more accurately than other callers, with TEMP2 and RetroSeq also performing well at this task,

reflecting the previous benchmarking on single genomes. TEFLoN, PopoolationTE, and

PopoolationTE2 struggled at this task due to many false negative predictions. When viewing the

entire SFS for Roo (Figure 6B) and Copia (Figure 6C) insertions, which were the two TEs with

the highest number of non-reference copies in this dataset, we found most callers predict the

proportion of alleles in each frequency category fairly accurately. Perhaps in part because this

dataset is comprised of a global sample of individuals rather than a local population, the vast

majority of TE alleles were rare in the sample, limiting our ability to assess allele frequency

prediction for high frequency alleles. However, combined with our previous benchmarking

results, we find that TEforest is uniquely capable of both recovering TE insertions and obtaining

accurate allele frequency estimates for downstream population genetic inference.

Figure 6: (**A**) Performance of TEforest compared to other TE callers at correctly predicting the allele frequencies of non-reference TE insertions in 13 fully homozygous individuals from the *Drosophila* synthetic population resource. The y axis represents the average difference in allele frequency predicted by each caller to the true allele frequency at that locus. Error bars represent the standard error for each caller. The site frequency spectrum of (**B**) roo and (**C**) jockey TE insertions, which had the highest number of TE insertions in these individuals.

TEforest outperforms other methods in reference TE detection

We also trained a model to detect the presence or absence of TEs that are found in the reference genome, calculating the same feature vector as the non-reference model for each reference TE region. Since the locations of reference TEs are predefined by the user, the reference TE model does not require candidate region or breakpoint identification steps. Similar to the non-reference

model, TEforest demonstrated superior accuracy in detecting the presence or absence of reference TEs compared to competing methods across the three tested read lengths and varying coverages (SFigure 10). Interestingly, this improvement in performance was due to a better balance between precision and recall compared to other models, rather than the increased recall observed in the non-reference model. With more informative reads available to detect reference TEs (there are often unique regions within each reference TE sequence where coverage changes depending on its presence or absence), TEforest, along with TEMP and TEMP2 (both of which use the same algorithm), maintained a high F_1 score across all tested coverages. In contrast, TEFLoN and PopoolationTE2 had lower recall at 20X coverage and below.

DISCUSSION

Long-read sequencing technologies are becoming increasingly affordable and useful for analyzing TEs (Chu et al. 2021; Han et al. 2022; Rech et al. 2022; Mohamed et al. 2023), but short-read sequencing remains the more accessible option for many researchers, especially for large-scale population studies. Moreover, the vast majority of population genomic datasets that are currently available to date consist of short-read sequences. To integrate recent advancements in long-read sequencing with existing population-scale short-read sequencing datasets, we developed TEforest—a machine learning tool for detecting and genotyping transposable element (TE) insertions using short-read data. By incorporating high-quality annotations from long-read *Drosophila melanogaster* genome assemblies in the training process, TEforest leverages the precision of long-read sequencing while taking advantage of the widespread availability of short-read data. Our comprehensive benchmarking, one of the most extensive real-data evaluations to date involving six different TE detection tools, demonstrates that TEforest not only outperforms existing methods but also maintains more stable performance across various read lengths and coverages. While other tools exhibit fluctuating performance depending on sequencing parameters, such as a surprising increase in performance with shorter read lengths, TEforest consistently delivers accurate detection and genotyping of TE insertions.

Our benchmarking assessments highlight the advantages of using real data over simulated data. While simulated datasets allow controlled testing of algorithms, currently do not capture the full complexity of real genomic data, including sequencing errors, structural variations near TE insertion breakpoints or within TE sequences, and the complexity of nested TE insertions.

Early testing on real data also shaped decisions made during the development of our algorithm,

such as accounting for the fact that many TE insertion breakpoints have small gaps between the

two breakpoints rather than the classic TSD signature (see the example insertion in Figure 1C).

Such details might have been overlooked if we developed the algorithm using a simulated

dataset.

In our study, we observed that in some cases PoPoolationTE and PoPoolationTE2, did

not perform as well as other methods both in simulated datasets (also reported in Chen *et al.*

(2023) and in our real datasets (see Vendrell-Mir *et al.* (2019) for a counter-example).

Conversely, TEMP2 and RetroSeq, which have shown strong performance in other studies, also

performed reasonably well in our benchmarks.

However, our results also suggest that the combination of TE callers that performs the

best on a given dataset will vary depending on the insert size, read length, sequencing coverage,

and other species-specific parameters, and thus may be difficult to choose in practice without

prior benchmarking. The benchmarks of previous studies have shown that TE detection tools

often perform variably across different species due to differences in genome structure and TE

content. For instance, tools like MELT (Gardner et al. 2017) originally developed for use in the

human genome performed worse when applied to *Drosophila* genomes due to the mislabeling of

related TE families, while TEMP2 and RetroSeq performed well in both contexts (Yu et al.

2021). Similarly, TEPID performed well when detecting simulated *Arabidopsis* insertions, but

declined in performance when applied to *Drosophila* (Verneret et al. 2024). A potential solution

to the variable performance of different methods depending on context is to use a combination of

methods to make more confident predictions. Similar to Xu *et al.* (Xu et al. 2023), we find that

certain combinations of TE detection methods can increase performance by increasing recall

without a severe decline in precision. We also note that when evaluating different combinations

of tools, we simply took the union of all TE insertions detected by a given set of callers, and it is

possible that more complex rules for combining TE call sets from different methods could be

more effective (e.g. requiring that all calls produced by a method with lower precision be

recovered by at least one additional caller). In addition, one could optimize a model or model-

ensemble to maximize precision at the cost of recall or vice-versa depending on the use case.

The tools XTea (Chu et al. 2021) and DeepMEI (Xu et al. 2023) also use machine

learning in the context of non-reference TE detection. XTea employs random forest with a less

comprehensive feature vector for short-read genotyping, while DeepMEI utilizes a convolutional neural network (CNN) for TE detection and genotyping. However, these tools are currently only available for use in human genomes and thus cannot be benchmarked against our method on this dataset.

The success of short-read TE detection approaches like TEforest is fundamentally dependent on the availability of high-quality reference genomes from which comprehensive TE annotations can be produced and for which short-read sequencing data can be obtained. Thankfully, these resources are readily available in our tested species *D. melanogaster* (Kaminker et al. 2002; Rech et al. 2022). The performance of our approach of training on TE annotations and short-read mapping patterns from such assemblies would degrade when using poorly assembled or annotated reference genomes or species with an incomplete library of TE consensus sequences. Bearing this in mind, along with the often-unpredictable performance of TE detection tools when tested in a new species for which high quality training/benchmarking data are not available, we recommend careful evaluation of TE detection tools when analyzing TEs in non-model species, perhaps by simulating TE insertions as realistically as possible for testing (Chen et al. 2023; Verneret et al. 2024). Despite these challenges, we expect the performance and utility of short-read TE detection tools to increase as genome assemblies become more complete (Li and Durbin 2024) and tools to automatically curate TE consensus sequences and annotate them in a reference genome become more advanced (Baril et al. 2024; Orozco-Arias et al. 2024).

Our study, along with others (see Xu *et al.* (2023)), found that shorter, fragmented TE insertions are more challenging to detect than full-length insertions. As shorter TE copies are often inactivated or made reliant upon the transposition machinery encoded by other TE copies, developing extensions of TEforest that can predict whether insertions are full-length or fragmented using reads mapping to TE sequences could provide useful information about TE dynamics. While this approach could detect deletions on the edge of a TE due to changes in read-mapping patterns, its utility would be limited by the length of short reads, as they may lack coverage of the internal regions of TE insertions needed to detect internal deletions. For example, the longest paired-end reads used in this study would reach ~400 bp into each side of the TE, while some TE families can exceed 10 kb in length. Another potential extension of TEforest involves detecting low-prevalence or somatic TE insertions, which are typically more

challenging to detect due to the low number of informative reads per insertion (Yu et al. 2021); training data for such a method could be obtained by combining reads from genomes known to have different genotypes for a TE insertion to produce a pooled sample where the TE insertion is present at the desired prevalence. A machine learning algorithm like TEforest could be useful in distinguishing true insertions from false positives, provided that appropriate training data is available.

TEforest represents a significant step forward in the detection and genotyping of transposable element (TE) insertions from short-read sequencing data. By combining machine learning techniques with comprehensive feature extraction from read alignments, it consistently outperforms existing methods. As genomic resources and annotations continue to improve, approaches like TEforest—which leverage high quality ground-truth TE insertion datasets to enhance the utility of less costly sequencing data—will become increasingly robust. In turn, this will enable broader and more in-depth studies on the evolutionary and functional impact of TEs.

ACKNOWLEDGEMENTS

We thank Marta Coronado Zamora and Josefa González for their assistance with the truth dataset. AD received support from the National Institute of General Medical Sciences of the National Institutes of Health under award numbers R35GM154969 and T32GM067553, and DRS received support from the National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM138286.

AUTHOR CONTRIBUTIONS

Austin Daigle: Conceptualization, Formal analysis, Software, Visualization, Writing—original draft, Writing—review & editing. Logan S. Whitehouse: Conceptualization, Formal analysis, Software, Writing—review & editing. Roy Zaho: Conceptualization, Resources, Writing—review & editing. Roy Zaho: Conceptualization, Resources, Writing—review & editing. Daniel R. Schrider: Conceptualization, Formal analysis, Visualization, Writing—review & editing, Funding acquisition.

DATA AVAILABILITY

The code used to train and use TEforest, along with the trained models and code used for

benchmarking, are available at https://github.com/SchriderLab/TEforest.git (figshare DOI:

10.6084/m9.figshare.28278599). The MCTE library of consensus TE sequences and annotations

are avaliable at the DIGITAL.CSIC repository and can be accessed at

http://doi.org/10.20350/DIGITALCSIC/13765 and

http://doi.org/10.20350/DIGITALCSIC/13894, respectively. The short-read sequences used for

this study are avaliable in the NCBI database under the BioProject accessions

PRJNA559813 and SRP011971.

REFERENCES

Adrion JR, Song MJ, Schrider DR, Hahn MW, Schaack S. 2017. Genome-wide estimates of transposable element insertion and deletion rates in *Drosophila melanogaster*. Genome Biol Evol. 9(5):1329–1340. doi:10.1093/gbe/evx050.

Baril T, Galbraith J, Hayward A. 2024. Earl Grey: A fully automated user-friendly transposable element annotation and analysis pipeline. Mol Biol Evol. 41(4):msae068. doi:10.1093/molbev/msae068.

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS, et al. 2018. Ten things you should know about transposable elements. Genome Biol. 19(1):199. doi:10.1186/s13059-018-1577-z.

Breiman L. 2001. Random forests. Mach Learn. 45(1):5–32. doi:10.1023/A:1010933404324.

Breiman L. 2002. Manual on setting up, using, and understanding random forests V3.1.

Casacuberta E, González J. 2013. The impact of transposable elements in environmental adaptation. Molecular Ecology. 22(6):1503–1517. doi:10.1111/mec.12170.

Chakraborty M, Emerson JJ, Macdonald SJ, Long AD. 2019. Structural variants exhibit widespread allelic heterogeneity and shape variation in complex traits. Nat Commun. 10(1):4872. doi:10.1038/s41467-019-12884-1.

Charlesworth B, Langley CH. 1989. The population genetics of *Drosophila* transposable elements. Ann Rev Genet. 23(Volume 23,):251–287. doi:10.1146/annurev.ge.23.120189.001343.

Chen J, Basting PJ, Han S, Garfinkel DJ, Bergman CM. 2023. Reproducible evaluation of transposable element detectors with McClintock 2 guides accurate inference of *Ty* insertion patterns in yeast. Mob DNA. 14(1):8. doi:10.1186/s13100-023-00296-4.

Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication
using fastp. iMeta. 2(2):e107. doi:10.1002/imt2.107.

Chu C, Borges-Monroy R, Viswanadham VV, Lee S, Li H, Lee EA, Park PJ. 2021.
Comprehensive identification of transposable element insertions using multiple sequencing
technologies. Nat Commun. 12(1):3836. doi:10.1038/s41467-021-24041-8.

Craig NL. 2007. Mobile DNA: an Introduction. In: Mobile DNA II. John Wiley & Sons, Ltd. p.
1–11. https://onlinelibrary.wiley.com/doi/abs/10.1128/9781555817954.ch1.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. GigaScience.
10(2):giab008. doi:10.1093/gigascience/giab008.

Drongitis D, Aniello F, Fucci L, Donizetti A. 2019. Roles of transposable elements in the
different layers of gene expression regulation. Int J Mol Sci. 20(22):5755.
doi:10.3390/ijms20225755.

Emerson JJ, Cardoso-Moreira M, Borevitz JO, Long M. 2008. Natural selection shapes genome-
wide patterns of copy-number polymorphism in *Drosophila melanogaster*. Science.
320(5883):1629–1631. doi:10.1126/science.1158078.

Eyre-Walker A, Woolfit M, Phelps T. 2006. The distribution of fitness effects of new deleterious
amino acid mutations in humans. Genetics. 173(2):891–900. doi:10.1534/genetics.106.057570.

Feschotte C. 2008. Transposable elements and the evolution of regulatory networks. Nat Rev
Genet. 9(5):397–405. doi:10.1038/nrg2337.

Finnegan DJ. 1992. Transposable elements. Curr Opin Genet Dev. 2(6):861–867.
doi:10.1016/S0959-437X(05)80108-X.

Gardner EJ, Lam VK, Harris DN, Chuang NT, Scott EC, Pittard WS, Mills RE, Consortium T
1000 GP, Devine SE. 2017. The Mobile Element Locator Tool (MELT): population-scale mobile
element discovery and biology. Genome Res. 27(11):1916–1929. doi:10.1101/gr.218032.116.

Han S, Dias GB, Basting PJ, Viswanatha R, Perrimon N, Bergman CM. 2022. Local assembly of
long reads enables phylogenomics of transposable elements in a polyploid cell line. Nucleic
Acids Research. 50(21):e124. doi:10.1093/nar/gkac794.

Hill T, Unckless RL. 2019. A deep learning approach for detecting copy number variation in
next-generation sequencing data. G3. 9(11):3575–3582. doi:10.1534/g3.119.400596.

Hoyt SJ, Storer JM, Hartley GA, Grady PGS, Gershman A, de Lima LG, Limouse C, Halabian
R, Wojenski L, Rodriguez M, et al. 2022. From telomere to telomere: The transcriptional and
epigenetic state of human repeat elements. Science. 376(6588):eabk3112.
doi:10.1126/science.abk3112.

Johri P, Aquadro CF, Beaumont M, Charlesworth B, Excoffier L, Eyre-Walker A, Keightley PD,
Lynch M, McVean G, Payseur BA, et al. 2022. Recommendations for improving statistical
inference in population genomics. PLoS Biol. 20(5):e3001669.
doi:10.1371/journal.pbio.3001669.

Kaminker JS, Bergman CM, Kronmiller B, Carlson J, Svirskas R, Patel S, Frise E, Wheeler DA,
Lewis SE, Rubin GM, et al. 2002. The transposable elements of the *Drosophila melanogaster*
euchromatin: a genomics perspective. Genome Biol. 3(12):research0084.1. doi:10.1186/gb-2002-
3-12-research0084.

Kapitonov VV, Jurka J. 2003. Molecular paleontology of transposable elements in the
Drosophila melanogaster genome. Proc Natl Acad Sci USA. 100(11):6569–6574.
doi:10.1073/pnas.0732024100.

Keane TM, Wong K, Adams DJ. 2013. RetroSeq: transposable element discovery from next-
generation sequencing data. Bioinformatics. 29(3):389–90. doi:10.1093/bioinformatics/bts697.

Keightley PD, Eyre-Walker A. 2007. Joint inference of the distribution of fitness effects of
deleterious mutations and population demography based on nucleotide polymorphism
frequencies. Genetics. 177(4):2251–2261. doi:10.1534/genetics.107.080663.

Kirov I, Merkulov P, Dudnikov M, Polkhovskaya E, Komakhin RA, Konstantinov Z, Gvaramiya
S, Ermolaev A, Kudryavtseva N, Gilyok M, et al. 2021. Transposons hidden in *Arabidopsis*
thaliana genome assembly gaps and mobilization of non-autonomous LTR retrotransposons
unravelled by Nanotei pipeline. Plants. 10(12):2681. doi:10.3390/plants10122681.

Kofler R, Betancourt AJ, Schlötterer C. 2012. Sequencing of pooled DNA samples (Pool-Seq)
uncovers complex dynamics of transposable element insertions in *Drosophila melanogaster*.
PLoS Genet. 8(1):e1002487. doi:10.1371/journal.pgen.1002487.

Kofler R, Gómez-Sánchez D, Schlötterer C. 2016. PoPoolationTE2: Comparative population
genomics of transposable elements using pool-seq. Mol Biol Evol. 33(10):2759–2764.
doi:10.1093/molbev/msw137.

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ.
2013. Software for computing and annotating genomic ranges. PLoS Comput Biol.
9(8):e1003118. doi:10.1371/journal.pcbi.1003118.

Lee YCG. 2015. The role of piRNA-mediated epigenetic silencing in the population dynamics of
transposable elements in *Drosophila melanogaster*. PLoS Genet. 11(6):e1005269.
doi:10.1371/journal.pgen.1005269.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics. 25(14):1754–1760. doi:10.1093/bioinformatics/btp324.

Li H, Durbin R. 2024. Genome assembly in the telomere-to-telomere era. Nat Rev Genet.
25(9):658–670. doi:10.1038/s41576-024-00718-w.

Linheiro RS, Bergman CM. 2012. Whole genome resequencing reveals natural target site
preferences of transposable elements in *Drosophila melanogaster*. PLoS One. 7(2):e30008.
doi:10.1371/journal.pone.0030008.

Lynch M. 2007. The origins of genome architecture. Sunderland, Mass: Sinauer Associates.

Makałowski W, Gotea V, Pande A, Makałowska I. 2019. Transposable elements: classification,
identification, and their use as a tool for comparative genomics. In: Anisimova M, editor.
Evolutionary Genomics: Statistical and Computational Methods. New York, NY: Springer. p.
177–207. https://doi.org/10.1007/978-1-4939-9074-0_6.

Mohamed M, Sabot F, Varoqui M, Mugat B, Audouin K, Pélisson A, Fiston-Lavier A-S,
Chambeyron S. 2023. TrEMOLO: accurate transposable element allele frequency estimation
using long-read sequencing data combining assembly and mapping-based approaches. Genome
Biol. 24(1):63. doi:10.1186/s13059-023-02911-2.

Montgomery E, Charlesworth B, Langley CH. 1987. A test for the role of natural selection in the
stabilization of transposable element copy number in a population of *Drosophila melanogaster*.
Genet Res. 49(1):31–41. doi:10.1017/S0016672300026707.

Montgomery EA, Huang SM, Langley CH, Judd BH. 1991. Chromosome rearrangement by
ectopic recombination in *Drosophila melanogaster*: genome structure and evolution. Genetics.
129(4):1085–1098. doi:10.1093/genetics/129.4.1085.

Nelson MG, Linheiro RS, Bergman CM. 2017. McClintock: an integrated pipeline for detecting
transposable element insertions in whole-genome shotgun sequencing data. G3. 7(8):2763.
doi:10.1534/g3.117.043893.

Orozco-Arias S, Sierra P, Durbin R, González J. 2024 Dec 9. MCHelper automatically curates
transposable element libraries across eukaryotic species. Genome Res.
doi:10.1101/gr.278821.123. https://genome.cshlp.org/content/early/2024/12/09/gr.278821.123.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer
P, Weiss R, Dubourg V, et al. 2011. Scikit-learn: Machine learning in python. JMLR.
12(85):2825–2830.

Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H, Jamilloux V,
Quesneville H, González J. 2022. Population-scale long-read sequencing uncovers transposable
elements associated with gene expression variation and adaptive signatures in Drosophila. Nat
Commun. 13(1):1948. doi:10.1038/s41467-022-29518-8.

Rishishwar L, Mariño-Ramírez L, Jordan IK. 2017. Benchmarking computational tools for
polymorphic transposable element detection. Briefings in Bioinformatics. 18(6):908–918.
doi:10.1093/bib/bbw072.

Schrader L, Schmitz J. 2019. The impact of transposable elements in adaptive evolution. Mol
Ecol. 28(6):1537–1549. doi:10.1111/mec.14794.

Shen W, Sipos B, Zhao L. 2024. SeqKit2: A Swiss army knife for sequence and alignment
processing. iMeta. 3(3):e191. doi:10.1002/imt2.191.

Singh ND, Petrov DA. 2004. Rapid sequence turnover at an intergenic locus in *Drosophila*. Mol
Biol Evol. 21(4):670–680. doi:10.1093/molbev/msh060.

Vasimuddin Md, Misra S, Li H, Aluru S. 2019. Efficient architecture-aware acceleration of
BWA-MEM for multicore systems. In: 2019 IEEE International Parallel and Distributed
Processing Symposium (IPDPS). p. 314–324. https://ieeexplore.ieee.org/document/8820962.

Vendrell-Mir P, Barteri F, Merenciano M, González J, Casacuberta JM, Castanera R. 2019. A
benchmark of transposon insertion detection tools using real data. Mobile DNA. 10(1):53.
doi:10.1186/s13100-019-0197-9.

Verneret M, Le VA, Faraut T, Turpin J, Lerat E. 2024. Particular sequence characteristics induce
bias in the detection of polymorphic transposable element insertions. :2024.09.25.614865.
doi:10.1101/2024.09.25.614865. https://www.biorxiv.org/content/10.1101/2024.09.25.614865v2.

Wells JN, Feschotte C. 2020. A field guide to eukaryotic transposable elements. Annu Rev Genet.
54(1):539–561. doi:10.1146/annurev-genet-040620-022145.

Williamson SH, Hernandez R, Fledel-Alon A, Zhu L, Nielsen R, Bustamante CD. 2005.
Simultaneous inference of selection and population growth from patterns of variation in the
human genome. Proc Natl Acad Sci USA. 102(22):7882–7887. doi:10.1073/pnas.0502300102.

Xu X, Huang Y, Wang X, Cheng J, Yuan H, Bu F. 2023. Identification of mobile element
insertion from whole genome sequencing data using deep neural network model.
:2023.03.07.531451. doi:10.1101/2023.03.07.531451.
https://www.biorxiv.org/content/10.1101/2023.03.07.531451v1.

Yu T, Huang X, Dou S, Tang X, Luo S, Theurkauf WE, Lu J, Weng Z. 2021. A benchmark and an
algorithm for detecting germline transposon insertions and measuring de novo transposon
insertion frequencies. Nucleic Acids Res. 49(8):e44. doi:10.1093/nar/gkab010.

Zhuang J, Wang J, Theurkauf W, Weng Z. 2014. TEMP: a computational method for analyzing
transposable element polymorphism in populations. Nucleic Acids Res. 42(11):6826–6838.
doi:10.1093/nar/gku323.