Investigating Gene Expression Noise Reduction by MicroRNAs and MiRISC Reinforcement by Self-Feedback Regulation of mRNA Degradation

Shuangmei Tian[1], Ziyu Zhao[1], Meharie G. Kassie[1], Fangyuan Zhang[2], Beibei Ren[3] and Degeng Wang[1*]

[1]: Department of Environmental Toxicology, and The Institute of Environmental and Human Health (TIEHH), Texas Tech University, Lubbock, TX 79416, USA.

[2]: Department of Mathematics and Statistics, Texas Tech University, Lubbock, TX 79409-1042, USA.

[3]: Department of Mechanical Engineering, Texas Tech University, Lubbock, TX 79409-1021, USA.

[*]: Correspondence: degeng.wang@ttu.edu.

Abstract

MicroRNA (miRNA) induced silencing complex (miRISC) is the targeting apparatus and arguably rate-limiting step of the miRNA-mediated regulatory subsystem – the major noise reducing though metabolically wasteful mechanism. Recently, we reported that miRISC channels miRNA-mediated regulatory activity back onto their own mRNAs to form negative self-feedback loops, a noise-reduction technique in engineering and synthetic/systems biology. Here, we describe mathematical modeling that predicts mRNA expression noise to correlate negatively with degradation rate (K_{deg}) and noise reduction by self-feedback control of K_{deg}. We also calculated K_{deg} and expression noise of mRNAs detected in a cutting-edge total-RNA single-cell RNA-seq (scRNA-seq) dataset. As predicted, miRNA-targeted mRNAs exhibited higher K_{deg} values in conjunction with lower inter-cell expression noise. Moreover, as predicted by our self-feedback loop model, miRISC mRNAs (AGO1/2/3 and TNRC6A/B/C) exhibited further reduced expression noise. In short, mathematical-modeling and total-RNA scRNA-seq data-analysis shed insight into operational trade-off between noise reduction and metabolic/energetic expenditure in producing miRNA-targeted mRNAs destined for enhanced degradation and translational inhibition, as well as negative self-feedback loop reinforcement of miRISC – the core of miRNA-mediated noise-reduction subsystem. To our knowledge, this is the first report of concurrent mRNA degradation and expression noise analyses and of noise reduction by self-feedback control of mRNA degradation.

Introduction

MicroRNA (miRNA) mediates a gene expression regulatory subsystem that functions throughout the plant and animal kingdoms, though the animal and plant sub-systems are very different from each other[1-4]. In animals, miRNA genes can be either independent transcription units (TU) or part of the introns of other genes[1,5,6]. A miRNA gene may encode a single miRNA hairpin precursor, or a cluster of multiple precursors. In canonical miRNA biogenesis pathway, the primary transcript is cleaved into the precursor hairpins that, upon nucleus export into the cytoplasm, are further processed by Dicer and loaded into the Argonaute (AGO) proteins. In non-canonical pathways, one or more steps are skipped[7-11]. AGO proteins then remove the complementary strand of the double-stranded precursor to form the mature miRNAs. The miRNA-AGO complex binds to cognate miRNA binding sites, usually in the 3'-UTR of the target mRNAs[12], leading to moderate translation inhibition, degradation enhancement or both.

The miRNA regulatory subsystem has three segments: 1) biogenesis of mature 22-nucleotide miRNAs described above; 2) formation of the miRISC targeting apparatus, arguably the rate-limiting and thus the most critical segment of this gene-expression noise-reduction subsystem; and 3) effectors for mRNA decay and translation inhibition. MiRISC consists of the miRNA-loaded AGO protein and AGO-recruited TNRC6 protein upon miRNA binding to its cognate sites on target mRNAs. The complex channels moderate inhibitory actions onto target mRNAs via TNRC6-recruitment of the effectors, with a single miRNA binding site conferring only ~20% inhibition.

Why do the cells maintain this regulatory subsystem that seemingly violates the cellular gene expression economics? It is wasteful to expend critical metabolic and energetic resources to produce the miRNA-targeted mRNAs but then render them translationally inhibited and under enhanced degradation pressure[13]. What are the operational advantages the cells gain as a trade-off for the wasted metabolic and energetic resources?

A prominent theory is miRNA-mediated reduction of cell-to-cell gene expression fluctuation, often termed noise, such as those incurred by the stochastic transcriptional bursting[14-16], thereby enhancing robustness of cellular processes[17]. That is, expression levels of targeted mRNAs and their proteins have reduced cell-to-cell fluctuations, enhancing homeostasis and cellular homogeneity. The importance of cellular homeostasis and homogeneity is exemplified by their disruption in cancers to enable cellular adaptability to spatiotemporally unpredictable environments, leading to uncontrollable clonal growth and metastasis[18]. However, the noise-reduction capacity of the sub-system is limited when acting alone, *i.e.*, open-loop regulation[19]. Consequently, miRNAs often partner with other regulators, especially TFs, to implement common system control strategies such as feedback, known as closed-loop control[20].

We previously reported that miRISC is itself controlled directly by a negative self-feedback loop[21,22] – a proven noise-reduction technique in engineering and synthetic/systems biology[23-27]. That is, the core miRISC apparatus is controlled by a negative closed-loop, while other miRNA-targeted mRNAs by either open loops or closed loops formed with other regulators.

We looked for opportunities to investigate whether and how this self-feedback loop reinforces the stability of miRISC mRNA expression.

Single-cell RNA-seq (scRNA-seq) is a powerful extension of the bulk RNA-seq technology that gains the capacity to analyze cell-to-cell gene expression noise, and cellular sub-populations[28,29]. ScRNA-seq methods can be roughly divided into two categories. First, the traditional scRNA-seq methods, exemplified by the 10x Chromium methods[30], are typically microfluidic droplet-based and sequence the ends of RNA molecules. They are capable of economical high, and still increasing, throughput analyses, currently tens of thousands of cells in one experiment. But they have low gene detection sensitivity[29]. Low expression level mRNAs, which unfortunately include many regulatory genes such as transcription factors (TF), are likely to evade their detection in most of the analyzed cells. Second, thus, a new category of methods recently emerged to complement them[31-34]. The new methods enhance gene detection sensitivity by covering full length RNAs. They are often plate based and have much lower throughput, currently analyzing hundreds of cells in an experiment. Successful applications of the methods to analyze miRNA-mediated regulatory activity, such as gene expression noise reduction, have been reported[35-37]. Additionally, many methods have a total RNA version, that is, simultaneously analyzing the transcriptome at nascent intron-containing and mature fully spliced RNA steps. To the best of our knowledge, total RNA scRNA-seq datasets have not been explored to advance our understanding of miRNA-mediated regulatory activities.

Thus, in this study, we explored a dataset of snapTotal-seq, the latest total-RNA scRNA-seq method capable of detecting over 10,000 genes at both nascent and mature RNA steps in a cell[34]. MiRNA-targeted mRNAs exhibited low expression noise in conjunction with enhanced degradation. We also observed a further reduction of expression noises of AGO1/2/3 and TNRC6A/B/C mRNAs, supporting the role of the self-feedback loop in reinforcing miRISC – the core of the miRNA-mediated noise reduction subsystem. Additionally, AGO1/2/3 mRNA expression noise reduction occurred at the mature mRNA step, whereas TNRC6A/B/C noise reduction occurred at the nascent RNA step and was further reinforced at the mature mRNA step.

Materials and Methods

SnapTotal-seq dataset and analysis. The snapTotal-seq dataset is publicly available at the NCBI GEO database under the accession number GSE202126. We downloaded the dataset and processed it following the procedure described in the original publication[34]. Briefly, the next generation sequencing (NGS) reads were mapped to human genome assembly (GRCh37) with the STAR aligner (version 2.5.3a). Then, uniquely mapped reads were mapped to the GENCODE gene annotation (version 19) with the htseq-count software. Nascent- and mature-RNA originated reads were distinguished by the presence of intronic sequences. Nascent and mature RNA expression data was normalized separately with respective total read counts of each cell. For details, please see the original publication[34].

Quantification of nascent and mature RNA inter-cell expression level noise. We used the coefficient of variation (CV) parameter to quantify inter-cell RNA expression noise, as CV

has been routinely used for this purpose. We examined the cross-cell distributions of nascent and mature RNA expression levels to determine how to calculate this parameter.

For mature mRNAs, we observed that the normalized read counts of individual genes across the cells already followed normal distributions. So, we calculated the standard deviation (σ) (SD) and the mean (μ), and then CV_{mat} (mat: mature) for each gene as the SD-to-mean ratio:

$$CV_{mat} = \sigma/\mu$$

At the nascent RNA step, due to stochastic transcriptional bursting, RNA copy-numbers across a cell population usually follow the Poisson distribution or Poisson distribution with zero spikes, also known as zero-inflated Poisson distribution[14-16]. Consistently, we observed that normalized read counts of a gene across the cells follow log-normal (LN) distributions. We thus $\log_2$-transformed the counts into a normal distribution and calculated the SD (σ) and mean (μ) of transformed data for each gene. The normalized read counts across the cells can then be notated as $LN(\mu, \sigma^2)$. The CV_{nas} (nas: nascent) of the original LN distribution can then be calculated in the standard procedure as below:

$$Mean_{LN} = 2^{(\mu + \sigma^2/2)}$$

$$Variance_{LN} = 2^{(2\mu + 2\sigma^2)} - 2^{(2\mu + \sigma^2)}$$

$$CV_{nas} = \sqrt{Variance_{LN}/Mean_{LN}^2} = \sqrt{2^{(\sigma^2 - 1)}}$$

Mathematical model of mRNA expression and inter-cell noise without feedback regulation. In the literature, mRNA expression is usually modeled as a first-order dynamic process to derive the relationship between mRNA expression level (R), production rate (P), degradation rate (K_{deg}), steady-state expression level (R*) and inter-cell expression level stability (S)[23,38]. When no feedback is involved, the model ($f_{unreg}(R)$, unreg: not self-feedback regulated) is shown below:

$$f_{unreg}(R) = \frac{dR}{dt} = P - K_{deg}R, \quad \text{with a steady-state level at } R^* = \frac{P}{K_{deg}}.$$

To experimentally quantify mRNA production rate P, there have been multiple approaches. Since RNA splicing co-occurs with and is faster than transcription[39-41], P can be simplified as transcription rate, which is measured by the GRO-seq technology. Alternatively, P is nascent RNA expression level (R_{nas}) multiplied by nascent RNA splicing rate (K_{sp}) and can be simplified as $K_{sp}R_{nas}$[38,42]. Recently, this has been the basis of the RNA velocity analysis facilitated by total-RNA scRNA-seq methods[33,34,38]. Thus, as in many other studies[33,34,38,42], we used $K_{sp}R_{nas}$ in this study as it was measured by snapTotal-seq. Thus, for mRNAs at steady-state level,

$$R^* = \frac{K_{sp}R_{nas}}{K_{deg}} \quad \text{and } K_{deg} = \frac{K_{sp}R_{nas}}{R^*}.$$

The mRNA expression-level stability without feedback regulation (S_{unreg}) can be calculated via Taylor expansion as:

$$S_{unreg} = abs(f'_{unreg}(R^*)) = abs(-K_{deg}) = K_{deg}.$$

Thus, higher K_{deg} predicts higher expression level stability.

Modeling of mRNA expression and inter-cell noise with self-feedback regulation. The self-feedback loop adds a regulatory element to the degradation rate (K_{deg}), highlighted by the red bold text in the model ($f_{reg}(R)$, reg: self-feedback regulated) below:

$$f_{reg}(R) = \frac{dR}{dt} = P - (K_{deg} + \textcolor{red}{\boldsymbol{aR}})R,$$ where the constant a accounts for the mRNA- to-protein stoichiometry ratio and the efficiency of miRNA-mediated mRNA degradation activity.

Then, we can compute the mRNA expression-level stability with self-feedback regulation (S_{reg}) via Taylor expansion as well as regulated-to-unregulated S-ratio as below:

$$\frac{S_{reg}}{S_{unreg}} = \frac{f'_{reg}(R^*)}{f'_{unreg}(R^*)} = \frac{-K_{deg}-2aR^*}{-K_{deg}} > 1.$$

Thus, the self-feedback loop should lead to enhanced inter-cell mRNA expression level stability S ($S_{reg} > S_{unreg}$).

Statistical Analysis. The R open-source software (version 4.0.2) was used for data analysis and plotting. Standard R functions were used for sample normalization, parameter calculation, linear regression and LOESS regression.

For a weighted linear regression, we first performed an un-weighted linear regression. Then use the reciprocal of the square of the residuals as the weights to perform the weighted linear regressions.

To compare the linear regression models, we used the ANOVA function, as one model is nested inside another models. The function outputs a F-ratio and a p-value, thus statistically quantifying the improvement from adding the second predictive variable.

Results

Different gene expression patterns at the transcription/nascent-RNA and mature RNA steps due to post-transcriptional regulation. In this study, we used the terms "transcription" and "nascent RNA" as inter-exchangeable terms to denote the gene expression step. Nascent RNA expression level R_{nas} is determined by transcription rate (TR) and nascent RNA splicing rate (K_{sp}) and, at steady state, equal to the TR/K_{sp} ratio. R_{nas} and TR are thus frequently used in the literature as a measure of each other. As described in the section of Materials and Methods, we downloaded the snapTotal-seq dataset from the NCBI GEO database, mapped the reads onto the human genome and used the presence of intron sequences to distinguish nascent RNAs from their mature RNA counterparts. We then normalized the nascent and mature RNA read counts across the cells.

It has been reported by us and others that gene expression is less selective at the transcription step than the mature mRNA step[42]. We tested this pattern with genes detected at both nascent RNA and mature mRNA steps in every cell in the snapTotal-seq dataset (Fig. 1). A

Figure 1. Change of expression patterns from nascent to mature RNA steps. Comparative analyses of nascent and mature RNA expressions are shown. Genes detected at both steps in all cells were used. **A:** A comparative boxplot of nascent/pre-splicing and mature RNA expression levels shows enhanced selectivity or dispersion level at the mature RNA step. **B:** A scatter plot of nascent RNA expression level versus transcription unit (TU) length shows positive correlation between the two. A LOESS regression curve is added to illustrate the relationship. The nascent RNAs whose mature RNA counterparts have >= 60 miRNA binding sites are highlighted. **C:** A scatter plot of mature mRNA expression level versus length is shown. A LOESS regression curve is added to illustrate the loss of the correlation observed in **B**. The mRNAs with >= 60 miRNA binding sites are highlighted.

comparative boxplot shows that the normalized read counts exhibited a narrower value range and reduced dispersion at the nascent RNA step compared to the mature spliced RNA step (Fig. 1A). That is, at the nascent RNA step, the gene expression levels were more uniform. In contrast, at the mature mRNA step, a portion of the genes were expressed at much higher levels, presumably to meet near-term cellular protein production needs, whereas another portion of the genes were expressed at much lower levels.

However, the normalized read count data missed a key factor – RNA length. Since the snapTotal-seq sequencing chemistry covers full-length RNA, the length becomes one of the determining factors for the read counts. The NGS read counts (K) are known to follow the Poisson distribution:

$$P_{(k)} = \lambda^K e^{-\lambda}/K!$$

For individual RNAs, the means (λ) of the distributions should be directly proportional to their lengths. Consistently, snapTotal-seq gene detection rates were higher for longer RNAs at both nascent and mature RNA steps.

RNA length gave us another opportunity to test the lower gene expression selectivity at the transcription step. The lower selectivity should make RNA length a more prominent determining factor for the read counts at this step than the mature RNA step. Not surprisingly, we observed a significant positive correlation between the nascent RNA read count and the transcription unit (TU) length – the length of the longest possible un-spliced RNA of a gene (Fig. 1B). However, the correlation essentially disappeared at the mature RNA step (Fig. 1C). Conceivably, additional regulatory actions on RNA splicing, nucleus export and RNA degradation led to higher gene expression selectivity at mature RNA step, obscuring the impact of RNA length.

Adjusting the read counts with RNA length. Nevertheless, the full-length RNA coverage by the snapTotal-seq sequencing chemistry imposed a need to adjust the read count data by RNA length, so that the read counts of RNAs of different lengths were comparable with one another. The read count of a RNA is determined by both its expression level, *i.e.*, copy number, and its length. With everything else remaining the same, the expected read count of a RNA, *i.e.*, the mean (λ) of the Poisson distribution, should be, as discussed above, directly

Figure 2. Analyzing noises observed in the snapTotal-Seq dataset. **A and B:** comparative scatter plots of RNA expression levels at the mature mRNA (**A**) and the nascent RNA (**B**) steps in the same pair of cells, illustrating a pattern of decreasing inter-cell fluctuation as expression level increases at both steps. **C and D:** scatter plot of RNA CV versus mean expression level at the mature mRNA (**C**) and the nascent RNA (**D**) steps. The pattern of decreasing inter-cell fluctuation, which is quantified by CV, as mean expression level increases is illustrated at both steps. Weighted linear regression line is shown to illustrate the pattern. CV: coefficient of variation.

proportional to the RNA length.

Given the obvious correlation between nascent RNA read count and TU length (Fig. 1B), we adjusted the nascent read count with a LOESS regression ($\log_2$(read-counts) versus $\log_2$(TU-length)). Consequently, the residuals of the regression were used as the adjusted read counts. Regarding the mature RNA step, due to the disappearance of the correlation, we used the read counts normalized by RNA length – the RPKM value popularly used in RNA-seq data analysis.

Mathematical model of gene expression noise. As discussed in the Materials and Methods section, we followed the conventional practice of modeling the mature mRNA expression without feedback control as a first-order dynamic process and derived relationship among the expression parameters. The model predicts that higher degradation activity, *i.e.*, the K_{deg} value, should lead to higher steady-state expression level (R*) stability (S_{unreg}). S is measured experimentally in the form of its reciprocal, that is, CV that quantifies inter-cell fluctuation of R* as shown below:

$$S_{unreg} = abs(f'_{unreg}(R^*)) = abs(-K_{deg}) = K_{deg}.$$
$$CV_{unreg} = 1/S_{unreg}.$$

Thus, higher K_{deg} predicts higher expression level stability and thus lower CV values[23,38]. We explored the snapTotal-seq dataset to estimate CV and K_{deg} and to test predicted relationship between the two.

Analyzing inter-cell gene expression noise. The inter-cell fluctuation of the gene expression levels has two contributing factors. The first is the gene expression noise due to stochasticity of the gene expression process – the target of this study. The second is the noise intrinsic to experimental procedures, which the NGS library preparation and sequencing processes are known to have. It is also known that noise from the second source is higher for low-expression-level RNAs than high-expression-level RNAs. As shown in figure 2A, the difference between the mature mRNA read counts of the same genes in two cells increases as the expression level decreases. And the same trend was observed for the nascent RNA read counts (Fig. 2B). Therefore, we analyzed the inter-cell RNA expression noises in the context of

Figure 3. Scatter plot of inter-cell expression level fluctuations of mature mRNAs versus those of nascent RNAs. Mean expression level adjusted CV values, *i.e.*, the residuals of the linear regressions in figures 2C and 2D, were used for the plot. A linear regression line is shown. The equation of the linear regression and the correlation coefficient of the two variables are also shown.

mean expression levels. As shown in figures 2C and 2D, the CV value decreases linearly as the mean expression level increases at both the mature mRNA and the nascent RNA steps, with the mature mRNA step having a steeper slope than the nascent step.

A comparative analysis of nascent and mature RNA expression noise was then performed (Fig. 3). Weighted linear regressions were conducted (CV versus expression levels), with the regression lines shown in figures 2C and 2D. We used the residuals of the regressions as adjusted CV and then plotted the scaled adjusted CV in a scatter plot (Fig. 3). As expected, a positive correlation was observed (Fig. 3). However, the correlation was moderate, with a correlation coefficient of 0.315, and the slope of the linear regression is only 0.33. Thus, post-transcriptional regulatory mechanisms must have reduced the impact of propagation of nascent RNA expression noise on the mature RNA expression noise. Otherwise, the correlation should be much stronger, and the linear regression slope should be much closer to 1.

Relationship between mRNA degradation activity and inter-cell expression noise. Having both nascent and mature RNA expression levels simultaneously allowed us to test our mathematical prediction. The nascent RNA expression levels were used in the literature in lieu of transcription and/or mRNA production rates in mRNA expression modeling. Therefore, we estimated the mature RNA K_{deg} as the mature to nascent RNA expression level ratio (see Materials and Methods). However, there are intrinsic complexities that contribute to mature RNA expression noise observed in scRNA-seq datasets. The propagation of nascent RNA expression noise, which variated from gene to gene, was discussed above (Fig. 3). The cellular heterogeneity such as cell cycle phases, among many other factors, undoubtedly further enhance the expression noise and complexity; for individual RNAs, different subpopulations of the cells might have different steady-state expression levels (R*). We tested whether higher degradation activity led to reduced inter-cell mature RNA expression noise and whether the trend was strong enough to overcome these interfering factors. Our

Figure 4. High degradation activity tends to reduce mature mRNA expression noise. **A:** Linear regression analysis of data shown in figure 2C, with mRNA K_{deg} added as the second predictive variable (CV versus (mean expression level) + K_{deg}). Predicted CV values are plotted versus the mean expression levels, with the data points color coded by K_{deg}. **B:** Linear regression analysis of data shown in figure 3, with mRNA K_{deg} added as the second predictive variable (CV_{mat} versus CV_{nas} + log2(K_{deg})). Predicted CV_{mat} values are plotted versus the CV_{nas} values, with the data points color coded by K_{deg}. Mat: mature. Nas: nascent.

approach was a linear regression of the data shown in figure 2D with K_{deg} added as the second predictive variable. The result is schematically shown in figure 4A. Despite these complexities, the analysis revealed a clear trend of decreasing CV values along increasing K_{deg} values. ANOVA comparison of the regression models determined that adding K_{deg} as the second predictive variable led to significant improvement (F_1 = 16.8, p-value = 4.37E-5).

Subsequently, we tested this trend via a comparative analysis of expression-level adjusted nascent and mature RNA CV values that, as shown in figure 3, had a moderately positive correlation with each other. Our approach was a linear regression with K_{deg} added as the second predictive variable. The result is schematically shown in figure 4B. The regression predicted a clear trend that mature RNAs with high K_{deg} values tended to have lower CV values relative to nascent RNAs (Fig. 4B). ANOVA comparison of the regression models determined that adding K_{deg} as the second predictive variable led to significant improvement (F_1 = 12.4, p-value = 0.00045).

Taken together, these results supported the notion that mature RNA degradation activity exerted prominent impact on inter-cell mature RNA expression noise, validating our mathematical model.

Reduced expression levels and inter-cell noises of miRNA-targeted mRNAs. Encouraged by these observations, we next explored whether miRNA-mediated target mRNA degradation and inter-cell expression noise reduction can be simultaneously assessed. Performing such analysis with traditional high-throughput scRNA-seq has been technically challenging due to the low sensitivity, _i.e._, low gene detection rate, of these scRNA-seq methods. MiRNA-targeted mRNAs are usually not expressed at high levels, thus evading detection by these methods. SnapTotal-seq substantially alleviates this low sensitivity issue. And it exhibits higher gene detection rate for long RNAs[34]. It is thus well-suited for studying regulatory activity mediated by miRNAs, whose target mRNAs also tend to be longer.

We calculated the count of unique evolutionarily conserved miRNA binding sites for each mRNA. Indeed, most top miRNA-targeted mRNAs, _i.e._, those with highest binding site counts, were detected by this scRNA-seq method. We identified 23 mRNAs with 60 or more

Figure 5. MiRNA targeted mRNAs have lower expression levels and expression noises, but their nascent RNA counterparts do not. **A:** Re-plot of figure 2C, with the data points color coded by miRNA binding site counts. **B:** Re-plot of figure 2D, with the data points color coded by miRNA binding site counts.

conserved binding sites. 16 of them were detected in every cell at the mRNA step, and 10 of them at the nascent RNA step. 8 of them were detected in every cell at both steps. As shown in figure 1C, the 8 miRNA-target mRNAs exhibited lower-than-expected expression levels. The pattern was specific for the mature mRNA step. At nascent RNA step, their counterpart RNAs exhibited normal expression levels (Fig. 1B).

To simultaneously analyze miRNA-mediated target mRNA degradation and expression noise reduction, we re-plotted the mature mRNA CV values versus length-adjusted mean expression level shown in figure 2C, but with the data points color-coded by the miRNA binding site counts (Fig. 5A). The plot revealed that mRNA expression levels decreased in a binding site count dependent manner, and so did the inter-cell expression noise. Both trends were evident without the need for regression analysis (Fig. 5A).

We directly analyzed miRNA-mediated target mRNA degradation via a comparative analysis of nascent and mature RNA expression levels. Our approach was a linear regression with $\log_2$(miRNA binding site count) added as the second predictive variable. The result is schematically shown in figure 6A. The regression predicted a clear trend that mature RNAs with high binding site counts tended to have lower expression levels compared to nascent RNAs (Fig. 6A). ANOVA comparison of the regression models demonstrated a significant improvement in the predictive power of the model when including the second predictive

Figure 6. Impact of miRNA-mediated regulation on target mRNA expression levels **(A)** and noise **(B)** revealed by linear regression analyses. **A:** Mature mRNA expression level versus nascent RNA expression level linear regression, with miRNA binding site count added as the second predictive variable. Predicted mature mRNA expression level was plotted versus the nascent RNA expression level, with the data points color coded by miRNA binding site counts. **B:** Linear regression analysis of data shown in figure 3, with miRNA binding site count added as the second predictive variable (CV_{mat} versus $CV_{nas} + \log_2$(Site Count)). Predicted CV_{mat} values are plotted versus the CV_{nas} values, with the data points color-coded by miRNA binding site count.

variable ($F_1 = 143.34$, p-value < 2.2E-16).

We also analyzed miRNA-mediated target mRNA expression noise reduction via a comparative analysis of nascent and mature RNA expression noise. We re-ran the two linear regressions shown in figures 3 and 4B, but incorporating $\log_2$(miRNA binding site count) as the second predictive variable. The result is schematically shown in figure 6B. The regression

analysis revealed a clear trend that mature RNAs with higher binding site counts exhibited lower expression noise compared to nascent RNAs (Fig. 6B). ANOVA comparison of the regression models determined that adding the second predictive variable led to significant improvement ($F_1 = 133$, p-value $< 2.2E-16$).

Thus, our combination of mathematical modeling and snapTotal-seq data analysis confirmed that miRNA-targeted mRNAs should have reduced inter-cell expression noise. To the best of our knowledge, this is the first report of using total-RNA scRNA-seq data to investigate miRNA-targeted mRNA degradation and expression noise reduction simultaneously. Next, we delved into the functional implications of our previous observation that major components of miRNA-mediated regulatory pathway themselves are top miRNA targets[21,22].

The miRISC negative self-feedback loop. We previously uncovered the AGO and TNRC6 negative self-feedback loops while analyzing miRNA binding site distribution[21,22]. Briefly, when human mRNAs were ranked in descending order by their unique miRNA binding site counts, AGO1/2/3 and TNRC6A/B/C were all highly ranked. AGO1 was ranked at the 5th, AGO2 the 6th, AGO3 the 11th (tied) and TNRC6B the 27th; TNRC6C and TNRC6A were respectively ranked, though not as high but both within the top 4%, at the 446th and the 800th. This dataset gave us an opportunity to assess the function of this miRISC self-feedback loop, as AGO1/2/3 and TNRC6A/B/C expressions were detected at both nascent and mature mRNA steps.

Figure 7. Reduced expression levels and noises of mRNAs with >= 60 miRNA binding sites. Re-plot of figure 2C, with mRNAs with >= 60 binding sites highlighted as blue data points. AGO1/2/3 mRNAs, which also have >60 binding sites, are highlighted as red data points. TNRC6A/B/C mRNA, which have < 60 binding sites, are highlighted as green data points.

As discussed in the Materials and Methods section, we mathematically modeled the AGO/TNRC6 negative self-feedback loop as adding a stabilizing element (αR) to K_{deg}. When higher-than-intended expression levels of a feedback-controlled mRNA leads to higher protein expression levels, the feedback path would channel higher miRNA-mediated degradation activity back onto the same mRNA, reducing expression levels back to normal level (R*). And the opposite would happen when the controlled mRNA is expressed at lower-than-expected levels. Impacts of the feedback element (αR) on S and CV are predicted below:

$$\frac{S_{reg}}{S_{unreg}} = \frac{f'_{reg}(R^*)}{f'_{unreg}(R^*)} = \frac{-K_{deg} - 2aR^*}{-K_{deg}} > 1.$$

$$\frac{CV_{reg}}{CV_{unreg}} = \frac{S_{unreg}}{S_{reg}} < 1.$$

Thus, the self-feedback loop should further reduce the inter-cell expression noise (CV) of AGO1/2/3 and TNRC6A/B/C mRNAs. As shown in figure 7, this was indeed observed. AGO1/2/3 mRNAs all have low noise levels in comparison to other mRNAs with 60 or more conserved miRNA binding sites and with similar mean expression levels. TNRC6A/B/C mRNAs, though having less than 60 binding sites, also have relatively low noise levels.

Figure 8. Nascent RNA counterparts of mRNAs with >= 60 miRNA binding sites have normal expression levels and noises. Re-plot of figure 2D. Counterparts of mRNAs with >= 60 binding sites are

Confirmation that miRNAs exert their regulatory activities at the mature mRNA step. Having the expression levels at nascent and mature mRNA steps simultaneously gave us an opportunity to determine the step, at which gene expression noise reduction occurs for individual genes. As discussed above, we already observed reduced expression levels of miRNA-targeted mRNAs (Fig. 1C), but not for their counterparts at the nascent RNA step (Fig. 1B). We investigated whether this pattern was also applicable for gene expression noise reduction, by re-plotting figure 2D and highlighting the counterparts for miRNA-targeted mRNAs (with >= 60 binding sites) as red data points (Fig. 8). As expected, the nascent-RNA counterparts exhibit normal gene expression noise levels (Fig. 8). And the pattern is generic to the whole transcriptome (Fig. 5B). The trends of decreasing expression levels and noise as miRNA binding site count increases observed at the mature RNA step (Fig. 5A) were absent at the nascent RNA step (Fig. 5B). The results demonstrated the power of total-RNA scRNA-seq in studying miRNA and other post-transcriptional regulatory mechanisms.

Differential control of AGO1/2/3 and TNRC6A/B/C RNA inter-cell expression noise. Next, we focused on the AGO1/2/3 and TNRC6A/B/C RNAs. As other mRNAs with 60 or more conserved miRNA binding sites, AGO1/2/3 mRNAs are controlled at the mature mRNA step. They did not show highly reduced expression levels and noises at the nascent RNA step (Fig. 9A) but did so at the mature mRNA step (Fig. 9B). However, TNRC6A/B/C already exhibited low inter-cell expression noise levels at the nascent RNA step (Fig. 9A). The self-feedback loop reinforces the low noise levels at the mature mRNA step (Fig. 9B).

Discussion

The importance of the miRNA-mediated gene expression regulation subsystem is underscored by, among many others, its evolutionary conservation[1-4], pathogenic mutations in diseases[43-45] and the recent award of the 2024 Nobel Prize in Physiology or Medicine to its discovery[46]. While entering this research area, we have expressed curiosity on why this subsystem is maintained so ubiquitously, despite that it seems metabolically and energetically wasteful[21,45,47]. Via the cell-computer analogy that we previously examined[48,49], we speculated its role in alleviation of operational latency caused by the time delay from transcription to translation.

One of the key functional advantages attributed to the miRNA regulatory mechanism is the enhancement of gene expression level stability. In this study, we took advantage of a total RNA scRNA-seq dataset, to the best of our knowledge for the first time, to simultaneously examine miRNA-mediated mRNA degradation and expression noise reduction. Although our analysis is limited to the mRNA step, it can be argued that the reduction in inter-cell mRNA expression noises is propagated to downstream steps, leading to reduced expression noise at

Figure 9. Inter-cell expression noise reductions of TNRC6 mRNAs occur at the nascent RNA step and were reinforced at mRNA step, whereas those of AGO mRNAs occur at the mRNA step. **A:** re-plot of figure 2D, with AGO1/2/3 mRNAs highlighted as red data points and TNRC6A/B/C mRNAs as blue data points. **B:** re-plot of figure 2C, with AGO1/2/3 mRNAs heighted as red data points and TNRC6A/B/C mRNA as blue data points.

the translation and protein steps.

This gain of operational advantage, in exchange for maintaining the miRNA-mediated gene expression regulatory subsystem, aligns with the evolutionary co-emergence of this subsystem with multicellularity, that is, its ubiquitous occurrence in such species.

Organismal-level functionalities and processes of multicellular species depend on cellular homogeneity and robust dynamic state transitions, defectives of which are often pathogenic. At the cellular level, stable steady-state gene expression level is central to cellular homogeneity. For instance, as discussed in the Introduction section, a major hallmark of cancer is the loss of this homogeneity to enable cancer cell clonal growth and metastasis. Not surprisingly, miRNAs have been shown to be globally depleted in cancers[50]. Furthermore, the DICER1 syndrome, caused by a heterozygous mutation of the DICER1 gene, is associated with elevated cancer risks[51]. Thus, to optimize multicellular operations, such organisms maintain the miRNA-mediated regulatory subsystem at the expense of the seemingly wasteful metabolic and energetic expenditure to produce miRNA-targeted mRNAs.

Our results also supported the noise reduction function of the miRISC negative self-feedback regulation loops. The AGO and the TNRC6 proteins form the miRNA targeting apparatus and the core of the miRISC complex, and their mRNAs are controlled by self-feedback loops[21,22]. The AGO proteins process the double-stranded precursors, which is passed onto them from DICER1, into mature miRNAs and orient them for target binding[52]. Upon targeting binding, the miRNA-AGO complex recruits the TNRC6 proteins, which subsequently recruit downstream effecter proteins[53]. Thus, AGO and TNRC6 channel the miRNA-mediated regulatory actions on the target mRNAs. Since AGO and TNRC6 mRNAs themselves are top miRNA targets, they channel significant regulatory actions back onto their own mRNAs to form close negative self-feedback loops. We observed that the inter-cell expression noise of AGO1/2/3 and TNRC6A/B/C mRNAs was further reduced, which is consistent with the feedback loop acting as a common noise reduction mechanism.

To the best of our knowledge, the miRISC negative self-feedback loop is the first that adds the feedback regulatory element onto mRNA degradation activity (K_{deg}). Most reported gene expression feedback loops add the regulatory elements onto transcription activity, and so are most of the loops in engineered regulatory circuits in synthetic biology. One exception is the feedback control of the nuclear export of its own nascent RNA by the HIV Rev protein, which reduces mRNA expression noise and is a critical HIV pathogenesis factor[54]. Additionally, the ribonucleoprotein particles (RNP) often feedback control the mRNAs for their own proteins[55]. However, none of feedback loops have been shown to directly add the regulatory element to mRNA K_{deg}. Nevertheless, the feedback control appears to be a ubiquitous regulatory mechanism throughout the entire gene expression process.

The differential control of AGO and TNRC6 mRNA inter-cell expression noise – AGOs at the mature mRNA step and TNRC6s at the nascent RNA step – is intriguing. Conceivably, it is related to the functional difference between the AGOs and the TNRC6s. Even though both are major miRISC components, they do have differences in other functional aspects. For instance, in addition to being part of the miRISC, TNRC6 proteins are considered the p-body scaffold proteins[56], while AGO proteins are not. Multiple RNA-silencing pathways converge on TNRC6[57]. Additionally, AGO1/2/3 proteins are well-known to be associated with ribosomes (both monosomes and polysomes)[58]; they were used to be also known/named as eukaryotic translation initiation factor 2Cs, EIF2C1/2/3. To the best of our knowledge, TNRC6 proteins have not been shown to do so.

However, another total RNA scRNA-seq dataset that traces the cells as they go through a dynamic physiological process is necessary to investigate another functional aspect of the self-feedback loop – ensuring smooth dynamic state transitions. In this context, a transition means a shift from one steady-state expression level to another. For example, when the transcription rate of a gene increases while the cognate mRNA K_{deg} remains the same, the mRNA steady-state expression level should increase accordingly. According to the control theory, a feedback loop should help to ensure a smooth transition process[27]. A dataset covering a cell population at a single condition or time-point is not sufficient for analyzing such dynamic processes.

In summary, this study took advantage of a total-RNA scRNA-seq dataset, whose NGS sequencing chemistry covers full-length RNAs. This enabled higher detection sensitivity, more than 10,000 genes in each cell, at both the nascent RNA and the mature mRNA steps of the gene expression process. We were able to investigate simultaneously the miRNA-mediated target mRNA degradation and inter-cell expression noise reduction, as well as the function of the miRISC negative self-feedback loop. This study demonstrated the power of the total RNA scRNA-seq technologies for such studies, paving the ground for comparative study of wild-type and isogenic mutant cells as well as cells at different stages of dynamic physiological processes.

Acknowledgement

This research was funded by NIGMS NIH, grant number R15GM147858, to D.W., and by Cancer Prevention and Research Institute of Texas (CPRIT), grant number RP220600, to D.W..

References

Bartel, D. P. Metazoan MicroRNAs. *Cell* **173**, 20-51 (2018). https://doi.org:10.1016/j.cell.2018.03.006

Dong, Q., Hu, B. & Zhang, C. microRNAs and Their Roles in Plant Development. *Front Plant Sci* **13**, 824240 (2022). https://doi.org:10.3389/fpls.2022.824240

Reinhart, B. J., Weinstein, E. G., Rhoades, M. W., Bartel, B. & Bartel, D. P. MicroRNAs in plants. *Genes Dev* **16**, 1616-1626 (2002). https://doi.org:10.1101/gad.1004402

Voinnet, O. Origin, biogenesis, and activity of plant microRNAs. *Cell* **136**, 669-687 (2009). https://doi.org:10.1016/j.cell.2009.01.046

Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol* **15**, 509-524 (2014). https://doi.org:10.1038/nrm3838

Kim, Y. K. & Kim, V. N. Processing of intronic microRNAs. *EMBO J* **26**, 775-783 (2007). https://doi.org:10.1038/sj.emboj.7601512

Ruby, J. G., Jan, C. H. & Bartel, D. P. Intronic microRNA precursors that bypass Drosha processing. *Nature* **448**, 83-86 (2007). https://doi.org:10.1038/nature05983

Berezikov, E., Chung, W. J., Willis, J., Cuppen, E. & Lai, E. C. Mammalian mirtron genes. *Mol Cell* **28**, 328-336 (2007). https://doi.org:10.1016/j.molcel.2007.09.028

Cifuentes, D. *et al.* A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science* **328**, 1694-1698 (2010). https://doi.org:10.1126/science.1190809

Cheloufi, S., Dos Santos, C. O., Chong, M. M. & Hannon, G. J. A dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature* **465**, 584-589 (2010). https://doi.org:10.1038/nature09092

Yang, J. S. *et al.* Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc Natl Acad Sci U S A* **107**, 15163-15168 (2010). https://doi.org:10.1073/pnas.1006432107

Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA target sites in mammalian mRNAs. *Elife* **4** (2015). https://doi.org:10.7554/eLife.05005

Ilker, E. & Hinczewski, M. Bioenergetic costs and the evolution of noise regulation by microRNAs. *Proc Natl Acad Sci U S A* **121**, e2308796121 (2024). https://doi.org:10.1073/pnas.2308796121

Senecal, A. *et al.* Transcription factors modulate c-Fos transcriptional bursts. *Cell Rep* **8**, 75-83 (2014). https://doi.org:10.1016/j.celrep.2014.05.053

Chong, S., Chen, C., Ge, H. & Xie, X. S. Mechanism of transcriptional bursting in bacteria. *Cell* **158**, 314-326 (2014). https://doi.org:10.1016/j.cell.2014.05.038

Khetan, N. *et al.* Single-cell RNA sequencing algorithms underestimate changes in transcriptional noise compared to single-molecule RNA imaging. *Cell Rep Methods* **4**, 100933 (2024). https://doi.org:10.1016/j.crmeth.2024.100933

Ebert, M. S. & Sharp, P. A. Roles for microRNAs in conferring robustness to biological processes. *Cell* **149**, 515-524 (2012). https://doi.org:10.1016/j.cell.2012.04.005

Dagogo-Jack, I. & Shaw, A. T. Tumour heterogeneity and resistance to cancer therapies. *Nat Rev Clin Oncol* **15**, 81-94 (2018). https://doi.org:10.1038/nrclinonc.2017.166

Schmiedel, J. M. *et al.* Gene expression. MicroRNA control of protein expression noise. *Science* **348**, 128-132 (2015). https://doi.org:10.1126/science.aaa1738

Tsang, J., Zhu, J. & van Oudenaarden, A. MicroRNA-mediated feedback and feedforward loops are recurrent network motifs in mammals. *Mol Cell* **26**, 753-767 (2007). https://doi.org:10.1016/j.molcel.2007.05.018

Wang, D. *et al.* Uncovering the cellular capacity for intensive and specific feedback self-control of the argonautes and MicroRNA targeting activity. *Nucleic Acids Res* **48**, 4681-4697 (2020). https://doi.org:10.1093/nar/gkaa209

Zhang, F. & Wang, D. The Pattern of microRNA Binding Site Distribution. *Genes (Basel)* **8** (2017). https://doi.org:10.3390/genes8110296

Becskei, A. & Serrano, L. Engineering stability in gene networks by autoregulation. *Nature* **405**, 590-593 (2000). https://doi.org:10.1038/35014651

Del Vecchio, D., Dy, A. J. & Qian, Y. Control theory meets synthetic biology. *J R Soc Interface* **13** (2016). https://doi.org:10.1098/rsif.2016.0380

Hasty, J., McMillen, D. & Collins, J. J. Engineered gene circuits. *Nature* **420**, 224-230 (2002). https://doi.org:10.1038/nature01257

Alon, U. Network motifs: theory and experimental approaches. *Nat Rev Genet* **8**, 450-461 (2007). https://doi.org:10.1038/nrg2102

Nise, N. S. *Control systems engineering*. 7th edn, (Wiley, 2014).

Heumos, L. *et al.* Best practices for single-cell analysis across modalities. *Nat Rev Genet* **24**, 550-572 (2023). https://doi.org:10.1038/s41576-023-00586-w

Jovic, D. *et al.* Single-cell RNA sequencing technologies and applications: A brief overview. *Clin Transl Med* **12**, e694 (2022). https://doi.org:10.1002/ctm2.694

Danielski, K. Guidance on Processing the 10x Genomics Single Cell Gene Expression Assay. *Methods Mol Biol* **2584**, 1-28 (2023). https://doi.org:10.1007/978-1-0716-2756-3_1

Sheng, K., Cao, W., Niu, Y., Deng, Q. & Zong, C. Effective detection of variation in single-cell transcriptomes using MATQ-seq. *Nat Methods* **14**, 267-270 (2017). https://doi.org:10.1038/nmeth.4145

Isakova, A., Neff, N. & Quake, S. R. Single-cell quantification of a broad RNA spectrum reveals unique noncoding patterns associated with cell types and states. *Proc Natl Acad Sci U S A* **118** (2021). https://doi.org:10.1073/pnas.2113568118

Salmen, F. *et al.* High-throughput total RNA sequencing in single cells using VASA-seq. *Nat Biotechnol* **40**, 1780-1793 (2022). https://doi.org:10.1038/s41587-022-01361-8

Niu, Y., Luo, J. & Zong, C. Single-cell total-RNA profiling unveils regulatory hubs of transcription factors. *Nat Commun* **15**, 5941 (2024). https://doi.org:10.1038/s41467-024-50291-3

Liu, W. & Shomron, N. Analysis of microRNA Regulation in Single Cells. *Methods Mol Biol* **2243**, 339-354 (2021). https://doi.org:10.1007/978-1-0716-1103-6_18

Liu, W. & Shomron, N. Analysis of MicroRNA Regulation and Gene Expression Variability in Single Cell Data. *J Pers Med* **12** (2022). https://doi.org:10.3390/jpm12101750

Wang, N. *et al.* Single-cell microRNA-mRNA co-sequencing reveals non-genetic heterogeneity and mechanisms of microRNA regulation. *Nat Commun* **10**, 95 (2019). https://doi.org:10.1038/s41467-018-07981-6

La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018). https://doi.org:10.1038/s41586-018-0414-6

Pandit, S., Wang, D. & Fu, X. D. Functional integration of transcriptional and RNA processing machineries. *Curr Opin Cell Biol* **20**, 260-265 (2008). https://doi.org:10.1016/j.ceb.2008.03.001

Rabani, M. *et al.* Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nat Biotechnol* **29**, 436-442 (2011). https://doi.org:10.1038/nbt.1861

Schwalb, B. *et al.* TT-seq maps the human transient transcriptome. *Science* **352**, 1225-1228 (2016). https://doi.org:10.1126/science.aad9841

Jiang, W. *et al.* A Multi-Parameter Analysis of Cellular Coordination of Major Transcriptome Regulation Mechanisms. *Sci Rep* **8**, 5742 (2018). https://doi.org:10.1038/s41598-018-24039-1

Bhattacharya, A. & Cui, Y. SomamiR 2.0: a database of cancer somatic mutations altering microRNA-ceRNA interactions. *Nucleic Acids Res* **44**, D1005-1010 (2016). https://doi.org:10.1093/nar/gkv1220

Hayes, J., Peruzzi, P. P. & Lawler, S. MicroRNAs in cancer: biomarkers, functions and therapy. *Trends in Molecular Medicine* **20**, 460-469 (2014). https://doi.org:10.1016/j.molmed.2014.06.005

Tian, S., Wang, J., Zhang, F. & Wang, D. Comparative Analysis of microRNA Binding Site Distribution and microRNA-Mediated Gene Expression Repression of Oncogenes and Tumor Suppressor Genes. *Genes (Basel)* **13** (2022). https://doi.org:10.3390/genes13030481

Burki, T. 2024 Nobel Prize awarded for work on microRNAs. *The Lancet* **404**, 1507-1508 (2024). https://doi.org:10.1016/S0140-6736(24)02303-1

Wang, T. *et al.* The Enrichment of miRNA-Targeted mRNAs in Translationally Less Active over More Active Polysomes. *Biology (Basel)* **12** (2023). https://doi.org:10.3390/biology12121536

Wang, D. Discrepancy between mRNA and protein abundance: insight from information retrieval process in computers. *Comput Biol Chem* **32**, 462-468 (2008). https://doi.org:10.1016/j.compbiolchem.2008.07.014

Wang, D. & Gribskov, M. Examining the architecture of cellular computing through a comparative study with a computer. *J R Soc Interface* **2**, 187-195 (2005). https://doi.org:10.1098/rsif.2005.0038

Lu, J. *et al.* MicroRNA expression profiles classify human cancers. *Nature* **435**, 834-838 (2005). https://doi.org:10.1038/nature03702

Gonzalez, I. A. *et al.* DICER1 tumor predisposition syndrome: an evolving story initiated with the pleuropulmonary blastoma. *Mod Pathol* **35**, 4-22 (2022). https://doi.org:10.1038/s41379-021-00905-8

Meister, G. Argonaute proteins: functional insights and emerging roles. *Nat Rev Genet* **14**, 447-459 (2013). https://doi.org:10.1038/nrg3462

Chen, C. Y., Zheng, D., Xia, Z. & Shyu, A. B. Ago-TNRC6 triggers microRNA-mediated decay by promoting two deadenylation steps. *Nat Struct Mol Biol* **16**, 1160-1166 (2009). https://doi.org:10.1038/nsmb.1709

Hansen, M. M. K. *et al.* A Post-Transcriptional Feedback Mechanism for Noise Suppression and Fate Stabilization. *Cell* **173**, 1609-1621 e1615 (2018). https://doi.org:10.1016/j.cell.2018.04.005

Stoiber, M. H. *et al.* Extensive cross-regulation of post-transcriptional regulatory networks in Drosophila. *Genome Res* **25**, 1692-1702 (2015). https://doi.org:10.1101/gr.182675.114

Eulalio, A., Behm-Ansmant, I. & Izaurralde, E. P bodies: at the crossroads of post-transcriptional pathways. *Nat Rev Mol Cell Biol* **8**, 9-22 (2007). https://doi.org:10.1038/nrm2080

Welte, T. *et al.* Convergence of multiple RNA-silencing pathways on GW182/TNRC6. *Mol Cell* **83**, 2478-2492 e2478 (2023). https://doi.org:10.1016/j.molcel.2023.06.001

Bose, M., Chatterjee, S., Chakrabarty, Y., Barman, B. & Bhattacharyya, S. N. Retrograde trafficking of Argonaute 2 acts as a rate-limiting step for de novo miRNP formation on endoplasmic reticulum-attached polysomes in mammalian cells. *Life Sci Alliance* **3** (2020). https://doi.org:10.26508/lsa.201800161