1 **Eukaryotic MAGs recovered from deep metagenomic sequencing of the seagrass,**
2 ***Zostera marina,* include a novel chytrid in the order Lobulomycetales**
3

4 Cassandra L. Ettinger[1,#], Jonathan A. Eisen[2,3,4], Jason E. Stajich[1,5]
5
[1]Department of Microbiology and Plant Pathology, University of California, Riverside, CA, United
States
[2]Genome Center, University of California, Davis, CA, United States
[3]Department of Evolution and Ecology, University of California, Davis, CA, United States
[4]Department of Medical Microbiology and Immunology, University of California, Davis, Davis,
CA, United States
[5]Institute for Integrative Genome Biology, University of California, Riverside, Riverside, CA,
United States
14
15
16 [#]**Corresponding Author:** Cassandra L. Ettinger, clettinger@ucdavis.edu

17
18

19 **Keywords:** *Zostera marina*, marine fungi, microbial eukaryotes, eelgrass, metagenomics,
20 genome assembly, genome annotation, Chytridiomycota, Bacillariophyta, Prymnesiophyceae,
21 Lobulomycetales

Abstract:

Fungi play pivotal roles in terrestrial ecosystems as decomposers, pathogens, and endophytes, yet their significance in marine environments is often understudied. Seagrasses, as globally distributed marine flowering plants, have critical ecological functions, but knowledge about their associated fungal communities remains relatively limited. Previous amplicon surveys of the fungal community associated with the seagrass, *Zostera marina* have revealed an abundance of potentially novel chytrids. In this study, we employed deep metagenomic sequencing to extract metagenome-assembled genomes (MAGs) from these chytrids and other microbial eukaryotes associated with *Z. marina* leaves. Our efforts resulted in the recovery of five eukaryotic MAGs, including a single fungal MAG in the order Loubulomycetales (65% BUSCO completeness), three MAGs representing diatoms in the family Bacillariaceae (93%, 70% and 31% BUSCO completeness) and a single MAG representing a haptophyte algae in the genus *Prymnesium* (40% BUSCO completeness). Whole-genome phylogenomic assessment of these MAGs suggests they all largely represent under sequenced, and possibly novel eukaryotic lineages. Of particular interest, the chytrid MAG was placed within the order Lobulomycetales, consistent with the identity of the dominant chytrid from previous *Z. marina* amplicon survey results. Annotation of this MAG yielded 5,650 gene models of which 77% shared homology to current databases. With-in these gene models, we predicted 121 carbohydrate-active enzymes and 393 secreted proteins (103 cytoplasmic effectors, 30 apoplastic effectors). Exploration of orthologs between the Lobulomycetales MAG and existing Chytridiomycota genomes have revealed a landscape of high-copy gene families related to host recognition and interaction. Further machine learning analyses based on carbohydrate-active enzyme composition predict that this MAG is a symbiont. Overall, these five eukaryotic MAGs represent substantial genomic novelty and valuable community resources, contributing to a deeper understanding of the roles of fungi and other microbial eukaryotes in the larger seagrass ecosystem.

Introduction:

Fungi are key contributors in terrestrial ecosystems, functioning as decomposers, pathogens, and endophytes, but their importance in marine environments remains underexplored [1–3]. While over 120,000 species of terrestrial fungi have been documented, only about 2,100 marine fungal species have been formally described, though the actual diversity is believed to be much higher [4–7]. Recent research has shed light on the distribution, activity, and biomass of marine fungi, with current estimates suggesting that fungi contribute 0.19-0.21 Gt C to the biomass of the open ocean [8–14]. However, host-associated fungal communities in marine ecosystems, including those associated with marine plants such as seagrasses, are still largely understudied.

Seagrasses are fully submerged flowering plants and vital foundation species in coastal ecosystems globally. *Zostera marina*, commonly known as eelgrass, is a widely distributed species that plays a crucial role in providing essential ecosystem services across coastal regions in the Northern Hemisphere [15–17]. Seagrasses like *Zostera marina* are home to a diverse range of microbial eukaryotes, including diatoms, dinoflagellates, algae, oomycetes, protists, and fungi [18–24]. Studies have shown that fungi represent only a small fraction of the total epiphytic eukaryotic community associated with *Z. marina*, with diatoms, oomycetes, and dinoflagellates among the dominant groups [25–27].

Although culture-based studies have explored fungal diversity associated with *Z. marina*, these approaches are limited by the challenges of growing marine fungi in laboratory settings [28–32]. Advances in DNA-based, culture-independent sequencing techniques have allowed for a more comprehensive characterization of seagrass-associated fungal communities, including their community assembly and biogeographical patterns [25,33–39]. Despite this progress, many seagrass-associated fungi, particularly those from Chytridiomycota lineages, remain difficult to taxonomically classify due to biases in reference databases, which are heavily skewed toward terrestrial fungi [25,36,38,39]. This is unsurprising as a limited number of lineages of marine

Chytridiomycota have been described through culture-based methods, and few have sequenced

genomes [6,7,40,41]. Nevertheless, marine Chytridiomycota are consistently recovered in DNA-

based surveys and are often dominant members of seagrass-associated fungal communities

[14,25,34,36,42].

Chytridiomycota in the order Lobulomycetales have previously been observed in high

abundance on and inside *Z. marina* leaves [36], and members of this group are part of the *Z.*

marina core mycobiome [25]. The Loubulomycetales are a relatively understudied group of

chytrid fungi that have been found to span terrestrial, freshwater and marine ecosystems as

both parasites and saprobes [43]. The two sequenced genomes from this group represent

Lobulomyces angularis, which was isolated from an acidic freshwater lake [44], and *Clydaea*

vesicula, which was isolated from soil underneath Eucalyptus trees [43]. During the initial

description of the Loubulomycetales, the authors noted that due to the difficulty in isolating

members of this group, even for 'chytrid specialists', that they expected most members would

need to be identified and described from molecular data [43].

While previous work identified high relative abundance of chytrid fungi associated with

seagrasses, particularly members of the Lobulomycetales, we still have little understanding of

the functional role of these enigmatic aquatic chytrids [25,36,45]. To address this, we used a

metagenomic sequencing approach on DNA that was previously extracted from *Zostera marina*

leaves in Ettinger and Eisen [36] and was reported to have a high relative abundance of

Lobulomycetales chytrid fungi. We then used this metagenomic sequencing data to: (i) generate

a set of refined eukaryotic metagenome-assembled genomes (MAGs), (ii) phylogenetically

place in the tree of life and functionally annotate these eukaryotic MAGs, and (iii) use

comparative genomics approaches to explore hypotheses of the possible function and ecology

of any fungal MAGs.

Methods:

Sequence generation

DNA was extracted from epiphytic washes from *Z. marina* leaves as part of previous work

focused on characterizing the mycobiome using high-throughput sequencing of the ITS2 region

[36]. Three DNA extracts from that work were selected based on predicted relative abundances

of an unclassified chytrid in the order Lobulomycetales (identified as amplicon sequence variant

SV8), with two samples (108A and 109A) having relatively high abundance of this putative taxa,

and one sample having a low abundance of this taxa (140A). These samples with variable

abundance of SV8 were chosen to aid binning algorithms which can utilize differential coverage

to aid in metagenomic bin recovery [46,47]. DNA was provided to the UC Davis Genome Center

DNA Technologies Core for library preparation and sequencing. DNA libraries were sequenced

on an Illumina HiSeq4000 to generate 150 bp paired-end reads.

Sequence processing

Sequence reads were trimmed using bbDuk v. 37.68 [48] with the following parameters: qtrim=rl

trimq=10 maq=10. We then mapped against and removed any reads from the metagenomes

matching the available genome for *Z. marina* v. 3.1 [49] using bowtie2 v. 2.4.5 [50] and

samtools v. 1.11 [51]. The remaining reads from all three metagenomic samples were co-

assembled using MEGAHIT v. 1.2.9 [52]. We ran Whokaryote v. 1.1.2 [53], which uses a

random forest classifier to predict the eukaryotic origin of contigs. Whokaryote was run with the

parameter --minsize 1 kbp and we chose to include predictions from Tiara v. 1.0.3 [53,54],

which uses deep-learning to further identify eukaryotic contigs.

The anvi'o v. 7.2 workflow [55] was used to identify and assess eukaryotic MAGs. This workflow

was executed twice: once for the entire co-assembly and once for the subset of the co-

assembly containing only the Whokaryote predicted eukaryotic contigs. Briefly, this workflow

involved obtaining read coverage for each metagenomic sample against the co-assembly using

bowtie2 v. 2.4.5 [50] and samtools v. 1.11 [51]. Next "anvi-gen-contigs-database" was used to

generate a contig database from the co-assembly and gene calls in the co-assembly were

predicted using Prodigal v. 2.6.3 [56]. Single-copy bacterial [57], archaeal [57], and protista [58]

genes were then predicted using HMMER v. 3.2.1 [59] and ribosomal RNA genes were

predicted using barrnap [60]. Each gene call was then assigned a putative taxonomy using Kaiju

v. 1.8.2 [61] with the NCBI BLAST non redundant protein database nr including fungi and

microbial eukaryotes v. 2020-05-25. For each metagenomic sample, an individual anvi'o profile

was constructed using "anvi-profile" for contigs >1 kbp with the "–cluster-contigs" option.

Several automatic binning algorithms were then employed to generate a preliminary set of

MAGs including MetaBAT2 v. 2.15, MaxBin v. 2.2.1, BinSanity v.0.5.4 and CONCOCT v. 1.1.0

[62–65]. All predicted MAG bins were then assessed with eukCC v. 2.0 [66] for eukaryotic

completeness and redundancy.

To obtain a final representative set of eukaryotic MAGs, dereplication was performed using

dRep v. 3.2.2 [67] by providing the eukCC completeness estimates. This step involved

combining bins identified from both anvi'o workflow runs (the results from the entire co-

assembly and the subset of eukaryotic contigs). The minimum completeness threshold for dRep

was set to 1 to ensure a comprehensive representation of eukaryotic MAGs, which often have

lower completeness compared to bacterial MAGs.

Representative MAGs with >30% eukaryotic completeness were assessed using anvi-refine. A

completeness threshold of >30% has been previously used as a cut-off for high-quality

eukaryotic draft MAGs [68]. Further assessment and decontamination of these eukaryotic MAGs

was performed using the BlobTools2 v. 4.1.4 [69] workflow. This workflow involves assigning

taxonomy based on searches against the UniProt Reference Proteomes database (v. 2021_04)

to each contig using diamond (v. 2.1.0) and against the nt database (downloaded 2023-03-21)

using command-line BLAST v. 2.2.30+ [70,71]. Next, coverage of each individual metagenomic

sample against each eukaryotic bin was calculated with Minimap2 v. 2.24 [72] and sorted using

samtools v. 1.11 [51]. Finally the BlobToolKit Viewer was used to visualize the GC content, read

coverage and predicted taxonomies of contigs to identify contaminants. To be conservative, any

contigs that were taxonomically identified as viral, bacterial, or archaeal were removed from the

eukaryotic bins.

After removing potential contamination, we ran EUKulele v. 1.0.2 [73] to assess and refine the

initial taxonomic assignments for each bin. We also used BUSCO v. 5.3.2 [74] to assess

completion of the de-contaminated bins in 'genome' mode using the eukaryotic_odb10 set, and

as appropriate the fungi_odb10 or stramenopiles_odb10 sets.

189
190   **Annotation**
191
Draft MAG assemblies were screened for contaminant vectors using the 'vecscreen' option in

the Automatic Assembly of the Fungi (AAFTF) pipeline v. 0.4.1 [75]. Repetitive regions were *de*

novo identified and soft masked prior to MAG annotation using RepeatModeler v. 2.0.1 [76] and

RepeatMasker v. 4-1-1 [77].

Draft eukaryotic MAGs were annotated using the Funannotate pipeline v. 1.8.14 [78].

Funannotate incorporates multiple software algorithms for eukaryotic gene prediction including

Augustus v. 3.4.0, GeneMark-ETS v. 4.71, GlimmerHMM v. 3.0.4, and SNAP v. 2013_11_29

[79–83], and then generates consensus gene models using EVidenceModeler v. 1.1.1 [84]. For

two MAGs (SGEUK-04 and SGEUK-05) where GeneMark-ETS was unable to run due to

assembly fragmentation, we used –weights genemark:0 during prediction to skip use of this

software. Funannotate also uses tRNAscan v. 2.0.11 to predict tRNAs [85]. Annotation of

consensus gene models by Funannotate is based on searches against Pfam v. 35.0 [86] and

dbCAN v. 9.0 [87,88] using HMMER v.3 [59] and against MEROPS v. 12.0 [89], eggNOG v.

2.1.9 [90], InterProScan v. 5.51-85.0 [91], and UniProt v. 2022_05 [92] using diamond v. 2.0.15

[93]. Transmembrane proteins are further predicted using Phobius v. 1.01 [94] and secreted

proteins using SignalP v. 5.0b [95]. For the fungal MAG (SGEUK-03), AntiSMASH v. 6.1.1 was

run to predict biosynthetic gene clusters [96] and EffectorP v. 3.0 was run on putative secreted

proteins to predict apoplastic and cytoplasmic effectors [97]. We also used CATAStrophy v.

0.1.0 [98], a classification method based on carbohydrate-active enzyme (CAZyme) patterns

from filamentous fungal plant pathogens, to predict the possible trophic strategy of SGEUK-03.

CATAStrophy was run in a Google Collab implementation using dbCAN v. 10 [87,88].

BUSCO v. 5.3.2 [74] was then used to assess completion of MAG annotations in 'protein' mode

using the eukaryotic_odb10 set, and as appropriate the fungi_odb10 or stramenopiles_odb10

sets.

218
219 **Phylogenetic assessment**
220
To examine the phylogenetic placement of the five eukaryotic MAGs, we carried out analysis of

three groups: stramenopiles (SGEUK-01, SGEUK-02, SGEUK-04), fungi (SGEUK-03), and

haptista (SGEUK-05). For each group, we obtained gene models from annotated genomes that

were publicly available in NCBI GenBank (Tables S1–S3). Given the limited availability of

annotated genomes for haptista (four genomes at the time of this work), we also incorporated

publicly available transcriptomic data from METdb [99–101].

For stramenopiles, the final dataset for phylogenomic analyses included SGEUK-01, SGEUK-

02, SGEUK-04, and 153 annotated genomes representing the following groups: Oomycota

(n=120), Bacillariophyta (n=11), Bigyra (n=8), Pelagophyceae (n=7), Eustigmatophyceae (n=3),

Phaeophyceae (n=2), Synurophyceae (n=1), and Xanthophyceae (n=1) (Table S1) [102–164].

For fungi, the final dataset for comparative genomics and phylogenomic placement included

SGEUK-03 and 98 annotated genomes from the following phyla: Chytridiomycota (n=71),

Blastocladiomycota (n=10), Neocallimastigomycota (n=8), Ascomycota (n=4),

Monoblepharidomycota (n=3), Olpidiomycota (n=1), and Basidiomycota (n=1) (Table S2) [165–

186]. For haptista, the final dataset for phylogenomic placement included SGEUK-05, 47

annotated genomes and transcriptomes from members of the orders Phaeocystales (n=14),

Isochrysidales (n=14), Prymnesiales (n=9), Pavlovales (n=7), Coccolithales (n=2),

Coccosphaerales (n=1), and Haptophyta *incertae sedis* (n=1) (Table S3) [101,162,187–189].

To place MAGs in a phylogenetic context, the PHYling_unified

(https://github.com/stajichlab/PHYling_unified) pipeline was used to generate a protein

alignment of each MAG to close relatives based on preliminary taxonomic assignment (i.e., the

datasets described in Table S1-S3). This pipeline utilizes HMMER v.3 [59] and ClipKIT [190] to

search for, build, and trim an alignment based on gene set. Here we used the most relative

BUSCO gene set for each dataset, e.g. the BUSCO fungi_odb10 for SGEUK-3,

eukaryotic_odb10 for SGEUK-05, and stramenopiles_odb10 for SGEUK-01, SGEUK-02, and

SGEUK-04. Maximum likelihood phylogenies were built from these alignments using IQ-TREE2

v.2.2.6 [191], with the -p option to indicate gene partitions [192] and the -m MFP option to run

ModelFinder Plus to identify and use the optimal evolutionary model for each partition [193]. The

resulting phylogenetic trees were imported into R v. 4.3.0 [194] and visualized using ggtree v.

3.8.2 [195].

Given that 18S rRNA sequences often do not end up in MAG bins [196] and to further refine

taxonomy for the fungal MAG (SGEUK-03), we used phyloflash v. 3.4 [197] to assemble and

identify the 18S rRNA gene sequences in the metagenomes. We then identified closely related

sequences to the 18S rRNA gene sequence for the fungal MAG using NCBI's Standard

Nucleotide BLAST with default settings [71]. These results were then used to guide a literature

search to identify additional sequences for inclusion during phylogenetic reconstruction (Table

S4) [43,198–209]. These 18S rRNA gene sequences were then aligned using SSU-ALIGN v.

0.1.1 [210] and trimmed using trimAl v. 1.4 with the -gappyout method [211]. A maximum

likelihood phylogeny was built from this alignment using IQ-TREE2 v.2.2.6 [191], with the -p

option to indicate gene partitions [192] and the -m MFP option to run ModelFinder Plus to

identify and use the optimal evolutionary model for each partition [193]. The resulting

phylogenetic tree was imported into R and visualized using ggtree v. 3.8.2 [195].

264
265   **Comparative genomics of chytrid MAG**
266
OrthoFinder v. 2.5.4 [212] was used to predict phylogenetic hierarchical orthogroups (HOGs), or

gene families, across Chytridiomycota and related lineages (Table S1). Given the little genomic

information available for close relatives of SGEUK-03 we focused our analyses on HOGs at the

root node (i.e., all HOGs). Due to the incomplete nature of MAGs, we chose to focus

comparative analyses on high-copy HOGs, and did not explore gene loss. High-copy HOGs (≥5

copies in SGEUK-03) were visualized in R using the pheatmap v. 1.0.12 package [213].

273
274   **Results**
275
276   **Successful recovery and preliminary annotation of five eukaryotic MAGs**
277
In total, we recovered five eukaryotic MAGs that were high-quality (>30% complete). Briefly, we

note that bins resulting from the co-assembly containing only Whokaryote identified eukaryotic

contigs had lower contamination, though at the expense of overall completion, compared to their

entire co-assembly counterparts. Among the five dereplicated MAGs, only one MAG (SGEUK-

05) was not obtained from the eukaryotic co-assembly.

To perform preliminary taxonomic assignment of MAGs, we used EUKulele which reports the

relative proportion of all the proteins within a MAG that have annotations that agree at a

particular taxonomic level. Using a conservative cut-off of >70% proportional agreement,

EUKulele predicted that we had three Bacillariophyta MAGs (SGEUK-01: 89.60%, SGEUK-02:

78.39%, SGEUK-04: 75.08%), one Eukaryota MAG (SGEUK-03: 94.24%), and one

Prymnesiophyceae MAG (SGEUK-05: 82.24%).

These eukaryotic draft MAGs were fairly fragmented (i.e., with total contigs ranging from 1,074

to 11,398) likely due to their relatively low mean coverage (i.e., ranging from 3.47x to 7.71x).

Briefly, we discuss the general characteristics of the assembly and annotation of each draft

MAG (Table 1):

●   SG-EUK01 (Bacillariophyta) spans 32.54 Mbp with 3.46x coverage, distributed across

2,172 contigs with an N50 of 20,103 bp. Initial EukCC completion and contamination

estimates were 84.5% and 5.7%, respectively. After removing 21 contigs flagged as

bacterial contamination by BlobTools, BUSCO analysis using the stramenopiles_odb10

dataset indicated a high completeness of 93.0%. Annotation with Funannotate resulted

in 13,603 gene models, including 13,520 mRNA genes and 83 tRNA genes, with 75.92%

of gene models having EggNog database hits. BUSCO protein estimates were

consistent with the genome completeness, showing 91.0% of the expected single-copy

orthologs present using the stramenopiles_odb10 set.

●   SG-EUK02 (Bacillariophyta) is 28.49 Mbp in length with 5.37x coverage, spanning 5,479

contigs and an N50 of 5,715 bp. EukCC estimated 64.39% completeness and 13.96%

contamination, lower in completeness but higher in contamination than SG-EUK01. We

subsequently removed 92 contigs identified as bacterial contamination using BlobTools.

BUSCO estimates show 70.0% completeness using the stramenopiles_odb10 dataset.

Annotation resulted in 10,889 gene models, including 10,868 mRNA genes and 21 tRNA

genes, with 76.43% of gene models having EggNog database hits. BUSCO protein

estimates are aligned with genome completeness, showing 69.0% of single-copy

orthologs present using the stramenopiles_odb10 set.

● SG-EUK03 (Eukaryota) is a smaller MAG at 12.11 Mbp with 7.71x coverage, distributed

across 1,074 contigs with an N50 of 15,102 bp. EukCC predicted completion and

contamination at 56.96% and 1.74%, respectively. After removing 10 bacterial contigs

identified by BlobTools, BUSCO analysis revealed moderate completeness, with 65.1%

using the eukaryota_odb10 dataset and 51.6% using the fungi_odb10 dataset.

Annotation resulted in 5,650 gene models, including 5,630 mRNA genes and 20 tRNA

genes, with 76.58% of gene models having EggNog database annotation hits. Notably,

393 genes (6.96%) were predicted to be secreted, including 133 potential effectors (103

cytoplasmic, 30 apoplastic), and 121 genes (2.14%) had CAZyme domains including 65

glycosyltransferases (GTs), 26 glycoside hydrolases (GHs), 13 carbohydrate esterases

(CEs), 8 polysaccharide lyases (PLs), 7 auxiliary activities (AAs), and 2 carbohydrate-

binding modules (CBMs) (Figure S1, Table S5). Based on CAZyme content,

CATAStrophy predicted that SGEUK-03 was most likely a symbiont. BUSCO protein

estimates indicated 65.9% completeness with eukaryota_odb10 and 57.0% with

fungi_odb10, outperforming the genome completeness estimates.

● SG-EUK04 (Bacillariophyta) spans 19.6 Mbp with 4.04x coverage, spread over 11,398

contigs with an N50 of 1,790 bp. EukCC estimated a relatively low completeness of

35.9% and contamination of 5.98%, lower in completeness compared to SG-EUK01 and

SG-EUK02. We removed 197 contigs identified as potential bacterial contaminants.

BUSCO estimates using the stramenopiles_odb10 dataset indicated that only 31.0% of

single-copy orthologs were present and complete. Annotation resulted in 8,359 gene

models, including 8,343 mRNA genes and 16 tRNA genes, with 64.77% of gene models

having EggNog database hits. BUSCO protein estimates are aligned with the genome

completeness, with 30.0% of expected single-copy orthologs present using the

stramenopiles_odb10 set.

● SG-EUK05 (Prymnesiophyceae) is the largest of the MAGs, spanning 59.4 Mbp with

5.12x coverage, distributed across 9,735 contigs with an N50 of 6,756 bp. Initial EukCC

estimates placed completion at 34.0% and contamination at 3.4%. Following the removal

of 197 bacterial contigs, BUSCO estimates showed that 40.0% of expected orthologs

were present using the eukaryota_odb10 dataset. Annotation resulted in 15,098 gene

models, including 15,075 mRNA genes and 23 tRNA genes, with 56.78% of gene

models having EggNog database annotation hits. Repeats represented 18.63% of the

genome length, much higher than for any of the other draft MAGs. BUSCO protein

estimates performed slightly worse than the genome completeness with only 34.5% of

the expected single-copy orthologs present and complete.

Overall, despite fragmentation, these five eukaryotic MAGs exhibited varying levels of genome

and annotation completeness. The Bacillariophyta MAGs, particularly SG-EUK01 and SG-

EUK02, displayed higher genome completeness, reflecting robust reconstructions of their

genomic landscapes. SG-EUK03 showed a more moderate genome completeness, but a more

robust annotation. SG-EUK04 and SG-EUK05 are more fragmented, though still represent

valuable insight into the genomes and functions of these microbial eukaryotes.

Phylogenetic placement of eukaryotic MAGs puts them in the diatom, fungal, and
haptophyte trees of life
358
Following MAG recovery and functional annotation, we constructed phylogenetic trees to refine

the taxonomic placement of the five draft MAGs. Using BUSCO protein sets and guided by

preliminary taxonomic assignments from EUKulele, we built three trees: one for Bacillariophyta,

one for fungi, and one for Haptista.

SGEUK-01, SGEUK-02, and SGEUK-04 were placed into the Bacillariophyta tree, and clustered

among diatom lineages in the Bacillariaceae family, with SG-EUK02 and SG-EUK04 forming a

clade that is sister to *Nitzschia inconspicua*, while SG-EUK01 formed a novel branch serving as

outgroup to the clade representing all members of the Bacillariaceae family (Figure 1). While

SGEUK-01 had very similar BUSCO completion estimates compared to annotated close

relatives, both SGEUK-02 and SGEUK-04 likely paint a more incomplete picture of the

functional landscape of these organisms. All three MAGs have genomes and annotations on the

smaller side with publicly available genomes for members of the Bacillariaceae family ranging

from 27.23 Mbp to 99.71 Mbp in size [123,214], and with available annotations ranging in gene

count from 11,972 to 39,085 genes [123,139].

SGEUK-03 was placed into the fungal tree, falling amongst Chytridiomycota lineages (Figure 2).

The MAG was placed sister to a clade containing *L. angularis* and *C. vesicula*, which both

belong to the order Lobulomycetales, an undersequenced group of chytrids. *Quaeritorhiza*

haematococci, a parasitic chytrid of algae [215], serves as an outgroup to the Lobulomycetales

clade containing these three taxa. It falls in the same taxonomic class as Lobulomycetales, the

Chytridiomycetes, but its phylogenetic placement in an order is uncertain. Regardless, it has

been shown to be sister to the Lobulomycetales in other phylogenetic studies [216]. The two

available Lobulomycetales genomes are similarly fragmented to SGEUK-03, with *L. angularis*

represented by 1253 scaffolds with an N50 of 26,617 bp and *C. vesicula* spanning 2291

scaffolds with an N50 of 14,665 bp [172]. However, both their genomes (24.65-24.94 Mbp) and

annotations (8737-9208 genes) are larger than SGEUK-03 which is in line with its lower BUSCO

completion scores (Figure 2).

To further confirm and improve placement of SGEUK-03 among the Lobulumycetales, we used

phyloFlash to recover an 18S rRNA gene from the metagenomic dataset likely belonging to this

MAG and then placed in in the context of available 18S rRNA genes representing this chytrid

lineage. This 18S rRNA gene phylogeny confirmed placement of SGEUK-03 among the

Lobulumycetales, and further placed it within the novel SW-I clade, an aquatic clade of

Lobuluymycetales largely described from DNA survey datasets (Figure 3). SGEUK-03 is sister

to a sequence from deep sea marine sediment [199] and is in a clade with other sequences

from marine surveys [198]. This taxonomic placement matches that previously obtained for the

dominant amplicon sequence variant (SV8) from ITS2 amplicon profiling of the same samples,

supporting that SGEUK-03 represents the chytrid fungus previously identified to be abundant on

Z. marina leaves [36].

SGEUK-05 was placed into the haptophyta tree, among the *Prymnesiales* order and sister to

the transcriptome of the phytoplankton, *Prymnesium parvum* (Figure 4). SGEUK-05 and *P.*

parvum were sister to a clade of transcriptomes from *Chrysochromulina polylepis* which are also

members of the *Prymnesiaceae* family. *Chrysochromulina* genomes are 59.07-65.75 Mbp [189],

though other members of the *Prymnesiales* have genomes up to ~167.68 Mbp [187]. Given the

relatively low completion of SGEUK-05, despite its larger size of 59.4 Mbp, it is likely we are

only capturing less than half of its genomic and functional space.

407
408   **Comparative genomics of SGEUK-03 reveals genes involved in host interaction**
409
We performed OrthoFinder analysis, comparing SGEUK-03 with other fungi with a focus on

comparisons to other members of the Lobulomycetales and Chytridiomycota. In total, we

identified 75,458 phylogenetically hierarchical orthogroups (HOGs) among all fungi, of which

44,619 were identified among all Chytridiomycota. Of those, 8192 were identified among the

Lobulomycetales, with a total of 4238 HOGs identified in SGEUK-03. Of the 5630 genes in

SGEUK-03, 4886 genes (86.8%) were assigned to HOGs. Given that SGEUK-03 represents an

incomplete MAG, we focused our analyses on high-copy number orthogroups (≥5 members) in

SGEUK-03 and compared these to the average copy numbers in other fungal groups (Figure 5).

In total there were 20 high-copy HOGs, of which five were unique to SGEUK-03. These included

three hypothetical proteins (of which two were predicted cytoplasmic effectors), a Hect-type E3

ubiquitin ligase and a Sel-1 like / Major Facilitator Superfamily (MFS) protein. Six HOGs were

shared exclusively with other Chytridiomycota or Lobulomycetales, including genes potentially

involved in host interactions, such as leucine rich repeat receptor like kinases (LRR-RLKs), a

galactose-binding like domain encoding gene, a tyrosinase, and a secreted chitin deacetylase

(NodB homology-domain containing protein). Additionally, three HOGs were shared with other

early-diverging fungal lineages, including a transcription factor, a GT2, and an LRR effector.

Finally, six HOGs were shared across distant fungi, including Dikarya, such as GH17,

transcription factors, and a glutaminase.

429
430 **Discussion**
431
432 **Draft MAGs represent key 'omics resources of under sequenced eukaryotic lineages**
433
We report the successful recovery of five eukaryotic MAGs associated with the leaves of the

seagrass *Z. marina*. The five draft MAGs represent under sequenced eukaryotic lineages

including three MAGs representing diatoms in the family Bacillariaceae (93%, 70% and 31%

BUSCO completeness), one MAG representing a haptophyte algae in the genus *Prymnesium*

(40% BUSCO completeness), and a single fungal MAG in the order Loubulomycetales (65%

BUSCO completeness). We annotated these five draft eukaryotic MAGs, providing an

invaluable functional resource to enable foundational insight into their roles in the marine

ecosystem. While all five MAGs represent relatively novel 'omics resources, we highlight

SGEUK-03 which we could only taxonomically place at the order level as part of the

Loubulomycetales, a group with only two sequenced genomes, of which both are distant

relatives of this MAG. Overall all five eukaryotic MAGs represent significant contributions of

genomic novelty and we expect comparative genomic and functional approaches by the

research community to further elucidate their evolution and ecology.

447
448   **Diatoms and Phytoplankton are critical, yet understudied, primary producers in seagrass**
449   **ecosystems**
450
Seagrass leaves serve as an ideal habitat and stable refuge for marine diatoms and algae. In

fact, diatoms, including members in the Bacillariaceae, make up the biggest component of the

microscopic epiphytic biomass associated with seagrass leaves [19,217–219], have the largest

relative abundance within the microbial eukaryotic community associated with *Z. marina* globally

[25] and are considered part of *Z. marina*'s core epiphytic leaf community [26,27]. Similarly,

phytoplankton and other microalgae are another important part of the epiphytic community of

seagrasses [18,24,220]. Both diatoms and microalgae make significant contributions to primary

productivity, are seasonally variable and serve as important food sources in seagrass bed

ecosystems [19,20,218,221–224]. In this work, seagrass leaves were free of a visible epiphytic

biomass layer, suggesting that the recovery of these MAGs might indicate a rather tight

association of these taxa with *Z. marina* leaf tissues. We hope these draft MAGs and their

annotations provide useful resources for the community to begin to interrogate their genomics,

possible functional roles, community dynamics in seagrass beds and the resilience of these

microbial eukaryotes in the face of global climate change.

465
466   **Loubulomycetales may have a biotrophic or symbiotic role in seagrass ecosystems**
467
SGEUK-03 clustered within Novel Clade SW-I in the Lobulumycetales (Figure 3). Known

isolated members of Novel Clade SW-I are chytrids that are parasites of freshwater algae and

diatoms [203,204,225]. While additional unclassified members of Novel Clade SW-I are only

known from amplicon-based marine surveys of deep sea sediments, open ocean and seagrass

ecosystems [36,198,199]. Overall, the Loubulomycetales are a relatively understudied group of

chytrid fungi that have been found to span terrestrial, freshwater and marine ecosystems as

both parasites and saprobes [43]. Given this diversity, and the absence of isolated

representatives from seagrass environments, the ecological role of SGEUK-03 remains unclear.

However, insights from its genome, specifically exploring effector and CAZyme profiles (Figure

S1, Table S5), provide clues to its possible functional role.

Effectors are critical for all plant-fungal interactions [226–229] including for symbiotic and

biotrophic fungi which still need to evade host plant recognition and suppress host defenses to

persist [230,231]. Additionally, parasitic chytrids of animals and plants have been observed to

have a repertoire of effectors proteins, as well as pathogenicity factors [166,170,232–234].

SGEUK-03 has a larger number of cytoplasmic effectors than apoplastic effectors, a pattern that

might be consistent with an intracellular role. The majority of predicted effectors were

hypothetical proteins with limited functional annotation. While the precise function of these

effectors remains unclear, their abundance in SGEUK-03 points to a host-associated role.

We can explore the CAZyme diversity of SGEUK-03 to further hone in on its functional role.

CAZyme-encoding genes vary among fungal species and likely reflect their adaptation to

different nutritional niches [235]. For example, plant pathogens have been found to have higher

numbers of CAZymes than saprophytic and animal pathogens [236]. Plant cell wall degrading

enzymes, such as GHs, CEs and PLs, in particular, are often employed by fungal plant

pathogens [237]. Necrotrophic pathogens deploy these cell wall-degrading enzymes to promote

host damage and colonization [238]. Whereas endophytic fungi and biotrophs tend to have

fewer of these cell wall degrading enzymes [238–241]. While CAZyme diversity has been widely

explored on land, particularly in the Dikaryan fungi, little characterization of these enzymes has

been done in chytrids, though initial reports suggest these fungi can have a wide range of

CAZymes [242].

SGEUK-03 had relatively few CAZymes overall, with a notable absence of enzymes reported to

be involved in the degradation of plant cell wall components such as cellulose (GH6, GH7,

GH12, GH45, GH3, GH61, CBM1), hemicellulose (GH10, GH11, GH74, GH27, GH36, GH26,

GH43, GH51, GH54, GH62, GH35, GH67, GH115), and pectin (GH28, GH78, PL1, PL3, PL4,

PL9) [236,239,243,244]. Instead, the glycosyl transferases (GTs) were the most represented

group (53.7% of CAZymes), with GT2 being the most frequent, a pattern perhaps consistent

with fungi that modify their own cell wall to evade detection, rather than degrade host cell walls.

Interestingly, the CAZyme profile of SGEUK-03 mirrors that of the obligate mutualistic symbiont

Rhizophagus irregularis, an arbuscular mycorrhizal fungus that engages in a mutualistic

relationship with land plants [245]. The relatively low CAZyme diversity of SGEUK-03, combined

with its effector content, suggest that it may engage in a similar strategy of minimal host

damage while exploiting host resources. This idea is further supported by the prediction from

CATAStrophy [98], which classified SGEUK-03 as a symbiont based on its CAZyme content.

However, it is important to note that CATAStrophy was trained on filamentous dikaryon fungi,

and its application to chytrids may be limited. Additionally, as the genome of SGEUK-03 is

incomplete, we cannot rule out the presence of additional CAZymes that could alter this

classification. Regardless, on the whole, the effector and CAZyme profiles of SGEUK-03 may

suggest that it has a symbiotic or biotrophic role in seagrass ecosystems.

518
519   **Are Lobulomycetales chytrids biotrophic parasites or symbionts of seagrasses**
520   **themselves?**
521

The frequent identification of *Lobulomycetales* species in *Z. marina* leaf epiphyte and endophyte

samples from previous amplicon surveys [25,36] supports the hypothesis that SGEUK-03 may

be closely associated with *Z. marina*. Microscopic observations of chytrids directly interacting

with seagrass tissues further support a possible symbiotic or weakly parasitic role [246].

However, given the diversity of hosts that chytrids can associate with, we cannot rule out the

possibility that SGEUK-03 may also be parasitizing a diatom or algal host that is itself

associated with *Z. marina*. While we saw no clear coverage pattern (e.g., 1-to-1) which might

suggest one of the other MAGs might be a host for SGEUK-03, another possibility could be that

SGEUK-03 is a generalist chytrid, capable of associating with multiple hosts including *Z. marina*

and members of its epiphytic community.

Chytrids are well known parasites of a broad range of phytoplankton, including diatoms, green

algae, dinoflagellates, and cyanobacteria and can display both high host specificity or broad

generalist strategies, depending on environmental conditions and host availability

[203,204,247–255]. A generalist strategy allows for infection of multiple species enabling

chytrids to persist despite fluctuations in available host communities, with trade-offs including

lower growth rates as a result of needing to maintain the genomic potential to evade diverse

host immune responses [251]. Members of the Lobulomycetales and marine fungi more

generally have been observed to exhibit some seasonality in their emergence which has been

hypothesized to relate to host-dynamics and environmental conditions [44,256–258]. Blooms of

parasitic chytrids, for example, frequently accompany an increased abundance of diatoms that

the chytrids infect and ultimately kill. Previous work has suggested that Lobulomycetales sp.

associated with seagrasses experience bloom-like dynamics, which could relate to seasonality,

environmental conditions or host availability [25,36]. It is possible that the presence of SGEUK-

03 in seagrass ecosystems is similarly influenced by seasonal patterns in host availability,

whether that host is seagrass itself or associated epiphytic algae or diatoms. Further work is

needed to confirm and explore the host-specificity of SGEUK-03.

SGEUK-03's high copy orthologs may represent a functional repertoire focused on host
association

To better understand its functional potential, particularly its interactions with its host, we used

comparative genomics approaches to examine SGEUK-03's highest copy orthologs. We found

twenty high-copy gene families with functions that indicate they may be involved in host or

environmental recognition, evasion of host immune responses, and nutrient acquisition.

Among the high-copy orthogroups, we identified two LRR-LRKs, which are receptor kinases that

detect external signals and are often implicated in non-self recognition. For example, LRR-LRKs

are best known for the role they play in plant and animal innate immunity [259,260]. Dramatic

gene family expansions of LRR-LRKs are observed in plants and animals, but these genes are

largely missing in dikaryon fungi, whose innate immune system is still not fully understood,

though a small number of LRR-LRKs have been reported in chytrids and zygomycetes [261–

263]. The HOGs that included LRR-LRKs were shared with other chytrids and were the most

abundant HOGs identified for SGEUK-03. Aquatic chytrids have motile zoospores which enable

the targeting of specific hosts and substrates, a capability that is absent in dikaryan spores

which lack a flagella [45]. For instance, chytrid zoospores can be attracted to dissolved

molecules in the environment which may facilitate host detection [264–266]. We speculate that

these LRR-LRKs may be involved in environmental sensing and host detection in SGEUK-03,

although a role in innate immunity, similar to that of plants and animals, could also be possible.

A galactose-binding domain protein, another high-copy gene family shared with chytrids, may

support early host recognition and adhesion. Galactose is an important sugar in seagrass cell

walls [267]. Additionally, a galactose-binding CBM plays an important role in host-fungal

interactions through mediating chemotaxis towards the host, attachment, and virulence in a

chytrid pathogen [268]. SGEUK-03 may be similarly using this gene family for sensing and

adhering to host tissues during colonization, establishing an interaction with the host early in the

infection process.

In addition to host recognition, SGEUK-03 may possess mechanisms to evade host immune

responses during and after invasion. Fungi have evolved several strategies to evade host

immune responses including modifying their cell wall composition and secreting effector

proteins to protect cell walls or help evade host detection [269,270]. SGEUK-03 possesses a

suite of effectors, including two unique secreted cytoplasmic effector families. In addition,

SGEUK-03 encodes a unique family of Hect-type E3 ubiquitin ligases. These ligases play critical

roles in regulating protein degradation through modification, which impacts cellular processes

such as stress responses, endocytosis, cell cycling, and pathogenicity [271–275] Such effectors

and ubiquitin ligases may provide SGEUK-03 with the ability to manipulate host cell machinery

and suppress immune responses.

Additionally, SGEUK-03 shares a gene family annotated as a chitin deacetylase (NodB

homology-domain containing protein) with other chytrids. Chitin deacetylases can modify the

fungal cell wall during plant invasion making it less recognizable to plant defense systems

[276,277]. Chitin deacetylases with NodB homology are also involved in invasion and signaling

by symbiotic fungi [278]. SGEUK-03 may similarly modify its cell wall to evade host immunity or

maintain host interactions.

SGEUK-03 has a glycoside hydrolase family 17 (GH17) enzyme gene family, which it shared

with distant fungal lineages, including members of Dikarya. GH17 enzymes are involved in the

modification of cell wall glucan in both fungal and plant cells, which is essential for the growth,

development and host-pathogen interactions [279–281]. On land, *Cladosporium fulvum*, a

hemibiotrophic fungal pathogen of tomatoes, secretes a GH17 which triggers release of a

damage-associated molecular pattern from the host cell wall triggers host innate immunity and

ultimately cell death [282]. Further, GH17 enzymes in fungi are abundant and thought to have

important roles in global carbon cycling in the open ocean [283]. For example, a marine

Cladosporium species secretes GH17 when utilizing laminarin [284], which is an algal and

diatom-derived β-glucan that's abundant in the ocean [285]. It is possible that GH17 enzymes in

SGEUK-03 could play a similar role in interacting with host cell walls, either during invasion or

as part of a defensive strategy.

Tyrosinases, a high-copy gene family of SGEUK-03 shared with other chytrid lineages, have

been previously reported as expanded and implicated in host-pathogen interactions [286,287].

These enzymes are involved in melanin biosynthesis, which contribute to fungal virulence by

protecting cells from UV radiation, reactive oxygen species (ROS), and hydrolytic enzymes that

degrade fungal cell walls [288–290]. Melanin production could provide SGEUK-03 with

protection from host defenses, as ROS accumulation is a common plant response to pathogen

recognition [238,291]. Therefore, the putative tyrosinases in SGEUK-03 might provide protection

from ROS-mediated host defenses, ensuring its survival within host tissues.

We also found that gene families with functions likely related to nutrient acquisition,

detoxification and stress responses including several transcription factors and transporter

genes. This included a unique Sel-1 like / MFS transporter family, which are a group of

transporters vital in cross-membrane transport of organic solutes. They are key in fungal

nutrient acquisition and may play roles in detoxification or efflux of harmful plant compounds

[292]. Overall, the observed suite of high-copy orthologs in SGEUK-03 supports a functional

landscape of gene families with putative annotations related to host or environmental

recognition, evasion, and interaction.

628
629 **Conclusion**

630
We successfully recovered five eukaryotic MAGs from DNA isolated from the leaves of the

seagrass *Zostera marina*, representing previously under-sequenced eukaryotic lineages. These

include three MAGs of diatoms from the family Bacillariaceae, one MAG from a haptophyte

algae in the genus *Prymnesium*, and one fungal MAG from the order Loubulomycetales. The

functional annotation of these draft MAGs offers a valuable resource for understanding their

ecological roles in the marine environment. Notably, we placed SGEUK-03 within the novel

clade SW-I of the Loubulomycetales and performed comparative genomic analysis, suggesting

a symbiotic or biotrophic lifestyle based on its effector and CAZyme repertoire. Further

exploration of high-copy orthogroups in SGEUK-03 revealed a suite of gene families potentially

involved in host or environmental interactions, recognition, and immune evasion. While

additional research is required to clarify host specificity and further characterize these MAGs,

our findings contribute to the growing knowledge of marine microbial eukaryotic diversity and

provide a genomic foundation for future studies on the ecological roles and evolutionary

trajectories of these enigmatic microorganisms.

645
646 **DNA Deposition**

647
The raw metagenomic sequencing data was deposited at GenBank under accession no.

PRJNA1140276. MAGs and annotations under this BioProject at DDBJ/ENA/GenBank under

accession no. JBKBCY000000000-JBKBDC000000000. The 18S rRNA gene obtained for

SGEUK-03 was deposited at GenBank under accession no. PQ146498. All code used in this

work has been deposited on Github ([casett/ZM_Euk_MAGs](#)) and archived in Zenodo (DOI:

[10.5281/zenodo.14278507](#)).

660 **Funding sources**

661
The sequencing data in this work was generated by grants from the UC Davis H. A. Lewin

Family Fellowship and the UC Davis Center for Population Biology to CLE. CLE was supported

by the National Science Foundation (NSF) under a NSF Ocean Sciences Postdoctoral

Fellowship (Award No. 2205744). JES is a CIFAR Fellow in the program Fungal Kingdom:

Threats and Opportunities and was partially supported by NSF awards EF-2125066 and IOS-

2134912. Computations were performed using the computer clusters and data storage

resources of the UC Riverside HPCC, which were funded by grants from NSF (MRI-2215705,

MRI-1429826) and NIH (1S10OD016290-01A1). The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

673
674 CLE conceived and designed the experiments, performed sampling, analyzed the data,

675 prepared figures and/or tables, wrote and reviewed drafts of the paper. JAE and JES reviewed

676 drafts of the paper.

Acknowledgements

We would like to thank Katherine Dynarski (ORCID: 0000-0001-5101-9666) and Sonia Ghose

(ORCID: 0000-0001-5667-6876) for their help with initial sample collection. We would also like

to thank John J. Stachowicz for use of his scientific sampling permit, California Department of

Fish and Wildlife Scientific Collecting Permit # SC 4874.

References

1. Peay KG, Kennedy PG, Talbot JM. Dimensions of biodiversity in the Earth mycobiome. Nat Rev Microbiol. 2016;14. doi:10.1038/nrmicro.2016.59

2. Amend A, Burgaud G, Cunliffe M, Edgcomb VP, Ettinger CL, Gutiérrez MH, et al. Fungi in the Marine Environment: Open Questions and Unsolved Problems. MBio. 2019;10. doi:10.1128/mBio.01189-18

3. Peng X, Amend AS, Baltar F, Blanco-Bercial L, Breyer E, Burgaud G, et al. Planktonic Marine Fungi: A Review. Journal of Geophysical Research: Biogeosciences. 2024;129: e2023JG007887. doi:10.1029/2023JG007887

4. Hawksworth DL, Lücking R. Fungal Diversity Revisited: 2.2 to 3.8 Million Species. Microbiology spectrum. 2017;5. doi:10.1128/microbiolspec.FUNK-0052-2016

5. Jones EBG. Are there more marine fungi to be described? Botanica Marina. 2011;54: 343–354. doi:10.1515/bot.2011.043

6. Jones EBG, Suetrong S, Sakayaroj J, Bahkali AH, Abdel-Wahab MA, Boekhout T, et al. Classification of marine Ascomycota, Basidiomycota, Blastocladiomycota and Chytridiomycota. Fungal Divers. 2015;73: 1–72. doi:10.1007/s13225-015-0339-4

7. Jones EBG, Gareth Jones EB, Pang K-L, Abdel-Wahab MA, Scholz B, Hyde KD, et al. An online resource for marine fungi. Fungal Diversity. 2019;96: 347–433. doi:10.1007/s13225-019-00426-5

8. Tisthammer KH, Cobian GM, Amend AS. Global biogeography of marine fungi is shaped by the environment. Fungal Ecology. 2016;19: 39–46. doi:10.1016/j.funeco.2015.09.003

9. Morales SE, Biswas A, Herndl GJ, Baltar F. Global Structuring of Phylogenetic and Functional Diversity of Pelagic Fungi by Depth and Temperature. Frontiers in Marine Science. 2019;6. doi:10.3389/fmars.2019.00131

10. Breyer E, Stix C, Kilker S, Roller BRK, Panagou F, Doebke C, et al. The Overlooked Contribution of Pelagic Fungi to Ocean Biomass. 2024. doi:10.2139/ssrn.4790633

11. Breyer E, Zhao Z, Herndl GJ, Baltar F. Global contribution of pelagic fungi to protein degradation in the ocean. Microbiome. 2022;10. doi:10.1186/s40168-022-01329-5

12. Baltar F, Zhao Z, Herndl GJ. Potential and expression of carbohydrate utilization by marine
fungi in the global ocean. Microbiome. 2021;9. doi:10.1186/s40168-021-01063-4

13. Peng X, Valentine DL. Diversity and N2O Production Potential of Fungi in an Oceanic
Oxygen Minimum Zone. Journal of fungi (Basel, Switzerland). 2021;7.
doi:10.3390/jof7030218

14. Hassett BT, Vonnahme TR, Peng X, Gareth Jones EB, Heuzé C. Global diversity and
geography of planktonic marine fungi. Botanica Marina. 2020;63: 121–139.
doi:10.1515/bot-2018-0113

15. Hemminga MA, Duarte CM. Seagrass Ecology. Cambridge University Press; 2000.
doi:10.1017/CBO9780511525551

16. Fourqurean JW, Duarte CM, Kennedy H, Marbà N, Holmer M, Mateo MA, et al. Seagrass
ecosystems as a globally significant carbon stock. Nature Geoscience. 2012;5: 505–509.
doi:10.1038/ngeo1477

17. Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, et al. A
Global Crisis for Seagrass Ecosystems. BioScience. 2006;56: 987. doi:10.1641/0006-
3568(2006)56[987:agcfse]2.0.co;2

18. Cummins SP, Roberts DE, Ajani P, Underwood AJ. Comparisons of assemblages of
phytoplankton between open water and seagrass habitats in a shallow coastal lagoon. Mar
Freshwater Res. 2004;55: 447–456. doi:10.1071/MF04017

19. Prazukin AV, Lee RI, Firsov YK, Kapranov SV. Vertical distribution of epiphytic diatoms in
relation to the eelgrass *Zostera noltii* canopy biomass and height. Aquat Bot. 2022;176:
103466. doi:10.1016/j.aquabot.2021.103466

20. Lebreton B, Richard P, Galois R, Radenac G, Pfléger C, Guillou G, et al. Trophic
importance of diatoms in an intertidal *Zostera noltii* seagrass bed: Evidence from stable
isotope and fatty acid analyses. Estuar Coast Shelf Sci. 2011;92: 140–153.
doi:10.1016/j.ecss.2010.12.027

21. Kolátková V, Mooney M, Kelly K, Hineva E, Gawryluk RMR, Elliott J. Eelgrass (*Zostera*
spp.) associated phytomyxids are host-specific congeneric parasites and predominant
eukaryotes in the eelgrass rhizosphere on a global scale. Environ Microbiol. 2023;25:
1522–1537. doi:10.1111/1462-2920.16376

22. Elliott JK, Simpson H, Teesdale A, Replogle A, Elliott M, Coats K, et al. A Novel
Phagomyxid Parasite Produces Sporangia in Root Hair Galls of Eelgrass (*Zostera marina*).
Protist. 2019;170: 64–81. doi:10.1016/j.protis.2018.12.001

23. Man In 't Veld WA, Karin CH, van Rijswick PCJ, Meffert JP, Boer E, Westenberg M, et al.
Multiple *Halophytophthora* spp. and *Phytophthora* spp. including *P. gemini, P. inundata* and
P. chesapeakensis sp. nov. isolated from the seagrass *Zostera marina* in the Northern
hemisphere. European Journal of Plant Pathology. 2019;153: 341–357.
doi:10.1007/s10658-018-1561-1

24. Armitage AR, Frankovich TA, Heck KL, Fourqurean JW. Experimental nutrient enrichment
causes complex changes in seagrass, microalgae, and macroalgae community structure in

754    Florida Bay. Estuaries. 2005;28: 422–434. doi:10.1007/BF02693924

755 25. Ettinger CL, Vann LE, Eisen JA. Global Diversity and Biogeography of the *Zostera marina*
756    Mycobiome. Appl Environ Microbiol. 2021;87. doi:10.1128/AEM.02795-20

757 26. Bengtsson MM, Bühler A, Brauer A, Dahlke S, Schubert H, Blindow I. Eelgrass Leaf
758    Surface Microbiomes Are Locally Variable and Highly Correlated with Epibiotic Eukaryotes.
759    Front Microbiol. 2017;8: 1312. doi:10.3389/fmicb.2017.01312

760 27. Segovia BT, Sanders-Smith R, Adamczyk EM, Forbes C, Hessing-Lewis M, O'Connor MI,
761    et al. Microeukaryotic Communities Associated With the Seagrass *Zostera marina* Are
762    Spatially Structured. J Eukaryot Microbiol. 2021;68: e12827. doi:10.1111/jeu.12827

763 28. Shoemaker G, Wyllie-Echeverria S. Occurrence of rhizomal endophytes in three temperate
764    northeast pacific seagrasses. Aquatic Botany. 2013;111: 71–73.
765    doi:10.1016/j.aquabot.2013.05.010

766 29. Kirichuk NN, Pivkin MV. Filamentous fungi associated with the seagrass *Zostera marina*
767    Linnaeus, 1753 of Rifovaya Bay (Peter the Great Bay, the Sea of Japan). Russian Journal
768    of Marine Biology. 2015;41: 351–355. doi:10.1134/s1063074015050053

769 30. Petersen L-E, Marner M, Labes A, Tasdemir D. Rapid Metabolome and Bioactivity Profiling
770    of Fungi Associated with the Leaf and Rhizosphere of the Baltic Seagrass *Zostera marina*.
771    Marine Drugs. 2019;17: 419. doi:10.3390/md17070419

772 31. Ettinger CL, Eisen JA. Fungi, bacteria and oomycota opportunistically isolated from the
773    seagrass, *Zostera marina*. PLoS One. 2020;15: e0236135.
774    doi:10.1371/journal.pone.0236135

775 32. Zhang ZF, Liu F, Liu LR, Li M, Cai L, Liu S, et al. Culturing the uncultured marine fungi in
776    the omics age: Opportunities and challenges. Fungal Biol Rev. 2024;48: 100353.
777    doi:10.1016/j.fbr.2023.100353

778 33. Hurtado-McCormick V, Kahlke T, Petrou K, Jeffries T, Ralph PJ, Seymour JR. Regional
779    and Microenvironmental Scale Characterization of the Seagrass Microbiome. Front
780    Microbiol. 2019;10: 1011. doi:10.3389/fmicb.2019.01011

781 34. Abdel-Wahab MA, Bahkali AH, Elgorban AM, Jones EBG. High-throughput amplicon
782    sequencing of fungi and microbial eukaryotes associated with the seagrass *Halophila*
783    *stipulacea* (Forssk.) Asch. from Al-Leith mangroves, Saudi Arabia. Mycol Prog. 2021;20:
784    1365–1381. doi:10.1007/s11557-021-01744-2

785 35. Randolph Quek ZB, Zahn G, Lee NLY, Ooi JLS, Lee JN, Huang D, et al. Biogeographic
786    structure of fungal communities in seagrass *Halophilia ovalis* across the Malay Peninsula.
787    Environ Microbiol Rep. 2021;13: 871–877. doi:10.1111/1758-2229.13003

788 36. Ettinger CL, Eisen JA. Characterization of the Mycobiome of the Seagrass, *Zostera marina*,
789    Reveals Putative Associations With Marine Chytrids. Front Microbiol. 2019;10: 2476.
790    doi:10.3389/fmicb.2019.02476

791 37. Wainwright BJ, Zahn GL, Arlyza IS, Amend AS. Seagrass-associated fungal communities
792    follow Wallace's line, but host genotype does not structure fungal community. Journal of

Biogeography. 2018;45: 762–770. doi:10.1111/jbi.13168

38. Wainwright BJ, Zahn GL, Zushi J, Lee NLY, Ooi JLS, Lee JN, et al. Seagrass associated
fungal communities show distance decay of similarity that has implications for seagrass
management and restoration. Ecology and Evolution. 2019;9: 11288–11297.
doi:10.1002/ece3.5631

39. Trevathan-Tackett SM, Allnutt TR, Sherman CDH, Richardson ME, Crowley TM, Macreadie
PI. Spatial variation of bacterial and fungal communities of estuarine seagrass leaf
microbiomes. Aquatic Microbial Ecology. 2020;84: 59–74. doi:10.3354/ame01926

40. Pang K-L, Gareth Jones EB, Abdel-Wahab MA, Adams SJ, Alves A, Azevedo E, et al.
Recent progress in marine mycological research in different countries, and prospects for
future developments worldwide. Botanica Marina. 2023;66: 239–269. doi:10.1515/bot-2023-
0015

41. Gonçalves MF, Esteves AC, Alves A. Marine Fungi: Opportunities and Challenges.
Encyclopedia. 2022;2: 559–577. doi:10.3390/encyclopedia2010037

42. Hassett BT, Ducluzeau A-LL, Collins RE, Gradinger R. Spatial distribution of aquatic marine
fungi across the western Arctic and sub-arctic. Environ Microbiol. 2017;19: 475–484.
doi:10.1111/1462-2920.13371

43. Simmons DR, James TY, Meyer AF, Longcore JE. Lobulomycetales, a new order in the
Chytridiomycota. Mycol Res. 2009;113. doi:10.1016/j.mycres.2008.11.019

44. Longcore JE. Morphology and Zoospore Ultrastructure of *Chytriomyces Angularis* Sp. Nov.
(Chytridiales). Mycologia. 1992. doi:10.1080/00275514.1992.12026158

45. Laundon D, Cunliffe M. A Call for a Better Understanding of Aquatic Chytrid Biology. Front
Fungal Biol. 2021;2: 708813. doi:10.3389/ffunb.2021.708813

46. Sharon I, Morowitz MJ, Thomas BC, Costello EK, Relman DA, Banfield JF. Time series
community genomics analysis reveals rapid shifts in bacterial species, strains, and phage
during infant gut colonization. Genome Res. 2013;23: 111–120. doi:10.1101/gr.142315.112

47. Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. Genome
sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple
metagenomes. Nat Biotechnol. 2013;31: 533–538. doi:10.1038/nbt.2579

48. Bushnell B. BBTools. 2022. Available: https://sourceforge.net/projects/bbmap/

49. Ma X, Olsen JL, Reusch TBH, Procaccini G, Kudrna D, Williams M, et al. Improved
chromosome-level genome assembly and annotation of the seagrass, *Zostera marina*
(eelgrass). F1000Res. 2021;10. doi:10.12688/f1000research.38156.1

50. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biology. 2009;10: R25.
doi:10.1186/gb-2009-10-3-r25

51. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.
doi:10.1093/bioinformatics/btp352

52. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution
for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.
2015;31: 1674–1676. doi:10.1093/bioinformatics/btv033

53. Pronk LJU, Medema MH. Whokaryote: distinguishing eukaryotic and prokaryotic contigs in
metagenomes based on gene structure. Microbial genomics. 2022;8.
doi:10.1099/mgen.0.000823

54. Karlicki M, Antonowicz S, Karnkowska A. Tiara: deep learning-based classification system
for eukaryotic sequences. Bioinformatics. 2022;38: 344–350.
doi:10.1093/bioinformatics/btab672

55. Eren AM, Murat Eren A, Esen ÖC, Quince C, Vineis JH, Morrison HG, et al. Anvi'o: an
advanced analysis and visualization platform for 'omics data. PeerJ. 2015;3: e1319.
doi:10.7717/peerj.1319

56. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:
119. doi:10.1186/1471-2105-11-119

57. Lee MD. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics. 2019;35:
4162–4164. doi:10.1093/bioinformatics/btz188

58. Delmont T. Assessing the completion of eukaryotic bins with anvi'o. 2018. Available:
http://merenlab.org/delmont-euk-scgs

59. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7: e1002195.
doi:10.1371/journal.pcbi.1002195

60. Seemann T. barrnap: BAsic Rapid Ribosomal RNA Predictor. Available:
https://github.com/tseemann/barrnap

61. Menzel P, Ng KL, Krogh A. Fast and sensitive taxonomic classification for metagenomics
with Kaiju. Nat Commun. 2016;7: 11257. doi:10.1038/ncomms11257

62. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning
metagenomic contigs by coverage and composition. Nat Methods. 2014;11: 1144–1146.
doi:10.1038/nmeth.3103

63. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning
algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ. 2019;7: e7359. doi:10.7717/peerj.7359

64. Graham ED, Heidelberg JF, Tully BJ. BinSanity: unsupervised clustering of environmental
microbial assemblies using coverage and affinity propagation. PeerJ. 2017;5: e3035.
doi:10.7717/peerj.3035

65. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. MaxBin: an automated binning
method to recover individual genomes from metagenomes using an expectation-
maximization algorithm. Microbiome. 2014;2: 1–18. doi:10.1186/2049-2618-2-26

66. Saary P, Mitchell AL, Finn RD. Estimating the quality of eukaryotic genomes recovered
from metagenomic analysis with EukCC. Genome Biol. 2020;21: 244. doi:10.1186/s13059-

020-02155-4

67. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic
comparisons that enables improved genome recovery from metagenomes through de-
replication. ISME J. 2017;11: 2864–2868. doi:10.1038/ismej.2017.126

68. Alexander H, Hu SK, Krinos AI, Pachiadaki M, Tully BJ, Neely CJ, et al. Eukaryotic
genomes from a global metagenomic data set illuminate trophic modes and biogeography
of ocean plankton. MBio. 2023;14: e0167623. doi:10.1128/mbio.01676-23

69. Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. BlobToolKit – Interactive Quality
Assessment of Genome Assemblies. G3 Genes|Genomes|Genetics. 2020;10: 1361–1374.
doi:10.1534/g3.119.400908

70. Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale using
DIAMOND. Nat Methods. 2021;18: 366–368. doi:10.1038/s41592-021-01101-x

71. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST :
architecture and applications. BMC Bioinformatics. 2009;10. doi:10.1186/1471-2105-10-421

72. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34:
3094–3100. doi:10.1093/bioinformatics/bty191

73. Krinos AI, Hu SK, Cohen NR, Alexander H. EUKulele: Taxonomic annotation of the unsung
eukaryotic microbes. Journal of Open Source Software. 2021;6: 2817.
doi:10.21105/joss.02817

74. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinformatics.
2015;31: 3210–3212. doi:10.1093/bioinformatics/btv351

75. Stajich J, Palmer J. stajichlab/AAFTF: v0.4.1. 2021. doi:10.5281/zenodo.3437300

76. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for
automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A.
2020;117: 9451–9457. doi:10.1073/pnas.1921046117

77. Smit AFA, Hubley R, Green P. RepeatMasker. 2013-2015. Available:
http://www.repeatmasker.org

78. Palmer JM, Stajich J. Funannotate v1.8.1: Eukaryotic genome annotation. 2020. Available:
http://doi.org/10.5281/zenodo.4054262

79. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic Acids Res. 2006;34: W435–9.
doi:10.1093/nar/gkl200

80. Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. Gene prediction in novel
fungal genomes using an *ab initio* algorithm with unsupervised training. Genome Research.
2008;18: 1979–1990. doi:10.1101/gr.081612.108

81. Majoros WH, Pertea M, Salzberg SL. TigrScan and GlimmerHMM: two open source ab
initio eukaryotic gene-finders. Bioinformatics. 2004;20: 2878–2879.

doi:10.1093/bioinformatics/bth315

82. Korf I. Gene finding in novel genomes. BMC Bioinformatics. 2004;5: 59. doi:10.1186/1471-
2105-5-59

83. Testa AC, Hane JK, Ellwood SR, Oliver RP. CodingQuarry: highly accurate hidden Markov
model gene prediction in fungal genomes using RNA-seq transcripts. BMC Genomics.
2015;16: 170. doi:10.1186/s12864-015-1344-4

84. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al. Automated eukaryotic
gene structure annotation using EVidenceModeler and the Program to Assemble Spliced
Alignments. Genome Biol. 2008;9: R7. doi:10.1186/gb-2008-9-1-r7

85. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA
genes in genomic sequence. Nucleic Acids Res. 1997;25: 955–964.
doi:10.1093/nar/25.5.955

86. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the
protein families database. Nucleic Acids Res. 2014;42: D222–30. doi:10.1093/nar/gkt1223

87. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-
active enzymes database (CAZy) in 2013. Nucleic Acids Res. 2014;42: D490–5.
doi:10.1093/nar/gkt1178

88. Huang L, Zhang H, Wu P, Entwistle S, Li X, Yohe T, et al. dbCAN-seq: a database of
carbohydrate-active enzyme (CAZyme) sequence and annotation. Nucleic Acids Res.
2018;46: D516–D521. doi:10.1093/nar/gkx894

89. Rawlings ND, Barrett AJ, Bateman A. Using the MEROPS Database for Proteolytic
Enzymes and Their Inhibitors and Substrates. Curr Protoc Bioinformatics. 2014;48: 1.25.1–
33. doi:10.1002/0471250953.bi0125s48

90. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. eggNOG
4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic,
prokaryotic and viral sequences. Nucleic Acids Res. 2016;44: D286–93.
doi:10.1093/nar/gkv1248

91. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-
scale protein function classification. Bioinformatics. 2014;30: 1236–1240.
doi:10.1093/bioinformatics/btu031

92. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, et al.
UniProtKB/Swiss-Prot, the Manually Annotated Section of the UniProt KnowledgeBase:
How to Use the Entry View. Methods Mol Biol. 2016;1374: 23–54. doi:10.1007/978-1-4939-
3167-5_2

93. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat
Methods. 2015;12: 59–60. doi:10.1038/nmeth.3176

94. Käll L, Krogh A, Sonnhammer ELL. A combined transmembrane topology and signal
peptide prediction method. J Mol Biol. 2004;338: 1027–1036.
doi:10.1016/j.jmb.2004.03.016

95. Armenteros JJA, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et al.
SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat
Biotechnol. 2019;37: 420–423. doi:10.1038/s41587-019-0036-z

96. Medema MH, Blin K, Cimermancic P, de Jager V, Zakrzewski P, Fischbach MA, et al.
antiSMASH: rapid identification, annotation and analysis of secondary metabolite
biosynthesis gene clusters in bacterial and fungal genome sequences. Nucleic Acids Res.
2011;39: W339–46. doi:10.1093/nar/gkr466

97. Sperschneider J, Dodds PN. EffectorP 3.0: Prediction of Apoplastic and Cytoplasmic
Effectors in Fungi and Oomycetes. Mol Plant Microbe Interact. 2022;35: 146–156.
doi:10.1094/MPMI-08-21-0201-R

98. Hane JK, Paxman J, Jones DAB, Oliver RP, de Wit P. "CATAStrophy," a Genome-Informed
Trophic Classification of Filamentous Plant Pathogens - How Many Different Types of
Filamentous Plant Pathogens Are There? Front Microbiol. 2019;10: 3088.
doi:10.3389/fmicb.2019.03088

99. Johnson LK, Alexander H, Brown CT. Re-assembly, quality evaluation, and annotation of
678 microbial eukaryotic reference transcriptomes. Gigascience. 2019;8.
doi:10.1093/gigascience/giy158

100. Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, et al. The Marine
Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): illuminating the
functional diversity of eukaryotic life in the oceans through transcriptome sequencing. PLoS
Biol. 2014;12: e1001889. doi:10.1371/journal.pbio.1001889

101. Niang G, Hoebeke M, Meng A, Liu X, Scheremetjew M, Finn R, et al. METdb: A
GENOMIC REFERENCE DATABASE FOR MARINE SPECIES. F1000Res. 2020;9.
doi:10.7490/f1000research.1118000.1

102. Scott P, Taylor P, Gardner J, Puértolas A, Panda P, Addison S, et al. *Phytophthora
aleatoria* sp. nov., associated with root and collar damage on *Pinus radiata* from nurseries
and plantations. Australas Plant Pathol. 2019;48: 313–321. doi:10.1007/s13313-019-
00631-5

103. Wawrzyniak I, Courtine D, Osman M, Hubans-Pierlot C, Cian A, Nourrisson C, et al.
Draft genome sequence of the intestinal parasite *Blastocystis* subtype 4-isolate WR1.
Genomics Data. 2015;4: 22–23. doi:10.1016/j.gdata.2015.01.009

104. Sambles C, Schlenzig A, O'Neill P, Grant M, Studholme DJ. Draft genome sequences of
Phytophthora kernoviae and *Phytophthora ramorum* lineage EU2 from Scotland. Genomics
Data. 2015;6: 193. doi:10.1016/j.gdata.2015.09.010

105. Studholme DJ, McDougal RL, Sambles C, Hansen E, Hardy G, Grant M, et al. Genome
sequences of six *Phytophthora* species associated with forests in New Zealand. Genomics
data. 2015;7. doi:10.1016/j.gdata.2015.11.015

106. Minardi D, Studholme DJ, van der Giezen M, Pretto T, Oidtmann B. New genotyping
method for the causative agent of crayfish plague (*Aphanomyces astaci*) based on whole
genome data. J Invertebr Pathol. 2018;156. doi:10.1016/j.jip.2018.06.002

107.   Dittami SM, Corre E, Brillet-Guéguen L, Lipinska AP, Pontoizeau N, Aite M, et al. The
989   genome of *Ectocarpus subulatus* - A highly stress-tolerant brown alga. Mar Genomics.
990   2020;52. doi:10.1016/j.margen.2020.100740

991   108.   Kiselev A, San Clemente H, Camborde L, Dumas B, Gaulin E. A Comprehensive
992   Assessment of the Secretome Responsible for Host Adaptation of the Legume Root
993   Pathogen *Aphanomyces euteiches* . Journal of Fungi. 2022;8: 88. doi:10.3390/jof8010088

994   109.   Armitage AD, Lysøe E, Nellist CF, Lewis LA, Cano LM, Harrison RJ, et al. Bioinformatic
995   characterisation of the effector repertoire of the strawberry pathogen *Phytophthora
996   cactorum* . PLoS One. 2018;13: e0202305. doi:10.1371/journal.pone.0202305

997   110.   Studholme DJ, Panda P, Von Stowasser ES, González M, Hill R, Sambles C, et al.
998   Genome sequencing of oomycete isolates from Chile supports the New Zealand origin of
999   *Phytophthora kernoviae* and makes available the first *Nothophytophthora* sp. genome. Mol
1000   Plant Pathol. 2019;20: 423–431. doi:10.1111/mpp.12765

1001   111.   Gann ER, Truchon AR, Papoulis SE, Dyhrman ST, Gobler CJ, Wilhelm SW.
1002   *Aureococcus anophagefferens* (Pelagophyceae) genomes improve evaluation of nutrient
1003   acquisition strategies involved in brown tide dynamics. J Phycol. 2022;58.
1004   doi:10.1111/jpy.13221

1005   112.   Adams TM, Armitage AD, Sobczyk MK, Bates HJ, Tabima JF, Kronmiller BA, et al.
1006   Genomic Investigation of the Strawberry Pathogen *Phytophthora fragariae* Indicates
1007   Pathogenicity Is Associated With Transcriptional Variation in Three Key Races. Front
1008   Microbiol. 2020;11: 516181. doi:10.3389/fmicb.2020.00490

1009   113.   Gaulin E, Pel MJC, Camborde L, San-Clemente H, Courbier S, Dupouy M-A, et al.
1010   Genomics analysis of *Aphanomyces* spp. identifies a new class of oomycete effector
1011   associated with host adaptation. BMC Biol. 2018;16: 1–21. doi:10.1186/s12915-018-0508-5

1012   114.   Bowler C, Allen AE, Badger JH, Grimwood J, Jabbari K, Kuo A, et al. The
1013   *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. Nature.
1014   2008;456: 239–244. doi:10.1038/nature07410

1015   115.   Haas BJ, Kamoun S, Zody MC, Jiang RHY, Handsaker RE, Cano LM, et al. Genome
1016   sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans* . Nature.
1017   2009;461: 393–398. doi:10.1038/nature08358

1018   116.   Cock JM, Sterck L, Rouzé P, Scornet D, Allen AE, Amoutzias G, et al. The *Ectocarpus*
1019   genome and the independent evolution of multicellularity in brown algae. Nature. 2010;465:
1020   617–621. doi:10.1038/nature09016

1021   117.   Mock T, Otillar RP, Strauss J, McMullan M, Paajanen P, Schmutz J, et al. Evolutionary
1022   genomics of the cold-adapted diatom *Fragilariopsis cylindrus* . Nature. 2017;541: 536–540.
1023   doi:10.1038/nature20803

1024   118.   Radakovits R, Jinkerson RE, Fuerstenberg SI, Tae H, Settlage RE, Boore JL, et al. Draft
1025   genome sequence and genetic transformation of the oleaginous alga *Nannochloropsis
1026   gaditana* . Nat Commun. 2012;3: 1–11. doi:10.1038/ncomms1688

119.   Osuna-Cruz CM, Bilcke G, Vancaester E, De Decker S, Bones AM, Winge P, et al. The

Seminavis robusta genome provides insights into the evolutionary adaptations of benthic
diatoms. Nat Commun. 2020;11: 1–13. doi:10.1038/s41467-020-17191-8

120.   Hackl T, Martin R, Barenhoff K, Duponchel S, Heider D, Fischer MG. Four high-quality
draft genome assemblies of the marine heterotrophic nanoflagellate *Cafeteria*
roenbergensis . Scientific Data. 2020;7: 1–9. doi:10.1038/s41597-020-0363-4

121.   Yang M, Duan S, Mei X, Huang H, Chen W, Liu Y, et al. The *Phytophthora cactorum*
genome provides insights into the adaptation to host defense compounds and fungicides.
Sci Rep. 2018;8: 1–11. doi:10.1038/s41598-018-24939-2

122.   Hongo Y, Kimura K, Takaki Y, Yoshida Y, Baba S, Kobayashi G, et al. The genome of
the diatom *Chaetoceros tenuissimus* carries an ancient integrated fragment of an extant
virus. Sci Rep. 2021;11: 1–13. doi:10.1038/s41598-021-00565-3

123.   Oliver A, Podell S, Pinowska A, Traller JC, Smith SR, McClure R, et al. Diploid genomic
architecture of *Nitzschia inconspicua* , an elite biomass production diatom. Sci Rep.
2021;11: 1–14. doi:10.1038/s41598-021-95106-3

124.   Davison JR, Rajwani R, Zhao G, Bewley CA. The genome of antibiotic-producing
colonies of the Pelagophyte alga *Chrysophaeum taylorii* reveals a diverse and non-
canonical capacity for secondary metabolism. Sci Rep. 2023;13: 1–10.
doi:10.1038/s41598-023-38042-8

125.   Guérin N, Ciccarella M, Flamant E, Frémont P, Mangenot S, Istace B, et al. Genomic
adaptation of the picoeukaryote *Pelagomonas calceolata* to iron-poor oceans revealed by
a chromosome-scale genome sequence. Communications Biology. 2022;5: 1–14.
doi:10.1038/s42003-022-03939-z

126.   Makkonen J, Vesterbacka A, Martin F, Jussila J, Diéguez-Uribeondo J, Kortet R, et al.
Mitochondrial genomes and comparative genomics of *Aphanomyces astaci* and
Aphanomyces invadans . Sci Rep. 2016;6: 1–9. doi:10.1038/srep36089

127.   Gobler CJ, Berry DL, Dyhrman ST, Wilhelm SW, Salamov A, Lobanov AV, et al. Niche of
harmful alga *Aureococcus anophagefferens* revealed through ecogenomics. Proc Natl
Acad Sci U S A. 2011;108. doi:10.1073/pnas.1016106108

128.   Suzuki S, Ota S, Yamagishi T, Tuji A, Yamaguchi H, Kawachi M. Rapid transcriptomic
and physiological changes in the freshwater pennate diatom *Mayamaea pseudoterrestris*
in response to copper exposure. DNA Res. 2022;29: dsac037. doi:10.1093/dnares/dsac037

129.   Fletcher K, Martin F, Isakeit T, Cavanaugh K, Magill C, Michelmore R. The genome of
the oomycete *Peronosclerospora sorghi* , a cosmopolitan pathogen of maize and sorghum,
is inflated with dispersed pseudogenes. G3 Genes|Genomes|Genetics. 2023;13: jkac340.
doi:10.1093/g3journal/jkac340

130.   Misner I, Blouin N, Leonard G, Richards TA, Lane CE. The Secreted Proteins of *Achlya*
hypogyna and *Thraustotheca clavata* Identify the Ancestral Oomycete Secretome and
Reveal Gene Acquisitions by Horizontal Gene Transfer. Genome Biol Evol. 2014;7: 120–
135. doi:10.1093/gbe/evu276

131.   Ali SS, Shao J, Lary DJ, Kronmiller BA, Shen D, Strem MD, et al. *Phytophthora*

megakarya and *Phytophthora palmivora* , Closely Related Causal Agents of Cacao Black
Pod Rot, Underwent Increases in Genome Sizes and Gene Numbers by Different
Mechanisms. Genome Biol Evol. 2017;9: 536–557. doi:10.1093/gbe/evx021

132.   Carpinelli EC, Telatin A, Vitulo N, Forcato C, D'Angelo M, Schiavon R, et al.
Chromosome Scale Genome Assembly and Transcriptome Profiling of *Nannochloropsis*
gaditana in Nitrogen Depletion. Mol Plant. 2014;7: 323–335. doi:10.1093/mp/sst120

133.   Thorpe P, Vetukuri RR, Hedley PE, Morris J, Whisson MA, Welsh LRJ, et al. Draft
genome assemblies for tree pathogens *Phytophthora pseudosyringae* and *Phytophthora*
boehmeriae . G3 Genes|Genomes|Genetics. 2021;11: jkab282.
doi:10.1093/g3journal/jkab282

134.   Lee Y, Cho K-S, Seo J-H, Sohn KH, Prokchorchik M. Improved Genome Sequence and
Gene Annotation Resource for the Potato Late Blight Pathogen *Phytophthora infestans* .
Mol Plant Microbe Interact. 2020. doi:10.1094/MPMI-02-20-0023-A

135.   Fletcher K, Shin O-H, Clark KJ, Feng C, Putman AI, Correll JC, et al. Ancestral
Chromosomes for Family Peronosporaceae Inferred from a Telomere-to-Telomere Genome
Assembly of *Peronospora effusa* . Mol Plant Microbe Interact. 2022. doi:10.1094/MPMI-09-
21-0227-R

136.   Carleson NC, Caroline M. Press, Grünwald NJ. High-Quality, Phased Genomes of
Phytophthora ramorum Clonal Lineages NA1 and EU1. Mol Plant Microbe Interact. 2022.
doi:10.1094/MPMI-11-21-0264-A

137.   Botkin J, Chanda AK, Martin FN, Hirsch CD. A Reference Genome Sequence Resource
for the Sugar Beet Root Rot Pathogen *Aphanomyces cochlioides* . Mol Plant Microbe
Interact. 2022;35. doi:10.1094/MPMI-11-21-0277-A

138.   Tanaka T, Maeda Y, Veluchamy A, Tanaka M, Abida H, Maréchal E, et al. Oil
Accumulation by the Oleaginous Diatom *Fistulifera solaris* as Revealed by the Genome
and Transcriptome. Plant Cell. 2015;27: 162–176. doi:10.1105/tpc.114.135194

139.   Basu S, Patil S, Mapleson D, Russo MT, Vitale L, Fevola C, et al. Finding a partner in
the ocean: molecular and evolutionary bases of the response to sexual cues in a planktonic
diatom. New Phytol. 2017;215: 140–156. doi:10.1111/nph.14557

140.   Armbrust EV, Berges JA, Bowler C, Green BR, Martinez D, Putnam NH, et al. The
Genome of the Diatom *Thalassiosira Pseudonana* : Ecology, Evolution, and Metabolism.
Science. 2004. doi:10.1126/science.1101156

141.   Tyler BM, Tripathy S, Zhang X, Dehal P, Jiang RH, Aerts A, et al. *Phytophthora* genome
sequences uncover evolutionary origins and mechanisms of pathogenesis. Science.
2006;313. doi:10.1126/science.1128796

142.   Rujirawat T, Patumcharoenpol P, Lohnoo T, Yingyong W, Lerksuthirat T,
Tangphatsornruang S, et al. Draft Genome Sequence of the Pathogenic Oomycete *Pythium*
insidiosum Strain Pi-S, Isolated from a Patient with Pythiosis. Genome Announc. 2015;3.
doi:10.1128/genomeA.00574-15

143.   Zepernick BN, Truchon AR, Gann ER, Wilhelm SW. Draft Genome Sequence of the

Freshwater Diatom *Fragilaria crotonensis* SAG 28.96. Microbiology resource announcements. 2022;11. doi:10.1128/mra.00289-22

144. Stajich JE, Vu AL, Judelson HS, Vogel GM, Gore MA, Carlson MO, et al. High-Quality Reference Genome Sequence for the Oomycete Vegetable Pathogen *Phytophthora capsici* Strain LT1534. Microbiology resource announcements. 2021;10. doi:10.1128/MRA.00295-21

145. Mahan KM, Polle JEW, McKie-Krisberg Z, Lipzen A, Kuo A, Grigoriev IV, et al. Annotated Genome Sequence of the High-Biomass-Producing Yellow-Green Alga *Tribonema minus* . Microbiology resource announcements. 2021;10. doi:10.1128/MRA.00327-21

146. Reghu RJ, Chellappan BV, Beena SH, Sasi A, Vasudevan SE, Nair AS. Draft Genome Sequence of the Oomycete *Globisporangium splendens* Strain rgcb-1. Microbiology resource announcements. 2020;9. doi:10.1128/MRA.01006-19

147. Morgan WR, Tartar A. Decontamination and Annotation of the Draft Genome Sequence of the Oomycete *Lagenidium giganteum* ARSEF 373. Microbiology resource announcements. 2023;12. doi:10.1128/mra.01346-22

148. Seddiki K, Godart F, Aiese CR, Sanseverino W, Barakat M, Ortet P, et al. Sequencing, De Novo Assembly, and Annotation of the Complete Genome of a New Thraustochytrid Species, Strain CCAP_4062/3. Genome Announc. 2018;6. doi:10.1128/genomeA.01335-17

149. Ohan JA, Hovde BT, Zhang XL, Davenport KW, Chertkov O, Han C, et al. Nuclear Genome Assembly of the Microalga *Nannochloropsis salina* CCMP1776. Microbiology resource announcements. 2019;8. doi:10.1128/MRA.00750-19

150. Lommer M, Specht M, Roy A-S, Kraemer L, Andreson R, Gutowska MA, et al. Genome and low-iron response of an oceanic diatom adapted to chronic iron limitation. Genome Biol. 2012;13: 1–21. doi:10.1186/gb-2012-13-7-r66

151. Sharma R, Xia X, Cano LM, Evangelisti E, Kemen E, Judelson H, et al. Genome analyses of the sunflower pathogen *Plasmopara halstedii* provide insights into effector evolution in downy mildews and *Phytophthora* . BMC Genomics. 2015;16: 1–23. doi:10.1186/s12864-015-1904-7

152. Fletcher K, Klosterman SJ, Derevnina L, Martin F, Bertier LD, Koike S, et al. Comparative genomics of downy mildews reveals potential adaptations to biotrophy. BMC Genomics. 2018;19: 1–23. doi:10.1186/s12864-018-5214-8

153. Engelbrecht J, Duong TA, Prabhu SA, Seedat M, van den Berg N. Genome of the destructive oomycete *Phytophthora cinnamomi* provides insights into its pathogenicity and adaptive potential. BMC Genomics. 2021;22: 1–15. doi:10.1186/s12864-021-07552-y

154. Fletcher K, Zhang L, Gil J, Han R, Cavanaugh K, Michelmore R. AFLAP: assembly-free linkage analysis pipeline using k-mers from genome sequencing data. Genome Biol. 2021;22: 1–26. doi:10.1186/s13059-021-02326-x

155. Liu H, Ma X, Yu H, Fang D, Li Y, Wang X, et al. Genomes and virulence difference between two physiological races of *Phytophthora nicotianae.* Gigascience. 2016;5:

s13742–016–0108–7. doi:10.1186/s13742-016-0108-7

156.   Denoeud F, Roussel M, Noel B, Wawrzyniak I, Da Silva C, Diogon M, et al. Genome
sequence of the stramenopile *Blastocystis*, a human anaerobic parasite. Genome Biol.
2011;12: 1–16. doi:10.1186/gb-2011-12-3-r29

157.   Krajaejun T, Patumcharoenpol P, Rujirawat T, Kittichotirat W, Tangphatsornruang S.
MGI short-read genome assemblies of *Pythium insidiosum* (reclassified as *Pythium*
periculosum) strains Pi057C3 and Pi050C3. BMC Res Notes. 2023;16: 1–4.
doi:10.1186/s13104-023-06587-6

158.   Jiang RHY, de Bruijn I, Haas BJ, Belmonte R, Löbach L, Christie J, et al. Distinctive
Expansion of Potential Virulence Genes in the Genome of the Oomycete Fish Pathogen
Saprolegnia parasitica. PLoS Genet. 2013;9: e1003272. doi:10.1371/journal.pgen.1003272

159.   Matson MEH, Liang Q, Lonardi S, Judelson HS. Karyotype variation, spontaneous
genome rearrangements affecting chemical insensitivity, and expression level
polymorphisms in the plant pathogen *Phytophthora infestans* revealed using its first
chromosome-scale assembly. PLoS Pathog. 2022;18: e1010869.
doi:10.1371/journal.ppat.1010869

160.   Gentekaki E, Curtis BA, Stairs CW, Klimeš V, Eliáš M, Salas-Leiva DE, et al. Extreme
genome diversity in the hyper-prevalent parasitic eukaryote *Blastocystis.* PLoS Biol.
2017;15: e2003769. doi:10.1371/journal.pbio.2003769

161.   Faure C, Veyssière M, Boëlle B, San Clemente H, Bouchez O, Lopez-Roques C, et al.
Long-Read Genome Sequence of the Sugar Beet Rhizosphere Mycoparasite *Pythium*
oligandrum. G3 Genes|Genomes|Genetics. 2020;10: 431–436. doi:10.1534/g3.119.400746

162.   Dorrell RG, Kuo A, Füssy Z, Richardson EH, Salamov A, Zarevski N, et al. Convergent
evolution and horizontal gene transfer in Arctic Ocean microalgae. Life Science Alliance.
2023;6. doi:10.26508/lsa.202201833

163.   Mohammadi M, Smith EA, Stanghellini ME, Kaundal R. Insights into the Host Specificity
of a New Oomycete Root Pathogen, *Pythium brassicum* P1: Whole Genome Sequencing
and Comparative Analysis Reveals Contracted Regulation of Metabolism, Protein Families,
and Distinct Pathogenicity Repertoire. Int J Mol Sci. 2021;22: 9002.
doi:10.3390/ijms22169002

164.   McMullan M, Gardiner A, Bailey K, Kemen E, Ward BJ, Cevik V, et al. Evidence for
suppression of immunity as a driver for genomic introgressions and host range expansion
in races of *Albugo candida*, a generalist parasite. 2015. doi:10.7554/eLife.04550

165.   Kämper J, Kahmann R, Bölker M, Ma L-J, Brefort T, Saville BJ, et al. Insights from the
genome of the biotrophic fungal plant pathogen *Ustilago maydis*. Nature. 2006;444: 97–
101. doi:10.1038/nature05248

166.   Farrer RA, Martel A, Verbrugghe E, Abouelleil A, Ducatelle R, Longcore JE, et al.
Genomic innovations linked to infection strategies across emerging pathogenic chytrid
fungi. Nat Commun. 2017;8: 1–11. doi:10.1038/ncomms14742

167.   Mondo SJ, Dannebaum RO, Kuo RC, Louie KB, Bewick AJ, LaButti K, et al. Widespread

adenine N6-methylation of active genes in fungi. Nat Genet. 2017;49: 964–968.
doi:10.1038/ng.3859

168.   Haitjema CH, Gilmore SP, Henske JK, Solomon KV, de Groot R, Kuo A, et al. A parts list
for fungal cellulosomes revealed by comparative genomics. Nature Microbiology. 2017;2:
1–8. doi:10.1038/nmicrobiol.2017.87

169.   Ahrendt SR, Quandt CA, Ciobanu D, Clum A, Salamov A, Andreopoulos B, et al.
Leveraging single-cell genomics to expand the fungal tree of life. Nature Microbiology.
2018;3: 1417–1428. doi:10.1038/s41564-018-0261-0

170.   van de Vossenberg BTLH, Warris S, Nguyen HDT, van Gent-Pelzer MPE, Joly DL, van
de Geest HC, et al. Comparative genomics of chytrid fungi reveal insights into the obligate
biotrophic and pathogenic lifestyle of *Synchytrium endobioticum*. Sci Rep. 2019;9: 1–14.
doi:10.1038/s41598-019-45128-9

171.   Chang Y, Rochon D 'ann, Sekimoto S, Wang Y, Chovatia M, Sandor L, et al. Genome-
scale phylogenetic analyses confirm *Olpidium* as the closest living zoosporic fungus to the
non-flagellated, terrestrial fungi. Sci Rep. 2021;11: 1–12. doi:10.1038/s41598-021-82607-4

172.   Amses KR, Simmons DR, Longcore JE, Mondo SJ, Seto K, Jerônimo GH, et al. Diploid-
dominant life cycles characterize the early evolution of Fungi. Proceedings of the National
Academy of Sciences. 2022;119: e2116841119. doi:10.1073/pnas.2116841119

173.   Ettinger CL, Ostovar T, Yacoub M, Ahrendt S, Hice RH, Federici BA, et al. Genomes
and transcriptomes help unravel the complex life cycle of the blastoclad fungus,
Coelomomyces lativittatus, an obligate parasite of mosquitoes and microcrustaceans.
Mycologia. 2023. doi:10.1080/00275514.2023.2228182

174.   Leonard G, Galindo LJ, Milner DS, Avelar GM, Gomes-Vieira AL, Gomes SL, et al. A
Genome Sequence Assembly of the Phototactic and Optogenetic Model Fungus
Blastocladiella emersonii Reveals a Diversified Nucleotide-Cyclase Repertoire. Genome
Biol Evol. 2022;14: evac157. doi:10.1093/gbe/evac157

175.   Chang Y, Wang S, Sekimoto S, Aerts AL, Choi C, Clum A, et al. Phylogenomic Analyses
Indicate that Early Fungi Evolved Digesting Cell Walls of Algal Ancestors of Land Plants.
Genome Biol Evol. 2015;7: 1590–1601. doi:10.1093/gbe/evv090

176.   Gómez Luciano LB, Tsai IJ, Chuma I, Tosa Y, Chen Y-H, Li J-Y, et al. Blast Fungal
Genomes Show Frequent Chromosomal Changes, Gene Gains and Losses, and Effector
Gene Turnover. Mol Biol Evol. 2019;36: 1148–1161. doi:10.1093/molbev/msz045

177.   Clemons R, Yacoub M, Faust E, Felipe Toledo L, Jenkinson TS, Carvalho T, et al.
Discovery of an endogenous DNA virus in the amphibian killing fungus and its association
with pathogen genotype and virulence. bioRxiv. 2024. p. 2023.03.16.532857.
doi:10.1101/2023.03.16.532857

178.   Cuomo CA, Güldener U, Xu J-R, Trail F, Gillian Turgeon B, Di Pietro A, et al. The
Fusarium graminearum Genome Reveals a Link Between Localized Polymorphism and
Pathogen Specialization. Science. 2007. doi:10.1126/science.1143708

179.   Youssef NH, Couger MB, Struchtemeyer CG, Liggenstoffer AS, Prade RA, Najar FZ, et

al. The genome of the anaerobic fungus *Orpinomyces* sp. strain C1A reveals the unique
evolutionary history of a remarkable plant biomass degrader. Appl Environ Microbiol.
2013;79: 4620–4634. doi:10.1128/AEM.00821-13

180.   Russ C, Franz Lang B, Chen Z, Gujja S, Shea T, Zeng Q, et al. Genome Sequence of
Spizellomyces punctatus. Genome Announc. 2016. doi:10.1128/genomea.00849-16

181.   Wilken SE, Monk JM, Leggieri PA, Lawson CE, Lankiewicz TS, Seppälä S, et al.
Experimentally Validated Reconstruction and Analysis of a Genome-Scale Metabolic Model
of an Anaerobic Neocallimastigomycota Fungus. mSystems. 2021.
doi:10.1128/msystems.00002-21

182.   Brown JL, Swift CL, Mondo SJ, Seppala S, Salamov A, Singan V, et al. Co☐cultivation of
the anaerobic fungus *Caecomyces churrovis* with *Methanobacterium bryantii* enhances
transcription of carbohydrate binding modules, dockerins, and pyruvate formate lyases on
specific substrates. Biotechnol Biofuels. 2021;14: 1–16. doi:10.1186/s13068-021-02083-w

183.   Goodwin SB, Ben M'Barek S, Dhillon B, Wittenberg AHJ, Crane CF, Hane JK, et al.
Finished Genome of the Fungal Wheat Pathogen *Mycosphaerella graminicola* Reveals
Dispensome Structure, Chromosome Plasticity, and Stealth Pathogenesis. PLoS Genet.
2011;7: e1002070. doi:10.1371/journal.pgen.1002070

184.   Joneson S, Stajich JE, Shiu S-H, Rosenblum EB. Genomic Transition to Pathogenicity in
Chytrid Fungi. PLoS Pathog. 2011;7: e1002338. doi:10.1371/journal.ppat.1002338

185.   Li Y, Li Y, Jin W, Sharpton TJ, Mackie RI, Cann I, et al. Combined Genomic,
Transcriptomic, Proteomic, and Physiological Characterization of the Growth of
Pecoramyces sp. F1 in Monoculture and Co-culture With a Syntrophic Methanogen. Front
Microbiol. 2019;10: 442669. doi:10.3389/fmicb.2019.00435

186.   Ettinger CL, Eisen JA, Stajich JE. Genomic streamlining of seagrass-associated
Colletotrichum sp. may be related to its adaptation to a marine monocot host. bioRxiv.
2024. p. 2024.12.17.629027. doi:10.1101/2024.12.17.629027

187.   Read BA, Kegel J, Klute MJ, Kuo A, Lefebvre SC, Maumus F, et al. Pan genome of the
phytoplankton *Emiliania* underpins its global distribution. Nature. 2013;499: 209–213.
doi:10.1038/nature12221

188.   Hulatt CJ, Wijffels RH, Posewitz MC. The Genome of the Haptophyte *Diacronema lutheri*
(*Pavlova lutheri*, Pavlovales): A Model for Lipid Biosynthesis in Eukaryotic Algae. Genome
Biol Evol. 2021;13: evab178. doi:10.1093/gbe/evab178

189.   Hovde BT, Deodato CR, Hunsperger HM, Ryken SA, Yost W, Jha RK, et al. Genome
Sequence and Transcriptome Analyses of *Chrysochromulina tobin*: Metabolic Tools for
Enhanced Algal Fitness in the Prominent Order Prymnesiales (Haptophyceae). PLoS
Genet. 2015;11: e1005469. doi:10.1371/journal.pgen.1005469

190.   Steenwyk JL, Buida TJ 3rd, Li Y, Shen X-X, Rokas A. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. PLoS Biol. 2020;18:
e3001007. doi:10.1371/journal.pbio.3001007

191.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et

al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Mol Biol Evol. 2020;37: 1530–1534. doi:10.1093/molbev/msaa015

192.   Chernomor O, von Haeseler A, Minh BQ. Terrace aware data structure for phylogenomic
inference from supermatrices. Syst Biol. 2016;65: 997–1008. doi:10.1093/sysbio/syw037

193.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat Methods. 2017;14: 587–589.
doi:10.1038/nmeth.4285

194.   The R Project for Statistical Computing. Available: https://www.r-project.org/

195.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods
Ecol Evol. 2017;8: 28–36. doi:10.1111/2041-210X.12628

196.   Mise K, Iwasaki W. Unexpected absence of ribosomal protein genes from metagenome-
assembled genomes. ISME Commun. 2022;2: 118. doi:10.1038/s43705-022-00204-6

197.   Gruber-Vodicka HR, Seah BKB, Pruesse E. phyloFlash: Rapid Small-Subunit rRNA
Profiling and Targeted Assembly from Metagenomes. mSystems. 2020;5.
doi:10.1128/mSystems.00920-20

198.   Majaneva M, Rintala J-M, Piisilä M, Fewer DP, Blomster J. Comparison of wintertime
eukaryotic community from sea ice and open water in the Baltic Sea, based on sequencing
of the 18S rRNA gene. Polar Biol. 2011;35: 875–889. doi:10.1007/s00300-011-1132-9

199.   Edgcomb VP, Beaudoin D, Gast R, Biddle JF, Teske A. Marine subsurface eukaryotes:
the fungal majority. Environ Microbiol. 2011;13: 172–183. doi:10.1111/j.1462-
2920.2010.02318.x

200.   Monchy S, Sanciu G, Jobard M, Rasconi S, Gerphagnon M, Chabé M, et al. Exploring
and quantifying fungal diversity in freshwater lake ecosystems using rDNA
cloning/sequencing and SSU tag pyrosequencing. Environ Microbiol. 2011;13: 1433–1453.
doi:10.1111/j.1462-2920.2011.02444.x

201.   Jobard M, Rasconi S, Solinhac L, Cauchie HM, Sime-Ngando T. Molecular and
morphological diversity of fungi and the associated functions in three European nearby
lakes. Environ Microbiol. 2012;14. doi:10.1111/j.1462-2920.2012.02771.x

202.   Genitsaris S, Kormas KA, Moustaka-Gouni M. Microscopic eukaryotes living in a dying
lake (Lake Koronia, Greece). FEMS Microbiol Ecol. 2009;69: 75–83. doi:10.1111/j.1574-
6941.2009.00686.x

203.   Van den Wyngaert S, Rojas-Jimenez K, Seto K, Kagami M, Grossart HP. Diversity and
Hidden Host Specificity of Chytrids Infecting Colonial Volvocacean Algae. J Eukaryot
Microbiol. 2018;65. doi:10.1111/jeu.12632

204.   Rad-Menéndez C, Gerphagnon M, Garvetto A, Arce P, Badis Y, Sime-Ngando T, et al.
Rediscovering *Zygorhizidium affluens* Canter: Molecular Taxonomy, Infectious Cycle, and
Cryopreservation of a Chytrid Infecting the Bloom-Forming Diatom *Asterionella formosa*.
Appl Environ Microbiol. 2018;84. doi:10.1128/AEM.01826-18

205.   Stoeck T, Taylor GT, Epstein SS. Novel eukaryotes from the permanently anoxic
1308   Cariaco Basin (Caribbean Sea). Appl Environ Microbiol. 2003;69.
1309   doi:10.1128/AEM.69.9.5656-5663.2003

1310   206.   Karpov SA, Kobseva AA, Mamkaeva MA, Mamkaeva KA, Mikhailov KV, Mirzaeva GS, et
1311   al. *Gromochytrium mamkaevae* gen. & sp. nov. and two new orders: Gromochytriales and
1312   Mesochytriales (Chytridiomycetes). 2014. doi:10.3767/003158514X680234

1313   207.   James TY, Letcher PM, Longcore JE, Mozley-Standridge SE, Porter D, Powell MJ, et al.
1314   A molecular phylogeny of the flagellated fungi (Chytridiomycota) and description of a new
1315   phylum (Blastocladiomycota). Mycologia. 2006. doi:10.1080/15572536.2006.11832616

1316   208.   Hussels C. Molekulare Charakterisierung pilzlicher Lebensgemeinschaften in
1317   Ackerboeden. Technische Fachhochschule Berlin. 2002.

1318   209.   Kupper FC, Maier I, Muller DG, Loiseaux-de Goer S, Guillou L. Phylogenetic affinities of
1319   two eukaryotic pathogens of marine macroalgae, *Eurychasma dicksonii* and *Chytridium*
1320   *polysiphoniae*. Cryptogamie, Algol. 2006;27: 165–184. Available:
1321   https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/cryptogamie-
1322   algologie2007v27f2a11.pdf

1323   210.   Nawrocki EP. Structural RNA Homology Search and Alignment Using Covariance
1324   Models. Washington University in Saint Louis. 2009. Available:
1325   https://www.proquest.com/docview/305017419

1326   211.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated
1327   alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25: 1972–
1328   1973. doi:10.1093/bioinformatics/btp348

1329   212.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
1330   genomics. Genome Biol. 2019;20: 238. doi:10.1186/s13059-019-1832-y

1331   213.   Kolde R. pheatmap: Pretty Heatmaps. 2018. Available:
1332   https://github.com/raivokolde/pheatmap

1333   214.   Onyshchenko A, Roberts WR, Ruck EC, Lewis JA, Alverson AJ. The genome of a
1334   nonphotosynthetic diatom provides insights into the metabolic shift to heterotrophy and
1335   constraints on the loss of photosynthesis. New Phytol. 2021;232: 1750–1764.
1336   doi:10.1111/nph.17673

1337   215.   Longcore JE, Qin S, Simmons DR, James TY. *Quaeritorhiza haematococci* is a new
1338   species of parasitic chytrid of the commercially grown alga, *Haematococcus pluvialis*.
1339   Mycologia. 2020;112: 606–615. doi:10.1080/00275514.2020.1730136

1340   216.   Simmons DR, Bonds AE, Castillo BT, Clemons RA, Glasco AD, Myers JM, et al. The
1341   Collection of Zoosporic Eufungi at the University of Michigan (CZEUM): introducing a new
1342   repository of barcoded and cultures. IMA Fungus. 2020;11: 20. doi:10.1186/s43008-020-
1343   00041-z

1344   217.   Chung M-H, Lee K-S. Species Composition of the Epiphytic Diatoms on the Leaf
1345   Tissues of Three *Zostera* Species Distributed on the Southern Coast of Korea. Algae. 23:
1346   75–81. doi:10.4490/algae.2008.23.1.075

218. Lebreton B, Richard P, Radenac G, Bordes M, Bréret M, Arnaud C, et al. Are epiphytes
a significant component of intertidal *Zostera noltii* beds? Aquat Bot. 2009;91: 82–90.
doi:10.1016/j.aquabot.2009.03.003

219. Atmaja PSP, Bengen DG, Madduppa HH. The Second Skin of Seagrass Leaves: A
Comparison of Microalgae Epiphytic Communities Between Two Different Species Across
Two Seagrass Meadows in Lesser Sunda Islands. Tropical Life Sciences Research.
2021;32: 97. doi:10.21315/tlsr2021.32.2.7

220. Pinckney JL, Micheli F. Microalgae on seagrass mimics: Does epiphyte community
structure differ from live seagrasses? J Exp Mar Bio Ecol. 1998;221: 59–70.
doi:10.1016/S0022-0981(97)00115-9

221. Erin Cox T, Cebrian J, Tabor M, West L, Krause JW. Do diatoms dominate benthic
production in shallow systems? A case study from a mixed seagrass bed. Limnology and
Oceanography Letters. 2020;5: 425–434. doi:10.1002/lol2.10167

222. Moncreiff CA, Sullivan M. J., Daehnick AE. Primary production dynamics in seagrass
beds of Mississippi Sound: the contributions of seagrass, epiphytic algae, sand microflora,
and phytoplankton. Marine Ecology-Progress Series. 2993;87: 161–161. Available:
https://www.jstor.org/stable/24831605

223. Coleman VL, Burkholder JM. Community structure and productivity of epiphytic
microalgae on eelgrass (*Zostera marina* L.) under water-column nitrate enrichment. J Exp
Mar Bio Ecol. 1994;179: 29–48. doi:10.1016/0022-0981(94)90015-9

224. Hasegawa N, Hori M, Mukai H. Seasonal shifts in seagrass bed primary producers in a
cold-temperate estuary: Dynamics of eelgrass *Zostera marina* and associated epiphytic
algae. Aquat Bot. 2007;86: 337–345. doi:10.1016/j.aquabot.2006.12.002

225. Frenken T, Alacid E, Berger SA, Bourne EC, Gerphagnon M, Grossart HP, et al.
Integrating chytrid fungal parasites into plankton ecology: research gaps and needs.
Environ Microbiol. 2017;19. doi:10.1111/1462-2920.13827

226. Wawra S, Fesel P, Widmer H, Timm M, Seibel J, Leson L, et al. The fungal-specific β-
glucan-binding lectin FGB1 alters cell-wall composition and suppresses glucan-triggered
immunity in plants. Nat Commun. 2016;7. doi:10.1038/ncomms13188

227. Martin F, Aerts A, Ahrén D, Brun A, Danchin EG, Duchaussoy F, et al. The genome of
Laccaria bicolor provides insights into mycorrhizal symbiosis. Nature. 2008;452.
doi:10.1038/nature06556

228. Plett JM, Kemppainen M, Kale SD, Kohler A, Legué V, Brun A, et al. A secreted effector
protein of *Laccaria bicolor* is required for symbiosis development. Curr Biol. 2011;21.
doi:10.1016/j.cub.2011.05.033

229. Voß S, Betz R, Heidt S, Corradi N, Requena N. RiCRN1, a Crinkler Effector From the
Arbuscular Mycorrhizal Fungus *Rhizophagus irregularis*, Functions in Arbuscule
Development. Front Microbiol. 2018;9. doi:10.3389/fmicb.2018.02068

230. Koeck M, Hardham AR, Dodds PN. The role of effectors of biotrophic and hemibiotrophic
fungi in infection. Cell Microbiol. 2011;13. doi:10.1111/j.1462-5822.2011.01665.x

231.   Tanaka S, Kahmann R. Cell wall-associated effectors of plant-colonizing fungi.
Mycologia. 2021;113. doi:10.1080/00275514.2020.1831293

232.   Thekkiniath J, Zabet-Moghaddam M, Kottapalli KR, Pasham MR, San FS, San FM.
Quantitative Proteomics of an Amphibian Pathogen, *Batrachochytrium dendrobatidis*,
following Exposure to Thyroid Hormone. PLoS One. 2015;10.
doi:10.1371/journal.pone.0123637

233.   Ellison AR, DiRenzo GV, McDonald CA, Lips KR, Zamudio KR. First in Vivo
Batrachochytrium dendrobatidis Transcriptomes Reveal Mechanisms of Host Exploitation,
Host-Specific Gene Expression, and Expressed Genotype Shifts. G3. 2017;7.
doi:10.1534/g3.116.035873

234.   de Vossenberg BTLH V, Prodhomme C, van Arkel G, van Gent-Pelzer MPE, Bergervoet
M, Brankovics B, et al. The *Synchytrium endobioticum* AvrSen1 Triggers a Hypersensitive
Response in Sen1 Potatoes While Natural Variants Evade Detection. Mol Plant Microbe
Interact. 2019;32. doi:10.1094/MPMI-05-19-0138-R

235.   Glass NL, Schmoll M, Cate JH, Coradetti S. Plant cell wall deconstruction by
ascomycete fungi. Annu Rev Microbiol. 2013;67. doi:10.1146/annurev-micro-092611-
150044

236.   Zhao Z, Liu H, Wang C, Xu JR. Comparative analysis of fungal genomes reveals
different plant cell wall degrading capacity in fungi. BMC Genomics. 2013;14.
doi:10.1186/1471-2164-14-274

237.   Ospina-Giraldo, Griffith JG, Laird EW, Mingora C. The CAZyome of *Phytophthora* spp.: a
comprehensive analysis of the gene complement coding for carbohydrate-active enzymes
in species of the genus *Phytophthora*. BMC Genomics. 2010;11. doi:10.1186/1471-2164-
11-525

238.   Rodriguez-Moreno L, Ebert MK, Bolton, Bphj T. Tools of the crook- infection strategies of
fungal plant pathogens. Plant J. 2018;93. doi:10.1111/tpj.13810

239.   Kubicek CP, Starr TL, Glass NL. Plant cell wall-degrading enzymes and their secretion
in plant-pathogenic fungi. Annu Rev Phytopathol. 2014;52. doi:10.1146/annurev-phyto-
102313-045831

240.   Lu H, Wei T, Lou H, Shu X, Chen Q. A Critical Review on Communication Mechanism
within Plant-Endophytic Fungi Interactions to Cope with Biotic and Abiotic Stresses. Journal
of fungi (Basel, Switzerland). 2021;7. doi:10.3390/jof7090719

241.   de Wit PJGM, Mehrabi R, Van den Burg HA, Stergiopoulos I. Fungal effector proteins:
past, present and future. Mol Plant Pathol. 2009;10: 735. doi:10.1111/j.1364-
3703.2009.00591.x

242.   Lange L, Barrett K, Pilgaard B, Gleason F, Tsang A. Enzymes of early-diverging,
zoosporic fungi. Appl Microbiol Biotechnol. 2019;103: 6885–6902. doi:10.1007/s00253-019-
09983-w

243.   Rosling A, Eshghi Sahraei S, Kalsoom Khan F, Desirò A, Bryson AE, Mondo SJ, et al.
Evolutionary history of arbuscular mycorrhizal fungi and genomic signatures of obligate

symbiosis. BMC Genomics. 2024;25: 529. doi:10.1186/s12864-024-10391-2

244.   Chang Y, Desirò A, Na H, Sandor L, Lipzen A, Clum A, et al. Phylogenomics of
Endogonaceae and evolution of mycorrhizas within Mucoromycota. New Phytol. 2019;222:
511–525. doi:10.1111/nph.15613

245.   Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, et al. Genome of an
arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. Proc Natl
Acad Sci U S A. 2013;110: 20117–20122. doi:10.1073/pnas.1313452110

246.   Newell SY, Fell JW. Mycoflora of Turtlegrass (*Thalassia testudinum* König) as Recorded
after Seawater Incubation. Botanica Marina. 1980;23: 265–276. doi:10.1515/bot-1980-
230409

247.   Poulin R, Krasnov BR, Mouillot D. Host specificity in phylogenetic and geographic space.
Trends Parasitol. 2011;27. doi:10.1016/j.pt.2011.05.003

248.   den Wyngaert S V, Ganzert L, Seto K, Rojas-Jimenez K, Agha R, Berger SA, et al.
Seasonality of parasitic and saprotrophic zoosporic fungi: linking sequence data to
ecological traits. ISME J. 2022;16. doi:10.1038/s41396-022-01267-y

249.   Kagami M, Seto K, Nozaki D, Nakamura T, Wakana H, Wurzbacher C. Single dominant
diatom can host diverse parasitic fungi with different degree of host specificity. Limnol
Oceanogr. 2021;66: 667–677. doi:10.1002/lno.11631

250.   Sassenhagen I, Irion S, Jardillier L, Moreira D, Christaki U. Protist Interactions and
Community Structure During Early Autumn in the Kerguelen Region (Southern Ocean).
Protist. 2020;171. doi:10.1016/j.protis.2019.125709

251.   Sassenhagen I, Langenheder S, Lindström ES. Infection strategies of different chytrids
in a diatom spring bloom. Freshwater Biology. 2023;68: 972–986. doi:10.1111/fwb.14079

252.   Haraldsson M, Gerphagnon M, Bazin P, Colombet J, Tecchio S, Sime-Ngando T, et al.
Microbial parasites make cyanobacteria blooms less of a trophic dead end than commonly
assumed. ISME J. 2018;12: 1008–1020. doi:10.1038/s41396-018-0045-9

253.   Garvetto A, Badis Y, Perrineau M-M, Rad-Menéndez C, Bresnan E, Gachon CMM.
Chytrid infecting the bloom-forming marine diatom *Skeletonema* sp.: Morphology,
phylogeny and distribution of a novel species within the Rhizophydiales. Fungal Biol.
2019;123: 471–480. doi:10.1016/j.funbio.2019.04.004

254.   Fernández-Valero AD, Reñé A, Timoneda N, Sampedro N, Garcés E. Dinoflagellate
hosts determine the community structure of marine Chytridiomycota: Demonstration of their
prominent interactions. Environ Microbiol. 2022;24. doi:10.1111/1462-2920.16182

255.   McKindles KM, Jorge AN, McKay RM, Davis TW, Bullerjahn GS. Isolation and
Characterization of *Rhizophydiales* sp. (Chytridiomycota), Obligate Parasite of *Planktothrix*
agardhii in a Laurentian Great Lakes Embayment. Appl Environ Microbiol. 2021;87.
doi:10.1128/AEM.02308-20

256.   Seto K, Degawa Y. *Cyclopsomyces plurioperculatus*: a new genus and species of
Lobulomycetales (Chytridiomycota, Chytridiomycetes) from Japan. Mycologia. 2015;107:

633–640. doi:10.3852/14-284

257.   Hassett BT, Gradinger R. Chytrids dominate arctic marine fungal communities. Environ Microbiol. 2016;18: 2001–2009. doi:10.1111/1462-2920.13216

258.   Rojas-Jimenez K, Rieck A, Wurzbacher C, Jürgens K, Labrenz M, Grossart H-P. A Salinity Threshold Separating Fungal Communities in the Baltic Sea. Front Microbiol. 2019;10: 680. doi:10.3389/fmicb.2019.00680

259.   Brubaker SW, Bonham KS, Zanoni I, Kagan JC. Innate immune pattern recognition: a cell biological perspective. Annu Rev Immunol. 2015;33. doi:10.1146/annurev-immunol-032414-112240

260.   Couto D, Zipfel C. Regulation of pattern recognition receptor signalling in plants. Nat Rev Immunol. 2016;16. doi:10.1038/nri.2016.77

261.   Soanes DM, Talbot NJ. Comparative genome analysis reveals an absence of leucine-rich repeat pattern-recognition receptor proteins in the kingdom Fungi. PLoS One. 2010;5: e12725. doi:10.1371/journal.pone.0012725

262.   Daskalov A. Emergence of the fungal immune system. iScience. 2023;26. doi:10.1016/j.isci.2023.106793

263.   Gaspar ML, Pawlowska TE. Innate immunity in fungi: Is regulated cell death involved? PLoS Pathog. 2022;18. doi:10.1371/journal.ppat.1010460

264.   Muehlstein LK, Amon JP, Leffler DL. Chemotaxis in the Marine Fungus *Rhizophydium littoreum*. Appl Environ Microbiol. 1988;54. doi:10.1128/aem.54.7.1668-1672.1988

265.   Moss AS, Reddy NS, Dortaj IM, Mj SF. Chemotaxis of the amphibian pathogen *Batrachochytrium dendrobatidis* and its response to a variety of attractants. Mycologia. 2008;100. doi:10.3852/mycologia.100.1.1

266.   Scholz B, Küpper FC, Vyverman W, Ólafsson HG, Karsten U. Chytridiomycosis of Marine Diatoms-The Role of Stress Physiology and Resistance in Parasite-Host Recognition and Accumulation of Defense Molecules. Mar Drugs. 2017;15. doi:10.3390/md15020026

267.   Pfeifer L, van Erven G, Sinclair EA, Duarte CM, Kabel MA. Profiling the cell walls of seagrasses from A (*Amphibolis*) to Z (*Zostera*). BMC Plant Biol. 2022;22. doi:10.1186/s12870-022-03447-6

268.   Wang Y, Verbrugghe E, Meuris L, Chiers K, Kelly M, Strubbe D, et al. Epidermal galactose spurs chytrid virulence and predicts amphibian colonization. Nat Commun. 2021;12: 5788. doi:10.1038/s41467-021-26127-9

269.   Rovenich H, Boshoven JC, Thomma BP. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. Curr Opin Plant Biol. 2014;20. doi:10.1016/j.pbi.2014.05.001

270.   Sánchez-Vallet A, Mesters JR, Thomma BP. The battle for chitin recognition in plant-microbe interactions. FEMS Microbiol Rev. 2015;39. doi:10.1093/femsre/fuu003

271.	Cao C, Xue C. More Than Just Cleaning: Ubiquitin-Mediated Proteolysis in Fungal Pathogenesis. Front Cell Infect Microbiol. 2021;11: 774613. doi:10.3389/fcimb.2021.774613

272.	Finley D, Ulrich HD, Sommer T, Kaiser P. The ubiquitin-proteasome system of *Saccharomyces cerevisiae*. Genetics. 2012;192. doi:10.1534/genetics.112.140467

273.	Liu TB, Xue C. The Ubiquitin-Proteasome System and F-box Proteins in Pathogenic Fungi. Mycobiology. 2011;39. doi:10.5941/MYCO.2011.39.4.243

274.	Pesin JA, Orr-Weaver TL. Regulation of APC/C activators in mitosis and meiosis. Annu Rev Cell Dev Biol. 2008;24. doi:10.1146/annurev.cellbio.041408.115949

275.	Rotin D, Kumar S. Physiological functions of the HECT family of ubiquitin ligases. Nat Rev Mol Cell Biol. 2009;10. doi:10.1038/nrm2690

276.	Grifoll-Romero L, Pascual S, Aragunde H, Biarnés X, Planas A. Chitin Deacetylases: Structures, Specificities, and Biotech Applications. Polymers . 2018;10. doi:10.3390/polym10040352

277.	Khokhani D, Carrera Carriel C, Vayla S, Irving TB, Stonoha-Arther C, Keller NP, et al. Deciphering the Chitin Code in Plant Symbiosis, Defense, and Microbial Networks. Annu Rev Microbiol. 2021;75: 583–607. doi:10.1146/annurev-micro-051921-114809

278.	Oldroyd GED. Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. Nat Rev Microbiol. 2013;11: 252–263. doi:10.1038/nrmicro2990

279.	Aspeborg H, Schrader J, Coutinho PM, Stam M, Kallas A, Djerbi S, et al. Carbohydrate-active enzymes involved in the secondary cell wall biogenesis in hybrid aspen. Plant Physiol. 2005;137. doi:10.1104/pp.104.055087

280.	Beauvais A, Latgé J-P. Special Issue: Fungal Cell Wall. Journal of Fungi. 2018;4: 91. doi:10.3390/jof4030091

281.	Henrissat B, Davies G. Structural and sequence-based classification of glycoside hydrolases. Curr Opin Struct Biol. 1997;7. doi:10.1016/s0959-440x(97)80072-3

282.	Ökmen B, Bachmann D, de Wit PJGM. A conserved GH17 glycosyl hydrolase from plant pathogenic Dothideomycetes releases a DAMP causing cell death in tomato. Mol Plant Pathol. 2019;20: 1710–1721. doi:10.1111/mpp.12872

283.	Chrismas N, Cunliffe M. Depth-dependent mycoplankton glycoside hydrolase gene activity in the open ocean-evidence from the Tara Oceans eukaryote metatranscriptomes. ISME J. 2020;14: 2361–2365. doi:10.1038/s41396-020-0687-2

284.	Cunliffe M, Hollingsworth A, Bain C, Sharma V, Taylor JD. Algal polysaccharide utilisation by saprotrophic planktonic marine fungi. Fungal Ecol. 2017;30: 135–138. doi:10.1016/j.funeco.2017.08.009

285.	Becker S, Tebben J, Coffinet S, Wiltshire K, Iversen MH, Harder T, et al. Laminarin is a major molecule in the marine carbon cycle. Proc Natl Acad Sci U S A. 2020;117. doi:10.1073/pnas.1917001117

1543   286.   Abramyan J, Stajich JE. Species-specific chitin-binding module 18 expansion in the
1544   amphibian pathogen *Batrachochytrium dendrobatidis*. MBio. 2012;3: e00150–12.
1545   doi:10.1128/mBio.00150-12

1546   287.   Powell AJ, Conant GC, Brown DE, Carbone I, Dean RA. Altered patterns of gene
1547   duplication and differential gene gain and loss in fungal pathogens. BMC Genomics.
1548   2008;9. doi:10.1186/1471-2164-9-147

1549   288.   Sexton AC, Howlett BJ. Parallels in fungal pathogenesis on plant and animal hosts.
1550   Eukaryot Cell. 2006;5. doi:10.1128/EC.00277-06

1551   289.   Rappleye CA, Goldman WE. Defining virulence genes in the dimorphic fungi. Annu Rev
1552   Microbiol. 2006;60. doi:10.1146/annurev.micro.59.030804.121055

1553   290.   Halaouli S, Asther M, Sigoillot J-C, Hamdi M, Lomascolo A. Fungal tyrosinases: new
1554   prospects in molecular characteristics, bioengineering and biotechnological applications. J
1555   Appl Microbiol. 2006;100: 219–232. doi:10.1111/j.1365-2672.2006.02866.x

1556   291.   Camejo D, Guzmán-Cedeño Á, Moreno A. Reactive oxygen species, essential
1557   molecules, during plant-pathogen interactions. Plant Physiol Biochem. 2016;103: 10–23.
1558   doi:10.1016/j.plaphy.2016.02.035

1559   292.   Perlin MH, Andrews J, San Toh S, editors. Essential Letters in the Fungal Alphabet:
1560   ABC and MFS Transporters and Their Roles in Survival and Pathogenicity. Advances in
1561   Genetics. Academic Press; 2014. pp. 201–253. doi:10.1016/B978-0-12-800271-1.00004-4

Figure 1. SGEUK-01, SGEUK-02 and SG-EUK04 are phylogenetically placed in the family

Bacillariaceae. From left to right, first, a maximum likelihood phylogeny that shows the

placement of SGEUK-01, SGEUK-02 and SG-EUK04 in the gyrista tree of life among diatom

lineages. This tree was generated using IQ-TREE2 on an alignment of BUSCO

stramenopiles_odb10 HMMs constructed using the PHYling_unified pipeline. Taxon labels in

the phylogeny are shown colored by their assigned taxonomic family. Next, in association with

this phylogeny, bar charts of the BUSCO "protein" completion status for the eukaryota_odb10

and stramenopiles_odb10 sets is shown. Bars show the percentage of genes found in each

genome annotation as a percentage of the total gene set and are colored by BUSCO status

(missing = gray, fragmented = yellow, complete and duplicated = green, complete and single

copy = blue).

Figure 2. SGEUK-03 is phylogenetically placed in the order Lobulomycetales. From left to right,

first, a maximum likelihood phylogeny that shows the placement of SGEUK-03 in the fungal tree

of life. This tree was generated using IQ-TREE2 on an alignment of BUSCO fungi_odb10 HMMs

constructed using the PHYling_unified pipeline. Taxon labels in the phylogeny are shown

colored by their assigned taxonomic order. Next, in association with this phylogeny, bar charts

of the BUSCO "protein" completion status for the eukaryota_odb10 and fungi_odb10 sets is

shown. Bars show the percentage of genes found in each genome annotation as a percentage

of the total gene set and are colored by BUSCO status (missing = gray, fragmented = yellow,

complete and duplicated = green, complete and single copy = blue).

1611

Figure 3. SGEUK-03 18S ribosomal RNA gene tree confirms placement in the

Lobulomycetales in Novel Clade SW-I. Given the lack of closely related genomes to SGEUK-03,

we used phyloFlash to assemble the 18S rRNA gene associated with SGEUK-03 and then

aligned it to close relatives using SSU-align. The alignment was trimmed with trimAl, and IQ-

TREE2 was run with ModelFinderPlus, to identify and use the optimal evolutionary model, and

1000 ultrafast bootstraps. Taxon labels in the phylogeny are colored and indicated by bars

based on their assigned taxonomic order. Additionally, Novel Clade SW-I is also indicated by a

bar. Displayed at each node as a circle in the tree are the ultrafast bootstrap values (e.g. a black

circle represents bootstrap values greater or equal to 90%, a grey circle represents bootstrap

values greater or equal to 70%, a white circle represents bootstrap values less than 70%).

Figure 4. SGEUK-05 is phylogenetically placed in the genus Prymnesium. From left to right,

first, a maximum likelihood phylogeny that shows the placement of SGEUK-05 in the haptista

tree of life among haptophyte lineages. This tree was generated using IQ-TREE2 on an

alignment of BUSCO stramenopiles_odb10 HMMs constructed using the PHYling_unified

pipeline. Taxon labels in the phylogeny are shown colored by their assigned taxonomic family.

Next, in association with this phylogeny, a bar chart of the BUSCO "protein" completion status

for the eukaryota_odb10 set is shown. Bars show the percentage of genes found in each

genome annotation as a percentage of the total gene set and are colored by BUSCO status

(missing = gray, fragmented = yellow, complete and duplicated = green, complete and single

copy = blue).

Figure 5. Only some high-copy orthogroups in SGEUK-3 are shared with close relatives. A

heatmap is depicted here visualizing the log-scaled copy numbers of phylogenetically

hierarchical orthogroups (HOGs) with 5 or more copies in SGEUK-03 in comparison to the

average copy number of the same HOGs in other fungal taxonomic groups (i.e., other members

of the order Lobulomycetales, other Chytridiomycota lineages, or other fungal phyla).

Additionally, whether these genes are predicted in the annotation of SGEUK-03 to be

membrane-bound, secreted or as plant-effector is indicated on the left of the heatmap. Further,

the dendrogram to the left of the heatmap clusters the HOGs by similarity in log-scaled counts

between the different HOGs.

Table 1. Eukaryotic metagenome-assembled genome assembly and annotation statistics. We identified five high-quality (> 30%

complete) eukaryotic MAGs associated with *Z. marina* leaves. Here, we list their proposed taxonomy, report on various genome and

annotation quality statistics, and provide the NCBI accession information for each MAG.

1675

Bin ID	Phylum	Putative taxonomy	EukCC Completion	Genome size (Mbp)	No. of contigs	N50 (bp)	GC content (%)	Mean coverage (x)	Repetitive regions (%)	No. of gene models	No. of mRNAs	No. of tRNAs	Genes with InterProScan hit (%)	Genes with eggNOG hit (%)	Genes with GO term (%)	Genes with CAZyme hit (#)	NCBI Accession
SGEUK-01	Gyrista	Bacillariaceae sp.	84.05	32.54	2172	20103	45.84	3.46	2.9	13603	13520	83	66.54	75.92	44.34	153	JBKBDC000000000
SGEUK-02	Gyrista	Bacillariaceae sp.	64.39	28.49	5479	5715	48.42	5.37	1.44	10889	10868	21	65.97	76.43	45.97	132	JBKBDB000000000
SGEUK-03	Chytridiomycota	Lobulomycetales sp.	56.96	12.11	1074	15102	48.57	7.71	1.7	5650	5630	20	77.38	76.58	58.48	121	JBKBDA000000000
SGEUK-04	Gyrista	Bacillariaceae sp.	35.9	19.6	11398	1790	48.43	4.04	1.55	8359	8343	16	53.08	64.77	34.08	44	JBKBCZ000000000
SGEUK-05	Haptista	Prymnesium sp.	34	59.4	9735	6756	53.36	5.12	18.63	15098	15075	23	54.65	56.78	36.01	175	JBKBCY000000000