Germline and somatic variation influence nuclear-mitochondrial crosstalk in tumourigenesis

Maëlle Daunesse[1,+], Vasavi Sundaram[1,+], Elissavet Kentepozidou[1,+], John Connelly[2,3,4,5,+], Craig J. Anderson[3,+], Liver Cancer Evolution Consortium, Núria López-Bigas[6,7,8,+], Colin A. Semple[3+], Duncan T. Odom[9,10,+], Martin S. Taylor[3,+], Maša Roller[1], Sarah J. Aitken[2,10,11,12,13,+,*], Paul Flicek[1,14,15+,*]

[1]   European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, UK
[2]   Medical Research Council Toxicology Unit, University of Cambridge, Cambridge, UK
[3]   Medical Research Council Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK
[4]   Edinburgh Pathology, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK
[5]   Laboratory Medicine, NHS Lothian, Edinburgh, UK
[6]   Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science and Technology, Barcelona, Spain
[7]   Universitat Pompeu Fabra (UPF), Barcelona, Spain
[8]   Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain
[9]   Division of Regulatory Genomics and Cancer Evolution (B270), German Cancer Research Center (DKFZ), Heidelberg, Germany
[10] Cancer Research UK - Cambridge Institute, University of Cambridge, Cambridge, UK
[11] Center of Molecular and Cellular Oncology, Yale Cancer Center, New Haven, USA
[12] Department of Pathology, Yale School of Medicine, New Haven, USA
[13] Department of Histopathology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK
[14] The Jackson Laboratory for Genomic Medicine, Farmington, USA
[15] Department of Genetics, University of Cambridge, Cambridge, UK

+   Liver Cancer Evolution Consortium

*   Corresponding authors

Abstract

Hepatocellular carcinoma is the most prevalent primary liver cancer and arises from hepatocytes, which are very metabolically active and contain abundant mitochondria. Despite extensive characterisation of nuclear driver genes and pathways, the role of crosstalk between the mitochondrial and nuclear genomes in mitochondrial mutagenesis remains poorly understood. To address this, we leveraged a chemical carcinogenesis model of liver cancer in four mouse strains. We analysed hundreds of tumours with paired whole-genome sequencing and total RNA sequencing to comprehensively characterise mitochondrial mutations and expression. Following *de novo* assembly of strain-specific mitochondrial genomes, we devised an accurate and efficient heteroplasmy detection approach and analysed mitochondrial DNA (mtDNA) content and transcription. Across strains, there was high concordance of the number of heteroplasmies, mutation signatures, and variant allele frequencies. In contrast, the heteroplasmy rate and genic locations differed between strains, suggesting strain-specificity in DNA repair mechanisms in early tumour development. Independent of age or environmental factors, tumours had lower mtDNA content than adjacent normal liver, suggesting mitochondrial loss during tumourigenesis. Additionally, within individual strains, mtDNA content differences were associated with nuclear driver gene choice. Our comprehensive characterisation of mitochondrial genomes and expression patterns demonstrates that both germline and somatic genetic variation influence tumour mtDNA content, mutation patterns, and gene expression.

Background

Hepatocellular carcinoma (HCC) is the most common type of primary liver cancer. It has the sixth highest incidence of all human cancers with five-year survival of only 25   12% (Bray et al. 2018). HCC typically occurs in the context of chronic liver disease (Llovet and Beaugrand 2003) caused by repeated damaging exposures, such as viral hepatitis infection, alcohol consumption, or metabolic dysfunction-associated steatotic liver disease (MASLD). Although there are considerable differences in HCC burden across the globe (Toh et al. 2023), how an individual's genetic variation interacts with 30   these known environmental risk factors is largely unstudied.

Cancer initiation and promotion, including that of HCC, is the product of acquired somatic mutations (Stratton et al. 2009). To identify such mutations within nuclear DNA, whole exome (Tomczak et al. 2015) and whole genome (Aaltonen et al. 2020; Zhang et al. 2019a) sequencing has been applied to large cohorts of liver cancers. 35   These efforts have led to the identification of multiple driver gene mutations in distinct oncogenic pathways involved in HCC development (Schulze et al. 2015; Zucman-Rossi et al. 2015; Fujimoto et al. 2016; Ng et al. 2021; Wu et al. 2023), as well a better understanding of somatic mosaicism in chronic liver disease (Brunner et al. 2019; Ng et al. 2021). Despite this extensive characterisation of the nuclear genome, the 40   contribution of mutations in the mitochondrial genome to liver tumourigenesis remains understudied.

Mitochondria are eukaryotic cytoplasmic organelles colloquially known as the "powerhouses of the cell". Mitochondria synthesise adenosine triphosphate (ATP) through oxidative phosphorylation to provide the energy required to support cellular 45   functions (Osellame 2012). Mammalian cells generally each have between 100 and 10,000 mitochondria depending on their function and developmental stage. Each mitochondrion contains one or more copies of its own extra-nuclear circular genome, referred to as mitochondrial DNA (mtDNA). In mice, as in humans, the mtDNA is approximately 16 kb in size and it contains 37 genes encoding two rRNAs, 22 tRNAs, 50   and 13 protein-coding genes that are components of the oxidative phosphorylation system. Mutations in mtDNA are mostly single nucleotide substitutions, insertions, or deletions (Kogelnik et al. 1996), typically occurring during mitochondrial genome replication and persisting in the mitochondrial population through repeated mitotic segregation. These mutational events can lead to multiple mtDNA variants in a cell or 55   individual: A phenomenon known as heteroplasmy.

Liver cells (hepatocytes) have very high ATP requirements to perform their multiple energy-consuming functions, including detoxification of blood and the metabolism of fat, proteins, and carbohydrates. Indeed, high-resolution imaging of mouse hepatocytes estimates the number of mitochondria to be between 1,000 and 3,000 per 60   cell (Parlakgül et al. 2024), making the liver tissue one of the most abundant sources of mtDNA. Deregulation of cellular metabolism is recognised as a core hallmark of cancer (Hanahan and Weinberg 2011; Hanahan 2022). More specifically, mitochondria are thought to play a central role in cancer development because malignant cells are characterised by increased cell proliferation, resistance to cell 65   death, and high energy consumption (Hertweck and Dasgupta 2017). Recent studies have suggested that disruptive mutations to mtDNA genes act as drivers in particular tumour types (Kim et al. 2022), and more generally, in normal tissues, both mtDNA copy number and heteroplasmy behaviour have been shown to be under nuclear genetic control in a large human genome wide association study (Gupta et al. 2023).

70　However, we still have a limited understanding of the mutagenic processes that occur in mtDNA during tumourigenesis and the functional impact of these mutations.

Given the substantial energy requirements of liver tissue and importance of metabolic alterations in HCC (Yang et al. 2023), mtDNA is a critical component of liver cancer genome analysis. Indeed, previous work has identified global heteroplasmy patterns,
75　expression changes, and other dysregulation in mtDNA associated with human liver tumourigenesis (Reznik et al. 2016; Yuan et al. 2020). Currently, the functional consequence of those heteroplasmies remains unclear; some studies showing positive selection for somatic mutations (Li et al. 2015; Grandhi et al. 2017) while others found neutral evolution (Liu et al. 2019; Yuan et al. 2020). There is also
80　evidence for decreased mitochondrial copy number (Reznik et al. 2016; Yuan et al. 2020) and decreased expression of mitochondrial genes (Reznik et al. 2017; Yuan et al. 2020) in human liver cancer compared to the tissue and cells of origin. In other tumour types, mtDNA mutation load has been identified as a potential biomarker of treatment response (Mahmood et al. 2024), as well both a positive (Gorelick et al.
85　2021) and negative (Ewing et al. 2024) overall survival in colorectal and ovarian cancer, respectively. However, these studies did not consider individuals' genetic background.

To investigate how individual genetic background affects mitochondria during tumourigenesis we used a well-characterised experimental model of liver cancer and
90　analysed nearly six hundred liver tumours spanning four genetically and evolutionary distinct mouse strains (Connor et al. 2018; Aitken et al. 2020, 2025). By controlling for both genetics and environment, we could determine the consequences of differences in genetic background on mtDNA mutations, content, and expression.

Results

95　We chemically induced liver tumours by treating 15-day-old mice with a single dose of diethylnitrosamine (DEN), and performed whole genome sequencing (WGS), total transcriptome sequencing (RNA-seq), and histopathological analysis (Connor et al. 2018; Aitken et al. 2020, 2025). For each strain, tumours were collected at a defined time point post DEN treatment and histologically confirmed as dysplastic nodules
100　(DNs). For this study, we included tumours (n=580) from inbred laboratory *Mus musculus* C3H/HeOuJ (C3H) mice (n=370 DNs) and C57BL/6J (BL6) mice (n=55 DNs); wild-derived inbred *Mus musculus castaneus* (CAST) mice (n=84 DNs); and wild-derived inbred *Mus caroli* (CAROLI) mice (n=71 DNs) (**Table S1**). The majority of the CAST genome separated from the common ancestor of BL6 and C3H
105　approximately 0.5 million years ago, although there are introgressions of the ancestral CAST genome in the BL6 and C3H genomes (Yang et al. 2011). CAROLI diverged from the common ancestor of the other three strains approximately 3 million years ago (Thybert et al. 2018). For clarity, these species, subspecies, and strains are all subsequently referred to as strains. We have recently shown that these germline
110　differences influence cancer susceptibility and somatic mutagenesis in the nuclear genome (Aitken et al. 2025). These strain-specific differences include driver gene preference within the MAPK pathway, genome stability (including whole genome duplication), and tumour latency which ranges from 25 weeks for C3H to 80 weeks for CAROLI.

De novo assembly of novel mitochondrial genomes

To ensure the robustness of our cross-strain analyses, we needed to use strain-matched mitochondrial genomes. Therefore, we assembled and annotated de novo mitochondrial genomes using our WGS data generated from matched non-tumour samples (Aitken et al. 2020, 2025) from C3H, CAST, and CAROLI (Methods) and used these mitochondrial genomes in conjunction with the annotated reference BL6 mitochondrial genome (NC_005089.1) (Bayona-Bafaluy et al. 2003). Our comparative analysis confirmed there are no structural rearrangements between these four mitochondrial genomes, they contain the same set of 37 genes (**Table S2**), and they are of comparable genomic size (C3H 16,301 bp; BL6 16,299 bp; CAST 16,299 bp; CAROLI 16,310 bp) (**Fig. 1**). Additionally, the sequence divergence between genomes reflects the evolutionary distance of the four strains (**Fig. S1a**): Specifically, closely related BL6 and C3H are 99.98% identical, whereas more divergent strains CAST and CAROLI exhibit lower sequence identity to BL6, at 97.67% and 87.21%, respectively (**Fig. S1b**).

To validate the accuracy of our *de novo* assemblies, we compared our assemblies to other available mtDNA assemblies (**Table 1**; Methods). Our genome assembly for CAST is 99.99% identical to the reference CAST mtDNA (NC_012387.1) with only two substitutions (positions 2,797 and 13,733). For CAROLI, our assembly is 98.88% identical to the reference *Mus caroli* mtDNA (NC_025268.1) and 99.99% identical to another *Mus caroli* mtDNA (MK166027.1). This difference likely arises from the fact that both our samples and those used for MK166027.1 were obtained from the same commercial provider, whereas the reference *Mus caroli* mitochondrial genome (NC_025268.1) was sequenced from a wild-caught individual (Rachal O'Neill, Johan Michaux, and Alex Greenwood personal communication). Although we were not able to identify a previous mtDNA assembly for our C3H strain (C3H/HeOuJ), comparison to a C3H/HeJ mitochondrial genome (EF108335.1) showed 99.98% identity, with only three substitutions (positions 8,898, 9,821, and 13,066). Taken together, these comparisons confirm that our *de novo* mitochondrial genome assemblies are high accuracy and consistent with published references, while also revealing minor sequence variations likely reflecting natural genetic drift.

We next examined the sequence conservation of the origin of replication for the light strand (ORI-L), an essential region for mtDNA maintenance in mammals (Wanrooij et al. 2012), which contains a highly conserved polyA sequence whose length has been associated with lifespan in mice (Hirose et al. 2018). In C3H, BL6, and CAST the ORI-L sequence is fully conserved, including an 11-adenine repeat (**Fig. S2**). However, in CAROLI, this region has evolved into a 10AG3A sequence, including a G-to-A change (**Fig. S2**). These differences at the CAROLI ORI-L locus are of particular interest given the longer lifespan and increased cancer resistance observed in CAROLI mice.

Figure 1 | Mitochondrial genome assemblies demonstrate similar genome structure, size, GC content, and sequencing coverage across strains. The C3H, BL6, CAST, and CAROLI mitochondrial genomes represented in CIRCOS plots. The outer layer of each plot represents the mitochondrial genomic position with gene annotation. The intermediate layer shows heteroplasmy variant allele frequency for somatic mutations (red) and recurrent heteroplasmies (blue) calibrated on the y-axis. The inner layer shows mitochondrial sequencing coverage for normal (external) and shift (internal) mapping (see **Fig. S3**). The genomic regions that are retained from each mapping approach are shaded blue (Methods).

Novel heteroplasmy detection workflow using whole genome sequencing

To detect heteroplasmies with high confidence across the entire mitochondrial genome, we implemented a novel heteroplasmy detection workflow, taking the circular nature of the mitochondrial genome into consideration. Our mapping approach (**Fig. S3a**) builds on a previous study (Zhang et al. 2019b), with a double alignment step (**Fig. S3b**), which reduces the detection of false positive somatic heteroplasmies, particularly those caused by the misalignment of mtDNA fragments integrated into the nuclear genome termed nuclear mitochondrial sequences (NUMTs).

First, we mapped whole genome sequencing reads to the mitochondrial genome. Uniquely mapping, valid, paired reads were then extracted and re-mapped to both the nuclear and mitochondrial genomes; only those that aligned to the mtDNA (and not to the nuclear genome) were extracted, thus ensuring that only true mitochondrial reads were considered for the next steps. These reads were then deduplicated and homoplasies were called and annotated (Methods; **Fig. S3a**), with an average sequencing depth of 5,166X at heteroplasmic loci. To account for the circularity of the mitochondrial genome, we created an alternate version of the mtDNA assembly for each of the four strains by shifting the linear starting position of the mtDNA by 8,150 bp (**Fig. S3b**). Then, the mapping approach described above was reapplied to this shifted genome, which results in more even read depth across the entire mitochondrial genome, in particular at the start and end of the mtDNA coordinate system which otherwise suffer from lower coverage due to unpaired read alignment (**Fig. 1**). A final heteroplasmy set was obtained by combining the heteroplasmies identified using the reference genome (nucleotide position 1,100 to 15,200) and the shifted genome (position 1 to 1,099, and 15,201 to the end of the genome) (**Fig. 1**; **Fig. S3b**).

Our novel approach of repeating the mapping using a shifted mtDNA reference (in addition to the classic reference) improved the coverage by an average of 1,500X in regions near the start and end of the mtDNA genome (**Fig. S5**). In addition to ensuring more uniform mapping coverage, this approach also significantly improves accuracy and specificity: In this dataset, we were able to exclude 310 false positive heteroplasmies that would otherwise be erroneously detected solely because of low coverage.

Strain-specific differences in mutational landscapes of tumour mitochondrial genomes

To identify and characterise the mitochondrial mutation landscape, we performed heteroplasmy detection on all normal (n=27) and tumour (n=580) samples for each of the four strains (Methods). Since germline selection can shape mtDNA diversity in humans (Wei et al. 2019), we first analysed the genomes of normal liver samples to identify germline variation/polymorphisms in each strain (**Fig. 1**). On average, each normal liver tissue genome had one heteroplasmy; since normal tissue is expected to be polyclonal, these heteroplasmies were considered to be germline in origin and therefore removed from the tumour set. Additionally, we identified that 49% of the heteroplasmy positions in tumours are strain-recurrent (Methods). The variant annotation of the strain-recurrent heteroplasmies revealed that 78% are synonymous and none were annotated as loss of function. Because these recurrent heteroplasmies

210 are likely to be germline or NUMT false positives, we also excluded them from downstream analysis.

After removing hypermutated tumours (n=7; Methods; **Fig. S6**), the remaining 573 tumour samples (C3H n=369; BL6 n=55; CAST n=80; CAROLI n=69) contained a total of 1,138 somatic heteroplasmies (C3H n=527; BL6 n=126; CAST n=340; CAROLI
215 n=145; **Table S4**), equating to a median of 1 to 4 (C3H median=1, BL6 and CAROLI 2, CAST 4) heteroplasmic sites per genome (**Fig. 2a**). Tumour samples are significantly more mutated than normal liver (p<0.001, Wilcoxon test). Heteroplasmies are positively correlated with nuclear somatic mutation rate in each strain (**Fig. S6**); accordingly, BL6 and CAST tumours have more mitochondrial heteroplasmies than
220 those from C3H and CAROLI, also reflecting the elevated levels of mutation in the nuclear DNA for these strains (Aitken et al. 2025). The mutational spectra signature is nearly identical across the four strains. The majority of mutations are C:G>T:A substitutions (~60%) and T:A>C:G substitutions (~30%) (**Fig. 2b**). This mutation profile does not reflect the DEN signatures that dominate the tumour nuclear genomes
225 (Aitken et al. 2025); instead, it closely matches the mtDNA signatures seen in human cancers (Lake et al. 2024; Yuan et al. 2020) which reportedly reflects mtDNA replication strand biases (Ju et al. 2014).

The majority of heteroplasmies (75%) have a low variant allele frequency (VAF; <2%), which is consistent with the accumulation of replication-coupled mutations (Ju et al.
230 2014). Given the gradual acquisition of such mutations, and the observation in humans that heteroplasmic single nucleotide variants can increase with age (Gupta et al. 2023), we considered whether this also applies in mice. We detected only a weak positive correlation between the number of heteroplasmies per sample and the age of the mouse at tumour isolation (Spearman's ρ=0.29, p<0.001; **Fig. S7a**). Similarly, VAF
235 is only very weakly correlated with age (Spearman's ρ=0.08, p<0.001; **Fig. S7b**).The absence of a stronger correlation in our cohort may arise because the experimental design imposed a fixed time point per strain (Aitken et al. 2025), which means that age is fully confounded with genetic background.

Next, we focussed specifically on mutations arising within the thirteen mitochondrial
240 protein-coding genes. We calculated sequence-composition adjusted mutation rate for each gene, which showed heterogeneity among the four strains (**Fig. 2c**). We find no significant difference between the genic distribution of heteroplasmies in C3H and BL6 (p=0.22), most of which are within the NADH dehydrogenase complex . Heteroplasmies in the other strains are less evenly distributed, with mitochondrially
245 encoded NADH dehydrogenase 6 (*Nd6*) overrepresented in CAST, and mitochondrially encoded NADH dehydrogenase 2 (*Nd2*) and mitochondrially encoded NADH dehydrogenase 4L (*Nd4l*) in CAROLI. The NADH dehydrogenase complex accounts for 39% of the mitochondrial genome, and somatic mutations have been associated with clinical outcome in diverse human cancers (Kurelac et al. 2013;
250 Koshikawa et al. 2017).

To understand the impact of these protein coding mutations we annotated their functional consequences and found a high proportion of missense mutations (~40%) in all four strains (**Fig. 2d**). This is consistent with previous work which found that, in contrast with the nuclear genome, mtDNA shows an enrichment for non-synonymous
255 variants, including in human liver cancer (Hertweck and Dasgupta 2017; Yuan et al. 2020). We found that CAST has a significantly higher proportion of heteroplasmies

with a loss of function annotation compared with the three other strains (Fisher's exact test p=0.033) (**Fig. 2d**).

260   We measured the relative rates of non-synonymous (dN) to synonymous (dS) mutations for each mitochondrial gene to infer selective pressure (**Table S3**). The resulting mtDNA-wide dN/dS scores suggest neutral evolution for missense heteroplasmies in C3H (1.0) and CAST (1.0). However, BL6 shows evidence of positive selection for missense heteroplasmies with nearly twice as many observed as expected by chance (dN/dS ratio 1.9; 95% confidence interval (CI) 1.01-3.57). BL6

265   genes with significant evidence for positive selection include *Nd2* (dN/dS ratio 4.7) and *Nd4* (dN/dS ratio 2.7). In contrast, the CAROLI mtDNA-wide score of 0.4 (95% CI 0.27-0.66) indicates higher purifying selection for mitochondrial mutations in this strain, with significant evidence for selection on *Nd2* (dN/dS ratio 0.2). We also estimated which of the non-synonymous heteroplasmies are likely to be deleterious by calculating SIFT

270   scores for each of the non-synonymous heteroplasmies (Kumar et al. 2009) (**Table S4**; Methods). For each strain, approximately three quarters of the missense heteroplasmies are predicted to be deleterious.

    In summary, the concordance of the number of heteroplasmies, mutation signatures, and VAF distribution across all tumours strongly suggests that the mutational

275   mechanisms of tumourigenesis are highly conserved across all strains in this study. However, the differences in heteroplasmy rate and genic location imply that strain-specific repair mechanisms or other differences in the dynamic process of tumourigenesis play an important role. These results illustrate the importance of background genotype on patterns of somatic mutations.

Figure 2 | Characterisation of heteroplasmies in normal liver and tumour samples. **(a)** Liver tumour (DN) genomes contain more mitochondrial heteroplasmies than normal liver tissue, which is replicated across all strains (C3H, BL6, CAST, CAROLI). P-values were calculated using a Wilcoxon test (ns = not significant, p>0.05; ** = p<0.01). **(b)** Percentage of heteroplasmies associated with each nucleotide substitution type (C>A, C>G, C>T, T>A, T>C, T>G) was calculated across all tumour samples on a per-strain basis; percentages include reverse complement mutations (e.g. T>C includes A>G). **(c)** Sequence-composition adjusted mutation rate (max normalised heteroplasmy per base pair) per gene, normalised by strain. The heatmap displays the heteroplasmy rates across mitochondrial genes, with darker shades indicating a higher mutation rate. **(d)** For each of the 1,138 somatic heteroplasmies (C3H n=527; BL6 n=126; CAST n=340; CAROLI n=145), the variant type was annotated (synonymous, missense, loss of function, tRNA, rRNA, and non-coding) and the proportions calculated per strain. Approximately 40% of variants are non-synonymous and may impact protein function.

Mitochondrial content is associated with tumour development and driver gene mutations

Having defined the mutational landscape of mtDNA mutations in DEN-induced tumours, we calculated changes in mtDNA content between normal liver and tumour sample (Methods, **Table S5**). We found that tumours have significantly lower mtDNA content than normal tissue in BL6 and CAST (**Fig. 3a**). C3H tumours show the same trend, but have too few normal samples to assess statistical significance, while CAROLI tumours demonstrate a different phenomenon described below. To determine whether the observed depletion in mitochondrial content is associated with (i) DEN-treatment, (ii) ageing (the normal samples used in **Fig. 3a** are 15 days old), or

(iii) tumourigenesis, we analysed two other BL6 sample sets. First, we collected background (non-tumour) liver tissue that was spatially distant to the DEN-induced tumours; by definition, these background tissues were also exposed to DEN when the mouse was 15 days old. Comparing paired DEN-exposed background liver to DEN-induced tumour from the same individuals (**Fig. 3b**), shows that tumours have significantly lower mtDNA content than the DEN-treated background samples ($p<0.0001$, Wilcoxon test). Furthermore, there is no significant difference in mitochondrial content between these DEN-exposed background liver samples and unexposed 15-day-old normal liver (**Fig. S7c**). This indicates that tumour development itself, rather than DEN treatment or age, is responsible for mitochondrial content loss. Second, we included a cohort of DEN-induced HCCs, which are histologically more advanced than the majority of dysplastic nodule tumours analysed in the rest of this study; this allowed us to investigate the effect of tumour stage on mtDNA content. We found that the mtDNA content of the HCC samples is comparable to the DNs (**Fig. 3b**), suggesting that while tumour development is associated with a decrease in mtDNA content, it is independent of tumour stage. These data are strikingly similar to the lower mtDNA content seen in human HCC relative to control liver tissue (Yuan et al. 2020), suggesting our murine model system recapitulates features of human disease.

A distinct feature of the CAROLI cohort is the high prevalence of whole-genome duplication (WGD) (Aitken et al. 2025), which was observed in 26 out of 69 CAROLI tumours. Analysing the mitochondrial content of these WGD tumours also revealed an approximate doubling of mtDNA content compared to diploid CAROLI tumours (**Fig. 3c**). In contrast, the small proportion of whole genome duplicated C3H tumours (11/369) do not have significantly higher mtDNA content than diploid tumours (**Fig. 3c**). The CAROLI results are consistent with a failure of karyokinesis, including preservation of both pairs of sister chromatids in the same nucleus (i.e. a single, 4n nucleus per cell) and increased cytoplasmic volume, thus maintaining the nuclear to mtDNA ratio. This hypothesis is supported by image analysis of digitised histopathology which found an approximate doubling of nuclear volume in CAROLI WGD tumours (Aitken et al. 2025), and shows strong parallels with human tumours, where WGD has been shown to evolve in concert with increased mitochondrial density (Kim et al. 2024).

Finally, we considered whether specific driver gene mutations in the nuclear genome had an impact on mtDNA content. This question is effectively intractable in human cohorts where there is often high diversity and multiple combinations of driver mutations present. In our study, the vast majority of tumours across all strains have an activating mutation in the MAPK pathway, but the genic distribution of these varies between strains. For that reason, we compared driver genes on a per strain basis and focused on tumours with one of the three most prevalent driver genes (Braf transforming gene (*Braf*), epidermal growth factor receptor (*Egfr*), and Harvey rat sarcoma virus oncogene (*Hras*)); CAROLI was omitted because there were too few diploid tumours to test statistical significance. In C3H, we find significantly higher mtDNA content in *Braf* driven tumours compared to *Hras* tumours ($p< 0.01$, Wilcoxon test) (**Fig. 3d**). By contrast, *Egfr* driven tumours had the highest mtDNA content in CAST tumours ($p<0.05$, Wilcoxon test). To our knowledge, this is the first demonstration of associations between the activation of particular driver genes and mtDNA content in tumours, though it has been established that MAPK signalling can affect many aspects of mitochondrial metabolism and biogenesis (Sarg et al. 2024).

355 In summary, mitochondrial content (i) decreases during tumourigenesis, (ii) is associated with specific driver gene mutations, and (iii) is correlated with whole genome duplication in some genetic backgrounds.

Figure 3 | Tumourigenesis alters mitochondrial content. **(a)** Across all samples,
360 tumours (DN) have less mtDNA content than normal liver samples, which is also significant in strain-specific analysis of BL6 and CAST. **(b)** Tumour (DN and HCC) genomes have significantly lower mtDNA content than adjacent, age-matched background liver tissue (DEN-exposed, non-tumour) from the same individual BL6 mouse, or normal (15-day-old, DEN-naive) liver, which indicates that it is the process
365 of tumourigenesis itself, rather than DEN treatment or tumour stage, that affects mtDNA content. **(c)** Whole genome duplicated tumours (4n) have significantly increased mtDNA content compared to diploid tumours (2n) in CAROLI, consistent with both increased nuclear and cytoplasmic volume. **(d)** Mitochondrial content is affected by an interaction of genetic background and somatic mutations: C3H tumours
370 with *Hras* mutations are significantly depleted in mitochondrial content compared with those with *Braf or Egfr* mutations, while CAST tumours with *Braf* mutations are depleted compared with *Egfr* driven tumours. For all plots p-values (p) were calculated using a Wilcoxon test (ns = p>0.05, * = p<0.05, ** = p<0.01, *** = p<0.001, **** = p<0.0001).

Tumourigenesis results in strain-specific mitochondrial expression profiles

To characterise the functional impact of mutations in the mitochondrial genome on cancer development we analysed the gene expression of all tumours with paired WGS and RNA-seq (n=378) and 52 normal liver transcriptomes (Methods). We first examined the overall expression of all thirteen mitochondrial genes (**Fig. 4a**) in normal and tumour samples across the four strains (**Table S6**). This revealed that C3H, CAST, and CAROLI have a significant loss of gene expression in tumour samples compared with normal samples, whereas BL6 tumours showed more muted changes. While the overall reduced mitochondrial gene expression is consistent with the trend for depletion of tumour mtDNA content (**Fig. 3a**), for a given tumour the correlation between mtDNA content and gene expression is relatively low for diploid tumours, with the exception of CAROLI (Spearman's ρ=0.42; p<0.01) (**Fig. S8**). WGD CAROLI tumours show stronger correlation (Spearman's ρ=0.57; p<0.01) than their diploid counterparts, which is also observed in the minority group of C3H WGD tumours (Spearman's ρ=0.43; p<0.01) (**Fig. S8**).

We also investigated the potential impact of specific driver genes on mitochondrial expression. Again, we focused our comparison on tumours that have one the three most prevalent oncogenes (*Braf*, *Egfr*, and *Hras*) and CAROLI samples were excluded due to insufficient tumour numbers. We found that CAST *Egfr* driven tumours show significantly higher mitochondrial expression than those with *Braf* drivers (**Fig. 4b**), consistent with the higher mtDNA content observed in CAST tumours (**Fig. 3d**).

To assess the mitochondrial gene expression changes in tumours, we performed a differential expression analysis of all 378 tumour RNA-seq libraries compared to normal adult liver samples (**Fig. 4c**). The C3H, CAST, and CAROLI tumours showed lower expression in tumours for most mitochondrial genes, especially those in the NADH complex, mitochondrially encoded ATP synthase 6 (*Atp6*), and mitochondrially encoded cytochrome b (*Cytb*). As with overall gene expression, above, BL6 shows a blunted pattern with fewer, and less significantly, differentially expressed genes.

Finally, we performed principal component analysis (PCA) using normalised mitochondrial expression of all normal and tumour samples for the four strains (**Fig. 4d**). Most striking is the tight clustering together of C3H, BL6, and CAST normal liver samples, with slight separation by age (**Fig. S10**); this is in contrast to the tumours which diverge between strains, but cluster within strain, partially accounting for PC1. The CAROLI samples behave differently, with normal and tumour samples clustering away from the other strains (accounting for PC2). The mitochondrial genes analysed here show a profile consistent with our transcriptome wide analysis of the same tumours (see Fig. 5a in Aitken et al. 2025 (Aitken et al. 2025)), where the first two principal components capture strain and tumour status, with CAROLI most divergent.

Here we have demonstrated that both genetic background and acquired somatic driver mutations in the nuclear genome impact mtDNA expression. The reduction in global mitochondrial gene expression in C3H, CAST, and CAROLI tumours is predominantly driven by reduced expression of the NADH complex genes, as well as *Atp6* and *Cytb*. The reduction of mtDNA expression observed in three of our four study strains has also been observed in human tumour samples, including liver cancer (Qiao et al. 2017).

Figure 4 | Tumour samples exhibit loss of mitochondrial gene expression. (a) Liver tumour (DN) samples have significantly lower mitochondrial gene expression than normal age-matched liver tissue in C3H, CAST, and CAROLI mice (Wilcoxon test, ns = p>0.05, *** = p<0.001, **** = p<0.0001; VST = variance stabilisation transformation). **(b)** In CAST mice, the expression level in tumour samples is associated with driver gene mutations, with *Egfr* mutated tumours having the highest expression levels. There was no correlation in other strains (**Fig. S9**). **(c)** Differential gene expression of mitochondrial genes was calculated between normal adult and tumour samples, per strain. As for total gene expression, the per-gene analysis highlights more marked changes in C3H, CAST, and CAROLI (Log2FC = log2 fold-change; p<0.01). Colour intensity indicates the magnitude of change. **(d)** Principal component analysis (PCA) of the VST mitochondrial expression read counts for all normal and tumour samples. Normal samples (crosses) cluster together, whereas tumour samples (circles) separate by strain.

Co-expression mtDNA genes with MAPK pathway genes across strains

We studied the functional interactions between mtDNA and nuclear genes using weighted gene correlation network analysis (WGCNA; Methods) to calculate co-expression networks for the tumours from each of the four strains. This resulted in a list of co-expressed nuclear genes for each mtDNA gene. We then performed a gene set enrichment analysis (GSEA) for each strain, ranked by the strength of overall co-expression of each mtDNA gene (**Fig. 5a**). As expected, the nuclear genes most enriched for co-expression with mtDNA genes are found in pathways related to mitochondrial and liver function; for example, oxidative phosphorylation and electron

transport chain in the OXPHOS system in mitochondria). In particular, one of the most enriched gene sets is the non-alcoholic fatty liver disease pathway, indicating that the mtDNA genes in our tumour samples are co-expressed with genes implicated in the pathogenesis of chronic liver disease.

450   We also generated co-expression networks for each of the strains (**Fig. 5b**) to identify nuclear genes that are strongly co-expressed with specific mitochondrial genes. All four strains show a strong connection between mitochondrial genes and nuclear genes found in the MAPK pathway: mitogen-activated protein kinase kinase 5 (*Map2k5*) for C3H, and mitogen-activated protein kinase associated protein 1 (*Mapkap1*) for BL6,
455   CAST and CAROLI. In addition, the latter three strains also show co-expression with phosphatidylinositol-4-phosphate 3-kinase catalytic subunit type 2 alpha (*Pik3c2a*) and cyclin dependent kinase 8 (*Cdk8*). This analysis identifies a correlation between tumour mitochondrial gene expression and genes that play a role in cancer development. Understanding these mechanisms could help identify therapeutic
460   strategies that target the mtDNA.

Figure 5 | Co-expression networks link mitochondrial genes with nuclear genes.
(**a**) Gene set enrichment analysis was performed using nuclear and mitochondrial genes identified from weighted gene co-expression network analysis (Methods). The
465   heatmap shows the enrichment for each gene, per strain, for the top five ranked pathways. Enrichment levels are indicated by the colour gradient, with darker shades representing higher enrichment. Statistically significant enrichments are marked with an asterisk (* = p<0.05). (**b**) mtDNA-centric co-expression networks of C3H, BL6, CAST, and CAROLI. The networks illustrate the co-expression connections
470   (grey/black edges) between mitochondrial genes (red nodes) and top nuclear genes (blue nodes); black edges indicating links that are at least 2-fold stronger than grey edges.

Discussion

The rising burden of hepatocellular carcinoma (HCC) reflects changing environmental
exposures and the increasing prevalence of chronic liver disease across diverse
populations. Since reprogramming of cellular metabolism is a core hallmark of cancer
(Hanahan and Weinberg 2011) that is characteristic of HCC, in this study we used a
model system of chemical carcinogenesis to induce liver tumours in four genetically
distinct mouse strains, and performed high resolution molecular mitochondrial
profiling. This approach addresses the critical need to understand the interactions
between carcinogenic exposures, mitochondrial function, and genetic variation. This
model system also enabled us to control for the genetic variability and uncertain
environmental exposures that characterise human studies.

We developed a novel, high-confidence heteroplasmy detection workflow that led to
the identification of two types of mutations. First, "strain-recurrent" heteroplasmies,
which are present in over 10% of tumour samples, appear to result from low-level
germline heteroplasmies enriched in tumour cells but lost in normal tissues, likely due
to random drift (He et al. 2010; Ju et al. 2014). Second, somatically acquired
heteroplasmies, which accumulate during tumour clonal expansion (Ju et al. 2014),
represent 51% of the identified heteroplasmies and are predominantly C>T and T>C
substitutions. This mutational spectrum aligns with those seen in human cancers (Ju
et al. 2014; Yuan et al. 2020) and likely arises from replication-coupled errors linked
to bidirectional initiation of mtDNA genome replication (Ju et al. 2014).

Tumour samples exhibit significantly higher mtDNA mutational burdens compared to
normal tissues, coupled with a striking depletion of mitochondrial content and gene
expression. These appear to be conserved features of liver tumour development,
rather than DEN treatment or age, since these differences are reproducible both within
and across strains. These results corroborate observations from human studies (Yuan
et al. 2020; Ju et al. 2014; Reznik et al. 2016; Zhang et al. 2019b), including an
association between WGD and mtDNA content (Kim et al. 2024).

We present evidence that particular driver gene variants in the nuclear genome are
linked to different levels of mitochondrial biogenesis. In addition, genome-wide co-
expression analysis identified strong correlations between mitochondrial protein-
coding genes and many nuclear genes, including an enrichment for MAPK pathway
genes underscoring the crosstalk between nuclear and mitochondrial genes in
tumourigenesis. Furthermore, these dramatic alterations to the mitochondrial genome
and expression patterns are evident even in the earliest stage of liver tumours
(dysplastic nodules) we have studied.

C3H mice, the strain most susceptible to cancer, exhibit the greatest depletion of
mitochondrial expression. In contrast, CAROLI mice, the most cancer resistant strain,
display a lower heteroplasmy burden and stronger purifying selection. The magnitude
of differential expression also broadly correlates with tumour susceptibility with C3H
showing the largest change and CAROLI the smallest. Previously, mitochondrial
content has been associated with clinical features, such as the degree of hepatic
steatosis (intracellular lipid accumulation) (Kamfar et al. 2016), and here we find that
differences in mtDNA content and expression are associated with specific nuclear
driver genes. For example, in CAST mice, *Egfr* driven tumours have the highest
mtDNA content (**Fig. 3d**), which is also reflected in the mitochondrial expression (**Fig.
4b**).

520   Although the study's design did not permit investigation of age-related mitochondrial changes, the absence of significant age effects across strains, despite varying tumour latencies, is intriguing. These findings suggest that certain mitochondrial features, such as expression changes, may represent universal responses to liver tumourigenesis, while others are strain-specific, likely influenced by evolutionary
525   distance or nuclear genotype. Such trends are also seen in the nuclear genome, where we have identified that interactions between germline variation and the environment are influential in tumour genome evolution (Aitken et al. 2025).

In summary, our study extends understanding of the mitochondrial contributions to tumourigenesis and underscores the importance of considering genetic background in
530   cancer research. The interplay between nuclear and mitochondrial genomes remains a promising avenue for future investigation, with potential for identifying novel therapeutic targets.

Methods

535   ### Tumourigenesis experiments

Liver tumours were induced, isolated, and processed as described previously (Connor et al. 2018; Aitken et al. 2020, 2025). Briefly, 15-day-old male mice were treated with a single intraperitoneal dose of diethylnitrosamine (DEN), and tumours isolated at fixed time points after DEN treatment (C3H 25 weeks, BL6 36 weeks, CAST 38 weeks,
540   CAROLI 78 weeks). Tumours bisected, with one half for histopathology and the other half processed for DNA and RNA extraction and whole genome (WGS) and total transcriptome sequencing (RNA-seq). Additional normal liver samples were processed for control experiments (**Table S1**).

Whole genome sequencing of normal liver and liver tumours are available in the
545   European Nucleotide Archive (ENA) under accession PRJEB37808 (C3H and CAST) (Aitken et al. 2020) and PRJEB15138 (BL6 and CAROLI) (Aitken et al. 2025). Total transcriptome sequencing of liver tissue and tumours is available at ArrayExpress (accession pending) (Aitken et al. 2025). Samples and associated metadata are listed in **Table S1**.

550   ### Novel *de novo* mitochondrial genome assembly and annotation for C3H, CAST, and CAROLI mice

The strains (reference genome assembly) used in this study are *Mus musculus musculus* C57BL/6J (GRCm39), *Mus musculus musculus* C3H/HeOuJ (C3H/HeJ), *Mus musculus castaneus* CAST/EiJ (CAST/EiJ) and *Mus caroli* CAROLI/EiJ
555   (CAROLI_EiJ) from Ensembl v.91 (Howe et al. 2021).

BL6 was analysed using the NC_005089.1 C57BL/6J reference mitochondrial genome accessed from NCBI. The C3H, CAST and CAROLI mitochondrial genomes were assembled *de novo* using NOVOplasty v. 3.8.1 (Dierckxsens et al. 2017, 2020) and have been deposited in GenBank (PV177089, PV231059, PV231060). For this, WGS
560   of all normal liver tissue samples were used for each strain (C3H=4; CAST =7; CAROLI=5).

For C3H and CAST, NOVOplasty was run on all the WGS libraries with default parameters using the *Cytb* gene sequence of NC_005089.1 as seed. CAROLI exhibits evolutionary divergence of *Cytb* gene and thus required a two-step process: first, the NC_005089.1 *Cytb* gene was used a as seed to create a preliminary assembly from which the longest conserved sequence between the preliminary assembly and NC_005089.1 was used as a seed for NOVOplasty to create the final assembly. All libraries for all samples returned circularised assemblies. For all strains, the assemblies for the various libraries were essentially identical (assemblies from two C3H libraries had a Y at one position that was an A in all others) and used to create strain-specific consensus circularised reference mitochondrial genomes assemblies for each strain. Gene annotation of the three mitochondrial genomes were performed using MITOS (Bernt et al. 2013). Start and stop positions for all genes were manually curated. The genes identified for the three strains are identical and matched BL6 gene annotation (**Fig. 1**).

To compare sequence similarity of the four mitochondrial genomes, multiple sequence alignment was performed using Clustal Omega (Madeira et al. 2024).

See **Table S2** for the final assembly and annotation.

Comparative mitochondrial genome analysis

For each strain, we aligned our *de novo* assembly and the corresponding sequences retrieved from GenBank using Clustal Omega with "sequence type" set to DNA. A summary of all sequences included in the comparative analysis is provided in **Table 1**.

Strain	GenBank ID	Length (bp)	Strain information
C3H	**PV177089.1**	**16,301**	***Mus musculus* strain C3H/HeOuJ (this study)**
C3H	EF108335.1	16,301	*Mus musculus* strain C3H/HeJ
CAST	**PV231059.1**	**16,299**	***Mus musculus castaneus* strain CAST/EiJ (this study)**
CAST	EF108342.1	16,299	*Mus musculus castaneus* strain CAST/EiJ Curated RefSeq NC_012387.1
CAROLI	**PV231060.1**	**16,310**	***Mus caroli* strain CAROLI/EiJ (this study)**
CAROLI	KJ530558.1	16,313	*Mus caroli* isolate R5231 mitochondrion Curated RefSeq NC_025268.1
CAROLI	MK166027.1	16,301	*Mus caroli*

Table 1 | **Mitochondrial genomes included in the comparative analysis**.

Heteroplasmy detection and annotation

We mapped the trimmed WGS libraries using the bwa-mem function from BWA (v.0.7.17) (Li and Durbin 2009), using the mitochondrial genomes as reference. Valid mapped pairs to the mitochondrial genome were then extracted using the view function from Samtools (v.1.9) (Li et al. 2009). The remaining reads were then re-mapped to the full genome (nuclear and mitochondrial) using bwa-mem. Valid mapped pairs to

the mitochondrial genome were extracted a second time. This approach allows potential NUclear Mitochondrial Sequences (NUMTs) to be filtered out, which otherwise tend to misalign to the mitochondrial genomes because of their high identity levels with the mitochondrial genome (Zhang et al. 2019b). Duplicates were removed using the MarkDuplicates function from Picard Tools (v.2.21.7) ([CSL STYLE ERROR: reference with no printed form.]).

For BL6, GRCm39 was used as the nuclear genome assembly to address a BL6 specific issue caused by a problematic NUMTs (Numts_002 in (Calabrese et al. 2012)) sequence located on 1: 24,650,616-24,655,269 (4,654bp long) that is 99.94% identical to MT:6,390-11,042 (4,653bp long). Although this region was improved in GRCm39, optimal mapping quality on the BL6 mitochondrial assembly required masking of the region on the nuclear genome. The masked sequence is within a non-coding region of the BL6 nuclear genome and is not conserved in the other strains. We are confident that masking this sequence did not have an impact on the heteroplasmy detection quality.

The mpileup function from Samtools was then used to generate pileup files from the BAM files with "-E -f" parameters to optimise the accuracy of the heteroplasmy detection (Wang et al. 2020). Heteroplasmy were detected using the mpileup2snp function of VarScan (v.2.4.4) (Koboldt et al. 2012). Finally, heteroplasmies were classified as synonymous, missense, loss of function, tRNA, rRNA, and non-coding using snpEff (v.5.1.2) using default parameters (Cingolani et al. 2012).

The mitochondrial genome is circular and must be arbitrarily cut to meet the input requirements of mapping alignment software. To mitigate biases caused by this linearisation (apparent low read coverage at the "ends"), we created a shifted version of each of the mitochondrial genomes with the starting position shifted by 8,150bp (**Fig. 1, Fig. S3b**). We re-applied the mapping approach described above using the shifted version of the mitochondrial genomes. The final mutation set was obtained by taking the mutation from the reference genome mapping from position MT:1,100-15,200 and the mutation from the shifted genome mapping from position MT:1-1,099 and MT:15,201-end of the genome. Re-mapping the libraries on a shifted mitochondrial genome ensures a deep coverage at every position of the mitochondrial genome.

A total of 3,382 heteroplasmies were initially detected, corresponding to 1,717 unique mitochondrial genomic positions.

To account for hypermutated samples (**Table S1, Fig. S6**), we first excluded any sample containing more than 50 heteroplasmies, which led to the removal of three samples (two from CAST and one from C3H). To further refine the filtering, we excluded samples with a heteroplasmy count exceeding three standard deviations from the strain-specific mean, resulting in the additional removal of four samples (two from CAST and two from CAROLI). After this filtering step, 2,232 heteroplasmies remained, covering 1,011 mitochondrial genomic positions (**Fig. 1**).

We further classified the remaining heteroplasmies into two categories: strain-recurrent and somatic. A heteroplasmy was defined as "strain-recurrent" if it was present in more than 10% of the strain's samples (or in more than 10 samples for C3H). After filtering out strain-recurrent heteroplasmies, 1,138 heteroplasmies

remained, corresponding to 975 unique mitochondrial genomic positions. See **Table S4** for the list of all identified heteroplasmies and associated annotations.

See GitLab (https://gitlab.com/maelle.daunesse/lce-mtdna.git) for complete snakemake pipeline.

Heteroplasmy analysis

Sequence-composition adjusted mutation rates were calculated as defined previously (Aitken et al. 2020) with modification to consider the mitochondrial genome:

$$\sum_{i}^{64} \left(\frac{n_{het\_i}}{n_i} \times \frac{n_{het\_i\_mtDNA}}{n_{i\_mtDNA}} \right)$$

Where n_{het_i} corresponds to the number of the trinucleotide i with a heteroplasmy in the gene of interest, n_i corresponds to the number of the trinucleotide i in the gene of interest, $n_{het\_i\_mtDNA}$ corresponds to the number of the trinucleotide i with a heteroplasmy in the mitochondrial genome and n_{i_mtDNA} corresponds to the number of the trinucleotide i in the mitochondrial genome.

dN/dS ratios were calculated for each gene using the dNdScv R package (v1.0) (Martincorena et al. 2017), which estimates the ratio of nonsynonymous to synonymous substitution rates while accounting for mutation rate variation and sequence composition biases. dNdScv employs a trinucleotide mutation model to correct for context-dependent mutational effects and was run using the "Vertebrate Mitochondrial Code" (`numcode = 2`) to appropriately model mitochondrial coding sequences. dN/dS scores are listed in **Table S3**.

SIFT scores were calculated for each missense annotated heteroplasmy using SIFT (v.6.2.1) (Kumar et al. 2009), with alignments calculated by NCBI BLAST+ (v.2.4.0) (Altschul et al. 1997) and UniRef90 (release 2025_01) as the reference protein database. SIFT scores are listed in **Table S4**.

See GitLab (https://gitlab.com/maelle.daunesse/lce-mtdna.git) for complete Rmarkdown notebooks.

Mitochondrial content calculation

We mapped the trimmed reads to the full genome (mtDNA + nuclear DNA) and duplicates were removed using the MarkDuplicates function from Picard. Finally, we used the idxstat function from Samtools to obtain the number of reads mapped to each chromosome.

Mitochondrial content was estimated using deduplicated libraries by calculating the fraction of mitochondrial reads in each of the WGS libraries (**Table S5**). We applied a formula that accounts for both cellularity and ploidy, as adapted from Reznik et al. 2016 (Reznik et al. 2016), which originally used a tumour purity metric (here replaced with cellularity). The cellularity of a tumour corresponds to the estimated fraction of sequenced cells that represent the dominant clonal expansion in the tumour (between 0 and 1); for normal samples, the cellularity was imputed as 1).

The mitochondrial content was calculated as follows:

675 $$mtDNA\ content\ = \frac{mtDNA\ reads}{nuclear\ reads} \times \frac{(Cellularity\ \times\ Ploidy)\ +\ (1\ -\ Cellularity) \times 2}{2}$$

This adjustment ensures the mitochondrial content reflects the cellular context, accounting for tumour heterogeneity and variations in ploidy.

Mitochondrial gene expression analysis

680 We mapped the trimmed RNA-seq libraries generated from normal liver tissue (15-day old and adult) and tumour samples using STAR (v.2.7) (Dobin et al. 2013), using the full genomes described above (nuclear and mitochondrial) as a reference. Read counts per gene were obtained using the featurecounts function from Subread (v.2.0.3) (Liao et al. 2014). Only the 14,779 genes that are orthologous in the four strains were kept for the next steps (**Table S6**).

685 Read counts from each library were then normalised using DEseq2 (v.1.38.2) (Love et al. 2014). We performed differential expression using normal adult and DN tumour samples with DESeq2 accounting for the strain, diagnosis, and cellularity in the design (**Table S6**). The PCA shown in **Fig. 4d** was obtained after variance stabilising transformations (VST). The 15-day old and adult normal samples cluster together (**Fig.**
690 **S10)**.

Co-expression of mitochondrial genes analysis

We performed a co-expression analysis using the WGCNA R package (v.1.71) (Langfelder and Horvath 2008). For each strain, a weighted gene co-expression network was built using both mitochondrial and nuclear genes using a soft thresholding
695 power (β) of 5. The resulting network assigns each pair of genes a weight ranging from 0 (low) to 1 (high) that corresponds to the connection strength between the two genes.

To determine the pathways with which the mitochondrial genes are co-expressed, we used a gene-set enrichment analysis (GSEA) approach. The GSEA analysis was performed using the co-expression weights as input to rank the genes, employing the
700 fgsea R package (v.1.24.0) (Korotkevich et al. 2021). Finally, we built a mtDNA-centric network using the 100 most connected nuclear genes to all the mitochondrial ones.

References

Aaltonen LA, Abascal F, Abeshouse A, Aburatani H, Adams DJ, Agrawal N, Ahn KS, Ahn S-M, Aikata H, Akbani R, et al. 2020. Pan-cancer analysis of whole
705 genomes. *Nature* **578**: 82–93.

Aitken SJ, Anderson CJ, Connor F, Pich O, Sundaram V, Feig C, Rayner TF, Lukk M, Aitken S, Luft J, et al. 2020. Pervasive lesion segregation shapes cancer genome evolution. *Nature* **583**: 265–270.

Aitken SJ, Connor F, Feig C, Rayner TF, Lukk M, Luft J, Aitken S, Arnedo-Pac C,
710 Hayes JF, Nicholson MD, et al. 2025. Genetic background sets the trajectory of cancer evolution. 2025.01.13.632787. https://www.biorxiv.org/content/10.1101/2025.01.13.632787v1 (Accessed April 24, 2025).

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997.
715    Gapped BLAST and PSI-BLAST: a new generation of protein database
    search programs. *Nucleic Acids Res* **25**: 3389–3402.

Bayona-Bafaluy MP, Acín-Pérez R, Mullikin JC, Park JS, Moreno-Loshuertos R, Hu
    P, Pérez-Martos A, Fernández-Silva P, Bai Y, Enríquez JA. 2003. Revisiting
    the mouse mitochondrial DNA sequence. *Nucleic Acids Research* **31**: 5349–
720    5355.

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J,
    Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan
    mitochondrial genome annotation. *Mol Phylogenet Evol* **69**: 313–319.

Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. 2018. Global
725    cancer statistics 2018: GLOBOCAN estimates of incidence and mortality
    worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for
    Clinicians* **68**: 394–424.

Brunner SF, Roberts ND, Wylie LA, Moore L, Aitken SJ, Davies SE, Sanders MA,
    Ellis P, Alder C, Hooks Y, et al. 2019. Somatic mutations and clonal dynamics
730    in healthy and cirrhotic human liver. *Nature* **574**: 538–542.

Calabrese FM, Simone D, Attimonelli M. 2012. Primates and mouse NumtS in the
    UCSC Genome Browser. *BMC Bioinformatics* **13 Suppl 4**: S15.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden
    DM. 2012. A program for annotating and predicting the effects of single
735    nucleotide polymorphisms, SnpEff. *Fly (Austin)* **6**: 80–92.

Connor F, Rayner TF, Aitken SJ, Feig C, Lukk M, Santoyo-Lopez J, Odom DT. 2018.
    Mutational landscape of a chemically-induced mouse model of liver cancer. *J
    Hepatol* **69**: 840–850.

Dierckxsens N, Mardulyn P, Smits G. 2017. NOVOPlasty: de novo assembly of
740    organelle genomes from whole genome data. *Nucleic Acids Res* **45**: e18.

Dierckxsens N, Mardulyn P, Smits G. 2020. Unraveling heteroplasmy patterns with
    NOVOPlasty. *NAR Genomics and Bioinformatics* **2**: lqz011.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson
    M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner.
745    *Bioinformatics* **29**: 15–21.

Ewing A, Meynert A, Silk R, Aitken S, Bendixsen DP, Churchman M, Brown SL,
    Hamdan A, Mattocks J, Grimes GR, et al. 2024. Divergent trajectories to
    structural diversity impact patient survival in high grade serous ovarian
    cancer. 2024.01.12.575376.
750    https://www.biorxiv.org/content/10.1101/2024.01.12.575376v1 (Accessed
    December 17, 2024).

Fujimoto A, Furuta M, Totoki Y, Tsunoda T, Kato M, Shiraishi Y, Tanaka H,
    Taniguchi H, Kawakami Y, Ueno M, et al. 2016. Whole-genome mutational

755   landscape and characterization of noncoding and structural mutations in liver cancer. *Nat Genet* **48**: 500–509.

Gorelick AN, Kim M, Chatila WK, La K, Hakimi AA, Berger MF, Taylor BS, Gammage PA, Reznik E. 2021. Respiratory complex and tissue lineage drive recurrent mutations in tumour mtDNA. *Nat Metab* **3**: 558–570.

Grandhi S, Bosworth C, Maddox W, Sensiba C, Akhavanfard S, Ni Y, LaFramboise
760   T. 2017. Heteroplasmic shifts in tumor mitochondrial genomes reveal tissue-specific signals of relaxed and positive selection. *Hum Mol Genet* **26**: 2912–2922.

Gupta R, Kanai M, Durham TJ, Tsuo K, McCoy JG, Kotrys AV, Zhou W, Chinnery PF, Karczewski KJ, Calvo SE, et al. 2023. Nuclear genetic control of mtDNA
765   copy number and heteroplasmy in humans. *Nature* **620**: 839–848.

Hanahan D. 2022. Hallmarks of Cancer: New Dimensions. *Cancer Discov* **12**: 31–46.

Hanahan D, Weinberg RA. 2011. Hallmarks of Cancer: The Next Generation. *Cell* **144**: 646–674.

He Y, Wu J, Dressman DC, Iacobuzio-Donahue C, Markowitz SD, Velculescu VE,
770   Diaz Jr LA, Kinzler KW, Vogelstein B, Papadopoulos N. 2010. Heteroplasmic mitochondrial DNA mutations in normal and tumour cells. *Nature* **464**: 610–614.

Hertweck KL, Dasgupta S. 2017. The Landscape of mtDNA Modifications in Cancer: A Tale of Two Cities. *Front Oncol* **7**: 262.

775   Hirose M, Schilf P, Gupta Y, Zarse K, Künstner A, Fähnrich A, Busch H, Yin J, Wright MN, Ziegler A, et al. 2018. Low-level mitochondrial heteroplasmy modulates DNA replication, glucose metabolism and lifespan in mice. *Sci Rep* **8**: 5872.

Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM,
780   Azov AG, Bennett R, Bhai J, et al. 2021. Ensembl 2021. *Nucleic Acids Research* **49**: D884–D891.

Ju YS, Alexandrov LB, Gerstung M, Martincorena I, Nik-Zainal S, Ramakrishna M, Davies HR, Papaemmanuil E, Gundem G, Shlien A, et al. 2014. Origins and functional consequences of somatic mitochondrial DNA mutations in human
785   cancer. *Elife* **3**.

Kamfar S, Alavian SM, Houshmand M, Yadegarazari R, Seifi Zarei B, Khalaj A, Shabab N, Saidijam M. 2016. Liver Mitochondrial DNA Copy Number and Deletion Levels May Contribute to Nonalcoholic Fatty Liver Disease Susceptibility. *Hepat Mon* **16**: e40774.

790   Kim M, Gorelick AN, Vàzquez-García I, Williams MJ, Salehi S, Shi H, Weiner AC, Ceglia N, Funnell T, Park T, et al. 2024. Single-cell mtDNA dynamics in tumors is driven by coregulation of nuclear and mitochondrial genomes. *Nat Genet* **56**: 889–899.

795   Kim M, Mahmood M, Reznik E, Gammage PA. 2022. Mitochondrial DNA is a major source of driver mutations in cancer. *Trends Cancer* **8**: 1046–1059.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* **22**: 568–576.

800   Kogelnik AM, Lott MT, Brown MD, Navathe SB, Wallace DC. 1996. MITOMAP: a human mitochondrial genome database. *Nucleic Acids Res* **24**: 177–179.

Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. 2021. Fast gene set enrichment analysis. 060012. https://www.biorxiv.org/content/10.1101/060012v3 (Accessed June 16, 2024).

805   Koshikawa N, Akimoto M, Hayashi J-I, Nagase H, Takenaga K. 2017. Association of predicted pathogenic mutations in mitochondrial ND genes with distant metastasis in NSCLC and colon cancer. *Sci Rep* **7**: 15535.

Kumar P, Henikoff S, Ng PC. 2009. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* **4**: 1073–
810   1081.

Kurelac I, MacKay A, Lambros MBK, Di Cesare E, Cenacchi G, Ceccarelli C, Morra I, Melcarne A, Morandi L, Calabrese FM, et al. 2013. Somatic complex I disruptive mitochondrial DNA mutations are modifiers of tumorigenesis that correlate with low genomic instability in pituitary adenomas. *Hum Mol Genet*
815   **22**: 226–238.

Lake NJ, Ma K, Liu W, Battle SL, Laricchia KM, Tiao G, Puiu D, Ng KK, Cohen J, Compton AG, et al. 2024. Quantifying constraint in the human mitochondrial genome. *Nature* **635**: 390–397.

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation
820   network analysis. *BMC Bioinformatics* **9**: 559.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The
825   Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li M, Schröder R, Ni S, Madea B, Stoneking M. 2015. Extensive tissue-related and allele-related mtDNA heteroplasmy suggests positive selection for somatic mutations. *Proc Natl Acad Sci USA* **112**: 2491–2496.

830   Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–930.

Liu Q, Lin D, Li M, Gu Z, Zhao Y. 2019. Evidence of Neutral Evolution of Mitochondrial DNA in Human Hepatocellular Carcinoma. *Genome Biol Evol* **11**: 2909–2916.

Llovet JM, Beaugrand M. 2003. Hepatocellular carcinoma: present status and future prospects. *J Hepatol* **38 Suppl 1**: S136-149.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. *Nucleic Acids Research* **52**: W521–W525.

Mahmood M, Liu EM, Shergold AL, Tolla E, Tait-Mulder J, Huerta-Uribe A, Shokry E, Young AL, Lilla S, Kim M, et al. 2024. Mitochondrial DNA mutations drive aerobic glycolysis to enhance checkpoint blockade response in melanoma. *Nat Cancer* **5**: 659–672.

Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Van Loo P, Davies H, Stratton MR, Campbell PJ. 2017. Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* **171**: 1029-1041.e21.

Ng SWK, Rouhani FJ, Brunner SF, Brzozowska N, Aitken SJ, Yang M, Abascal F, Moore L, Nikitopoulou E, Chappell L, et al. 2021. Convergent somatic mutations in metabolism genes in chronic liver disease. *Nature* **598**: 473–478.

Osellame LD. 2012. Cellular and molecular mechanisms of mitochondrial function. *Best Pract Res Clin Endocrinol Metab* **26**: 711–723.

Parlakgül G, Pang S, Artico LL, Min N, Cagampan E, Villa R, Goncalves RLS, Lee GY, Xu CS, Hotamışlıgil GS, et al. 2024. Spatial mapping of hepatic ER and mitochondria architecture reveals zonated remodeling in fasting and obesity. *Nat Commun* **15**: 3982.

Qiao L, Ru G, Mao Z, Wang C, Nie Z, Li Q, Huang-yang Y, Zhu L, Liang X, Yu J, et al. 2017. Mitochondrial DNA depletion, mitochondrial mutations and high TFAM expression in hepatocellular carcinoma. *Oncotarget* **8**: 84373–84383.

Reznik E, Miller ML, Şenbabaoğlu Y, Riaz N, Sarungbam J, Tickoo SK, Al-Ahmadie HA, Lee W, Seshan VE, Hakimi AA, et al. 2016. Mitochondrial DNA copy number variation across human cancers ed. C.V. Dang. *eLife* **5**: e10769.

Reznik E, Wang Q, La K, Schultz N, Sander C. 2017. Mitochondrial respiratory gene expression is suppressed in many cancers ed. C.V. Dang. *eLife* **6**: e21592.

Sarg NH, Zaher DM, Abu Jayab NN, Mostafa SH, Ismail HH, Omar HA. 2024. The interplay of p38 MAPK signaling and mitochondrial metabolism, a dynamic target in cancer and pathological contexts. *Biochem Pharmacol* **225**: 116307.

Schulze K, Imbeaud S, Letouzé E, Alexandrov LB, Calderaro J, Rebouissou S, Couchy G, Meiller C, Shinde J, Soysouvanh F, et al. 2015. Exome

sequencing of hepatocellular carcinomas identifies new mutational signatures and potential therapeutic targets. *Nat Genet* **47**: 505–511.

875 Stratton MR, Campbell PJ, Futreal PA. 2009. The cancer genome. *Nature* **458**: 719–724.

Thybert D, Roller M, Navarro FCP, Fiddes I, Streeter I, Feig C, Martin-Galvez D, Kolmogorov M, Janoušek V, Akanni W, et al. 2018. Repeat associated mechanisms of genome evolution and function revealed by the Mus caroli and Mus pahari genomes. *Genome Res* **28**: 448–459.

880 Toh MR, Wong EYT, Wong SH, Ng AWT, Loo L-H, Chow PK-H, Ngeow J. 2023. Global Epidemiology and Genetics of Hepatocellular Carcinoma. *Gastroenterology* **164**: 766–782.

Tomczak K, Czerwińska P, Wiznerowicz M. 2015. The Cancer Genome Atlas (TCGA): an immeasurable source of knowledge. *Contemp Oncol (Pozn)* **19**: 885 A68-77.

Wang D, Xiang H, Ning C, Liu H, Liu J-F, Zhao X. 2020. Mitochondrial DNA enrichment reduced NUMT contamination in porcine NGS analyses. *Brief Bioinform* **21**: 1368–1377.

Wanrooij S, Miralles Fusté J, Stewart JB, Wanrooij PH, Samuelsson T, Larsson N-G, 890 Gustafsson CM, Falkenberg M. 2012. In vivo mutagenesis reveals that OriL is essential for mitochondrial DNA replication. *EMBO Rep* **13**: 1130–1137.

Wei W, Tuna S, Keogh MJ, Smith KR, Aitman TJ, Beales PL, Bennett DL, Gale DP, Bitner-Glindzicz MAK, Black GC, et al. 2019. Germline selection shapes human mitochondrial DNA diversity. *Science* **364**: eaau6520.

895 Wu H, Eckhardt CM, Baccarelli AA. 2023. Molecular mechanisms of environmental exposures and human disease. *Nat Rev Genet* **24**: 332–344.

Yang F, Hilakivi-Clarke L, Shaha A, Wang Y, Wang X, Deng Y, Lai J, Kang N. 2023. Metabolic reprogramming and its clinical implication for liver cancer. *Hepatology* **78**: 1602–1624.

900 Yang H, Wang JR, Didion JP, Buus RJ, Bell TA, Welsh CE, Bonhomme F, Yu AH-T, Nachman MW, Pialek J, et al. 2011. Subspecific origin and haplotype diversity in the laboratory mouse. *Nat Genet* **43**: 648–655.

Yuan Y, Ju YS, Kim Y, Li J, Wang Y, Yoon CJ, Yang Y, Martincorena I, Creighton CJ, Weinstein JN, et al. 2020. Comprehensive molecular characterization of 905 mitochondrial genomes in human cancers. *Nat Genet* **52**: 342–352.

Zhang J, Bajari R, Andric D, Gerthoffert F, Lepsa A, Nahal-Bose H, Stein LD, Ferretti V. 2019a. The International Cancer Genome Consortium Data Portal. *Nat Biotechnol* **37**: 367–369.

Zhang R, Nakahira K, Choi AMK, Gu Z. 2019b. Heteroplasmy concordance between 910 mitochondrial DNA and RNA. *Sci Rep* **9**: 12942.

Zucman-Rossi J, Villanueva A, Nault J-C, Llovet JM. 2015. Genetic Landscape and Biomarkers of Hepatocellular Carcinoma. *Gastroenterology* **149**: 1226-1239.e4.

Picard Tools - By Broad Institute. https://broadinstitute.github.io/picard/ (Accessed December 19, 2024).

Computational analysis environment and code availability

The analysis pipeline including snakemake pipelines to process the data, subsequent scripts and R notebook to generate the figures can be obtained without restriction from the repository https://gitlab.com/maelle.daunesse/lce-mtdna.git.

920 ## Data availability

Sequencing data in this study are described in Aitken et al. 2020 (Aitken et al. 2020) and Aitken et al. 2025 (Aitken et al. 2025). Whole genome sequencing of normal tissues and and liver tumours are available in the European Nucleotide Archive (ENA) under accession PRJEB37808 (Aitken et al. 2020) and PRJEB15138 (Aitken et al. 925 2020, 2025). Total transcriptome sequencing of liver tissue and tumours is available at ArrayExpress (accession pending) (Aitken et al. 2025). Whole genome sequencing alignment for reference and shift mapping of the mitochondrial genomes of liver tissue and tumours are available in NCBI under accession PRJNA1225162. The mitochondrial genomes assembled for C3H, CAST, and CAROLI in this study have 930 been deposited in GenBank (PV177089, PV231059, PV231060).

Ethics

Animal experimentation was carried out in accordance with the Animals (Scientific Procedures) Act 1986 (United Kingdom) and with the approval of the Cancer Research UK Cambridge Institute Animal Welfare and Ethical Review Body (AWERB). Full 935 details of the experimental procedures leading to the data generation are provided in Aitken et al. 2020 (Aitken et al. 2020) and Aitken et al. 2025 (Aitken et al. 2025).

Acknowledgements and funding

We thank Andrew Yates (EMBL-EBI) for assistance with data management, and members of the Aitken Lab (H. Ahmed Nur, C. Arnedo-Pac, L. Cocker, E. Reilly) for 940 comments on the manuscript.

This work was supported by European Molecular Biology Laboratory core funding, Cancer Research UK Cambridge Institute core award (20412), MRC Human Genetics Unit core funding programme grants (MC_UU_00007/11, MC_UU_00007/16, MC_UU_00035/1, MC_UU_00035/2), and MRC Toxicology Unit core funding 945 programme grants (RG94521 and MC_PC_24012). IRB Barcelona is a recipient of a Severo Ochoa Centre of Excellence Award from the Spanish Ministry of Science, Innovation and Universities (MICINN, Government of Spain) and is supported by CERCA (Generalitat de Catalunya). Support was also provided from specific research grants: Cancer Research UK strategic award (22398); Wellcome Trust 950 (WT108749/Z/15/Z, WT202878/B/16/Z, 202878/Z/16/Z); PID2021-126568OB-I00 (CHEMOHEALTH) project, funded by the Spanish Ministry of Science (MCIN), AEI /10.13039/501100011033; ERC (615584, 788937); Helmholtz Society (DKFZ abteiling B270). J.C. was supported by a Wellcome Trust PhD Training Fellowship for Clinicians

955 (WT223088/Z/21/Z) as part of the Edinburgh Clinical Academic Track (ECAT) programme. M.D. and E.K. were supported by the EMBL International PhD Programme. V.S. received an EIPOD fellowship. S.J.A. received a Wellcome Trust PhD Training Fellowship for Clinicians (WT106563/Z/14/Z and WT106563/Z/14/A), National Institute for Health Research (NIHR) Clinical Lectureship, and CRUK Clinician Scientist Fellowship (RCCCSF-May23/100001).

Author contributions

960 M.D., S.J.A., and P.F. conceptualised the project. D.T.O. supervised experimental data collection. M.D. and P.F. performed primary genomic and sequence data analysis. V.S., E.K. J.C., and C.J.A. provided computational support. N.L.B., C.S., M.S.T., and M.R. advised on specific analysis components. S.J.A. and P.F. supervised 965 the work. M.D., S.J.A., and P.F. wrote the paper. All authors had the opportunity to edit the manuscript. All authors approved the final manuscript.

The Liver Cancer Evolution Consortium members are:

Sarah J. Aitken, Stuart Aitken, Craig J. Anderson, Claudia Arnedo-Pac, John Connelly, Frances Connor, Maëlle Daunesse, Ruben M. Drews, Ailith Ewing, Christine Feig, 970 Paul Flicek, Paul A. Ginno, Vera B. Kaiser, Elissavet Kentepozidou, Erika López-Arribillaga, Núria López-Bigas, Juliet Luft, Margus Lukk, Duncan T. Odom, Oriol Pich, Tim F. Rayner, Colin A. Semple, Inés Sentís, Vasavi Sundaram, Lana Talmane, Martin S. Taylor & Jan C. Verberg

Data and materials

975 Correspondence, and data and materials requests should be addressed to: Sarah J. Aitken sarah.aitken@yale.edu and Paul Flicek pf273@cam.ac.uk

Conflict of interest

J.C. has received honorarium from Roche Diagnostics. S.J.A. has received funding from AstraZeneca for a PhD studentship. P.F. is a member of the Scientific Advisory 980 Board of Fabric Genomics, Inc.