Stimulation of the *Caulobacter crescentus* surface sensing pathway by deletion of a specialized minor pilin-like gene

Farah Obeid Charrouf[1], Gregory B Whitfield[1], Courtney K Ellison[2], Yves V Brun[1*].

1: Département de Microbiologie, Infectiologie et Immunologie, Université de Montréal, 2900 Boulevard Édouard-Montpetit, Montréal, Québec H3T 1J4, Canada.

2: Department of Microbiology, University of Georgia, Athens, GA, USA.

*: corresponding author: yves.brun@umontreal.ca

Abstract:

Bacteria colonize surfaces through complex mechanisms of surface sensing. Pili are dynamic bacterial appendages that play an important role in this process. In *Caulobacter crescentus*, tension on retracting, surface-bound pili triggers the rapid synthesis of the adhesive holdfast, which permanently attaches cells to surfaces. However, the detailed mechanisms of pilus-mediated surface sensing are unclear. In this study, we used a genetic screen to isolate mutants with altered pilus activity to identify genes that may be involved in pilus-mediated surface-sensing. This screen identified *cpaL*, whose deletion led to reduced piliation levels, and surprisingly, a threefold increase in surface adhesion due to increased holdfast production. To understand this finding, we compared holdfast synthesis in wild-type and *cpaL* mutant cells under conditions that block pilus retraction. While this treatment increased holdfast production in wild-type cells by triggering the surface-sensing pathway, no increase was observed in the *cpaL* mutant, suggesting that mutation of *cpaL* maximally stimulates surface-sensing. Furthermore, when the *cpaL* mutant was grown in a medium that blocks the surface sensing pathway, cells exhibited decreased surface attachment and holdfast production, consistent with a role for CpaL in pilus-dependent surface sensing in *C. crescentus*. To better understand the function of CpaL, we analyzed its predicted structure, which suggested that CpaL is a minor pilin fused to a mechanosensitive von Willebrand factor type A (vWA) domain that could be accommodated at the pilus tip. These results collectively position CpaL as a strong candidate for a mechanosensory element in pilus-mediated surface sensing.

Importance:

Surface sensing is a crucial mechanism that allows bacteria to change their behaviors to adapt to life on a surface. Surface recognition by bacteria is the initial step toward surface colonization and biofilm formation. In *Caulobacter crescentus*, tight adherence (Tad) pili play a key role in surface recognition and adaptation. However, the mechanism of pilus-mediated surface sensing and the proteins that influence this process remain unknown. Here, we demonstrate that CpaL, a potential pilus tip mechanosensory protein, could be the major element of Tad pilus-mediated surface attachment and colonization in *C. crescentus*. CpaL plays an important role in the regulation of holdfast synthesis and production upon surface contact. By identifying CpaL as a key player in the process of surface recognition, our work offers valuable insights into the mechanisms of bacterial adhesion.

Introduction:

Surface recognition and attachment are essential steps in the process of bacterial surface colonization to form biofilms. Biofilms are cohesive, multicellular microbial communities wherein resident bacteria are protected from harsh environmental conditions, including variations in osmolarity, pH, nutrient accessibility, shear forces, or exposure to antibacterial agents (1, 2). Bacteria are known to sense mechanical or physical cues upon contact with solid substrata, allowing them to respond with phenotypic changes that transform free-swimming planktonic bacterial cells into surface-adherent cells that form biofilms (3–5). The process by which bacterial cells sense a surface via mechanical stimuli and convert this signal into downstream cellular processes is called surface-sensing (6, 7). A great deal of effort has been expended in deciphering the mechanisms of surface sensing in bacteria, establishing a pivotal role for extracellular appendages such as flagella and pili in this process (8–11).

Pili are thread-like, proteinaceous appendages that physically interact with their surroundings and are critical for many important cellular processes such as adherence, aggregation, biofilm formation, horizontal gene transfer, and virulence in some pathogenic species (12, 13). Pili are composed of thousands of repeats of a small protein subunit, the major pilin, and less abundant minor pilin(s) (12, 14). A subset of pili, called type IV pili (T4P), are characterized by their dynamic activity, exhibiting the ability to extend away from the cell surface and subsequently retract back into the bacterial cell by polymerization and depolymerization of the pilus fiber, respectively. This is achieved by a complex membrane-spanning machine that draws from a pool of pilins in the inner membrane to assemble the fiber, which passes through an outer membrane channel (14). Based primarily on differences in the motor components of the T4P machinery, recent phylogenetic analyses have divided the T4P into

three subclasses: T4aP, T4bP, and T4cP (11, 15, 16). Among these, T4cP, also known as tight

adherence (Tad) pili, are thought to have evolved from an archaeal ancestor. Tad pili are broadly

distributed among bacteria, including in the freshwater bacterium *Caulobacter crescentus*, where

they have been implicated in adhesion and surface-sensing (11, 17).

C. crescentus exhibits a dimorphic life cycle, wherein each cell division produces a

nonmotile stalked cell and a motile swarmer cell harboring multiple Tad pili and one flagellum at

the same pole (18). The swarmer cell subsequently undergoes differentiation into a stalked cell

as it developmentally progresses through the cell cycle, whereupon it synthesizes a

compositionally complex adhesin called holdfast, which mediates permanent surface attachment

(10, 18, 19). Interestingly, surface contact by a swarmer cell can hasten the differentiation

process by rapidly triggering multiple processes, starting with the retraction of the pili into the

cell, cessation of further pilus activity, ejection of the flagellum, and finally synthesis of the

holdfast (11). The rapid, surface-contact mediated production of holdfast requires the presence of

the pilus, and physical obstruction of pilus retraction leads to rapid holdfast synthesis even in the

absence of surface contact, implying that tension on retracting, surface-bound pili may be a cue

to sense surface contact (11, 20). However, the mechanistic details of how Tad pili sense surface

contact, convey the signal across the cell envelope, and translate it into an output capable of

upregulating surface-associated behaviors, remain poorly understood.

In this study, we address these knowledge gaps in our mechanistic understanding of Tad

pilus-mediated mechanosensing. We performed a genetic screen that identified mutations in

cpaL, whose deletion led to significantly reduced pilus synthesis. However, despite reduced pilus

production, we found that this mutant showed increased surface adherence and holdfast

synthesis. Stimulation of holdfast production was not observed when a *cpaL* mutant was cultured

in a defined medium that nutritionally restricts surface sensing. In addition, holdfast synthesis

did not increase when pilus retraction was blocked in the *cpaL* mutant to constitutively stimulate

surface sensing. Finally, we examined the predicted structure of CpaL using AlphaFold3 and

found that CpaL is comprised of a pilin-like module connected to a von Willebrand factor type A

(vWA) domain, a fold that is often implicated in mechanosensing. Further modelling suggests

that CpaL could form a complex with the minor pilins CpaJ and CpaK, that can be

accommodated at the distal end of the pilus. Collectively, these data suggest that CpaL plays an

important role in the pilus-dependent surface sensing pathway of *C. crescentus*.

Results:

A screen for mutations conferring resistance to the pilus-dependant phage ΦCbK identifies

the *cpaL* gene

117   Tad pili exhibit dynamic cycles of extension and retraction, which play a crucial role in

surface-sensing in *C. crescentus* (11). To understand this process, we sought to identify genes in

C. crescentus that impact pilus activity using a forward genetics approach to identify mutants

that are resistant to the pilus-dependent phage ΦCbK. We reasoned that impaired pilus activity,

e.g. through impaired biosynthesis or dynamics, might increase resistance to the phage ΦCbK.

To carry out our screen and all subsequent analyses, we used *C. crescentus pilA-cys* as the parent

background, where WT cells are modified to incorporate a cysteine residue in the major pilin

PilA, allowing for pilus labeling and modulation (11, 21). A pooled transposon mutant library

was generated in the parent strain and then mixed with ΦCbK, and phage resistant mutants were

isolated. The ability of these mutants to elaborate a pilus was analyzed via microscopy by

labeling pili with a thiol-reactive, maleimide-conjugated fluorophore. Mutants that produced pili

and yet were ΦCbK-resistant were chosen for further study, and the site of transposon insertion

was determined (Figure S1) (11, 21). Among these mutants, five transposon insertions mapped to

cpaL (*CCNA_0199*) (Figure S1). The gene *cpaL* is located outside the main pilus gene cluster

and has previously been shown to contribute to phage sensitivity in *C. crescentus* (22).

132   We generated an unmarked, in-frame deletion of *cpaL* and assessed the sensitivity of this

mutant to ΦCbK. We spotted serial dilutions of ΦCbK onto growth plates with *C. crescentus*

incorporated into the top agar and analyzed the formation of plaques due to phage infection

(Figure 1A). The phage-susceptible parent strain exhibited the formation of clear plaques up to

the 10^{-5} phage dilution, while the pilus-deficient mutant Δ*pilA* displayed complete resistance to

ΦCbK. In contrast, the Δ*cpaL* mutant demonstrated intermediate resistance to ΦCbK, with

formation of cloudy plaques visible only up to the 10^{-1} phage dilution. The ΦCbK resistance

phenotype in Δ*cpaL* was completely reversed when complemented with a wild-type copy of

cpaL on a replicating plasmid (Figure 1A). The growth curves of all the analyzed strains were

comparable, indicating that ΦCbK resistance in the Δ*cpaL* mutant is not due to a variation in

growth rate (Figure S2). Together, these results indicate that *cpaL* plays an important role in

pilus biosynthesis.

The Δ*cpaL* mutant produces fewer pili per cell, but with an increased average length

Since the Δ*cpaL* mutant was less sensitive to the pilus-dependent phage ΦCbK, we hypothesized

that this could be due to a decrease in the amount of pilus activity within the population. To

visualize pilus activity, we observed the internalization of externally labeled pilins into the cell

during pilus retraction, which results in cells with fluorescent bodies in *C. crescentus* (Figure 1B)

(11). We quantified the proportion of fluorescent cell bodies among synchronized swarmer cells

stained for pili, comparing the parent and mutant strains. Approximately 23.5% of synchronized

swarmer cells of the parent strain had fluorescent cell bodies after labeling, consistent with

previous reports (11, 23). In contrast, less than 3% of the Δ*cpaL* synchronized swarmer cells

displayed fluorescent cell bodies (Figure 1B and C). Moreover, approximately 6% of parent cells

had visible pili, whereas piliated cells were rarely observed for the Δ*cpaL* mutant (Figure 1B and

D). The Δ*pilA* mutant lacking pili served as a negative control, showing no fluorescence. These

results indicate that the Δ*cpaL* mutant exhibits significantly less pilus activity than the parental

strain, but that those pili that are produced can retract. The dynamic activity of pili produced by

the Δ*cpaL* mutant was examined by time-lapse microscopy, which revealed a distribution of

dynamic behaviors that is consistent with what has been observed previously for *C. crescentus*

(Movies S1-S10) (11).

Next, we incubated synchronized parental and Δ*cpaL* swarmer cells with PEG5000-

maleimide (PEG5000-mal) to block pilus retraction, through its reaction with the modified PilA

cysteine residue, while simultaneously labeling pili before imaging, as described previously (11).

We found that around 53% of the parental swarmer cell population exhibited piliated cells

(Figure 2A and B). In contrast, around 6% of swarmer cells from the Δ*cpaL* mutant population

produced pili (Figure 2A and B). Finally, the production of pili was restored to parental levels

when the Δ*cpaL* mutant was complemented with plasmid-borne wild-type *cpaL* (Figure S3).

Next, we analyzed the number of pili produced by each piliated cell after blocking and

found that the Δ*cpaL* mutant produced only one pilus per piliated cell, in contrast to the parent

strain, where 29% of piliated cells elaborated more than one pilus (Figure 2A and C). In addition,

the single pilus produced by the Δ*cpaL* mutant had an average length of 1.81 µm, which is

significantly longer than the pili produced by the parent strain (1.02 µm; consistent with previous

reports) (11) (Figure 2A and D). These findings reinforce the importance of CpaL for Tad pilus

biosynthesis in *C. crescentus,* perhaps contributing to the initiation or regulation of pilus

formation.

Deletion of *cpaL* increases surface adhesion and holdfast production

It has previously been demonstrated that pili play a crucial role in surface-sensing and

adhesion in *C. crescentus* by rapidly stimulating synthesis of the holdfast upon surface-contact

(11, 20). Therefore, we expected that the Δ*cpaL* mutant would exhibit a surface attachment

defect due to reduced pilus synthesis. To test this hypothesis, we used a surface attachment assay

(20) wherein cells were spotted onto a glass coverslip and incubated for 30 min to allow

attachment to the surface. The coverslip was then washed to remove unattached cells, and the

surface-adherent cells were imaged by microscopy. In this assay, holdfast-deficient cells do not

attach, while the Δ*pilA* mutant exhibits around 54% reduction in surface binding compared to the

pilA-cys holdfast positive (HF+) strain (Figure 3A). In contrast, the Δ*cpaL* mutant exhibited a

three-fold increase in surface binding compared to the *pilA-cys* HF+ strain (Figure 3A, S4).

To determine whether the elevated surface adherence of the Δ*cpaL* mutant was due to

increased holdfast production, we imaged holdfasts on agarose pads using a fluorescently labeled

wheat-germ agglutinin lectin (AF488-WGA) that binds specifically to the *N*-acetyl-glucosamine

moiety present in holdfasts (11, 24). We imaged on soft PYE agarose pads since they minimize

the stimulation of holdfast production via the surface-sensing pathway (19, 25). Holdfasts were

detected in 55% of the cells in the Δ*pilA* mutant and in the *pilA-cys* HF+ strain (Figure 3B). In

contrast, 74% of the cells in the Δ*cpaL* mutant population had a holdfast (Figure 3B). Surface

binding and holdfast production were restored to the levels of the *pilA-cys* HF+ strain when the

Δ*cpaL* mutant was complemented with *cpaL* (Figure S5A and B). These results suggest that the

increased surface adherence of the Δ*cpaL* mutant is attributable to elevated holdfast production,

suggesting a role for CpaL in either the developmental or surface-sensing pathway of holdfast

production.

CpaL plays a role in surface-sensing

The Δ*cpaL* mutant exhibits increased surface adhesion and holdfast synthesis in the

complex medium PYE (Figure 3A and B). In PYE, *C. crescentus* holdfast synthesis is regulated

by both the developmental pathway, where holdfast is produced during differentiation from

swarmer to stalked cells, and the surface-sensing pathway, where holdfast production is rapidly

stimulated upon surface contact, hastening cell differentiation. In contrast, *C. crescentus* cells

grown in the defined medium M2G do not undergo surface-contact mediated holdfast synthesis

(19).

To determine whether the increased surface attachment and holdfast synthesis phenotypes

of the Δ*cpaL* mutant are related to the developmental or surface sensing pathway of holdfast

synthesis, we performed surface attachment and holdfast quantification assays in defined M2G

medium, using identical conditions as in PYE (19). First, we found that pilus production in the

Δ*cpaL* mutant was comparable between M2G and PYE, with a low proportion of cells

synthesizing a single long pilus, a defect that was complemented by a plasmid copy of *cpaL*

(Figure S3). However, in M2G medium, the Δ*cpaL* mutant exhibited a 48% reduction in surface

attachment compared to the *pilA-cys* HF+ strain, as expected for a mutant with reduced pilus

production (Figure 3C). Furthermore, approximately 42% of both the parental and Δ*cpaL* mutant

populations produced holdfast, with no significant difference detected between these strains

(Figure 3D). These results are in stark contrast to the increased surface adhesion and holdfast

synthesis phenotypes of the Δ*cpaL* mutant seen in PYE. Since surface contact does not stimulate

holdfast production in M2G, these results suggest that increased holdfast synthesis and surface

attachment of the Δ*cpaL* mutant in PYE results from a role of CpaL in pilus-mediated surface-

sensing.

To further examine the interaction of cells with a surface, we monitored the timing of

holdfast synthesis in single cells in response to surface contact in PYE medium using a PDMS

microfluidic device, where cells were introduced into a well of the device and allowed to adhere

to a glass cover slip in the presence of AF488-WGA. The AF488-WGA allowed us to track the

formation of holdfast in individual cells upon surface contact. While the *pilA*-cys HF+ strain

showed rapid holdfast synthesis upon surface contact, within an average of approximately 26.13

± 2.88 sec, the Δ*cpaL* mutant was comparatively slower in producing holdfast upon surface

contact (106.4 ± 17.45 sec, Figure 4 A and B). The median holdfast synthesis times for the *pilA-*

cys HF+ mutant and the Δ*cpaL-pilA-cys* HF+ mutant were 10 sec and 20 sec, respectively. These

results indicate that *cpaL* plays a role in regulating holdfast production through the surface

sensing pathway.

The obstruction of pilus retraction does not affect surface attachment and holdfast

production in the Δ*cpaL* mutant

The above results suggest that the deletion of *cpaL* elevates holdfast production through

stimulation of the surface sensing pathway. To test if surface sensing could be further stimulated

in the Δ*cpaL* mutant, we blocked pilus retraction using PEG5000-mal. This treatment has

previously been shown to stimulate the production of holdfast via the surface sensing pathway,

even in the absence of a surface (11). Specifically, we compared surface attachment and holdfast

production among the different strains with and without PEG5000-mal to block pilus retraction.

Blocking pilus retraction in the *pilA-cys* HF+ strain caused a strong increase in the percentage of

surface-attached cells compared to the HF+ strain lacking the *pilA*-cys mutation and to the

unblocked *pilA*-cys HF+ negative control (Figure 5A), consistent with previous results (11). In

contrast, while the Δ*cpaL pilA-cys* mutant exhibited greater surface attachment compared to

pilA-cys HF+ (Figure 3A), attachment of this mutant could not be further stimulated by

PEG5000-mal (Figure 5C).

We next measured holdfast production in the presence of PEG5000-mal. Holdfast

production was stimulated by PEG5000-mal in the *pilA-cys* HF+ strain compared to the HF+

strain without the *pilA-cys* mutation and the *pilA-cys* HF+ mutant in the absence of PEG5000-

mal (Figure 5B), consistent with previous results (11). However, blocking pilus retraction in the

Δ*cpaL pilA-cys* HF+ mutant did not further increase holdfast production compared to the

untreated condition, or compared to the Δ*cpaL* HF+ mutant without the *pilA-cys* mutation, in

either the blocked or unblocked conditions (Figure 5D).

Collectively, our results indicate that the absence of *cpaL* leads to a constitutive

stimulation of the surface-sensing pathway and suggest that the *cpaL* gene plays a role not only

in pilus biogenesis but also in some aspect of the surface-sensing mechanism.

262   CpaL is predicted to adopt a pilin-like module linked to a vWA domain

To better understand the function of CpaL, we performed several sequence-based

bioinformatic analyses to identify conserved domains and sequence features of CpaL. The

predicted 3D structure of CpaL was generated using the AlphaFold3 server (26) (Figure 6A, S6A

and B). The predicted structure was then submitted to the Dali server (27) to identify

experimentally determined protein structures in the Protein Data Bank (PDB) that exhibit

structural similarity to the predicted CpaL model (Figure 6A). The top structural analogs of

CpaL determined by Dali are all von Willebrand factor type A (vWA)-domain containing

proteins that play a role in pilus or fimbrial synthesis in bacteria (Table S4). The vWA domain is

a widely distributed structural motif that, in bacteria, is characterized by its ability to mediate

surface adhesion, adherence to host-derived proteins (28, 29), and mechanosensing (6) in

conjunction with T4P.

The N-terminus of CpaL contains a domain of approximately 120 residues predicted to

adopt a pilin-like fold (residues 27-147): a long α-helix followed by a β-sheet consisting of three

β-strands (Figure 6A and B, dark grey), giving this module its characteristic lollipop shape (16).

The predicted pilin-like module of CpaL shares structural similarity with the T4P pilin

THHA1221 from *Thermus thermophilus* (PDB: 4BHR) and the minor pilin PilX from *Neisseria*

meningitidis (PDB: 2OPD), as determined by Dali, as well as the predicted structures of the *C.*

crescentus Tad minor pilins CpaK and CpaJ (Figure S9B). Sequence analysis of this module

revealed a potential cleavage site for the prepilin peptidase CpaA, which consists of a stretch of

hydrophilic residues followed by the consensus sequence G/A - X_4 - F/E, ending with a stretch of

predominantly hydrophobic residues (16, 28) (Figure S9A).

The vWA-like domain is present at the C-terminus of CpaL and is formed by two

sections of the polypeptide chain separated by a β-rich domain (residues 148-203 and 384-626)

(Figure 6A and B, blue). The vWA domain of CpaL is predicted to adopt a classical Rossmann

fold that is broadly conserved across all vWA-like domains (30). Alignment of the vWA domain

of CpaL with the vWA domain of the top Dali hit, SpaC of *L. rhamnosus* GG (PDB ID: 6M48)

(Table S4), revealed significant structural similarity in the core Rossmann fold of the vWA

domains of these proteins, with a 1.039 Å root mean square deviation (RMSD) across 118 atom

pairs. The vWA domain of CpaL also contains a sequence motif known as the metal ion-

dependant adhesion-site (MIDAS) motif (Figure 6A and B, S10A and B), which is often present

in vWA domains, where it coordinates a divalent metal ion (31). This motif is characterized by

the DxS/TxS/T, T, and D signature (32), which in CpaL corresponds to D_{160}, T_{162}, S_{164}, T_{470}, and

D_{515}, respectively, located at the top of the central β-sheet, where they are predicted to coordinate

an Mg^{2+} ion (Figure 6A and B, S10A and B). The primary sequence that comprises the CpaL

vWA domain flanks an intervening β-rich region (residues 204-384, purple) that is projected

away from the MIDAS motif and contains two consecutive six-stranded β-barrels (Figure 6A and

B). A Dali search using only the structure of the β-rich region did not return any hits with strong

similarity to this region of CpaL.

CpaL is predicted to assemble at the pilus tip

The presence of a pilin-like domain in the predicted structure of CpaL (Figure 6A and

S9B) and the similarity of this structure to pilus tip-localized minor pilins (Table S4) suggests

that CpaL may function in the context of the pilus tip. To examine this, we utilized AlpahFold3

(26) to predict a putative Tad pilus tip complex composed of CpaL, the minor pilins CpaJ and

CpaK, and ten copies of the major pilin subunit PilA to simulate a pilus filament. The sequences

of the predicted or known signal sequences of each of these proteins were removed prior to the

prediction. This yielded a structure in which CpaJ, CpaK, and the pilin-like domain of CpaL

form a trimeric complex (Figure 6C, S7) that sits atop the ten PilA subunits (Figure 6C, S8A and

B), which are themselves arranged similarly to the recently resolved structure of the PilA

filament (Figure S8C; PDB 8U1K) (33). The transmembrane helices of CpaJ, CpaK, and CpaL

are incorporated into the distal end of the filament-like PilA structure, while the vWA domain of

CpaL projects away from the rest of the complex on the opposite side (Figure 6C, S8A and B).

Importantly, predicted aligned error (PAE) scores for the model, particularly in the region

comprising the interaction between CpaJ, CpaK, and the pilin-like domain of CpaL, are low

(Figure 6D, S7C and S8A), suggesting that this could represent a biological arrangement of these

proteins.

CpaL functions differently than the other minor pilins CpaJ and CpaK

Because of the similarity of CpaL's pilin-like domain to the minor pilins CpaJ and CpaK

(Figure S9B) and its predicted assembly with these minor pilins to form a complex at the tip of

the pilus (Figure 6C and D, S7 and S8), we sought to determine if the minor pilins CpaJ and

CpaK play a similar role in the surface sensing pathway. We tested the effect of Δ*cpaJ* and

Δ*cpaK* deletions on surface adhesion and holdfast synthesis and found that deletion of these two

minor pilin genes phenocopied Δ*pilA* cells – they were resistant to the pilus-dependent phage

ΦCbK (Figure 7A), lacked pili (Figure 7B), had significantly reduced surface adhesion (Figure

7C), and produced holdfasts similarly to WT (Figure 7D). Thus, we conclude that the function of

CpaL is different from the function of the minor pilins CpaJ and CpaK.

In summary, CpaL is predicted to adopt an N-terminal pilin-like fold that may allow it to

incorporate into the tip of the pilus filament alongside the minor pilins CpaJ and CpaK, while its

C-terminal vWA domain may play a role in adhesion and/or surface-sensing.

Discussion:

In this study, we demonstrate the importance of CpaL for pilus biosynthesis in *C.*

crescentus and show that the absence of CpaL influences holdfast production, likely through

stimulation of the surface-sensing pathway. In *C. crescentus,* the absence of pili reduces

adherence to a solid substratum (11). However, while deletion of *cpaL* dramatically reduces pilus

production (Figure 2), this mutation leads to an unexpected increase in surface attachment and

holdfast synthesis (Figure 3A and B). These results prompted us to explore whether CpaL may

perform a role in pilus-mediated surface sensing. To examine this, we cultured *C. crescentus* in

M2G minimal medium, which prevents holdfast synthesis via the surface sensing pathway but

still allows for developmentally-regulated holdfast synthesis (19). Under these conditions, we

found that the Δ*cpaL* mutant exhibits reduced attachment to glass, as would be expected for a

mutant with impaired pilus biosynthesis (Figure 3C), and no longer produces increased levels of

holdfast (Figure 3D). Together, these results suggest that deletion of *cpaL* stimulates holdfast

biosynthesis and attachment through the surface-sensing pathway. This hypothesis is supported

by the absence of further stimulation of holdfast production in the Δ*cpaL* mutant after artificially

blocking pilus retraction with PEG5000-mal, which strongly boosts holdfast production and

attachment in the *pilA-cys* HF+ strain (Figure 4A and B). Thus, it appears that the surface-

sensing pathway is stimulated constitutively in the Δ*cpaL* mutant, suggesting that CpaL plays a

role in this pathway.

To gain insight into how CpaL may be performing seemingly opposing roles in pilus

production and surface sensing, we examined the predicted structure of CpaL using Alphafold3.

Analysis of its structure identified an N-terminal pilin-like module with a structural arrangement

that is characteristic of pilin proteins (16) (Figure 6A, B, S9A, and B), with particular structural

homology to the major pilin THHA1221 from *T. thermophilus* (PDB: 4BHR) (34) and the minor

pilin PilX from *N. meningitidis* (PDB: 2OPD) (35) (Figure S9B). The pilin-like module of CpaL

also shares similarity with the predicted structures of the *C. crescentus* minor pilins, CpaJ and

CpaK (Figure S9B). A potential cleavage site for the pre-pilin peptidase CpaA was identified in

the pilin domain of CpaL (Figure S9A), which suggests that CpaL could be processed to a

mature form to facilitate incorporation into the pilus fiber, as occurs for *C. crescentus* PilA, and

potentially also for CpaJ and CpaK (Figure S9A). Position +5 after the peptidase cleavage site in

T4aP and T4bP pilins is a highly conserved Glu that forms a salt bridge with the N-terminal

amine of the previously-incorporated pilin subunit, which is critical for fibre assembly.

Exceptions to this rule are noted for a subset of minor pilins, the GspK orthologs, which instead

have a non-polar residue at position +5. It is hypothesized that the absence of the conserved Glu5

in these minor pilins is indicative that they are the first component to comprise the nascent

filament(16). However, archaeal pilins lack the conserved Glu5, and although this residue is

present in bacterial Tad pilins, the structure of the *C. crescentus* PilA filament demonstrated that

Glu5 points towards the solvent and does not participate in intermolecular interactions (33).

Instead, the adjacent Tyr6 is oriented below the N-terminal Ala1 of the previously-incorporated

pilin subunit, maintaining the helical register of the filament. Therefore, Tyr6 in Tad pilins may

fulfill a parallel function to Glu5 in T4aP and T4bP pilins. However, CpaL possesses an Ala at

position +6, which may indicate that, like the GspK orthologs, CpaL is the first subunit to

comprise the nascent filament. Indeed, in our AlphaFold3-predicted structure of CpaL at the

pilus tip (Figure 6C), CpaL is oriented as the first molecule in the filament. These predicted

structural features collectively suggest that CpaL could be a Tad minor pilin in *C. crescentus*,

albeit of unusual size and architecture. Indeed, the CpaL ortholog from the Tad pilus system of

A. actinomycetemcomitans, TadG, has previously been identified as a constituent of pilus fiber

preparations (36–38).

Minor pilins with atypical domain architectures have recently been structurally resolved

in the T4P system of *S. sanguinis* (28, 39), where the minor pilins PilB and PilC were found to

have a vWA domain and a glycan binding domain, respectively, linked to a pilin module. These

studies have further demonstrated that the tripartite minor pilin complex of *S. sanguinis*,

composed of PilA, PilB, and PilC, forms an "open wings" architecture that can only be

accommodated at the tip of the pilus fiber (39, 40). The incorporation of such a minor pilin

complex at the pilus tip is consistent with what is observed in the T4aP systems of *P. aeruginosa*

and *M. xanthus*. In these species, an inner membrane complex composed of several minor pilins,

as well as the vWA domain containing non-pilin subunit PilY1, recruits major pilin subunits to

initiate pilus fiber assembly. This minor pilin-PilY1 complex is ultimately incorporated into the

tip of the growing pilus fiber after successful nucleation of pilus assembly (41, 42). Thus, CpaL,

together with the putative Tad minor pilins CpaJ and CpaK, could similarly form a complex that

is incorporated at the tip of the pilus fiber in *C. crescentus*. Indeed, our structural predictions

with AlphaFold3 suggest that CpaL forms a tripartite complex with CpaJ and CpaK (Figure S7

and S8) that could be accommodated at the distal end of the pilus filament (Figure 6C and S8).

Based on this hypothesis, it is possible that the reduced pilus production phenotype of the Δ*cpaL*

mutant is due to disruption of this complex, resulting in fewer pilus nucleation events. Infrequent

nucleation could also explain why the pili of the Δ*cpaL* mutant are longer on average: infrequent

pilus nucleation events would result in a larger pool of major pilin subunits in the inner

membrane, drawn from to assemble fewer pili (typically one per piliated cell). However, CpaL

does not simply play the role of a minor pilin since deletion of *cpaJ* or *cpaK* yields cells without

pili and without the stimulation of surface adhesion and holdfast synthesis seen for Δ*cpaL.*

In addition to a pilin-like module, CpaL also has a predicted vWA domain (6, 30, 43) (Figure

6A, B, S6A, B, and S11). T4P-associated proteins in Gram-positive and Gram-negative bacteria

harboring vWA domains are often directly involved in surface adhesion (9, 28, 29). Many vWA

domains contain a five-residue motif called the MIDAS motif, which is often required to mediate

surface adherence (28). For example, the MIDAS motif from the T4P associated protein PilC1 in

Kingella kingae is required for adherence to host tissues as well as for twitching motility (44).

CpaL is also predicted to contain a canonical MIDAS motif that coordinate a metal ion Mg^{2+}

(Figure 6A, B, and S11A, B), supporting CpaL's role in surface adhesion. Additionally, it is

well-documented that vWA domain containing proteins have a mechanosensory function in

eukaryotes (30). Thus, it is thought that vWA domains could also play a role in T4P-mediated

mechanosensing in bacteria (6, 9). Indeed, in the pilus-tip associated protein PilY1 in *P.*

aeruginosa, the vWA domain undergoes sustained conformational changes when force is applied

by AFM. This mechanosensitivity is perturbed by the mutation of a critical disulphide-bond in

the protein, which also reduces surface-sensing behaviours in *P. aeruginosa*, suggesting a link

between force-induced conformational changes in PilY1 and surface sensing (6, 9). Given the

CpaL's role in surface sensing, its predicted structure with a vWA domain, and its hypothesized

localization to the pilus tip, a similar mechanism could therefore be at play in CpaL-mediated

surface sensing in *C. crescentus*.

Despite the similarities of CpaL to PilB in *S. sanguinis* and PilY1 in *P. aeruginosa*, the

increased surface attachment of the *C. crescentus* Δ*cpaL* mutant stands in contrast to what has

been reported for the *pilB* and *pilY1* mutants, which exhibit decreased biofilm formation,

consistent with their loss of pilus production (45, 46). In fact, loss of pili in *C. crescentus*

through deletion of the major pilin *pilA* also reduces biofilm formation through abolishment of

pilus-mediated surface-sensing (11). However, deletion of *cpaL* stimulates the production of

holdfast via the surface-sensing pathway, similar to what is observed when pilus retraction is

artificially blocked. It has previously been shown that binding of bulky PEG5000-mal adducts to

the pilus fiber prevents pilus retraction by sterically occluding the entrance of the modified pilus

filament into the CpaC secretin pore in the outer membrane, thereby stimulating the surface

sensing pathway (20). Similarly, stimulation of surface sensing was observed when a glycine to

aspartate mutation was introduced into CpaC in the predicted outer lip of the pilus secretin pore

(20), again pointing to interactions between the pilus filament and the secretin as key to the

mechanism of surface sensing. Based on our results, we hypothesize that CpaL mediates the

surface sensing pathway by modulating how the pilus fiber interfaces with CpaC and rest of the

Tad pilus secretion machinery. Evidence from *M. xanthus* and *P. aeruginosa* suggests that pilus

tip proteins remain assembled as a priming complex through successive rounds of pilus

extension and retraction, and that this complex acts as a plug in the secretin pore to prevent full

retraction of the pilus into the inner membrane (47) (48). In these species, pilus retraction may

position PilY1 within the secretin pore, where force-induced conformational changes in PilY1

due to surface contact (9) may hypothetically be sensed. By analogy, conformational changes in

CpaL due to surface contact may similarly be sensed by CpaC to stimulate the surface sensing

pathway in *C. crescentus*. However, in the absence of CpaL, the pilus tip may interact with the

secretin in such a way that surface-sensing is constitutively triggered. Further investigation is

required to better understand the exact role of CpaL in the surface-sensing mechanism.

In conclusion, our results demonstrate the importance of CpaL for pilus formation and for

regulation of holdfast biosynthesis through the surface-sensing pathway. Structural predictions of

CpaL reveal similarity to the mechanosensitive vWA domain and possible incorporation at the

tip of the pilus fiber via a pilin-like module. Together, these results implicate CpaL as a

candidate mechanosensor for the Tad pilus in *C. crescentus*.

 Material and Methods:

 Bacterial strains, plasmids, and growth conditions:

The bacterial strains used in this study are listed in Table S1. *Caulobacter crescentus*

strains were grown at 30°C in peptone-yeast extract (PYE) (49) or in defined M2 medium

supplemented with 0.2% (w/v) glucose (M2G) (50). PYE was supplemented with 5 µg/ml

kanamycin (Kan), where appropriate. Commercial, chemically competent *Escherichia coli* DH5-

α (Bioline and New England Biolabs) was used for plasmid construction and was grown at 37°C

in lysogeny broth (LB) supplemented with 25 µg/ml Kan, where appropriate.

Plasmids (Table S2) were transferred to *C. crescentus* by electroporation, transduction

with ΦCr30 phage lysates, or conjugation with S17-1 *E. coli* as described previously (51). In-

frame deletion strains were made by double homologous recombination using pNPTS-derived

plasmids as previously described (52). Briefly, plasmids were introduced into *C. crescentus*, and

then two-step recombination was performed using kanamycin resistance to select for single

crossover followed by sucrose resistance to identify plasmid excision events. All mutants were

validated by sequencing to confirm the presence of the deletion.

Plasmid construction:

The *cpaL* complementation construct was made using the high-copy-number vector

pBXMCS-2 (53) with *cpaL* under the control of a xylose-inducible promoter. Because

expression from this promoter is leaky, no xylose was added to growth media for *cpaL*

expression. The *cpaL* open reading frame was amplified from *C. crescentus* NA1000 genomic

DNA using the cpaL-F and cpaL-R primers (Table S3). The *cpaL* PCR fragment was digested

using NdeI and EcoRI and ligated into pBXMCS-2 digested by the same enzymes. Clones with a

positive insert were verified by Sanger sequencing.

Screen of gene for pilus retraction:

500 µl of an overnight culture *of pilA-cys Caulobacter* cells were mixed with 50 µl of

exponential *E. coli* carrying pFD1 (Rubin et al. PNAS 1999) and briefly vortexed. The mixture

was pipetted onto a 0.22 µm filter on a vacuum manifold to concentrate cells to encourage

conjugation, and the filter containing the cell concentrate was placed cell-side up on a plain PYE

plate and incubated overnight at room temperature. Filter-concentrated cells were resuspended in

500 µl of PYE medium and mixed with 500 µl of undiluted ~10^{10} pfu/ml ϕCbK phage stock

(MOI of ~10^6) and incubated at room temperature for 10 min. 100 µl aliquots of cell/phage

mixture were then plated on PYE plates containing kanamycin to select for transposon mutants

and nalidixic acid to select against *E. coli* cells and grown at 30°C until colonies appeared (1-2

days). 184 colonies from kanamycin/nalidixic acid plates were inoculated into 96-well plate

wells containing 200 µl of PYE and grown overnight at 30°C. Cells in wells were mixed with 20

µl of DMSO and stored at -80°C for long-term storage.

Individual mutants were then inoculated into 3 ml of PYE medium and grown overnight

at 30°C. To assess pilus phenotypes, overnight cultures were labeled with 25 µg/ml AF488-mal

and incubated at room temperature for 5 minutes. To remove excess dye, cultures were

centrifuged at 7,500 *x g* for 1 min, the supernatant was removed, and cells were resuspended in

100 µl fresh PYE before imaging. 30 mutants exhibited labeled T4P and/or fluorescent cell

bodies indicative of T4P production and were further characterized. Irradiated ϕCr30 lysates of

isolated transposon mutants were generated and transduced into the parent strain YB8288 to

confirm that phage resistance and T4P phenotypes were dependent on transposon insertions.

Strains exhibiting complete or partial phage resistance along with T4P fluorescence phenotypes

where further characterized. Genomic DNA was extracted from each strain and digested using

the restriction enzyme Sau3AI at 37°C for 1 hr followed by heat inactivation at 65°C for 20 min.

Digested DNA was then ligated in 100 μl reactions to promote self-annealing using T4 phage

ligase at room temperature for 1 hr. Ligations were then used as templates for inverse PCR

reactions using primers Mariner F and Mariner R targeting the transposon element and

sequenced using the same primers. Sequences adjacent to the transposon element were mapped

to the *Caulobacter crescentus* genome to identify transposon insertion sites.

Phage sensitivity assays:

Phage sensitivity assays were performed using the pilus-specific phage ΦCbK as

described previously (23). Briefly, 200 μl of *C. crescentus* stationary phase culture was mixed

with 3 ml of 0.5% (w/v) soft PYE agar. The mixture was spread over a 1.5% (w/v) PYE agar

plate and incubated at room temperature for 1 h to solidify. A tenfold serial dilution series of

ΦCbK was prepared in PYE, and 5 μl of each dilution was spotted on top of the agar plate. The

plates were grown for 2 days at 30°C before imaging using a ChemiDoc MP (BioRad).

Growth rate analysis:

Growth of *C. crescentus* strains was measured in 24-well polystyrene plates (Falcon)

using a SpectraMax iD3 microplate reader (Molecular Devices). Stationary-phase cultures were

diluted to an optical density at 600 nm (OD_{600}) of 0.05, and 1 ml of this culture was added to the

wells of the microplate in triplicate and incubated for 24 h at 30°C under shaking. The OD_{600} was

measured at 30 min intervals to generate growth curves (OD_{600} versus time).

Synchronization of *C. crescentus* populations:

C. crescentus populations were synchronized to enrich for pilus-producing swarmer cells,

enabling facile quantification of pilus characteristics. The swarmer cells were synchronized and

collected as described previously (54) with some modifications. To do this, *C. crescentus*

cultures were grown to an OD_{600} of ~ 0.15-0.3, and 2 ml of this culture was centrifuged at 5,400

$\times g$ for 5 min. The supernatant was removed, and the cell pellet was resuspended in 280 µl of

PYE. Following this, 120 µl of polyvinylpyrrolidone colloidal silica solution (Percoll, Sigma)

was added and mixed by gentle inversion. This mixture was centrifuged at 11,000 $\times g$ for 15

min, generating two bands of distinct cell populations: stalked cells in the upper band and

swarmer cells in the lower band. 15 µl of the swarmer cell band was removed and added to 85 µl

of PYE. This solution of synchronized swarmer cells was washed with 100 µl of PYE, before

proceeding to pilus labeling, as described below.

Pilin labeling, blocking, imaging, and quantification:

The pili of *C. crescentus* were labelled as described previously (11). Briefly, 25 µg/ml of

Alexa Fluor 488 C_5 Maleimide (AF488-maleimide, ThermoFisher Scientific) was added to 100

µl of synchronized swarmer cell culture, prepared as described above, and incubated for 5 min at

room temperature. Labelled cultures were centrifuged for 1 min at 5,400 $\times g$, and the pellet was

washed once with 100 µl of PYE to remove excess dye. The labelled cell pellets were

resuspended in 7 to 10 µl of PYE, 0.5 µl of which was spotted onto a 1% agarose PYE pad

(SeaKem LE, Lonza Bioscience). The agarose pad was sandwiched between glass coverslips for

imaging.

To artificially block pilus retraction, methoxy-polyethylene glycol maleimide (PEG5000-

maleimide, Sigma) with an average molecular weight of 5 kDa was used. 500 μM PEG5000-mal

was added to 100 μl of synchronized swarmer cell cultures immediately prior to the addition of

25 μg/ml AF488-mal. Cultures were further prepared and imaged as described above.

For observing pilus dynamics, *C. crescentus* cells were grown to an OD_{600} of ~ 0.15-0.3,

labelled with AF488-maleimide as mentioned above, and 0.5 μl of the labelled sample was

spotted onto a 1% agarose PYE or M2G medium pad as appropriate (SeaKem LE, Lonza

Bioscience). The agarose pad was sandwiched between glass coverslips for imaging. Pilus

dynamics were detected by time-lapse microscopy every 3 sec for 1 min.

Imaging was performed using a Nikon Ti-E inverted fluorescence microscope with Plan

Apo 60X or 100X objectives, a GFP filter cube, an Andor iXon3 DU885 EM CCD camera, and

Nikon NIS Elements imaging software.

The number of pili per cell, the percentage of piliated cells in the whole cell population,

and the percentage of cells with fluorescent cell bodies were quantified manually using ImageJ

software (55).

Surface binding assay and holdfast quantification:

Surface binding assays and holdfast quantification were carried out in parallel on the

same samples. For each strain to be analyzed, a single colony was isolated and used to inoculate

3 ml of PYE or M2G medium, as appropriate. Tenfold serial dilutions of this culture (10^{-1} to 10^{-4}

) were prepared and incubated overnight at 30°C. The OD_{600} of each culture was measured, and

the culture with an OD_{600} of ~ 0.05 was selected. Different cultures with an OD_{600} of ~ 0.05 were

normalised exactly to $OD_{600} = 0.05$ before proceeding.

For each culture prepared as described above, 15 µl was transferred onto a glass coverslip

and incubated in dark and humid conditions (to prevent desiccation and effects from varying

light levels) for 30 min at 30°C. After incubation, coverslips were extensively washed in water,

and a 1% PYE or M2G agarose pad, as appropriate, was added on top of the cells on the

coverslip for microscopy analysis.

To quantify holdfast production, cells from the same culture preparations described above

($OD_{600} = 0.05$) were labelled with 0.5 µg/ml AF488-WGA (Wheat germ agglutinin lectin,

ThermoFisher Scientific) for 1 min before spotting them onto a 1% agarose PYE or M2G pad, as

appropriate, for microscopy analysis. WGA binds specifically to the N-acetylglucosamine

residues present in the holdfast polysaccharide (24).

Cells attached to the coverslip, and labelled holdfast, were imaged using a Nikon Ti-E

fluorescence microscope as described above. The percentage of cells attached to the coverslip,

and the percentage of cells with holdfast, were quantified manually from microscopy images

using ImageJ software (55).

To quantify surface binding and holdfast production from cells with blocked pili, 500 µM

PEG5000-mal was added to cell cultures and incubated for 5 min before performing the

experiments described above.

Timing of holdfast synthesis after surface contact:

Microfluidic well devices were constructed from PDMS (Polydimethylsiloxane) as described

previously (11, 25). Briefly, a 10:1 mixture of PDMS prepolymer:curing agent was poured into a

sterile petri plate, then the plate was placed for a few hours under a vacuum until the bubbles

were removed. The plates were placed for 3h in an oven at 65°C. A rectangle of 25 x 35 mm was

cut from the PDMS plates, with fluid access holes of 3 mm in diameter punched at 5 mm

intervals. Next, the glass coverslip and the PDMS cast were assembled after plasma treatment

and put overnight in an oven at 65°C.

For sample preparation, 200 ml of bacterial cell culture of $OD_{600} = 0.6\text{-}0.8$ grown in PYE

was diluted in 800 ml of PYE containing 0.5 mg/ml AF488-WGA (Wheat germ agglutinin lectin,

ThermoFisher Scientific) for holdfast labeling, then 10-15 μl of the above mixture was added to

one well of the PDMS device.

Time-lapse videos were taken every 5 sec for 30 min at the glass-liquid interface using a

Nikon Ti-E fluorescence microscope with Apo 60X objective, a GFP filter cube, an Andor iXon3

DU885 EM CCD camera, and Nikon NIS Elements imaging software. The attaching cells were

detected using the phase contrast channel while holdfast synthesis was detected using the GFP

channel. The time difference between holdfast synthesis and cell-surface attachment was

determined using ImageJ software (55).

CpaL structural and functional predictions:

Prediction of protein domains, their global distribution, and associated architectures was

done using the Dali (27) server which was used for comparing protein structures in 3D after

generation of the predicted structure of CpaL using AlphaFold3 (26). The structure of CpaL in

complex with CpaJ and CpaK at the pilus tip was predicted using AlphaFold3 (26). Molecular

visualization of 3D protein structures was done using the ChimeraX software (56) and

AlphaFold3 (26). The GenBank accession number for the primary sequence of CpaL is

WP_010918088.1 and the UniProt Primary accession number is A0A0H3C404. The

superposition of the vWA domain of CpaL with the vWA domain of SpaC (PDB ID: 6M48-A)

was performed using ChimeraX software (56), while the superposition of the AlphaFold3-

predicted PilA helical structure with the experimentally-determined structure of the pilus

filament (PDB 8U1K), and calculation of alignment statistics, was done using MM-align (57).

<u>Acknowledgment:</u>

This work was supported by a Fonds de Recherche du Québec, Nature et Technologies (FRQNT)

doctoral training scholarship, a Bourse de Mérite from the Faculté de Médecine at the Université de

Montréal, and a Bourse de formation à la recherche Mitacs to F.O.C.; a Natural Sciences and Engineering

Research Council of Canada (NSERC) and FRQNT postdoctoral fellowship to G.B.W.; fellowship

1342962 from the National Science Foundation (NSF) to C.K.E.; and the Canada 150 Research Chair in

Bacterial Cell Biology and grant R35GM122556 from the National Institutes of Health to Y.V.B. We also

thank members of the Brun Lab for their feedback and critical review of the manuscript.

References:
631

1. Sharma D, Misba L, Khan AU. 2019. Antibiotics versus biofilm: an emerging battleground in

microbial communities. Antimicrobial Resistance & Infection Control 8:76.

2. Berne C, Ellison CK, Ducret A, Brun YV. 2018. Bacterial adhesion at the single-cell level. 10.

Nat Rev Microbiol 16:616–627.

3. Kuchma SL, O'Toole GA. 2022. Surface-Induced cAMP Signaling Requires Multiple Features

of the Pseudomonas aeruginosa Type IV Pili. Journal of Bacteriology 204:e00186-22.

4. Gordon VD, Wang L. 2019. Bacterial mechanosensing: the force will be with you, always.

Journal of Cell Science 132:jcs227694.

5. Ellison C, Brun YV. 2015. Mechanosensing: A Regulation Sensation. Current Biology

25:R113–R115.

6. Webster SS, Wong GCL, O'Toole GA. 2022. The Power of Touch: Type 4 Pili, the von

Willebrand A Domain, and Surface Sensing by Pseudomonas aeruginosa. Journal of

Bacteriology 204:e00084-22.

7. Laventie B-J, Jenal U. 2020. Surface Sensing and Adaptation in Bacteria. Annual Review of

Microbiology 74:735–760.

8. Ellison CK, Kan J, Chlebek JL, Hummels KR, Panis G, Viollier PH, Biais N, Dalia AB, Brun YV.

2019. A bifunctional ATPase drives tad pilus extension and retraction. Science Advances

5:eaay2591.

9. Webster SS, Mathelié-Guinlet M, Verissimo AF, Schultz D, Viljoen A, Lee CK, Schmidt WC,

Wong GCL, Dufrêne YF, O'Toole GA. Force-Induced Changes of PilY1 Drive Surface Sensing

by Pseudomonas aeruginosa. mBio 13:e03754-21.

10. Hug I, Deshpande S, Sprecher KS, Pfohl T, Jenal U. 2017. Second messenger–mediated

tactile response by a bacterial rotary motor. Science 358:531–534.

11. Ellison CK, Kan J, Dillard RS, Kysela DT, Ducret A, Berne C, Hampton CM, Ke Z, Wright ER,

Biais N, Dalia AB, Brun YV. 2017. Obstruction of pilus retraction stimulates bacterial

surface sensing. Science 358:535–538.

12. Ellison CK, Whitfield GB, Brun YV. 2022. Type IV Pili: dynamic bacterial nanomachines. FEMS

Microbiology Reviews 46:fuab053.

13. Hospenthal MK, Costa TRD, Waksman G. 2017. A comprehensive guide to pilus biogenesis

in Gram-negative bacteria. 6. Nat Rev Microbiol 15:365–379.

14. Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional

consequences. Nat Rev Microbiol 17:429–440.

15. Denise R, Abby SS, Rocha EPC. 2019. Diversification of the type IV filament superfamily into

machines for adhesion, protein secretion, DNA uptake, and motility. PLOS Biology

17:e3000390.

16. Giltner CL, Nguyen Y, Burrows LL. 2012. Type IV Pilin Proteins: Versatile Molecular Modules.

Microbiol Mol Biol Rev 76:740–772.

17. Sangermani M, Hug I, Sauter N, Pfohl T, Jenal U. 2019. Tad Pili Play a Dynamic Role in

Caulobacter crescentus Surface Colonization. mBio 10:e01237-19.

18. Li G, Brown PJB, Tang JX, Xu J, Quardokus EM, Fuqua C, Brun YV. 2012. Surface contact

stimulates the just-in-time deployment of bacterial adhesins. Molecular Microbiology

83:41–51.

19. Berne C, Ellison CK, Agarwal R, Severin GB, Fiebig A, Morton RI, Waters CM, Brun YV. 2018.

Feedback regulation of Caulobacter crescentus holdfast synthesis by flagellum assembly

via the holdfast inhibitor HfiA. Molecular Microbiology 110:219–238.

20. Snyder RA, Ellison CK, Severin GB, Whitfield GB, Waters CM, Brun YV. 2020. Surface sensing

stimulates cellular differentiation in Caulobacter crescentus. Proceedings of the National

Academy of Sciences 117:17984–17991.

21. Ellison CK, Dalia TN, Dalia AB, Brun YV. 2019. Real-time microscopy and physical

perturbation of bacterial pili using maleimide-conjugated molecules. 6. Nat Protoc

14:1803–1819.

22. Christen M, Beusch C, Bösch Y, Cerletti D, Flores-Tinoco CE, Del Medico L, Tschan F, Christen

B. 2016. Quantitative Selection Analysis of Bacteriophage φCbK Susceptibility in

Caulobacter crescentus. Journal of Molecular Biology 428:419–430.

23. Ellison CK, Rusch DB, Brun YV. 2019. Flagellar Mutants Have Reduced Pilus Synthesis in

Caulobacter crescentus. J Bacteriol 201:e00031-19.

24. Merker RI, Smit J. 1988. Characterization of the Adhesive Holdfast of Marine and

Freshwater Caulobacters. Applied and Environmental Microbiology 54:2078–2085.

25. Hoffman MD, Zucker LI, Brown PJB, Kysela DT, Brun YV, Jacobson SC. 2015. Timescales and

Frequencies of Reversible and Irreversible Adhesion Events of Single Bacterial Cells. Anal

Chem 87:12032–12039.

26. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L,

Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D,

Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A,

Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB, Gladman H,

Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S, Stecula A,

Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P,

Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of biomolecular

interactions with AlphaFold 3. Nature 630:493–500.

27. Holm L. 2020. Using Dali for Protein Structure Comparison. Methods Mol Biol 2112:29–42.

28. Raynaud C, Sheppard D, Berry J-L, Gurung I, Pelicic V. 2021. PilB from Streptococcus

sanguinis is a bimodular type IV pilin with a direct role in adhesion. PNAS 118.

29. Kant A, Palva A, von Ossowski I, Krishnan V. 2020. Crystal structure of lactobacillar SpaC

reveals an atypical five-domain pilus tip adhesin: Exposing its substrate-binding and

assembly in SpaCBA pili. Journal of Structural Biology 211:107571.

30. Springer TA. 2014. von Willebrand factor, Jedi knight of the bloodstream. Blood 124:1412–

1425.

31. Shimaoka M, Takagi J, Springer TA. 2002. Conformational Regulation of Integrin Structure

and Function. Annual Review of Biophysics and Biomolecular Structure 31:485–516.

32. Lee J-O, Rieu P, Arnaout MA, Liddington R. 1995. Crystal structure of the A domain from the

a subunit of integrin CR3 (CD11 b/CD18). Cell 80:631–638.

33. Sonani RR, Sanchez JC, Baumgardt JK, Kundra S, Wright ER, Craig L, Egelman EH. 2023. Tad

and toxin-coregulated pilus structures reveal unexpected diversity in bacterial type IV pili.

Proc Natl Acad Sci U S A 120:e2316668120.

34. Karuppiah V, Thistlethwaite A, Derrick JP. 2016. Structures of type IV pilins from Thermus

thermophilus demonstrate similarities with type II secretion system pseudopilins. J Struct

Biol 196:375–384.

35. Helaine S, Dyer DH, Nassif X, Pelicic V, Forest KT. 2007. 3D structure/function analysis of PilX

reveals how minor pilins can modulate the virulence properties of type IV pili. Proceedings

of the National Academy of Sciences 104:15888–15893.

36. Inoue T, Tanimoto I, Ohta H, Kato K, Murayama Y, Fukui K. 1998. Molecular Characterization

of Low-Molecular-Weight Component Protein, Flp, in Actinobacillus

actinomycetemcomitans Fimbriae. Microbiology and Immunology 42:253–258.

37. Inouye T, Ohta H, Kokeguchi S, Fukui K, Kato K. 1990. Colonial variation and fimbriation of

Actinobacillus actinomycetemcomitans. FEMS Microbiology Letters 69:13–17.

38. Harano K, Yamanaka A, Okuda K. 1995. An antiserum to a synthetic fimbrial peptide of

Actinobacillus actinomycetemcomitans blocked adhesion of the microorganism. FEMS

Microbiology Letters 130:279–285.

39. Shahin M, Sheppard D, Raynaud C, Berry J-L, Gurung I, Silva LM, Feizi T, Liu Y, Pelicic V.

2023. Characterization of a glycan-binding complex of minor pilins completes the analysis

of Streptococcus sanguinis type 4 pili subunits. Proceedings of the National Academy of

Sciences 120:e2216237120.

40. Anger R, Pieulle L, Shahin M, Valette O, Le Guenno H, Kosta A, Pelicic V, Fronzes R. 2023.

Structure of a heteropolymeric type 4 pilus from a monoderm bacterium. 1. Nat Commun

14:7143.

41. Nguyen Y, Harvey H, Sugiman-Marangos S, Bell SD, Buensuceso RNC, Junop MS, Burrows LL.

2015. Structural and Functional Studies of the Pseudomonas aeruginosa Minor Pilin, PilE*.

Journal of Biological Chemistry 290:26856–26865.

42. Nguyen Y, Sugiman-Marangos S, Harvey H, Bell SD, Charlton CL, Junop MS, Burrows LL.

2015. Pseudomonas aeruginosa Minor Pilins Prime Type IVa Pilus Assembly and Promote

Surface Display of the PilY1 Adhesin*. Journal of Biological Chemistry 290:601–611.

43. Bharati KP, Prashanth UR. 2011. Von Willebrand Disease: An Overview. Indian J Pharm Sci

73:7–16.

44. Sacharok AL, Porsch EA, St. Geme JW. 2022. The Kingella kingae PilC1 MIDAS Motif Is

Essential for Type IV Pilus Adhesive Activity and Twitching Motility. Infection and Immunity

91:e00338-22.

45. Gurung I, Spielman I, Davies MR, Lala R, Gaustad P, Biais N, Pelicic V. 2016. Functional

analysis of an unusual type IV pilus in the Gram-positive Streptococcus sanguinis.

Molecular Microbiology 99:380–392.

46. Orans J, Johnson MDL, Coggan KA, Sperlazza JR, Heiniger RW, Wolfgang MC, Redinbo MR.

2010. Crystal structure analysis reveals Pseudomonas PilY1 as an essential calcium-

dependent regulator of bacterial surface motility. Proceedings of the National Academy of

Sciences 107:1065–1070.

47. Treuner-Lange A, Chang Y-W, Glatter T, Herfurth M, Lindow S, Chreifi G, Jensen GJ, Søgaard-

Andersen L. 2020. PilY1 and minor pilins form a complex priming the type IVa pilus in

Myxococcus xanthus. 1. Nat Commun 11:5054.

48. Guo S, Chang Y, Brun YV, Howell PL, Burrows LL, Liu J. 2024. PilY1 regulates the dynamic

architecture of the type IV pilus machine in Pseudomonas aeruginosa. Nat Commun

15:9382.

49. Poindexter JS. 1964. BIOLOGICAL PROPERTIES AND CLASSIFICATION OF THE CAULOBACTER

GROUP. Bacteriol Rev 28:231–295.

50. Johnson RC, Ely B. 1977. ISOLATION OF SPONTANEOUSLY DERIVED MUTANTS OF

CAULOBACTER CRESCENTUS. Genetics 86:25–32.

51. Ely B. 1991. [17] Genetics of Caulobacter crescentus, p. 372–384. *In* Methods in

Enzymology. Academic Press.

52. Ried JL, Collmer A. 1987. An nptI-sacB-sacR cartridge for constructing directed, unmarked

mutations in Gram-negative bacteria by marker exchange-eviction mutagenesis. Gene

57:239–246.

53. Thanbichler M, Iniesta AA, Shapiro L. 2007. A comprehensive set of plasmids for vanillate-

and xylose-inducible gene expression in Caulobacter crescentus. Nucleic Acids Res

35:e137.

54. Schrader JM, Shapiro L. 2015. Synchronization of Caulobacter Crescentus for Investigation

of the Bacterial Cell Cycle. J Vis Exp 52633.

55. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nat Methods 9:671–675.

56. Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. UCSF

ChimeraX: Tools for structure building and analysis. Protein Sci 32:e4792.

57. Mukherjee S, Zhang Y. 2009. MM-align: a quick algorithm for aligning multiple-chain protein

complex structures using iterative dynamic programming. Nucleic Acids Res 37:e83.

58. Marks ME, Castro-Rojas CM, Teiling C, Du L, Kapatral V, Walunas TL, Crosson S. 2010. The

Genetic Basis of Laboratory Adaptation in Caulobacter crescentus. J Bacteriol 192:3678–

3688.

59. Skerker JM, Shapiro L. 2000. Identification and cell cycle control of a novel pilus system in

Caulobacter crescentus. EMBO J 19:3223–3234.

786
787

Strain designation	Genotype and method of construction	Source or reference
Caulobacter crescentus		
YB8220	NA1000 *hfsA+ pilA*^{T36C}	(11)
YB6374	NA1000 *hfsA+ ΔpilA*	(11)
YB8644	NA1000 *hfsA+ pilA*^{T36C} *ΔcpaL* (Transduced strain YB8215 with lysate made from strain YB8287)	This study
YB8454	NA1000 *pilA*^{T36C} *ΔcpaK* (Electroporated plasmid pNPTS138ΔcpaK from strain 8453 into strain 8288.	This study
YB10278	NA1000 *pilA*^{T36C} *ΔcpaJ* (Electroporated plasmid pNPTS138ΔcpaJ from strain 10277 into strain 8288.	
FC764	NA1000 *hsfA+*	(58)
YB8215	NA1000 *hfsA+ ΔcpaL* (electroporated plasmid from strain YB8206 into strain FC764)	This study
YB8288	NA1000 *pilA*^{T36C}	(11)
LS3118	NA1000 *ΔpilA*	(59)
YB8294	NA1000 *pilA*^{T36C} *ΔcpaL* (Electroporated	This study

	strain YB8288 with plasmid from strain Y8206)	
YB9770	NA1000 $pilA^{T36C}$ pBXMCS-2 (Electroporated strain YB8288 with plasmid pBXMCS-2)	This study
YB9771	NA1000 $\Delta pilA$ pBXMCS-2 (Electroporated strain LS3118 with plasmid pBXMCS-2)	This study
YB9772	NA1000 $pilA^{T36C}$ $\Delta cpaL$ pBXMCS-2 (Electroporated strain YB8294 with plasmid pBXMCS-2)	This study
YB9742	NA1000 $pilA^{T36C}$ $\Delta cpaL$ pBXMCS-2 $cpaL$ (YB9741 Electroporated strain YB8294 with plasmid from YB9741)	This study
YB9773	NA1000 $hfsA+$ $pilA^{T36C}$ pBXMCS-2 (Electroporated strain YB8220 with plasmid pBXMCS-2)	This study
YB9774	NA1000 $hfsA+$ $pilA^{T36C}$ $\Delta cpaL$ pBXMCS-2 (Electroporated strain YB8644 with plasmid pBXMCS-2)	This study
YB9775	NA1000 $hfsA+$ $\Delta pilA$ pBXMCS-2 (Electroporated strain YB6374 with plasmid pBXMCS-2)	This study

YB9776	NA1000 *hfsA+ pilA*^{T36C} *ΔcpaL* pBXMCS-2 *cpaL* (Plasmid from YB9741 electroporated into strain YB8644 by electroporation)	This study
YB9777	NA1000 *hfsA+ pilA*^{T36C} *ΔcpaJ* (Electroporated strain YB 8220 with plasmid pNPTS138 *ΔcpaJ* from YB10277)	This study
YB9778	NA1000 *hfsA+ pilA*^{T36C} *ΔcpaK* (Electroporated strain YB 8220 with plasmid pNPTS138 *ΔcpaK* from YB 8453)	This study
Escherichia coli		
YB4030	S17-1/pNPTS138 Δ*pilA*	(19)
YB8206	α-select/pNPTS139 Δ*cpaL*	This study
YB8286	α-Select/pNPTS139 *pilA*^{T36C}	(11)
YB8453	α-select/pNPTS138 Δ*cpak*	This study
YB102777	α-select/pNPTS138 Δ*cpaJ*	This study
YB9741	NEB 5-alpha/pBXMCS-2 *cpaL*	This study

794 **Table S2: Plasmids used in this study**

Plasmid	Description	Source or reference
pNPTS138	Litmus 38 derivative, *oriT sacB* Kan^r	M. R. K. Alley
pNPTS139	Litmus 39 derivative, *oriT sacB* Kan^r	M. R. K. Alley
pNPTS139 *pilA*^{T36C}	pNPTS139 containing 500-bp fragments upstream and downstream of *pilA*^{T36C} point mutation	(11)
pNPTS138 Δ*pilA*	pNPTS138 containing 480 bp fragments upstream and downstream of *pilA*	(19)
pNPTS139 Δ*cpaL*	pNPTS139 containing 500-bp fragments upstream and downstream of *cpaL*	This study
pNPTS138 Δ*cpaJ*	pNPTS138 containing 500 bp fragments upstream and downstream of *cpaJ*	This study
pNPTS138 Δ*cpaK*	pNPTS138 containing 500 bp fragments upstream and downstream of *cpaK*	This study
pBXMCS-2	High copy replicating plasmid, xylose inducible promotor, Kan^r	(53)
pBXMCS-2 *cpaL*	pBXMCS-2 containing *cpaL* open reading frame	This study

798   **Table S3: Primers used in this study**

Primer name	Sequence*	Function
cpaL-UP F	**TTC TGG ATC CAC GAT** <u>CGA GCA CCC GCT GGT CCA</u>	pNPTS139Δ*cpaL* construction
cpaL-UP R	<u>TCT CAG ACG GCG GGC GAA A</u>	
cpaL-DN F	<u>TCT TTC CAG GCC ATT GCT C</u>	
cpaL-DN R	**AGC TTC CTG CAG GAT** <u>GCA GGA CGG CAA TTC GGT</u>	
cpaL-F	GCA **CAT ATG** <u>ATG CCG ACC AAG AGC CGT TT</u>	pBXMCS-2 *cpaL* construction
cpaL-R	GGG **GAA TTC** <u>CTA TCT GGC GAT CCG CAG GT</u>	
cpaJ-UP F	**GCC AAG CTT CTC TGC AGG AT**T CGA GCC GGT TTC CGC GAA G	pNPTS138Δ*cpaJ* construction
cpaJ-UP R	*GGT GAA CGA GGT GGC GGC GTA* <u>GAT GAG GGA GCG TGA AGC CA</u>	
cpaJ-DN F	<u>TAC GCC GCC ACC TCG TTC ACC</u>	
cpaJ-DN R	**GCG AAT TCG TGG ATC CAG AT**<u>G AGG TGT AGA CAT AGC TCG CC</u>	
cpaK-UP F	**GC GAA TTC TGG ATC CAC GAT** CAC GTC TCC GGC TGG CGC GTT	pNPTS138Δ*cpaK* construction
cpaK-UP R	*T CTC GGC CTT* <u>GCG CCA GAA GCT GGA AAG GGG GCG</u>	
cpaK-DN F	**C TTC TGG CGC** <u>AAG GCC GAG ACG ATC CTC TGG TCG ACG</u>	

cpaK-DN R	**CA GAA AGC TTC CTG CAG GAT** <u>GGT CGC GGG CTT GAT CGG CCC</u>	
Mariner F	ACG GTA TCG ATA AGC TTG ATA TCG A	
Mariner R	CAG AGT TGT TTC TGA AAC ATG GCA	

799   *Restriction sites and regions of plasmid complementarity to facilitate Gibson assembly are in

800   bold; regions of complementary to the target amplicon are underlined; regions of reverse

801   complementarity (to facilitate allele assembly) are italicized.

803 **Table S4: The top five PDB hits identified by the DALI server for the predicted structure of CpaL**

PDB hit	Description	Species	Z-score	rmsd	lali	nres	%id
6m48-A	SpaC	*Lactobacillus rhamnosus GG*	20.3	4.1	241	813	16
2ww8-A	Cell wall surface anchor family protein; Pilus adhesion RrgA	*Streptococcus pneumoniae*	19.9	13.2	254	815	18
6to1-A	Minor fimbrium subunit Mfa5	*Porphyromonas gingivalis*	18.9	4	232	566	21
7b7p-A	Type IV pilus biogenesis protein PilB	*Streptococcus sanguinis*	18.7	4.8	244	420	17
7w6b-A	von Willebrand factor type A domain protein	*Streptococcus oralis*	18.3	5.5	212	782	21

Rmsd: Root mean square deviation, calculated across only the aligned portions of CpaL and the

hit; lali: The number of residues of the hit protein that were aligned with CpaL; nres: the total

number of residues in the hit protein; %id: percentage identity between CpaL and the hit protein.

Figure legends:

Figure S1: Genetic screen for resistance to a pilus-dependent phage identifies *cpaL* as a

mediator of pilus activity. Schematic of the forward genetic screen used to identify mutants that

are resistant to the pilus-dependent *Caulobacter* phage, φCbK, despite having pili. A *C.*

crescentus Mariner transposon mutant library was generated and mixed with φCbK and grown

on plates. Phage resistant mutants were isolated and imaged for pilus synthesis. Red bars indicate

the position of transposon insertions identified within the *cpaL* (CCNA_00199) gene coding

sequence (gray arrow). Scale bar, 10 μm.

Figure 1: Deletion of *cpaL* reduces sensitivity to the pilus-specific phage ΦCbK by reducing

pilus activity. A: Top agar phage sensitivity assays for the parent strain (NA1000 *pilA*-cys),

Δ*pilA* (phage-resistant strain lacking pili), the Δ*cpaL* mutant, and the plasmid-complemented

Δ*cpaL* mutant. Serial dilutions of the phage were spotted on plates with top agar containing each

bacterial strain. ND, no dilution; EV, empty vector. **B:** Representative microscopy images for

synchronized swarmer cells of the parent strain (NA1000 *pilA*-cys) and the isogenic Δ*cpaL*

mutant, labeled with the AF488-maleimide dye (green) that reacts with the engineered cysteine

residue in the major pilin, PilA. Scale bars, 10 μm. White arrows indicate cells with labeled pili,

red arrows indicate fluorescent cell bodies. **C:** Quantification of the percentage of cells with

fluorescent cell bodies in synchronized populations of the indicated strains labelled with AF488-

maleimide. **D:** Quantification of the percentage of piliated cells in synchronized swarmer cells of

the indicated strains labelled with AF488-maleimide dye (green). Results are the mean of four

independent biological replicates, with at least 500 cells analyzed per replicate. Error bars

represent the standard error of the mean (SEM). Statistical comparisons were made using

Tukey's multiple comparisons test. ***, P <0.001; **, P < 0.01; ns, no significant difference.

Figure S2: *Caulobacter* **cell growth is not affected by the absence of** *cpaL***.** Growth curves of

the indicated strains of *C. crescentus* grown in PYE. Data were collected every 30 min and are

the average of six independent biological replicates. Error bars represent the standard error of the

mean.

Figure 2: Deletion of *cpaL* **reduces the number of pili produced per cell while increasing**

their average length. A: Representative microscopy images of synchronized swarmer cells of

the parent strain (NA1000 *pilA*-cys) and the isogenic Δ*cpaL* mutant labeled with AF488-

maleimide (green) that reacts with the engineered cysteine residue in the major pilin, PilA, in

conjunction with pilus retraction blocking using PEG5000-mal, which reacts with the same

cysteine residue. White arrows indicate cells with a single pilus per cell. Orange arrows indicate

cells with two or more pili per cell. Scale bars, 10 μm. **B:** Quantification of the percentage of

piliated cells in synchronized populations of the indicated strains when pilus retraction was

blocked with PEG5000-mal. Results are the mean of eight independent biological replicates,

with at least 200 cells analyzed per replicate. Error bars represent the standard error of the mean.

Statistical comparisons were made using Tukey's multiple comparisons test. **C:** Quantification of

the number of pili produced per piliated cell of the indicated strains, measured in synchronized

populations after blocking pilus retraction with PEG5000-mal. Results are the mean of four

independent biological replicates with at least 200 cells analyzed per replicate. Error bars

represent the standard error of the mean. **D:** Average length of pili produced by synchronized

swarmer cells of the indicated strains after blocking with PEG5000-mal. 100 pili were measured

for each strain. Error bars indicate the minimum to maximum range of lengths. Statistical

comparison was made using a two-tailed unpaired T-test. ****, $P < 0.0001$; **, $P < 0.01$.

Figure S3: Representative microscopy images for the strains used in the phage assay in

PYE and M2G medium. Synchronized swarmer cells were blocked for pilus retraction with

PEG5000-mal and labeled with AF488-maleimide (green). Tested strains included the parent

(EV: empty vector), the Δ*cpaL* (EV) mutant, the Δ*pilA* (EV) mutant, and the plasmid-

complemented Δ*cpaL* mutant. Scale Bar, 10 μm.

Figure 3: The deletion of *cpaL* increases surface attachment and holdfast production in

PYE but not in M2G medium. **A:** Quantification of the attachment of cells grown in the

complex medium PYE to a glass coverslip after 30 min of incubation relative to the strain *pilA*-

cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast positive). **B:** Quantification of the percentage of cells

producing holdfast in the population in the complex medium PYE on an agarose pad. **C:**

Quantification of the attachment of cells grown in the minimal medium M2G to a glass coverslip

after 30 min of incubation relative to the strain *pilA*-cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast

positive). **D:** Quantification of the percentage of cells producing holdfast in the population in the

minimal medium M2G on an agarose pad. Data are the mean of four independent biological

replicates. Error bars indicate the standard error of the mean (SEM). Statistical comparisons were

made using Tukey's multiple comparisons test. ****, $P < 0.0001$; **, $P < 0.01$; *, $P < 0.05$.

Figure S4: Representative microscopy images for cells attached to the surface. Cells are grown in the complex medium PYE and incubated for 30 min to a glass coverslip (HF+: Holdfast positive). Holdfasts are labeled with the AF488 conjugated wheat germ agglutinin (AF488-WGA). Scale Bar, 10 μm.

Figure S5: Plasmid complementation of the Δ*cpaL* mutant rescues its increased attachment and holdfast production phenotypes. A: Quantification of the attachment of cells grown in the complex medium PYE to a glass coverslip after 30 min of incubation **B:** Quantification of the percentage of cells producing holdfast in the population in the complex medium PYE on an agarose pad. Data are the mean of three independent biological replicates. Error bars indicate the standard error of the mean. Statistical comparisons were made using Tukey's multiple comparisons test. [****], $P < 0.0001$; [***], $P < 0.001$; [**], $P < 0.01$.

Figure 4: The absence of CpaL leads to a delay in holdfast production after surface contact. **A:** Logarithmic violin plot showing the time of holdfast synthesis after surface contact (sec), for there independent replicates of *pilA-cys* HF+ (n=119) and Δ*cpaL-pilA-cys* HF+ (n=163). Statistical comparison was made using a two-tailed unpaired T-test. ****, $P < 0.0001$. **B:** Table showing the mean and the median of the strains used in the violin plot.

Figure 5: The Δ*cpaL* mutant is insensitive to the addition of PEG5000-mal: A: Quantification of the attachment of cells to a glass coverslip after 30 min of incubation relative to the strain WT HF+ (NA1000, HF+: Holdfast positive). PEG5000-mal was added to cultures 5 min before cells were added to the coverslip to block pili retraction in *pilA-cys* strains. **B:**

Quantification of the percentage of cells producing holdfast in the population after 5 min of incubation with PEG5000-mal on agarose pad. The percentage of cells producing holdfast in the population was quantified after labeling holdfast with the AF488 conjugated wheat germ agglutinin (AF488-WGA). **C:** Quantification of the attachment of cells to a glass coverslip after 30 min of incubation relative to the strain Δ*cpaL* HF+ (NA1000, HF+: Holdfast positive). PEG5000-mal was added to cultures 5 min before cells were added to the coverslip to block pili retraction in *pilA-cys* strains. **D:** Quantification of the percentage of cells producing holdfast in the population after 5 min of incubation with PEG5000-mal on agarose pad. The percentage of cells producing holdfast in the population was quantified after labeling holdfast with the AF488 conjugated wheat germ agglutinin (AF488-WGA). Data are the mean of three independent biological replicates. Error bars indicate the standard error of the mean. Statistical comparisons were made using Tukey's multiple comparisons test. ****, P <0.0001.

Figure 6: CpaL is predicted to contain a pilin-like module and a vWA domain. A: Ribbon diagram showing the predicted structure of CpaL generated using AlphaFold3. Domains and coloring are as described in panel B. The inset depicts the MIDAS motif in the vWA-like domain which is composed of five residues (D160, T162, S164, T470, and D515, green) with a predicted Mg^{2+} ion (gray) incorporated among those residues. **B:** Schematic representation of the CpaL domain organization. The first region at the N-terminus is predicted to be a signal peptide (residues 1-26, yellow), followed by a pilin-like module (residues 27-147, dark grey). A single von Willebrand Factor A-like domain (vWA) is divided between two distinct segments of the CpaL sequence (residues 148-203 and 384-626, blue), between which are two tandem β-rich domains (residues 204-383, pink). **C:** Top-ranked structure of a complex composed of the minor

pilins CpaJ (blue), CpaK (green), CpaL (red), and ten copies of the major pilin subunit PilA

(orange) predicted by AlphaFold3. The predicted or known signal sequences of each protein

were removed prior to the prediction. **D:** The predicted aligned error (PAE) scores for the model

depicted in panel C. The PAE indicates the positional error in angstroms (Å) for a given pair of

residues across all protein chains in the model.

Figure S6: A: Representative of the five predicted model structures of CpaL predicted by

Alphafold3. Structures are ranked according to the predicted template modeling (pTM) score and

are colored according to the predicted local distance difference test (pLDDT) score, which

indicates per-residue model confidence for the protein chain. The colors represented the

measurement of the confidentiality in the prediction of the structure and are according to the

predicted local distance difference test (pLDDT) score. **B:** Confidence in the prediction of the

structure is indicated by the predicted aligned error (PAE) scores, which indicate positional error

in angstroms for a given pair of residues within the protein chain.

Figure S7: A: Top-ranked structure of a complex composed of the minor pilins CpaJ (blue),

CpaK (green), and CpaL (red), predicted by AlphaFold3. The predicted or known signal

sequences of each protein were removed prior to the prediction. **B:** The predicted aligned error

(PAE) scores for the model depicted in panel A. The PAE indicates the positional error in

angstroms (Å) for a given pair of residues across all protein chains in the model. **C:** Structures of

a complex composed of CpaJ, CpaK, and CpaL, with their predicted signal sequences removed,

predicted by AlphaFold3. Structures are aligned and shown from the same orientation. Structures

are ranked according to the predicted template modeling (pTM) score and are colored according

to the predicted local distance difference test (pLDDT) score, which indicates per-residue model

confidence for the individual protein chains within the complex. **D:** Confidence in the prediction

of the complex is indicated by the predicted aligned error (PAE) scores, which indicate

positional error in angstroms for a given pair of residues across all protein chains.

Figure S8: A: Structures of a complex composed of CpaJ, CpaK, CpaL, and ten PilA subunits,

with their predicted signal sequences removed, predicted by AlphaFold3. Structures are aligned

and shown from the same orientation. Structures are ranked according to the predicted template

modeling (pTM) score and are colored according to the predicted local distance difference test

(pLDDT) score, which indicates per-residue model confidence for the individual protein chains

within the complex. **B:** Confidence in the prediction of the complex is indicated by the predicted

aligned error (PAE) scores, which indicate positional error in angstroms for a given pair of

residues across all protein chains. **C:** Alignment of the structure of the *C. crescentus* PilA

filament determined by cryo-EM (from PDB 8U1K) and the structure of the PilA filament as

determined by AlphaFold3 (from the prediction in panel A).

Figure S9: CpaL has a pilin-like module with a predicted prepilin peptidase (CpaA)

cleavage site. A: N-terminal sequence of PilA, minor pilins CpaK and CpaJ, and CpaL.

Hydrophilic residues are highlighted in orange, and hydrophobic residues are highlighted in blue.

The consensus sequence (G/A-X-X-X-F/E) for recognition by the prepilin peptidase CpaA is

shown in a grey box. The potential CpaA cleavage site is indicated with a red arrow. **B:**

Structural comparison of the predicted pilin-like module of CpaL, CpaJ, and CpaK, with their

predicted signal sequences removed, and the predicted and experimentally determined structure

of the T4P THHA1221 from Thermus thermophilus (PDB:4BHR) and the minor pilin PilX from

Neisseria meningitidis T4aP (PDB ID: 1AY2).

Figure S10: Structural comparison of the vWA domain of CpaL and the SpaC protein of

***Lactobacillus rhamnosus* GG. A:** Structural superposition of the predicted vWA domain of

CpaL (blue, residues 148-203, and 383-626) with the vWA domain of SpaC (pink, residues 129-

386) determined by X-ray crystallography (PDB ID: 6M48-A). **B:** Magnified image of the box in

panel A depicting the MIDAS motif regions of CpaL (blue) and SpaC (pink). The Mg^{2+} ion that

co-crystallized with SpaC as well as the Mg^{2+} ion predicted by Alhphafold3 for the vWA domain

of CpaL are represented by two large grey spheres, while the water molecule involved in Mg^{2+}

ion coordination by the SpaC are depicted as a small red sphere. RMSD: 1.039 Å between 118

atom pairs; RMSD.: 9.981 Å across all 229 atoms.

Figure 7: The deletion of minor pilin genes *cpaJ* and *cpaK* phenocopies Δ*pilA*. A: Top agar

phage sensitivity assays. Serial dilutions of the phage were spotted on plates with top agar

containing each bacterial strain. ND, no dilution. **B:** Representative microscopy images of

synchronized swarmer cells that were blocked for pilus retraction with PEG5000-mal and labeled

with AF488-maleimide (green). Scale Bar, 10 μm. **C:** Quantification of the attachment of cells

grown in the complex medium PYE to a glass coverslip after 30 min of incubation relative to the

strain *pilA*-cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast positive, Parent: NA1000 HF-). **D:**

Quantification of the percentage of cells producing holdfast in the population in the complex

medium PYE on an agarose pad. Data are the mean of three independent biological replicates.

Error bars indicate the standard error of the mean (SEM). Statistical comparisons were made

using Tukey's multiple comparisons test. ****, P <0.0001.

Movies

Movie S1: Time-lapse movie of labeled non-synchronized parent cells showing pili extension-

retraction dynamics after labeling with AF488-maleimide (green). Capture rate is 3 sec per

frame. Frame rate is 5 fps. Scale Bar, 5 μm.

Movie S2: Time-lapse movies of labeled non-synchronized *cpaL* mutant cells showing pili

extension-retraction dynamics after labeling with AF488-maleimide (green). Capture rate is 3 sec

per frame. Frame rate is 5 fps. Scale Bar, 5 μm.

1002

1003 **Movie S3:** A representative time-lapse movie of cells producing holdfast upon surface contact

1004 for the parent in the presence of AF488-WGA. The cell bodies are in gray and the holdfasts are

1005 in green. Capture rate is 5 sec per frame. Frame rate is 25 fps. Scale Bar, 5 μm.

1006

1007 **Movie S4:** A representative time-lapse movie of cells producing holdfast upon surface contact

1008 for the *cpaL* mutant in the presence of AF488-WGA. The cell bodies are in gray and the

1009 holdfasts are in green. Capture rate is 5 sec per frame. Frame rate is 25 fps. Scale Bar, 5 μm.

1010

Figure S1

Figure S1: Genetic screen for resistance to a pilus-dependent phage identifies *cpaL* as a mediator of pilus activity. Schematic of the forward genetic screen used to identify mutants that are resistant to the pilus-dependent *Caulobacter* phage, φCbK, despite having pili. A *C. crescentus Mariner* transposon mutant library was generated and mixed with φCbK and grown on plates. Phage resistant mutants were isolated and imaged for pilus synthesis. Red bars indicate the position of transposon insertions identified within the *cpaL* (CCNA_00199) gene coding sequence (gray arrow). Scale bar, 10 µm.

Figure 1

Figure 1: **Deletion of *cpaL* reduces sensitivity to the pilus-specific phage ΦCbK by reducing pilus activity. A:** Top agar phage sensitivity assays for the parent strain (NA1000 *pilA*-cys), Δ*pilA* (phage-resistant strain lacking pili), the Δ*cpaL* mutant, and the plasmid-complemented Δ*cpaL* mutant. Serial dilutions of the phage were spotted on plates with top agar containing each bacterial strain. ND, no dilution; EV, empty vector. **B:** Representative microscopy images for synchronized swarmer cells of the parent strain (NA1000 *pilA*-cys) and the isogenic Δ*cpaL* mutant, labeled with the AF488-maleimide dye (green) that reacts with the engineered cysteine residue in the major pilin, PilA. Scale bars, 10 µm. White arrows indicate cells with labeled pili, red arrows indicate fluorescent cell bodies. **C:** Quantification of the percentage of cells with fluorescent cell bodies in synchronized populations of the indicated strains labelled with AF488-maleimide. **D:** Quantification of the percentage of piliated cells in synchronized swarmer cells of the indicated strains labelled with AF488-maleimide dye (green). Results are the mean of four independent biological replicates, with at least 500 cells analyzed per replicate. Error bars represent the standard error of the mean (SEM). Statistical comparisons were made using Tukey's multiple comparisons test. ***, P <0.001; **, P < 0.01; ns, no significant difference.

Figure S2

Figure S2: *Caulobacter* **cell growth is not affected by the absence of** *cpaL*
Growth curves of the indicated strains of *C. crescentus* grown in PYE. Data were collected every 30 min and are the average of six independent biological replicates. Error bars represent the standard error of the mean.

Figure 2

Figure 2: Deletion of *cpaL* reduces the number of pili produced per cell while increasing their average length. A: Representative microscopy images of synchronized swarmer cells of the parent strain (NA1000 *pilA*-cys) and the isogenic Δ*cpaL* mutant labeled with AF488-maleimide (green) that reacts with the engineered cysteine residue in the major pilin, PilA, in conjunction with pilus retraction blocking using PEG5000-mal, which reacts with the same cysteine residue. White arrows indicate cells with a single pilus per cell. Orange arrows indicate cells with two or more pili per cell. Scale bars, 10 μm. **B:** Quantification of the percentage of piliated cells in synchronized populations of the indicated strains when pilus retraction was blocked with PEG5000-mal. Results are the mean of eight independent biological replicates, with at least 200 cells analyzed per replicate. Error bars represent the standard error of the mean. Statistical comparisons were made using Tukey's multiple comparisons test. **C:** Quantification of the number of pili produced per piliated cell of the indicated strains, measured in synchronized populations after blocking pilus retraction with PEG5000-mal. Results are the mean of four independent biological replicates with at least 200 cells analyzed per replicate. Error bars represent the standard error of the mean. **D:** Average length of pili produced by synchronized swarmer cells of the indicated strains after blocking with PEG5000-mal. 100 pili were measured for each strain. Error bars indicate the minimum to maximum range of lengths. Statistical comparison was made using a two-tailed unpaired T-test. ****, P <0.0001; **, P < 0.01.

Figure S3

PYE	Parent (EV)	Δ*cpaL* (EV)	Δ*pilA* (EV)	Δ*cpaL* pBMXCS-2::*cpaL*

Figure S3: Representative microscopy images for the strains used in the phage assay in PYE and M2G medium. Synchronized swarmer cells were blocked for pilus retraction with PEG5000-mal and labeled with AF488-maleimide (green). Tested strains included the parent (EV: empty vector), the Δ*cpaL* (EV) mutant, the Δ*pilA* (EV) mutant, and the plasmid-complemented Δ*cpaL* mutant. Scale Bar, 10 μm.

Figure 3

Figure 3: The deletion of *cpaL* increases surface attachment and holdfast production in PYE but not in M2G medium. **A:** Quantification of the attachment of cells grown in the complex medium PYE to a glass coverslip after 30 min of incubation relative to the strain *pilA*-cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast positive). **B:** Quantification of the percentage of cells producing holdfast in the population in the complex medium PYE on an agarose pad. **C:** Quantification of the attachment of cells grown in the minimal medium M2G to a glass coverslip after 30 min of incubation relative to the strain *pilA*-cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast positive). **D:** Quantification of the percentage of cells producing holdfast in the population in the minimal medium M2G on an agarose pad. Data are the mean of four independent biological replicates. Error bars indicate the standard error of the mean (SEM). Statistical comparisons were made using Tukey's multiple comparisons test. ****, P <0.0001; **, P <0.01; *, P <0.05.

Figure S4

Parent HF-

Δ*cpaL-pilA-cys* HF-

Parent HF+

Δ*cpaL-pilA-cys* HF+

Figure S4: Representative microscopy images for the attached cells to the surface. Cells are grown in the complex medium PYE and incubated for 30 min to a glass coverslip (HF+: Holdfast positive). Holdfasts are labeled with the AF488 conjugated wheat germ agglutinin (AF488-WGA). Scale Bar, 10 µm.

Figure S5

Figure S5: Plasmid complementation of the ΔcpaL mutant rescues its increased attachment and holdfast production phenotypes. A: Quantification of the attachment of cells grown in the complex medium PYE to a glass coverslip after 30 min of incubation **B:** Quantification of the percentage of cells producing holdfast in the population in the complex medium PYE on an agarose pad. Data are the mean of three independent biological replicates. Error bars indicate the standard error of the mean. Statistical comparisons were made using Tukey's multiple comparisons test. [****], P <0.0001; [***], P <0.001; [**], P <0.01.

Figure 4

A

B

Strain	Mean ± SD (sec)	Median
pilA-cys HF+	26.13 ± 2.88	10
Δ*cpaL-pilA-cys* HF+	106.4 ± 17.45	20

Figure 4: The absence of CpaL leads to a delay in holdfast production after surface contact. A: Logarithmic violin plot showing the time of holdfast synthesis after surface contact (sec), for there independent replicates of *pilA-cys* HF+ strain (n=119) and Δ*cpaL-pilA-cys* HF+ strain (n=163) data. Statistical comparison was made using a two-tailed unpaired T-test. ****, P <0.0001. **B:** Table showing the mean and the median of the strains used in the violin plot.

Figure 5

Figure 5: The addition of PEG5000-mal only affects the *pilA-cys* mutant: A: Quantification of the attachment of cells to a glass coverslip after 30 min of incubation relative to the strain WT HF+ (NA1000, HF+: Holdfast positive). PEG5000-mal was added to cultures 5 min before cells were added to the coverslip to block pili retraction in *pilA-cys* strains. **B:** Quantification of the percentage of cells producing holdfast in the population after 5 min of incubation with PEG5000-mal on agarose pad. The percentage of cells producing holdfast in the population were quantified after labeling holdfast with the AF488 conjugated wheat germ agglutinin (AF488-WGA). **C:** Quantification of the attachment of cells to a glass coverslip after 30 min of incubation relative to the strain Δ*cpaL* HF+ (NA1000, HF+: Holdfast positive). PEG5000-mal was added to cultures 5 min before cells were added to the coverslip to block pili retraction in *pilA-cys* strains. **D:** Quantification of the percentage of cells producing holdfast in the population after 5 min of incubation with PEG5000-mal on agarose pad. The percentage of cells producing holdfast in the population were quantified after labeling holdfast with the AF488 conjugated wheat germ agglutinin (AF488-WGA). Data are the mean of three independent biological replicates. Error bars indicate the standard error of the mean. Statistical comparisons were made using Tukey's multiple comparisons test. ****, P <0.0001.

Figure 6

Figure 6: CpaL is predicted to contain a pilin-like module and a vWA domain. **A:** Ribbon diagram showing the predicted structure of CpaL generated using AlphaFold3. Domains and coloring are as described in panel B. The inset depicts the MIDAS motif in the vWA-like domain which is composed of five residues (D160, T162, S164, T470, and D515, green) with a predicted Mg^{2+} ion (gray) incorporated among those residues. **B:** Schematic representation of the CpaL domain organization. The first region at the N-terminus is predicted to be a signal peptide (residues 1-26, yellow), followed by a pilin-like module (residues 27-147, dark grey). A single von Willebrand Factor A-like domain (vWA) is divided between two distinct segments of the CpaL sequence (residues 148-203 and 384-626, blue), between which are two tandem β-rich domains (residues 204-383, pink). **C:** Top-ranked structure of a complex composed of the minor pilins CpaJ (blue), CpaK (green), CpaL (red), and ten copies of the major pilin subunit PilA (orange) predicted by AlphaFold3. The predicted or known signal sequences of each protein were removed prior to the prediction. **D:** The predicted aligned error (PAE) scores for the model depicted in panel C. The PAE indicates the positional error in angstroms (Å) for a given pair of residues across all protein chains in the model.

Figure S6

A

Rank 1   Rank 2   Rank 3   Rank 4   Rank 5

pLDDT confidence

low   high

B

PAE (Å)

0   15   30

Figure S6: A: Representative of the five predicted model structures of CpaL predicted by Alphafold3. Structures are ranked according to the predicted template modeling (pTM) score and are colored according to the predicted local distance difference test (pLDDT) score, which indicates per-residue model confidence for the protein chain. The colors represented the measurement of the confidentiality in the prediction of the structure and are according to the predicted local distance difference test (pLDDT) score. **B:** Confidence in the prediction of the structure is indicated by the predicted aligned error (PAE) scores, which indicate positional error in angstroms for a given pair of residues within the protein chain.

Figure S7

Figure S7: A: Top-ranked structure of a complex composed of the minor pilins CpaJ (blue), CpaK (green), and CpaL (red), predicted by AlphaFold3. The predicted or known signal sequences of each protein were removed prior to the prediction. **B:** The predicted aligned error (PAE) scores for the model depicted in panel A. The PAE indicates the positional error in angstroms (Å) for a given pair of residues across all protein chains in the model. **C:** Structures of a complex composed of CpaJ, CpaK, and CpaL, with their predicted signal sequences removed, predicted by AlphaFold3. Structures are aligned and shown from the same orientation. Structures are ranked according to the predicted template modeling (pTM) score and are colored according to the predicted local distance difference test (pLDDT) score, which indicates per-residue model confidence for the individual protein chains within the complex. **D:** Confidence in the prediction of the complex is indicated by the predicted aligned error (PAE)

Figure S8

A Rank 1   Rank 2   Rank 3   Rank 4   Rank 5

pLDDT confidence
low — high

B

PAE (Å)
0   15   30

C

PilA filament cryo-EM
structure (PDB 8U1K)

r.m.s.d. = 3.10 Å
TM-score = 0.877

AlphaFold3 PilA
filament prediction

Figure S8: A: Structures of a complex composed of CpaJ, CpaK, CpaL, and ten PilA subunits, with their predicted signal sequences removed, predicted by AlphaFold3. Structures are aligned and shown from the same orientation. Structures are ranked according to the predicted template modeling (pTM) score and are colored according to the predicted local distance difference test (pLDDT) score, which indicates per-residue model confidence for the individual protein chains within the complex. **B:** Confidence in the prediction of the complex is indicated by the predicted aligned error (PAE) scores, which indicate positional error in angstroms for a given pair of residues across all protein chains. **C:** Alignment of the structure of the *C. crescentus* PilA filament determined by cryo-EM (from PDB 8U1K) and the structure of the PilA filament as determined by AlphaFold3 (from the prediction in panel A).

Figure S9

A

PilA	MTKFVTRFLKDESG ATAIEYGLIVALIAVVIVTAVTTLGTNL
CpaK	MSRRRPLSSFWRDRFG VSAVEFALIAPVMIVMYCGLAEVTQ
CpaJ	MASRSLIARARLRLARSAGRFARADEG ATAVEFALVAIPFLMLLFAIIELGLVFL
CpaL	MPTKSRFLRRLSEGVAAFARRLRRDDRGA IAIQEALLALPLSILLFGLLDVGRLSL

■ Hydrophilic   ■ Hydrophobic   G/A XXXXF/E: consensus signal sequence

B

Pillin THHA1221
Thermus thermophilus
(Predicted)

Minor pilin PilX
Neisseria meningitidis
(Predicted)

Pilin-like CpaL
C. crescentus
(Predicted)

Minor pilin CpaJ
C. crescentus
(Predicted)

Minor pilin CpaK
C. crescentus
(Predicted)

Pillin THHA1221
Thermus thermophilus
(4BHR)

Minor pilin PilX
Neisseria meningitidis
(2OPD)

Figure S9: CpaL has a pilin-like module with a predicted prepilin peptidase (CpaA) cleavage site. A: N-terminal sequence of PilA, minor pilins CpaK and CpaJ, and CpaL. Hydrophilic residues are highlighted in orange, and hydrophobic residues are highlighted in blue. The consensus sequence (G/A-X-X-X-F/E) for recognition by the prepilin peptidase CpaA is shown in a grey box. The potential CpaA cleavage site is indicated with a red arrow. **B:** Structural comparison of the predicted pilin-like module of CpaL, CpaJ, and CpaK, with their predicted signal sequences removed, and the predicted and experimentally determined structure of the T4P THHA1221 from Thermus thermophilus (PDB:4BHR) and the minor pilin PilX from *Neisseria meningitidis* T4aP (PDB ID: 1AY2).

Figure S10

A

B

MIDAS region

T470
T229

D143
D160

D258

Mg²⁺

D515

Mg²⁺

S147
S164

S143
T162

■ Predicted vWA domain CpaL   ■ vWA domain of SpaC. PDB ID: 6M48-A

RMSD.: 1.039 Å between 118 atoms pairs
RMSD.: 9.981 Å across all 229 atoms

Figure S10: Structural comparison of the vWA domain of CpaL and the SpaC protein of *Lactobacillus rhamnosus* GG. A: Structural superposition of the predicted vWA domain of CpaL (blue, residues 148-203, and 383-626) with the vWA domain of SpaC (pink, residues 129-386) determined by X-ray crystallography (PDB ID: 6M48-A). **B:** Magnified image of the box in panel A depicting the MIDAS motif regions of CpaL (blue) and SpaC (pink). The Mg²⁺ ion that co-crystallized with SpaC as well as the Mg²⁺ ion predicted by Alhphafold3 for the vWA domain of CpaL are represented by two large grey spheres, while the water molecule involved in Mg²⁺ ion coordination by the SpaC are depicted as a small red sphere. RMSD: 1.039 Å between 118 atom pairs; RMSD.: 9.981 Å across all 229 atoms.

Figure 7

A

B

Figure 7: The deletion of minor pilin genes *cpaJ* and *cpaK* behaves differently from the deletion of the minor pilin *cpaL*. A: Top agar phage sensitivity assays. Serial dilutions of the phage were spotted on plates with top agar containing each bacterial strain. ND, no dilution. **B:** Representative microscopy images of synchronized swarmer cells that were blocked for pilus retraction with PEG5000-mal and labeled with AF488-maleimide (green). Scale Bar, 10 μm. **C:** Quantification of the attachment of cells grown in the complex medium PYE to a glass coverslip after 30 min of incubation relative to the strain *pilA*-cys HF+ (NA1000 *pilA*-cys, HF+: Holdfast positive, Parent: NA1000 HF-). **D:** Quantification of the percentage of cells producing holdfast in the population in the complex medium PYE on an agarose pad. Data are the mean of three independent biological replicates. Error bars indicate the standard error of the mean (SEM). Statistical comparisons were made using Tukey's multiple comparisons test. ****, P <0.0001.