Comparative pangenomics unveils distinct host adaptation levels

and conserved biosynthetic potential in microbiome *Clostridia*

Lucas De Vrieze[1,2], Jan Aerts[3], Joleen Masschelein[1,2, #]

[1] Laboratory for Biomolecular Discovery & Engineering, Department of Biology, KU Leuven,

Heverlee, Belgium

[2] VIB-KU Leuven Center for Microbiology, Heverlee, Belgium

[3] Augmented Intelligence for Data Analytics, Department of Biosystems, KU Leuven, Heverlee,

Belgium

[#]Corresponding author, joleen.masschelein@kuleuven.be

Email addresses:

Lucas De Vrieze: lucas.devrieze@kuleuven.be

Jan Aerts: jan.aerts@kuleuven.be

Joleen Masschelein: joleen.masschelein@kuleuven.be

Abstract

Bacterial species adopt various lifestyles to thrive in diverse ecological niches, ranging from living freely in the soil to being part of various human and animal microflora. However, the impact of these adaptation processes on their genomes and metabolisms remains largely unexplored beyond the genus level. Investigating these evolutionary dynamics for higher taxonomic levels can enhance our understanding of the relationship between host adaptation and their functional capabilities. In this regard, we examined the evolutionary trajectories and metabolic capabilities of the *Clostridia* class, which display a variety of lifestyles and are of high importance for industry, medicine and microbiome research. First, we uncover that the clostridial orders have significantly different adaptation rates. Second, we show that the *Oscillospirales* order has undergone extensive genomic and functional specialisation toward a host-associated lifestyle, while the *Lachnospirales* order tends to be at a lower level of host association, retaining a remarkably high number of free-living trait genes and a high degree of metabolic versatility. Third, we reveal substantial differences in genomic architecture and metabolic versatility between the clostridial orders and link these to the progressing stages of host adaptation. Additionally, we identify widely conserved biosynthetic gene clusters, highlighting untapped biosynthetic potential of evolutionary significance. Hence, the beyond-genus level analyses in this study provide valuable new insights with implications for the biological sciences and biotechnology.

Keywords

Introduction

Microorganisms constantly interact with their environment, adapting not only through short-term regulatory and metabolic responses, but also through long-term genomic changes. The continuous acquisition, mutation and deletion of genes is closely tied to environmental selection pressures, giving rise to various lifestyles that vary in their dependence on external factors. These selection pressures drive evolutionary processes, such as host adaptation and genome streamlining, which can comprehensively reshape microbial genomes. This, in turn, has a profound impact on their functional and metabolic versatility, and hence their ability to survive in different ecological niches (Dufresne, Garczarek and Partensky 2005, Siguier, Gourbeyre and Chandler 2014, Marais, Calteau and Tenaillon 2008).

Species in the *Clostridia* class exhibit a remarkable variety of lifestyles and phenotypes, ranging from thermophiles and psychrophiles to acidophiles and pathogens. They occupy diverse ecological niches, including soil, sediments and the intestinal tracts of humans and animals. This versatility underlies their increasing industrial and medical relevance, exemplified by industrially relevant species like *C. acetobutylicum* and *C. beijerinckii* (Birgen et al. 2021, Patakova et al. 2019, Roberts et al. 2010) and notorious pathogens like *C. botulinum*, *C. tetani* and *C. difficile* (Cohen et al. 2017, Smith, Hill and Raphael 2015, Sandhu and McBride 2018). Moreover, *Clostridium* species were the first anaerobes discovered to produce biosynthetic gene cluster (BGC)-encoded secondary metabolites with potent biological activities (Pahalagedara et al. 2020, Letzel, Pidot and Hertweck 2013, Behnken and Hertweck 2012, Sugimoto et al. 2019).

Despite the abundance and relevance of clostridial species, only few comparative studies have explored their evolutionary history and ecological adaptations beyond the species level. Previous higher-level comparative studies focused at most on a few individual species or a single genus. For example, a study of *Clostridium sensu stricto* found evidence of a highly plastic genome, harbouring a relatively large metabolic core and many specialised genetic capabilities that underpin the mainly free-living nature of this genus (Udaondo, Duque and Ramos 2017). Another study mainly focusing on *Clostridium sensu stricto* identified divergent evolution of the shikimate pathway in pathogenic clostridia and extensive gene mobility among their toxin genes (Cruz-Morales et al. 2019). A last study focusing on the gut-associated genus *Blautia* revealed

that many gene exchange events are associated with metabolic functions, likely driven by the dynamically

changing host environment. Moreover, horizontal gene transfer may have caused a metabolic dependence

on other gut bacteria (Maturana and Cárdenas 2021).

In this work, we present a comprehensive study of the interplay between host adaptation and

genomic architecture for the taxonomic class of *Clostridia*. By combining pangenomic and phylogenetic

analyses, we characterised the evolutionary dynamics within different orders and inferred lifestyles for each

order's ancestors. In parallel, we assessed the metabolic versatility of each species and linked this with their

evolutionary history. We also screened all genomes for genomic hallmarks and mobile genetic elements

(MGEs) that are related with progressing host adaptation. Finally, we zoomed in on secondary metabolite

BGCs, assessing their conservation and evolutionary patterns.

75 Materials and methods

76 *Data gathering, QC and filtering*

All genome assemblies were retrieved from RefSeq using the search query "Clostridia" in the NCBI Datasets

CLI (v16.2.0) (O'Leary et al. 2024), excluding MAGs and atypical assemblies (# = 10599 as of 22/05/2025). The

resulting set was filtered by number of contigs (<= 200), N50 (>= 50 kb), completeness (>= 90%) and

contamination (<= 10%), and dereplicated at species level using skDER (v1.3.1) (Salamzade and Kalan 2023)

(ANI cutoff: 96%, AF cutoff: > 50%; **Add. File 2: Figure S1**), resulting in 1010 representative assemblies. For

consistency, all assemblies were reannotated with Bakta (v1.9.1) (Schwengers et al. 2021) and taxonomically

verified using GTDB-tk (v2.4.1) (Chaumeil et al. 2022). To ensure sufficient statistical power in downstream

analyses, only taxonomic orders with at least 30 representative species were retained.

85 *Metabolic versatility analysis*

86 The metabolic versatility of all taxa was assessed by KEGG module completeness as determined by the

87 MicrobeAnnotator (v2.0.5) pipeline (Ruiz-Perez, Conrad and Konstantinidis 2021) in light mode at default

88 settings. The module completenesses were clustered using Ward's hierarchical clustering and plotted using

89 in-house Python scripts. Optimal leaf ordering was enabled while plotting the clustermap.

Pangenome construction and analysis

A pangenome was constructed for the full genome set using PPanGGOLiN (v2.0.5) (Gautreau et al. 2020) with gene defragmentation enabled. To account for the high diversity in this class-level genome set, clustering was carried out with a relaxed sequence identity threshold of 40% and an alignment coverage cutoff of 40%, while the pangenome was partitioned using a core genome threshold of 90% and an accessory genome threshold of 0.099% (or 1/1010), as appeared appropriate from our parameter benchmark (**Add. File 3**). Presence/absence matrices for the different clostridial orders were extracted from the full class-level matrix by selecting for gene families present in at least one taxon from that order. The same pangenome partitioning thresholds were applied for consistency.

Phylogeny construction

A phylogenetic tree for the full clostridial genome set was constructed with PhyloPhlAn (v3.0.67) (Asnicar et al. 2020) with a custom gene marker database derived from the representative protein sequences of the earlier identified core gene families. The tree was rooted using *B. subtilis* 168 as outgroup (RefSeq: GCF_000009045.1). The phylogeny was computed at medium diversity setting using DIAMOND (Buchfink, Reuter and Drost 2021) as the mapping algorithm, MAFFT (Katoh and Standley 2013) as the alignment algorithm, Trimal (Capella-Gutiérrez, Silla-Martínez and Gabaldón 2009) as the alignment trimmer, and IQ-TREE2 (Minh et al. 2020) as the tree constructor. 10.000 ultrafast bootstraps (Minh, Nguyen and von Haeseler 2013) were requested by manually adding the "-bb 10000" flag to the IQ-TREE entry in the PhyloPhlAn configuration file. The outgroup was trimmed from the rooted phylogeny using the R package *ape*, after which the tree was visualised with iTol (v6.9) (Letunic and Bork 2024) and ggtree (v3.11.1) (Yu et al. 2017).

Functional annotation of the pangenome

The protein family reference sequences of the pangenome were functionally annotated using eggNOG-mapper (v2.1.12) (Cantalapiedra et al. 2021). COG category counts and fractions were calculated using pandas (v2.2.1) and plotted using ggplot2 (v3.5.1).

Genomic divergence rate analysis

A genomic divergence rate analysis estimates the rate of gene exchange within a clade and statistically

compares these rates for significant differences. It requires the phylogenies of each examined clade to have

an identical time scale. Therefore, a subtree was extracted for each order from the full phylogeny using *ape*.

Gene exchange (GE) models were inferred for each order using *Panstripe* (Tonkin-Hill et al. 2023). All options

were left at their defaults, except that Gaussian distributions were employed to fit the underlying generalised

linear models (GLMs). Cumulative GE plots and statistical comparisons of the GE models were carried out

using the appropriate *Panstripe* functions.

Ancestral state reconstruction

To track the evolutionary origin of gene exchanges, the ancestral states of all gene families were reconstructed

from the full core genome phylogeny and the full pangenome presence/absence matrix. This was done

according to Wagner parsimony with equal penalty scores for both gene gains and losses using Count (v10.4)

(Csurös 2010). KOG annotations of the genes exchanged at the tree nodes of interest were retrieved from the

earlier made eggNOG annotations and visualised using in-house R scripts.

Mobilome analysis

Pseudogenes were detected using Pseudofinder (v1.1.0) (Syberg-Olsen et al. 2022) at default settings. The

NCBI Protein database filtered for taxon ID 186801 (the *Clostridia* class) was used as reference database and

indexed using DIAMOND (v2.0.14). Insertion elements (ISEs) were inferred using ISEscan (v1.7.2.3) (Xie and

Tang 2017) at default settings. Biosynthetic gene clusters (BGCs) were identified using antiSMASH (v8.0.0)

(Blin et al. 2025) at default settings. Prophages were found by running PHASTEST (Wishart et al. 2023) in lite

mode using the image from Docker Hub (digest 2477a2f3b6b8). Phage infectivity graphs were made using

supervenn (v0.5.0). All mobile element counts were normalised by dividing by the genome length and

statistically compared using the Brunner-Munzel test ($\alpha = 0.05$) in scipy. Significance annotations were added

using *statannotations* (v0.6).

BGC clustering analysis

All BGCs identified during the mobilome analysis were clustered using BiG-SCAPE (v1.1.8) (Navarro-Muñoz et

al. 2020), adding MIBiG reference BGCs (flag '--mibig'), mixing all BGC classes (flag '--mix'), and including

singletons (flag '--include_singletons'). A BiG-SCAPE distance cutoff value of 0.3 was applied as appeared

appropriate from parameter benchmarking (**Add. File 4**) and the resulting similarity networks. To integrate

phylogroup metadata in CytoScape (v3.10.2), the unidirectional network links in the output network files of

BiG-SCAPE were annotated by order or MIBiG and converted into bidirectional links by adding the reverse

link. BGC gene alignments were generated using clinker (v0.0.28) (Gilchrist and Chooi 2021).

Results

Composing the genome set

We downloaded all available *Clostridia* genomes from NCBI. After applying quality controls and species

dereplication, we had a collection of 1010 genomes. These sequences have been collected from various

isolation sources, including both open ecosystems as well as host-associated systems like human or animal

microbiomes. After taxonomic classification using GTDB-tk, we selected four orders that together cover a

large share of the *Clostridia* class, while being sufficiently large for strong statistical power (# >= 30) (**Add. File**

2: Figure S2): *Clostridiales*, *Lachnospirales*, *Oscillospirales* and *Peptostreptococcales*. The *Clostridiales*

comprise mainly free-living species from *Clostridium sensu stricto*. The *Lachnospirales* and the *Oscillospirales*

contain prevalent constituents of the human gut microbiome (the *Clostridium XIVa* and *Clostridium IV* species

groups, resp.). The *Peptostreptococcales* are commonly found as part of various microflora in the human body

as well (e.g. *C. difficile*).

Pangenome partition sizes and functional annotation indicate high genome plasticity

To investigate the evolutionary dynamics, we first captured the gene conservation in all genomes through the

construction of a pangenome for our full genome set. Using a carefully selected set of clustering and

partitioning thresholds (**Add. File 3**), the pangenomes for each assessed group of taxa were constructed and

partitioned (**Table 1**). The recurringly large shell and cloud partitions suggest that there is a high level of

diversity among all orders, meaning that each genome is likely to harbour unique gene families. This finding

highlights the high level of genomic plasticity that is typical for species living in highly dynamic environments

like open ecosystems or human-associated microflora.

Table 1. Number of gene families in the pangenome partitions for each examined order, both in absolute numbers and referenced to the total number of gene families in each order's pangenome.

	All genomes (# = 1010)	*Clostridiales* (# = 209)	*Lachnospirales* (# = 349)	*Oscillospirales* (# = 164)	*Peptostreptococcales* (# = 94)
Core (> 90%)	353	690	574	465	444
	0.09 %	0.79 %	0.36 %	0.59 %	0.81 %
Shell (0.1 % - 90%)	158,627	39,179	63,580	30,325	21,451
	39.78 %	44.64 %	40.06 %	38.43 %	39.04 %
Cloud (< 0.1%)	239,797	47,903	94,546	48,109	33,053
	60.16 %	54.58 %	59.58 %	60.98 %	60.15 %

After establishing the pangenome partitions, we took a closer look at the functional differences

between the orders (**Figure 1**). The functional profile is similar for most core genomes and is dominated by

the ribosomal proteins (COG category J). Nevertheless, a notable difference is the larger share of signal

transduction mechanisms in the core genome of *Clostridiales* (COG category T). Such mechanisms are a

hallmark of a free-living lifestyle as they facilitate more complex sensing of the environment, enabling more

specialised responses. In almost all orders, the majority of the shell and cloud gene families have no known

function (COG categories S and -), suggesting the presence of mobile genetic elements (MGEs), highly

specialised functions, and/or rapidly evolving proteins that fall outside the scope of our current

understanding.

Figure 1. COG functional annotation profiles for all pangenome partitions of each examined clostridial order. Fractions are referenced to the number of gene families in a partition. The functional profiles are similar, except for a larger share of signal transduction genes in the core genome of the *Clostridiales* order, which indicates a free-living lifestyle.

KEGG pathway completeness analysis uncovers two distinct levels of metabolic versatility

A KEGG pathway completeness analysis gauges the metabolic versatility of an organism by assessing the completeness of metabolic pathway modules as documented in the KEGG Pathway database, based on the genomic presence of the module's constituent genes. Remarkably, two distinct levels of pathway completeness were observed for our full genome set. 394 out of 1010 organisms have visibly more complete pathway modules, indicating a richer metabolism (**Figure 2**). This discrepancy is not driven by a limited number of pathways but is the result of a genome-wide trend, as multiple modules tend to be more complete. This aligns with earlier observations of a remarkably high metabolic versatility within the *Bacillota* phylum, especially within the *Clostridium* genus (Pascal Andreu et al. 2023, Vieira-Silva et al. 2016).

Interestingly, the *Lachnospirales* order has a higher fraction of versatile organisms than the other orders (55% vs. 25% - 35%; **Add. File 2: Supp. Fig. S3**). Moreover, only this order comprises genera and lineages that are (almost) all metabolically versatile (*Enterocloster, Lacrimispora Butyvibrio,* one of two *Blautia* lineages) Also, this versatility appears independent of genome size, except for the *Clostridiales* order

Figure 2. Ward's clustered heatmap of KEGG module completeness, annotated by taxonomic order. Metabolically versatile genomes recurringly show a general trend in higher module completenesses. Summarising, the key differing modules comprise glycolysis-related and BCAA biosynthesis pathways, as well as Lys biosynthesis, the pentose phosphate pathway and other carbon metabolisms. For a figure with all KEGG module labels, see Add. File 2: Fig. S5.

(**Add. File 2: Supp. Fig. S4**). Multiple clostridial genera contain both versatile and non-versatile species,

confirming an intra-genus divide that has previously been reported (Vieira-Silva et al. 2016).

Notably, the most prominent differences in module completeness are found in core metabolic

processes: the central carbon metabolism (glycolysis, pentose phosphate pathway) and branched-chain

amino acid and Lys biosynthesis. A versatile *Lachnospirales* species thus has a higher metabolic independence

than one would expect from a host-associated bacterium. In turn, this more developed primary metabolism

may support a more extensive secondary metabolism, providing additional advantages for the microbe.

Genomic divergence rate analysis reveals significantly different evolutionary dynamics

The higher metabolic versatility of the *Lachnospirales* suggests that this order has a divergent evolutionary

trajectory than its other host-associated counterparts *Oscillospirales* and *Peptostreptococcales*. To assess

whether this discrepancy indeed may result from distinct evolutionary dynamics, gene exchange (GE) models

were fitted for each order separately and statistically compared to each other using *Panstripe*.

Panstripe uses a compound generalised linear model (GLM) to relate phylogenetic branch lengths

with the number of genes exchanged at those branches. Using an additional GLM term, internal nodes of the

tree can be differentiated from terminal ones, yielding core and tip gene exchange rates, respectively.

Exchange rates significantly different from 0 signify an association between branch length and the number of

gene exchange events, and thus active gene exchange. The core rate is to be interpreted as the overall gene

exchange rate underpinning the general core structure of the phylogeny, while the tip rate relates to short-

term gene mobility processes such as transposon or phage activity. However, in highly diverse genome sets

like the one in this work, an orthologous gene family may have been split into multiple gene families during

pangenome construction, artificially inflating both core and tip gene exchange rates. Consequently, *Panstripe*

is modelling here a more general genomic divergence (GD) rate, capturing both intrinsic gene evolution as

well as gene exchanges, rather than only the latter. Nevertheless, we argue that this more general genomic

divergence rate is still sufficient to address our research question, as it provides a measure of the pace at

which genomes have been evolving.

This GD modelling revealed several interesting findings about the evolutionary dynamics of the four

orders (**Table 2**). First, the *Oscillospirales* have the lowest core divergence rate, which is congruent with the

observation that the streamlined genomes of host-associated species evolve at a reduced rate (Dufresne,

Garczarek and Partensky 2005, Marais, Calteau and Tenaillon 2008). In contrast, the *Lachnospirales* have

significantly higher core and tip divergence rates, indicating a higher level of both long-term gene divergence

and short-term gene mobility, respectively. Strikingly, this contradicts its host-associated status since higher

levels of gene divergence and mobility imply still ongoing host adaptation (see also **Figure G**) (Siguier,

Gourbeyre and Chandler 2014). Furthermore, the tip divergence rate of the *Peptostreptococcales* is higher as

well, however, the core rates are not. One possible explanation is that this concerns increased gene mobility

at the onset of host adaptation. Another possibility is that the *Peptostreptococcales* are reaching the end of

the host conversion process, which reflects in dropping adaptation rates and the last to be eliminated gene

mobility.

Table 2. Fitted core and tip divergence rates of the GE models for each order, including p-values of all pairwise rate comparisons. *Oscillospirales* have a significantly lower divergence rate, consistent with its host-associated nature, whereas *Lachnospirales* display a high divergence rate inconsistent with its presumed host-associated status. This inconsistency can be explained by *Lachnospirales* tending to be in the earlier stages of host adaptation

		Tip divergence rate			
		Clostridiales 363.39 ± 42.28	*Lachnospirales* 621.03± 35.88	*Oscillospirales* 445.80 ± 35.63	*Peptostreptococcales* 537.80 ± 67.47
Core divergence rate	*Clostridiales* 5140.29 ± 603.74	/	$9.8 \cdot 10^{-6}$ ****	0.150	0.0255 *
	Lachnospirales 5995.25 ± 547.81	0.314	/	$3.37 \cdot 10^{-3}$ **	0.306
	Oscillospirales 4218.27 ± 412.98	0.210	0.0199 *	/	0.186
	Peptostreptococcales 5222.06 ± 684.72	0.928	0.399	0.177	/

Ancestral state reconstruction uncovers different ancestral lifestyles for the host-associated orders

An ancestral state reconstruction (ASR) infers the clade-specific genes that have been passed on from the

common ancestor to its descendants. Based on the annotation of these reconstructed ancestral genes, we

can infer the lifestyle of this common ancestor, pinpointing the historic origin of lifestyle differences. The ASR

of this phylogeny shows that a relatively high number of genes have been gained and lost in the vicinity of

the root node of each order (**Figure**).

796 genes appear to have been present since the clostridial last common ancestor (LCA). These are

active in several central subsystems like glycolysis, tRNA biosynthesis and the ribosomes. In addition, they

include genes for flagellar assembly, two-component signalling (2CS) systems and DNA repair systems. Hence,

the last common ancestor of the clostridial lineage appears a well-equipped free-living microbe.

Regarding the most recent common ancestors of each order, some interesting contrasts can be

observed in both the number and functional annotation of the exchanged genes (**Add. File 5**). The MRCA of

Clostridiales gained 88 genes. Only about half of these families is annotated, which are mostly associated with

metabolism, signal transduction and cell motility, aligning with a free-living lifestyle. In contrast, it lost only

19 genes, of which the annotated share is again mostly metabolism-related. Notably, the ancestor of a large

subcluster containing the *Clostridium sensu stricto* clade has gained 182 genes, which are mostly associated

with metabolism, and DNA replication and repair, indicating that this subcluster even further refined and

expanded its ancestral free-living capacities.

The *Oscillospirales* MRCA gained 42 genes and lost 38, mostly related to metabolism in both cases.

Interestingly, the direct ancestor of this node has lost a considerable number of gene families. These 143 lost

genes were involved in cell motility, cofactor metabolism and signal transduction pathways. These losses align

with the reduction in free-living capabilities that typically occurs when a microorganism transitions to a host-

associated lifestyle.

In contrast, the *Lachnospirales* MRCA has gained 62 genes and lost 40. The gained genes are

associated with amino acid and lipid metabolism, and DNA replication and repair, while the lost ones are

mostly unannotated. Also here, there is a descendant that has exchanged a considerable number of genes:

the MRCA of the *Clostridium XIVa* clade. This descendant gained 133 genes and lost 121. Gained

functionalities affect several metabolic systems, including secondary metabolite biosynthesis, while lost

functionalities are linked with cell motility and signal transduction. Both the number of exchanged genes and

the combination of gaining metabolic capabilities with losing free-living traits suggest that the *Lachnospirales*

are at a different level of host association than the *Oscillospirales*.

Lastly, the *Peptostreptococcales* ancestral lineages exchanged relatively few genes. For example, the

MRCA gained 49 genes and lost only 7. The gained genes are involved in several metabolic subsystems, such

as amino acid metabolism and glycan biosynthesis. This low level of gene exchange may indicate that the

Peptostreptococcales are at the end of the host conversion process or only have just started it.

Figure 3 Wagner parsimony gene gain (A) and loss (B) profiles for the full genome set (coloured branches), and absolute number and KEGG BRITE level B annotation profile of exchanged genes for a selection of ancestral nodes (inset pie charts).

Focused mobilome screen confirms different levels of host adaptation

Both the GD and the ASR analyses indicate varying levels of host association among the clostridial orders.

Moreover, the higher observed metabolic versatility of *Lachnospirales* may be related with its lower levels of

host association. To further support our identification of these varying stages of host adaptation, we screened

our entire genome set for the abundance of a set of genomic hallmarks tied with progressing host adaptation.

More specifically, the transition from a free-living to a host-associated lifestyle in bacteria is thought

to be coupled with a genome streamlining process driven by insertion sequences (ISes) and intrinsic

mutations (Siguier, Gourbeyre and Chandler 2014). ISes are a type of intracellular mobile genetic elements

(MGEs) that employ transposases for their mobility. When bacteria move from a nutritionally poor free-living

environment to a nutritionally rich host-associated setting, many genes lose their essentiality due to the

constant availability of metabolites. This causes a reduced selection pressure on non-essential genes,

enabling higher mutation rates as well as IS insertions into these genes. Both these phenomena drive

pseudogene formation (Marais, Calteau and Tenaillon 2008). Over time, the accumulation and multiplication

of IS regions promotes genome rearrangements and deletions, ultimately yielding smaller streamlined

genomes tailored to their new host-associated lifestyle (**Figure G**) (Siguier, Gourbeyre and Chandler 2014).

To explore this hypothesis, we screened our full clostridial genome set for the abundance of the

following MGEs and genomic features: (1) genome sizes, (2) ISes, (3) pseudogenes, (4) prophage regions, and

(5) biosynthetic gene clusters. (1), (2) and (3) are key hallmarks of the genome streamlining process. (4) may

reflect a shift in environmental niche. (5) may provide an indication of the metabolic versatility of the

organism, as the necessary precursor metabolites must be available.

The results align strikingly well with our previous interpretations. First, *Oscillospirales* and

Peptostreptococcales genomes are significantly smaller than *Clostridiales* and *Lachnospirales* (**Figure A**).

Second, *Lachnospirales* and *Peptostreptococcales* genomes harbour significantly more ISes than

Clostridiales, which, in turn, harbour more than *Oscillospirales* genomes (**Figure B**). This observation aligns

with our hypothesis that *Lachnospirales* is in the early stages of host adaptation and thus is undergoing an IS

expansion. Furthermore, the smaller genomes and lower number of ISes of the *Peptostreptococcales* indicate

that this latter group is slightly more advanced in this transition process than the *Lachnospirales*.

A deeper analysis of the IS families revealed some substantial differences in their distribution among

the species orders (**Add. File 2: Figure S6**). For example, family IS1634 is only present in *Lachnospirales* and

Peptostreptococcales, while family IS5 is only present in *Oscillospirales*. Notably, families IS110, IS66 and ISL3

are more prevalent in *Lachnospirales* species. These IS families do not have a specific insertion site preference

(Siguier, Gourbeyre and Chandler 2014, Siguier et al. 2015), so their abundances may merely be the result of

IS expansion during the host conversion process.

Third, pseudogenes are significantly less abundant in *Clostridiales* than in *Peptostreptococcales,*

which at their turn harbour less than *Oscillospirales* and *Lachnospirales.* (**Figure C**). This implies that both gut-

associated clusters have undergone or are undergoing more intense pseudogene formation.

Fourth, the number of BGCs is significantly higher in *Clostridiales* and *Peptostreptococcales*, with

Lachnospirales and *Oscillospirales* at comparable lower levels (**Figure D**). This hints that *Clostridiales* and

Peptostreptococcales species maintain a greater capacity for specialised metabolism, whereas the other two

orders seem to have eliminated a substantial share of their BGCs. Given that other genomic features indicate

that *Lachnospirales* is only in the early stages of host adaptation, it may seem that BGCs are among the first

to be eliminated under altered selection pressures despite the maintenance of non-BGC metabolic versatility.

However, this may also reflect our current knowledge gap for BGCs from gut-associated species. Regarding

the BGC classes, the most substantial differences in abundance are observed in the cyclic lactone autoinducer

(CLA), NRPS(-like), and RiPP-like classes. In addition, there is a consistent presence of ranthipeptides and

terpene precursors across all orders (**Add. File 2: Figure S7**).

Fifth, *Lachnospirales* have significantly less prophages than the other orders (**Figure E**). Regarding

infection targets, *Lachnospirales* are infected by the highest number of prophage types, while

Peptostreptococcales by the lowest. Interestingly, about 75% of the detected prophage types infect clostridial

species from multiple orders, with 39 even infecting species from all orders (**Figure F**). Besides, some

prophages carry passenger genes for methyltransferases, restriction enzymes, toxin(-antitoxin system)s and

transporters (including beta-lactam efflux pumps and other resistance proteins), highlighting a mechanism by

which clostridial species may acquire new genes (**Add. File 6**). Hence, while there is a clear difference in the

prophages present, it remains unclear whether this merely reflects a niche shift or that these prophages carry

genes that have become essential for the bacterium's survival due to the host adaptation.

To conclude, the abundances of these genomic features show again that the clostridial species orders

are in various stages of the host adaptation process (**Figure G**).

Figure 4. **Screening of the genomic hallmarks of host dependence. (A) – (F)** Comparison of the abundance of a genomic feature across all orders. Feature counts are normalised by dividing by the genome length in Mb. The applied statistical test is the Brunner-Munzel test (α = 0.05). (A) genome size, (B) insertion sequence elements, (C) pseudogenes, (D) biosynthetic gene clusters, (E) prophage regions, (F) number of infecting phages by host order. **(G) Schematical overview of the impact of the genome streamlining process on the bacterial genomic architecture**, including the inferred position of each clostridial order in this process. Figure inspired on (Siguier, Gourbeyre and Chandler 2014). (1) The (ancestral) free-living genome has only few ISes. (2) As a result of an environmental bottleneck, IS expansion occurs, leading to an increased abundance of ISes, pseudogenes and mutations. Genomic rearrangements occur. (3) With time, the repetitive regions of the ISes facilitate the deletion of non-essential genomic regions. (4) Eventually, extensive deletion leads to the elimination of the ISes. (5) Finally, the genome reaches its streamlined form, with a gene content tailored for life in its host environment. (6) Upon contact with IS-carrying strains or phages, ISes may be reintroduced.

318 *BGC clustering analysis reveals strongly conserved BGCs*

319 The species in our clostridial dataset have faced varying selection pressures. Hence, any BGC that is conserved

320 despite these varying pressures, is likely to provide potent properties that offer an evolutionary advantage

321 large enough to justify its conservation. Such high-potential BGCs should emerge in a BGC clustering analysis.

322   Our clustering revealed that the five largest BGC groups are two terpene-precursor groups and three

323 ranthipeptide groups. Both *Clostridiales* and *Lachnospirales* harbour one terpene-precursor and one or two

324 ranthipeptide BGC groups (**Figure**). Alignment of gene clusters from each of these three largest ranthipeptide

325 groups shows that the core biosynthetic genes, the SCIFF precursor and the RRE gene, are conserved across

326 all groups (**Add. File 2: Figure S8)**. Moreover, these ranthipeptide groups are well-distributed across the full

327 phylogeny, with every species coding maximum one ranthipeptide (**Add. File 2: Figure S9**). Interestingly, two

328 of these ranthipeptide BGCs have both been experimentally shown to participate in quorum sensing activities

329 (Chen et al. 2020). The fact that these two BGCs belong to the same ranthipeptide group in our analysis

330 suggests that similar quorum sensing functions may be associated with this entire ranthipeptide group.

331   The smaller identified BGC groups are cyclic lactone autoinducers and deazapurines, next to more

332 smaller ranthipeptides and terpene precursor groups. We coclustered our BGC set with the MIBiG database,

333 a highly curated and standardised repository of known BGCs (Zdouc et al. 2025). Interestingly, very few MIBiG

334 BGCs cocluster with our clostridial BGCs, underscoring the novelty of these conserved ranthipeptides and

335 terpene precursors, and their high potential for new functional significance. We also observed a large group

336 of singleton BGCs (**Add. File 7**), which may comprise BGCs acquired more recently by horizontal gene transfer

337 or gene rearrangement. While these singletons have not undergone sufficient evolutionary selection to refine

338 their functional significance, any potent and biologically relevant activity cannot be excluded.

Figure 5. BiG-SCAPE similarity networks of all BGCs detected in our full genome set (threshold 0.3), annotated (A) by clostridial order or as MIBiG entry, and (B) by BGC type. The five largest groups on the top left are either terpene precursors or ranthipeptides. Only groups with at least five members are shown.

Discussion

This work presents the first comparative pangenomics analysis of multiple orders of the *Clostridia* class and is the first to report about their intriguingly different host adaptation tendencies. Moreover, it uncovered a remarkably large repertoire of novel BGCs, including several highly conserved ranthipeptides.

A major finding of this study is the varying stages in host adaptation in each taxonomic order. By using multiple independent approaches in parallel, we were able to position each order on the host association spectrum. In particular, the early-stage host association trend of the *Lachnospirales* highlights a new important nuance of the current host-associated status of this prevalent clade of gut symbionts. Much of the genomic and metabolic diversity of this clade may have been lumped by the generic label "host-associated".

Bacteria in the early stages of host association likely still retain part of their free-living gene repertoire. Indeed,

the KEGG pathway completeness assessment has shown that the *Lachnospirales* are more likely to be

metabolically versatile. Due to their higher innate capacity to produce more complex metabolites, these

species would be excellent starting points for the development of anaerobic chassis organisms for

heterologous expression of potent BGCs. Moreover, they have a high potential as therapeutic delivery

vehicles in gut environments, as they are adapting to exactly those environments (Kubiak and Minton 2015).

Another key outcome of this study is the comprehensive identification of clostridial BGCs, providing

the first comparative view of the natural product repertoire of clostridial bacteria. Despite extensive host

adaptation and a long track record of exposure to varying selection pressures, several ranthipeptide and

terpene precursor BGCs have been widely conserved. The lack of MIBiG entries coclustering with these BGCs

underscores their novelty, tapping further into the unexplored richness of BGCs from the anaerobic realm.

These novel conserved BGCs are prime targets for further experimental characterisation.

Nevertheless, there are some limitations to be considered for this study. First, since we conducted a

high-level analysis averaging counts from species orders, the conclusions presented here are to be considered

general tendencies of each order and should not be extrapolated to individual species without further

analysis. For example, it may be possible that a *Lachnospirales* species is already highly adapted to its host

because of higher selection pressures, or because it started the host adaptation process early compared to

its close relatives. Second, we dereplicated our dataset to ensure an equal species representation in the

pangenome. Non-representative genomes may harbour even more novel diversity, both at the pangenome

level as well as at the BGC level.

Overall, the combined application of comparative genomics, phylogenetic techniques and narrow-

focus genomic feature screens at higher taxonomic levels offers a flexible approach to investigate host

adaptation trends. We uncovered counterintuitively different levels of host adaptation in clostridial species

clusters, which translate into different levels of metabolic versatility. Taken together, the findings of this study

are an encouraging starter for the systematic investigation of adaptation levels in other "host-associated"

bacteria classes next to *Clostridia*. Moreover, in the light of natural product research, our findings pave the

way for (1) establishing clostridial species as new anaerobic host organisms for a diverse range of applications,

as well as for (2) expanding the natural product repertoire further into the anaerobic realm.

Data availability

377   All genome assemblies analysed in this study were retrieved from NCBI RefSeq. Accession codes and relevant

378   metadata can be found in Add. File 1.

Code availability

380   The full analysis pipeline has been deposited at GitHub (https://github.com/LucoDevro/closHomer) and at

381   Zenodo (https://doi.org/10.5281/zenodo.14719962).

Author contribution statement

383   LDV and JM conceived this study. LDV and JM designed the research. LDV performed the data analysis,

384   implemented the workflow, and gathered the results. LDV analysed and interpreted the data, with assistance

385   from JA and JM. LDV wrote the article, which was revised by LDV, JA and JM.

Acknowledgements

387   We thank Jan Aerts for his helpful advice on improving our data visualisations.

Funding statement

389   This work was funded by the European Commission through the European Research Council (MiStiC,

390   101078461).

Competing interest statement

392   We declare no competing interests.

References

Asnicar, Francesco, Thomas, Andrew Maltez, Beghini, Francesco, Mengoni, Claudia, Manara, Serena,
Manghi, Paolo, et al., 'Precise Phylogenetic Analysis of Microbial Isolates and Genomes from
Metagenomes Using PhyloPhlAn 3.0', *Nature Communications*, 11/1 (2020), 2500
Behnken, Swantje, and Hertweck, Christian, 'Anaerobic Bacteria as Producers of Antibiotics', *Applied
Microbiology and Biotechnology*, 96/1 (2012), 61–67
Birgen, Cansu, Degnes, Kristin F., Markussen, Sidsel, Wentzel, Alexander, and Sletta, Håvard, 'Butanol
Production from Lignocellulosic Sugars by Clostridium Beijerinckii in Microbioreactors', *Biotechnology
for Biofuels*, 14/1 (2021)
Blin, Kai, Shaw, Simon, Vader, Lisa, Szenei, Judit, Reitz, Zachary L, Augustijn, Hannah E, et al., 'AntiSMASH
8.0: Extended Gene Cluster Detection Capabilities and Analyses of Chemistry, Enzymology, and
Regulation', *Nucleic Acids Research*, 2025
Buchfink, Benjamin, Reuter, Klaus, and Drost, Hajk-Georg, 'Sensitive Protein Alignments at Tree-of-Life Scale
Using DIAMOND', *Nature Methods*, 18/4 (2021), 366–68

Cantalapiedra, Carlos P, Hernández-Plaza, Ana, Letunic, Ivica, Bork, Peer, and Huerta-Cepas, Jaime, 'EggNOG-Mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale', *Molecular Biology and Evolution*, 38/12 (2021), 5825–29

Capella-Gutiérrez, Salvador, Silla-Martínez, José M., and Gabaldón, Toni, 'TrimAl: A Tool for Automated Alignment Trimming in Large-Scale Phylogenetic Analyses', *Bioinformatics*, 25/15 (2009), 1972–73

Chaumeil, Pierre-Alain, Mussig, Aaron J, Hugenholtz, Philip, and Parks, Donovan H, 'GTDB-Tk v2: Memory Friendly Classification with the Genome Taxonomy Database', *Bioinformatics*, 38/23 (2022), 5315–16

Chen, Yunliang, Yang, Yunpeng, Ji, Xinjian, Zhao, Ran, Li, Guoquan, Gu, Yang, et al., 'The SCIFF-Derived Ranthipeptides Participate in Quorum Sensing in Solventogenic Clostridia', *Biotechnology Journal*, 15/10 (2020)

Cohen, Jonathan E., Wang, Rong, Shen, Rong-Fong, Wu, Wells W., and Keller, James E., 'Comparative Pathogenomics of Clostridium Tetani', *PLOS ONE*, 12/8 (2017), e0182909

Cruz-Morales, Pablo, Orellana, Camila A, Moutafis, George, Moonen, Glenn, Rincon, Gonzalo, Nielsen, Lars K, et al., 'Revisiting the Evolution and Taxonomy of Clostridia, a Phylogenomic Update', *Genome Biology and Evolution*, 11/7 (2019), 2035–44

Csűrös, Miklós, 'Count: Evolutionary Analysis of Phylogenetic Profiles with Parsimony and Likelihood', *Bioinformatics*, 26/15 (2010), 1910–12

Dufresne, Alexis, Garczarek, Laurence, and Partensky, Frédéric, 'Accelerated Evolution Associated with Genome Reduction in a Free-Living Prokaryote', *Genome Biology*, 6/2 (2005), R14

Gautreau, Guillaume, Bazin, Adelme, Gachet, Mathieu, Planel, Rémi, Burlot, Laura, Dubois, Mathieu, et al., 'PPanGGOLiN: Depicting Microbial Diversity via a Partitioned Pangenome Graph', *PLOS Computational Biology*, 16/3 (2020), e1007732

Gilchrist, Cameron L.M., and Chooi, Yit Heng, 'Clinker & Clustermap.Js: Automatic Generation of Gene Cluster Comparison Figures', *Bioinformatics*, 37/16 (2021), 2473–75

Katoh, K., and Standley, D. M., 'MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability', *Molecular Biology and Evolution*, 30/4 (2013), 772–80

Kubiak, Aleksandra M., and Minton, Nigel P., 'The Potential of Clostridial Spores as Therapeutic Delivery Vehicles in Tumour Therapy', *Research in Microbiology*, 166/4 (2015), 244–54

Letunic, Ivica, and Bork, Peer, 'Interactive Tree of Life (ITOL) v6: Recent Updates to the Phylogenetic Tree Display and Annotation Tool', *Nucleic Acids Research*, 52/W1 (2024), W78–82

Letzel, Anne Catrin, Pidot, Sacha J., and Hertweck, Christian, 'A Genomic Approach to the Cryptic Secondary Metabolome of the Anaerobic World', *Natural Product Reports*, 30/3 (2013), 392–428

Marais, Gabriel A.B., Calteau, Alexandra, and Tenaillon, Olivier, 'Mutation Rate and Genome Reduction in Endosymbiotic and Free-Living Bacteria', *Genetica*, 134/2 (2008), 205–10

Maturana, José Luis, and Cárdenas, Juan P., 'Insights on the Evolutionary Genomics of the Blautia Genus: Potential New Species and Genetic Content Among Lineages', *Frontiers in Microbiology*, 12 (2021)

Minh, B. Q., Nguyen, M. A. T., and von Haeseler, A., 'Ultrafast Approximation for Phylogenetic Bootstrap', *Molecular Biology and Evolution*, 30/5 (2013), 1188–95

Minh, Bui Quang, Schmidt, Heiko A, Chernomor, Olga, Schrempf, Dominik, Woodhams, Michael D, von Haeseler, Arndt, et al., 'IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era', *Molecular Biology and Evolution*, 37/5 (2020), 1530–34

Navarro-Muñoz, Jorge C., Selem-Mojica, Nelly, Mullowney, Michael W., Kautsar, Satria A., Tryon, James H., Parkinson, Elizabeth I., et al., 'A Computational Framework to Explore Large-Scale Biosynthetic Diversity', *Nature Chemical Biology*, 16/1 (2020), 60–68

O'Leary, Nuala A., Cox, Eric, Holmes, J. Bradley, Anderson, W. Ray, Falk, Robert, Hem, Vichet, et al., 'Exploring and Retrieving Sequence and Metadata for Species across the Tree of Life with NCBI Datasets', *Scientific Data*, 11/1 (2024), 732

Pahalagedara, Amila Srilal Nawarathna Weligala, Flint, Steve, Palmer, Jon, Brightwell, Gale, and Gupta, Tanushree Barua, 'Antimicrobial Production by Strictly Anaerobic Clostridium Spp.', *International Journal of Antimicrobial Agents*, 55/5 (2020)

Pascal Andreu, Victòria, Augustijn, Hannah E., Chen, Lianmin, Zhernakova, Alexandra, Fu, Jingyuan,
Fischbach, Michael A., et al., 'GutSMASH Predicts Specialized Primary Metabolic Pathways from the
Human Gut Microbiota', *Nature Biotechnology*, 41/10 (2023), 1416–23

Patakova, Petra, Branska, Barbora, Sedlar, Karel, Vasylkivska, Maryna, Jureckova, Katerina, Kolek, Jan, et al.,
'Acidogenesis, Solventogenesis, Metabolic Stress Response and Life Cycle Changes in Clostridium
Beijerinckii NRRL B-598 at the Transcriptomic Level', *Scientific Reports*, 9/1 (2019)

Roberts, Seth B, Gowen, Christopher M, Brooks, J Paul, and Fong, Stephen S, 'Genome-Scale Metabolic
Analysis of Clostridium Thermocellum for Bioethanol Production', *BMC Systems Biology*, 4/1 (2010), 31
<https://bmcsystbiol.biomedcentral.com/articles/10.1186/1752-0509-4-31>

Ruiz-Perez, Carlos A., Conrad, Roth E., and Konstantinidis, Konstantinos T., 'MicrobeAnnotator: A User-
Friendly, Comprehensive Functional Annotation Pipeline for Microbial Genomes', *BMC Bioinformatics*,
22/1 (2021), 11

Salamzade, Rauf, and Kalan, Lindsay R., 'SkDER: Microbial Genome Dereplication Approaches for
Comparative and Metagenomic Applications', *BioRxiv*, 2023

Sandhu, Brindar K., and McBride, Shonna M., 'Clostridioides Difficile', *Trends in Microbiology*, 26/12 (2018),
1049–50

Schwengers, Oliver, Jelonek, Lukas, Dieckmann, Marius Alfred, Beyvers, Sebastian, Blom, Jochen, and
Goesmann, Alexander, 'Bakta: Rapid and Standardized Annotation of Bacterial Genomes via
Alignment-Free Sequence Identification', *Microbial Genomics*, 7/11 (2021)

Siguier, Patricia, Gourbeyre, Edith, and Chandler, Mick, 'Bacterial Insertion Sequences: Their Genomic
Impact and Diversity', *FEMS Microbiology Reviews*, 38/5 (2014), 865–91

Siguier, Patricia, Gourbeyre, Edith, Varani, Alessandro, Ton-Hoang, Bao, and Chandler, Mick, 'Everyman's
Guide to Bacterial Insertion Sequences', *Microbiology Spectrum*, 3/2 (2015)

Smith, Theresa J., Hill, Karen K., and Raphael, Brian H., 'Historical and Current Perspectives on Clostridium
Botulinum Diversity', *Research in Microbiology*, 166/4 (2015), 290–302

Sugimoto, Yuki, Camacho, Francine R., Wang, Shuo, Chankhamjon, Pranatchareeya, Odabas, Arman, Biswas,
Abhishek, et al., 'A Metagenomic Strategy for Harnessing the Chemical Repertoire of the Human
Microbiome', *Science*, 366/6471 (2019)

Syberg-Olsen, Mitchell J, Garber, Arkadiy I, Keeling, Patrick J, McCutcheon, John P, and Husnik, Filip,
'Pseudofinder: Detection of Pseudogenes in Prokaryotic Genomes', *Molecular Biology and Evolution*,
39/7 (2022)

Tonkin-Hill, Gerry, Gladstone, Rebecca A., Pöntinen, Anna K., Arredondo-Alonso, Sergio, Bentley, Stephen D.,
and Corander, Jukka, 'Robust Analysis of Prokaryotic Pangenome Gene Gain and Loss Rates with
Panstripe', *Genome Research*, 33/1 (2023), 129–40

Udaondo, Zulema, Duque, Estrella, and Ramos, Juan Luis, 'The Pangenome of the Genus Clostridium',
Environmental Microbiology, 19/7 (2017), 2588–2603

Vieira-Silva, Sara, Falony, Gwen, Darzi, Youssef, Lima-Mendez, Gipsi, Garcia Yunta, Roberto, Okuda, Shujiro,
et al., 'Species-Function Relationships Shape Ecological Properties of the Human Gut Microbiome',
Nature Microbiology, 1/8 (2016)

Wishart, David S, Han, Scott, Saha, Sukanta, Oler, Eponine, Peters, Harrison, Grant, Jason R, et al.,
'PHASTEST: Faster than PHASTER, Better than PHAST', *Nucleic Acids Research*, 51/W1 (2023), W443–50

Xie, Zhiqun, and Tang, Haixu, 'ISEScan: Automated Identification of Insertion Sequence Elements in
Prokaryotic Genomes', *Bioinformatics*, 33/21 (2017), 3340–47

Yu, Guangchuang, Smith, David K., Zhu, Huachen, Guan, Yi, and Lam, Tommy Tsan-Yuk, 'Ggtree: An R
Package for Visualization and Annotation of Phylogenetic Trees with Their Covariates and Other
Associated Data', *Methods in Ecology and Evolution*, 8/1 (2017), 28–36

Zdouc, Mitja M, Blin, Kai, Louwen, Nico L L, Navarro, Jorge, Loureiro, Catarina, Bader, Chantal D, et al.,
'MIBiG 4.0: Advancing Biosynthetic Gene Cluster Curation through Global Collaboration', *Nucleic Acids
Research*, 53/D1 (2025), D678–90