Title
Alphavirus replicons encoding IFN-γ enhance cancer virotherapy by overcoming macrophage-
mediated suppression

Authors and affiliations

Laura Horvathova[1]*, Priscilla Kinderman[2]*, Thijs Janzen[3], Baukje Nynke Hoogeboom[1], Franz J.
Weissing[3], Nadine van Montfoort[2], Toos Daemen[1], Darshak K. Bhatt[1,4]

1. Department of Medical Microbiology and Infection Prevention, University Medical Center
Groningen, University of Groningen, Groningen, The Netherlands
2. Department of Gastroenterology and Hepatology, Leiden University Medical Center, Leiden, The
Netherlands
3. Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The
Netherlands
4. Lead contact: Darshak K. Bhatt (d.bhatt@umcg.nl)

* Equal contributions

Summary

Interference by tumor-associated macrophages may significantly reduce the efficacy of therapeutic
viruses designed to infect cancer cells and activate antitumor T-cells. Using a computational model,
we hypothesized that viruses encoding a T cell-stimulating signal, like IFN-γ, could overcome this
barrier. We engineered an alphavirus-based replicon expressing IFN-γ and evaluated its effect in
various human-derived tumor-immune coculture systems and an *in vivo* murine model. While
alphavirus replicons do not replicate in macrophages, macrophages acted as a barrier, limiting tumor
infection in a frequency-dependent but phenotype-independent manner. Nonetheless, T-cell
activation occurred even when only a fraction of infected tumor cells expressed IFN-γ, regardless of
macrophage presence, frequency, or phenotype. Additionally, viral stimulation drove macrophage-
repolarization towards a pro-inflammatory phenotype favoring T-cell activation. These findings
highlight a strategy for optimizing virotherapy in macrophage-rich tumors by designing viruses that
stimulate T-cell activation, ensuring therapeutic efficacy.

Keywords
alphavirus replicons, tumor-associated macrophages, therapeutic resistance, computational model,
tumor-immune coculture

Introduction

Tumor-associated macrophages are a predominant component of the tumor immune infiltrate across many cancers[1]. These macrophages exhibit context-dependent phenotypic plasticity[2,3] and function to either support or suppress antitumor immunity[2–5]. In particular, they perform phagocytosis, act as antigen-presenting cells, express checkpoint regulators, and secrete an array of cytokines that shape cancer progression and response to immunotherapy[4,6,7].

Cancer virotherapy is a form of immunotherapy where viruses are used to infect cancer cells, and subsequently boost anticancer T cell responses[8]. Virus-induced cancer cell death causes the release of danger signals and antigens in the tumor microenvironment[9]. Ideally, these signals are immunogenic, enhancing the recruitment and proinflammatory activation of immune cells into the tumor. However, tumor-associated macrophages may hinder the therapeutic efficacy of virotherapy by capturing virus particles, thereby significantly reducing the availability of the virus for infecting cancer cells[10–13]. This effect could be particularly pronounced when macrophages are non-permissive for viral replication, as it restricts tumor cell infection and the subsequent release of immunostimulatory signals and tumor antigens, thereby indirectly suppressing downstream anticancer T-cell activity. Tumor-associated macrophages further limit oncolysis through innate antiviral signaling[10–13] and T-cell suppression by regulatory mechanisms[3,14]. Variation in tumor composition[15,16], particularly in terms of macrophage frequency and phenotype can lead to variability in therapeutic responses[17–19]. Enhancing T-cell-mediated antitumor responses using virotherapy thus remains challenging, especially in macrophage-rich tumors. Therefore, development of therapeutic viruses capable of strongly stimulating T-cell immune responses is crucial to overcome the limitations posed by non-permissive macrophages.

Semliki Forest virus (SFV)-based replicons are emerging as promising candidates for immunogenic virotherapy. SFV is a positive-stranded RNA virus, belonging to the *Alphavirus* genus. The RNA genome of SFV functions as a replicon, encoding non-structural viral proteins capable of viral-RNA translation and replication. Previous efforts in developing recombinant SFV particles (rSFV) have focused on improving safety by deleting viral genes that code for structural proteins[20]. This creates suicidal virus particles capable of a single round of infection. We and others have demonstrated that rSFV particles can successfully infect and express viral genes in a wide range of cancer cells and healthy stromal cells, however, macrophages are non-permissive to SFV infections and virus-encoded protein translation[21–23]. Furthermore, the immunogenicity of rSFV has been improved by encoding antigens[24–27], cytokines[22,28], or antibodies[29] to boost humoral and cellular immune responses in the context of cancer therapy and vaccination against infectious diseases. Our group has demonstrated phase-1 and phase 2 clinical safety, immunogenic potential and clinical efficacy of an rSFV-based therapeutic vaccine encoding antigens of human papillomavirus in patients with cervical cancer[30,31]. So far, studies using rSFV as an anticancer agent have only been conducted in murine models[22,28,29]. Recently our group showed that rSFV can be engineered to express immunogenic human cytokines or chemokines with an enhanced potential to recruit and activate T-cells in different human cancer models[23].

In this study, we implemented a combined theoretical and experimental approach to design an immunogenic-rSFV therapy capable of T-cell activation and macrophage polarization. First, we evaluated whether indeed macrophages limit rSFV infection in cancer cells growing in monolayer (2D, two-dimensional) or spheroid-based (3D, three-dimensional) tumor-macrophage cocultures. Second, we employed a computational model to assess whether rSFV-encoding immunogenic signals can promote anticancer T-cell responses and tumor eradication despite the presence of macrophages. Third, we analyzed literature to identify interferon-gamma (IFN-γ) as a pro-

inflammatory cytokine capable of both macrophage and T-cell activation. We further evaluated the
correlation between intra-tumoral IFN-γ signature and activation of macrophages and T-cells using
data of cervical and pancreatic cancer patients available from The Cancer Genome Atlas (TCGA)[1]. For
experimental validation, we engineered rSFV to express IFN-γ upon infection of cancer cells and
employed both monolayer and spheroid-based tumor-immune cocultures to evaluate T-cell
activation. In these tumor-immune cocultures, we introduced macrophages of either a naïve (M$_{naive}$)
or an IL-4-induced phenotype (M$_{IL4}$). Finally in a proof-of-concept study, we tested the efficacy of
rSFV in a pancreatic cancer mouse model. These models allowed us to assess how tumor infection
and T-cell activation are influenced by macrophages and to evaluate the immunogenic potential of
virus-encoded IFN-γ.

Results

Effect of macrophage frequency and phenotype on rSFV-mediated tumor infection

As a proof-of-concept, we performed all analyses in the context of two independent solid tumor
types, i.e. cervical and pancreatic cancer, with innate differences in immune signaling
(**Supplementary Figure 1**). rSFV-mediated infection of cancer cells leads to expression of encoded
transgenes but not production of progeny virus particles (as illustrated in **Figure 1A**). To validate our
findings in an experimental setup, we employed an *in vitro* cancer-macrophage coculture model in
either a monolayer or spheroid-based spatial organization (**Figure 1B**). This allowed us to have
control over factors such as the frequency and phenotype of macrophages and cancer cells. Using
real-time microscopy-based imaging, we studied the effect of macrophages in regulating rSFV
infection of a pancreatic cancer cell line (PANC-1). Figure 1C-D shows microscopy images resulting
from the spheroid (**Figure 1C**) or monolayer (**Figure 1D**) coculture setup consisting of cancer cells and
a varying frequency of macrophages. With an increasing frequency of macrophages (stained red in
the images) present in the coculture, we observed a decrease in virus infection (stained green,
expressing virus-encoded green fluorescent protein, GFP). We performed the cancer-macrophage
coculture in various setups, either varying only the frequency of macrophages (green bar) or the
frequencies of both macrophages and cancer cells (purple bar) (**Figure 1E**). In particular, we
confirmed[21,32] that virus-encoded GFP expression is restricted to cancer cells but not macrophages.
The decrease in the number of infected cells corresponds to a decrease in the absolute number of
target cancer cells and to a relative increase in the number of macrophages (**Figure 1F-G**), indicating
that presence of macrophages limits the effect of virotherapy.

We used non-polarized macrophages for the experiments presented in Figure 1. Since tumor-
associated macrophages can exhibit diverse phenotypic states, we specifically addressed the
contribution of two phenotypes most frequently found in the tumor microenvironment. For this, we
polarized them to either a classically activated anti-tumoral phenotype (M$_{class}$ or M1-like) or an
alternatively activated pro-tumoral phenotype (M$_{alter}$ or M2-like) through cytokine stimulation
(**Figure 2A**). Here, we used either a combination of LPS and IFN-γ to generate classically activated
macrophages (M$_{LPS+IFNγ}$), or IL-4 to generate alternatively activated macrophages (M$_{IL4}$). Pro-
inflammatory cell surface marker proteins such as CD80 and CD86 are upregulated in classically
activated M$_{LPS+IFNγ}$ macrophages, whereas regulatory markers like CD206 are more abundant in
alternatively activated M$_{IL4}$ macrophages (**Figure 2B**) also corresponding to a distinct cellular state
(**Figure 2C**). We then assessed if the macrophage phenotype, in particular the alternatively activated
M$_{IL4}$ phenotype that is most frequently found in tumors, influenced tumor infection by rSFV in
monolayer or spheroid cocultures (**Figure 2D-E**). We observed that the number of GFP+ cells reduced

with an increase in the number of macrophages, however, independent of the macrophage
phenotype (**Figure 2F-G**), as there were no differences observed in the magnitude of reduction in
GFP+ cancer cells between conditions that had either $M_{Naïve}$ or M_{IL4} polarized macrophages. This was
also the case when macrophages were cocultured with a cervical cancer cell line (Ca-Ski) (**Figure 2H**).

Notably, cytokine-driven macrophage polarization had no impact on the expression of genes
encoding SFV-entry receptors (**Supplementary Figure 2**). Furthermore, blocking JAK-STAT signaling to
examine the role of type-I interferon in antiviral effects failed to improve rSFV infection in cancer
cells (**Supplementary Figure 3**). Thus, the frequency of non-permissive macrophages, rather than
antiviral signaling or their polarized phenotype, primarily contributed to the observed reduction in
infection.

Model predictions regarding the effectiveness of immunogenic-rSFV therapy

In a previous study, we developed a spatiotemporal model to assess the effect of anticancer T-cell
responses in response to immunogenic signals released upon virotherapy[33,34]. **Figure 3A** (left image)
depicts a snapshot of a computer simulation with a tumor containing uninfected cancer cells, virus-
infected cancer cells, and macrophages. Upon infection-caused cell death, the model assumes that
virus-induced immunogenic signals are released in the neighborhood (as depicted in **Figure 3A** right
and illustrated in **Figure 3B**) stimulating anticancer T-cell response. Using this model, we assessed the
efficacy of rSFV encoding immunogenic signals as a non-replicating suicidal virotherapy. To estimate
therapeutic success, we assessed the probability of tumor eradication caused by rSFV-infection and
anticancer T-cell cytotoxicity. Panels **C** to **E** show, for five scenarios regarding the percentage of
initially infected tumor cells (0-50%), how the probability of tumor eradication is related to the IFN-γ
level produced per infected cell (**Figure 3C**), the degree of T-cell cytotoxicity (**Figure 3D**), and the
frequency of macrophages at the time of rSFV therapy (**Figure 3E**). Generally, 30% initially infected
cells are sufficient to promote tumor eradication even at low levels of immunogenic signal
production. A minimum T-cell cytotoxicity level of 3 (corresponding to 3 target cells killed per
cytotoxic T-cell per day) is required for an effective therapeutic outcome, as increasing numbers do
not correspond with increased tumor eradication. Therapeutic outcome was found to be mainly
determined by the percentage of initially infected cells producing immunogenic signals and not by
the macrophage frequency. To guide the selection of effective immunogenic signals for rSFV-based
therapy, we next consulted the literature to identify candidates with strong T-cell-activating
potential.

The association between IFN-γ and macrophage-T-cell activation

We analyzed current literature to screen potential immunomodulatory cytokines produced and/or
consumed by T-cells and macrophages. By re-evaluating previously published data[35], we found that
IFN-γ produced by T-cells represents a unique antitumoral cytokine that has the potential to
stimulate both macrophages and T-cells through JAK-STAT signaling pathway (**Figure 4A**) with an
effective concentration ranging from 3 to 40 pM (**Figure 4B**).

Based on cervical and pancreatic cancer patient data from TCGA[1,36], we analyzed the correlation
between the IFN-γ signature in tumor samples with the frequency of macrophages and T-cells and
the phenotype of macrophages (**Figure 4C**). The IFN-γ signature applied here was derived from the
TCGA pan-cancer immunogenomic analysis[1,36]. **Figure 4D** shows an overview of the correlation
between the intratumoral IFN-γ signature and different phenotypes of macrophages and cytotoxic
CD8 T-cells. The IFN-γ signature was found to positively correlate with a classically activated (M_{class} or
M1-like) pro-inflammatory macrophage phenotype in both cervical (CESC) and pancreatic (PAAD)
tumor samples (**Figure 4E, 4G**) but not with an alternatively activated (M_{alter} or M2-like)

immunosuppressive macrophage phenotype (**Figure 4F, 4H**). As T-cells are one of the primary cells
producing IFN-γ, we analyzed if there was a correlation between intra-tumoral CD8 T-cells and a
particular macrophage phenotype (**Figure 4I**). We observed a positive correlation between classically
activated (M_{class}) proinflammatory macrophages and the frequency of intra-tumoral CD8 T-cells
(**Figure 4J, 4L**). Inversely, there was a negative correlation between alternatively activated (M_{alter})
immunosuppressive macrophages and CD8 T-cell frequency (**Figure 4K, 4M**). These findings highlight
a link between IFN-γ–driven macrophage polarization and CD8 T-cell infiltration, prompting further
investigation into whether rSFV-delivered IFN-γ could enhance this interaction in tumor–immune
cocultures.

T cell activation by rSFV in the presence of macrophages

We treated monolayer cocultures of macrophages and cancer cells with rSFV and added peripheral
blood mononuclear cells (PBMCs) to the culture to introduce T-cells and evaluate their activation
(**Figure 5A**). Through flow cytometry, we quantified expression of immune activation (CD69) and
cytotoxic-degranulation (CD107a) markers as a proxy of CD4 (helper) and CD8 (cytotoxic) T-cell
activation. These markers were chosen from a broader panel of T-cell activation and regulation
markers (**Supplementary Figure 4,5**). T-cells were considered fully activated when simultaneously
positive for CD38, CD69, and CD107a expression. **Figure 5B and 5C** illustrate the population
distribution of activated CD4 and CD8 T-cells in different scenarios of rSFV infection and tumor-
immune cocultures consisting of either PANC-1 cells or Ca-Ski cells in culture with M_{naive} or M_{IL4}
macrophages in different frequencies. The upper right quadrant in each flow cytometry panel
indicates the population of fully activated T-cells as characterized by the expression of CD69 and
CD107a.

We observed a decrease in CD4 but not CD8 T-cell activation as a result of coculturing PBMCs solely
with either M_{naive} or M_{IL4} macrophages (black circle/line, **Figure 5D**, 6E leftmost panel). PBMC
coculture with cancer cells alone or in the presence of increasing frequency of either M_{naive} or M_{IL4}
macrophages also led to a similar decrease in CD4 T-cell activation when compared to PBMCs alone
(NT = no treatment, black circle/line, **Figure 5D**). rSFV particles themselves in the absence of target
cancer cells did not have any immunomodulatory effects on CD4 or CD8 T-cell activation directly
(green circle/line, **Figure 5D, 5E**, leftmost panel). Upon rSFV-GFP therapy, i.e. not encoding IFN-γ,
CD8 T-cell activation was only observed in the case of infecting Ca-Ski cells but not PANC-1 cells and
was independent of macrophage presence or phenotype (green circle/line, **Figure 5E**, rightmost
panel). Upon rSFV-IFN-γ (rSFV-encoding IFN-γ) treatment, however, both CD4 and CD8 T-cells were
activated independently of infecting Ca-Ski cells or PANC-1 cells (red circle/line, **Figure 5D, 5E**). Here,
CD4 T-cell activation increased with an increasing number of either M_{naive} or M_{IL4} macrophages
present in the coculture (**Figure 5D**). We confirmed that rSFV-encoded IFN-γ mediates T cell
activation by inhibiting JAK-STAT signaling in T cells. Treatment with ruxolitinib, in combination with
rSFV-IFN-γ, resulted in a reduced frequency of activated T cells as compared to when treated with
rSFV-IFN-γ alone (**Supplementary Figure 6**).

Next, we assessed T-cell activation in spheroid-based tumor-immune cocultures (**Figure 6A**). In
contrast to the results from the monolayer-based coculture system, we observed a decrease in the
number of activated CD4 T-cells with an increasing frequency of macrophages in the non-rSFV-
treated coculture (black circle/line, **Figure 6B**). However, similar to the results from the monolayer
coculture, we observed that rSFV-GFP particles only induced CD8 T-cell activation in the case of
infecting Ca-Ski cells but not PANC-1 cells (green circle/line, **Figure 6C**, rightmost panels). Moreover,
rSFV-IFN-γ also led to a higher frequency of activated CD4 and CD8 T-cells independent of whether it

infected Ca-Ski cells or PANC-1 cells (red circle/line, **Figure 6B, 6C**). Here, CD4 and CD8 T-cell
activation even improved further with an increasing frequency of either M_{naive} or M_{IL4} macrophages.

Multimodal activation of macrophages by rSFV-IFN-γ

Recently our group has shown that rSFV replicon virus particles have a direct influence on
macrophage phenotype and can lead to their pro-inflammatory activation, in part by JAK-STAT and
NF-κB signaling, independent of their initial state[32]. Consequently, we evaluated if macrophages are
also stimulated by infected cancer cells alone and if there is any additional effect in combination with
virus-mediated activation (**Figure 7A**). To quantify this response, we assessed the change in
expression of costimulatory molecules (CD80, CD86) on macrophages involved in T-cell regulation in
addition to CD206 expression that is often associated with a regulatory phenotype and poor cancer
prognosis. IL-4 polarized (M_{IL4}) regulatory macrophages were found to upregulate CD86 and CD80
cell-surface expression upon stimulation with either only rSFV particles (**Figure 7B**) or only PANC-1
infected cells expressing virus-encoded IFN-γ (**Figure 7C**) or both (**Figure 7D**). The upregulation of
CD80 and CD86 was the highest when macrophages received simultaneous stimulation by virus
particles and infected cells producing virus-encoded IFN-γ. Importantly, significant downregulation of
cell-surface CD206 expression was observed when macrophages were stimulated with infected cells
producing virus-encoded IFN-γ but not GFP. Notably, IL-4 polarized macrophages upregulated the T-
cell inhibitory ligand PD-L1 in the presence of cancer cells infected with rSFV-IFN-γ (**Supplementary
Figure 7**). This upregulation is likely driven by canonical IFN-γ signaling pathways. However it is
important to highlight that this phenomenon did not coincide with a reduction of T-cell activation
phenotype as shown in **Figure 5** and **6**.

We further confirmed that both naïve (M_{naive}) and regulatory (M_{IL4}) macrophages upregulate CD80
and CD86 cell-surface expression upon stimulation with rSFV replicon particles (**Figure 7F**).
Moreover, CD206 downregulation and a strong CD80 and CD86 upregulation were observed when
either of the macrophage types was stimulated simultaneously with rSFV particles and infected
cancer cells expressing virus-encoded IFN-γ (**Figure 7G-J**). This observation holds true for both PANC-
1 (panel **G-H**) and Ca-Ski (**panel I-J**) cancer cell lines.

Additionally, we confirmed that rSFV-encoded IFN-γ mediates macrophage activation through the
JAK-STAT pathway by evaluating macrophage phenotypes in the presence of the JAK inhibitor
ruxolitinib. We observed that rSFV-IFN-γ, but not rSFV-Flt3L (control stimuli for myeloid cell
activation) or rSFV-GFP mediated, infection of cancer cells led macrophage stimulation as measured
by increased expression of co-stimulatory, anti-tumoral markers (CD80, CD86) and reduced levels of
pro-tumoral markers (CD206). However, treatment with ruxolitinib attenuated these changes,
indicating that IFN-γ-driven macrophage activation is at least partially dependent on JAK-STAT
signaling (**Supplementary Figure 7**).

Assessing the in vivo efficacy of rSFV-IFN-γ

Finally in a proof-of-concept experiment, we studied the effect of intratumoral administration of
rSFV-IFN-γ on tumor growth and the tumor immune microenvironment (TME) in mice bearing
KPC3[56–58] pancreatic tumors (**Figure 8**). As demonstrated in (**Supplementary Figure 8**), these tumors
are rich in macrophages. When tumors were palpable, mice received three intratumoral injections of
PBS, rSFV-GFP, or rSFV-IFN-γ at doses of 10^7 or 10^8 particles on days 12, 13 and 14 post-tumor
implantation (illustrated in **Figure 8A**), Seven days after the last injection, the immune cells in tumor
and spleen were analyzed by flow cytometry. Treatment with rSFV-IFN-γ at both doses markedly
showed a reduction in tumor size compared to PBS and rSFV-GFP, with the highest efficacy observed
at the 10^8 dose (**Figure 8B, Supplementary Figure 9**). Notably, the higher dose of rSFV-GFP (10^8) also

led to a measurable reduction in tumor size compared to PBS, indicating that replicon treatment
alone can elicit some degree of anti-tumor activity.

Flow cytometric analysis of tumors (**Figure 8C-N**) collected 7 days after the final injection revealed
enhanced infiltration of innate and adaptive immune cells in the rSFV-IFN-γ–treated groups.
Importantly, while total macrophage frequency **Figure 8C**) was not different between groups, a clear
shift in macrophage polarization was observed: pro-tumoral CD206$^+$ macrophages were reduced
significantly (**Figure 8D, Supplementary Figure 9B**), and anti-tumoral MHCII$^+$ macrophages were
enriched (**Figure 8E**) in the rSFV-IFN-γ–treated tumors. These changes were dose- and stimuli
dependent and most pronounced at the higher IFN-y dose. Furthermore, increased frequencies of NK
cells, and CD8$^+$ T-cell were found in the rSFV-IFN-γ groups compared to controls (**Figure 8F, I and L**).
NK cells and CD8+ T cells expressed more markers associated with recent antigen exposure, including
CD69, CD44 and PD-1 across all rSFV-treated groups, compared to PBS-treated mice (**Figure 8 G, J, K,**
N).

Second, the systemic effects were analyzed in the spleen. This analysis revealed an increased depot
of CD3$^+$ T-cells (Figure 8O) in mice treated specifically by the rSFV-IFN-y virus, whereas general rSFV
treatment showed increased numbers of CD44+ CD4+ and CD8+ T-cell subsets (**Figure 8Q and 8R**).
These data indicate that intratumoral rSFV treatment not only modified the local immune subsets in
the TME, but also activated T-cells systemically.

Collectively, our data demonstrates that intratumoral administration of rSFV-IFN-γ can slow down
tumor growth and modulate the TME of immune-suppressive tumors towards a more immune-
infiltrated phenotype.

Discussion

In this study, we investigated how macrophages impact the therapeutic efficacy of rSFV-based
anticancer virotherapy. Our findings show that macrophages, regardless of their phenotype, do not
translate rSFV-encoded proteins[37] and simultaneously limit tumor cell infection. This highlights a
significant barrier to effective virotherapy, as the presence of macrophages can limit the subsequent
immunogenic activity of the virus, potentially reducing the overall therapeutic outcome[17,33]. Notably,
tumor-associated macrophages often adopt immunosuppressive phenotypes that support tumor
progression and contribute to an overall inhibitory tumor microenvironment. Leveraging a
computational modelling approach, we hypothesized that arming therapeutic viruses to express
immunogenic signals like IFN-γ could stimulate robust T-cell activation and re-polarize macrophages,
tipping the balance towards tumor eradication. This was experimentally validated in both *in vitro*
human-based tumor-immune cocultures and an *in vivo* murine tumor model. Overall, our results
demonstrate the potential of a combined theoretical and experimental approach in improving the
immunogenic potential of anticancer virotherapy.

Our computational modeling results provided insights into the spatiotemporal dynamics of rSFV-IFN-
γ therapy and its potential despite an abundant presence of macrophages. Our model demonstrated
that the predicted release of IFN-γ at effective concentrations is vital for achieving robust T-cell
activation and subsequent tumor eradication. Specifically, the model underscored the importance of
achieving the effective concentration threshold via either increasing the number of infected cells in
the tumor or by improving IFN-γ production by individual infected cells; in both cases ensuring
activation of anticancer cytotoxic T-cells to mount a strong antitumor response. The model also
highlighted the importance of T-cell cytotoxicity in the success of rSFV-IFN-γ therapy. A minimally

required cytotoxicity rate of three target cells killed per day by each cytotoxic T-cell, comparable to what is noted in the literature[38], was found to be essential for effective tumor eradication. This emphasizes that alongside the production of IFN-γ, the intrinsic killing efficiency of T-cells is a critical factor in the overall therapeutic outcome. Importantly, the model predicted that the therapeutic success is largely independent of the number of macrophages present at the time of treatment. This suggests that rSFV-IFN-γ may effectively overcome the influence of macrophages non-permissive for SFV, provided that the thresholds for infected cells and T-cell cytotoxicity are met.

These theoretical predictions were substantiated *in vivo* using the murine KPC3 pancreatic tumor model, which is an aggressive, immunologically "cold" tumor[56–58] known for its dense macrophage-rich barrier and poor T-cell infiltration. Intratumoral administration of rSFV-IFN-γ led to a marked delay in tumor growth compared to controls. Flow cytometry analysis of tumor-infiltrating and systemic immune cells confirmed an increased presence and activation of cytotoxic CD8+ T-cells, CD4+ T-helper cells, NK cells, and a shift in macrophage phenotype from immunosuppressive to pro-inflammatory populations. This immune reprogramming likely resulted from intra-tumoral IFN-γ expression, macrophage repolarization, or both. Notably, while rSFV-IFN-γ was most effective, even high-dose rSFV-GFP showed partial tumor growth delay, suggesting that viral infection and innate immune stimulation alone contribute to therapeutic benefit. These findings collectively underscore the potential of rSFV-IFN-γ virotherapy to enhance antitumor responses by leveraging the synergistic effects of targeted viral infection and potent immune activation, regardless of the presence of macrophages.

Based on our empirical findings, we confirm that rSFV while not infecting macrophages leads to their immunogenic activation. Importantly, as the frequency of macrophages in coculture with cancer cells increased, the infectivity of the cancer cells decreased in both monolayer and spheroid-based settings. We confirmed that macrophages themselves did not allow rSFV-encoded transgene expression, which can likely be attributed to innately active antiviral signaling pathways. This is in line with our previous work[32] and the observations made by Olupe Kurena and colleagues[37]. Multiple oncolytic viruses have been shown to activate macrophages during therapy, both through direct virus-uptake and signaling from infected cancer cells[39–43]. Specifically for replication-competent oncolytic viruses, macrophage activation has also been attributed to therapeutic resistance due to virus-uptake by macrophages, antiviral signaling (e.g. via TNF-α or IFN pathway) and killing of infected cancer cells to inhibit virus replication[42–44]. In case of replication-deficient virotherapy like rSFV, previous studies have demonstrated that sensing of viral RNA by various endosomal toll-like receptors (TLRs) and cytosolic retinoic acid-inducible gene I (RIG-I) receptor can result in degradation of viral RNA to downregulate viral-protein expression in macrophages[9]. However, TLR and RIG-I mediated sensing also may lead to the proinflammatory activation of macrophages[45,46]. In line with our previous study[32], we observed that macrophages upregulate CD80 and CD86 surface protein expression upon viral stimulation, corresponding to a stronger co-stimulatory signaling for antigen presentation. This is attributed partially via activation of JAK-STAT and NF-κB signaling in macrophages[32]. A similar macrophage activation profile was observed independent of their initial naïve (M_{naive}) or regulatory (M_{IL4}) macrophage phenotype. Various groups have demonstrated so far that although macrophages demonstrate functionally distinct phenotypes[47,48], they are capable of switching through these phenotypes in response to environmental stimuli within 48 hours[49,50]. Despite their adaptability, macrophages maintain their non-permissive nature towards rSFV-mediated transgene expression independent of their initial phenotype, their expression of rSFV-entry receptors, and their switch towards a pro-inflammatory phenotype upon rSFV-mediated stimulation.

Engineering rSFV to encode IFN-γ in infected cancer cells enhanced macrophage polarization to a
proinflammatory phenotype, is consistent with the increased frequency of anti-tumoral macrophages
observed *in vivo* following rSFV-IFN-γ treatment. This aligns with previous findings where rSFV driven
IFN-γ expression induced a strong antitumor immune response and inhibited tumor growth in a
murine orthotopic breast cancer model, by promoting macrophage activation and remodeling the
tumor microenvironment[51]. In our study, macrophages upregulated pro-inflammatory markers
(CD80, CD86) and downregulated pro-tumoral markers (CD206) to a significantly higher magnitude
when in the presence of infected cancer cells expressing rSFV-encoded IFN-γ, as compared to being
stimulated by virus alone or rSFV-GFP and rSFV-Flt3L infected cancer cells, where Flt3L served as an
myeloid-stimulating control. We assume this to be a result of the bimodal stimulation by virus-
particles and extracellular sensing of IFN-γ. Previous studies have demonstrated that either type-I or
type-II IFNs are required for macrophages to polarize towards a strongly pro-inflammatory
phenotype when stimulated by stress signals like extracellular release of heat-shock proteins or
danger signal like bacterial lipopolysaccharides and viral RNA[48,49,52]. Therefore, it is likely that the
intracellular release of rSFV-RNA upon virus uptake in the macrophages and extracellular IFN-γ
produced by neighboring infected cancer cells results in their proinflammatory activation. We indeed
confirmed that macrophage activation by rSFV-encoded IFN-γ can be reversed upon blocking JAK-
STAT signaling. Furthermore, as we did not observe rSFV-mediated transgene expression by
macrophages, we consider that rSFV-encoded IFN-γ expression is limited to infected cancer cells and
that IFN-γ is not sensed in an autocrine manner by macrophages.

Our study demonstrates that rSFV-encoding IFN-γ significantly stimulates T-cell responses,
independent of the influence of macrophages. This was observed across both monolayer (2D) and
spheroid (3D) cocultures, as well as *in vivo* highlighting the robustness of this approach. In monolayer
cocultures, the presence of rSFV-IFN-γ led to pronounced activation of both CD4 and CD8 T-cells,
even in the presence of high macrophage frequencies. Here, T-cell activation relied on IFN-γ or
macrophage polarization and was diminished upon inhibition of JAK-STAT signaling. Notably,
macrophages, despite their non-permissiveness to SFV infection and potential for antiviral signaling,
did not impede T-cell activation induced by rSFV-IFN-γ. This finding underscores the potential of IFN-
γ as a powerful immunostimulatory cytokine that can override the suppressive effects of
macrophages in the tumor microenvironment. Furthermore, the impact of macrophage phenotype
and frequency on T-cell responses varied between monolayer and spheroid cocultures. In monolayer
systems, macrophages reduced CD4 but not CD8 T-cell activation, suggesting a differential regulatory
effect on helper versus cytotoxic T-cells. However, in spheroid cocultures, an increase in macrophage
frequency led to a reduction in activation of both CD4 and CD8 T-cells. Despite these variations, rSFV-
IFN-γ consistently promoted strong T-cell responses regardless of either a naïve (M_{naive}) or regulatory
(M_{IL4}) macrophage phenotype. Finally, in both monolayer and spheroid cocultures, a high frequency
of macrophages correlated to a stronger CD4 T-cell activation by rSFV-IFN-γ, which may be attributed
to rSFV-mediated proinflammatory activation of the macrophages. In addition to transgenic IFN-γ
expression, differences in innate cancer cell-mediated immune signaling were also found to influence
both macrophage and T-cell activation, as noted in a previous study[23]. Nevertheless, IFN-γ expression
in infected PANC-1 and Ca-Ski cells proved effective in inducing robust immune responses.
Observations of Meissner et al. corroborate our results, that encoding IFN-γ by a non-replicating
virotherapy, influenza A virus in their case, can promote tumor eradication through enhanced
immune responses and not primarily through virus-mediated oncolysis[53]. Indeed, a replication-
competent virotherapy causing extensive virus-mediated oncolysis in combination with IFN-γ
mediated immune stimulation can also lead to tumor eradication albeit with a risk of virus
persistence[54].

In conclusion, our study presents a compelling case for the use of rSFV-encoding IFN-γ as a means to
enhance antitumor immune responses, effectively overcoming macrophage-mediated regulation of
virus-infection and T-cell activation. The combination of computational modeling with experimental
validation in diverse human tumor models, provides evidence supporting this therapeutic strategy.
Future studies should explore the clinical translation of these findings, assessing the efficacy of rSFV-
IFN-γ in preclinical models like patient-derived organoids to test its therapeutic potential in a patient-
specific manner. The significant implications of our findings lie in their potential to inform the design
of anticancer virotherapies, especially for tumors characterized by high macrophage infiltration and
immune suppression.

Limitations of the study

The primary aim of our study was to assess and improve the efficacy of rSFV replicons in the context
of tumor-associated macrophages in a human model. While our study demonstrates the potential of
rSFV-encoding IFN-γ in enhancing T-cell immune responses, several limitations must be
acknowledged. One significant limitation of our study is the lack of direct measurement of T-cell-
mediated tumor cell killing *in vitro*, due to the unavailability of a tumor-specific T-cell model.
However, the observed tumor growth inhibition and increased immune cell infiltration *in vivo*
following rSFV-IFN-γ treatment provide indirect evidence supporting the functional relevance of the
T-cell activation seen in our coculture systems. Furthermore, our study does not fully account for the
complex interactions within the tumor immune microenvironment, focusing primarily on T-cell
responses. This narrow focus may overlook the broader spectrum of immune cell dynamics and the
potential contributions of other immune cells, such as dendritic cells. Additionally, the *in vitro*
coculture models, while informative, do not fully replicate the *in vivo* human tumor
microenvironment. Expanding the study to test different virotherapies in the context of *in vivo*
humanized murine models with xenograft tumors or patient-derived organoids representing diverse
tumor and HLA-types will be essential to validate these findings and to fully understand the
therapeutic potential of immunogenic virotherapy in a clinical setting.

Acknowledgements

This study was partially funded by Stichting De Cock-Hadders (grant 2020-35) awarded to DB and by
Stichting Overleven met alvleesklierkanker (grant SOAK 22.02) awarded to NvM. Part of the work has
been performed at the UMCG Imaging and Microscopy Center (UMIC) and the Flow Cytometry
Research Unit (FCU) which is sponsored by UMCG. We thank the Center for Information Technology
of the University of Groningen for their support and for providing access to the Hábrók high-
performance computing cluster. We acknowledge the Flow Cytometry Facility and animal facility at
LUMC for their support with the *in vivo* study and immunological analyses.

Author contributions

All authors made substantial contributions to the manuscript.

LH: Writing and editing manuscript, Data collection and analysis, Validation, Visualization

PK: Writing and editing manuscript, Data collection and analysis, Validation, Visualization

TJ: Conceptualization, Writing and editing manuscript, Software, Data analysis, Visualization,
Supervision

BNH: Writing and editing manuscript, Data collection and analysis, Validation, Visualization

FJW: Conceptualization, Writing and editing manuscript, Supervision

NM: Conceptualization, Writing and editing manuscript, Supervision, Funding acquisition

TD: Conceptualization, Writing and editing manuscript, Supervision, Funding acquisition

DB: Conceptualization, Writing and editing manuscript, Data collection and analysis, Software,
Visualization, Supervision, Funding acquisition

Declaration of interests

Toos Daemen is co-founder of ViciniVax, a spin-off from the University of Groningen, developing
therapeutic cancer vaccines.

Figure titles and legends

Figure 1: Effect of macrophage frequency on rSFV-mediated tumor infection. (A) Illustration
explaining the mode of action of rSFV encoding either GFP or an immunogenic signal used as a non-
replicating "suicidal" virotherapy. **(B)** Schematic of the experimental setup for studying rSFV infection
(MOI-10) in a pancreatic cancer cell line (PANC-1) cocultured with macrophages in both monolayer
(2D) and spheroid (3D) spatial organizations. The frequency of macrophages and cancer cells were
controlled at the time of initiating the coculture. The coculture setup showing representative images
of **(C)** spheroids or **(D)** monolayers composed of cancer cells, infected cells (green, GFP+ cells), and
varying frequencies of macrophages (red, stained with FarRed dye) after 24 hours post infection with
rSFV encoding GFP (rSFV-GFP). **(E)** Diagram illustrating experimental setups with varying frequencies
of macrophages and cancer cells. Top row (variable setup): Frequency of macrophages and cancer
cells are varied. Bottom row (constant setup): Frequency of macrophages is varied while cancer cells
are kept constant. The grey cells indicate empty space that can be occupied by either cancer cells or
macrophages. Quantitative analysis of GFP expression indicating virus infection in the **(F)** spheroid
and **(G)** monolayer coculture setups. The plots represent data from 4 replicates of respective
coculture methods. Data are presented as mean values $\pm$ SD.

Figure 2: Effect of macrophage phenotype on rSFV-mediated tumor infection. (A) Schematic of the
experimental setup for polarizing peripheral blood derived macrophages to either a naïve (M_{naive}) or
or an anti-tumoral ($M_{LPS+IFN\gamma}$ or M1-like), or a pro-tumoral (M_{IL4} or M2-like) phenotype using
cytokines. **(B)** The surface expression of immunomodulatory proteins involved in anti-tumoral (CD80,
CD86) and pro-tumoral (CD206, CD163) macrophage activity. **(C)** Principle component analysis of the
differences in the phenotypes of macrophages from various healthy blood donors (n=5).
Representative images of rSFV infection (green, GFP+ cells) in a pancreatic cancer cell line (PANC-1)
cocultured with M_{naive} and M_{IL4} macrophages in both spheroid **(D)** and monolayer **(E)** spatial
organizations. Quantitative analysis of GFP expression indicating virus infection (MOI-10) in the **(F)**
spheroid and **(G)** monolayer coculture of PANC-1 cells with M_{naive} and M_{IL4} macrophages. **(H)**
Quantification of GFP expression in monolayer coculture of Ca-Ski cancer cells with M_{naive} and M_{IL4}
macrophages. The plots represent data from 4 replicates of respective coculture methods. Data are
presented as mean values $\pm$ SD.

Figure 3: Predicting the outcomes of immunogenic-rSFV therapy using a spatiotemporal model of tumor development. (A) Snapshot of a simulation. Left: Spatial structure of a tumor containing uninfected cancer cells (red), virotherapy-infected cancer cells (green), and macrophages (blue). Right: Distribution of infection-induced immunogenic signals (magenta) released from dying infected cancer cells, which stimulate a cytotoxic anti-cancer T-cell response. **(B)** Time trajectory illustrating the dynamics of a suicidal rSFV-therapy. Initially, uninfected cancer cells are targeted by the virus and weakly attacked by T-cells. Over time, virus-killed cancer cells release immunogenic signals, boosting T-cell cytotoxicity. Finally, the boosting effect wanes as immunogenic signal-levels diminish. **(C)** Effect of the amount of immunogenic signal produced per infected cells on the probability of tumor eradication. The colors indicate five simulation scenarios differing in the percentage of infected cells at the time of infection. **(D)** The effect of anticancer T-cell cytotoxicity rates on the probability of tumor eradication. **(E)** The effect of macrophage frequency at the time of rSFV therapy on the probability of tumor eradication. When not specifically under consideration, the parameter values were kept at their default values: the amount of immunogenic signal released was set to 0.25, the T-cell cytotoxicity rate was set to 5, and the number of macrophages was set to 2500. Colored dotted lines indicate the mean values, colored envelopes indicate the 95% confidence interval obtained via bootstrapping. Each panel represents 10000 simulations for respective parameter combinations.

Figure 4: Impact of IFN-γ on tumor associated macrophage and T-cells in the tumor microenvironment. (A) Schematic of cytokine production and reception by T-cells and macrophages (MØ), highlighting IFN-γ, IL-10, and IL-4 as key cytokines involved in influencing both cell types. Cytokine interactions are color-coded based on their impact on tumor progression: pro-tumoral (purple) and antitumoral (orange). **(B)** The effective concentration range of various cytokines (EC$_{50}$) with the points indicating the maximum and minimum experimentally reported limits (as per [35]). **(C)** Analysis process of cervical (CESC, 300 samples) and pancreatic (PAAD, 151 samples) cancer patient datasets stored at The Cancer Genome Atlas (TCGA) by using the CRI iAtlas online platform to correlate the presence of macrophage types with T-cells and IFN-γ signature. **(D)** Correlation heatmap of intra-tumoral IFN-γ signature with presence of various macrophage phenotypes. **(E-H)** Scatter plots showing the relationship between the intra-tumoral IFN-γ signature with the presence of classical (M$_{class}$ or M1-like) and alternatively activated (M$_{alter}$ or M2-like) macrophages in cervical or pancreatic cancer patients. **(I)** Correlation heatmap of intra-tumoral fraction of cytotoxic (CD8) T-cells with presence of various macrophage phenotypes. **(J-M)** Scatter plots showing the relationship between the intra-tumoral fraction of CD8 T-cells with the presence of M$_{class}$ or M$_{alter}$ macrophages in cervical or pancreatic cancer patients. In E-F and J-M, bivariate linear regression was performed to assess correlation, with r (correlation coefficient) and p-values indicated per plot.

Figure 5: Evaluation of T-cell activation in tumor-immune monolayer cocultures by rSFV. (A) Schematic representation of the experimental setup consisting of tumor-immune monolayer cocultures of cancer cells and macrophages treated with rSFV (MOI-10), followed by the addition of PBMCs to introduce T-cells and evaluate their activation. Flow cytometry analysis to measure activation of **(B)** CD4 helper T-cells and **(C)** CD8 cytotoxic T-cells based on the expression of CD69 and CD107a proteins, marked in the upper right quadrant in each panel. rSFV-GFP treated cocultures are indicated in green, rSFV-IFN-γ treated cocultures in red and non-infected (NT) in black. The representative plots in (B-C) illustrate T-cell activation in the context of PANC-1 cancer cells in coculture with either M$_{naive}$ or M$_{IL4}$ macrophages in different frequencies. Quantification of **(D)** CD4 helper T-cell or **(E)** CD8 cytotoxic T-cell activation in the tumor-immune cocultures by rSFV. The plots represent data from 4 replicates. Data are presented as mean values ± SD.

Figure 6: Evaluation of T-cell activation in tumor-immune spheroid co-cultures upon rSFV infection.
(A) Schematic representation of the experimental setup consisting of tumor-immune spheroid
cocultures of cancer cells and macrophages treated with rSFV, followed by the addition of PBMCs to
introduce T-cells and evaluate their activation. Flow cytometry analysis to measure activation of T-
cells as explained in Figure 6. Quantification of **(B)** CD4 helper T-cell or **(C)** CD8 cytotoxic T-cell
activation in the tumor-immune cocultures by rSFV. rSFV-GFP treated cocultures are indicated in
green and rSFV-IFN-γ treated cocultures in red. The plots represent data from 4 replicates. Data are
presented as mean values ± SD.

Figure 7: Influence of rSFV infection on macrophage phenotype. (A) Experimental setup to study
changes in macrophage phenotype upon stimulation by rSFV (MOI-10). This includes studying the
effect of **(B)** either rSFV-particles alone (rSFV row), **(C)** or by infected cells alone, **(D)** or by both. **(E)**
Heatmap of change in cell-surface expression of proteins involved in either pro-tumoral (CD206) or
anti-tumoral (CD80, CD86) function of M_{IL4} macrophages cocultured or not with PANC-1 cancer cells
upon stimulation resulting from rSFV particles and/or infected cells. Heatmap of M_{naive} and M_{IL4}
macrophage cell-surface protein expression upon stimulation with **(F)** rSFV particles alone (bottom
two rows), or both virus particles and infected **(G-H)** PANC-1 or **(I-J)** Ca-Ski cancer cells. The plots
represent mean value data from 4 replicates.

Figure 8: Intratumoral virotherapy with rSFV-IFN-γ induces anti-tumoral macrophage polarization
and immune cell infiltration in KPC3 tumor-bearing mice. (A) C57BL/6 mice were subcutaneously
injected with KPC3 tumor cells (2×10^5) on day 0, and received intra-tumoral virotherapy with PBS
(untreated control) or rSFV-replicons encoding either GFP or IFN-γ at doses of 10^7 or 10^8 particles per
injection on days 12, 13 and 14. (B) Individual tumor growth across treatment groups, with untreated
(black filled circles), SFV-GFP 10^7 (green empty circles), SFV-GFP 10^8 (green filled circles), SFV-IFN-γ
10^7 (red empty circles), and SFV-IFN-γ 10^8 (red filled circles). (C–N) Flow cytometric analysis of
percentage tumor-infiltrating immune cells 7 days post-treatment: (C) total F4/80+ macrophages, (D)
pro-tumoral CD206+ macrophages, (E) anti-tumoral MHCII+ macrophages, (F) NK cells, (G) CD69+ NK
cells, (H) KLRG1+ NK cells, (I) CD8+ T cells, (J) CD44+ CD8+ T cells, (K) PD-1+ CD8+ T cells, (L) CD4+ T
cells, (M) CD44+ CD4+ T cells, (N) PD-1+ CD4+ T cells. (O–R) Flow cytometric analysis of splenic
immune cells: (O) CD3+ T cells, (P) CD69+ NK cells, (Q) CD44+ CD8+ T cells, (R) CD44+ CD4+ T cells. Each
symbol represents an individual mouse (n=6–7 per group), with bars indicating mean ± SEM. To
evaluate differences in tumor growth and immune cell infiltration following virotherapy across
treatment groups, data were analyzed using one-way ANOVA to assess statistical significance, with p-
value indicated per comparison. The individual tumor growth curves for respective groups are shown
in **supplementary figure 9** and the flow cytometry gating strategy is illustrated in **supplementary**
figure 10.

STAR Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
anti-human CD4 (OKT4) FITC	Biolegend	317408
anti-human CD8 (SK11) APC-Cy7	Biolegend	344714
anti-human CD38 (HIT2) BV421	Biolegend	303526
anti-human CD69 (FN50) APC	Biolegend	310910

anti-human CD107a (H4A3) PE	Biolegend	328608
anti-human CD223 (LAG-3) (11C3C65) PE	Biolegend	369306
anti-human CD152 (CTLA-4) (BNI3) BV421	Biolegend	369606
anti-human CD279 (PD-1) (A17188B) APC	Biolegend	621610
anti-human CD11b (ICRF44) PE	Biolegend	301306
anti-human CD206 (15-2) APC-Fire	Biolegend	321133
anti-human CD80 (2D10) APC	Biolegend	374204
anti-human CD86 (BU63) FITC	Biolegend	328608
anti-human CD163 (GHI/61) PE	Biolegend	333605
anti-human CD68 (Y1/82A) PE-Cy7	Biolegend	333815
anti-human CD206 (MMR) APC	Biolegend	321109
anti-human CD80 (2D10) BV421	Biolegend	305221
anti-murine CD45.2 (104) FITC	Biolegend	109806
anti murine CD45.2 (104) AF700	Biolegend	109822
anti-murine CD3 (145-2C11) PE-CF594	BD Biosciences	562286
anti-murine CD8α (53-6.7) AF700	eBioscience	56-0081-82
anti-murine NK1.1 (PK136) BV650	BD Biosciences	564143
anti-murine CD4 (RM4-5) APC	BioLegend	100547
anti-murine F4/80 (BM8) PE-Cy7	Biolegend	123114
anti-murine CD206 (C068C2) BV711	Biolegend	141727
anti-murine Ly6G (1A8) BV786	Biolegend	127645
anti murine Ly6C (HK1.4) BV605	Biolegend	128036
Anti mouse/human CD44 (IM7) BV785	Biolegend	103059
Anti mouse CD69 (H1.2F4) FITC	eBioscience	11-0691-82
Anti mouse KLRG1 PE-Cy7	eBioscience	25-5893-82
Anti mouse CD279 (PD-1) (28F.1A12) PE	Biolegend	135205
anti-murine MHC-II (M5/114) BV421	BD Biosciences	562564
Zombie aqua viability dye	Biolegend	423102
Bacterial and virus strains		
Plasmid (Helper-2 and SFV-transgene) constructs to produce recombinant SFV replicon particles	Liljestrom lab, Karolinska Institutet Sweden	NA
E coli K12 JM110	Agilent	200239
rSFV-GFP	This study (in-house)	NA
rSFV-IFN-γ	This study (in-house)	NA
Biological samples		
Human buffy coats for PBMC isolation from HLA-A*02:01-typed healthy donors	Sanquin, Nijmegen, The Netherlands	NA
Chemicals, peptides, and recombinant proteins		
Restriction enzyme PspOMI	ThermoFischer Scientific	ER0131
Restriction enzyme XmaI	ThermoFischer Scientific	ER0171
Restriction enzyme SpeI	ThermoFischer Scientific	ER1251
SP6 polymerase	Amersham Pharmacia Biotech, Piscataway, US	NA

Celltrace FarRed dye	Invitrogen	C34572
Ruxolitinib	Stem Cell Technologies	73402
α-chymotrypsin	Sigma Aldrich	C4129
Aprotinin	Sigma Aldrich	A1153
Macrophage colony stimulating factor (M-CSF)	Biolegend	572903
Interleukin-4 (IL-4)	Biolegend	574004
Interferon-gamma (IFN-γ)	Biolegend	570204
Lipopolysaccharides (LPS) from *E. coli* O128:B12	Sigma Aldrich	L2755
LIberase TL	Sigma Aldrich	05401020001
Mouse BD Fc block (anti-mouse CD16 and CD32)	BD Biosciences	553142
Hydrogen peroxidase	VWR	7422478
ELK powder	Campina	NA
Rat, anti-mouse F4/80 antibody	Sanbio	CI:A3-1
Biotinylated rabbit anti-rat antibody	Agilent	NA
VECTASTAIN Elite ABC HRP Kit	Vector laboratories	PK-6100
Entellan	Sigma Aldrich	107960
Critical commercial assays		
NA		
Deposited data		
NA		
Experimental models: Cell lines		
Hamster, BHK-21 cells	ATCC	ccl-10
Human, PANC-1 cells	ATCC	crl-1469
Human, Ca-Ski	ATCC	crl-1550
Murine, KPC3 cells	Nadine van Montfoort lab, Leiden University Medical Center, Leiden	RRID:CVCL_A9ZK
Experimental models: Organisms/strains		
C57BL/6J mice	Charles River Laboratories	C57BL/6NCrl
Oligonucleotides		
NA		
Recombinant DNA		
Human IFN-gamma gene	This paper	NA
Murine IFN-gamma gene	This paper	NA
Software and algorithms		
Graphpad Prism	Graphpad	https://www.graphpad.com/
Incucyte® Software Modules	Sartorious	https://www.sartorius.com/
Qpath	Qpath	https://qupath.github.io/
FlowJo	FlowJo	https://www.flowjo.com/
Rstudio	Posit (Rstudio)	http://www.posit.co/

RawGraphs	RawGraphs	https://www.rawgraphs.io/
Other		
Nunclon Sphera 96 well plates	ThermoFischer Scientific	174925
Oncolytic Virus Immune simulator (OVI) model	This paper	https://github.com/rugtres/OVI/tree/macrophage
Superfrost Plus slides	VWR	48311-703

Resource availability

Lead contact: Further information and requests for resources and reagents should be directed to and
will be fulfilled by the lead contact Darshak K. Bhatt (d.bhatt@umcg.nl)

Material availability: All unique/stable reagents generated in the study are available upon request,
except where restricted by institutional agreements or MTAs.

Data and code availability:

• All data reported in this paper will be shared by the lead contact upon request.
• Any additional information required to reanalyze the data reported in this paper is available
from the lead contact upon request.
• The code used for this work and an executable version of the Oncolytic Virus Immune
simulator (OVI) can be found at https://github.com/rugtres/OVI/tree/macrophage.

Method details

TCGA analysis: We used the online CRI iAtlas portal to analyze TCGA data of cervical and pancreatic
cancer patients[1,36]. We selected the TCGA cohort of cervical cancer (CESC) patients (n=300) and
pancreatic cancer (PAAD) patients (n=151). Performed Pearson correlation to analyze the
relationship between IFN-γ signature, cytotoxic CD8 T-cells, and pro-tumoral (M2-like or M_{alter}) or
anti-tumoral (M1-like or M_{class}) macrophages. IFN-γ signature and immune cell subset quantifications
of the tumor samples were performed previously as a part of the immune landscape analysis of
cancer[1,36].

Computational model: In our previous work[33,34], we developed a spatiotemporal model for the effect
of anticancer virotherapy on tumor development to evaluate how anticancer T-cell responses
triggered by immunogenic signals affect the therapeutic outcome. In the current study, we employed
the model to simulate a tumor environment with three cell types: stromal macrophages, uninfected
infection-sensitive cancer cells, and infected cancer cells. We assumed that infection-sensitive cancer
cells and stromal macrophages are initially distributed randomly. Cells can divide, change status, or
die, with varying rates across cell types. Cancer cells can be infected by a therapeutic virus, which
targets and kills them, while stromal macrophages are non-permissive to infection. Since rSFV is a
non-replicating virotherapy, no further viral spread occurs in the model. We incorporated a cell-
specific immune response to assess how virus-induced immunogenic signals influence therapeutic
outcomes. The model assumes that infection-induced cell death induces the release of immunogenic
signals, which diffuses and subsequently activates T-cells to kill cancer cells, with cytotoxicity of the
activated T-cells depending on the local concentration of immunogenic signals. Key variables
assessed in this study include the percentage of infected cancer cells, amount of immunogenic signal
released per infected cell, T-cell cytotoxicity rate, and macrophage frequency. Immunogenic
molecules are initially absent and increase in concentration (λ) in the grid cell upon infection-induced
cell death. Immunogenic signals disperse to neighboring cells via diffusion and diminishes over time
due to evaporation. In our previous studies, the model revealed that therapeutic outcomes are
probabilistic. To account for this, we ran 10,000 simulations per parameter combination to capture
stochastic variations in the results. Since rSFV is a suicidal virotherapy and does not replicate, the
model predicted two possible outcomes at 1,000 days: (i) total tumor eradication, where all cancer
cells are eliminated; or (ii) tumor persistence, where cancer cells survive due to insufficient virus-
and/or immune-mediated killing. For this study, we focused on the probability of total tumor

eradication as the primary outcome. A detailed description of the model can be found in references
33 and 34. The code used for this work and an executable version of the Oncolytic Virus Immune
simulator (OVI) can be found at https://github.com/rugtres/OVI/tree/macrophage.

Cell culture: BHK-21, Ca-Ski, and Panc-1 cells were cultured in RPMI 1640 medium supplemented
with 10% fetal bovine serum (FBS), 100 U/ml penicillin, and 100 µg/ml streptomycin (P/S, Thermo
Scientific). Since, PANC-1 and Ca-Ski cells are HLA-A*02:01-restricted, we performed the cancer-
immune coculture assays with peripheral blood mononuclear cells (PBMCs) derived from HLA-
A*02:01-matched healthy donors (Sanquin, Netherlands). Cancer cell-macrophage cocultures were
cultured in RPMI 1640 medium supplemented with 10% FBS and 100 U/ml penicillin, 100 µg/ml
streptomycin. Cocultures with PBMCs were similarly cultured in RPMI 1640 medium supplemented
with 10% FBS and 100 U/ml penicillin, 100 µg/ml streptomycin. For spheroid cocultures, cells were
seeded in NuncSphera round-bottom 96-well plates (Thermo Fischer) and centrifuged for 10 minutes
at 1500 RPM. All cells were maintained in an incubator at 37°C with 5% CO_2.

Monocyte derived macrophage differentiation and polarization: Peripheral blood derived monocytes
were differentiated to macrophages and polarized in RPMI 1640 medium supplemented with 10%
FBS, 100 U/ml penicillin, 100 µg/ml streptomycin, 1mM sodium pyruvate (Thermo Scientific), non-
essential amino-acids (NEAA, Thermo Fisher), and 100 ng/mL macrophage colony-stimulating factor
(M-CSF, BioLegend San Diego, CA, USA). Briefly, PBMCs from healthy donors were cultured for 2
hours to allow monocyte adherence. Afterward, non-adherent cells were gently removed. The
adhered cells were stimulated with medium containing 100 ng/ml M-CSF. The medium containing M-
CSF was refreshed on 3 days post culture. Macrophages were replated on 7 days of culture. 24 hours
later, macrophages were polarized to either a M_{IL4} state with medium containing 20 ng/mL IL-4
(BioLegend, San Diego, CA, USA), or a M1-like state with medium containing 20 ng/ml IFN-γ and 100
ng/ml lipopolysaccharides (LPS), or were kept unstimulated (M_{naive}).

Design, production and titer determination of rSFV: We have previously designed rSFV-replicon
particles expressing immunogenic transgenes and capable of a single round of infection[23,55]. Briefly,
enhanced green fluorescent protein (GFP) and IFN-γ (human and murine versions) as transgenes
were ordered as a DNA construct (Eurofins Genomics, Ebensburg, Germany) and were cloned in the
SFV-replicon backbone plasmid (pSFV) by using PspOMI and XmaI as restriction sites and *E. coli*
JM110 as the competent cell chassis. Sanger sequencing was performed on isolated clones to
validate insertion (Eurofins Genomics). pSFV containing either GFP or IFN-γ and a SFV-Helper-2
plasmid (hSFV) were linearized with SpeI digestion (Life Technologies) for *in vitro* RNA synthesis by
SP6 polymerase reaction (Amersham Pharmacia Biotech, Piscataway, US). Next, pSFV-transgene RNA
and hSFV RNA were mixed in a 2:1 ratio and co-transfected in BHK21 cells in the presence of
electroporation buffer using the BioRad Gene Pulser II system (2 pulses, 850 V/25 µF; Biorad,
Hercules, U.S.A.). After electroporation, the cells were cultured in RPMI 1640 media supplemented
with 5% FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin for 48 hours at 30°C with 5% CO_2. The
rSFV-transgene particles were purified by discontinuous sucrose density gradient ultracentrifugation
and stored in TNE buffer as aliquots at -80°C. Before use, all rSFV particles were activated by the
addition of 1:20 volume 10 mg/ml α-chymotrypsin (Sigma Chemical, St. Louis, US) and 2 mM $CaCl_2$ for
30 minutes to cleave the mutated spike proteins. After which, the α-chymotrypsin was inactivated by
the addition of 1:2 volume 2 mg/ml aprotinin (Sigma Chemical). Finally, the titer determination of
rSFV-particles was performed as described previously. Briefly, rSFV-particles were titrated by serial
dilution on monolayers of BHK-21 cells cultured in LabTek slides. After infection and incubation for 24
hours, the cells were fixed in 10% (w/v) acetone and further stained for nsP3 using a primary
polyclonal rabbit-anti-nsP3 antibody (1:2000 dilution), whilst a secondary Cy3-labeled animal-anti-

rabbit antibody (1:200 dilution) was used to amplify the signal. Positive cells were counted using
fluorescence microscopy, and the titers were determined.

Evaluating virus-infection in macrophage-cancer cell cocultures: M_{naive} or M_{IL4} macrophages were
differentiated from human PBMCs as described above. Different frequencies of M0 or M_{IL4}
macrophages were cocultured with either a constant or variable frequency of Panc-1 or Ca-Ski cancer
cells. When specified, macrophages were stained with CellTrace™ Far Red dye (ThermoFischer
Scientific, C34564) following the recommended protocol from the manufacturer. After overnight
incubation, cocultures were infected with an MOI of 10 (multiplicity-of-infection) of rSFV-GFP. The
ability of rSFV-particles to infect cells and express GFP was monitored over 24 hours by Incucyte-
based brightfield and fluorescence microscopy. The number of GFP-positive cells served as a measure
of the rSFV-GFP particle's infectivity.

Evaluating macrophage activation by virus particles and infected cancer cells: M_{naive} or M_{IL4}
macrophages were differentiated from human PBMCs as described above. M_{naive} or M_{IL4}
macrophages were cocultured in a 1:2 ratio with Panc-1 or Ca-Ski cells in a 24-well plate. After
overnight incubation, cocultures were infected with MOI-10 of rSFV-GFP or rSFV-IFN-γ. 24 hours
post-infection (HPI), all cells were collected and processed for Flow cytometry-based analysis of
macrophage activation markers. The gating strategy is illustrated in **Supplementary Figure 7**.

Evaluating T-cell activation in tumor-immune cocultures: M_{naive} or M_{IL4} macrophages were
differentiated as described above, from HLA-A*02:01 typed healthy donors. They were cocultured
with Panc-1 cells or Ca-Ski cells for 24 hours in 2 tumor cells to 1 macrophage cell ratio (15 000 cells
per well) in a treated 96-well plate. Cocultures were then infected with MOI-10 of rSFV-particles
encoding GFP or IFN-γ. 8 hours post-infection, freshly thawed PBMCs (from the same donor as used
for generating macrophages) were added to the cocultures (75 000 cells per well). 18 hours post
PBMCs addition, all cells were collected and stained for CD4 and CD8 T-cell population, as well as
immune activation cell surface markers, and analyzed *via* Flow cytometry. The gating strategy is
illustrated in **Supplementary Figure 4**.

Animal experiments: *In vivo* experiments were conducted as described previously[56–58]. In brief, male
C57BL/6J mice (6–8 weeks old; were obtained from Charles River Laboratories and housed in
individually ventilated cages (3-4 mice/cage) at the LUMC animal facility. The experiment was
reviewed, ethically approved, and registered by the institutional Animal Welfare Body of Leiden
University Medical Center. The experiment was carried out in compliance with Dutch and EU
regulations on animal experimentation under project license AVD1160020187004, issued by the
competent authority on animal experiments in the Netherlands. After one week of acclimatization,
mice were subcutaneously injected in the right flank with 2×10^5 KPC3 cells in 100 μL PBS/0.1% BSA.
Once tumors became palpable, mice were stratified into groups with similar average tumor volumes
and treated intratumorally under isoflurane anesthesia with 10^7 or 10^8 infectious particles of rSFV-
GFP, rSFV-IFN-γ, or PBS (30 μL) on three consecutive days. In contrast to the previous experiments,
rSFV-encoding murine-IFN-γ was used. Tumor volume, welfare and body weight were monitored
thrice weekly using calipers (tumor volume = length × width × height), blinded when possible. Mice
were sacrificed for immune profiling 7 days post-treatment or earlier if they reached human
endpoint, as described to be tumors exceeding 1000mm^3, signs of tumor ulceration or non-
experimental related complications (n=1 mouse, that was excluded due to fighting wounds). Tumors
were digested with Liberase TL for 15 min at 37°C, and processed into single-cell suspensions. Cells
were stained with Zombie Aqua viability dye, incubated with Mouse BD Fc Block™ (anti-Mouse
CD16/CD32), stained for cell surface markers and fixed in 1% paraformaldehyde. Flow cytometry was

performed on a BD LSRFortessa X20 at the LUMC Flow Cytometry Facility and analyzed using FlowJo
v10. The gating strategy is illustrated in **Supplementary Figure 10**.

Immunohistochemistry staining: Formalin fixed tumors were embedded in paraffin and sectioned at
4μm before placing on Superfrost® Plus slides (VWR). Sections were dried overnight at 37°C and
stored at 4°C until staining. Slides were deparaffinized and endogenous peroxidase was blocked with
0,3% hydrogen peroxidase (VWR) in methanol for 20 minutes. Rehydration was followed by 16
minute boiling in Trypsin/Calcium chloride solution (pH 7,4) followed by blocking with 0,2% ELK
(Campina) in PBS/0,1% BSA. Overnight incubation with primary rat anti-mouse F4/80 (Cl:A3-1,
Sanbio) was performed at RT in humidified box. The next morning, samples were incubated for 30
minutes at RT with biotinylated rabbit anti-rat (Agilent), followed by 30 minutes incubation with
avidin-biotin complex (VECTASTAIN® Elite® ABC HRP Kit; Vector Laboratories). Peroxidase activity was
detected using the 2-component liquid DAB+ system (Agilent) according to the manufacturer's
instructions for 5 min. Counterstaining was performed using hematoxylin (Sigma Aldrich) for 15
seconds, followed by dehydration and mounting using Entellan (Sigma Aldrich). Slides were scanned
using Zeiss Axio Scan Z.1 and processed with QuPath.

Quantification and statistical analysis: Experimental data represents the mean ± SEM of the number
of replicates. Statistical analysis of the data is specified in the respective figure legends when
applicable. For linear correlation, bivariate linear regression was performed, with r (correlation
coefficient) and p-values indicated per plot. one-way ANOVA was also used to compare groups and
assess statistical significance, with p-value indicated in the plots per comparison. Graphs were made
using Rstudio, RawGraphs, and Graphpad Prism 10.

Supplementary file

All the supplementary figures can be found in the supplementary document.

References

1. Thorsson, V., Gibbs, D.L., Brown, S.D., Wolf, D., Bortone, D.S., Ou Yang, T.-H., Porta-Pardo, E.,
Gao, G.F., Plaisier, C.L., Eddy, J.A., et al. (2018). The Immune Landscape of Cancer. Immunity *48*,
812-830.e14. https://doi.org/10.1016/j.immuni.2018.03.023.

2. Park, M.D., Silvin, A., Ginhoux, F., and Merad, M. (2022). Macrophages in health and disease. Cell
185, 4259–4279. https://doi.org/10.1016/j.cell.2022.10.007.

3. Biswas, S.K., and Mantovani, A. (2010). Macrophage plasticity and interaction with lymphocyte
subsets: cancer as a paradigm. Nat Immunol *11*, 889–896. https://doi.org/10.1038/ni.1937.

4. Fridman, W.H., Zitvogel, L., Sautès–Fridman, C., and Kroemer, G. (2017). The immune contexture
in cancer prognosis and treatment. Nat Rev Clin Oncol *14*, 717–734.
https://doi.org/10.1038/nrclinonc.2017.101.

5. Cassetta, L., and Pollard, J.W. (2023). A timeline of tumour-associated macrophage biology. Nat
Rev Cancer *23*, 238–257. https://doi.org/10.1038/s41568-022-00547-1.

6. Gooden, M.J.M., de Bock, G.H., Leffers, N., Daemen, T., and Nijman, H.W. (2011). The prognostic
influence of tumour-infiltrating lymphocytes in cancer: a systematic review with meta-analysis.
Br J Cancer *105*, 93–103. https://doi.org/10.1038/bjc.2011.189.

7. Brummel, K., Eerkens, A.L., De Bruyn, M., and Nijman, H.W. (2023). Tumour-infiltrating
lymphocytes: from prognosis to treatment selection. Br J Cancer *128*, 451–458.
https://doi.org/10.1038/s41416-022-02119-4.

8. Russell, S.J., Peng, K.-W., and Bell, J.C. (2012). Oncolytic virotherapy. Nat Biotechnol *30*, 658–670.
https://doi.org/10.1038/nbt.2287.

9. Bhatt, D.K., and Daemen, T. (2024). Molecular Circuits of Immune Sensing and Response to
Oncolytic Virotherapy. IJMS *25*, 4691. https://doi.org/10.3390/ijms25094691.

10. Denton, N., Chen, C.-Y., Scott, T., and Cripe, T. (2016). Tumor-Associated Macrophages in
Oncolytic Virotherapy: Friend or Foe? Biomedicines *4*, 13.
https://doi.org/10.3390/biomedicines4030013.

11. Jakeman, P.G., Hills, T.E., Fisher, K.D., and Seymour, L.W. (2015). Macrophages and their
interactions with oncolytic viruses. Current Opinion in Pharmacology *24*, 23–29.
https://doi.org/10.1016/j.coph.2015.06.007.

12. Mealiea, D., and McCart, J.A. (2022). Cutting both ways: the innate immune response to
oncolytic virotherapy. Cancer Gene Ther *29*, 629–646. https://doi.org/10.1038/s41417-021-
00351-3.

13. Hofman, L., Lawler, S.E., and Lamfers, M.L.M. (2021). The Multifaceted Role of Macrophages in
Oncolytic Virotherapy. Viruses *13*, 1570. https://doi.org/10.3390/v13081570.

14. DeNardo, D.G., and Ruffell, B. (2019). Macrophages as regulators of tumour immunity and
immunotherapy. Nat Rev Immunol *19*, 369–382. https://doi.org/10.1038/s41577-019-0127-6.

15. Bhatt, D.K., Wekema, L., Carvalho Barros, L.R., Chammas, R., and Daemen, T. (2021). A
systematic analysis on the clinical safety and efficacy of onco-virotherapy. Molecular Therapy -
Oncolytics *23*, 239–253. https://doi.org/10.1016/j.omto.2021.09.008.

16. Li, Z., Jiang, Z., Zhang, Y., Huang, X., and Liu, Q. (2020). Efficacy and Safety of Oncolytic Viruses in
Randomized Controlled Trials: A Systematic Review and Meta-Analysis. Cancers (Basel) *12*.
https://doi.org/10.3390/cancers12061416.

17. Bhatt, D.K., Chammas, R., and Daemen, T. (2021). Resistance Mechanisms Influencing Oncolytic
Virotherapy, a Systematic Analysis. Vaccines *9*, 1166. https://doi.org/10.3390/vaccines9101166.

18. de Gruijl, T.D., Janssen, A.B., and van Beusechem, V.W. (2015). Arming oncolytic viruses to
leverage antitumor immunity. Expert Opin Biol Ther *15*, 959–971.
https://doi.org/10.1517/14712598.2015.1044433.

19. Kaufman, H.L., Kohlhapp, F.J., and Zloza, A. (2015). Oncolytic viruses: a new class of
immunotherapy drugs. Nature Reviews Drug Discovery *14*, 642–662.
https://doi.org/10.1038/nrd4663.

20. Smerdou, C., and Liljeström, P. (1999). Two-helper RNA system for production of recombinant
Semliki forest virus particles. J Virol *73*, 1092–1098. https://doi.org/10.1128/JVI.73.2.1092-
1098.1999.

21. Kurena, B., Müller, E., Christopoulos, P.F., Johnsen, I.B., Stankovic, B., Øynebråten, I., Corthay, A.,
and Zajakina, A. (2017). Generation and Functional In Vitro Analysis of Semliki Forest Virus
Vectors Encoding TNF-α and IFN-γ. Front. Immunol. *8*, 1667.
https://doi.org/10.3389/fimmu.2017.01667.

22. Sánchez-Paulete, A.R., Teijeira, Á., Quetglas, J.I., Rodríguez-Ruiz, M.E., Sánchez-Arráez, Á.,
Labiano, S., Etxeberria, I., Azpilikueta, A., Bolaños, E., Ballesteros-Briones, M.C., et al. (2018).
Intratumoral Immunotherapy with XCL1 and sFlt3L Encoded in Recombinant Semliki Forest Virus-
Derived Vectors Fosters Dendritic Cell-Mediated T-cell Cross-Priming. Cancer Res. *78*, 6643–
6654. https://doi.org/10.1158/0008-5472.CAN-18-0933.

23. Bhatt, D.K., Meuleman, S.L., Hoogeboom, B.N., and Daemen, T. (2024). Oncolytic alphavirus
replicons mediated recruitment and activation of T cells. iScience *27*, 109253.
https://doi.org/10.1016/j.isci.2024.109253.

24. Draghiciu, O., Boerma, A., Hoogeboom, B.N., Nijman, H.W., and Daemen, T. (2015). A rationally
designed combined treatment with an alphavirus-based cancer vaccine, sunitinib and low-dose
tumor irradiation completely blocks tumor development. Oncoimmunology *4*, e1029699.
https://doi.org/10.1080/2162402X.2015.1029699.

25. Ip, P.P., Boerma, A., Regts, J., Meijerhof, T., Wilschut, J., Nijman, H.W., and Daemen, T. (2014).
Alphavirus-based vaccines encoding nonstructural proteins of hepatitis C virus induce robust and
protective T-cell responses. Mol Ther *22*, 881–890. https://doi.org/10.1038/mt.2013.287.

26. Knudsen, M.L., Ljungberg, K., Tatoud, R., Weber, J., Esteban, M., and Liljeström, P. (2015).
Alphavirus replicon DNA expressing HIV antigens is an excellent prime for boosting with
recombinant modified vaccinia Ankara (MVA) or with HIV gp140 protein antigen. PLoS ONE *10*,
e0117042. https://doi.org/10.1371/journal.pone.0117042.

27. Öhlund, P., García-Arriaza, J., Zusinaite, E., Szurgot, I., Männik, A., Kraus, A., Ustav, M., Merits, A.,
Esteban, M., Liljeström, P., et al. (2018). DNA-launched RNA replicon vaccines induce potent anti-
Ebolavirus immune responses that can be further improved by a recombinant MVA boost. Sci
Rep 8, 12459. https://doi.org/10.1038/s41598-018-31003-6.

28. Quetglas, J.I., Labiano, S., Aznar, M.Á., Bolaños, E., Azpilikueta, A., Rodriguez, I., Casales, E.,
Sánchez-Paulete, A.R., Segura, V., Smerdou, C., et al. (2015). Virotherapy with a Semliki Forest
Virus–Based Vector Encoding IL12 Synergizes with PD-1/PD-L1 Blockade. Cancer Immunology
Research 3, 449–454. https://doi.org/10.1158/2326-6066.CIR-14-0216.

29. Ballesteros-Briones, M.C., Martisova, E., Casales, E., Silva-Pilipich, N., Buñuales, M., Galindo, J.,
Mancheño, U., Gorraiz, M., Lasarte, J.J., Kochan, G., et al. (2019). Short-Term Local Expression of
a PD-L1 Blocking Antibody from a Self-Replicating RNA Vector Induces Potent Antitumor
Responses. Mol Ther 27, 1892–1905. https://doi.org/10.1016/j.ymthe.2019.09.016.

30. Komdeur, F.L., Singh, A., van de Wall, S., Meulenberg, J.J.M., Boerma, A., Hoogeboom, B.N.,
Paijens, S.T., Oyarce, C., de Bruyn, M., Schuuring, E., et al. (2021). First-in-Human Phase I Clinical
Trial of an SFV-Based RNA Replicon Cancer Vaccine against HPV-Induced Cancers. Mol Ther 29,
611–625. https://doi.org/10.1016/j.ymthe.2020.11.002.

31. Eerkens, A.L., Esajas, M.D., Brummel, K., Vledder, A., Van Rooij, N., Plat, A., Avalos Haro, S.B.,
Paijens, S.T., Slagter-Menkema, L., Schuuring, E., et al. (2025). Vvax001, a Therapeutic Vaccine
for Patients with HPV16-positive High-grade Cervical Intraepithelial Neoplasia: a Phase II Trial.
Clinical Cancer Research. https://doi.org/10.1158/1078-0432.CCR-24-1662.

32. Chen, S. (2023). Manipulating immune cells — a therapeutic strategy for cancer and
inflammatory disease: https://doi.org/10.33612/diss.822658140.

33. Bhatt, D.K., Janzen, T., Daemen, T., and Weissing, F.J. (2022). Modelling the spatial dynamics of
oncolytic virotherapy in the presence of virus-resistant tumour cells. PLoS Comput Biol 18,
e1010076. https://doi.org/10.1371/journal.pcbi.1010076.

34. Bhatt, D.K., Janzen, T., Daemen, T., and Weissing, F.J. (2024). Effects of virus-induced
immunogenic cues on oncolytic virotherapy. Sci Rep 14, 28861. https://doi.org/10.1038/s41598-
024-80542-8.

35. Altan-Bonnet, G., and Mukherjee, R. (2019). Cytokine-mediated communication: a quantitative
appraisal of immune complexity. Nat Rev Immunol 19, 205–217.
https://doi.org/10.1038/s41577-019-0131-x.

36. Eddy, J.A., Thorsson, V., Lamb, A.E., Gibbs, D.L., Heimann, C., Yu, J.X., Chung, V., Chae, Y., Dang,
K., Vincent, B.G., et al. (2020). CRI iAtlas: an interactive portal for immuno-oncology research.
F1000Res 9, 1028. https://doi.org/10.12688/f1000research.25141.1.

37. Kurena, B., Müller, E., Christopoulos, P.F., Johnsen, I.B., Stankovic, B., Øynebråten, I., Corthay, A.,
and Zajakina, A. (2017). Generation and Functional In Vitro Analysis of Semliki Forest Virus
Vectors Encoding TNF-α and IFN-γ. Frontiers in Immunology 8.
https://doi.org/10.3389/fimmu.2017.01667.

38. Weigelin, B., den Boer, A.Th., Wagena, E., Broen, K., Dolstra, H., de Boer, R.J., Figdor, C.G.,
Textor, J., and Friedl, P. (2021). Cytotoxic T cells are able to efficiently eliminate cancer cells by
additive cytotoxicity. Nat Commun 12, 5217. https://doi.org/10.1038/s41467-021-25282-3.

39. Masemann, D., Köther, K., Kuhlencord, M., Varga, G., Roth, J., Lichty, B.D., Rapp, U.R., Wixler, V.,
and Ludwig, S. (2018). Oncolytic influenza virus infection restores immunocompetence of lung
tumor-associated alveolar macrophages. OncoImmunology 7, e1423171.
https://doi.org/10.1080/2162402X.2017.1423171.

40. Pulaski, B.A., Smyth, M.J., and Ostrand-Rosenberg, S. (2002). Interferon-gamma-dependent
phagocytic cells are a critical component of innate immunity against metastatic mammary
carcinoma. Cancer Res 62, 4406–4412.

41. Van Den Bossche, W.B.L., Kleijn, A., Teunissen, C.E., Voerman, J.S.A., Teodosio, C., Noske, D.P.,
Van Dongen, J.J.M., Dirven, C.M.F., and Lamfers, M.L.M. (2018). Oncolytic virotherapy in
glioblastoma patients induces a tumor macrophage phenotypic shift leading to an altered
glioblastoma microenvironment. Neuro-Oncology 20, 1494–1504.
https://doi.org/10.1093/neuonc/noy082.

42. Meisen, W.H., Wohleb, E.S., Jaime-Ramirez, A.C., Bolyard, C., Yoo, J.Y., Russell, L., Hardcastle, J.,
Dubin, S., Muili, K., Yu, J., et al. (2015). The Impact of Macrophage- and Microglia-Secreted TNFα
on Oncolytic HSV-1 Therapy in the Glioblastoma Tumor Microenvironment. Clin Cancer Res 21,
3274–3285. https://doi.org/10.1158/1078-0432.CCR-14-3118.

43. Saha, D., Martuza, R.L., and Rabkin, S.D. (2017). Macrophage Polarization Contributes to
Glioblastoma Eradication by Combination Immunovirotherapy and Immune Checkpoint
Blockade. Cancer Cell 32, 253-267.e5. https://doi.org/10.1016/j.ccell.2017.07.006.

44. Fulci, G., Dmitrieva, N., Gianni, D., Fontana, E.J., Pan, X., Lu, Y., Kaufman, C.S., Kaur, B., Lawler,
S.E., Lee, R.J., et al. (2007). Depletion of Peripheral Macrophages and Brain Microglia Increases
Brain Tumor Titers of Oncolytic Viruses. Cancer Research 67, 9398–9406.
https://doi.org/10.1158/0008-5472.CAN-07-1063.

45. Thoresen, D., Wang, W., Galls, D., Guo, R., Xu, L., and Pyle, A.M. (2021). The molecular
mechanism of RIG-I activation and signaling. Immunol Rev 304, 154–168.
https://doi.org/10.1111/imr.13022.

46. Rojas, J.J., Sampath, P., Bonilla, B., Ashley, A., Hou, W., Byrd, D., and Thorne, S.H. (2016).
Manipulating TLR Signaling Increases the Anti-tumor T Cell Response Induced by Viral Cancer
Therapies. Cell Rep 15, 264–273. https://doi.org/10.1016/j.celrep.2016.03.017.

47. Muñoz-Rojas, A.R., Kelsey, I., Pappalardo, J.L., Chen, M., and Miller-Jensen, K. (2021). Co-
stimulation with opposing macrophage polarization cues leads to orthogonal secretion programs
in individual cells. Nat Commun 12, 301. https://doi.org/10.1038/s41467-020-20540-2.

48. Piccolo, V., Curina, A., Genua, M., Ghisletti, S., Simonatto, M., Sabò, A., Amati, B., Ostuni, R., and
Natoli, G. (2017). Opposing macrophage polarization programs show extensive epigenomic and
transcriptional cross-talk. Nat Immunol 18, 530–540. https://doi.org/10.1038/ni.3710.

49. Stout, R.D., Jiang, C., Matta, B., Tietzel, I., Watkins, S.K., and Suttles, J. (2005). Macrophages
Sequentially Change Their Functional Phenotype in Response to Changes in Microenvironmental
Influences. The Journal of Immunology 175, 342–349.
https://doi.org/10.4049/jimmunol.175.1.342.

50. Lavin, Y., Winter, D., Blecher-Gonen, R., David, E., Keren-Shaul, H., Merad, M., Jung, S., and Amit,
I. (2014). Tissue-Resident Macrophage Enhancer Landscapes Are Shaped by the Local
Microenvironment. Cell 159, 1312–1326. https://doi.org/10.1016/j.cell.2014.11.018.

51. Trofimova, O., Korotkaja, K., Skrastina, D., Jansons, J., Spunde, K., Isaguliants, M., and Zajakina, A.
(2021). Alphavirus-Driven Interferon Gamma (IFNg) Expression Inhibits Tumor Growth in
Orthotopic 4T1 Breast Cancer Model. Vaccines (Basel) *9*, 1247.
https://doi.org/10.3390/vaccines9111247.

52. Müller, E., Speth, M., Christopoulos, P.F., Lunde, A., Avdagic, A., Øynebråten, I., and Corthay, A.
(2018). Both Type I and Type II Interferons Can Activate Antitumor M1 Macrophages When
Combined With TLR Stimulation. Front. Immunol. *9*, 2520.
https://doi.org/10.3389/fimmu.2018.02520.

53. Meissner, R., Wixler, V., Wulfert, F.P., Jacob, J.C., Hale, B.G., Robeck, T., Masemann, D.,
Boergeling, Y., and Ludwig, S. (2023). Replication-incompetent influenza A viruses armed with
IFN-γ effectively mediate immune modulation and tumor destruction in mice harboring lung
cancer. Molecular Therapy - Oncolytics *31*, 100741.
https://doi.org/10.1016/j.omto.2023.100741.

54. Bourgeois-Daigneault, M.-C., Roy, D.G., Falls, T., Twumasi-Boateng, K., St-Germain, L.E.,
Marguerie, M., Garcia, V., Selman, M., Jennings, V.A., Pettigrew, J., et al. (2016). Oncolytic
vesicular stomatitis virus expressing interferon-σ has enhanced therapeutic activity. Molecular
Therapy - Oncolytics *3*, 16001. https://doi.org/10.1038/mto.2016.1.

55. Singh, A., Koutsoumpli, G., Van De Wall, S., and Daemen, T. (2019). An alphavirus-based
therapeutic cancer vaccine: from design to clinical trial. Cancer Immunol Immunother *68*, 849–
859. https://doi.org/10.1007/s00262-018-2276-z.

56. Groeneveldt, C., Kinderman, P., Van Den Wollenberg, D.J.M., Van Den Oever, R.L., Middelburg, J.,
Mustafa, D.A.M., Hoeben, R.C., Van Der Burg, S.H., Van Hall, T., and Van Montfoort, N. (2020).
Preconditioning of the tumor microenvironment with oncolytic reovirus converts CD3-bispecific
antibody treatment into effective immunotherapy. J Immunother Cancer *8*, e001191.
https://doi.org/10.1136/jitc-2020-001191.

57. Groeneveldt, C., van Ginkel, J.Q., Kinderman, P., Sluijter, M., Griffioen, L., Labrie, C., van den
Wollenberg, D.J.M., Hoeben, R.C., van der Burg, S.H., ten Dijke, P., et al. (2023). Intertumoral
Differences Dictate the Outcome of TGF-β Blockade on the Efficacy of Viro-Immunotherapy.
Cancer Research Communications *3*, 325–337. https://doi.org/10.1158/2767-9764.CRC-23-0019.

58. Groeneveldt, C., Kinderman, P., Griffioen, L., Rensing, O., Labrie, C., van den Wollenberg, D.J.M.,
Hoeben, R.C., Coffey, M., Loghmani, H., Verdegaal, E.M.E., et al. (2024). Neutralizing Antibodies
Impair the Oncolytic Efficacy of Reovirus but Permit Effective Combination with T cell–Based
Immunotherapies. Cancer Immunology Research *12*, 334–349. https://doi.org/10.1158/2326-
6066.CIR-23-0480.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

(A) rSFV **+** Monolayer coculture of macrophages and cancer cells $\xrightarrow{\text{8 hrs}}$ **+** PBMCs $\xrightarrow{\text{16 hrs}}$ Flow cytometry to measure T cell activation

(B) CD4+ T cells

(C) CD8+ T cells

(D) %CD69+CD107a+CD38+ (CD4+ PBMCs)

Coculture with PANC-1 — Coculture with Ca-Ski

rSFV-IFN-γ, rSFV-GFP, NT

(E) %CD69+CD107a+CD38+ (CD8+ PBMCs)

Coculture with PANC-1 — Coculture with Ca-Ski

rSFV-IFN-γ, rSFV-GFP, NT

Figure 6

Figure 7

Figure 8