Title: Mobile genetic elements that shape microbial diversity and functions in thawing permafrost soils

Running title: The Stordalen Mire Mobilome

Authors

Jiarong Guo[1,2,8], Samuel Aroney[3,8], Guillermo Dominguez-Huerta[1,2,8], Derek Smith[3,8], Dean Vik[2,8], Carlos Owusu Ansah[1,8], Akbar Adjie Pratama[1,8], Sergei Solonenko[1,8], Funing Tian[1,8], Cristina Howard-Varona[1,8], ZhiPing Zhong[1,2,5], Aminata Fofana[1,8], Garrett Smith[2], Suzanne B. Hodgkins[1,8], Dylan Cronin[1,8], EMERGE Field Teams 2010-2019, EMERGE Coordinators, Ben J. Woodcroft[3,8], Gene W. Tyson[3,8], Virginia I. Rich[1,8], Matthew B. Sullivan[1,2,5,7,8]*, Simon Roux[6,8]*, Sarah C. Bagby[3,8]*

Affiliations

[1]Department of Microbiology, The Ohio State University; Columbus, OH, USA.
[2]Center of Microbiome Science, The Ohio State University; Columbus, OH, USA.
[3]Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of Technology, Translational Research Institute; Woolloongabba, QLD, Australia.
[4]Department of Biology, Case Western Reserve University; Cleveland, OH, USA.
[5]Byrd Polar and Climate Research Center, Ohio State University; Columbus, OH, USA.
[6]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory; Berkeley, CA, USA.
[7]Department of Civil, Environmental, and Geodetic Engineering, The Ohio State University; Columbus, OH, USA.
[8]National Science Foundation EMERGE Biology Integration Institute, The Ohio State University, Columbus, OH, USA
*Co-corresponding authors. MBS: sullivan.948@osu.edu, SR: sroux@lbl.gov, SCB: scb126@case.edu

One Sentence Summary

By developing microbiome-ready analytics and applying them to thawing permafrost microbiomes, we identified ~2.1 million diverse mobile genetic element (MGE) recombinases (integrons, insertion sequences, transposons, conjugative elements, phage-like elements) and revealed their function and activity to establish baseline ecological data and a framework for MGE study in complex natural systems.

Abstract

Microbiomes shape ecosystems through functional profiles influenced by gene gains and losses. While culture-based experiments demonstrate that mobile genetic elements (MGEs) can mediate gene flux, quantitative field data remains scarce. Here we leverage large-scale soil meta-omic data to develop and apply analytics for studying MGEs. In our model permafrost-thaw ecosystem, Stordalen Mire, we identify ~2.1 million MGE recombinases across 89 microbial phyla to assess ecological distributions, affected functions, past mobility, and current activity. This revealed

45 MGEs shaping natural genetic diversity via lineage-specific engagement, and functions including
46 canonical defense elements alongside diverse regulatory and metabolic genes affecting carbon flux
47 and nutrient cycling. These findings and this systematic meta-omic framework open new avenues
48 to better investigate MGE diversity, activity, mobility, and impacts in nature.
49
50

Main text

Microbiomes are increasingly recognized for their roles in biogeochemical and nutrient
cycling across diverse systems (*1–4*). Mobile genetic elements (MGEs) likely alter these microbial
impacts since culture-based studies show impacts such as horizontal gene transfer and/or genomic
rearrangement (*5*, *6*), as well as MGEs encoding key niche-defining genes involved in
antimicrobial resistance (AMR) (*7–9*), production of public goods (*10*), or defense against
microbes, phages or other mobile elements (*11–13*). Further, MGEs are highly abundant and
diverse with recent estimates of ~25 MGEs per genome across ~84K cultured bacterial genomes,
with broad MGE distributions across phyla (*8*). However, outside of cultivated isolates, the
ecological and evolutionary effects of MGEs in nature remain poorly understood. This is because
the same diversity and activity that drive MGE impacts also hamper metagenomic investigation—
MGEs lacked well-developed marker genes until recently (*8*), microbial populations must be
deeply and regularly sequenced to capture elements that move between subpopulations over time,
and metagenomic assemblies tend to break when the same MGE is sampled in multiple contexts
(*14*, *15*). As a result, with rare host-microbiome exceptions (*16*), MGEs in nature have largely been
studied only one type at a time (*17–19*), which limits context for understanding their collective
prevalence, distribution, and ecological impacts.
The potential scope for MGE impacts in natural systems is vast. MGEs can influence host
phenotypes and niches through genes they carry (cargo) and/or the host genomic context of MGE
insertion (**Fig. 1A**). MGE movements cause gains or losses of function and/or regulatory shifts
(*20*, *21*) that should scale and diversify with the number of such movements (**Fig. 1B**). The wider
an MGE's host range is, the greater is its range of partners—both hosts and other MGEs—for
exchanging genetic modules (*22*). But which genes are affected, and what are the consequences?
Do MGEs substantially affect just the social functions intensively investigated in cultivated
isolates (reviewed above), or might recently observed plasmid-borne genes affecting host
"domestic" activities (e.g., stress response, nitrogen fixation) (*23*) be the tip of the iceberg for
MGE-influenced processes in nature? Because such domestic activities directly affect microbial
ecosystem outputs and adaptability, *in natura* mobilome activity and impacts constitute a
knowledge gap that critically impairs our ability to understand and model microbial ecosystem
processes and response to change.
Here, we combine recent literature-provided MGE recombinase identification approaches
(*8*), a sufficiently large longitudinal metagenomic dataset, and novel analytics development to
yield broad new insights into their ecology and ecosystem impacts. We examined ~5.8 terabases
of sequencing data collected over eight years along a permafrost peatland thaw gradient in
Stordalen Mire, Sweden (**Fig. 2A**), a climate-critical, long-term ecological research site (*24–26*).
We reasoned that a site undergoing rapid, landscape-scale change might be an observatory for
mobilome-mediated microbiome adaptation to changes in a range of key environmental variables.
From these data, we identified, characterized, and genomically and ecologically contextualized
~2.1M MGE recombinases to not only deepen our understanding of how the microbiome of
thawing permafrost evolves, but also establish a conceptual and analytic foundation for studying

all known, major MGE types in any complex system.
92
MGE recombinases from permafrost thaw gradient microbiomes
Stordalen Mire, as introduced in companion papers (*27–29*), is a climate-critical permafrost
peatland, extensively studied for decades to assess the interplay of vegetation, microbiota, and
climate along a three-stage thaw gradient of palsa, bog, and fen habitats that represent,
respectively, the least to most thawed soils. Prior to our work, ~5.8 terabases of metagenomic and
metatranscriptomic sequencing data had been generated from 8 years of field sampling (**Fig. 2A,**
table S1). We first identified and quantified all major MGEs in these data via recently developed
hidden Markov models (HMMs) and MGE identification protocols that leverage recombinases as
MGE hallmark genes (*8*) (**table S2**). Specifically, we screened 131.5M genes predicted from
30.7M short-read metagenomic contigs, and this identified 2,107,204 MGE recombinases
(1,130,776 unique protein sequences) derived from 1,790,672 contigs (**Fig. 2B,C&D, fig. S1, fig.**
S2A&B, fig. S3A&B, table S3 item A). On average, this represented 17.3 MGE recombinases
per million base pairs (Mbp) of microbiome sequence data, dwarfing the representation found for
even the most abundant KEGG gene family, the stress-response sigma factor *rpoE* (2.4
copies/Mbp; **Fig. 2C**).
We next classified these abundant recombinases into four previously-established types (*8*):
(i) integrons, (ii) insertion sequences and transposons, herein "IS_Tn", (iii) phage / phage-like
elements, herein "Phage", and (iv) conjugative elements and mobility islands, herein "CE".
Classification facilitates evaluation of ecological impacts, since these types are known to differ in
length, mode of propagation, and gene content. Prior work evaluating 187 metagenomes available
in 2010 and >80K bacterial and archaeal isolate genomes (*8, 30*) found IS_Tn dominant in highly
abundant MGE pools. In Stordalen Mire, we also found thatIS_Tn was the most commonly
recovered type (58.6%), distantly followed by Phage (16.1%), CE (2.8%), and integron (1.2%)
(the remaining 21.3% were ambiguous; see Methods; **fig. S2A**). Ecologically, at this high level of
recombinase types, this rank-order pattern remained remarkably constant across time (8 years of
peak growing season sampling) and space (within and between the study site's three habitat types)
(**Fig. 2D, fig. S2B, fig. S2C**). The same pattern largely persisted across phyla for both MAG- and
contig-level taxonomic assignments (216,720 and 804,456 MGE recombinases, respectively,
assigned to a total of 89 phyla with 93.7% agreement in assignments for the subset shared; **Fig.**
3A,B, **fig. S4A,B, table S3 item A**). MGE distribution broadly followed host lineage relative
abundances, with most MGEs in ecosystem-dominant bacterial and archaeal phyla such as
Acidobacteriota, Actinomycetota, Pseudomonadota, Verrucomicrobiota, and Halobacteriota (*25*)
(**Fig. 3**, **fig. S4&S5**). Echoing isolate-based observations (*8*), the virtual absence of integrons in
the abundant Actinomycetota stands out against a background of low cross-phylum variation in
the frequency of each MGE type in our Stordalen microbiomes.
Next, we wondered how the MGE frequency of ~25 MGEs/genome measured in isolates
would compare with estimates from natural populations, given the potential for MGEs to be lost
during laboratory cultivation due to reduced selective pressure in clonal isolates. To this end, we
quantified MGEs per genome via two approaches applied to our Stordalen microbiomes. First, we
used a contig-based estimate, comparing identified MGE recombinase counts to total genome
counts estimated using a standard single-copy marker gene approach for each sample (*31, 32*).
This revealed a median of 36 MGE recombinases per genome in Stordalen microbiomes (**Fig. 2E**).
Separately, we considered only the subset of MGEs that were binned into medium/high-quality
metagenome-assembled genomes (MAGs) (≥70% completeness, ≤10% contamination) (**table S3**

item **B**). Scaling per-MAG MGE counts by MAG completeness revealed a median of 23 MGE recombinases per genome (**Fig. 2E**). Comparing the two approaches, we note that only 15.9% of MGE-bearing contigs were binned, even though Stordalen MAGs capture a remarkable 73% of sequence reads at genus level (**REF temporal paper**). Thus, we interpret the MAG-based estimate as a lower bound since it empirically captures only a subset of the contig-observed MGEs, and dynamic rearrangements in MGE-bearing genomic regions are a known challenge for short-read metagenome assembly and binning (*33*, *15*, *34*).

To empirically quantify the impact of assembly and binning challenges on short-read counts, we performed long-read sequencing on six samples and evaluated MGE recombinase recovery across the assembly and binning process. In total, these data suggested that short-read assembly and binning recovered 35–60% of MGE recombinases, with IS_Tn recombinases disproportionately under-recovered (**Fig. 2F**, **fig. S6&S7, Supplementary text**). Using this benchmarking to scale our MAG-based counts, we obtain a revised estimate of 38–66 MGE recombinases per genome *in situ*. We interpret this number as comparable with the contig-based estimate and consistent with the hypothesis that isolates might lose MGEs during lab cultivation.

MGE impacts range far beyond biotic social interactions

We next wondered whether MGEs affected genes of ecological importance, and how these fit into microbiome activity in Stordalen Mire. Prior genome-resolved metagenomic analyses (*25*, *28*) show that intercellular interactions and public goods matter in this system: cross-feeding relationships between carbon generalists and methanogens are persistent and strongly associated with methane flux, and extracellular degradation of polysaccharides is a bottleneck in carbon flow. Beyond microbes, there are also abundant viral community members that harbor auxiliary metabolic genes that likely mediate exopolysaccharide breakdown (*26*). Such outward-directed social functions are expected to be affected by MGEs, as are some stress responses (*23*), but these are only a small slice of microbial activities. Each cell must also manage its own routine "domestic" processes (e.g., genome maintenance, gene expression, nutrient homeostasis) (*35–38*), functions that shape physiology and ecosystem outputs but that have been little studied in the mobilome to date.

Thus, we asked which Stordalen Mire functions could be confidently detected as mobilome-affected (i.e., encoded in MGEs or interrupted by them) and how far beyond biotic interactions the mobilome's direct influence extends. To ensure that host gene activities were not misattributed to MGEs, we adopted a highly conservative approach for curating MGE boundaries. While this permitted assessment of only ~0.8% of the total 2.1M MGE recombinases identified, annotation identified putative functions for genes within conservatively defined boundaries of 5,288 cargo-carrying MGEs (900 CEs, 551 integrons, and 3,837 phage) and for host genes interrupted by 11,295 IS_Tns. (**fig. S1, Methods**). Among this set of MGEs, our manually curated annotation identified high-confidence functional predictions for 3,631 MGE-affected genes with 1,199 distinct KEGG annotations (**table S3 item C-F**, **table S4, Supplementary text**).

As expected, this reduced set of curated MGE-affected genes included many related to host biotic interactions (n=1377; **Fig. 4, table S4**). This included diverse phage defense systems (379 carried, 168 interrupted) that can abort infection, degrade phage DNA by restriction-modification, or inhibit phage DNA synthesis by nucleotide depletion (dCTP deaminase) (*12*, *39*). MGEs also impact genes for YD-repeat-containing proteins (10 carried, 29 interrupted), a contact-dependent growth inhibitor thought to shape responses to high population densities (*40*). In social interactions mediated by public goods, MGEs impact genes encoding carbohydrate-active enzymes

(CAZymes) involved in synthesis or degradation of extracellular polysaccharides (154 carried, 125 interrupted). The former produce public goods and molecules involved in cell-phage and cell-cell interactions, while the latter can become public goods through direct secretion (extracellular enzymes) or outer membrane vesicle budding (periplasmic enzymes) (*41*). Separately, MGEs appear to impact synthesis and transport of other public goods (39 carried, 38 interrupted), predominantly siderophores. Finally, MGEs can impact genes that encode a public service (2 carried, 13 interrupted), the detoxification of diffusible peroxides (*42*).

Beyond biotic interactions, we identified diverse MGEs seemingly affecting domestic processes. For example, we uncovered MGEs affecting central carbon metabolism genes (6 carried, 44 interrupted; **table S4**). Intriguingly, even in this small subset of conservatively-curated MGEs, there were three direct connections to methane metabolism: the key hydrogenotrophic metabolism gene formate dehydrogenase (*43*) was carried by a Phage from Halobacteriota (*Methanoregula* sp003136075), a group of methanogens active at Stordalen Mire (*24*, *25*) (**REF Temporal Paper**), and IS_Tns interrupted a methane monooxygenase (*pmoC*) subunit in a Pseudomonadota (unbinned Alphaproteobacterial contig) and the methylamine utilization gene *mauG* in a Verrucomicrobiota (unbinned UBA11358 contig). Thus, even within this highly conservative subset of MGE-affected genes, we find genomic evidence that MGEs may alter Stordalen methane fluxes both directly (by changing genomic potential for methanogenesis and methanotrophy) and indirectly (by changing the distribution of carbon-processing pathways that control the flow of substrates to methanogens).

MGEs also might influence other major nutrient cycles, trace metal cycles, and cellular housekeeping functions. In nitrogen cycling, we found MGEs influencing nitrogen processing (26 carried, 58 interrupted; **Fig. 4, table S4**) across the thaw gradient, predominantly in amino acid metabolism but also touching other nitrogen processes, including fixation. Because fixed nitrogen is thought to be a partially "leaky" public good (*44*), IS_Tn interruption of an Acidobacterial nitrogen fixation regulator (*45*) illustrates the potential for MGEs influencing growth beyond the host cell by controlling nitrogen. The CAZyme genes highlighted above include 10 chitinase genes carried and 1 interrupted by MGEs; chitin is hypothesized to serve as a key alternative nitrogen source at our N-limited site (*28*). In sulfur cycling, we found MGEs that affect genes putatively related to inorganic sulfur limitation (19 carried, 15 interrupted), including sulfatases that might degrade sulfated polysaccharides (*36*) and cysteine desulfurases and selenocysteine lyases that might scavenge thiol groups and reduce sulfur demand (*46*, *47*). Beyond macronutrient cycling, MGEs impact iron cycling (39 carried, 37 interrupted) via siderophore production and transport, roles highlighted above in connection with public goods. Finally, MGEs also encode core housekeeping genes involved in genome maintenance: DNA repair, predominantly *radC* (89 carried, 49 interrupted; genome replication, predominantly *dinB* and *polC* (79 carreid, 15 interrupted); regulation of host gene transcription (RNA polymerase subunits, transcriptional regulators; 174 carried, 135 interrupted), and translation (ribosomal proteins and one elongation factor; 32 carried, 14 interrupted); and proteome maintenance (predominantly chaperonin, ubiquitin activation, and proteases; 104 carried, 53 interrupted).

Critically, even though our stringent analysis criteria reduced the set of MGE-affected genes analyzed to 0.02% of the total MGEs identified, we still find abundant examples of mobilome-affected social and domestic functions beyond previous observations (*48*), and add many further examples of MGE-impacted functions including those that influence carbon flow through this climate-critical ecosystem, both directly (exopolysaccharide degradation, central carbon metabolism) and indirectly (via nitrogen, sulfur, and/or iron availability to support carbon

flux). Although research to date has focused on MGE impacts on biotic interactions, MGE impacts on domestic functions appear relatively common, at the level of both the functions (~51% domestic vs. ~38% social; the remainder play both roles, support MGE maintenance and replication, or are uncategorized) and the MGEs themselves (~55% affect at least one domestic function). If this conservative subset is representative of the Stordalen mobilome as a whole, then as sequencing methods, functional annotation, and databases improve, we would expect ~1.1M of the full ~2.1M MGE recombinases to influence domestic functions, with significant implications for nutrient cycling, as well as microbiome activity, ecology, and evolution.

MGE recombinase mobility and activity

To assess ecological and evolutionary change in Stordalen Mire MGEs, we evaluated their mobility and activity from three different viewpoints: (i) snapshots of metatranscriptomic MGE recombinase expression as a proxy for mobilization underway, (ii) genomic neighborhood change as a marker of completed MGE movement between insertion sites, and (iii) partial carriage of an MGE by the host population as the result of evolution and selection over longer time scales. Integrating results across these analyses can help highlight individual MGE dynamics. For example, high recombinase expression rates together with low rates of genomic neighborhood change for a given MGE would suggest a strong preference among insertion sites. Our overall findings are as follows.

First, we asked whether MGE recombinases were "transcriptionally active" as measured via read mapping from 50 available metatranscriptomes against contigs from paired metagenomes (**table S5**), taking ribosomal gene mapping rates as a baseline for constitutive transcriptional activity of actively growing cells (*49*) (**table S6**). This revealed transcriptional activity in ~3.4–56.4% (median: 27.8%) of ribosomal genes and ~0.04–9.1% (median: 1.2%) of MGE recombinase genes (**Fig. 5A, table S3 item G**), making MGE recombinases the second-most-transcribed non-ribosomal genes in Stordalen Mire (**Fig. 5A**). Stratified by recombinase type, we found roughly equal levels of transcriptional activity for IS_Tn and Phage recombinases versus ~five- to ten-fold lower integron and CE recombinase transcription (**Fig. 5B**). Ecologically, IS_Tn recombinase transcription increased along the thaw gradient (significant from palsa to fen, $p \leq 0.01$, Wilcoxon rank-sum test) (**Fig. 5C, fig.S8B**), suggesting increased MGE activity in the more-disturbed fen. Taxonomically, transcriptionally active MGE recombinases were detected across hosts spanning 30 phyla with activity largely proportional to MGE- and phylum abundance (**Fig. 5D, fig. S8C**). However, one of these, the archaeal phylum *Halobacteriota,* was unique in that it is neither abundant nor a MGE recombinase-rich phylum, and yet was the most active phylum for MGE recombinase transcription. Given that Stordalen Mire *Halobacteriota* are known to be methanogens that are active and selected for in thawed fen and bog habitats (*24, 50*), we speculate that their disproportionate MGE recombinase activity might be adaptive.

Second, where MGE recombinases were observed multiple times, we assessed the genomic context (300 bp upstream and downstream) surrounding each instance. We then interpreted this genomic context as a proxy for intra/inter-genomic mobility, either of the recombinase itself or of other recombinase-proximal genes (see examples of changing neighborhoods in **fig. S9, Methods**). From the original ~2.1M MGE recombinases, we identified 74,257 MGE recombinase clusters (100% average amino acid identity [AAI])) observed more than once in our dataset. Among these, most had identical genomic neighborhoods across observations, but some (0.1-10%, depending on MGE type) did show neighborhood variation (**Fig. 5B**, **table S3 item H**). Neighborhood variability was most common among IS_Tn recombinases (~10%), and much less so for Phage (~1%),

integron (0.2%) and CE (0.1%) recombinases. This was consistent among years and habitats, with larger variation detected across diverse host phyla (up to 37% in Desulfobacterota_G) (**fig. S10**). Mindful that hypervariable islands in cellular genomes can often be impacted by multiple MGE types, we quantified co-occurring IS_Tn insertions, which could cause the neighborhood to change while the focal MGE remains stable. Fewer than 1 in 5 of our changed-genomic-neighborhood examples was due to co-occurring IS_Tn insertions (**fig. S9**). Considering our approach only detects insertion in immediate neighborhoods of MGE recombinase, this is a lower bound estimate of mobility at hypervariable multiple-MGE genomic islands among MGE context changes at Stordalen.

Third, we sought to assess what fraction of a population contains any given MGE recombinase by evaluating read-mapping-based coverage of the recombinase against that of its surrounding genomic context (see Methods). Where MGE recombinase coverage was significantly lower, we infer that only a subset of the host population's genotypes encoded the MGE recombinase, presumably a function of MGE mobility. Overall, across MGE types, we found 8,033 recombinase clusters (100% AAI) that had ≥20 kbp of genomic context. Of these, ~5–23% showed variation in this genomic context consistent with at least part of the cellular population having the MGE and part not having the MGE (**Fig. 5B, table S3 item I**). These findings were again broadly consistent across years, habitats, and well-sampled lineages (**fig. S11B-D**). In spite of spatio-temporal consistencies, there were notable variations in intrapopulation penetrance between MGE types, with IS_Tn and Phage particularly prone to varying across individuals within a host population (**fig. S11D**). Because MGE sizes vary and read mapping for lesser-sampled taxa is noisy, these numbers represent conservative lower bounds on active MGE penetrance into cellular population genomes. Even so, the extent of partial carriage observed highlights MGEs' role in what amounts to large ongoing *in situ* knockout experiments: among 372 IS_Tn elements that both interrupt a protein-coding gene and were amenable to partial carriage analysis, 81 (~22%) variably interrupt or leave intact host protein-coding genes. Together, these results suggest that MGE penetrance within member genomes of any given population differentially provide individuals access to genetic diversity, while shielding the population from loss of function due to MGE insertion (*51*, *52*).

Conclusions

MGEs have long been known from cultivated isolate genomes to be important drivers of strain-level variation and key functional adaptations, particularly in biomedical contexts related to AMR (*7–9*). However, virtually no data exist for diverse MGE types in natural settings. Our analyses presented here for Stordalen Mire microbiomes illuminate the mechanisms and scope of MGE impact on a climate-sensitive ecosystem by showing that, well beyond biotic interactions, the mobilome affects functions intrinsic to a wide range of physiologically and ecologically significant processes. Strikingly, the IS_Tn elements that heavily dominate recovered MGE recombinases in our dataset are also the most active by two of our three measures (i.e., transcription and genomic neighborhood change) (**Fig. 5B**), highlighting the need for future studies to examine context-dependent MGE effects like modulation of host gene expression downstream of MGE insertion sites.

As improved methods for studying the genomic potential of microbes in nature come on-line, the field has progressed from community-level taxonomic and functional descriptions towards species-resolved links between taxa and their functions (*25*, *31*). Continued advances in sequencing and analytical approaches are increasing this resolution to reveal intra-population

variation, early speciation, and selection (*53*) with recent MGE typing efforts (*8*) and further analytical advances made here finally bringing a quantitative understanding of the mobilome within reach, even in complex natural systems. This work benchmarks sequencing artifacts and provides empirical baselines and inferred impacts for all major MGEs in a natural soil system. These findings suggest that ~1% of MGE recombinases are active, genomes contain 36–66 MGE recombinases on average, and that MGEs impact (via carriage or interruption) diverse ecosystem-relevant functions. As technologies advance, the mobilome will come more fully into focus, and, given this new toolkit and baseline findings, we predict it will help reveal a subtler layer of change that could serve as the canary in the coalmine for early detection of change.

References

(See the full list at the end of this doc)

Acknowledgements

Non-author contribution: we thank Ohio Supercomputer Center for providing computing resources.

Funding: this work is supported by following sources of funding:
National Science Foundation, Biology Integration Institutes Program, Award # 202207. The work conducted by the U.S. Department of Energy Joint Genome Institute (JGI; https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231. This work by JGI includes BER Support Science Proposal 503530 (used for the majority of samples sequenced by JGI), and Facilities Integrating Collaborations for User Science (FICUS) initiative proposal 503547 (used for sequencing samples from the SIP experiment).

Author contribution: J.G., S.R., M.B.S., and S.C.B. conceptualized, designed, coordinated the study. S.A., J.G. performed long read and short read comparison. S.A., A.F., J.G. performed MGE recombinase host lineage analyses. G.D.-H., D.S., D.V., S.C.B., A.A.P., G.S., F.T., C.H.-V., S.S., Z.Z., C.O.A., J.G. performed MGE impacted function analyses. D.C., S.A., S.B.H. performed metagenome assembly and binning. J.G., S.R., M.B.S., and S.C.B. drafted the manuscript with contribution from B.J.W., G.W.T., V.I.R. All authors read, edited, commented on, and approved the final manuscript.

Competing interests: The authors declare that they have no competing interests.
Data and material availability: The authors declare that all the data in this study are freely and publicly available without restriction. The metagenomic and metatranscriptomic raw reads are available in SRA / ENA under Umbrella BioProject PRJNA888099 with individual accessions cited in **table S1**. The data provided by the DOE Joint Genome Institute were generated under proposals "10.46936/10.25585/60001148" and "10.46936/fics.proj.2017.49950/60006215". MAGs used in this study are available in Zenodo with https://doi.org/10.5281/zenodo.13984630 (main set of 36,419 MAGs) and https://doi.org/10.5281/zenodo.14207415 (733 short-read, long-read, and hybrid MAGs from six paired samples). The paired long read and short read data from six samples are available in PRJNA888099 with accessions cited in **table S7**. Processed data

including tables for MGE recombinase identification, activity by metatranscriptome, mobility by
genomic neighborhood change, and partial carriage, and MGE impacted functions are available
through supplementary tables and a large file data package shared through CERN's Zenodo
repository (https://zenodo.org/records/14287562). The code produced in this study is also
included in the above repository, which is an archive of the latest version of the project's GitHub
repository (https://github.com/emerge-bii/mobilome-submit-202407).
373
374

Supplementary Materials
376 Methods and Materials
377 Supplementary Text
378 Figs. S1 - S11
379 Tables S1 - S7
380 References (*54 - 102*)
381
382

Figures
384

385
386
Figure 1. MGE cargo and context determine the impact of MGE activity on host coding
capacity and population genetic diversity. A. An MGE may or may not carry cargo and may
insert either intergenically, potentially altering transcriptional regulation downstream, or into a
host gene, causing function loss. MGE and recombinase gene product, red and white waves;
cargo gene and product, solid teal; host gene, black; host gene product, white diamond.
Disrupted genes yield no product (X); genes downstream of MGE insertions may or may not be
transcribed (?). **B.** Each cycle of excision and integration may shift the MGE and its insertion
site and/or cargo between contexts, generating new genetic diversity. Intracellular movement
(solid arrows, teal cells) may interrupt (X) or influence gene regulation of (?) a new host gene
(diamonds); intrapopulation movement (dashed arrows, blue cells) lets the MGE interrupt or

influence the same or different host genes in a new genetic background; interpopulation
movement (dot-dashed arrows, yellow cells) lets the MGE interrupt or influence a new suite of
genes; and MGE loss (gray dotted arrows) may remove or leave behind cargo genes and
potentially restore function of previously disrupted host genes. Not depicted: capture of host
genes as new cargo genes.

**Figure 2. The Stordalen Mire model ecosystem and its mobile genetic element (MGE)
recombinases. A.** Long-term study site (image credit: NASA) with large datasets (gray inset)
along a palsa-bog-fen thaw gradient. **B.** MGE recombinase percentage in all genes from
Stordalen contigs (≥1,500 bp). **C.** Counts of most abundant genes: MGE recombinase; *rpoE*,

RNA polymerase sigma 70 factor; *drrA*, antibiotic transport system ATP binding protein; *pknB*,
serine/threonine protein kinase; *ABC.CD.P*, putative ABC transport system permease. **D.** MGE
recombinase type distributions: "IS_Tn", insertion sequence/transposon; "Phage", phage and
phage-like elements; "CE", conjugative elements/mobility islands; "Integron", integron;
"ambiguous", recombinases found in >1 MGE types. **E.** MGE estimates per genome for
Stordalen Mire contigs (median=36) or MAGs (median=23) and from cultured isolates in
Khedkar et al. (median=25; red dashed line). Filled circles, points >1.5 times the interquartile
range below the first quartile or above the third quartile. **F.** Comparison of MGE recombinase
recovery between short-read and hybrid assemblies (n = 17), with recombinase matching at
100% protein identity. "Short read missed": recombinase found only in hybrid MAGs; "short
read recovered": recombinase found in both hybrid and short-read MAGs (recovery rates shown
as percentages on top of bars).

**Figure 3. MGE recombinase type distribution across microbial host lineages (MAG-based)
of the full Stordalen Mire short read dataset. A.** Number of each MGE type found in each
phylum, as identified from contigs binned into MAGs. MGE recombinase counts shown for the
top 12 phyla (out of 46 total), ordered by average MGE recombinase number per genome
(indicated to the left of each bar with "+" indicating Archaea). **B.** Number of MGE recombinases
identified per genome grouped by MGE type and host phylum. **C.** Number of MGE
recombinases (y-axis) against estimated number of genomes per phylum (x-axis). Dotted line
represents the global phylum-averaged number of MGE recombinases per genome.

442
Figure 4. Host functions impacted by MGEs in Stordalen Mire. Distribution across habitats,
host phyla, and MGE types of selected nutrient cycling pathways (yellow highlights in cell,
groups 1, 4, 6, and 7), housekeeping functions (red highlight, group 5), and signaling and social
functions (blue highlights, groups 1–3) impacted by MGEs whose boundaries could be
conservatively established. Colored boxes indicate positive detection of an MGE-affected
function in a habitat, phylum, or MGE type. Phyla are ordered from left to right by increasing
per-genome MGE count (see fig. 3). CE, INT, and phage carry impacted functions as cargo
genes; IS_Tn interrupt host genes, causing loss of function.

451
452
453
454

455
Figure 5. MGE recombinase activity. A. Top panel: Genes detected most frequently as active
based on metatranscriptome read mapping. Gene categories from right to left: *cspA*, major cold
shock protein gene; MGE recombinase; *rpoE*, RNA polymerase sigma-70 factor gene; *hupB*,
DNA binding protein HU-beta gene; *glnA*, glutamine synthetase gene; RP, ribosomal protein
genes. Bottom panel: percentage of genes active based on the full set of genes identified in the
matching metagenome. Wilcoxon rank-sum test of statistical significance; ****, *P* value≤0.0001.
B. Three facets of MGE recombinase activity/mobility-- transcription (left), genomic
neighborhood stability (middle), partial carriage by host population (right) – are conceptually
presented (upper panel) and quantified and summarized per MGE recombinase type (lower
panel). The x-axis is the percentage of unique MGE recombinase (100% identity at protein level)
detected with activity / mobility within each type. **C**. Comparison of ratio of active MGE

recombinase, "IS_Tn" , and "Phage" recombinases across three habitats, normalized by the ratio
of active ribosomal genes for the same sample. **D.** Normalized active ratio of "IS_Tn" and
"Phage" recombinase across host phyla with ≥5 active "IS_Tn" or "Phage" recombinases, and
ribosomal genes. "CE" and "Integron" recombinases are not shown here due to low counts.
471
472

Supplementary Materials

Methods and Materials

Data collection, metagenome assembly and binning, gene annotation and taxonomy assignment

Our study site, Stordalen Mire, is a long-term ecological research site in northern Sweden. Large sampling and sequencing efforts since 2010 have yielded ~4.9 Tb of metagenomic sequence data from 615 readsets from 408 bulk soil samples that span space (across the site's three habitat types, n = 188 palsa, 200 bog, 227 fen; average ~8 Gb/sample) and time (summer peak growing season, 2010–2017) (**table S1**) with details described in another parallel study (*28*) and previously (*25, 26, 54*). The IsoGenie Consortium and EMERGE Biology Integration Institute have developed extensive resources from Stordalen Mire metagenome data, including assembly across all samples to produce 30,710,180 contigs (≥1500 bp) and 36,419 medium to high-quality metagenome-assembled genomes (MAGs; completeness ≥ 70% and contamination ≤10%, CheckM2 v1.0.2) (*55*) that clustered into 2,048 populations (95% Average Nucleotide Identity [ANI]) per community standards (*56*). Select samples have also been sequenced using metatranscriptomics to assess gene expression (50 spatio-temporally diverse samples; average ~18 Gb per sample). For this work, we additionally performed paired long- and short-read DNA sequencing on six samples, chosen to be representative of the larger dataset, as a control to assess short-read MGE recovery.

The protocol used for sample preparation and sequencing for short-read metagenomes has been described previously (*25, 26, 54*). Briefly, nucleic acids were extracted according to a published protocol (*57*), except for the MAG source samples from substrate addition incubations ("201607_SubstrateInc_9to19"), which were extracted with the DNeasy PowerSoil Kit (cat# 12988-100, Qiagen). Libraries were prepared with either KAPA Hyperprep, TruSeq, or Nextera XT kits for short-read metagenomes; SMRTbell Express Template Prep Kit 2.0 for long-read metagenomes; or TruSeq Stranded mRNA kit for metatranscriptomes. Reads were sequenced with either Illumina HiSeq (2000 or 2500), NovaSeq 6000, or NextSeq 500 for short-read metagenomes; PacBio Sequel IIe for long-read metagenomes; or NovaSeq 6000 for metatranscriptomes.

Short-read sequencing data from individual samples were trimmed by Trimmomatic (v.0.36) (*58*), assembled by SPAdes with --meta option (v.3.12) (*59*), and then binned by MetaBAT2 (*60*), GroopM2 (*61*), and MaxBin2 (*62*) (Cronin.v2), or an ensemble approach (*28*). Those sequenced at JGI were processed by JGI's standard assembly and binning pipeline. Briefly, reads were quality-controlled, filtered, and trimmed using the BBTools package (*63*), assembled with SPAdes with --meta option (*59*), and binned with MetaBAT2 (*60*). The bins produced from this effort were combined with 9,406 bins from Woodcroft 2018 (*25*) and 11,120 bins from stable isotope probing (SIP) experiments with field peat soil and labeled litter (sequenced with NovaSeq at JGI and assembled with MEGAHIT v1.1.3) (*64*), and then filtered

using 70% completeness and 10% contamination CheckM2 (v1.0.2) (*55*) cutoffs to form the
36,419 metagenome-assembled genomes (MAGs; consisting of 4,193 from Cronin.v1, 27,423
from Cronin.v2, 847 from JGI, 1,543 from Woodcroft 2018 and 2,413 from SIP). Next, bins
obtained from the same metagenome were de-duplicated using Galah (95% ANI, commit
e9e59a8) (*56*), and all bins across the entire dataset were further dereplicated at 95% ANI using
Galah (commit e9e59a8), producing 2,048 population level clusters. Further, the contigs
encoding a recombinase were annotated with DRAM (--min_contig_size 1500) (*65*) and
taxonomically assigned at phylum level with CAT (*66*) requiring a minimal contig length of 3kb.
The contigs in MAGs were assigned to the MAG taxon (obtained with GTDB-tk) (*67*), since
taxonomic assignments based on whole MAGs are typically more reliable than single-contig
assignments.
524
525   *MGE recombinase identification:*
526   Recombinase identification was based on a framework recently developed to robustly
identify and classify MGE recombinases from microbial isolate genomes (*8*). Here, we applied
the same recombinase HMMs profiles and suggested cut-offs to identify MGEs in metagenomic
assemblies. In brief, open reading frames were predicted from assembled contigs by prodigal
(v2.6.3) using the "-p meta" option (*68*), then recombinases were identified among the predicted
protein sequences by hmmsearch (v3.3.2) (*69*) using the 68 recombinase subfamily HMMs
(**table S2**) and the "--cut-ga" option. Identified recombinases were assigned to the respective
MGE recombinase type corresponding to the HMM with best score – as suggested previously
(*8*).
535
536   *Annotation-based inclusion/exclusion criteria:*
537   MGE recombinase candidates were further quality controlled as recommended previously
(*8*) by using eggnog-mapper (v2.1.9, default settings) (*70*). First, genes annotated as invertase,
replication protein, DNA repair, DNA binding, homologous recombination, viral exonucleases,
or reverse transcriptase (interpreted to be spurious hits to other host genes related to DNA
manipulation) were removed, except for the ones also annotated with the Pfam catalytic domains
of "transposase" or "helicase" that were retained after manual curation. This removed 0.68% of
the identified possible recombinases (**fig. S1**). Second, sequences annotated by eggnog-mapper
as CRISPR, defense response to virus, or maintenance of DNA repeat elements were removed to
filter out spurious hits thought to be similar to Cas family recombinases, but not considered as an
MGE recombinase. This removed 0.46% of the identified possible recombinases. Third, we
discarded predicted recombinases that contained only an HTH (DNA binding) domain, as well as
relaxase hits for which Pfam annotates a helix domain, but without the catalytic domain required
for recombinase activity. This removed 0.07% of the identified possible recombinases. Fourth,
recombinases matching a "Cell" type recombinase HMMs (and thus unlikely to be encoded by
an MGE) were removed. This removed 2.52% of the identified candidate recombinases, and
yielded a final total of 2,107,204 MGE recombinases (**table S3** item A). Lastly, some
recombinase subfamilies/HMMs include recombinases from different MGE types, which
prevents assignment for that HMM to a specific MGE type. These recombinases with ambiguous
origin accounted for 20.51% of the identified recombinases. They were *included* in the analyses
of total MGE recombinases, but *excluded* in analyses of individual MGE types, which focused
on four recombinase-based MGE types: "IS_Tn" (insertion sequence [IS] and transposon [Tn]),
"CE" (conjugative elements including conjugative element and mobility islands), "PhL" (phage

and phage-like elements) and "Integron" (integron) (**fig. S1**). To assess the diversity of these
these MGE recombinases, we considered recombinase sequences as "redundant" when they
clustered with other sequences at 100% AAI using CD-HIT (v4.8.1) (*71*) with "-c 1 -g 0 -G 1 -l
30 -d 0 -M 0". This yielded 1,130,776 clusters that we considered as unique recombinases.
563
Identification of cargo genes/interrupted genes associated with MGEs
Once MGE recombinases were identified, we surveyed their genomic context to assess
potential cargo and/or interrupted genes. This was done using MGE-specific approaches as
follows.
Integron: All integron-containing contigs were screened for cargo genes (also known as
"gene cassettes") using integron-finder (v2.0.2) (*72*) with default parameters. Briefly, integron-
finder identifies integrons by using integron specific integrase HMMs to identify integron
integrases (*intI*), and a covariance model to identify the integron attachment site (*attC*), as the
two bounding regions of an integron sequence. Only integrons encoding the *intI* and at least one
attC treated as "complete" by integron-finder were retained. There are two types of integron
structures that are known in terms of the orientation of *intI* and *attC*: 1) a regular type with *intI*
and *attC* on opposite strands and facing outwards, 2) another type with *intI* and *attC* on the same
strand, called inverted integrase integron (*72*). Of the 686 "complete" integrons we identified,
681 integrons also met these orientation expectations and the 5 that did not were not considered
further. The gene cassettes delimited by *intI* and *attCs* in these integrons were treated as cargo
genes associated with integrons (*73*), resulting in integron regions with various lengths
(minimum: 983bp, median: 2,310bp, mean: 3,092bp, and maximum: 14,195bp). Problematically,
most contigs (97.4% or 25,953 out of 26,634) encoding Khedkar's "Integron" type recombinase
were not considered complete by integron-finder. To test this, we also replaced the integron
integrase HMM in the integron-finder tool with that from Khedkar et al. (*8*) and found it
produced nearly identical results and did not yield additional instances of predicted complete
integrons. One explanation could be that *attC*, not integron integrase, is limitation, since the
flanking *attC* sequences are identified using a database of known *attC* sequences and these could
miss novel *attC* sequences used by integrons in natural microbiomes. Given this, we interpret the
large discrepancy between Khedkar's and integron-finder's integrons to represent one of the
following scenarios: (i) degraded or domesticated integrons that lack *attC* sites (*74*, *75*), (ii)
integrons that lacks *attC* sites due to fragmented assembly, or (iii) intact and likely complete
integrons that have novel *attC* sites. Thus our analyses will under-estimate the extent of cargo
genes carried by integrons in Stordalen Mire.
Conjugative Element (CE): CE are made of three key components – a relaxosome
(includes the CE-type recombinase), a conjugation couple protein, and a type IV secretion
system (T4SS) – that can be used as hallmark genes that are more conserved than other CE genes
(*76*). To identify cargo genes associated with CE, the CE hallmark genes were identified in CE
recombinase-encoding contigs by conjscan implemented in MacSyFinder (v2.0rc7) with "--
models CONJScan/Plasmids all --db-type gembase" and "--models CONJScan/Chromosome all
--db-type gembase" with gene protein sequences as input (*76*, *77*). Conjscan specifically
establishes 8 types of CEs using hallmark genes that are core across all 8 CE types, and
accessory genes that are diagnostic for specific CE types. CE identified by conjscan from
Stordalen Mire sequence data were those that matched a relaxase, a couple protein (an ATPase
that couples MGE with the T4SS), a T4SS ATPase, and an additional structural gene from T4SS.
For the purposes of identifying cargo genes, matches of one model, MOB, were removed since it

only requires the relaxase and thus does not identify the boundaries to establish a complete CE.
As a result, we identified 1,017 near complete CEs. As recommended, we required the distance
between the hallmark and accessory genes to be <=30 genes except for the relaxase which could
be as much as 60 genes away from any other hallmark and accessory genes (*76*). In a small
number (13 of 1,017) of instances conjscan identified >1 CE in a region. For these, we manually
separated the identified hallmark and accessory genes into subgroups according to gene
proximity and required each subgroup to include all hallmark genes (relaxase, conjugation
couple protein and T4SS ATPase) and at least one accessory T4SS gene. Once such "complete"
CEs were identified, the genes between the leftmost and rightmost hallmark and accessory genes
were assumed to be cargo genes carried by the CE (except for hallmark and accessory genes
themselves). This resulted in CE regions with various lengths (minimum: 7907, median: 16,290,
mean: 23,383, and maximum: 81,148 bp). Overall, 1.6% of the 62,660 CE type recombinase
genes identified by our analyses were considered near-complete CE by conjscan. Again, this was
not a function of improved HMMs as we tested replacing the relaxase HMMs in conjscan by
those from Khedkar et al. (*8*) and found it produced similar results. Instead we interpret this large
discrepancy to suggest that most CEs in these data are degraded, fragmented from poor
assembly, or can not be formally detected as complete because of limitation and bias in hallmark
gene databases.
Insertion sequence / Transposon: To identify "IS_Tn" recombinase interrupted genes, up
to 3,000 bp upstream and downstream of "IS_Tn" recombinases were collected from each
contig, requiring a minimal length of at least 50 bp on each end and at least 500 bp across the
two ends combined (1,063,800 IS_Tn recombinases in total meeting the requirements), and then
aligned by minimap2 (v.2.22, default settings) (*78*) to the coding sequence of representatives of
protein clusters (PCs) obtained by MMseqs2 at 95% identity (v4.8.1) (*79*) from all contigs
(≥1500 bp) used in this study. The corresponding coding sequence was considered interrupted if
the following four conditions were met: 1) upstream and downstream alignments combined
covered ≥50% of the coding sequence of the reference PC; 2) if there is a gap between upstream
and downstream alignments of the coding sequence, the gap is ≤10% of the coding sequence
length; 3) if there is an overlap between upstream and downstream alignments on the coding
sequence, the overlap is ≤10% of the shorter alignment; and 4) if there are multiple alignments in
either upstream or downstream to the coding sequence, the pair with largest coverage of coding
sequence is chosen. This identified 5,342 PCs putatively interrupted by a total of 12,332 IS_Tn
type recombinases.
Phage / phage-like elements: All contigs encoding "Phage" recombinase were screened
for temperate phage by VirSorter2 (v2.2.3) following VS2 SOP step 1 (v3.0) (*80*, *81*), and the
predicted phage sequences trimmed by checkV (default setting, version 0.7.0) (*82*). Next, the
checkV trimmed sequences encoding a "Phage" recombinase were considered as phage/phage-
like regions. To identify cargo genes, i.e. host(-like) genes encoded on phage/phage-like
elements, we opted for a conservative approach in which we reduced the dataset for "cargo gene
inspection" to include only phage/phage-like regions with one "Phage" recombinase, and use the
leftmost and rightmost checkV called "viral gene" or "Phage" recombinases as the boundaries.
This identified 4,277 temperate phages with "Phage" recombinases and at least one extra gene
other than two at ends. Meanwhile, 347 phage/phage-like elements were removed for having
more than one "Phage" recombinase, in order to avoid cases where multiple contiguous
temperate phages or degraded ones yielded a single prophage prediction, which could lead to
host genes being mistaken as cargo genes. This resulted in 3930 conserved phage/phage-like

regions (minimum: 1386bp, median: 7293bp, mean: 13,779bp, and maximum: 352,302bp).
Annotations of cargo/interrupted genes: Insertion sequence or transposon interrupted
genes and genes within other MGEs were functionally annotated using KEGG and Pfam
databases in DRAMv for phages/phage-like elements or DRAM for others MGEs (v1.2.3, --
min_contig_size 1500) (*65*), and further evaluated by comparison with best hits in NCBI NR
(downloaded on 10/04/2020) using DIAMOND (v2.0.8.146, blastp bit-score ≥50) (*83*). A subset
of the genes of specific interest to the Stordalen Mire thawing permafrost system were selected
and manually curated for display (**table S4**) including Transport, Carbon/Nitrogen/Sulfur
cycling, Signaling, Antiviral defense, and Genome maintenance and expression.
660
Evaluation of short read assembly and binning artifacts for recovering MGE recombinases
using long reads.
Assembly and binning of paired long reads and short reads: Soil samples (6 total) were
collected at the Stordalen Mire site from 2019 from two depths (cm below ground) across three
habitats (Palsa, Bog, and Fen). DNA extraction and Illumina NovaSeq sequencing was
performed as described above. In addition, DNA sequencing was performed from the same DNA
extractions using PacBio Sequel IIe HiFi sequencing platform at JGI. PacBio data was processed
at JGI to form filtered CCS (Circular Consensus Sequencing) reads. Assemblies were generated
through two methods: short-only and hybrid short-read plus long-read (herein: hybrid). Short-
only was assembled with metaSPAdes v3.15.4 (*59*) using Aviary v0.5.3 (*84*) with default
parameters. Hybrid assembly was performed using Aviary v0.5.3 with default parameters. This
involved a step-down procedure with long-read assembly through metaFlye v2.9-b1768 (*85*),
followed by short-read polishing by Racon v1.4.3 (*86*), Pilon v1.24 (*87*) and then Racon again.
Next, reads that didn't map to high-quality metaFlye contigs were hybrid assembled with
metaSPAdes v3.15.4 and binned out with MetaBAT2 v2.1.5 (*60*). For each bin, the reads within
the bin were hybrid assembled using Unicycler v0.4.8 (*88*). The high-coverage metaFlye contigs
and Unicycler contigs were then combined to form the assembly fasta file. Genome recovery was
performed using Aviary v0.5.3 with samples chosen for differential abundance binning by Bin
Chicken v0.4.2 (*89*) using SingleM metapackage S3.0.5 (*90*). This involved initial read mapping
through CoverM v0.6.1 (*91*) using minimap2 v2.18 (*78*) and binning by MetaBAT v2.1.5 (*92*),
MetaBAT2 v2.1.5 (*60*), VAMB v3.0.2 (*93*), SemiBin v1.3.1 (*94*), Rosella v0.4.2 (*95*),
CONCOCT v1.1.0 (*96*) and MaxBin2 v2.2.7 (*62*). Genomes were analyzed using CheckM2
v1.0.2 (*55*) and clustered at 95% ANI using Galah v0.4.0 (*56*).
Assessment of MGE recombinase recovery by short-read assembly and binning: We
considered the high quality hybrid MAGs as the best available references and used these to
assess the recovery of MGE recombinases in short-read contigs and MAGs. A total of 17 high-
quality MAGs (completeness ≥95%, contaminant ≤5% and at most 100 contigs) from hybrid
assembly were selected and paired with 17 MAGs recovered from short-read assembly through
the 95% ANI clusters (the highest ANI were chosen if there were many) for comparison. The
short-read MAGs included 5 medium quality (≥70% completeness, ≤10% contamination) and 12
high quality (≥95% completeness, ≤5% contamination).
Evaluation of these 17 MAGs were carried out as follows. First, each pair of hybrid
MAG and short-read MAG were compared in terms of total number of MGE recombinases, as
well as unique and shared MGE recombinases at 100% AAI by CD-HIT (v4.8.1) with "-c 1 -g 0
-G 1 -l 30 -d 0 -M 0" (*71*). Second, we evaluated the impact of sequencing depth, assembly, and
binning to assess which step was most problematically leading to under-detection of MGE

recombinase identification in short reads. Initially, we mapped short-read assembled contigs
(sheared into 500 bp fragments) to hybrid MAGs from the same sample using coverM v0.6.1
requiring each read to be 75% aligned with 95% identity (*91*). Third, these data were used to
assign MGE recombinases in the 17 hybrid MAGs into one of three categories: (i) those
recovered by short-read MAGs (termed "covered"), (ii) those not fully covered by short-read
contigs, ie, missed by short-read assembly (termed "missed assembly"), and (iii) those fully
covered by short-read contigs, ie, recovered by short read assembly but missed by short-read
binning (termed "missed binning"). Fourth, the coverage of each gene was generated by dirseq
v0.4.3 (*25*) using the BAM file from above and used to assess the relative abundances of each
category as a metric for the impact on MGE-representation in the short-read assemblies.
With a better understanding for which step was failing most often for MGE recombinase
recovery in short-read assemblies, we sought next to evaluate what factor(s) might be causing the
failed capture events. First, we evaluated sequencing depth since low coverage might cause
breakages in short-read assembly graphs. This was done by mapping trimmed short reads to
hybrid MAGs using coverM v0.6.1 requiring each read 75% aligned with 95% identity,
calculating the read coverage of each gene by dirseq v0.4.3, and comparing coverages among the
three categories of MGE recombinases. Second, we compared the genomic neighborhood
number among the three categories, reasoning that perhaps indels were an issue. Genomic
neighborhood number was calculated by clustering genomic neighborhoods of each unique MGE
recombinase (see below) at 90% ANI. To test this, nucleotide diversity for each base position
was calculated as $1 - [$(number of "A" bases / total bases) $^2 + $ (number of "C" bases / total
bases) $^2 + $ (number of "T" bases / total bases) $^2 + $ (number of "G" bases / total bases) $^2 + $
(number of "Deletelion" bases / total bases) $^2 + $ (number of "Insertion" bases / total bases) $^2]$,
adopting the nucleotide diversity definition from inStrain (*53*) with addition of deletion and
insertion variants. The variant count at the base position was generated by pysamstats (v1.1.2,
https://github.com/alimanfoo/pysamstats) with "--type=variation" option. Third, we assessed
nucleotide diversity of each MGE recombinase, reasoning that perhaps elevated nucleotide
diversity could lead to short-read assembly graph breakages. To test this, for each of the 17
MAG pairs, MGE recombinases only identified in hybrid MAG were compared to short-read
assembled contigs to track whether they were assembled by short reads and then whether they
were binned in MAGs from other MAG clusters. The tracking of MGE recombinases in contigs
was accomplished with "MMseqs map" (v13.45111) with default setting and then filtered for hits
with 100% identity in protein sequence (*79*).

MGE recombinase per genome estimation from contigs and MAGs
To estimate MGE recombinase number per genome from contigs, the median count of
ribosomal genes was first used to estimate the number of genomes in each assembly. To do this,
ribosomal genes were extracted from contig annotations by DRAM (*65*) using KEGG IDs (**table
S6**), and the average number of recombinase genes per genome was calculated by dividing the
total number of recombinases in the sample by the estimated total number of genomes. We next
linked individual recombinase-encoding contigs to MAGs built from the same sample. To that
end, we used exact string searching to identify MGE-recombinase-encoding contigs in the de-
replicated MAGs from the same sample. Through this process, 106,017 out of a total 665,700
(15.9%) MGE-recombinases-encoding contigs ≥3kbp could be linked to a MAG. To account for
partial MAG recovery, the number of MGE recombinases per genome estimated from MAGs
was calculated as the number of MGE recombinases in a given MAG divided by MAG

completeness.
744
Genomic neighborhoods of MGE recombinases and mobility evaluation

MGE mobility was assessed by comparing the genomic neighborhoods of individual
recombinases. For this analysis, we limited the input data to recombinases that occurred ≥2 times
in the total dataset (assembled contigs per sample). This was achieved by clustering the
recombinases at 100% AAI using CD-HIT v4.8.1 with "-c 1 -g 0 -G 1 -l 30 -d 0 -M 0" (*71*), only
keeping the clusters with at least two recombinases. For each MGE recombinase, 300 bp
upstream and downstream of the recombinase were considered as its immediate genomic
neighborhoods, excluding those near the contig ends (< 300bp). For each recombinase cluster,
pairwise comparison of neighborhoods was performed among all member recombinases using
both neighborhoods (ie, upstream vs. upstream, and downstream vs. downstream), using
BLAST+ (v2.8.1+) with "-task blastn" (*97*) and bitscore ≥50. Average nucleotide identity (ANI)
for each neighborhood comparison was calculated as the "sequence identity from blastn output"
multiplied by the "alignment coverages". For each comparison, we labeled the neighborhood
with the higher ANI value as "high_ani" and the neighborhood with lower ANI as "low_ani".
For some MGEs, we expect the neighborhood on the MGE side of the recombinase to be part of
the MGE and thus remain identical after integration into a new locus, so we used "low_ani"
values (i.e. the lower ANI between the upstream and downstream region of the pair) to evaluate
whether the neighborhood of an MGE changed or not. An MGE recombinase pair was
considered to be in different genomic locations if "low_ani" was <1%, and a recombinase cluster
was treated as active if one pairwise comparisons showed different neighborhoods. This
approach only used immediate neighborhood (within 300bp) since a requirement of
neighborhood with longer range would remove more recombinases from this analysis due to the
short average contig length. However, as a sensitivity analysis, we also tested the above with 600
bp and 1000 bp neighborhoods. These analyses showed that the size of the genomic
neighborhood (300 - 1000bp) had very little impact on the fraction of MGE recombinases
identified as mobile via this changing genomic neighborhood metric (**fig. S10**).
771
MGE recombinase activity by metatranscriptome

MGE activity was assessed via documenting which MGE recombinases were detectably
transcribed in metatranscriptomes. Leveraging 81 paired metatranscriptomes and metagenomes,
we evaluated MGE recombinase transcriptional activity by mapping metatranscriptome reads to
contigs assembled from metagenomes using transcriptM v0.3.1 (*98*) with ORF predictions from
DRAM (see above) as the "--gff" input, which also removed tRNA, rRNA, tmRNA, and PhiX
from the reads. To ensure the quality of RNAseq data (reverse stranded), we only analyzed 58
pairs with ≥80% of reads mapped to the coding reverse strand of ORFs in contigs, based on
dirseq v0.4.3 (*25*) with BAM and GFF files from above as inputs (**table S5**). We only considered
contigs ≥10 kbp to avoid spurious ORF predictions from short contigs. Next we defined genes
with ≥90% of gene length covered as expressed using pydirseq v0.1.0-alpha with BAM and GFF
files from above as inputs (*99*). To normalize for the differences in sequencing depth and overall
activity of the microbial community across samples, we used the average percentage of
ribosomal genes (**table S6**) detected as active as the baseline activity to normalize other genes
(termed "normalized active ratio"), since ribosomal genes are expected to be constitutively
expressed in active cells. This "normalized active ratio" thus provides an estimation of the
number of genes detected as expressed within a functional category (e.g. recombinases) relative

to an estimated total number of genomes detected as transcriptionally active in the same sample.
790
MGE recombinase partial carriage by host population
As a result of MGE movement within the host population and an equilibrium of the co-
evolution of MGEs and their hosts (*100*), we noticed that some MGE recombinases were present
in only a fraction of the contigs assigned to a given host population (termed "partial carriage").
To quantitatively evaluate this, trimmed metagenomic reads were mapped to contigs from the
same sample using coverM v0.6.1 with default setting (*91*), then the resulting BAM file was
filtered requiring ≥90% each read aligned with ≥95% identity, and the filtered BAM file was
used as input to "samtools depth" (v1.17) with "-a" option to get the read depth coverage at each
position (*101*). The first and last 150bp of each contig were not considered since read depth
coverage near the ends are expected to be artificially lower. We defined partial carriage as cases
where the coverage of the MGE recombinase was at least one standard deviation lower than the
larger one (requiring a minimal coverage of 1x) of the average coverages of the upstream and
downstream contig regions (Cohen's d ≥1). To ensure enough host regions were considered for
reliable mapping-based estimation of genome coverage, the MGE recombinase total set was
filtered to 23,647 MGE recombinases with ≥20 kb upstream and downstream. Although, again
sensitivity analyses were conducted, and similar patterns were observed when analyzing 2.5 kbp
instead of 20 kbp regions (**fig. S11**), which suggests that this conservative cutoff did not
substantially influence the detection of partial carriage in these data.
We expect that this method should give us a lower-bound estimate of partial carriage,
since it is not able to detect cases for which (i) the hosts have low read coverage (rare), (ii) the
carriage ratio is lower than but close to 1, (iii) host genome coverage has a wide range of
variation in different genomic locations due to strain variation within the population, or (iv) the
MGE recombinase is on the edge of a contig.
814
Statistics
All statistics and visualization were performed in R v4.2.1 (*102*). The R packages and
versions used can be found in https://github.com/emerge-bii/mobilome-submit-202407. For
statistical significance between groups, Wilcoxon rank sum test was used for two groups and
Kruskal Wallis test was used for >2 groups as implemented in "compare_means" in ggpubr
package v0.6.0, with *P values* adjusted by Benjamini-Hochberg adjustment for multiple
comparisons.
822
823 # Supplementary Text
824
Short reads under-detect active MGE recombinases, especially insertion sequences.
As a technical control for sequencing artifacts, we sought to investigate the impact of short
read assembly on MGE recombinase recovery from metagenomes as compared to that recovered
from short- plus long-read hybrid assemblies. To evaluate this, we leveraged paired short-read
and high-fidelity long-read sequencing data from 6 permafrost thaw soil samples collected in
2019 from Stordalen Mire. In this analysis, we sought to use high quality hybrid MAGs as the
best available references to understand the impact of short read assembly and binning on MGE
recombinase recovery.
We identified 17 high-quality hybrid-assembled MAGs (≥95% completeness, ≤5%
contamination, <100 contigs) that also had representation in the short-read assemblies (clustered

at 95% ANI). As expected, we found fewer MGE recombinases in short-read MAGs than in
hybrid-assembled MAGs (n = 312 versus 937, respectively, across all MGE types). While MGE
recombinase type recovery varied, IS_Tn type was consistently the least recovered (**fig. S6B**).
When directly comparing the recombinases detected in each pair of short-read and hybrid-
assembled MAGs (100% protein identity clustering), the short-read MAGs recovered 57% CE,
35% IS_Tn, and 60% Phage recombinases in the 17 selected hybrid MAGs (**Fig. 2F**) (Integron
recombinases were not included because of their low number, 27 in total in the shared MAGs),
for an overall average of 36% (17–92% across individual MAGs, all MGE types combined) per
hybrid MAG (**fig. S6C**). Therefore, MGE recombinase recovery rate varies and is highly
dependent on host taxon, but IS_Tn recombinases are the main type of recombinases under-
detected in short-read datasets.
We next investigated why some MGE recombinases were recovered by hybrid-assembled
MAGs, but missed by short-read MAGs. This revealed that virtually all hybrid-MAG MGE
recombinases were found in short reads, but not short-read assembled contigs, suggesting that
short read sequencing depth was not an issue, but assembly challenges prevented hybrid-
assembled MGE recombinases from being captured in the short-read MAGs (**fig. S7A&B**). We
posited that assembly could fail either due to hypervariability of the recombinase locus itself
and/or the surrounding genomic context. We found that recombinases missed by short read
assembly ("missed assembly") have significantly higher numbers of genomic neighborhoods (p
≤0.0001) and nucleotide diversity (p ≤0.001, Wilcoxon rank-sum test) compared to those
assembled and/or binned (**fig. S7C&D**), which we interpret as confirming that hypervariable
regions "break" short read assemblies due to mixed variation at that locus. For MGE
recombinases successfully assembled from short reads, we found a relatively low number of
inconsistent binning (<5%, **fig.S7E**). This is consistent with previous studies (*14*, *15*) showing
short read assembly and, to a lesser extent, binning under-detects MGE recombinases generally
with simulated data. With the majority of the metagenomic data generated with short read
sequencing, this means that the estimation of MGE prevalence and activity/mobility should be
seen as a lower bound.
863
Conservative MGE detection and impacted gene curation.
To understand the capacity of the mobilome to affect microbial function across Stordalen,
we sought to identify the genes whose distribution is influenced by mobilome activity, a group
defined by both the cargo genes carried by an MGE and the host gene context into which the
MGE is inserted. Because calling the precise boundaries of MGEs remains particularly
challenging in metagenomic assemblies, we opted for a conservative approach, treating the
outermost MGE hallmark genes as element boundaries and the genes between these bounds as
cargo (Methods). Although this approach undercounts cargo genes, it can be seen as a lower
bound. In our analyses – across integrons, CEs, and PhLs – we identified 90,219 cargo genes in
all, annotated with 4,322 different KEGG functions (**table S3** item C,D,E,F). We took a similarly
conservative approach to counting IS_Tn-impacted host genes, counting only those examples
where the genomic neighborhood immediately upstream and downstream of IS/transposons
could be robustly identified as the N- and C-termini of the same gene. Again, this approach will
undercount IS_Tn-impacted host genes, as for instance insertion just upstream of a gene may
influence or even block transcriptional regulation of that gene. Overall, we identified 12,332
interrupted host genes, annotated with 866 different KEGG functions. Then we selected and
manually curated a subset (1,220) of the KEGG functions of specific interest to the Stordalen

Mire thawing permafrost system for display based on functional categories, specifically
Transport, Carbon/Nitrogen/Sulfur cycling, Signaling, Antiviral defense, Genome maintenance
and expression, and prioritizing functions for which MGE impact was detected in multiple
samples rather than single observations (**Fig.4**, **table S4**).

Supplemental Figures

Figure S1. Flowchart of MGE recombinase and MGE boundary identification. A total of
~2.2M recombinases were identified using 68 HMMs, filtered in four steps, and then the
resultant ~2.1M MGE recombinases were classified into 'types' according to their HMM
affiliation using criteria established previously (*17*). These identification and typing steps are
collectively denoted by the background gray shading. The numbers and percentages of
recombinases involved in each step are shown in each box. Within each MGE recombinase type,
below the gray shaded area, various strategies to conservatively identify boundaries were used
(see tools listed in figure) largely by using hallmark genes annotated by these tools as
conservative boundaries (see Methods). In the remainder of the study, the ~2.1M MGE
recombinases identified were used for host lineage and activity/mobility inferences, whereas
only the boundary-constrained used for MGE impacted metabolism analyses.

902
Figure S2.recombinase_overview. Overview of MGE recombinase distribution from
Stordalen Mire. A. Distribution of contig-identified MGE recombinases by MGE type, as used
in the main text, as well as MGE recombinase family and subfamily (see a full list described in
table S2). **B.** Unique MGE recombinase (clustered by 100% AAI) by habitat. The numbers at the
top of bars are the number of unique MGE recombinases. **C.** Total MGE recombinase
distribution by habitat and year. The numbers at the top of bars are the number of total MGE
recombinases.

Figure S3.recombinase_by_subfamily. MGE recombinase distribution based on subfamily.
A. MGE recombinase subfamily distribution by habitat. The numbers at the top of bars are the number of total MGE recombinases. B. MGE recombinase subfamily distribution by habitat and year.

Figure S4.host_lineage.contig. MGE host lineage based on MGE recombinase identified from contigs. MGE recombinase encoding contigs (>=3kb) were evaluated for taxonomic

patterns as a complement to MAG-based analyses in Figure 3. **A.** Number of each MGE type
found in each phyla where the top 15 phyla (out of 89 in total) accounting for 84.4% of total
MGE recombinases are shown, ordered by MGE recombinase count per genome (the number to
the left of each bar).The genome number is estimated from ribosomal genes (see Methods).
Phyla names are colored per Figure 3. **B.** The number of MGE recombinases identified per
genome grouped by MGE type and host phyla. **C.** Phylum distribution of the overall microbial
community (estimated from ribosomal gene encoding contigs) and MGE recombinase types. **D**.
MGE recombinase number per genome per phylum. The dashed red lines are the average MGE
recombinase number per genome across all phyla, and the blue dashed lines are the average
across the phyla shown in the plot.
951
952

953
Fig.S5.host_lineage.contig_vs_mag. MGE recombinase and genome number distribution across phyla. A. MGE recombinase number per genome per phylum from MAGs. The dashed red lines are the average MGE recombinase number per genome across all phyla, and the blue dashed lines are the average across the phyla shown in the plot. **B.** Number of MGE

recombinases against number of genomes per phyla using contigs. Dotted line represents the
global average number of MGE recombinases per genome. The genome number is estimated
from ribosomal genes.The right panel is a zoomin of the area with cluttered dots in the left panel.
C. Number of MGE recombinases against number of genomes per phyla using MAGs. The right
panel is a zoomin of the area with clustered dots in the left panel.

1004
Figure S6.mag.short_vs_hybrid.biology A six sample assessment of short-read assembly
and binning artifacts for recovering MGE recombinases. A. Because short-read assemblies
can struggle with hypervariable and/or repeat regions of genomes, six metagenomic samples
were re-sequenced using high-fidelity long-read sequencing to develop a set of high-quality
MAGs (>= 95% completeness, <=5% contamination, and <100 contigs) that could be compared
for recombinase signal to short-read assembled MAGs from the same species cluster (>=95%
ANI). A total of 17 such pairs resulted and were compared. **B.** Recombinase-centric patterns in
recovery, where each circle represents the total recombinase number identified in the short read
versus matching hybrid MAG and dashed lines are the 1:1 lines. The inset in CE is zoomin of
clustered dots at lower left. **C.** Recombinase-centric patterns in recovery, where MGE
recombinases are considered either recovered by short read or missed (i.e., if they were only
found in hybrid MAGs) based upon searching via 100% identical protein sequence matches. **D.**
Lineage-centric patterns in MGE recombinase recovery. MAG name color matches color in
panel A. The numbers to the right of the bars are the percentages of total unique MGE
recombinases recovered by short read MAGs relative to the total.
1020
1021

1023
1024
Figure S7.mag.short_vs_hybrid.techinical Technical assessment for why short-read MAGs miss some MGE recombinases. In all panels, "covered" represents MGE recombinases assembled and binned both in short-read and hybrid MAGs, "missed assembly" represents those in hybrid MAGs missing in short-read MAGs and short-read contigs, "missed binning" represents those in hybrid MAGs missing in short-read MAGs, but captured in short-read contigs (statistical significance per Fig.1). **A.** Short reads mapped to the full hybrid MAGs dataset of MGE recombinases evaluated by sequencing depth (left panel, vertical coverage) and breadth (right panel, horizontal coverage). The red dashed line at y=10 (left panel) approximates the

minimal depth coverage needed for genome assembly considering the normal distribution of per
bp position of depth coverages. **B.** Short-read-contig-based depth (left panel) and breadth (right
panel) coverage per MGE recombinase to assess whether MGE recombinases were missed due to
assembly or binning. **C.** To evaluate whether highly variable MGE-recombinase-caused genomic
neighborhood context could prevent assembly, we quantified such variation as genomic context
number via a clustering approach (see Methods). **D.** Nucleotide variation (pi, see Methods)
across three MGE recombinase 'capture scenarios'. **E.** Logic workflow to establish the fraction
of short- (left panel) or hybrid- (right panel) MAG-only recombinases that are mis-binned in
other MAGs. Each step is a yes or no answered question, and numbers represent MGE
recombinase counts in our dataset.
1043

1044
1045
Figure S8.metat_activity. MGE recombinase activity accessed by metatranscriptomics. This
figure complements Fig.5. **A.** Normalized active ratio among IS_Tn and Phage type recombinase
types across host phyla. The rest of panels are MGE recombinase metatranscriptomics activity by

active OTU (clustered at 100% identity in protein sequence) percentage. Each OTU with >=1
active MGE recombinase is considered as active. **B.** within different years from 2011 to 2017; **C.**
within each habitat; **D.** within host phyla (only those >= 10 OTUs are shown; the X axis is
ordered by the average of active OTU percentage across MGE recombinase types).

1061
1062 **Fig.S9.genomic_nh.example Example of unstable genomic neighborhoods of MGE**
1063 **recombinases.** The labels at the left of genome maps are habitat, year, and contig length
1064 (separated by "|"), followed by host phyla. **A**. IS_Tn type (transposase); **B**. Phage type (int, short
1065 for integrase); **C**. CE type (relaxase), and **D**. Integron type (intI) recombinase.

1066
1067
1068
1069

1070
Figure S10.genomic_nh.by_recombinase_otu. MGE recombinase genomic neighborhood
stability. MGE activity could lead to their genomic neighborhood instability. To measure MGE
recombinase genomic neighborhood stability, MGE recombinases are clustered at 100% AAI,
and neighborhoods of 300 bp up/down-stream of recombinase are extracted. If different
neighborhoods (<1% ANI) are detected within a cluster, the recombinase cluster is considered as
unstable. The proportion of unstable MGE recombinase OTU were summarized: **A.** within each
MGE type across neighborhood length cutoff ranging from 300 - 1000bp; **B.** within each year;
C. within each habitat; and **D.** within host phyla (only those w/ >=10 OTUs are shown; the X
axis is ordered by the average across MGE recombinase types). "CE" and "Integron" are not
shown here due to lack of data.

Figure S11.partial_carriage MGE recombinase partial carriage by microbial populations.
MGE activity could lead to partial carriage of MGE in microbial populations. Here MGE
recombinase partial carriage is defined as significantly lower coverage of an MGE recombinase
than of its contig (only large contigs >=20kb upstream and downstream of a MGE recombinase
are used to reduced the spurious coverage estimates from short contigs), and further MGE
recombinase OTUs (100% AAI) with >= 1 partially carried MGE recombinase are considered as

partially carried OTUs. The percentage of partially carried MGE recombinase OTUs were measured: **A.** within each MGE type across neighborhood length cutoff ranging from 2500 - 20,000bp; **B.** within each year; **C.** within each habitat; and **D.** within host phyla (only those w/ >= 10 OTUs are shown; the X axis is ordered by the average across MGE recombinase types).

References

1. T. W. Crowther, J. van den Hoogen, J. Wan, M. A. Mayes, A. D. Keiser, L. Mo, C. Averill, D. S. Maynard, The global soil community and its influence on biogeochemistry. *Science*, doi: 10.1126/science.aav0550 (2019).

2. L. Paoli, H.-J. Ruscheweyh, C. C. Forneris, F. Hubrich, S. Kautsar, A. Bhushan, A. Lotti, Q. Clayssen, G. Salazar, A. Milanese, C. I. Carlström, C. Papadopoulou, D. Gehrig, M. Karasikov, H. Mustafa, M. Larralde, L. M. Carroll, P. Sánchez, A. A. Zayed, D. R. Cronin, S. G. Acinas, P. Bork, C. Bowler, T. O. Delmont, J. M. Gasol, A. D. Gossert, A. Kahles, M. B. Sullivan, P. Wincker, G. Zeller, S. L. Robinson, J. Piel, S. Sunagawa, Biosynthetic potential of the global ocean microbiome. *Nature* **607**, 111–118 (2022).

3. S. Nayfach, Z. J. Shi, R. Seshadri, K. S. Pollard, N. C. Kyrpides, New insights from uncultivated genomes of the global human gut microbiome. *Nature* **568**, 505–510 (2019).

4. P. J. Turnbaugh, M. Hamady, T. Yatsunenko, B. L. Cantarel, A. Duncan, R. E. Ley, M. L. Sogin, W. J. Jones, B. A. Roe, J. P. Affourtit, M. Egholm, B. Henrissat, A. C. Heath, R. Knight, J. I. Gordon, A core gut microbiome in obese and lean twins. *Nature* **457**, 480–484 (2009).

5. M. Haudiquet, J. M. de Sousa, M. Touchon, E. P. C. Rocha, Selfish, promiscuous and sometimes useful: how mobile genetic elements drive horizontal gene transfer in microbial populations. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **377**, 20210234 (2022).

6. M. G. Durrant, M. M. Li, B. A. Siranosian, S. B. Montgomery, A. S. Bhatt, A Bioinformatic Analysis of Integrative Mobile Genetic Elements Highlights Their Role in Bacterial Adaptation. *Cell Host Microbe* **27**, 140-153.e9 (2020).

7. S. R. Partridge, S. M. Kwong, N. Firth, S. O. Jensen, Mobile Genetic Elements Associated with Antimicrobial Resistance. *Clin. Microbiol. Rev.* **31**, e00088-17 (2018).

8. S. Khedkar, G. Smyshlyaev, I. Letunic, O. M. Maistrenko, L. P. Coelho, A. Orakov, S. K. Forslund, F. Hildebrand, M. Luetge, T. S. B. Schmidt, O. Barabas, P. Bork, Landscape of mobile genetic elements and their antibiotic resistance cargo in prokaryotic genomes. *Nucleic Acids Res.* **50**, 3155 (2022).

9. J. Zhang, Y. Xu, M. Wang, X. Li, Z. Liu, D. Kuang, Z. Deng, H.-Y. Ou, J. Qu, Mobilizable plasmids drive the spread of antimicrobial resistance genes and virulence genes in Klebsiella pneumoniae. *Genome Med.* **15**, 106 (2023).

10. D. J. Rankin, E. P. C. Rocha, S. P. Brown, What traits are carried on mobile genetic elements, and why? *Heredity* **106**, 1–10 (2011).

11. A. Dragoš, A. J. C. Andersen, C. N. Lozano-Andrade, P. J. Kempen, Á. T. Kovács, M. L. Strube,
Phages carry interbacterial weapons encoded by biosynthetic gene clusters. *Curr. Biol. CB* **31**,
3479-3489.e5 (2021).

12. E. P. C. Rocha, D. Bikard, Microbial defenses against mobile genetic elements and viruses: Who
defends whom from what? *PLoS Biol.* **20**, e3001514 (2022).

13. S. Benler, G. Faure, H. Altae-Tran, S. Shmakov, F. Zhang, E. Koonin, Cargo Genes of Tn7-Like
Transposons Comprise an Enormous Diversity of Defense Systems, Mobile Genetic Elements, and
Antibiotic Resistance Genes. *mBio* **12**, e02938-21.

14. J. J. Kerkvliet, A. Bossers, J. G. Kers, R. Meneses, R. Willems, A. C. Schürch, Metagenomic
assembly is the main bottleneck in the identification of mobile genetic elements. *PeerJ* **12**, e16695
(2024).

15. F. Maguire, B. Jia, K. L. Gray, W. Y. V. Lau, R. G. Beiko, F. S. L. Brinkman, Metagenome-
assembled genome binning methods with short reads disproportionately fail for plasmids and
genomic Islands. *Microb. Genomics* **6**, mgen000436 (2020).

16. I. L. Brito, S. Yilmaz, K. Huang, L. Xu, S. D. Jupiter, A. P. Jenkins, W. Naisilisili, M. Tamminen,
C. S. Smillie, J. R. Wortman, B. W. Birren, R. J. Xavier, P. C. Blainey, A. K. Singh, D. Gevers, E.
J. Alm, Mobile genes in the human microbiome are structured from global to individual scales.
Nature **535**, 435–439 (2016).

17. S. Roux, J. R. Brum, B. E. Dutilh, S. Sunagawa, M. B. Duhaime, A. Loy, B. T. Poulos, N.
Solonenko, E. Lara, J. Poulain, S. Pesant, S. Kandels-Lewis, C. Dimier, M. Picheral, S. Searson, C.
Cruaud, A. Alberti, C. M. Duarte, J. M. Gasol, D. Vaqué, Tara Oceans Coordinators, P. Bork, S. G.
Acinas, P. Wincker, M. B. Sullivan, Ecogenomics and potential biogeochemical impacts of globally
abundant ocean viruses. *Nature* **537**, 689–693 (2016).

18. J. Tamayo-Leiva, J. Alcorta, F. Sepúlveda, S. Fuentes-Alburquenque, J. I. Arroyo, J. E. González-
Pastor, B. Díez, Structure and dispersion of the conjugative mobilome in surface ocean
bacterioplankton.

19. Y. Yi, S. Liu, Y. Hao, Q. Sun, X. Lei, Y. Wang, J. Wang, M. Zhang, S. Tang, Q. Tang, Y. Zhang,
X. Liu, Y. Wang, X. Xiao, H. Jian, A systematic analysis of marine lysogens and proviruses. *Nat.
Commun.* **14**, 1–15 (2023).

20. P. Prentki, B. Teter, M. Chandler, D. J. Galas, Functional promoters created by the insertion of
transposable element IS1. *J. Mol. Biol.* **191**, 383–393 (1986).

21. A. Lipszyc, M. Szuplewska, D. Bartosik, How Do Transposable Elements Activate Expression of
Transcriptionally Silent Antibiotic Resistance Genes? *Int. J. Mol. Sci.* **23**, 8063 (2022).

22. A. J. Weisberg, J. H. Chang, Mobile Genetic Element Flexibility as an Underlying Principle to
Bacterial Evolution. *Annu. Rev. Microbiol.* **77**, 603–624 (2023).

23. S. S. Finks, J. B. H. Martiny, Plasmid-Encoded Traits Vary across Environments. *mBio* **14**,
e0319122 (2023).

24. C. K. McCalley, B. J. Woodcroft, S. B. Hodgkins, R. A. Wehr, E.-H. Kim, R. Mondav, P. M. Crill,

J. P. Chanton, V. I. Rich, G. W. Tyson, S. R. Saleska, Methane dynamics regulated by microbial
community response to permafrost thaw. *Nature* **514**, 478–481 (2014).

25. B. J. Woodcroft, C. M. Singleton, J. A. Boyd, P. N. Evans, J. B. Emerson, A. A. F. Zayed, R. D.
Hoelzle, T. O. Lamberton, C. K. McCalley, S. B. Hodgkins, R. M. Wilson, S. O. Purvine, C. D.
Nicora, C. Li, S. Frolking, J. P. Chanton, P. M. Crill, S. R. Saleska, V. I. Rich, G. W. Tyson,
Genome-centric view of carbon processing in thawing permafrost. *Nature* **560**, 49–54 (2018).

26. J. B. Emerson, S. Roux, J. R. Brum, B. Bolduc, B. J. Woodcroft, H. B. Jang, C. M. Singleton, L. M.
Solden, A. E. Naas, J. A. Boyd, S. B. Hodgkins, R. M. Wilson, G. Trubl, C. Li, S. Frolking, P. B.
Pope, K. C. Wrighton, P. M. Crill, J. P. Chanton, S. R. Saleska, G. W. Tyson, V. I. Rich, M. B.
Sullivan, Host-linked soil viral ecology along a permafrost thaw gradient. *Nat. Microbiol.* **3**, 870–
880 (2018).

27. A. A. Pratama, G. Domínguez-Huerta, J. M. Wainaina, A. A. Zayed, B. Bolduc, F. Tian, J.
Ellenbogen, J. Guo, L. Solden, EMERGE Field Team, EMERGE Coordinators, J. H. Kuhn, K. C.
Wrighton, M. B. Sullivan, RNA virus ecogenomics along a subarctic permafrost thaw gradient.
(2025).

28. D. Cronin, H. Holland-Moritz, D. Smith, S. T. N. Aroney, S. Hodgkins, M. Borton, Y.-F. Li, K.
Healy, I. Field &, A. Teams 2010-2017, E. I. Coordinators, M. M. Tfaily, P. Crill, C. K.
McCalley, K. Wrighton, R. Varner, G. W. Tyson, B. Woodcroft, S. C. Bagby, J. Ernakovich, V. I.
Rich, Stable states in an unstable landscape: microbial resistance at the front line of climate change.
bioRxiv [Preprint] (2025). https://doi.org/10.1101/2025.02.07.636677.

29. R. K. Varner, D. R. Cronin, P. Crill, M. Palace, C. K. McCalley, J. Deng, C. Herrick, M. Kuhn, S.
B. Hodgkins, K. McArthur, J. D. Singer, B. Bolduc, Y.-F. Li, T. A. to A. (A2A) P. Team, E. I.
Coordinators, C. Li, G. Tyson, S. Frolking, J. P. Chanton, A. Persson, S. R. Saleska, V. I. Rich,
From Archaea to the atmosphere: remotely sensing Arctic methane. bioRxiv [Preprint] (2025).
https://doi.org/10.1101/2025.02.13.638097.

30. R. K. Aziz, M. Breitbart, R. A. Edwards, Transposases are the most abundant, most ubiquitous
genes in nature. *Nucleic Acids Res.* **38**, 4207–4217 (2010).

31. C. He, R. Keren, M. L. Whittaker, I. F. Farag, J. A. Doudna, J. H. D. Cate, J. F. Banfield, Genome-
resolved metagenomics reveals site-specific diversity of episymbiotic CPR bacteria and DPANN
archaea in groundwater ecosystems. *Nat. Microbiol.* **6**, 354–365 (2021).

32. S. Roux, B. G. Paul, S. C. Bagby, S. Nayfach, M. A. Allen, G. Attwood, R. Cavicchioli, L.
Chistoserdova, R. J. Gruninger, S. J. Hallam, M. E. Hernandez, M. Hess, W.-T. Liu, T. A.
McAllister, M. A. O'Malley, X. Peng, V. I. Rich, S. R. Saleska, E. A. Eloe-Fadrosh, Ecology and
molecular targets of hypermutation in the global microbiome. *Nat. Commun.* **12**, 3076 (2021).

33. O. Zablocki, M. Michelsen, M. Burris, N. Solonenko, J. Warwick-Dugdale, R. Ghosh, J. Pett-Ridge,
M. B. Sullivan, B. Temperton, VirION2: a short- and long-read sequencing and informatics
workflow to study the genomic diversity of viruses in nature. *PeerJ* **9**, e11088 (2021).

34. M. L. Coleman, M. B. Sullivan, A. C. Martiny, C. Steglich, K. Barry, E. F. DeLong, S. W.
Chisholm, Genomic Islands and the Ecology and Evolution of Prochlorococcus. *Science*, doi:
10.1126/science.1122050 (2006).

35.   K. J. Romanowicz, B. C. Crump, G. W. Kling, Genomic evidence that microbial carbon degradation
1215   is dominated by iron redox metabolism in thawing permafrost. *ISME Commun.* **3**, 1–11 (2023).

1216   36.   M. Pester, K.-H. Knorr, M. W. Friedrich, M. Wagner, A. Loy, Sulfate-reducing microorganisms in
1217   wetlands – fameless actors in carbon cycling and climate change. *Front. Microbiol.* **3**, 19769
1218   (2012).

1219   37.   J. Limpens, F. Berendse, C. Blodau, J. G. Canadell, C. Freeman, J. Holden, N. Roulet, H. Rydin, G.
1220   Schaepman-Strub, Peatlands and the carbon cycle: from local processes to global implications – a
1221   synthesis. *Biogeosciences* **5**, 1475–1491 (2008).

1222   38.   R. AminiTabrizi, R. M. Wilson, J. D. Fudyma, S. B. Hodgkins, H. M. Heyman, V. I. Rich, S. R.
1223   Saleska, J. P. Chanton, M. M. Tfaily, Controls on Soil Organic Matter Degradation and Subsequent
1224   Greenhouse Gas Emissions Across a Permafrost Thaw Gradient in Northern Sweden. *Front. Earth*
1225   *Sci.* **8**, 557961 (2020).

1226   39.   N. Tal, R. Sorek, SnapShot: Bacterial immunity. *Cell* **185**, 578-578.e1 (2022).

1227   40.   S. Koskiniemi, J. G. Lamoureux, K. C. Nikolakakis, C. t'Kint de Roodenbeke, M. D. Kaplan, D. A.
1228   Low, C. S. Hayes, Rhs proteins from diverse bacteria mediate intercellular competition. *Proc. Natl.*
1229   *Acad. Sci.* **110**, 7032–7037 (2013).

1230   41.   S. Rakoff-Nahoum, M. J. Coyne, L. E. Comstock, An ecological network of polysaccharide
1231   utilization among human intestinal symbionts. *Curr. Biol. CB* **24**, 40–49 (2014).

1232   42.   J. J. Morris, R. E. Lenski, E. R. Zinser, The Black Queen Hypothesis: Evolution of Dependencies
1233   through Adaptive Gene Loss. *mBio* **3**, 10.1128/mbio.00036-12 (2012).

1234   43.   M. F. Abdul Halim, D. R. Fonseca, T. D. Niehaus, K. C. Costa, Functionally redundant formate
1235   dehydrogenases enable formate-dependent growth in *Methanococcus maripaludis*. *J. Biol. Chem.*
1236   **300**, 105550 (2024).

1237   44.   B. A. Lerch, D. A. Smith, T. Koffel, S. C. Bagby, K. C. Abbott, How public can public goods be?
1238   Environmental context shapes the evolutionary ecology of partially private goods. *PLOS Comput.*
1239   *Biol.* **18**, e1010666 (2022).

1240   45.   B. J. Kapili, S. E. Barnett, D. H. Buckley, A. E. Dekas, Evidence for phylogenetically and
1241   catabolically diverse active diazotrophs in deep-sea sediment. *ISME J.* **14**, 971–983 (2020).

1242   46.   R. Hidese, H. Mihara, N. Esaki, Bacterial cysteine desulfurases: versatile key players in biosynthetic
1243   pathways of sulfur-containing biofactors. *Appl. Microbiol. Biotechnol.* **91**, 47–61 (2011).

1244   47.   Exploring the selenium-over-sulfur substrate specificity and kinetics of a bacterial selenocysteine
1245   lyase. *Biochimie* **182**, 166–176 (2021).

1246   48.   S. Benler, E. V. Koonin, Recruitment of Mobile Genetic Elements for Diverse Cellular Functions in
1247   Prokaryotes. *Front. Mol. Biosci.* **9**, 821197 (2022).

1248   49.   E. A. Franzosa, X. C. Morgan, N. Segata, L. Waldron, J. Reyes, A. M. Earl, G. Giannoukos, M. R.
1249   Boylan, D. Ciulla, D. Gevers, J. Izard, W. S. Garrett, A. T. Chan, C. Huttenhower, Relating the
1250   metatranscriptome and metagenome of the human gut. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E2329-

1251   2338 (2014).

1252   50.   R. Mondav, B. J. Woodcroft, E.-H. Kim, C. K. McCalley, S. B. Hodgkins, P. M. Crill, J. Chanton,
1253   G. B. Hurst, N. C. VerBerkmoes, S. R. Saleska, P. Hugenholtz, V. I. Rich, G. W. Tyson, Discovery
1254   of a novel methanogen prevalent in thawing permafrost. *Nat. Commun.* **5**, 1–7 (2014).

1255   51.   A. Bernheim, R. Sorek, The pan-immune system of bacteria: antiviral defence as a community
1256   resource. *Nat. Rev. Microbiol.* **18**, 113–119 (2020).

1257   52.   O. X. Cordero, L.-A. Ventouras, E. F. DeLong, M. F. Polz, Public good dynamics drive evolution of
1258   iron acquisition strategies in natural bacterioplankton populations. *Proc. Natl. Acad. Sci. U. S. A.*
1259   **109**, 20059–20064 (2012).

1260   53.   M. R. Olm, A. Crits-Christoph, K. Bouma-Gregson, B. A. Firek, M. J. Morowitz, J. F. Banfield,
1261   inStrain profiles population microdiversity from metagenomic data and sensitively detects shared
1262   microbial strains. *Nat. Biotechnol.* **39**, 727–736 (2021).

1263   54.   B. B. McGivern, D. R. Cronin, J. B. Ellenbogen, M. A. Borton, E. L. Knutson, V. Freire-Zapata, J.
1264   A. Bouranis, L. Bernhardt, A. I. Hernandez, R. M. Flynn, R. Woyda, A. B. Cory, R. M. Wilson, J.
1265   P. Chanton, B. J. Woodcroft, J. G. Ernakovich, M. M. Tfaily, M. B. Sullivan, G. W. Tyson, V. I.
1266   Rich, A. E. Hagerman, K. C. Wrighton, Microbial polyphenol metabolism is part of the thawing
1267   permafrost carbon cycle. *Nat. Microbiol.* **9**, 1454–1466 (2024).

1268   55.   A. Chklovski, D. H. Parks, B. J. Woodcroft, G. W. Tyson, CheckM2: a rapid, scalable and accurate
1269   tool for assessing microbial genome quality using machine learning. *Nat. Methods* **20**, 1203–1212
1270   (2023).

1271   56.   S. T. N. Aroney, A. P. Camargo, G. W. Tyson, B. J. Woodcroft, Galah: More scalable dereplication
1272   for metagenome assembled genomes, version v0.4.0, Zenodo (2024);
1273   https://doi.org/10.5281/zenodo.10526086.

1274   57.   Y.-F. Li, I. 1 P. Coordinators, V. Rich, R. Mondav, Modified Nucleic Acid Extraction Protocol for
1275   Humic-Rich Permafrost Peat Samples. (2024).

1276   58.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data.
1277   *Bioinformatics* **30**, 2114–2120 (2014).

1278   59.   S. Nurk, D. Meleshko, A. Korobeynikov, P. A. Pevzner, metaSPAdes: a new versatile metagenomic
1279   assembler. *Genome Res.* **27**, 824–834 (2017).

1280   60.   D. D. Kang, F. Li, E. Kirton, A. Thomas, R. Egan, H. An, Z. Wang, MetaBAT 2: an adaptive
1281   binning algorithm for robust and efficient genome reconstruction from metagenome assemblies.
1282   *PeerJ* **7**, e7359 (2019).

1283   61.   M. Imelfort, D. Parks, B. J. Woodcroft, P. Dennis, P. Hugenholtz, G. W. Tyson, GroopM: an
1284   automated tool for the recovery of population genomes from related metagenomes. *PeerJ* **2**, e603
1285   (2014).

1286   62.   Y.-W. Wu, B. A. Simmons, S. W. Singer, MaxBin 2.0: an automated binning algorithm to recover
1287   genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2016).

63. B. Bushnell, J. Rood, E. Singer, BBMerge – Accurate paired shotgun read merging via overlap.
PLOS ONE **12**, e0185056 (2017).

64. D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: an ultra-fast single-node solution
for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**,
1674–1676 (2015).

65. M. Shaffer, M. A. Borton, B. B. McGivern, A. A. Zayed, S. L. La Rosa, L. M. Solden, P. Liu, A. B.
Narrowe, J. Rodríguez-Ramos, B. Bolduc, M. C. Gazitúa, R. A. Daly, G. J. Smith, D. R. Vik, P. B.
Pope, M. B. Sullivan, S. Roux, K. C. Wrighton, DRAM for distilling microbial metabolism to
automate the curation of microbiome function. *Nucleic Acids Res.* **48**, 8883–8900 (2020).

66. F. A. B. von Meijenfeldt, K. Arkhipova, D. D. Cambuy, F. H. Coutinho, B. E. Dutilh, Robust
taxonomic classification of uncharted microbial sequences and bins with CAT and BAT. *Genome*
Biol. **20**, 1–14 (2019).

67. P.-A. Chaumeil, A. J. Mussig, P. Hugenholtz, D. H. Parks, GTDB-Tk: a toolkit to classify genomes
with the Genome Taxonomy Database.

68. D. Hyatt, G.-L. Chen, P. F. LoCascio, M. L. Land, F. W. Larimer, L. J. Hauser, Prodigal:
prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**,
119 (2010).

69. S. R. Eddy, Accelerated Profile HMM Searches. *PLOS Comput. Biol.* **7**, e1002195 (2011).

70. C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas, eggNOG-mapper v2:
Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale.
Mol. Biol. Evol. **38**, 5825–5829 (2021).

71. W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

72. B. Néron, E. Littner, M. Haudiquet, A. Perrin, J. Cury, E. P. C. Rocha, IntegronFinder 2.0:
Identification and Analysis of Integrons across Bacteria, with a Focus on Antibiotic Resistance in
Klebsiella. *Microorganisms* **10**, 700 (2022).

73. Y. Boucher, M. Labbate, J. E. Koenig, H. W. Stokes, Integrons: mobilizable platforms that promote
genetic diversity in bacteria. *Trends Microbiol.* **15**, 301–309 (2007).

74. J. Cury, T. Jové, M. Touchon, B. Néron, E. P. Rocha, Identification and analysis of integrons and
cassette arrays in bacterial genomes. *Nucleic Acids Res.* **44**, 4539–4550 (2016).

75. C. Loot, G. A. Millot, E. Richard, E. Littner, C. Vit, F. Lemoine, B. Néron, J. Cury, B. Darracq, T.
Niault, D. Lapaillerie, V. Parissi, E. P. C. Rocha, D. Mazel, Integron cassettes integrate into
bacterial genomes via widespread non-classical attG sites. *Nat. Microbiol.* **9**, 228–240 (2024).

76. J. Cury, S. S. Abby, O. Doppelt-Azeroual, B. Néron, E. P. C. Rocha, Identifying Conjugative
Plasmids and Integrative Conjugative Elements with CONJscan. *Methods Mol. Biol. Clifton NJ*
2075, 265–283 (2020).

77. S. S. Abby, B. Néron, H. Ménager, M. Touchon, E. P. C. Rocha, MacSyFinder: A Program to Mine

Genomes for Molecular Systems with an Application to CRISPR-Cas Systems. *PLOS ONE* **9**,
e110726 (2014).

78. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100
(2018).

79. M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for the analysis of
massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

80. J. Guo, B. Bolduc, A. A. Zayed, A. Varsani, G. Dominguez-Huerta, T. O. Delmont, A. A. Pratama,
M. C. Gazitúa, D. Vik, M. B. Sullivan, S. Roux, VirSorter2: a multi-classifier, expert-guided
approach to detect diverse DNA and RNA viruses. *Microbiome* **9**, 37 (2021).

81. J. Guo, D. Vik, A. A. Pratama, S. Roux, M. Sullivan, Viral sequence identification SOP with
VirSorter2. (2021).

82. S. Nayfach, A. P. Camargo, F. Schulz, E. Eloe-Fadrosh, S. Roux, N. C. Kyrpides, CheckV assesses
the quality and completeness of metagenome-assembled viral genomes. *Nat. Biotechnol.* **39**, 578–
585 (2021).

83. B. Buchfink, C. Xie, D. H. Huson, Fast and sensitive protein alignment using DIAMOND. *Nat Meth*
12, 59–60 (2015).

84. R. J. P. Newell, S. T. N. Aroney, J. Zaugg, P. Sternes, G. W. Tyson, B. J. Woodcroft, Aviary:
Hybrid assembly and genome recovery from metagenomes with Aviary, version v0.9.0, Zenodo
(2024); https://doi.org/10.5281/zenodo.10806928.

85. M. Kolmogorov, D. M. Bickhart, B. Behsaz, A. Gurevich, M. Rayko, S. B. Shin, K. Kuhn, J. Yuan,
E. Polevikov, T. P. L. Smith, P. A. Pevzner, metaFlye: scalable long-read metagenome assembly
using repeat graphs. *Nat. Methods* **17**, 1103–1110 (2020).

86. R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly from long
uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

87. B. J. Walker, T. Abeel, T. Shea, M. Priest, A. Abouelliel, S. Sakthikumar, C. A. Cuomo, Q. Zeng, J.
Wortman, S. K. Young, A. M. Earl, Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. *PLOS ONE* **9**, e112963 (2014).

88. R. R. Wick, L. M. Judd, C. L. Gorrie, K. E. Holt, Unicycler: Resolving bacterial genome assemblies
from short and long sequencing reads. *PLOS Comput. Biol.* **13**, e1005595 (2017).

89. S. T. N. Aroney, R. J. P. Newell, G. W. Tyson, B. J. Woodcroft, Bin Chicken: targeted
metagenomic coassembly for the efficient recovery of novel genomes. bioRxiv [Preprint] (2024).
https://doi.org/10.1101/2024.11.24.625082.

90. B. J. Woodcroft, Default SingleM reference "metapackage" data, version 3.0.5, Zenodo (2022);
https://doi.org/10.5281/zenodo.7130825.

91. S. T. N. Aroney, R. J. P. Newell, J. Nissen, A. P. Camargo, G. W. Tyson, B. J. Woodcroft, CoverM:
Read coverage calculator for metagenomics, version v0.7.0, Zenodo (2024);
https://doi.org/10.5281/zenodo.10531254.

92. D. D. Kang, J. Froula, R. Egan, Z. Wang, MetaBAT, an efficient tool for accurately reconstructing
single genomes from complex microbial communities. *PeerJ* **3**, e1165 (2015).

93. J. N. Nissen, J. Johansen, R. L. Allesøe, C. K. Sønderby, J. J. A. Armenteros, C. H. Grønbech, L. J.
Jensen, H. B. Nielsen, T. N. Petersen, O. Winther, S. Rasmussen, Improved metagenome binning
and assembly using deep variational autoencoders. *Nat. Biotechnol.* **39**, 555–560 (2021).

94. S. Pan, C. Zhu, X.-M. Zhao, L. P. Coelho, A deep siamese neural network improves metagenome-
assembled genomes in microbiome datasets across different environments. *Nat. Commun.* **13**, 2326
(2022).

95. R. J. P. Newell, G. W. Tyson, B. J. Woodcroft, Rosella: Metagenomic binning using UMAP and
HDBSCAN, version v0.5.3, Zenodo (2024); https://doi.org/10.5281/zenodo.10460259.

96. J. Alneberg, B. S. Bjarnason, I. de Bruijn, M. Schirmer, J. Quick, U. Z. Ijaz, L. Lahti, N. J. Loman,
A. F. Andersson, C. Quince, Binning metagenomic contigs by coverage and composition. *Nat.*
Methods **11**, 1144–1146 (2014).

97. S. F. Altschul, T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, D. J. Lipman, Gapped
BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*
25, 3389–3402 (1997).

98. P. R. Sternes, G. W. Tyson, TranscriptM: A metatranscriptome bioinformatics pipeline including
metagenome contamination correction, version v0.4, Zenodo (2024);
https://doi.org/10.5281/zenodo.11157285.

99. J. Guo, pydirseq: a tool to estimate per gene depth and breadth coverage from bam files, version
v0.1.0-alpha, Zenodo (2024); https://doi.org/10.5281/ZENODO.12683197.

100. E. V. Koonin, K. S. Makarova, Y. I. Wolf, M. Krupovic, Evolutionary entanglement of mobile
genetic elements and host defence systems: guns for hire. *Nat. Rev. Genet.* **21**, 119–131 (2020).

101. P. Danecek, J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane,
S. A. McCarthy, R. M. Davies, H. Li, Twelve years of SAMtools and BCFtools. *GigaScience* **10**,
giab008 (2021).

102. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for
Statistical Computing, Vienna, Austria, 2014; http://www.R-project.org/).