Defined cellular reprogramming of androgen receptor-active prostate cancer to neuroendocrine prostate cancer

Shan Li[1,¥], Kai Song[2,¥], Huiyun Sun[1,3,¥], Yong Tao[1], Arthur Huang[4], Vipul Bhatia[1], Brian Hanratty[1], Radhika A. Patel[1], Henry W. Long[5,6], Colm Morrissey[4], Michael C. Haffner[1,7,8], Peter S. Nelson[1,7,8,9], Thomas G. Graeber[2,10,11,12*], John K. Lee[1,2,11,12,13,14,15*]

[1] Human Biology Division, Fred Hutchinson Cancer Center, Seattle, WA, 98109, USA.
[2] Department of Molecular and Medical Pharmacology, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[3] Molecular Engineering and Sciences Institute, University of Washington, Seattle, WA, 98195, USA.
[4] Department of Urology, University of Washington, Seattle, WA, 98195, USA.
[5] Department of Medical Oncology, Dana-Farber Cancer Institute, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, 02115, USA.
[6] Center for Functional Epigenetics, Dana-Farber Cancer Institute, Boston, MA, 02115, USA.
[7] Department of Pathology and Laboratory Medicine, University of Washington, Seattle, WA. 98195, USA.
[8] Clinical Research Division, Fred Hutchinson Cancer Center, Seattle, WA 98109, USA.
[9] Division of Medical Oncology, University of Washington, Seattle, WA, 98195, USA.
[10] Crump Institute for Molecular Imaging, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[11] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[12] Jonsson Comprehensive Cancer Center, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[13] Division of Hematology/Oncology, Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[14] Institute of Urologic Oncology, Department of Urology, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA
[15] Department of Molecular and Medical Pharmacology, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90095, USA

[¥] These authors contributed equally to this work.

[*] Co-corresponding authors:
Thomas G. Graeber   tgraeber@mednet.ucla.edu
John K. Lee   jklee@mednet.ucla.edu

Shan Li   sli23@fredhutch.org
Kai Song   kaisong@ucla.edu
Huiyun Sun   hsun@fredhutch.org
Yong Tao   ytao@fredhutch.org
Arthur Huang   arthurhuang2000@gmail.com
Vipul Bhatia   vbhatia@fredhutch.org
Brian Hanratty   bhanratt@fredhutch.org
Radhika A. Patel   rpatel2@fredhutch.org
Henry W. Long   henry_long@dfci.harvard.edu
Colm Morrissey   cmorriss@uw.edu
Michael C. Haffner   mhaffner@fredhutch.org
Peter S. Nelson   pnelson@fredhutch.org

SUMMARY

Neuroendocrine prostate cancer (NEPC) arises primarily through neuroendocrine

transdifferentiation (NEtD) as an adaptive mechanism of therapeutic resistance. Models to

define the functional effects of putative drivers of this process on androgen receptor (AR)

signaling and NE cancer lineage programs are lacking. We adapted a genetically defined

strategy from the field of cellular reprogramming to directly convert AR-active prostate cancer

(ARPC) to AR-independent NEPC using candidate factors. We delineated critical roles of the

pioneer factors ASCL1 and NeuroD1 in NEtD and uncovered their abilities to silence AR

expression and signaling by remodeling chromatin at the somatically acquired AR enhancer and

global AR binding sites with enhancer activity. We also elucidated the dynamic temporal

changes in the transcriptomic and epigenomic landscapes of cells undergoing acute lineage

conversion from ARPC to NEPC which should inform future therapeutic development. Further,

we distinguished the activities of ASCL1 and NeuroD1 from the inactivation of RE-1 silencing

transcription factor (REST), a master suppressor of a major neuronal gene program, in

establishing a NEPC lineage state and in modulating the expression of genes associated with

major histocompatibility complex class I (MHC I) antigen processing and presentation. These

findings provide important, clinically relevant insights into the biological processes driving NEtD

of prostate cancer.

Keywords: ASCL1, NeuroD1, neuroendocrine transdifferentiation, cellular reprogramming,

neuroendocrine prostate cancer, androgen receptor, lineage plasticity

INTRODUCTION

Prostate cancer (PC) is a hormonally-driven disease in which androgen receptor (AR) signaling

plays a fundamental role in defining a luminal epithelial cancer lineage and promoting cell

survival and proliferation. Thus, androgen deprivation therapy (ADT) and AR signaling inhibitors

(ARSIs) represent standard and effective frontline treatments for advanced PC. While the

majority of PCs initially respond to combined treatment with ADT and an ARSI, therapeutic

resistance ultimately develops over the course of months to years and leads to a disease state

known as castration-resistant PC (CRPC) [1,2]. The mechanisms by which CRPC subvert ADT

and ARSIs are numerous, but most are centered on aberrant reactivation of AR signaling

through AR gene mutations, amplification, alternative splicing, and enhancer duplication [2-5].

Beyond reactivation of the targeted AR pathway, clinical evidence implicates lineage plasticity or

a shift in cell identity as a more extreme adaptive mechanism of resistance in PC and other

epithelial malignancies. A prime example is treatment-related transdifferentiation of AR-active

PC (ARPC) to neuroendocrine PC (NEPC) in which the AR program is extinguished and

supplanted with a NE program (NEtD) [6,7], thereby leading to a cancer lineage state that no

longer depends on AR signaling and is resistant to ADT and ARSIs.

Multiple genetic and molecular events have been associated with the process of NEtD, including

amplification of *MYCN*, loss of RE1-silencing transcription factor (REST) activity, and loss of

PTEN, *TP53*, and *RB1* [8-13]. While our prior studies using human prostate epithelial

transformation models have demonstrated that overexpression of *MYCN* and constitutively

active myristoylated *AKT1* [8] or the PARCB factors (dominant-negative *TP53*, myristoylated

AKT1, short hairpin targeting *RB1*, *MYC*, and *BCL2*) can initiate NEPC [9], these experimental

systems start with AR-null basal prostate epithelial cells. Thus, they do not address how AR-

enforced luminal epithelial lineage commitment may be repressed and bypassed which is a

critical step in the NEtD of PC. The leading biological model currently available to investigate

NEtD is the LTL-331 ARPC patient-derived xenograft (PDX) model which reproducibly

undergoes transdifferentiation after passage in castrated mice to the LTL-331R NEPC tumor

model [14]. Repeated biopsies and molecular analyses of tumors from this system have provided

insights into molecular programs that are associated with NEtD [15]. Yet, this model remains

experimentally unwieldy for functional studies, and other more tractable approaches have been

lacking.

Another challenge has been the inconsistent definition of NEPC applied in the field. Clinically,

NEPC consists of a heterogeneous group of neuroendocrine tumors primarily defined by

histologic morphology that includes a range of subtypes from well-differentiated carcinoid to

aggressive large-cell and small-cell carcinomas [16]. The expression of NE markers like

chromogranin A (CHGA), synaptophysin (SYP), or CD56 (neural cell adhesion molecule 1 or

NCAM1) are often assayed by immunohistochemistry (IHC) to support or confirm a diagnosis of

NEPC. In general, individual NE markers expression can be heterogeneous, and combined

marker scores are more robust [17]. In experimental studies, phenotypic features such as NE

marker expression or the emergence of dendrite-like projections from PC cells are broadly used

as indicators of NE transdifferentiation of PC. The use of these sparse phenotypic features to

define NEPC may be problematic as they may not reflect a *bona fide* cancer lineage state

divergent from ARPC. A prime example is amphicrine prostate cancer (AMPC) which co-

expresses AR and NE markers [18]. Despite the expression of NE markers, AMPC models such

as VCaP remain sensitive to the ARSI enzalutamide, indicating a functional dependence on the

AR-driven lineage program.

Lineage-defining transcription factors and transcription factor networks specify cell states by

regulating chromatin accessibility and gene expression programs. The basic helix-loop-helix

(bHLH) proteins Achaete-scute homolog 1 (ASCL1) and neurogenic differentiation 1 (NeuroD1)

are pioneer transcriptional factors that are necessary for lineage specification during neuronal

development [19,20]. In cancer, ASCL1 and NeuroD1 define distinct subtypes of small cell lung

cancer (SCLC) [21] and NEPC [22] with differential transcriptional, epigenetic, cell surface proteomic,

and drug sensitivity profiles [23-25]. The role of ASCL1 as a driver of neuroendocrine

transformation has been supported by the genetically engineered *TCKO* mouse model of high-

grade neuroendocrine lung cancer where *Ascl1* knockout prevents tumor formation [21]. Further,

knockdown of *ASCL1* or *NEUROD1* in established human SCLC or NEPC cell line models

significantly impairs cell viability [26,27], indicating that these factors are needed for sustained

survival and growth. Insights into the functional role of these pioneer transcription factors in

mediating the acute lineage conversion from ARPC to NEPC are limited. Recent work has

shown that ASCL1 expression may be induced by ARSI therapy in PC and ASCL1 activates a

neuronal stem cell-like lineage program through chromatin remodeling mediated by the

polycomb repressive complex 2 (PRC2) [28].

In this study, we developed a genetically defined strategy adapted from the field of cellular

reprogramming to directly convert ARPC to NEPC. We identified key functional roles for ASCL1

and NeuroD1 in NE transdifferentiation, specifically new findings uncovering their ability to

silence the AR program through chromatin remodeling of the AR enhancer which is somatically

amplified in PC and global AR binding sites with enhancer activity. In contrast, we found that

loss of REST activity induces the expression of NE markers but is insufficient to drive a NE

cancer lineage state. Repeated sampling over the reprogramming time course revealed a wide

range of temporal changes in the transcriptomic and epigenomic landscapes associated with

the lineage conversion from ARPC to NEPC. Lastly, we found that ASCL1 and NeuroD1

expression, but not the loss of REST activity, induces the downregulation of class I major

histocompatibility complex (MHC) expression in PC, providing a basis for the lineage-specific

potential for immune evasion associated with NEPC.

RESULTS

Candidate factors directly reprogram ARPC to NEPC and bypass the dependence on AR

signaling

To functionally define factors required for the conversion of ARPC to NEPC, we established a

NEtD assay wherein candidate factors were introduced into the human ARPC cell lines LNCaP

and C4-2B after which they were propagated in media conditions permissive to the growth of

NEPC and analyzed for a set of phenotypic markers of ARPC and NEPC after two weeks

(**Figure 1A**). Initially, we introduced a pool of lentiviruses (LVs) expressing eight factors

associated with NEPC including dominant-negative *TP53* R175H, a short hairpin targeting *RB1*

(shRB1), *MYCN*, *ASCL1*, *SRRM4*, *NR0B2*, *BCL2*, and *KRAS* G12V. Loss of *TP53* and *RB1* as

well as amplification of *MYCN* have been identified as common genetic features associated with

NEPC [8,9,29,30]. SRRM4 is a splicing regulator that has been shown to inactivate the RE-1

silencing transcription factor (REST) which functions to suppress a major neural gene program.

Loss of REST has been implicated as a common feature in NEPC that has been reported to

functionally drive NEtD of PC [12,31]. In addition, the pioneer neural transcription factor ASCL1 is

expressed in most NEPC and appears to define a transcriptional and epigenetic subtype of the

disease [22]. NR0B2, otherwise known as short heterodimer partner (SHP), is an orphan nuclear

receptor that is highly expressed in NEPC that interacts with AR and inhibits AR activity [32].

Lastly, BCL2 and KRAS G12V were included to potentially reduce apoptosis and enhance

reprogramming efficiencies [33,34].

The introduction of this candidate factor LV pool at a multiplicity-of-infection (MOI) of four for

each factor in this assay led to a significant downregulation of AR and the AR target NK3

homeobox 1 (NKX3-1) as well as the upregulation of NE markers including forkhead box A2

(FOXA2), SYP, NCAM1, insulinoma-associated protein 1 (INSM1), and POU class 3 homeobox

2 (POU3F2 or BRN2) by immunoblot analysis (**Figure 1B**). Furthermore, immunocytochemical

analysis of the C4-2B and LNCaP cell lines transduced with the candidate factor LV pool

demonstrated a prominent loss of nuclear AR expression and increased SYP expression

(**Figure 1C**). The AR$^{low/-}$NE$^+$ cancer phenotype induced by the candidate factor LV pool was

durable based on IHC studies in tumors obtained after subcutaneous xenografting in non-

castrate male NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wjl} SzJ* (NSG) mice (**Figure S1A**).

We next sought to determine whether the PC cells reprogrammed with the candidate factor LV

pool were functionally independent of AR signaling. We implemented a potent AR-dependent

negative selection strategy (**Figure 1D**) wherein C4-2B cells were transduced with LV encoding

AR response elements (ARE) upstream of a FK506 binding protein-caspase 8 (FKBP-Casp8)

fusion. A clonal C4-2B ARE-FKBP-Casp8 line was established and transduced with either a

control LV expressing green fluorescent protein (GFP) or the candidate factor LV pool and

subjected to selection with 1 nM R1881 (to induce AR signaling and expression of FKBP-Casp8

in AR$^+$ cells) and AP20187 which is a synthetic FK506 analog (to activate FKBP-Casp8 and

induce apoptosis). We identified no viable, adherent cells in the control LV condition after five

days of selection. In contrast, we appreciated viable, adherent cells in the candidate factor LV

pool condition and ~70% of cells demonstrated functional AR bypass as they could survive

stringent selection against AR signaling (**Figures 1E and S1B**). These finding indicated that a

discrete set of factors is capable of reprogramming human ARPC, in the context of negative

selection for non-reprogrammed cells (modeling highly stringent AR inhibitor therapy), to a

cellular state that recapitulates critical phenotypic and functional features of NEPC.

The pioneer neural transcription factor ASCL1 suppresses the AR program during NEtD

but requires additional genetic alterations to maintain a proliferative phenotype

To specify the functional contribution of each candidate factor in the LV pool to NE

reprogramming, we employed a leave-one-out approach in which each factor was iteratively left

out from the LV pool in the C4-2B (**Figure 2A**) and LNCaP **(Figure S2)** cell lines. Based on

immunoblot analyses, we observed several established regulatory interactions that have been

previously identified in cancer. These included 1) ectopic expression of N-Myc leading to

elevated levels of NCAM1 in neuroblastoma [35], 2) repression of POU3F2 expression by Rb1 in

retinoblastoma [36], 3) and enhanced SOX2 expression with Rb1 inactivation in PC [13]. Most

strikingly, we observed that the exclusion of ASCL1 diminished the decrease in AR and NKX3-1

expression and likewise abrogated the otherwise induced FOXA2, NCAM1, and INSM1

expression, implicating ASCL1 as a critical player in NE transdifferentiation. In contrast, the

omission of SRRM4 restored REST expression and suppressed SYP but had little effect on AR

or other NE markers. These data are consistent with prior reports in which ectopic introduction

of SRRM4 in PC cell lines induced SYP and chromogranin B expression but did not directly

affect AR expression [31,37]. Removal of NR0B2, BCL2, and KRAS G12V had no significant

effects on AR and NE marker expression (**Figures 2A and S2A**).

Next, we employed an orthogonal factor reconstitution strategy to identify the minimal set of

factors needed to induce NEtD in the ARPC lines. Immunoblot analyses showed that ASCL1

expression alone downregulated AR markers and induced NE marker expression **(Figure 2B)**.

On the other hand, the PRNB factors (dominant-negative TP53 H175R, shRB1, MYCN, and

BCL2) were insufficient to enforce NEtD (**Figure 2B**). SRRM4 alone imparted a partial effect on

phenotype as it led to a loss of REST expression and upregulated SYP without reducing AR

markers (**Figure 2B)**. As *TP53*, *RB1*, *MYCN*, and *BCL2* alterations are common in NEPC and

other aggressive NE carcinomas we questioned why they appeared to be dispensable in our

NEtD assay. To confirm whether all factors were introduced into the cells, we encoded each

lentiviral construct with a unique 10-nucleotide barcode and performed single-cell DNA amplicon

sequencing [38] 72 hours after transduction. Of 3,870 cells analyzed, 3,785 cells (97.8%)

demonstrated reads from all barcodes, while 84 cells (2.2%) were missing one barcode and 1

cell was missing two barcodes (**Figure S2B**). We measured cell proliferation for each of the

factor reconstitution conditions as an indicator of proliferative fitness in our experimental system.

PRNB significantly enhanced proliferation while ASCL1 was detrimental to cell viability over time

in the ARPC lines (**Figure 2C and Figure S2C**). This effect of ASCL1 is consistent with prior

reports that ASCL1 promotes cell cycle exit and terminal differentiation when overexpressed in

neural progenitor cells and glioblastoma stem cells [39]. Notably, SRRM4 promoted proliferation

but alone was insufficient to rescue the effects of ASCL1 without PRNB. These data support the

concept that the appropriate genetic and proliferative context may be an important determinant

of ASCL1's impact on the emergence of NEPC and highlight the idea that overall fitness may be

a combination of proliferation- and lineage-linked benefit. In other words, this mechanism of

therapeutic escape may require a change in lineage without a substantial decrease in

proliferative fitness.

NeuroD1 is competent to induce neuroendocrine lineage reprogramming

We have previously observed that most NEPCs can be assigned to ASCL1^{high}, NeuroD1^{high}, or

mixed ASCL1/NeuroD1 groups based on transcriptional profiling [26]. These findings have also

been described by other groups wherein ASCL1 and NeuroD1 appear to define molecular

subtypes of NEPC [22] and the neuroendocrine subtypes of small cell lung cancer [21]. To

determine whether NeuroD1 can also drive lineage conversion from ARPC to NEPC, we

performed the reprogramming assay using the C4-2B and LNCaP ARPC lines by replacing

ASCL1 with NeuroD1 or combining NeuroD1 with ASCL1. The addition of NeuroD1 to PRNB

and SRRM4 substantially diminished AR markers and increased NE marker expression (**Figure**

3A). We discovered that NeuroD1 induced ASCL1 expression and heightened overall NE

marker panel expression. Bulk RNA-seq gene expression analysis of the reprogrammed cell

lines with respect to established 22-gene AR signature and NE signature scores revealed a

significant de-enrichment of the AR gene signature and enrichment of the NE gene signature in

the PRNBSA (PRNB, SRRM4, and ASCL1) and PRNBSN (PRNB, SRRM4, and NeuroD1)

conditions relative to the GFP control (**Figure 3B**). As C4-2B is derived from the LNCaP cell line,

we also confirmed the reproducibility of these findings in the independent ARPC cell line MDA

PCa 2b using this cellular reprogramming assay (**Figure S3A, C**).

Several neural transcription factors including FOXA2, INSM1, POU3F2, and SOX2 have also

been described as documented drivers or regulators of NE transdifferentiation in PC and other

cancer types [13,40,41]. We sought to determine whether these factors are independently

competent to induce NE lineage reprogramming. We introduced PRNB, SRRM4, and each of

the above listed neural transcription factors individually to the C4-2B line in the NEtD assay. In

contrast to ASCL1 and NeuroD1, these factors were by themselves incapable of NE lineage

reprogramming (**Figure S3B**). As ASCL1 and NeuroD1 also appear to regulate the expression

of FOXA2 and INSM1, ASCL1 and NeuroD1 may sit atop a hierarchy of multiple NE-associated

transcription factors required in combination to establish and maintain a NEPC transcriptional

regulatory network.

To confirm the relevance of our NE transdifferentiation models in PC and to use omic-scale data

to minimize the effect of individual marker heterogeneity, we performed partial least squares-

discriminant analysis (PLS-DA) of RNA-seq gene expression data from previously characterized

human metastatic CRPC patient specimens annotated as ARPC or NEPC [42] and projected the

C4-2B factor-reconstituted samples onto them (**Figure 3C**). This analysis showed that PRNB

alone did not induce a substantial shift away from the ARPC phenotype. Further, the addition of

PRNB and SRRM4 (PRNBS) resulted in a minor, intermediate shift toward the NEPC phenotype.

Only upon the addition of ASCL1 or NeuroD1 to PRNB (PRNBA and PRNBN) or PRNBS

(PRNBSA and PRNBSN) did we observe a dramatic gene expression program-based shift from

ARPC to NEPC. Similar results were observed in in MDA PCa 2b cell line (**Figure S3D**). These

data highlight the relative lineage impacting importance of these factors in NEtD of PC.

SRRM4 promotes alternative splicing and inactivates REST through the inclusion of a neural

exon that generates the REST-4 isoform [31,43]. We confirmed the presence of alternatively

spliced *REST* mRNA in C4-2B reprogramming conditions that included SRRM4 (**Figure S4A**).

Further, we performed PLS-DA analysis on alternative splicing data from Beltran et al., 2016

and projected the C4-2B reprogrammed samples onto them. This analysis indicated that

SRRM4 drives the alternative splicing phenotype associated with NEPC with a marginal

contribution from ASCL1 or NeuroD1 (**Figure S4B**).

Dynamic changes in cancer phenotype, transcriptional and epigenetic landscapes during

NE reprogramming

Having established a defined and tractable system that recapitulates NE transdifferentiation of

PC, we sought to investigate the dynamic molecular and regulatory changes underlying this shift

in cancer lineage state. We applied the NE reprogramming assay to the ARPC C4-2B line and

temporally sampled multiple timepoints within the 14-day reprogramming period (D2, D3, D4,

D5, D8, D11, and D14) for downstream analyses (**Figure 4A**). Immunoblot analyses revealed a

reduced expression of AR markers (AR and NKX3-1) as early as D2, with expression of the NE

markers INSM1 and FOXA2 evident shortly thereafter on D4-5 (**Figure 4B**). These findings

indicate that downregulation of the AR program precedes the establishment of the NE program

in our experimental system. Transcriptome profiling by bulk RNA-seq of the C4-2B cells

reprogrammed with PRNBSA or PRNBSN recapitulated the findings from the immunoblot

analyses where downregulation of *AR* expression preceded the induction of *INSM1* expression

(**Figures 4C and D**). Notably, PCA analysis of the time course RNA-seq gene expression data

from C4-2B cells reprogrammed with PRNBSA or PRNBSN, but not PRNBS, showed an arc-like

trajectory during NEtD, indicating dynamic changes in gene expression programs during ARPC

to NEPC reprogramming (**Figures S5A-C**). This arc-like trajectory is also broadly observed in

normal and cancer development and cellular differentiation [44-47].

We next used PLS-DA projections from the time course RNA-seq gene expression data to

further investigate the temporal nature of the phenotypic shift from ARPC to NEPC during NEtD.

We distinguished three general phases of reprogramming in the C4-2B cells modified with

PRNBSA and PRNBSN that could be defined as "early" on D2, "intermediate" from D3-5, and

"late" from D8-14 (**Figure 4E**). In contrast, these phases were not apparent in C4-2B cells

transduced with PRNB and SRRM4 (**Figure S5D**), further supporting the previous finding that

ASCL1 or NEUROD1 is needed for full lineage shift via NEtD.

Lineage reprogramming of ARPC to NEPC is largely driven by epigenetic dysregulation and

reprogramming as few genetic alterations apart from the enrichment of *TP53* and *RB1*

mutations and genomic amplification of *MYCN* have been identified in NEPC [30,42,48,49]. To

investigate global changes in the epigenetic landscape during the transition from ARPC to

NEPC, we performed Cleavage Under Targets & Release Using Nuclease (CUT&RUN) analysis

on PRNBSA reprogrammed C4-2B cells over the 14-day time course to examine genome-wide

changes in histone modifications, including monomethylation of histone H3 at lysine 4

(H3K4me1) which is commonly associated with gene enhancers, acetylation of histone H3 on

lysine 27 (H3K27ac) which indicates both active promoters and enhancers [50,51], and

trimethylation of histone H3 at lysine 4 (H3K4me3) which marks active gene promoters.

Unsupervised principal component analysis (PCA) showed that these histone modifications shift

significantly during the time course (**Figure 4F**), indicating that dynamic epigenetic changes

occur during NEtD.

As ASCL1 is a pioneer transcription factor, we queried whether there may be time-dependent

changes in putative ASCL1-mediated gene regulation during the acute conversion of ARPC to

NEPC. We integrated the RNA-seq gene expression data and ASCL1 CUT&RUN genome

occupancy data over the 14-day reprogramming period to identify genes both in proximity to

ASCL1 binding and upregulated in PRNBSA compared to PRNBS. Gene set enrichment

analysis (GSEA) revealed several pathways enriched early during NE reprogramming including

the CDC25 pathway, TGFβ1 pathway, and estradiol response pathway (**Figure 5A**). In contrast,

ASCL1 targets, PRC2 targets including those of the core subunits embryonic ectoderm

development (EED) and SUZ12, and neuronal fate commitment pathways were significantly

enriched later in the NE transdifferentiation process. The pathways enriched early may

represent components of an unstable transcriptional state prior to the establishment of a stable

NEPC lineage, consistent with our observation of an arc-like trajectory.

Next, we evaluated the dynamic activities of active enhancers, super-enhancers, and active

promoters in the NE reprogramming assay. We defined active enhancers based on overlapping

H3K4me1 and H3K27ac peaks and characterized super-enhancers by applying the Rank

Ordering of Super-Enhancer (ROSE) algorithm to H3K27ac peaks over the reprogramming time

course [52]. This analysis revealed a substantial temporal shift in active enhancers (**Figure S6A**)

and super-enhancers (**Figure 5B**). Gene ontology (GO) analysis of downregulated active

enhancer- and super-enhancer-associated genes over time revealed a highly significant

enrichment of androgen response genes including the Hallmark Androgen Response gene set

(**Figures S6B and 5C**). In contrast, genes associated with upregulated active enhancers over

time were enriched for endocrine therapy resistance pathways and targets of the PRC2 complex

core subunits SUZ12 and enhancer of zeste homolog 2 (EZH2) (**Figure S6C**). We also

examined active promoters by determining the overlap of H3K4me3 and H3K27ac peaks over

the reprogramming time course (**Figure S7A**). Genes associated with downregulated active

promoters over time were enriched in androgen response and AR regulated targets (**Figure**

S7B). On the other hand, GO analysis of upregulated active promoters over time did not

highlight any gene sets of statistical significance.

ASCL1/NeuroD1 inhibits AR expression by remodeling the somatically acquired AR

enhancer

To further investigate how ASCL1 and NeuroD1 may suppress AR expression and signaling

during NE reprogramming, we performed single-cell (sc) multiome RNA-seq and ATAC-seq

analysis on D14 after reprogramming of C4-2B cells with PRNB, PRNBS, PRNBSA, and

PRNBSN. The PRNB and PRNBS conditions were included as controls as these factor

combinations have no apparent effect on the AR program. Pseudotime analysis revealed a

progression from PRNB to PRNBS to PRNBSA to PRNBSN which was associated with a

heightened NEPC score and reduced *AR* expression (**Figure S8A-D**). In addition, we observed

two distinct clusters within the PRNBSA and PRNBSN conditions indicating heterogeneous cell

populations likely due to differences in expression levels of *ASCL1* and *NEUROD1* (**Figures**

S8E, F).

Most strikingly, the integration analysis of RNA-seq and ATAC-seq data identified a significant

reduction in chromatin accessibility of the somatically acquired *AR* enhancer, but not the *AR*

gene body, that was associated with reduced *AR* expression in the PRNBSA and PRNBSN

conditions (**Figure 6A**). In contrast, the reduction in chromatin accessibility of the somatically

acquired *AR* enhancer was not seen in the control PRNB and PRNBSN conditions (**Figure 6A**).

The somatically acquired enhancer of *AR* and its amplification independent of the *AR* gene body

has previously been shown to be a noncoding driver of CRPC [5]. Attesting to the clinical

relevance of this finding, we identified a hepatic metastasis from a patient with lethal mCRPC

who underwent rapid autopsy as part of the University of Washington Tissue Acquisition

Necropsy (UW TAN) program that showed a mixed population of $AR^+/ASCL1^-$ and $AR^-/ASCL1^+$

tumors cells by immunofluorescence staining (**Figure 6B**). Similar to the findings from our NE

reprogramming studies, scATAC-seq analysis of this metastatic tumor showed loss of chromatin

accessibility at the enhancer of *AR* but not the *AR* gene body in $AR^-/ASCL1^+$ tumor cells

compared to $AR^+/ASCL1^-$ cells (**Figure 6C**).

We next sought to investigate the chromatin accessibility of genome-wide AR binding sites

(ARBS) with enhancer activity. ARBS with enhancer activity have previously been characterized

using Self-Transcribing Active Regulatory Regions sequencing (STARR-seq) in the LNCaP cell

line and classified as constitutively active enhancers, inducible enhancers, and inactive

enhancers based on their responsiveness to the androgen dihydrotestosterone [53]. We evaluated

the chromatin accessibility at these ARBS and identified a global reduction in the accessibility of

ARBS with enhancer activity from least reprogrammed (PRNB) to most reprogrammed

(PRNBSA or PRNBSN) cell lines (**Figure 6D**). In contrast, a random selection of genomic sites

of similar intervals across all chromosomes, as a control, did not show reduction in chromatin

accessibility. These results further demonstrate the acute epigenetic rewiring of PC during NEtD

to silence an AR-driven PC lineage program.

ASCL1/NeuroD1 but not loss of REST via SRRM4 activity establish a NE transcriptional

network

Lineage-defining transcription factors including neural lineage transcriptions factors have been

reported to auto-activate their own expression by binding to super-enhancers to establish a

positive feedback loop required to sustain a lineage program [54]. We investigated the potential

for self-regulation of ASCL1 and NeuroD1 in our NE reprogramming assay based on differential

RNA-seq read alignment of exogenous proviral transcripts and endogenous transcripts (**Figure**

S9A). We specifically leveraged the presence of the synonymous single-nucleotide variant

NM_004316.4(ASCL1);c.627C>G encoded in the ASCL1 lentivirus to distinguish exogenous

and endogenous reads. The introduction of either exogenous ASCL1 or NeuroD1 to PRNB or

PRNBS resulted in the induction of endogenous *ASCL1* expression (**Figure S9B**). This was

also confirmed by the presence of reads mapping to the 5' and 3' untranslated regions (UTRs)

which are present in endogenous transcripts but not encoded in exogenous proviral transcripts.

Using a similar strategy of mapping reads to the 5' and 3' UTR, we also found that exogeneous

NeuroD1 could activate endogenous *NEUROD1* expression (**Figure S9C**).

We then investigated how the different conditions in our NE reprogramming assay may impact

the establishment of a transcriptional regulatory network (TRN) underlying an NEPC lineage

state. We first focused on the differential expression of transcription factors due to SRRM4

(PRNBS vs. PRNB), ASCL1 (PRNBSA vs. PRNBS), and NeuroD1 (PRNBSN vs. PRNBS). We

observed a significant induction of NE-associated transcription factors including INSM1, SOX2,

SOX4, NK2 homeobox 2 (NKX2-2), and FOXA2 related to ASCL1 and NeuroD1 (**Figure 7A**). In

stark contrast, not a single transcription factor demonstrated statistically significant differential

expression upon the addition of SRRM4 to PRNB. Consistent with these findings, footprinting

analysis of ATAC-seq data identified a significant enrichment in binding motifs associated with

these NE-associated transcription factors in conditions of enforced ASCL1 and NeuroD1

expression but not SRRM4 expression (**Figure 7B**). Global chromatin accessibility profiles are

also informative as they can differentiate lineage states in normal tissues and cancer. When we

projected pseudo-bulk scATAC-seq data of reprogrammed C4-2B lines on the PCA space

defined by ATAC-seq data of previously published ARPC and NEPC PDX models [22], we found

that C4-2B lines reprogrammed with PRNBSA and PRNBSN clustered with the NEPC PDXs

while those reprogrammed with PRNB and PRNBS were more similar to ARPC PDXs (**Figure**

7C). Further, we found that the expression of NEPC-associated cell surface targets (*DLL3*,

L1CAM, *CEACAM5*, *RET* and *SEZ6*) were significantly increased and the ARPC-associated cell

surface targets (*FOLH1, PSCA, STEAP1*, and *TACSTD2*) were significantly decreased upon the

addition of ASCL1 or NeuroD1. In contrast, SRRM4 alone did not have a prominent effect on

the expression of these cell surface targets (**Figure 7D**). Taken together, these findings indicate

that, unlike ASCL1 and NeuroD1, SRRM4 activity and the resultant loss of REST repressor

activity alone may be insufficient to drive a NEPC lineage due to the absence of active

transcriptional and epigenetic reprogramming.

Downregulation of major histocompatibility complex class I (MHC I) in NEPC is

functionally attributable to ASCL1/NeuroD1-driven programs

The downregulation of MHC I antigen presentation machinery by cancer to overcome immune

surveillance and avoid immune destruction is an established hallmark of cancer. In particular,

this phenomenon has been associated with cancers with neuroendocrine differentiation [55]. In a

recent study, examination of SCLC data sets showed that tumors with low NE scores

demonstrated higher expression of MHC I genes while those with high NE scores showed

decreased MHC I and reduced immune infiltration [56]. Further, cancer cells are capable of co-

opting PRC2 in a lineage-specific fashion to silence the MHC I antigen processing and

presentation pathway [57]. Mechanistically, mutation of the catalytic domain or pharmacologic

inhibition of the PRC2 core subunit EZH2 was shown to impair MHC I repression, highlighting

the critical role of PRC2 in this process.

We therefore asked whether a similar phenomenon may be operative in NEPC and could be

functionally attributable to lineage-defining ASCL1 and/or NeuroD1 expression. We first

examined a published RNA-seq gene expression dataset of mCRPC [42] including CRPC-Ad

(adenocarcinoma) and CRPC-NE (NEPC) patient specimens and found that NEPC was

associated with reduced MHC I antigen processing and presentation gene signature scores as

well as lower *B2M* (beta-2-microglobulin, a critical component of MHC I) expression (**Figures**

8A and S10). A similar analysis using published data from tumors in the UW TAN PC rapid

autopsy cohort also showed a reduced overall expression of *B2M* in the NEPC samples (**Figure**

8B). We then evaluated RNA-seq gene expression data from our reprogrammed C4-2B lines

and correlated each condition and NE signature score with MHC I antigen processing and

presentation gene signature scores. Consistent with prior literature, we discovered a reduction

in MHC I gene expression with increasing NE differentiation (**Figure 8C**). Downregulation of

MHC I gene expression started as early as D2 along with AR pathway repression (**Figure S11**).

Furthermore, the addition of SRRM4 to PRNB did not have an appreciable effect on MHC I

expression, indicating that loss of REST does not itself contribute to the downregulation of MHC

I. In contrast, conditions where ASCL1 or NeuroD1 were added did demonstrate reduced

expression of MHC I gene sets. We found that *B2M* expression was especially downregulated in

conditions reprogrammed with NeuroD1 which were also associated with prominent NE

signature scores (**Figure 8C**). These findings were in line with integrated scRNA-seq and

scATAC-seq data from reprogrammed C4-2B cells showing an inverse correlation between

NEPC score and *B2M* expression (**Figure 8D-F**) To confirm these findings at the protein level,

we performed flow cytometry on reprogrammed C4-2B cells and showed that the PRNBSN

condition was associated with a significant reduction in B2M positivity relative to the GFP control

condition (29.2% vs. 60.9%) (**Figure 8G**). In addition, we observed that cells with diminished

MHC I gene expression including B2M showed high EZH2 expression in both C4-2B and MDA

PCa 2b cell lines (**Figure 8C and Figure S12**). Together, these findings functionally attribute

the lineage-specific downregulation of MHC I in NEPC (relative to ARPC) to the lineage-defining

transcription factors ASCL1 and NeuroD1.

DISCUSSION

AR signaling is pivotal in defining the luminal epithelial lineage of most PCs. However, potent therapies have been developed that apply stringent selective pressure against AR signaling, leading to the increased therapy-driven clinical presentation of lineage plasticity in PC. One such example is the conversion of ARPC to NEPC where the epithelial cancer state enforced by the AR program is replaced by a neural/NE cancer lineage [7]. Alternative AR-null PC lineages have also been identified including double-negative prostate cancer (DNPC) which lacks both AR and NE marker expression [18,58]. Further, this lineage plasticity in the setting of targeted therapies has also been observed in other epithelial cancer types such as epidermal growth factor receptor (EGFR)-mutant lung adenocarcinoma in which transformation to SCLC occurs in up to 15% of patients treated with EGFR tyrosine kinase inhibitors [59]. However, the mechanisms by which AR (or EGFR) expression and signaling are effectively silenced during this transition have been poorly understood. The ability to functionally characterize these dynamic processes in a defined manner by reverse genetics has been limited by the lack of available model systems allowing for rapid manipulation and repeated sampling over time. Furthermore, due to documented heterogeneity, the application of narrow phenotypic markers to define NEPC has also been problematic and complicated our understanding of this PC lineage state.

We established an *in vitro* system in which we can reproducibly and durably reprogram established human ARPC cell lines to NEPC with defined genetic factors over a specified time course. This tractable system recapitulates key features of the process of NEtD including global, dynamic transcriptional and epigenetic reprogramming. Importantly, we uncovered the ability of the pioneer neural transcription factors ASCL1 and NeuroD1 to suppress AR expression and signaling by chromatin remodeling of the somatically acquired enhancer of AR and global ARBS with enhancer activity. These findings show that the activity of a new lineage-defining

transcription factor can concurrently silence the prior cancer lineage program while establishing

a divergent lineage. An immediate question that emerges from these findings is whether AR and

ASCL1 or NeuroD1 activity is mutually exclusive in individual PC cells. Emerging single-cell

sequencing data sets of human and mouse models of NEPC indicate that this may be the case

[60,61]. The competence of ASCL1 and/or NeuroD1 to silence the AR program in prostate cancer

may be a major constraint to NEtD, providing a potential basis for why NEPC only occurs in 15%

of therapy-driven mCRPC cases. How the expression of these neural/NE transcription factors is

initially induced in PC also remains to be determined and warrants further investigation.

Our studies have also distinguished the relative contributions of SRRM4 and ASCL1 or

NeuroD1 to NE reprogramming of PC, and yield insight into the problematic nature of using

sparse protein (IHC) or RNA analysis based on only a few markers to define NEPC lineage

states. In our reprogramming studies, we find that SRRM4 activity results in the loss of REST

which activates neural/NE gene expression including SYP and CHGA. However, unlike ASCL1

or NeuroD1, SRRM4 does not induce a new transcriptional regulatory network based on

transcription factor expression and inferred transcription factor activity analyses, does not elicit

epigenomic shifts consistent with NEtD, does not affect AR expression and signaling, and does

not alter the expression of ARPC-associated cell surface targets. Thus, our findings indicate that

a reliance on a sparse set of NE markers, including highly canonical markers, to define NEPC

can be flawed as their expression may be independent of true NEPC lineage states. A major

clinical consideration is that mCRPCs that express select NE markers but have not undergone

NEtD may still be responsive to therapies intended for ARPC including ARSIs and PSMA

radioligand therapy.

In our NE reprogramming assay, we have found that ASCL1 and NeuroD1 but not other NE-

associated transcription factors are competent to induce NEtD of PC. This supports a

transcription regulatory network model where ASCL1 and NeuroD1 sit atop a hierarchy of NE-

associated transcription factors necessary to establish self-sustaining NEPC transcriptional

convergence. We have shown that ASCL1 and NeuroD1 are capable of self-regulation but the

full set of regulatory interactions between these and other NE-associated transcription factors

are unknown in NEPC. Future studies to define these relationships through genetic loss-of-

function screening using this and other NEtD assays elucidate these functional interactions,

identify potential vulnerabilities, and promote therapeutic development to inhibit NEtD under the

pressures of otherwise effective targeted therapies.

Prior work identified suppression of MHC I antigen processing and presentation in

neuroendocrine cancers and a functional association with lineage-specific PRC2 activity [55-57].

Because our studies directly implicated ASCL1 and NeuroD1 in lineage reprogramming of

ARPC to NEPC as well as the induction of PRC2 activity, we examined the expression of MHC I

pathway gene sets and discovered that MHC I pathway genes were suppressed in the effective

ASCL1- or NeuroD1- driven reprogramming conditions. In contrast, the addition of SRRM4 and

subsequent loss of REST activity had little to no effect on the MHC I pathway. These findings

support that downregulation of the MHC I antigen processing and presentation pathway in

NEPC is functionally attributable to ASCL1 and NeuroD1 activity. Immune checkpoint inhibitors

have demonstrated limited utility in mCRPC except in the case of DNA mismatch repair

deficiency [62]. Thus, our results support therapeutic efforts to induce immunogenic responses in

NEPC using combinations with epigenetic inhibitors such as those targeting EZH2, histone

deacetylases, or demethylases to enhance MHC I expression [63,64].

Overall, this study provides important mechanistic details of NEtD of PC by highlighting key

lineage-defining activities of neural pioneer transcription factors in this process. These findings

have important implications on the molecular definition of NEPC and therapeutic considerations

related to the application of targeted agents and immune-based therapies.

METHODS

Cell lines

LNCaP (Cat# CRL-1740, RRID: CVCL_0395), C4-2B (Cat# CRL-3315, RRID: CVCL_4784),

MDA PCa 2b (Cat# CRL-2422, RRID: CVCL_4748) cell lines were purchased from ATCC. All

cell lines were validated by short tandem repeat analysis after receipt and on a yearly basis.

LNCaP and C4-2B were maintained in RPMI medium supplemented with 10% FBS, 100 U/mL

penicillin and 100 µg/mL streptomycin, and 4 mmol/L GlutaMAX. MDA PCa 2b was maintained

in F-12K medium supplemented with 20% FBS, 25 ng/ml cholera toxin, 10 ng/ml mouse

epidermal growth factor, 0.005 mM phosphoethanolamine, 100 pg/ml hydrocortisone, 1X

insulin-transferrin-selenium (ITS-G), 100 U/mL penicillin and 100 µg/mL streptomycin, and 4

mmol/L GlutaMAX.

Lentiviral constructs and lentivirus production

Double-barcoded lentiviral vectors used as a lentiviral backbone vector were generated from

FU-CGW by sequentially inserting matched 10-nucleotide barcodes into the PacI site distal to

the HIV FLAP using the Quick Ligation Kit (New England Biolabs) and PCR amplification of the

WPRE sequence and barcode with insertion into the ClaI site proximal to the 3' LTR by HiFi

DNA Assembly (New England Biolabs). U6 promoter and shRNA cassettes were isolated by

digesting pLKO.1 TRC shRNA clones with PspXI and EcorRI and were inserted into the

digested double-barcoded plasmid as previously described [8,9]. Lentiviral vectors expressing

dominant-negative TP53, BCL2, MYC, SRRM4, NR0B2, KRAS, ASCL1 and NeuroD1 were

generated by PCR amplification of the open reading frames and subcloning into the EcoRI site

of double-barcoded lentiviral vectors by NEBuilder HiFi DNA Assembly (New England Biolabs).

Lentiviruses were prepared and titered as previously described [8].

Immunoblotting

Whole cell extracts were collected in 9 M urea lysis buffer and quantified using the Pierce Rapid

Gold BCA Protein Assay Kit (Thermo Fisher Scientific). Protein samples were fractionated by

SDS-PAGE using Bolt Bis-Tris Plus gels and transferred to a nitrocellulose membrane using

Bolt Transfer Buffer according to the manufacturer's instructions (Invitrogen). Membranes were

blocked with 5% non-fat milk in PBST (DPBS + 0.5% Tween 20) for 30 minutes while shaking at

room temperature, then incubated with primary antibodies at 4°C overnight. Membranes were

washed three times for 5 minutes with PBST and incubated with horseradish peroxidase (HRP)

conjugated anti-mouse or anti-rabbit secondary antibodies for 1 hour at room temperature. Blots

were washed three times for 5 minutes each with PBST and developed with Immobilon Western

Chemiluminescent HRP Substrate (EMD Millipore) for 3 minutes at room temperature. Blot

images were acquired using a ChemiDoc Imaging System (Bio-Rad).

Immunohistochemistry (IHC) and immunofluorescence (IF) studies

Cells and xenografts were fixed in 10% buffered formalin for 12 hours and then embedded in

HistoGel (Thermo Fisher Scientific) and paraffin, sectioned, and mounted on glass slides

(Thermo Fisher Scientific). IHC was performed as previously described [26]. In brief, slides

containing formalin-fixed paraffin-embedded sections were deparaffinized in xylene and

rehydrated in 100%, 95%, and 70% ethanol. Slides were then heated in antigen retrieval buffer

(0.2 mol/L citric acid and 0.2 mol/L sodium citrate) within a pressure cooker followed by PBS

wash. Slides were blocked with 2.5% horse serum for 30 minutes and then incubated with

primary antibody diluted in 2.5% horse serum overnight at 4°C. HRP was detected with

ImmPRESS-HRP anti-mouse or anti-rabbit IgG peroxidase detection kits (Vector Laboratories)

and staining was visualized with DAB peroxidase substrate (Dako). Slides were counterstained

with hematoxylin and dehydrated for mounting.

Immunofluorescence studies using specific antibodies were carried out on archival formalin

fixed paraffin embedded tissues. In brief, 5 μm paraffin sections were de-waxed and rehydrated

following standard protocols. Antigen retrieval consisted of steaming for 40 minutes in Target

Retrieval Solution (Agilent). Slides were then washed and equilibrated in TBS-Tween buffer

(Sigma) for 10 minutes. Primary antibodies were applied at at 37°C for 60 minutes. Sequential

dual-immunofluorescence labeling was carried out using Tyramide SuperBoost kits (Thermo

Fisher).

Neuroendocrine transdifferentiation assay

Cells were seeded in 6-well tissue culture plates at a density of 3×10^5 cells per mL in 3 mL of

RPMI medium supplemented with 10% FBS, 100 U/mL penicillin and 100 μg/mL streptomycin,

and 4 mmol/L GlutaMAX. Cells were transduced approximately 4-6 hours after seeding at a

defined multiplicity of infection (MOI) of 4 for each lentivirus. 72 hours after transduction, cells

were trypsinized, washed, and transferred to 100 mm tissue culture plates in 15 mL of neural

stem cell media (N-SCM) consisting of Advanced DMEM/F12 medium supplemented with 1X

serum-free B27, 10 ng/mL recombinant human bFGF, 10 ng/mL recombinant human EGF, 100

U/mL penicillin and 100 μg/mL streptomycin, and 4 mmol/L GlutaMAX. Media were replenished

every 3-4 days. Cells were collected 14 days post-transduction for analysis.

Mouse studies

All animal care and studies were performed in accordance with an approved Fred Hutchinson

Cancer Center Institutional Animal Care and Use Committee protocol and Comparative

Medicine regulations. Eight-week-old male NSG (NOD-SCID-IL2Rγ-null, RRID:BCBC_4142)

mice were obtained from the Jackson Laboratory. A total of 1×10^6 cells from each C4-2B

reprogramming condition were suspended in 100 μL of cold Matrigel (Corning) and implanted by

injection subcutaneously into NSG mice.

Single-cell DNA amplicon sequencing library preparation and sequencing

A custom panel was designed for the Mission Bio Tapestri to amplify segments of ten mouse genes at two exons each, the 5' and 3' lentiviral barcodes, and lentiviral GFP. Libraries were generated from C4-2B cells transduced with PRNBSA LV pool at MOI of 4 for each lentivirus using the Mission Bio Tapestri Single-cell DNA Custom Kit according to the manufacturer's recommendations. Single cells (3,000 to 4,000 cells per µl) were resuspended in Tapestri cell buffer and encapsulated using a Tapestri microfluidics cartridge, lysed, and barcoded. Barcoded samples were subjected to targted PCR amplification and PCR products were removed from individual droplets, purified with KAPA Pure Beads (Roche Molecular Systems), and used as a template for PCR to incorporate Illumina P7 indices. PCR products were purified by KAPA Pure Beads, and quantified by Qubit dsDNA High Sensitivity Assay (ThermoFisher Scientific). Sample quality was assessed by Agilent TapeStation analysis. Libraries were pooled and sequenced on an Illumina MiSeq or HiSeq 2500 with 150 bp paired-end reads in the Fred Hutchinson Cancer Center Genomics Shared Resource.

Cell proliferation assay

For cell proliferation assays, the CellTiter-Glo® 2.0 Cell Viability Assay system (Promega) was used according to the manufacturer's instructions. 5×10^4 cells were plated into each well in a 96-well and 25 µL per well of CellTiter-Glo® 2.0 reagent was added at each endpoint for analysis. After a 30 minute incubation at room temperature, luminescence was measured using a BioTek Synergy H1 Multimode Reader (Agilent). Six replicate wells per time point were used to obtain measures of cell proliferation.

RNA-seq

Total RNA was extracted with a Purelink[TM] RNA mini kit (Invitrogen). cDNA libraries were

prepared from isolated RNA using the Illumina Stranded mRNA prep kit. High-throughput

sequencing with 50 bp paired-end reads was performed using an Illumina NovaSeq 6000 SP.

Sequencing reads were mapped to human genome reference hg38 and gene expression was

quantified and normalized using the Toil RNA-seq pipeline (v4.1.2). Sequence adapters were

trimmed using CutAdapt (v1.9) [65]. Trimmed sequences were then aligned to human reference

genome GRCh38 using STAR (v2.4.2a) [66] and gene expression quantification was performed

using RSEM v1.2.25 [67]. RSEM expected counts were subsequently normalized by upper-

quartile (UQ) normalization for downstream analysis.

The enrichment scores were calculated in R using the GSVA package using a previously

published set of 8 AR-regulated genes for AR-activity scores and the 14 genes in the NEURO I

and NEURO II gene sets for NEURO scores[18]. Differential gene expression analysis was

performed using the R package DEseq2 (v1.36.0) RNA-seq data from Beltran et al., 2016 was

processed through the Toil RNA-seq pipeline (v4.1.2). Gene expression data was projected

onto the Beltran et al., 2016 PCA space to examine reprogramming towards the NEPC

phenotype. Gene signature scores were calculated based on RNA-seq expression data using

the R package GSVA [68]. The neuroendocrine score was previously described based on CRPC

patient data [42]. Briefly, 70 genes were used to construct the score based on differential

expression or methylation between CRPC-NE and CRPC-Ad or relevance to CRPC-NE based

on the literature. The mean expression of the 70 genes was calculated to form the NE score.

The scores were visualized in heatmap format using ComplexHeatmap (v2.12.0) [69].

Differential RNA alternative splicing analysis

Differential RNA alternative splicing (AS) between vector transduction conditions was calculated

from bulk RNA-sequencing using the rMATS (version 4.1.1)[70]. Gene and transcript annotations

were sourced from Ensembl GRCH38 release 104[71]. The program was run under the statoff

mode to obtain the percent spliced in (PSI) matrix for all C42B samples. AS events that were

defined by both junction counts and exon counts were used for downstream analysis. The PSI

values of all five AS events: skipped exon (SE), alternative 5' splice site (A5SS), alternative 3'

splice site (A3SS), mutually exclusive exons (MXE) and retained intron (RI) were aggregated

together to form a large matrix for PCA. CRPC-Adeno and CRPC-NE Beltran et al 2016 patient

RNAseq data was processed using the same rMATS workflow for AS calling. C42B was

projected onto Betlran et al 2016 PSI PCA space for examining the remodeling of AS by

oncogene transduction towards the NE direction.

CUT&RUN

Cells were harvested by centrifugation at 1200 rpm for 3 minutes in a swinging bucket rotor and

washed twice in ice cold wash buffer (20 mM HEPES pH 7.5; 150 mM NaCl; 0.5 mM

Spermidine; 1 tablet of Roche complete EDTA-free protease inhibitor). Concanavalin A-coated

beads (Bangs Laboratories) were activated by resuspension and washed in binding buffer (20

mM HEPES-KOH at pH 7.9; 10 mM KCl, 1 mM $CaCl_2$; 1 mM $MnCl_2$). Cells were mixed with

activated beads at a ratio of 9:1 at room temperature for 10 minutes. The cell and bead mixtures

were resuspended in antibody solution (2 mM EDTA; 0.05% digitonin in wash buffer).

Corresponding antibodies were added at desired concentrations to 150 µl of cell and bead

mixture and rotated overnight at 4ºC. Libraries were prepared using the AutoCUT&RUN

protocol (https://www.protocols.io/view/autocut-run-genome-wide-profiling-of-chromatin-pro-

6qpvre6zblmk/v1). High-throughput sequencing with 50 bp paired-end reads was performed on

CUT&RUN libraries using an Illumina NextSeq.

Single-cell multiome RNA-seq and ATAC-seq

Cells were collected 14 days post transduction and processed per manufacturer's instructions

using the 10x Chromium Next GEM Single Cell Multiome ATAC + Gene Expression platform to

generate libraries. Single-cell RNA-seq libraries and single-cell ATAC-seq libraries were

sequenced on an Illumina NovaSeq 6000 targeting 35,000 read pairs/nuclei for RNA-seq and

40,000 read pairs/nucleus for ATAC-seq.

Single-cell RNA-seq and ATAC-seq analysis

Single-cell multiome ATAC- and RNA-sequencing reads were processed through the

CellRanger ARC analysis pipeline. RNA-sequencing filtered feature count matrices and ATAC-

sequencing fragment paths from each sample were loaded individually loaded into R with

Signac (version 1.7.0)[72]. The data was filtered for total RNA count between 1000 and 50000,

total feature RNA count between 500 and 10000, percent mitochondria RNA less than 20%,

total ATAC fragment count between 1000 and 50000, nucleosome signal score less than 2, and

a TSS enrichment score above 1.5. ATACseq peaks were called using MACS2 [73]. Peaks on

nonstandard chromosomes and in genomic blacklist regions of hg38 were removed. ATAC

counts were quantified for all peaks across all samples.

Single-cell trajectory analysis

Filtered Seurat object was loaded into Monocle3 (version 1.2.9) to calculate Single-cell

trajectory [74,75]. First 50 PCA dimensions were used in the preprocessing step. Cells were

aligned by regressing on percent mitochondria RNA.

CUT&RUN data processing and analyses

Peak calls for H3K4me1, H3K4me3 and H3K27ac from CUT&RUN data at different timepoints

were performed using SEACR (v1.3) [76]. Bedgraph files were prepared using bedtools (v2.27.1)

to serve as the inputs for SEACR. IgG control conditions at each timepoint were used to define

empirical thresholds for peak calling. The "norm" parameter was used in SEACR to normalize

the data. Peaks were called in "stringent" mode. Active promoters were defined by checking if

H3K4me3 peaks overlap with H3K27ac signals using the bedtools intersect -wa option. The

resulting regions within +/- 5 kb of transcription start sites (TSS) were retained. TSS regions

were calculated by applying GTFtools (v0.8.5) [77] to the gencode v39 (GRCh38.p13) GTF file.

Active enhancers were obtained by overlapping H3K4me1 peaks with H3K27ac peaks, followed

by subtracting the +/- 5 kb TSS regions using the bedtools subtract –A option. Super-enhancers

were called using ROSE (v0.1) [52,78] on H3K27ac CUT&RUN data. STITCHING_DISTANCE was

set to 12.5 kb. TSS_EXCLUSION_ZONE_SIZE was set to 2.5 kb. hg38 was used as the

reference genome. The resulting genomic regions of active promoters, active enhancers and

super-enhancers were loaded into DiffBind [79] for differential binding analysis. Counts that

mapped to the genomic regions were normalized using the Relative Log Expression (RLE)

method. For super-enhancers, the counts of member regions that belong to each parental broad

genomic region were aggregated together to form the overall count. The correlations between

time and genomic regions were calculated using Kendall test followed by Benjamini-Hochberg

P-value adjustment. Genomic regions were mapped to genes using ChIPseeker (v1.32.0) [80].

Genomic regions and genes demonstrating significant changes along with time were visualized

in heatmap format using ComplexHeatmaps (v2.12.0)[69] and analyzed by Gene Ontology using

hypergeometric test with MSigDB gene sets [81].

AR binding site analyses

Inducible, constitutively active, and inactive AR binding sites (ARBS) were obtained from a

previously published study [53]. A random set of genomic regions with the same length of ARBS

(700 bp) was generated by randomly selecting 50 start sites for each chromosome. Fragments

of scATAC-seq data were quantified against these sites. The significance of the difference

between conditions was calculated based on the log transformed counts using Welch's t-test.

AR enhancer analyses

The genomic coordinates of the somatically acquired AR enhancer were obtained from a previously published study [5]. scATAC-seq data were examined at the same overall genomic region to determine the location of the AR enhancer in our model system and in the prostate cancer hepatic metastasis. The correlation between scATAC chromatin accessibility and scRNA gene expression in reprogrammed cells was visualized using CoveragePlot in the R package Signac (version 1.7.0) [72].

Statistical analyses

Data analysis was performed on GraphPad Prism 9 (GraphPad Software, Inc.). Quantitative PCR results were analyzed in Excel. Statistical significance was determined using the unpaired two-tailed Student t test, unpaired two-tailed Welch t test where the variances are shown to be different via F-test, one-way ANOVA, or two-way ANOVA. Only two-tailed tests were used. Results are depicted as mean + SD unless stated otherwise. All P values of <0.05, <0.01, <0.001, and <0.0001 were considered significant. Pearson correlation coefficient was used to determine correlation between genes. The symbols used to represent the P values were: ns, nonsignificant for $P > 0.05$; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$; ****, $P \leq 0.0001$. The test used in each statistical analysis is specified in the figure legends.

ACKNOWLEDGEMENTS

We would like to thank the Fred Hutch Genomics Shared Resource, Comparative Medicine

Shared Resource, and Experimental Histopathology Shared Resource (supported by NIH/NCI

Cancer Center Support Grant P30CA015704). We also acknowledge support by the Pacific

Northwest Prostate Cancer SPORE (NCI P50CA97186) and the Institute for Prostate Cancer

Research. SL was supported by a Department of Defense Prostate Cancer Research Program

Early Investigator Research Award (W81XWH-20-1-0083). The work was also supported by NIH

DP2 CA271301 (J.K.L.); NCI R01 CA23715 (J.K.L., M.C.H., P.S.N.); NCI R01 CA266452 (J.K.L.,

P.S.N., M.C.H.); NCI R01 CA222877 (T.G.G.); NCI P50CA092131 (T.G.G., J.K.L.); NCI

P01CA163227 (P.S.N., H.W.L.); the W.M. Keck Foundation (T.G.G.); the V Foundation (M.C.H.);

the UCLA Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research

(T.G.G.); Hal Gaba Director's Fund for Cancer Stem Cell Research (T.G.G.); and Prostate

Cancer Foundation Challenge Awards (J.K.L., P.S.N.).

AUTHOR CONTRIBUTIONS

Conceptualization, S.L., K.S., H.S., T.G.G., and J.K.L.; Methodology, S.L. and J.K.L.;

Investigation, S.L., K.S., H.S., Y.T., A.H., V.B., B.H., R.A.P., H.W.L., C.M., M.C.H.; Writing –

Original Draft, S.L. and J.K.L.; Writing – Review & Editing, P.S.N., T.G.G., and J.K.L.;

Resources, C.M., P.S.N.; Supervision, T.G.G. and J.K.L.

DECLARATION OF INTERESTS

J.K.L. has served as a consultant for Hierax Therapeutics and has equity in, an invention

licensed to, and a sponsored research agreement with PromiCell Therapeutics. T.G.G. reports

receiving an honorarium from Amgen, having consulting and equity agreements with Auron

Therapeutics, Boundless Bio Coherus BioSciences and Trethera Corporation. The lab of T.G.G.

has completed a research agreement with ImmunoActiva.

FIGURE LEGENDS

Figure 1. Direct reprogramming of ARPC to NEPC with functional bypass of a

dependence on AR signaling. (**A**) Experimental schema of the reprogramming of ARPC cells

to NEPC through the introduction of a candidate factor lentiviral (LV) pool and propagation in

neural stem cell media (N-SCM) permissive to NEPC growth. (**B**) Immunoblot analysis of the

C4-2B and LNCaP cell lines transduced with either green fluorescent protein (GFP) or the LV

pool and subjected to the reprogramming assay. Lysates were collected on day 14. (**C**)

Immunocytochemical analysis of the C4-2B and LNCaP lines at day 14. (**D**) Experimental

schema for the stringent negative selection of AR$^+$ cell populations based on the AR-dependent

expression of an inducible FKBP-Casp8 fusion protein that dimerizes and induces caspase

activation and apoptosis in the presence of the FK506 analog AP20817. (**E**) Relative cell

viability determined by CellTiter-Glo assay of treated versus untreated groups after five days

(n=4 each). P-value was assessed by Student's t-test. **** denotes p <0.0001.

Figure 2. The pioneer neural transcription factor ASCL1 suppresses AR expression and

drives NE transdifferentiation in prostate cancer. Immunoblot analysis of (**A**) leave-one-out

conditions and (**B**) factor reconstitution conditions in reprogramming studies using the C4-2B

cell line. PRNB represents dominant-negative TP53 H175R, shRB1, MYCN, and BCL2. Lysates

were collected on day 14. (**C**) Relative cell viability over time determined by CellTiter-Glo assay

of C4-2B cells reprogrammed with various factor combinations (n=4 per condition). * denotes p

<0.05, ** denotes p <0.01.

Figure 3. ASCL1 and NeuroD1 are competent to induce NE lineage reprogramming of

prostate cancer. (A) Immunoblot analysis of C4-2B and LNCaP cells reprogrammed with GFP

or PRNB, SRRM4, and ASCL1 and/or NeuroD1. Lysates were collected on day 14. (**B**)

Heatmap of RNA-seq gene expression from reprogrammed C4-2B cell line conditions showing

22-gene AR and NE signature scores and select genes associated with the AR program and NE

programs (NEURO I and NEURO II). UQ: upper quartile normalization. (**C**) Partial least

squares-discriminant analysis (PLS-DA) plot based on RNA-seq gene expression of

reprogrammed C4-2B cell line conditions (black shapes) projected onto human ARPC (red dots)

and NEPC (blue dots) samples from Beltran et al., 2016. Ellipses represent 95% confidence

level for multivariate t-distributions defined by ARPC (red) and NEPC (blue) data.

Figure 4. Dynamic changes in cancer phenotype and transcriptional and epigenetic

landscapes during acute reprogramming of ARPC to NEPC. (**A**) Experimental schema for

the temporal investigation of the reprogramming of ARPC to NEPC by immunoblot, RNA-seq,

and CUT&RUN analyses. (**B**) Immunoblot analysis of C4-2B cells modified with PRNB and

SRRM4 (PRNBS), PRNBS and NeuroD1 (PRNBSN), or PRNBS and ASCL1 (PRNBSA) that

were collected at various timepoints during the 14-day reprogramming period. Heatmaps of

RNA-seq gene expression from C4-2B cells reprogrammed with (**C**) PRNBSA or (**D**) PRNBSN

over the 14-day reprogramming period showing 22-gene NE signature scores and select genes

associated with the AR program and NE programs (NEURO I and NEURO II). (**E**) PLS-DA plot

based on RNA-seq gene expression of C4-2B cells reprogrammed with PRNBSA (top) or

PRNBSN (bottom) at different timepoints (black shapes) projected onto human ARPC (red dots)

and NEPC (blue dots) samples from Beltran et al., 2016. Ellipses represent 95% confidence of

t-distribution defined by ARPC (red) and NEPC (blue) data. (**F**) Principal component analysis

(PCA) of CUT&RUN enrichment for H3K4me3, H3K4me1 and H3K27Ac signal in C4-2B cells

reprogrammed with PRNBSA over time.

Figure 5. Time-dependent changes in ASCL1-regulated gene programs and super-

enhancer organization during NE transdifferentiation of prostate cancer. (**A**) Heatmap of

MSigDB gene sets enriched over time with ASCL1-regulated genes identified by integrating

CUT&RUN and RNA-seq data from C4-2B cells reprogrammed with PRNBSA over the 14-day

reprogramming period. (**B**) Heatmap showing dynamic changes in super-enhancer regions in

C4-2B cells reprogrammed with PRNBSA over the 14-day reprogramming period. Genomic

regions shown were obtained by setting a p-value cutoff of 1e-03 on Kendall correlation

between day and peak activity. RLE: Relative Log Expression normalization. (**C**) Plot showing

Gene Ontology (GO) enrichment of Hallmark gene sets associated with decreasing super

enhancer signals from D2 to D14 in C4-2B cells reprogrammed with PRNBSA.

Figure 6. ASCL1/NeuroD1 inhibit AR expression by remodeling chromatin accessibility at

the somatically acquired AR enhancer and global AR binding sites with enhancer activity.

(**A**) Tracks showing chromatin accessibility peaks at the AR locus and AR gene expression

(violin plot) from pseudo-bulk analysis of scATAC-seq and scRNA-seq data from C4-2B cells

reprogrammed with PRNB, PRNBS, PRNBSA, and PRNBSN. Tracks in the region of the

somatically acquired AR enhancer are magnified. (**B**) Immunofluorescence analysis of a human

prostate cancer hepatic metastasis with ASCL1 staining (top), AR staining (middle), and an

overlay of ASCL1 and AR staining (bottom) shown. DAPI was used as a nuclear counterstain.

(**C**) Tracks showing chromatin accessibility at the AR locus by pseudo-bulk analysis of scATAC-

seq data from the $AR^-/ASCL1^+$ and $AR^+/ASCL1^-$ tumor cell populations present in the human

prostate cancer hepatic metastasis shown in **B**. Tracks in the region of the somatically acquired

AR enhancer are magnified. (**D**) Schematic representation of different classes of AR enhancers

(top) is shown. The distribution of chromatin accessibility at AR bindings sites (ARBS) of

different enhancer activity and at random genomic regions of the same interval are shown.

Black dots represent the means. ns denotes $p > 0.05$, ** denotes $p < 0.01$, *** denotes $p < 0.001$.

Figure 7. ASCL1/NeuroD1 induce the expression and activity of NE-associated

transcription factors. (**A**) Volcano plots showing the differential expression of genes encoding

transcriptions factors in pairwise comparisons of RNA-seq gene expression data of C4-2B cells

reprogrammed with PRNBS vs. PRNB, PRNBSA vs. PRNBS, and PRNBSN vs. PRNBS. Red

dots represent genes with a -$\log_{10}$(FDR) > 2. FC represents fold-change. (**B**) Plots of differential

transcription factor motif activity from footprinting analysis of pseudo-bulk scATAC-seq data

from C4-2B cells reprogrammed with PRNBS vs. PRNB, PRNBSA vs. PRNBS, and PRNBSN vs.

PRNBS. Note that labelled motifs in the second and third panel do not show up in the first panel.

(**C**) PCA of chromatin accessibility signals from pseudo-bulk scATAC-seq of reprogrammed C4-

2B cell line conditions projected on previously published ATAC-seq data from NEPC and ARPC

PDX models. (**D**) Heatmap of RNA-seq gene expression data showing cell surface targets

relevant to ARPC and NEPC in reprogrammed C4-2B cell conditions.

Figure 8. Downregulation of MHC class I antigen processing and presentation genes in

NEPC is functionally attributable to ASCL1/NeuroD1. (**A**) Heatmap of RNA-seq gene

expression data showing the NE signature and MHC class I pathway scores (top) and select

genes (bottom) in ARPC and NEPC from Beltran et al., 2016. (**B**) Heatmap of RNA-seq gene

expression data showing the AR and NE signature scores and select genes in AR$^+$/NE$^-$ and AR$^-$

/NE$^+$ metastatic prostate cancers from the University of Washington Tissue Acquisition

Necropsy (UW TAN) program. (**C**) Heatmap of RNA-seq gene expression data from

reprogrammed C4-2B cell line conditions showing the NE signature and MHC class I pathway

scores (top) and select MHC class I genes including B2M (bottom). UQ: upper quartile

normalization. (**D**) UMAP analysis of scRNA-seq and scATAC-seq data from C4-2B cells

reprogrammed with PRNB, PRNBS, PRNBSA or PRNBSN using the Seurat package. UMAP

plots of reprogrammed cells colored based on (**E**) NEPC score and (**F**) B2M gene expression.

(**G**) Flow cytometry histogram plots showing the percentage of B2M positive cells in

reprogrammed C4-2B cells.

REFERENCES

1. Chen, C.D., Welsbie, D.S., Tran, C., Baek, S.H., Chen, R., Vessella, R., Rosenfeld, M.G., and Sawyers, C.L. (2004). Molecular determinants of resistance to antiandrogen therapy. Nature Medicine *10*, 33-39. 10.1038/nm972.

2. Watson, P.A., Arora, V.K., and Sawyers, C.L. (2015). Emerging mechanisms of resistance to androgen receptor inhibitors in prostate cancer. Nat Rev Cancer *15*, 701-711. 10.1038/nrc4016.

3. Hu, R., Dunn, T.A., Wei, S.Z., Isharwal, S., Veltri, R.W., Humphreys, E., Han, M., Partin, A.W., Vessella, R.L., Isaacs, W.B., et al. (2009). Ligand-Independent Androgen Receptor Variants Derived from Splicing of Cryptic Exons Signify Hormone-Refractory Prostate Cancer. Cancer Research *69*, 16-22. 10.1158/0008-5472.Can-08-2764.

4. Viswanathan, S.R., Ha, G., Hoff, A.M., Wala, J.A., Carrot-Zhang, J., Whelan, C.W., Haradhvala, N.J., Freeman, S.S., Reed, S.C., Rhoades, J., et al. (2018). Structural Alterations Driving Castration-Resistant Prostate Cancer Revealed by Linked-Read Genome Sequencing. Cell *174*, 433-+. 10.1016/j.cell.2018.05.036.

5. Takeda, D.Y., Spisak, S., Seo, J.H., Bell, C., O'Connor, E., Korthauer, K., Ribli, D., Csabai, I., Solymosi, N., Szallasi, Z., et al. (2018). A Somatically Acquired Enhancer of the Androgen Receptor Is a Noncoding Driver in Advanced Prostate Cancer. Cell *174*, 422-432 e413. 10.1016/j.cell.2018.05.037.

6. Beltran, H., Rickman, D.S., Park, K., Chae, S.S., Sboner, A., MacDonald, T.Y., Wang, Y., Sheikh, K.L., Terry, S., Tagawa, S.T., et al. (2011). Molecular characterization of neuroendocrine prostate cancer and identification of new drug targets. Cancer discovery *1*, 487-495. 10.1158/2159-8290.Cd-11-0130.

7. Beltran, H., Hruszkewycz, A., Scher, H.I., Hildesheim, J., Isaacs, J., Yu, E.Y., Kelly, K., Lin, D., Dicker, A., Arnold, J., et al. (2019). The Role of Lineage Plasticity in Prostate Cancer Therapy Resistance. Clin Cancer Res *25*, 6916-6924. 10.1158/1078-0432.Ccr-19-1423.

8. Lee, J.K., Phillips, J.W., Smith, B.A., Park, J.W., Stoyanova, T., McCaffrey, E.F., Baertsch, R., Sokolov, A., Meyerowitz, J.G., Mathis, C., et al. (2016). N-Myc Drives Neuroendocrine Prostate Cancer Initiated from Human Prostate Epithelial Cells. Cancer cell *29*, 536-547. 10.1016/j.ccell.2016.03.001.

9. Park, J.W., Lee, J.K., Sheu, K.M., Wang, L., Balanis, N.G., Nguyen, K., Smith, B.A., Cheng, C., Tsai, B.L., Cheng, D., et al. (2018). Reprogramming normal human epithelial tissues to a common, lethal neuroendocrine cancer lineage. Science (New York, N.Y.) *362*, 91-95. 10.1126/science.aat5749.

10. Nyquist, M.D., Corella, A., Coleman, I., De Sarkar, N., Kaipainen, A., Ha, G., Gulati, R.,
Ang, L., Chatterjee, P., Lucas, J., et al. (2020). Combined TP53 and RB1 Loss Promotes
Prostate Cancer Resistance to a Spectrum of Therapeutics and Confers Vulnerability to
Replication Stress. Cell Rep *31*. ARTN 107669
10.1016/j.celrep.2020.107669.

11. Berger, A., Brady, N.J., Bareja, R., Robinson, B., Conteduca, V., Augello, M.A., Ruca, L.,
Ahmed, A., Dardenne, E., Lu, X.D., et al. (2019). N-Myc mediated epigenetic
reprogramming drives lineage plasticity in advanced prostate cancer. J Clin Invest *129*,
3924-3940. 10.1172/Jci127961.

12. Zhang, X.T., Coleman, I.M., Brown, L.G., True, L.D., Kollath, L., Lucas, J.M., Lam, H.M.,
Dumpit, R., Corey, E., Chery, L., et al. (2015). SRRM4 Expression and the Loss of
REST Activity May Promote the Emergence of the Neuroendocrine Phenotype in
Castration-Resistant Prostate Cancer. Clin Cancer Res *21*, 4698-4708. 10.1158/1078-
0432.Ccr-15-0157.

13. Mu, P., Zhang, Z., Benelli, M., Karthaus, W.R., Hoover, E., Chen, C.C., Wongvipat, J.,
Ku, S.Y., Gao, D., Cao, Z., et al. (2017). SOX2 promotes lineage plasticity and
antiandrogen resistance in TP53- and RB1-deficient prostate cancer. Science (New York,
N.Y.) *355*, 84-88. 10.1126/science.aah4307.

14. Lin, D., Wyatt, A.W., Xue, H., Wang, Y., Dong, X., Haegert, A., Wu, R., Brahmbhatt, S.,
Mo, F., Jong, L., et al. (2014). High fidelity patient-derived xenografts for accelerating
prostate cancer discovery and drug development. Cancer Res *74*, 1272-1283.
10.1158/0008-5472.CAN-13-2921-T.

15. Akamatsu, S., Wyatt, Alexander W., Lin, D., Lysakowski, S., Zhang, F., Kim, S., Tse, C.,
Wang, K., Mo, F., Haegert, A., et al. (2015). The Placental Gene PEG10 Promotes
Progression of Neuroendocrine Prostate Cancer. Cell reports *12*, 922-936.
https://doi.org/10.1016/j.celrep.2015.07.012.

16. Epstein, J.I., Amin, M.B., Beltran, H., Lotan, T.L., Mosquera, J.M., Reuter, V.E.,
Robinson, B.D., Troncoso, P., and Rubin, M.A. (2014). Proposed Morphologic
Classification of Prostate Cancer With Neuroendocrine Differentiation. Am J Surg Pathol
38, 756-767. 10.1097/Pas.0000000000000208.

17. Balanis, N.G., Sheu, K.M., Esedebe, F.N., Patel, S.J., Smith, B.A., Park, J.W., Alhani, S.,
Gomperts, B.N., Huang, J., Witte, O.N., and Graeber, T.G. (2019). Pan-cancer
Convergence to a Small-Cell Neuroendocrine Phenotype that Shares Susceptibilities
with Hematological Malignancies. Cancer Cell *36*, 17-+. 10.1016/j.ccell.2019.06.005.

18. Labrecque, M.P., Coleman, I.M., Brown, L.G., True, L.D., Kollath, L., Lakely, B., Nguyen,
H.M., Yang, Y.C., da Costa, R.M.G., Kaipainen, A., et al. (2019). Molecular profiling

stratifies diverse phenotypes of treatment-refractory metastatic castration-resistant
prostate cancer. J Clin Invest *129*, 4492-4505. 10.1172/Jci128212.

19.   Merkens, L., Sailer, V., Lessel, D., Janzen, E., Greimeier, S., Kirfel, J., Perner, S., Pantel,
K., Werner, S., and von Amsberg, G. (2022). Aggressive variants of prostate cancer:
underlying mechanisms of neuroendocrine transdifferentiation. J Exp Clin Canc Res *41*.
ARTN 46
10.1186/s13046-022-02255-y.

20.   Raposo, A.A.S.F., Vasconcelos, F.F., Drechsel, D., Marie, C., Johnston, C., Dolle, D.,
Bithell, A., Gillotin, S., van den Berg, D.L.C., Ettwiller, L., et al. (2015). Ascl1
Coordinately Regulates Gene Expression and the Chromatin Landscape during
Neurogenesis. Cell Rep *10*, 1544-1556. 10.1016/j.celrep.2015.02.025.

21.   Borromeo, M.D., Savage, T.K., Kollipara, R.K., He, M., Augustyn, A., Osborne, J.K.,
Girard, L., Minna, J.D., Gazdar, A.F., Cobb, M.H., and Johnson, J.E. (2016). ASCL1 and
NEUROD1 Reveal Heterogeneity in Pulmonary Neuroendocrine Tumors and Regulate
Distinct Genetic Programs. Cell Rep *16*, 1259-1272. 10.1016/j.celrep.2016.06.081.

22.   Cejas, P., Xie, Y.T., Font-Tello, A., Lim, K., Syamala, S., Qiu, X.T., Tewari, A.K., Shah,
N., Nguyen, H.M., Patel, R.A., et al. (2021). Subtype heterogeneity and epigenetic
convergence in neuroendocrine prostate cancer. Nat Commun *12*. ARTN 5775
10.1038/s41467-021-26042-z.

23.   Costanzo, F., Diez, M.M., Nunez, G.S., Diaz-Hernandez, J.I., Robles, C.M.G., Perez,
J.D., Compe, E., Ricci, R., Li, T.K., Coin, F., et al. (2022). Promoters of ASCL1-and
NEUROD1-dependent genes are specific targets of lurbinectedin in SCLC cells. Embo
Mol Med *14*. ARTN e14841
10.15252/emmm.202114841.

24.   Baine, M.K., Hsieh, M.S., Lai, W.V., Egger, J.V., Jungbluth, A.A., Daneshbod, Y., Beras,
A., Spencer, R., Lopardo, J., Bodd, F., et al. (2020). SCLC Subtypes Defined by ASCL1,
NEUROD1, POU2F3, and YAP1: A Comprehensive Immunohistochemical and
Histopathologic Characterization. J Thorac Oncol *15*, 1823-1835.
10.1016/j.jtho.2020.09.009.

25.   Szeitz, B., Megyesfalvi, Z., Woldmar, N., Valko, Z., Schwendenwein, A., Barany, N.,
Paku, S., Laszlo, V., Kiss, H., Bugyik, E., et al. (2022). In-depth proteomic analysis
reveals unique subtype-specific signatures in human small-cell lung cancer. Clin Transl
Med *12*. ARTN e1060
10.1002/ctm2.1060.

26.   DeLucia, D.C., Cardillo, T.M., Ang, L., Labrecque, M.P., Zhang, A., Hopkins, J.E., De
Sarkar, N., Coleman, I., da Costa, R.M.G., Corey, E., et al. (2021). Regulation of

CEACAM5 and Therapeutic Efficacy of an Anti-CEACAM5-SN38 Antibody-drug
Conjugate in Neuroendocrine Prostate Cancer. Clin Cancer Res *27*, 759-774.
10.1158/1078-0432.CCR-20-3396.

27.   Augustyn, A., Borromeo, M., Wang, T., Fujimoto, J., Shao, C., Dospoy, P.D., Lee, V.,
Tan, C., Sullivan, J.P., Larsen, J.E., et al. (2014). ASCL1 is a lineage oncogene
providing therapeutic targets for high-grade neuroendocrine lung cancers. Proceedings
of the National Academy of Sciences of the United States of America *111*, 14788-14793.
10.1073/pnas.1410419111.

28.   Nouruzi, S., Ganguli, D., Tabrizian, N., Kobelev, M., Sivak, O., Namekawa, T., Thaper,
D., Baca, S.C., Freedman, M.L., Aguda, A., et al. (2022). ASCL1 activates neuronal
stem cell-like lineage programming through remodeling of the chromatin landscape in
prostate cancer. Nat Commun *13*. ARTN 2282
10.1038/s41467-022-29963-5.

29.   Ku, S.Y., Rosario, S., Wang, Y., Mu, P., Seshadri, M., Goodrich, Z.W., Goodrich, M.M.,
Labbe, D.P., Gomez, E.C., Wang, J., et al. (2017). Rb1 and Trp53 cooperate to
suppress prostate cancer lineage plasticity, metastasis, and antiandrogen resistance.
Science (New York, N.Y.) *355*, 78-83. 10.1126/science.aah4199.

30.   Dardenne, E., Beltran, H., Benelli, M., Gayvert, K., Berger, A., Puca, L., Cyrta, J.,
Sboner, A., Noorzad, Z., MacDonald, T., et al. (2016). N-Myc Induces an EZH2-
Mediated Transcriptional Program Driving Neuroendocrine Prostate Cancer. Cancer cell
30, 563-577. 10.1016/j.ccell.2016.09.005.

31.   Li, Y.A., Donmez, N., Sahinalp, C., Xie, N., Wang, Y.W., Xue, H., Mo, F., Beltran, H.,
Gleave, M., Wang, Y.Z., et al. (2017). SRRM4 Drives Neuroendocrine
Transdifferentiation of Prostate Adenocarcinoma Under Androgen Receptor Pathway
Inhibition. Eur Urol *71*, 68-78. 10.1016/j.eururo.2016.04.028.

32.   Gobinet, J., Auzou, G., Nicolas, J.C., Sultan, C., and Jalaguier, S. (2001).
Characterization of the interaction between androgen receptor and a new transcriptional
inhibitor, SHP. Biochemistry *40*, 15369-15377. 10.1021/bi011384o.

33.   Babos, K.N., Galloway, K.E., Kisler, K., Zitting, M., Li, Y.C., Shi, Y.X., Quintino, B., Chow,
R.H., Zlokovic, B.V., and Ichida, J.K. (2019). Mitigating Antagonism between
Transcription and Proliferation Allows Near-Deterministic Cellular Reprogramming. Cell
Stem Cell *25*, 486-+. 10.1016/j.stem.2019.08.005.

34.   Jouravel, N., Sablin, E., Arnold, L.A., Guy, R.K., and Fletterick, R.J. (2007). Interaction
between the androgen receptor and a segment of its corepressor SHP. Acta Crystallogr
D *63*, 1198-1200. 10.1107/S0907444907045702.

35.   Judware, R., and Culp, L.A. (1995). Over-expression of transfected N-myc oncogene in
human SKNSH neuroblastoma cells down-regulates expression of beta 1 integrin
subunit. Oncogene *11*, 2599-2607.

36.   Cobrinik, D., Francis, R.O., Abramson, D.H., and Lee, T.C. (2006). Rb induces a
proliferative arrest and curtails Brn-2 expression in retinoblastoma cells. Mol Cancer *5*,
72. 10.1186/1476-4598-5-72.

37.   Li, Y.N., Chen, R.Q., Bowden, M., Mo, F., Lin, Y.Y., Gleave, M., Collins, C., and Dong,
X.S. (2017). Establishment of a neuroendocrine prostate cancer model driven by the
RNA splicing factor SRRM4. Oncotarget *8*, 66878-66888. 10.18632/oncotarget.19916.

38.   Li, S., Wong, A., Sun, H., Bhatia, V., Javier, G., Jana, S., Montgomery, R.B., Wright, J.L.,
Lam, H.M., Hsieh, A.C., et al. (2023). Combinatorial genetic strategy accelerates the
discovery of cancer genotype-phenotype associations. bioRxiv.
10.1101/2023.04.12.536652.

39.   Vue, T.Y., Kollipara, R.K., Borromeo, M.D., Smith, T., Mashimo, T., Burns, D.K., Bachoo,
R.M., and Johnson, J.E. (2020). ASCL1 regulates neurodevelopmental transcription
factors and cell cycle genes in brain tumors of glioma mouse models. Glia *68*, 2613-
2630. 10.1002/glia.23873.

40.   Han, M., Li, F., Zhang, Y., Dai, P., He, J., Li, Y., Zhu, Y., Zheng, J., Huang, H., Bai, F.,
and Gao, D. (2022). FOXA2 drives lineage plasticity and KIT pathway activation in
neuroendocrine prostate cancer. Cancer cell *40*, 1306-1323.e1308.
10.1016/j.ccell.2022.10.011.

41.   Bishop, J.L., Thaper, D., Vahid, S., Davies, A., Ketola, K., Kuruma, H., Jama, R., Nip,
K.M., Angeles, A., Johnson, F., et al. (2017). The Master Neural Transcription Factor
BRN2 Is an Androgen Receptor-Suppressed Driver of Neuroendocrine Differentiation in
Prostate Cancer. Cancer discovery *7*, 54-71. 10.1158/2159-8290.Cd-15-1263.

42.   Beltran, H., Prandi, D., Mosquera, J.M., Benelli, M., Puca, L., Cyrta, J., Marotz, C.,
Giannopoulou, E., Chakravarthi, B.V., Varambally, S., et al. (2016). Divergent clonal
evolution of castration-resistant neuroendocrine prostate cancer. Nat Med *22*, 298-305.
10.1038/nm.4045.

43.   Ohnishi, T., Shirane, M., and Nakayama, K.I. (2017). SRRM4-dependent neuron-specific
alternative splicing of protrudin transcripts regulates neurite outgrowth. Sci Rep-Uk *7*.
ARTN 41130
10.1038/srep41130.

44.   Tsoi, J., Robert, L., Paraiso, K., Galvan, C., Sheu, K.M., Lay, J., Wong, D.J.L., Atefi, M.,
Shirazi, R., Wang, X.Y., et al. (2018). Multi-stage Differentiation Defines Melanoma
Subtypes with Differential Vulnerability to Drug-Induced Iron-Dependent Oxidative Stress.
Cancer Cell *33*, 890-+. 10.1016/j.ccell.2018.03.017.

45.   Richard, A., Boullu, L., Herbach, U., Bonnafoux, A., Morin, V., Vallin, E., Guillemin, A.,
Gao, N.P., Gunawan, R., Cosette, J., et al. (2016). Single-Cell-Based Analysis Highlights
a Surge in Cell-to-Cell Molecular Variability Preceding Irreversible Commitment in a
Differentiation Process. Plos Biol *14*. ARTN e1002585
10.1371/journal.pbio.1002585.

46.   Basili, D., Zhang, J.L., Herbert, J., Krol, K., Denslow, N.D., Martyniuk, C.J., Falciani, F.,
and Antczak, P. (2018). In Silico Computational Transcriptomics Reveals Novel
Endocrine Disruptors in Largemouth Bass (Micropterus salmoides). Environ Sci Technol
52, 7553-7565. 10.1021/acs.est.8b02805.

47.   Jiang, S., Williams, K., Kong, X.D., Zeng, W.H., Nguyen, N.V., Ma, X.Y., Tawil, R.,
Yokomori, K., and Mortazavi, A. (2020). Single-nucleus RNA-seq identifies divergent
populations of FSHD2 myotube nuclei. Plos Genet *16*. ARTN e1008754
10.1371/journal.pgen.1008754.

48.   Karanikolas, B.D.W., Figueiredo, M.L., and Wu, L. (2010). Comprehensive Evaluation of
the Role of EZH2 in the Growth, Invasion, and Aggression of a Panel of Prostate Cancer
Cell Lines. Prostate *70*, 675-688. 10.1002/pros.21112.

49.   Cyrta, J., Augspach, A., De Filippo, M.R., Prandi, D., Thienger, P., Benelli, M., Cooley,
V., Bareja, R., Wilkes, D., Chae, S.S., et al. (2020). Role of specialized composition of
SWI/SNF complexes in prostate cancer lineage plasticity. Nat Commun *11*. ARTN 5549
10.1038/s41467-020-19328-1.

50.   Crispatzu, G., Rehimi, R., Pachano, T., Bleckwehl, T., Cruz-Molina, S., Xiao, C., Mahabir,
E., Bazzi, H., and Rada-Iglesias, A. (2021). The chromatin, topological and regulatory
properties of pluripotency-associated poised enhancers are conserved in vivo. Nat
Commun *12*. ARTN 4344
10.1038/s41467-021-24641-4.

51.   Stepniak, K., Machnicka, M.A., Mieczkowski, J., Macioszek, A., Wojtas, B., Gielniewski,
B., Poleszak, K., Perycz, M., Krol, S.K., Guzik, R., et al. (2021). Mapping chromatin
accessibility and active regulatory elements reveals pathological mechanisms in human
gliomas (vol 12, 3621, 2021). Nat Commun *12*. ARTN 7218
10.1038/s41467-021-27448-5.

52.   Whyte, W.A., Orlando, D.A., Hnisz, D., Abraham, B.J., Lin, C.Y., Kagey, M.H., Rahl, P.B.,
Lee, T.I., and Young, R.A. (2013). Master Transcription Factors and Mediator Establish

Super-Enhancers at Key Cell Identity Genes. Cell *153*, 307-319.
10.1016/j.cell.2013.03.035.

53.   Huang, C.C.F., Lingadahalli, S., Morova, T., Ozturan, D., Hu, E., Yu, I.P.L., Linder, S.,
Hoogstraat, M., Stelloo, S., Sar, F., et al. (2021). Functional mapping of androgen
receptor enhancer activity. Genome Biol *22*. ARTN 149
10.1186/s13059-021-02339-6.

54.   Saint-Andre, V., Federation, A.J., Lin, C.Y., Abraham, B.J., Reddy, J., Lee, T.I., Bradner,
J.E., and Young, R.A. (2016). Models of human core transcriptional regulatory circuitries.
Genome Res *26*, 385-396. 10.1101/gr.197590.115.

55.   Dhatchinamoorthy, K., Colbert, J.D., and Rock, K.L. (2021). Cancer Immune Evasion
Through Loss of MHC Class I Antigen Presentation. Front Immunol *12*. ARTN 636568
10.3389/fimmu.2021.636568.

56.   Cai, L., Liu, H.Y., Huang, F., Fujimoto, J., Girard, L., Chen, J., Li, Y.W., Zhang, Y.A., Deb,
D., Stastny, V., et al. (2021). Cell-autonomous immune gene expression is repressed in
pulmonary neuroendocrine cells and small cell lung cancer. Commun Biol *4*. ARTN 314
10.1038/s42003-021-01842-7.

57.   Burr, M.L., Sparbier, C.E., Chan, K.L., Chan, Y.C., Kersbergen, A., Lam, E.Y.N., Azidis-
Yates, E., Vassiliadis, D., Bell, C.C., Gilan, O., et al. (2019). An Evolutionarily Conserved
Function of Polycomb Silences the MHC Class I Antigen Presentation Pathway and
Enables Immune Evasion in Cancer. Cancer Cell *36*, 385-+. 10.1016/j.ccell.2019.08.008.

58.   Bluemn, E.G., Coleman, I.M., Lucas, J.M., Coleman, R.T., Hernandez-Lopez, S.,
Tharakan, R., Bianchi-Frias, D., Dumpit, R.F., Kaipainen, A., Corella, A.N., et al. (2017).
Androgen Receptor Pathway-Independent Prostate Cancer Is Sustained through FGF
Signaling. Cancer Cell *32*, 474-+. 10.1016/j.ccell.2017.09.003.

59.   Sequist, L.V., Waltman, B.A., Dias-Santagata, D., Digumarthy, S., Turke, A.B., Fidias, P.,
Bergethon, K., Shaw, A.T., Gettinger, S., Cosper, A.K., et al. (2011). Genotypic and
Histological Evolution of Lung Cancers Acquiring Resistance to EGFR Inhibitors. Sci
Transl Med *3*. ARTN 75ra26
10.1126/scitranslmed.3002003.

60.   Wang, Z.W., Wang, T., Hong, D.N., Dong, B.J., Wang, Y., Huang, H.Q., Zhang, W.H.,
Lian, B.J., Ji, B.Y., Shi, H.Q., et al. (2022). Single-cell transcriptional regulation and
genetic evolution of neuroendocrine prostate cancer. Iscience *25*. ARTN 104576
10.1016/j.isci.2022.104576.

61.   Brady, N.J., Bagadion, A.M., Singh, R., Conteduca, V., Van Emmenis, L., Arceci, E.,
Pakula, H., Carelli, R., Khani, F., Bakht, M., et al. (2021). Temporal evolution of cellular
heterogeneity during the progression to advanced AR-negative prostate cancer. Nature
communications *12*, 3372. 10.1038/s41467-021-23780-y.

62.   Abida, W., Cheng, M.L., Armenia, J., Middha, S., Autio, K.A., Vargas, H.A., Rathkopf, D.,
Morris, M.J., Danila, D.C., Slovin, S.F., et al. (2019). Analysis of the Prevalence of
Microsatellite Instability in Prostate Cancer and Response to Immune Checkpoint
Blockade. JAMA oncology *5*, 471-478. 10.1001/jamaoncol.2018.5801.

63.   Hiatt, J.B., Sandborg, H., Garrison, S.M., Arnold, H.U., Liao, S.Y., Norton, J.P., Friesen,
T.J., Wu, F.N., Sutherland, K.D., Rienhoff, H.Y., et al. (2022). Inhibition of LSD1 with
Bomedemstat Sensitizes Small Cell Lung Cancer to Immune Checkpoint Blockade and
T-Cell Killing. Clin Cancer Res *28*, 4551-4564. 10.1158/1078-0432.Ccr-22-1128.

64.   Nguyen, E.M., Taniguchi, H., Chan, J.M., Zhan, Y.Q.A., Chen, X.P., Qiu, J., de
Stanchina, E., Allaj, V., Shah, N.S., Uddin, F., et al. (2022). Targeting Lysine-Specific
Demethylase 1 Rescues Major Histocompatibility Complex Class I Antigen Presentation
and Overcomes Programmed Death- Ligand 1 Blockade Resistance in SCLC. J Thorac
Oncol *17*, 1014-1031. 10.1016/j.jtho.2022.05.014.

65.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal *17*, 10-12.

66.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics (Oxford, England) *29*, 15-21. 10.1093/bioinformatics/bts635.

67.   Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq
data with or without a reference genome. Bmc Bioinformatics *12*. Artn 323
10.1186/1471-2105-12-323.

68.   Hanzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis
for microarray and RNA-Seq data. Bmc Bioinformatics *14*. Artn 7
10.1186/1471-2105-14-7.

69.   Gu, Z.G., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. Bioinformatics *32*, 2847-2849.
10.1093/bioinformatics/btw313.

70.   Shen, S.H., Park, J.W., Lu, Z.X., Lin, L., Henry, M.D., Wu, Y.N., Zhou, Q., and Xing, Y.
(2014). rMATS: Robust and flexible detection of differential alternative splicing from

1207   replicate RNA-Seq data. P Natl Acad Sci USA *111*, E5593-E5601.
1208   10.1073/pnas.1419161111.

1209   71.   Howe, K.L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean,
1210   I.M., Azov, A.G., Bennett, R., Bhai, J., et al. (2021). Ensembl 2021. Nucleic Acids Res
1211   *49*, D884-D891. 10.1093/nar/gkaa942.

1212   72.   Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2021). Single-cell
1213   chromatin state analysis with Signac. Nat Methods *18*, 1333-+. 10.1038/s41592-021-
1214   01282-5.

1215   73.   Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
1216   Nussbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based
1217   Analysis of ChIP-Seq (MACS). Genome Biol *9*. ARTN R137
1218   10.1186/gb-2008-9-9-r137.

1219   74.   Cao, J.Y., Spielmann, M., Qiu, X.J., Huang, X.F., Ibrahim, D.M., Hill, A.J., Zhang, F.,
1220   Mundlos, S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional
1221   landscape of mammalian organogenesis. Nature *566*, 496-+. 10.1038/s41586-019-0969-
1222   x.

1223   75.   Qiu, X.J., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).
1224   Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*,
1225   979-+. 10.1038/Nmeth.4402.

1226   76.   Meers, M.P., Tenenbaum, D., and Henikoff, S. (2019). Peak calling by Sparse
1227   Enrichment Analysis for CUT&RUN chromatin profiling. Epigenet Chromatin *12*. ARTN
1228   42
1229   10.1186/s13072-019-0287-4.

1230   77.   Li, H.-D., Lin, C.-X., and Zheng, J. (2022). GTFtools: a software package for analyzing
1231   various features of gene models. Bioinformatics (Oxford, England) *38*, 4806-4808.
1232   10.1093/bioinformatics/btac561.

1233   78.   Loven, J., Hoke, H.A., Lin, C.Y., Lau, A., Orlando, D.A., Vakoc, C.R., Bradner, J.E., Lee,
1234   T.I., and Young, R.A. (2013). Selective Inhibition of Tumor Oncogenes by Disruption of
1235   Super-Enhancers. Cell *153*, 320-334. 10.1016/j.cell.2013.03.036.

1236   79.   Ross-Innes, C.S., Stark, R., Teschendorff, A.E., Holmes, K.A., Ali, H.R., Dunning, M.J.,
1237   Brown, G.D., Gojis, O., Ellis, I.O., Green, A.R., et al. (2012). Differential oestrogen
1238   receptor binding is associated with clinical outcome in breast cancer. Nature *481*, 389-
1239   U177. 10.1038/nature10730.

1240   80.   Yu, G.C., Wang, L.G., and He, Q.Y. (2015). ChIPseeker: an R/Bioconductor package for
1241   ChIP peak annotation, comparison and visualization. Bioinformatics *31*, 2382-2383.
1242   10.1093/bioinformatics/btv145.

1243   81.   Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
1244   Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene
1245   set enrichment analysis: A knowledge-based approach for interpreting genome-wide
1246   expression profiles. P Natl Acad Sci USA *102*, 15545-15550. 10.1073/pnas.0506580102.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8