Wood ants on the edge: how do the characteristics of linear edges effect the population dynamics of an edge specialist?

Jacob A. Podesta[1], Catherine L. Parr[2], Kelly R. Redeker[1] and Elva J. H. Robinson[1]

[1]Department of Biology, University of York, York, UK

[2]School of Environmental Sciences, University of Liverpool, Liverpool, UK

Abstract

Landscape structure may facilitate or inhibit the dispersal of species by presenting barriers or opportunities. Slow-dispersing edge specialists are likely to be most affected by local topography and land management practices, because they require two different adjacent habitats within their limited dispersal range. The northern hairy wood ant, *Formica lugubris,* requires adjacent forest and open habitats to access both aphid honeydew from trees, and direct sunlight for thermoregulation. In managed forests, canopy gaps that provide this combination are often linear and of anthropogenic origin, such as paths, firebreaks, and roads (collectively 'rides'), and their orientation determines shade distribution. Using population data spanning 10 years, we ask how properties of rides, such as bearing and width, affect the distribution and dispersal of wood ant populations in the North York Moors, UK. Regardless of ride orientation, nests were more abundant on the north side of rides, where they receive more sunlight, and nests were generally larger on narrower rides, which are more shaded. We found that *F. lugubris* dispersed approximately 3 times quicker along rides that were oriented north-south or east-west compared with NE-SW or NW-SE, although N-S bearings were used more symmetrically than E-W bearings, where the north side was predominantly used. Our results clearly show that ride characteristics (width, orientation), resulting from forestry practices, can influence the distribution and dispersal of an ecosystem engineer woodland species. Given that wood ants can suppress defoliating pests, these findings have potential forest management implications and may also apply to other forest-dwelling edge specialists.

Keywords

Dispersal, Habitat fragmentation, Forest edges, Social insects, Population dynamics, Fire management

Highlights

- Slow dispersing edge specialists are strongly affected by landscape structure
- We used a 10-year dataset on edge specialist wood ants to quantify these effects
- Wood ant nest size and distribution is shaped by access to insolation
- Wood ants disperse 3 times quicker on N-S and E-W rides than intermediate bearings
- The orientation of linear features can aid or hinder edge specialist dispersal.

Introduction

The ability of species to move through a landscape can have major implications for the conservation of biodiversity, and for preventing species loss in a changing climate (Mendenhall *et al.*, 2014; Lenoir and Svenning, 2015). How porous a landscape is to individual movement will also affect the dispersal of species, and this porosity may be determined by the contrast between one habitat and the next. For example, low contrast edges between forest patches and agricultural land in Central American countryside ecosystems are more porous to bats than high contrast edges between forest patches and open water, and forest patches in the agricultural ecosystem harbour higher bat species richness as a result (Mendenhall *et al.*, 2014). The permeability of habitats to species may become more important as climate change induced range shifts along latitudinal or altitudinal gradients become more apparent (Lenoir and Svenning, 2015).

Natural linear features such as rivers, coasts, habitat edges, and mountain ranges, or anthropogenic ones such as roads, railways and canals can be major facilitators of species movement, providing corridors of suitable habitat for certain species through otherwise inhospitable areas. For example, wind-dispersed seeds travel nearly four times the distance along linear forest disturbances than in undisturbed forest (Roberts *et al.*, 2018) while, in a meta-analysis, 70% of studies indicate that either plant abundance or diversity is higher along linear gaps and verges than habitat interiors (Suárez-Esteban *et al.*, 2016). By facilitating the dispersal of food plants, linear features can indirectly promote the dispersal of invertebrate herbivores. Ragworts (*Senecio* spp.) disperse along man-made linear features (e.g., railways), and ragwort-dependant cinnabar moths (*Tyria jacobaeae*) show increased dispersal and egg-laying success in areas with roads and valleys compared with areas without linear features (Brunzel et al., 2004), suggesting that invertebrates also benefit from linear features.

The creation of linear features can be a conservation tool, as in the use of habitat corridors to connect smaller areas of protected land and reserves (Bennett, 1990; Brito et al., 2017). However, habitat corridors can affect generalist and specialist species differently: landscape corridors connecting areas of fragmented forest in south-eastern Estonia facilitated the dispersal of generalist plant species more than forest specialist species (Liira and Paal, 2013). The presence and structure of edges thus has conservation potential, but the implications for species movement and survival may differ depending on the target species.

The widespread anthropogenic changes in natural forests, and the creation of large areas of plantation forests, mean that human activities determine the present distribution and fragmentation of most forests. This makes the study of how fragmentation and edge habitats impact species dispersal important for conserving species in increasingly anthropogenic habitats. The structure of forest fragments and distribution of edges is especially important in the context of anthropogenic forests because the history of the edges themselves can cause differential effects on different species and their movement. Anthropogenic edges, which are maintained by recent human disturbance, are less permeable to forest specialist carabids (Magura et al., 2017) and fungi (Ruete et al., 2017) than edges maintained by natural processes. In the context of plantation forest or managed woodland, this highlights the degree of influence that edge creation in forests can have of on species, including edge specialists.

Plantation forests represent a significant portion of forested area in the UK (Defra, 2021) and the management of these forests for timber necessitates canopy gaps (known as rides) to provide access for industrial activity, in addition to firebreaks, geographical features and land ownership boundaries that constrain where trees can be planted. Many of the resulting canopy gaps are linear, here collectively termed 'rides'. These rides can harbour fauna (Oxbrough et al., 2006) and flora (Smith et al., 2007) distinct from the interior of stands, providing refuges for species that might otherwise be unable to tolerate the planation environment. Sunlight is an important factor in the differences in species richness and composition between canopy gaps and the interior of stands (Sparks et al., 1996; Smith et al., 2007). The capacity of rides to harbour distinct communities is dependent on ride characteristics, in turn dependent on management practices during ride creation that determine their width and orientation, with downstream effects on community composition (Oxbrough et al., 2006; Smith et al., 2007). Ride orientation also influences the suitability on rides for certain species and locations: affecting the abundance of ground spiders (Lycosidae) but not ground beetles (Carabidae), for example (Carter et al., 1991). While these studies have focused on whole taxonomic groups, we might predict different responses from individual species that are edge specialists.

Certain species are more abundant in edge and transitional habitats between forest and open area because edges allow these species to benefit from two habitats at once. As a result, highly mobile vertebrates able to use multiple habitats, such as the noisy miner (*Manorina melanocephala*; Major *et al.*, 2001; Taylor *et al.*, 2008; Loyn, 1987) and white-tailed deer (*Odocoileus virginianus*; Williamson and Hirth, 1985), can be edge specialists. Similarly, plants may benefit from improved seed dispersal or pollination opportunities provided by the edge habitat (Lamb and Mallik, 2003) while maintaining the benefits of shelter and nutrient availability (Weathers *et al.*, 2001) from nearby forest. Highly mobile species commonly found on edges may merely be 'edge-tolerant' generalists exploiting their relative advantage over specialists of the bordering habitats (Ries and Sisk, 2010) rather than having any particular adaptive preference for edges. On the other hand, poor dispersers or sessile species that are more abundant in edge habitats are likely to be dependent on the edge habitat, because of their greater sensitivity to edges (Ewers and Didham, 2005); they will spend multiple generations under its influence and would therefore locally die out if the edge habitat was unfavourable for them. These species may be properly called edge specialists.

The population dynamics of edge specialists are affected by certain properties of edges. For example, the presence of complexities in the forests edge (corners, projections) predict noisy miner (*M. melanocephala*) occupation (Taylor *et al.*, 2008). Both edge orientation and width affect butterfly community diversity and composition, with shadier, south-facing edges (southern hemisphere) supporting greater forest butterfly diversity (van Schalkwyk *et al.*, 2022) Conversely, leaf miner, parasitoid, and plant communities appear less sensitive to light availability related to edge orientation (Bernaschini *et al.*, 2020). While edges clearly impact species distributions, the effect of ride width and especially ride bearing on dispersal are not fully understood.

Formica lugubris, a wood ant in the *Formica rufa* group, is a forest edge specialist and a poor disperser. UK populations of *F. lugubris* establish new nests by budding: a mated queen and some workers leave the maternal nest on foot to establish new nests (Maeder *et al.*, 2016). As a result, dispersal is local, and even when abundant suitable habitat is available, population margins extend by less than 50m per year (Procter, 2015). These ants are reliant on edge habitat because they require honeydew from arboreal aphids as their main source of carbohydrate but they also require sunlight to regulate the temperature of their nest (Domisch *et al.*, 2016). These dual requirements mean that *F. lugubris* nests are typically found within 10m of canopy gaps and on the north side of gaps (Procter *et al.*, 2015; Sudd et al., 1977). Their potential for dispersal and population expansion through forest is therefore dependent on the availability of sunlight. In managed forestry land this is largely determined by management decisions regulating the properties of canopy gaps in the form of firebreaks, access roads, footpaths and felled areas.

The orientation of linear canopy openings has the potential to affect both the abundance and dispersal ability of wood ants. In the northern hemisphere, south-facing sites tend to get more insolation. Therefore, along canopy gaps that run approximately east-west, the shadier (south) side of the ride will receive little insolation, so we would predict that wood ants would be less able to utilise nesting sites on the south side of these rides, while canopy gaps that run north-south would be occupied equally on either side, but to a lesser degree, due to the reduced hours in direct sunlight (Figure S.2.). Finally, we would predict that the width of a ride could affect wood ant abundance: narrow rides where the canopy nearly closes would be shadier, so we would expect fewer, larger nests on narrow rides and more, smaller nests on wider rides.

We conducted a survey of canopy gaps and edge characteristics in plantation forest of the North York Moors, and combined pre-existing long-term monitoring data with new data on wood ant populations at the study sites to answer the question: How do characteristics of rides affect the population dynamics of *F. lugubris*? We tested the following hypotheses:

H1 - The orientation of anthropogenic canopy gaps (degrees from N; 0-90°) affects wood ant nest a) abundance and b) total volume along those canopy gaps.

H2 - The orientation of anthropogenic canopy gaps (degrees from N; 0-90°) affects wood ant dispersal rate along those canopy gaps.

Methods

Ride selection: abundance and volume measures

Our study site was the North York Moors, UK, where *Formica lugubris* is abundant, populations are expanding into previously unoccupied areas, and high-quality nest distribution data is available from the years 2011, 2013, 2018-2022 (Procter, 2016; Holgate, 2021; Podesta 2023). Rides suitable for testing the hypothesis that canopy gap orientation will affect wood ant nest abundance and volume (H1) were identified prior to the field survey using pre-existing long-term data on wood ant populations in the North York Moors (Procter *et al.*, 2015; see Supplementary Methods S.1 and Figure S.6). A ride was defined as a linear feature causing a break in the canopy at least 5m wide, starting and ending where it intersected other rides or reached the end of the forest. Examples of rides included linear canopy gaps due to roads and footpaths, as well as gaps caused by forest management such as areas of row thinning (strips of trees harvested at intervals, leaving most of the crop standing) or firebreaks. We measured the length of the wood ant-occupied portion of the ride from the oldest nest on it to the nest furthest from the source population. The population margins and the source were identified using long-term data on the wood ant populations in the North York

Moors (Procter, 2016; Holgate, 2021; Podesta, 2023). For rides with nests along their entire length, this measurement equalled the total length of the ride. When mapping the margins, a margin was defined as the last occupied area after which no nest was encountered for 200m; this is far greater than typical dispersal distance for this species, consistent with the long-term data on these populations (Procter *et al.*, 2015). Rides were considered to have ended if the ride entered a stand of different tree species, a stand of different age/tree height, or rounded a corner (ride bearing changes by more than 15 degrees). We measured the length and bearing of the ride using the *Measure tool* in ArcGIS Pro version 2.3.3, connecting the first and last nest on the ride.

Selecting rides for calculating rate of expansion

In order to determine whether ride bearing affects wood ant dispersal, we required a subset of rides at the population margins, where population margin expansion could have occurred during our study period (since 2011) to calculate the average yearly population margins expansion along rides. In addition to the selection criteria in the previous section, margin rides were also required to meet the following criteria:

1. Have had potential for population expansion between 2011 and 2022 i.e., must have uncolonized length that does not extend into unsuitable habitat like open farmland.
2. Expansion has been unidirectional, from a single point of origin, to exclude situations where sections of the ride has been colonised from two origins (due to intersecting two rides with ant presence), as this might inflate the measures of margins expansion on this ride.
3. Expansion has not been interrupted by factors such as areas of forest on the ride being harvested during the period for which we have data, causing the ride to become unsuitable for wood ants. In cases where the expansion had been interrupted, we used the data up to the interruption. For example, a ride that was surveyed from 2011-21, but that was clear-felled in 2019, can provide us with an eight-year average expansion rate rather than ten.

Suitable rides were identified from three forests within the North York Moors, totalling 50 rides, 29 of which were on the margins and suitable for testing margins expansion, each with record of the bearing of the ride, length, mean width, the number of nests on each side of the ride. For the 29 rides on the margins, we also had two measures of the rate of margin change calculated from long-term population data: abundance change (nests/year) and distance change (m/year).

Nest abundance and volume measurements

For each ride, the surveyor walked down one side of the ride and up the other, recording nest presence and volume. Nests more than 5m from the edge were excluded to ensure all surveyed nests were directly using the ride canopy gap. In areas where the understory was relatively open,

spotting nests up to 5m from the edge did not present a challenge but in areas where the understory was dense, some nests were difficult to spot. Where the understory was too thick to confidently spot nests up to 5m from the edge, a second transect was walked, parallel to the first, 2.5m from the edge (behind obscuring vegetation) to minimise the risk of missing nests. At each nest, we recorded the location (latitude and longitude) and on which side of the ride (east or west/north or south) the nest was, giving each nest a unique ID code to prevent recounting. Additionally, nest width, length and height were measured to calculate nest volume as a proxy for number of ants (Chen and Robinson, 2013).

Ride details were also recorded: these included ride width (between opposite points where the canopy is directly overhead) and canopy height (measured using *Toolbox* android app calibrated against a gun clinometer) on both sides of the ride and ride bearing (expressed as degrees east or west of north). We measured each of these ride variables at the beginning, midpoint and end of each ride and averaged them across the three to provide ride characteristics. Latitude and longitude were recorded at the start and end of each ride so that the length of the ride could be measured retrospectively using ArcGIS Pro version 2.3.3.

Calculating rate of margins expansion along rides

From the long-term data, we calculated two measures of margins change a) distance change and b) number of nests. The distance the margin changed along a ride was the distance in metres between the location of the last nest along a ride recorded in the oldest long-term data and the location of the last nest on the same ride surveyed in 2022. The change in the number of nests was defined as the number of nests between the last nest at the date of the earliest survey in the long-term data and the last nest along the ride in the 2022 survey. Although both rate of distance change and the rate of nest abundance change can be calculated from this data, rate of nest abundance change is a less reliable measure of expansion than the rate of distance change for two reasons. Firstly, there is a five-year interval between measurements in the long-term data and while all population margins were mapped at each survey date, the entire population was not re-mapped on each survey (Procter *et al.*, 2015). This means that, in many cases, we have no evidence of continuous occupation, nor of the number of nests that may have been established and subsequently abandoned in the intervening period. Secondly, these populations of *F. lugubris* are polydomous, a life history trait whereby a single 'colony' builds and occupies many nest mounds, which share workers and food resources for mutual support. As a result, in areas where the wood ants have recently colonised, some small nests may be abandoned and the occupying ants absorbed into neighbouring nests in the polydomous network (Burns *et al.*, 2020) with the effect that the actual numbers of nests may fluctuate. On the other hand, the predicted rate of distance change was low, owing to the slow

dispersal of this species (Procter *et al.*, 2015) whereas we anticipated that the change in the number of nests would be large and more easily detectable.

Statistical methods

Modelling the relationship between ride characteristics and population measures

To test H1 (i.e., the effect of ride characteristics (bearing, width, and side) on the abundance and relative volume of nests) we fitted generalised additive models using the mgcv package version 1.8-41 (Wood, 2011) in R (version 4.2.1). Because our orientation data were circular and we could not make assumptions about the shape of the relationship between orientation and our wood ant population measures, the additional flexibility of a GAM was preferred over a GLM. We fitted models including all combinations of the ride characteristics (bearing, width, and side), including site and ride ID as random effects, and used the minimum AIC to select the most parsimonious model for each response variable (Supplementary Methods S.2). Because of the circular nature of ride bearing (-90° and 90° relative to north are equivalent), we included ride bearing smoothing term as a cyclic p-spline, whereas mean width was included as a p-spline only. We plotted response vs. fitted values, fitted vs. observed and histograms of the residuals to inspect the model fit, and predicted from the models for visualisation. The GAMs provided estimations of degrees of freedom. For nest abundance, cyclic p-splines were fitted for both east and west sides of rides from predicted values of a point count generalised additive model using ride bearing as a predictor offset by ride length, to account for the zero inflation of count data, with the formula nest abundance ≈ side + s(bearing) + s(width) of the negative binomial family (Link function = log, offset = log(length). For nest volume, the same approach was used, with the formula volume per nest ≈ side + s(bearing) + s(width). For all our models, we calculated deviance explained, as deviance explained is favoured over R^2 for non-Gaussian models (Table 1, model 1; the nest abundance model) as a more robust measure of explanatory power (Wood, 2006).

Pooling bearing into categories for margins growth analysis

Our final dataset resulted in 29 rides available to test H2 (rate of margin change). The bearing data that we collected was continuous, and this was used to fit the GAM model for margins growth as described above. Inspection of the data, however, suggested a bigger difference between rides bearing near the cardinal points (N, S, E, W) and rides bearing near the intercardinals (NE, SW etc.) than between N-S and E-W rides. This suggests that, to test pooled bearing, bins should be E-W +/- 22.5°, N-S +/- 22.5°, and intercardinal +/- 22.5° (Figure S.1.), in order to appropriately capture the variability in the data. This variability has sound ecological justification: assuming that there is a minimum threshold of daily sunlight exposure for nesting sites to be viable for the wood ant nests to

persist, we can be sure that this threshold lies somewhere below 50% of summer day lengths, as both sides of N-S rides are often utilised by ants. This threshold is also greater than 0% of the daily hours of sunlight because wood ants were often entirely absent from the south side of E-W rides. As such, going around the compass from north, the hours of available light on the north side will gradually increase from 50% up to c100% after 90°. Meanwhile, the south side will decrease from 50% to nearly 0%, passing below the threshold for wood ants to utilise potential nesting sites somewhere between. This reduction in potential nesting sites and the total availability of sunlight on the ride will have negative effects on wood ant populations and result in slower population growth leading to slower margins expansion.

Results

We found that nests were mor abundant on the north sides of rides and varied considerably with ride orientation; more nests were found on rides that ran E-W or N-S and fewer on rides that were oriented NE-SW or NW-SE (intercardinal). Furthermore, we found that the rate (number of nests) and the distance of dispersal of wood ant nests was lowest along rides intercardinal rides, and three times greater on cardinal rides. On the other hand, no consistent effect was found of ride orientation on nest volume.

Ride bearing distribution

A total of 371 nests were recorded along all studied rides. One of the rides on the margins was also at the true edge of a forest stand and therefore the ride had no measurable width and only one side on which nests could occur. This ride was dropped from analyses that incorporated side or ride width as variables. Ride orientation covered the full range of possible bearings but rides approximately 15-30° west of north and 70-80° east of north were more abundant than other orientations (Figure 1).

Nest aspect distribution

The aspect (direction of the long face/lower edge) of nest mounds was overwhelmingly south-facing (Figure 1) with a circular mean of 191.47°, significantly different from a uniform distribution (Rayleigh's test, r = 197.47, var = 0.46, p < 0.0001). This is what would be expected if nests are constructed to present a large surface area to the sun and maximise insolation.

Does the orientation of canopy gaps affect wood ant nest abundance and volume?

The most parsimonious model of nest abundance included only side, bearing and width (Figure 2; Table 1). There was a significant interaction between ride bearing, and the side of the ride on which

nests are most abundant (Figure 2), with nests being most abundant on the East sides of rides that have bearings around the NW-SE axis and most abundant on the West side of rides that have bearings around the NE-SW axis. For rides that have bearings on the NS axis, no difference in abundance between sides was seen. Overall, the number of nests per side was compensated by the other (Figure 2): when the east side has few nests (at bearings that cause the east side to be shaded), the west side has more and vice-versa, so the total sum of nests per ride varied little with bearing. Ride width did not have a significant effect on nest abundance (Table 1).

Figure 1. Left: The frequency of rides at different bearings was determined by the availability of suitable rides at the field sites; ride bearings include most possible directions without excessively favouring one direction. Right: The aspect (direction of the lowest side of the nest) of nests was most commonly just west of south (180-225 degrees).

Table 1. Generalised additive models to predict several wood ant population measures using ride characteristics. Models 1 and 2 included the side of the ride that nests were on as a 'by' term (fitting the two sides as separate splines) so that we could test the significance of each smooths (difference from line with gradient of 0) for data from the east side of rides and the west, and test the smooths against one another (East:West). Both R^2 and deviance explained are included, as deviance explained is preferred for the non-Gaussian model 1. Expanded table in Supplementary Material (Table S.1.).

Model	Dependant variable	'By' interaction	Smoothing term		Effective degrees of freedom	Test statistic (F unless specified)	P-Value	N	R^2 (Deviance explained)	Notes
1	Nest abundance	Side	Ride Bearing	Side = East	4	17.741 (X^2)	0.001	98	0.054 (31%)	Nest abundance offset by ride length (Figure 2)
				Side = West	4	16.883 (X^2)	0.002			
				East:West	4	17.75 (X^2)	0.001			
			Ride Width		1.142	0.407 (X^2)	0.557			
2	Mean nest volume	Side	Ride Bearing	Side = East	5	3.476	0.008	69	0.267 (40.5%)	(Figure 2)
				Side = West	5	1.059	0.393			
				East:West	5	0.877	0.502			
			Ride Width		1.823	5.487	0.006			
3	Margins nest abundance change per year	None	Ride Bearing		4	4.443	0.082	28	0.163 (41.1%)	(Figure 4)
			Ride Width		4	2.061	0.126			
4	Margins distance change per year	None	Ride Bearing		4	1.934	0.148	28	0.256 (50.8%)	(Figure 4)
			Ride Width		4	1.011	0.428			
			Site		1.154	1.53	0.098			

Figure 2. Left: the total volume of all nests on each side of rides of different bearings. The lines of predicted values from a generalised additive model (GAM; R^2 = 0.267, deviance explained = 40.5%) were plotted for values of ride bearing (lines) alongside the original data (points) and 95%CIs (shaded areas). Although the east side smooth was significantly different from a line at the intercept of gradient = 0 ($p < 0.01$), there was no significant difference between the east and west smooths and no clear pattern is discernible from the plot. Mean width was also significant as a smoothing term (Figure 3; Table 1). Right: the number of nests per side of a ride (E or W) for rides of a range of bearings. The points represent the real data, the lines are predicted values and the shaded areas are 95%CIs. Both east and west smooths were significantly different from 0 ($p < 0.005$) and significantly different from each other ($p = 0.001$).

The most parsimonious model of nest volume included site, bearing and width as smoothing terms, however the inclusion of site only marginally improved explanatory power and did not justify a more complex model, so was dropped (Table S.1; Figure S.3). In both models, the ride bearing significantly affected volume only for nests on the East side, and relationship between ride bearing and nest volume on the East and West sides did not differ significantly from one another (Figure 2; Table 1). The asymmetry of this result, and the lack of difference between the two sides, suggests that ride bearing is a poor predictor of mean nest volume for these data.

Mean ride width was significant as a smoothing term ($p < 0.01$; Table 1) and a plot of model predictions (Figure 3) shows that the mean volume of nests is lower on wider rides but does not continue to decrease as ride width exceeds 8m. A relatively small number of large nests on rides narrower 6m may be major contributors to this effect; however, this small number of nests were distributed across all three sites.

Does the orientation of canopy gaps affect wood ant dispersal rate?

We used two measures of population margins growth: the change in the number of nests at the margin and the distance that the margins moved per year over the study period. The model that best explained the change in nest number per year included ride bearing and width (Figure 4; Table 1). Neither term was below the significance threshold of 0.05 (ride bearing $p = 0.082$, width $p = 0.126$) and the model explained only approximately 15% of the variation. Overall, we see a similar shaped pattern in the rate of nest number change (Figure 4) as we saw for nest abundance on each side of rides (Figure 2), with minima for rides bearing approximately NW-SE and NE-SW (intercardinal), though not reaching statistical significance. While this model has poor predictive power, it does suggest that bearing may play some role in the rate of change in nest number.

Figure 3. Generalised additive model (GAM) with the formula volume per nest ≈ s(bearing) + s(width), R^2 = 0.18, deviance explained = 28.7% were plotted for values of width (line) alongside the original data (points) and 95%CIs (shaded area), with the points representing field data. The smooth of mean width was significantly different from the intercept (p < 0.01), whereas ride bearing alone (i.e. not split by side) was not.

Figure 4. Upper: a generalised additive model (GAM) was fit with the formula nest abundance change per year ≈ s(bearing) + s(width), R^2 = 0.163, deviance explained = 41.1% and the predictions for values of ride bearing were plotted (solid line) with 95%CIs (shaded areas). Ride bearing was not significant as a smoothing term and the model has little explanatory power. Ride width is not correlated with bearing. Including width as a smoothing term lower predicted values than the observed values due to the fixing of ride width at the mean value for non-Gaussian data. Lower: A generalised additive model was fit with the formula margins change ≈ s(bearing) + s(width) + s(site), R^2 = 0.256, deviance explained = 50.8%. Empirical data (points), model predictions (lines) and 95%CIs (shaded areas) are shown. None of the smoothing terms were significant, but site was a significant random effect and its inclusion resulted in the most parsimonious model.

The median rate of nest abundance change was greater on rides with cardinal orientations than those with intercardinal ones (approximately 0.5 nests/year; Figure 5), but this difference did not reach statistical significance (Welch's t-test, t = 2.02, df = 21.37, p = 0.056). A further split in the data was tested to N-S, E-W, NW-SE and NE-SW against each other, but no statistically significant difference was found between them for either variable (Figures S.4 and S.5).

The distance that the margins moved per year over the study period was also included as a response variable in a GAM and the best model includes site, ride bearing and ride as smoothing terms (Figure 4; Table 1). This model performed better than the nest number change model (R^2 = 0.256) and the overall pattern was the same (minima on rides near intercardinal directions), but none of the smoothing terms were significant (p > 0.05; Table 1). Model predictions showed that rides in Cropton Forest experienced more wood ant population growth than either of the other two sites.

Overall, rides running in intercardinal resulted in lower population margins growth (Figure 5). We found that the rides that bear in cardinal directions experienced three times the annual margins expansion than intercardinal ones (15m/year compared to 5m/year; Welch's t-test, t = 2.122, df = 22.497, p = 0.045; Figure 5; Table 2) and the effect size was medium (Hedges' g =0.72).

Table 2. Dispersal range was measured as rate of change of the number of nests at the margin, and distance of the final nest along a ride from the original (2011) population extent to the final (2022) population extent. We made compared rides of varied orientation relative to north; cardinal (N-S and E-W +/- 22.5°) and intercardinal (NW-SE and NE-SW+/- 22.5°) bearings.

Measure	Bins	Tests	Degrees of freedom	Test statistic	P-value	Effect size (Hedges' g)
Rate of change in nest number at margins (Figure 5)	2	Welch's *t*-test	21.37	t = 2.02	0.056	0.68
Rate of distance change at margins (Figure 5)	2	Welch's *t*-test	22.50	t = 2.12	0.045	0.72

Figure 5. Ride data for change in nest number (A; Welch's *t*-test, t = 2.02, p = 0.056), and distance of margin change (B; Welch's *t*-test, t = 2.12, p = 0.045), comparing cardinal (22.5° +/- N, S, E or W) and intercardinal (22.5° +/- NE, NW, SE or SW) bearings (Table 2 for statistical details).

Discussion

Summary of results

We aimed to determine the effect of anthropogenic canopy gap characteristics on populations of an edge specialist, the northern hairy wood ant *Formica lugubris*. Our results clearly show that the orientation and width of linear canopy gaps affect *F. lugubris* distribution and dispersal, shaping their population dynamics. Nests were more abundant on the northern side of rides for all orientations, as predicted, but orientation did not similarly affect nest size in a consistent way. In general, narrower rides had larger nests, while wider rides were populated by smaller nests, consistent with predictions that nests receiving less insolation are larger (Chen and Robinson 2014). Finally, we found that, as predicted, the orientation of rides affected *F. lugubris* dispersal. Over the preceding 10 years, populations of *F. lugubris* dispersed more quickly along rides that run in cardinal directions (N, E, S, W) than ones that run in intercardinal directions (NE, SE, SW, NW).

The importance of edge characteristics

We showed that the orientation of edge habitats relative to north is an important factor in edge quality for *Formica lugubris* and is sufficient to have measurable effect on nest abundance and population margin expansion. Ride width is also a factor in edge quality for these wood ants; we found larger nests in narrower rides. In general, larger nests are found in shadier areas because larger nests are better able to thermoregulate (Chen and Robinson, 2014). The strong south-facing bias in nest aspect we found confirms that our other results are likely to driven by insolation, as sunlight is important for many aspects of wood ant biology. Nests lacking direct sunlight, especially small nests, may initiate foraging later (Rosengren *et al.*, 1987), have slower brood maturation (Kadochová and Frouz, 2013) and therefore potentially slower colony growth.

Dispersal along rides

In addition to ride bearing affecting wood ant population density (nest volume and abundance), it also affects the ability of the wood ants to disperse through these plantation forests, with most rapid dispersal along rides oriented along the cardinal bearings (N-S or E-W). Due to the generally slow dispersal of *F. lugubris*, the cardinal ride dispersal distance of 15m/year is equivalent to three times the rate of dispersal in intercardinal rides (5m/year). We hypothesise that the slower population margin expansion observed for rides with bearings in intercardinal directions is due to a minimum threshold of insolation for newly established nests of *F. lugubris* to be viable. Above this threshold, a wood ant nest's outcomes continue to improve. We suggest that the south side of a ride with an intercardinal bearing receives insolation below this minimum threshold, due to being shaded for most of the day by the trees on the southern side, while the north side will receive less insolation

than the north side of a ride running east-west. This means nests on these bearings are in a 'worst of both worlds' environment resulting in fewer individuals and a reduced ability to form new nests in the polydomous network. Polydomy allows for food and labour sharing between nests that can subsidise a new nest while it grows big enough to establish foraging (Ellis and Robinson, 2015). Newly established nests in shadier areas may require extra support from neighbouring nests (Rosengren *et al.*, 1987), so being further from the centre of this network may leave new nests more vulnerable to abandonment. The combination of lower rates of nest foundation and a higher failure rate for nests that have been established further from the parental nests, could explain the reduced margins growth that we measured in shadier, intercardinal rides.

Management implications

Given that wood ants increase biodiversity in plantation forests and may provide additional pest control benefits (Laine and Niemela, 1980; Karhu and Neuvonen 1998; Zingg *et al.*, 2018), the results that we present here may be useful in planning management of plantations. Wood ants can reduce defoliating pest burden (Karhu and Neuvonen 1998), influence tree growth via their effects on aphids (Kilpeläinen *et al.*, 2009), compete with other insectivores (Haemig, 1992), and change the spatial distribution of soil resources (Podesta, 2023), so the effects of ride characteristics on their dispersal have major implications for conservation and forest management. When planning new areas of forestry adjacent to existing populations of wood ants, or in areas where translocation of wood ants' nests is being considered (Nielsen *et al.*, 2018), we would recommend cutting firebreaks, access tracks etc. in cardinal directions, to maximise the ability of wood ants to disperse. The width of the ride should also be considered when managing forest to promote species richness or dispersal. Previous work has suggested that canopy gaps should be at least 15m wide in order to harbour species that would be shaded out in the forest interior (Oxbrough *et al.,* 2006; Smith *et al.,* 2007), and our data suggest that wood ants similarly have a minimum threshold of insolation (also impacted by ride width) that allows them to thrive and disperse.

Wider ecological implications

The work presented here adds to a body of research showing that linear canopy gaps in plantation forests have important implications for forest dwelling species. The characteristics of rides, such as width, affect the diversity or abundance of butterflies (Greatorex-Davies *et al.*, 1993), spiders (Oxbrough *et al.*, 2006; Carter *et al.,* 1991), ground beetles (Carter *et al.,* 1991), vascular plants (Sparks *et al.,* 1996; Smith *et al.,* 2007), and bryophytes (Smith *et al.,* 2007). We have shown that, in addition to the effects that ride characteristics can have on the diversity and abundance of forest species in plantation, the orientation of rides can influence the dispersal ability of an edge specialist,

opening up the question of whether other species with similar habitat requirements might be similarly affected.

Managing plantations in a manner favourable to wood ant dispersal (by cutting firebreaks and other necessary linear gaps along cardinal directions) could be beneficial to other species and biodiversity in economically productive forests in two ways. Firstly, the presence of wood ants themselves can be beneficial to other species. The nests of wood ants are home to many myrmecophilous species, with over 100 myrmecophile species known from wood ants' nests (Robinson *et al.*, 2016; Parmentier *et al.,* 2014). Additionally, wood ants can play an important role in the dispersal of seed, including some declining and charismatic wildflowers like the endangered *Melampyrum cristatum* (crested cow-wheat) which is dispersed by wood ants and is likely range-limited by the availability of dispersers (Stachnowicz, 2013). Secondly, other species with similar habitat requirements and dispersal ability to the wood ants would benefit from the same rides as dispersal corridors as the wood ants. Species that might benefit from the same corridor design as wood ants could include wildflowers that likewise require direct sunlight but benefit from other conditions provided by the forest edge such as protection from wind, acidic soil or edge related nutrient influx (Weathers *et al.*, 2001).

On the other hand, species movement can be detrimental; there are well documented negative economic (Cuthbert *et al.*, 2022) and environmental (Pyšek *et al.*, 2020) effects of species invasions. Edge specialists with the potential to benefit from improved dispersal due to edge orientation may also be pests, for example, the larvae of *Gilpinia virens,* a pine sawfly and common pest of *Pinus sylvestris*, are found in higher numbers in stands with relatively open canopies (Gawęda and Grodzki, 2020). As a result, pests such as *G. virens* may also benefit from the presence of canopy gaps cut in favourable, cardinal directions, although it should be noted that it does not necessarily follow that a preference for canopy gaps will result in the same response to different gap characteristics as we see in wood ants.

Conclusions

The potential to promote dispersal of both rare and desirable species as well as pests of canopy gaps in plantation forests makes forest edges and linear canopy gaps worthy of more detailed study. Very little work has investigated the effects of gap orientation on the dispersal ability of plants or animals that might utilise plantation forests despite the necessity of gaps in plantations forests as fire breaks or for access for logging machinery, and the control that forestry planners have over the placement of fire breaks and other linear canopy gaps. The increase in demand for timber products and forested areas to reach carbon offset goals will likely lead to an increase in the area and intensity of

plantation forests in the future (McEwan et al., 2020). If these economic and environmental goals are not to come at the detriment of biodiversity or resilience, care must be taken to ensure the development of forestry practices that allow wildlife to proliferate in plantation forests, for the ecosystem services they can provide and for the biodiversity that they represent.

Acknowledgements

We thank Duncan Procter and Megan Holgate for additional data that contributed to the work and Forest Research for site access permission. We also thank Imre Sándor Piross and Mark Hodson for advice with the analysis and manuscript respectively. The project was funded by the NERC ACCE2 DTP.

References

Bennett, A.F. (1990) 'Habitat corridors and the conservation of small mammals in a fragmented forest environment', *Landscape Ecology*, 4(2–3), pp. 109–122.

Bernaschini, M.L., Valladares, G. and Salvo, A. (2020) 'Edge effects on insect–plant food webs: assessing the influence of geographical orientation and microclimatic conditions', *Ecological Entomology*, 45(4).

Brito, T.F. *et al.* (2017) 'Forest reserves and riparian corridors help maintain orchid bee (Hymenoptera: Euglossini) communities in oil palm plantations in Brazil', *Apidologie*, 48(5), pp. 575–587.

Brunzel, S., Elligsen, H. and Frankl, R. (2004) 'Distribution of the Cinnabar moth *Tyria jacobaeae* L. at landscape scale: Use of linear landscape structures in egg laying on larval hostplant exposures', *Landscape Ecology*, 19(1).

Burns, D.D.R. *et al.* (2020) 'A longitudinal study of nest occupancy, trail networks and foraging in a polydomous wood ant population', *Insectes Sociaux*, 67(3), pp. 419–427.

Carter, C. I. (1991). 'Paper 4: Ride Orientation and Invertebrate Activity', *Edge management in woodlands*, 28, pp. 17-21.

Chen, Y.H. and Robinson, E.J.H. (2013) 'A comparison of mark-release-recapture methods for estimating colony size in the wood ant *Formica lugubris*', *Insectes Sociaux*, 60(3), pp. 351–359.

Chen, Y.H. and Robinson, E.J.H. (2014) 'The relationship between canopy cover and colony size of the wood ant *Formica lugubris*-implications for the thermal effects on a keystone ant species', *PLoS ONE*, 9(12).

Cuthbert, R.N. *et al.* (2022) 'Are the "100 of the world's worst" invasive species also the costliest?', *Biological Invasions*, 24(7).

Defra (2021) *The England Trees Action Plan 2021-2024*.

Domisch, T., Risch, A. and Robinson, E.J.H. (2016) Chapter 7 'Wood ant ecology and conservation', in J. Stockan and E.J.H. Robinson (eds). Cambridge University Press.

Ellis, S. and Robinson, E.J.H. (2014) 'Polydomy in red wood ants', *Insectes Sociaux*, 61(2), pp. 111–122.

Ellis, S. and Robinson, E.J.H. (2015) 'The role of non-foraging nests in polydomous wood ant colonies', *PLoS ONE*, 10(10).

Ewers, R. M. and Didham, R. K. (2005) 'Confounding factors in the detection of species responses to habitat fragmentation', *Biological Reviews*, 81(1), pp. 117-142.

Gawęda, P. and Grodzki, W. (2020) ' The influence of stand and habitat characteristics on the occurrence of pine sawflies *Diprion pini* L. and *Gilpinia virens* (Klug) (Hymenoptera, Diprionidae) in selected areas of northern Poland ', *Forest Research Papers*, 81(3).

Greatorex-Davies, J. N., Sparks, T. H., Hall, M. L., and Marss, R. H. (1993). 'The influence of shade on butterflies in rides of coniferised lowland woods in southern England and implications for conservation management', *Biological Conservation*, 63(1), 31-41.

Haemig, P.D. (1992) 'Competition between Ants and Birds in a Swedish Forest', *Oikos*, 65(3).

Holgate, M.K. (2021) 'The effects of land use change on the population expansion of a poor disperser, the wood ant *Formica lugubris'*, PhD Thesis, University of York.

Kadochová, Š. and Frouz, J. (2013) 'Thermoregulation strategies in ants in comparison to other social insects, with a focus on *Formica rufa'*, *F1000Research*, 2(May), p. 280.

Karhu, K.J. and Neuvonen, S. (1998) 'Wood ants and a geometrid defoliator of birch: Predation outweighs beneficial effects through the host plant', *Oecologia*, 113(4), pp. 509–516.

Kilpeläinen, J., et al., Does the mutualism between wood ants (*Formica rufa* group) and Cinara aphids affect Norway spruce growth? *Forest Ecology and Management*, 2009. **257**(1): p. 238-243

Laine, K.J. and Niemela, P. (1980) 'The Influence of Ants on the Survival of Mountain Birches During an *Oporinia autumnata* (Lep., Geometridae) Outbreak', *Oecologia (Berl.)*.

Lamb, E.G. and Mallik, A.U. (2003) 'Plant species traits across a riparian-zone/forest ecotone', *Journal of Vegetation Science*, 14(6).

Lenoir, J. and Svenning, J.C. (2015) 'Climate-related range shifts - a global multidimensional synthesis and new research directions', *Ecography*, 38(1).

Liira, J. and Paal, T. (2013) 'Do forest-dwelling plant species disperse along landscape corridors?', *Plant Ecology*, 214(3).

Loyn, R.H. (1987) 'Effects of patch area and habitat on bird abundances, species numbers and tree health in fragmented Victoria forests.', *Nature conservation: the role of remnants of native vegetation* [Preprint].

Maeder, A. *et al.* (2016) 'Wood ant reproductive biology and social systems', in J.A. Stockan and E.J.H.E. Robinson (eds) *Wood Ant Ecology and Conservation*. Cambridge University Press (Ecology, Biodiversity and Conservation), pp. 37–50.

Magura, T., Lövei, G.L. and Tóthmérész, B. (2017) 'Edge responses are different in edges under natural versus anthropogenic influence: a meta-analysis using ground beetles', *Ecology and Evolution*, 7(3).

Major, R.E., Christie, F.J. and Gowing, G. (2001) 'Influence of remnant and landscape attributes on Australian woodland bird communities', *Biological Conservation*, 102(1).

McEwan, A. *et al.* (2020) 'Past, present and future of industrial plantation forestry and implication on future timber harvesting technology', *Journal of Forestry Research*. Northeast Forestry University, pp. 339–351.

Mendenhall, C.D. *et al.* (2014) 'Predicting biodiversity change and averting collapse in agricultural landscapes', *Nature*, 509(7499).

Nielsen, J.S. *et al.* (2018) 'Experiences in transplanting wood ants into plantations for integrated pest management', *Sociobiology*, 65(3), pp. 403–414.

Oxbrough, A. G., Gittings, T., O'Halloran, J., Giller, P. S., and Kelly, T. C. (2006). 'The influence of open space on ground-dwelling spider assemblages within plantation forests', *Forest Ecology and Management*, 237(1-3), 404-417.

Parmentier, T., W. Dekoninck and T. Wenseleers (2014) 'A highly diverse microcosm in a hostile world: a review on the associates of red wood ants (*Formica rufa* group)', *Insectes Sociaux* 61(3): 229-237.

Podesta, J. (2023). Interactions between the wood ant *Formica lugubris* and plantation forests: how ants affect soil properties and how forest management affects the dispersal of an ecosystem engineer. PhD thesis, University of York.

Procter, D.S. (2016) 'The effects of forest cover change and polydomous colony organisation on the wood ant *Formica lugubris*', PhD Thesis, University of York.

Procter, D.S. *et al.* (2015) 'Do non-native conifer plantations provide benefits for a native forest specialist, the wood ant *Formica lugubris*?', *Forest Ecology and Management*, 357, pp. 22–32.

Pyšek, P. *et al.* (2020) 'Scientists' warning on invasive alien species', *Biological Reviews*, 95(6).

Ries, L. and Sisk, T.D. (2010) 'What is an edge species? The implications of sensitivity to habitat edges', *Oikos*, 119(10).

Robinson, E.J.H., Stockan, J.A. and Iason, G.R. (2016) 'Wood ants and their interaction with other organisms', in J.A. Stockan and E.J.H. Robinson (eds) *Wood Ant Ecology and Conservation*. Cambridge: Cambridge University Press, pp. 177–199.

Roberts, D., Ciuti, S., Barber, Q. E., Willier, C., and Nielsen, S. E. (2018) 'Accelerated seed dispersal along linear disturbances in the Canadian oil sands region', *Scientific Reports*, 8(1), 4828.

Rosengren, R. *et al.* (1987) 'Phenology and causation of nest heating and thermoregulation in red wood ants of the *Formica rufa* group studied in coniferous forest habitats in southern Finland.', *Annales Zoologici Fennici*, 24(2), pp. 147–155.

Ruete, A. *et al.* (2017) 'Contrasting long-term effects of transient anthropogenic edges and forest fragment size on generalist and specialist deadwood-dwelling fungi', *Journal of Applied Ecology*, 54(4), pp. 1142–1151.

Smith, G. F., Iremonger, S., Kelly, D. L., O'Donoghue, S., and Mitchell, F. J. (2007). Enhancing vegetation diversity in glades, rides and roads in plantation forests. *Biological Conservation*, 136(2), 283-294.

Sparks, T. H., Greatorex-Davies, J. N., Mountford, J. O., Hall, M. L., and Marrs, R. H. (1996) 'The effects of shade on the plant communities of rides in plantation woodland and implications for butterfly conservation', *Forest Ecology and Management*, 80(1-3), 197-207.

Stachnowicz, W. (2013) 'Melampyrum Cristatum L. – A Rare River Corridor Plant in Wielkopolska and Poland', *Biodiversity Research and Conservation*, 32(1).

Suárez-Esteban, A. *et al.* (2016) 'Can anthropogenic linear gaps increase plant abundance and diversity?', *Landscape Ecology*.

Sudd, J.H. *et al.* (1977) 'The distribution of wood-ants (*Formica lugubris* Zetterstedt) in a northern English forest', *Ecological Entomology*, 2(4).

Taylor, R.S., Oldland, J.M. and Clarke, M.F. (2008) 'Edge geometry influences patch-level habitat use by an edge specialist in south-eastern Australia', *Landscape Ecology*, 23(4), pp. 377–389.

van Schalkwyk, J. *et al.* (2022) 'Corridor width and orientation are complementary design variables for butterflies in conservation corridors', *Landscape Ecology* [Preprint].

Weathers, K.C., Cadenasso, M.L. and Pickett, S.T.A. (2001) 'Forest edges as nutrient and pollutant concentrators: Potential synergisms between fragmentation, forest canopies, and the atmosphere', *Conservation Biology*, 15(6).

Williamson, S. and Hirth, D. (1985) 'An Evaluation of Edge Use by White-tailed Deer', *Wildlife Society Bulletin*, 13(3).

Wood, S.N. (2006) 'Generalized Additive Models: An Introduction with R', Chapman and Hall/CRC.

Wood, S.N. (2011) 'Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models', *Journal of the Royal Statistical Society (B)*, 73(1), pp. 3–36.

Zingg, S. *et al.* (2018) 'The negative effect of wood ant presence on tick abundance', *Parasites and Vectors*, 11(1).