Wood ants on the edge: how do the characteristics of linear edges effect the population dynamics of an edge specialist?

Jacob A. Podesta[1], Catherine L. Parr[2], Kelly R. Redeker[1] and Elva J. H. Robinson[1*]

[1]Department of Biology, University of York, York, UK

[2]School of Environmental Sciences, University of Liverpool, Liverpool, UK

Abstract

Landscape structure modulates species dispersal by presenting barriers or opportunities. Slow-dispersing edge specialists, e.g. the northern hairy wood ant (*Formica lugubris*), are likely to be most affected by topography and land management practices, because they require adjacent contrasting habitats, e.g. to access both food and sunlight. In managed forests, canopy gaps are often linear and anthropogenic, such as paths, firebreaks, and roads (collectively 'rides'), and their orientation determines shade distribution. Using data spanning 10 years, we ask how ride orientation and width affect the distribution and dispersal of three *F. lugubris* populations in the North York Moors, UK. Ride orientation clearly affected nest abundance, with a higher nest density on rides oriented north-south (N-S) or east-west (E-W) (cardinal directions) than on those oriented NE-SW or NW-SE (intercardinal directions). Conforming to predictions based on sunlight availability, N-S oriented rides were occupied more symmetrically than E-W ones, where the north side was used predominantly. Nests were generally larger on narrower rides. Ride orientation also clearly affected dispersal: wood ants dispersed c15 m/year along rides oriented in cardinal directions, compared with only c5 m/year on rides oriented intercardinally. Our results show that ride characteristics (width, orientation), resulting directly from forestry practices, influence the distribution and dispersal of an ecosystem engineer woodland species; this may also apply to other forest-edge specialists. As wood ants can suppress defoliating pests, these findings could benefit forest management; forest planners could encourage wood ant colonisation of plantation forest by ensuring linear features contribute to north-south and east-west connectivity.

* Corresponding author email address: Elva.Robinson@york.ac.uk (Elva J. H. Robinson).

Keywords

Dispersal, Habitat fragmentation, Forest edges, Social insects, Population dynamics, Fire management

Introduction

The ability of species to move through a landscape can have major implications for the conservation of biodiversity, and for preventing species loss in a changing climate (Mendenhall *et al.*, 2014; Lenoir and Svenning, 2015). How porous a landscape is to individual movement will also affect the dispersal of species, and this porosity may be determined by the contrast between one habitat and the next. For example, low contrast edges between forest patches and agricultural land in Central American countryside ecosystems are more porous to bats than high contrast edges between forest patches and open water, and forest patches in the agricultural ecosystem harbour higher bat species richness as a result (Mendenhall *et al.*, 2014). The permeability of habitats to species may become more important as climate change induced range shifts along latitudinal or altitudinal gradients become more apparent (Lenoir and Svenning, 2015).

Natural linear features such as rivers, coasts, habitat edges, and mountain ranges, or anthropogenic ones such as roads, railways and canals can be both barriers (Shepard *et al.,* 2008) or facilitators of species movement, providing corridors of suitable habitat for certain species through otherwise inhospitable areas. For example, wind-dispersed seeds travel nearly four times the distance along linear forest disturbances than in undisturbed forest (Roberts *et al.*, 2018) while, in a meta-analysis, 70 % of studies indicate that either plant abundance or diversity is higher along linear gaps and verges than habitat interiors (Suárez-Esteban *et al.*, 2016). By facilitating the dispersal of food plants, linear features can indirectly promote the dispersal of invertebrate herbivores. Ragworts (*Senecio* spp.) disperse along man-made linear features (e.g., railways), and ragwort-dependant cinnabar moths (*Tyria jacobaeae*) show increased dispersal and egg-laying success in areas with roads and valleys compared with areas without linear features (Brunzel et al., 2004), suggesting that invertebrates also benefit from linear features.

The creation of linear features can be a conservation tool, as in the use of habitat corridors to connect smaller areas of protected land and reserves (Bennett, 1990; Brito *et al.*, 2017). However, habitat corridors can affect generalist and specialist species differently: landscape corridors connecting areas of fragmented forest in south-eastern Estonia facilitated the dispersal of generalist plant species more than forest specialist species (Liira and Paal, 2013). The presence and structure of edges thus has conservation potential, but the implications for species movement and survival may differ depending on the target species.

The widespread anthropogenic changes in natural forests, and the creation of large areas of plantation forests, mean that human activities determine the present distribution and fragmentation of most

forests. This makes the study of how fragmentation and edge habitats impact species dispersal
important for conserving species in increasingly anthropogenic habitats. The structure of forest
fragments and distribution of edges is especially important in the context of anthropogenic forests
because the history of the edges themselves can cause differential effects on different species and their
movement. Anthropogenic edges, which are maintained by recent human disturbance, are less
permeable to forest specialist carabids (Magura *et al*., 2017) and fungi (Ruete *et al.*, 2017) than edges
maintained by natural processes. In the context of plantation forest or managed woodland, this
highlights the degree of influence that edge creation in forests can have of on species, including edge
specialists.

Plantation forests represent a significant portion of forested area in the UK (Defra, 2021) and the
management of these forests for timber necessitates canopy gaps (known as rides) to provide access
for industrial activity, in addition to firebreaks, geographical features and land ownership boundaries
that constrain where trees can be planted. Many of the resulting canopy gaps are linear, here
collectively termed 'rides'. These rides can harbour fauna (Oxbrough *et al.,* 2006) and flora (Smith *et*
al., 2007) distinct from the interior of stands, providing refuges for species that might otherwise be
unable to tolerate the plantation environment. Sunlight is an important factor in the differences in
species richness and composition between canopy gaps and the interior of stands (Sparks *et al.,* 1996;
Smith *et al.,* 2007). The capacity of rides to harbour distinct communities is dependent on ride
characteristics, in turn dependent on management practices during ride creation that determine their
width and orientation, with downstream effects on community composition (Oxbrough *et al.*, 2006;
Smith *et al.*, 2007). Ride orientation also influences the suitability on rides for certain species and
locations: affecting the abundance of ground spiders (Lycosidae) but not ground beetles (Carabidae),
for example (Carter *et al.*, 1991). While these studies have focused on whole taxonomic groups, we
might predict different responses from individual species that are edge specialists.

Certain species are more abundant in edge and transitional habitats between forest and open area
because edges allow these species to benefit from two habitats at once. As a result, highly mobile
vertebrates able to use multiple habitats, such as the noisy miner (*Manorina melanocephala*; Major *et*
al., 2001; Taylor *et al*., 2008; Loyn, 1987) and white-tailed deer (*Odocoileus virginianus*; Williamson
and Hirth, 1985), can be edge specialists. Similarly, plants may benefit from improved seed dispersal
or pollination opportunities provided by the edge habitat (Lamb and Mallik, 2003) while maintaining
the benefits of shelter and nutrient availability (Weathers *et al*., 2001) from nearby forest. Highly
mobile species commonly found on edges may merely be 'edge-tolerant' generalists exploiting their
relative advantage over specialists of the bordering habitats (Ries and Sisk, 2010) rather than having
any particular adaptive preference for edges. On the other hand, poor dispersers or sessile species that
are more abundant in edge habitats are likely to be dependent on the edge habitat, because of their
greater sensitivity to edges (Ewers and Didham, 2005); they will spend multiple generations under its

influence and would therefore locally die out if the edge habitat was unfavourable for them. These
species may be properly called edge specialists.

The population dynamics of edge specialists are affected by certain properties of edges. For example,
the presence of complexities in the forests edge (corners, projections) predict noisy miner (*M.*
melanocephala) occupation (Taylor *et al.*, 2008). Both edge orientation and width affect butterfly
community diversity and composition, with shadier, south-facing edges (southern hemisphere)
supporting greater forest butterfly diversity (van Schalkwyk *et al.*, 2022) Conversely, leaf miner,
parasitoid, and plant communities appear less sensitive to light availability related to edge orientation
(Bernaschini *et al.*, 2020). While edges clearly impact species distributions, the effect of ride width
and especially ride bearing on dispersal are not fully understood.

Formica lugubris, a wood ant in the *Formica rufa* group, is a forest edge specialist and a poor
disperser in much of its range, including the UK (Ellis and Robinson, 2014). UK populations of *F.*
lugubris establish new nests by budding: a mated queen and some workers leave the maternal nest on
foot to establish new nests (Maeder *et al.*, 2016). As a result, dispersal is local, and even when
abundant suitable habitat is available, population margins extend by an average of 5 m per year
(Procter *et al.*, 2015). These ants are reliant on edge habitat because they require honeydew from
arboreal aphids as their main source of carbohydrate but they also require sunlight to regulate the
temperature of their nest (Rosengren *et al.,* 1987; Domisch *et al.*, 2016). These dual requirements
mean that *F. lugubris* nests are typically found within 10m of canopy gaps and on the north side of
gaps (Procter *et al.*, 2015; Sudd et al., 1977). Their potential for dispersal and population expansion
through forest is therefore dependent on the availability of sunlight. In managed forestry land this is
largely determined by management decisions regulating the properties of canopy gaps in the form of
firebreaks, access roads, footpaths and felled areas.

The orientation of linear canopy openings has the potential to affect both the abundance and dispersal
ability of wood ants. In the northern hemisphere, south-facing sites tend to get more insolation.
Therefore, along canopy gaps that run approximately east-west, the shadier (south) side of the ride
will receive little insolation, so we would predict that wood ants would be less able to utilise nesting
sites on the south side of these rides, while canopy gaps that run north-south would be occupied
equally on either side, but to a lesser degree, due to the reduced hours in direct sunlight (Figure S.2.).
Additionally, wood ant nests found in shadier locations are typically larger; an adaptive response
allowing them to thermoregulate more effectively (Chen and Robinson, 2014). As a result, we expect
that nests on rides oriented north-south, or nests on the south side of rides running east-west, will be
larger than their counterparts on the north side, on average. Finally, we would predict that the width of
a ride could affect wood ant abundance: narrow rides where the canopy nearly closes would be

shadier, so we would expect larger but less numerous nests on narrow rides and more numerous
smaller nests on wider rides.

We conducted a survey of canopy gaps and edge characteristics in plantation forest of the North York
Moors at three sites (Cropton Forest, Broxa Forest and East Moor Wood; Figure S.6), and combined
pre-existing long-term monitoring data with new data on wood ant populations at the study sites to
answer the question: How do characteristics of rides affect the population dynamics of *F. lugubris*?
We tested the following hypotheses:

H1 - The orientation of anthropogenic canopy gaps (degrees from N; 0-90°) affects wood ant nest a)
abundance and b) average volume of nests along those canopy gaps.

H2 - The orientation of anthropogenic canopy gaps (degrees from N; 0-90°) affects wood ant dispersal
rate along those canopy gaps.

147 Methods

148 Ride selection: abundance and volume measures

Our study site was the North York Moors, UK, where *Formica lugubris* is abundant, populations are
expanding into previously unoccupied areas, and high-quality nest distribution data is available from
the years 2011, 2013, 2018-2022 (Procter, 2016; Holgate, 2021; Podesta 2023). Rides suitable for
testing the hypothesis that canopy gap orientation will affect wood ant nest abundance and volume
(H1) were identified prior to the field survey using pre-existing long-term data on wood ant
populations in the North York Moors (Procter *et al.*, 2015; see Supplementary Methods S.1 and
Figure S.6). A ride was defined as a linear feature causing a break in the canopy at least 5m wide,
starting and ending where it intersected other rides or reached the end of the forest. Examples of rides
included linear canopy gaps due to roads and footpaths, as well as gaps caused by forest management
such as areas of row thinning (strips of trees harvested at intervals, leaving most of the crop standing)
or firebreaks. We measured the length of the wood ant-occupied portion of the ride from the nest
closest to the source population to the nest furthest from it. The population margins and the source
were identified using long-term data on the wood ant populations in the North York Moors (Procter,
2016; Holgate, 2021; Podesta, 2023). For rides with nests along their entire length, this measurement
equalled the total length of the ride. The total length of all of the rides surveyed was 4133 m. When
mapping the margins, a margin was defined as the last occupied area after which no nest was
encountered for 200 m; this is far greater than typical dispersal distance for this species, consistent
with the long-term data on these populations (Procter *et al.*, 2015). Rides were considered to have
ended if the ride entered a stand of different tree species, a stand of different age/tree height, or
rounded a corner (ride bearing changes by more than 15 degrees). We measured the length and

bearing of the ride using the *Measure tool* in ArcGIS Pro version 2.3.3, connecting the first and last
nest on the ride.

Selecting rides for calculating rate of expansion

In order to determine whether ride bearing affects wood ant dispersal, we required a subset of rides at
the population margins, where population margin expansion could have occurred during our study
period (since 2011) to calculate the average yearly population margins expansion along rides. In
addition to the selection criteria in the previous section, margin rides were also required to meet the
following criteria:

1. Have had potential for population expansion between 2011 and 2022 i.e., must have
uncolonized length that does not extend into unsuitable habitat like open farmland.
2. Expansion has been unidirectional, from a single point of origin, to exclude situations where
sections of the ride has been colonised from two origins (due to intersecting two rides with
ant presence), as this might inflate the measures of margins expansion on this ride.
3. Expansion has not been interrupted by factors such as areas of forest on the ride being
harvested during the period for which we have data, causing the ride to become unsuitable for
wood ants. In cases where the expansion had been interrupted, we used the data up to the
interruption. For example, a ride that was surveyed from 2011-21, but that was clear-felled in
2019, can provide us with an eight-year average expansion rate rather than ten.

Suitable rides were identified from three forests within the North York Moors, totalling 50 rides, 29 of
which were on the margins and suitable for testing margins expansion, each with record of the bearing
of the ride, length, mean width, the number of nests on each side of the ride. For the 29 rides on the
margins, we also had two measures of the rate of margin change calculated from long-term population
data: abundance change (nests/year) and distance change (m/year). One of the rides on the margins
was also at the true edge of a forest stand and therefore the ride had no measurable width and only one
side on which nests could occur. This ride was dropped from analyses that incorporated side or ride
width as variables.

Nest abundance and volume measurements

For each ride, the surveyor walked down one side of the ride and up the other, recording nest presence
and volume. Nests more than 5m from the edge were excluded to ensure all surveyed nests were
directly using the ride canopy gap. In areas where the understory was relatively open, spotting nests
up to 5m from the edge did not present a challenge but in areas where the understory was dense, some
nests were difficult to spot. Where the understory was too thick to confidently spot nests up to 5m
from the edge, a second transect was walked, parallel to the first, 2.5 m from the edge (behind
obscuring vegetation) to minimise the risk of missing nests. At each nest, we recorded the location
(latitude and longitude) and on which side of the ride (east or west/north or south) the nest was, giving

each nest a unique ID code to prevent recounting. Additionally, nest width, length and height were
measured to calculate nest volume as a proxy for number of ants (Chen and Robinson, 2013).

Ride details were also recorded: these included ride width (the shortest straight line distance between
sides of the ride where the canopy is directly overhead and excluding the understory of bushes and
seedlings), providing a distance between the crowns of trees, and canopy height (measured using
Toolbox android app calibrated against a gun clinometer) on both sides of the ride and ride bearing
(expressed as degrees east or west of north).　We measured each of these ride variables at the
beginning, midpoint and end of each ride and averaged them across the three to provide ride
characteristics. Latitude and longitude were recorded at the start and end of each ride so that the
length of the ride could be measured retrospectively using ArcGIS Pro version 2.3.3.

Calculating rate of margins expansion along rides

From the long-term data, we calculated two measures of margins change a) distance change and b)
number of nests. The distance the margin changed along a ride was the distance in metres between the
location of the last nest along a ride recorded in the oldest long-term data and the location of the last
nest on the same ride surveyed in 2022. The change in the number of nests was defined as the number
of nests between the last nest at the date of the earliest survey in the long-term data and the last nest
along the ride in the 2022 survey. Although both rate of distance change and the rate of nest
abundance change can be calculated from this data, rate of nest abundance change is a less reliable
measure of expansion than the rate of distance change for two reasons. Firstly, there is a five-year
interval between measurements in the long-term data and while all population margins were mapped
at each survey date, the entire population was not re-mapped on each survey (Procter *et al.*, 2015).
This means that, in many cases, we have no evidence of continuous occupation, nor of the number of
nests that may have been established and subsequently abandoned in the intervening period. Secondly,
these populations of *F. lugubris* are polydomous, a life history trait whereby a single 'colony' builds
and occupies many nest mounds, which share workers and food resources for mutual support. As a
result, in areas where the wood ants have recently colonised, some small nests may be abandoned and
the occupying ants absorbed into neighbouring nests in the polydomous network (Burns *et al.*, 2020)
with the effect that the actual numbers of nests may fluctuate. On the other hand, the predicted rate of
distance change was low, owing to the slow dispersal of this species (Procter *et al.*, 2015) whereas we
anticipated that the change in the number of nests would be large and more easily detectable. Note
also that the polydomous social organisation of this population means that new nests being established
along rides do not necessarily represent new colonies, but rather more likely are an expansion of
existing colony networks, although these nests may later become independent.

Statistical methods

Modelling the relationship between ride characteristics and population measures

To test H1 (i.e., the effect of ride characteristics (bearing, width, and side) on the abundance and relative volume of nests) we fitted generalised additive models using the mgcv package version 1.8-41 (Wood, 2011) in R (version 4.2.1). Because our orientation data were circular and we could not make assumptions about the shape of the relationship between orientation and our wood ant population measures, the additional flexibility of a GAM was preferred over a GLM. We fitted models including all combinations of the ride characteristics (bearing, width, and side), including site and ride ID as random effects, and used the minimum AIC to select the most parsimonious model for each response variable (Supplementary Methods S.2). Because of the circular nature of ride bearing (-90^{o} and 90^{o} relative to north are equivalent), we included ride bearing smoothing term as a cyclic p-spline, whereas mean width was included as a p-spline only. We plotted response vs. fitted values, fitted vs. observed and histograms of the residuals to inspect the model fit, and predicted from the models for visualisation. The GAMs provided estimations of degrees of freedom. For nest abundance, cyclic p-splines were fitted for both east and west sides of rides from predicted values of a point count generalised additive model using ride bearing as a predictor offset by ride length, to account for the zero inflation of count data, with the formula nest abundance $\approx$ side + s(bearing) + s(width) of the negative binomial family (Link function = log, offset = log(length). For nest volume, the same approach was used, with the formula volume per nest $\approx$ side + s(bearing) + s(width). For all our models, we calculated deviance explained, as deviance explained is favoured over R^2 for non-Gaussian models (Table 1, model 1; the nest abundance model) as a more robust measure of explanatory power (Wood, 2006).

Pooling bearing into categories for margins growth analysis

Our final dataset resulted in 29 rides available to test H2 (rate of margin change). The bearing data that we collected was continuous, and this was used to fit the GAM model for margins growth as described above. Inspection of the data, however, suggested a bigger difference between rides bearing near the cardinal points (N, S, E, W) and rides bearing near the intercardinals (NE, SW etc.) than between N-S and E-W rides. This suggests that, to test pooled bearing, bins should be E-W +/- 22.5°, N-S +/- 22.5°, and intercardinal +/- 22.5° (Figure S.1.), in order to appropriately capture the variability in the data. This variability has sound ecological justification: assuming that there is a minimum threshold of daily sunlight exposure for nesting sites to be viable for the wood ant nests to persist, we can be sure that this threshold lies somewhere below 50% of summer day lengths, as both sides of N-S rides are often utilised by ants. This threshold is also greater than 0% of the daily hours of sunlight because wood ants were often entirely absent from the south side of E-W rides. As such, going around the compass from north, the hours of available light on the north side will gradually increase from 50% up to nearly 100% after 90^{o}. Meanwhile, the south side will decrease from 50% to

273 nearly 0%, passing below the threshold for wood ants to utilise potential nesting sites somewhere

274 between. This reduction in potential nesting sites and the total availability of sunlight on the ride will

275 have negative effects on wood ant populations and result in slower population growth leading to

276 slower margins expansion.

Results

Summary of results

We found that nests were more abundant on the north sides of rides and varied considerably with ride

orientation; more nests were found, per metre, on rides that ran E-W or N-S and fewer on rides that

were oriented NE-SW or NW-SE (intercardinal). Furthermore, we found that the rate (number of

nests) and the distance of dispersal of wood ant nests was lowest along rides intercardinal rides, and

three times greater on cardinal rides. On the other hand, no consistent effect was found of ride

orientation on nest volume.

Ride bearing and nest aspect distribution

286 A total of 371 nests were recorded along all studied rides. Ride orientation covered the full range of

287 possible bearings but rides approximately 15-30° west of north and 70-80° east of north were more

288 abundant than other orientations (Figure 1). The aspect (direction of the long face/lower edge) of nest

289 mounds was overwhelmingly south-facing (Figure 1) with a circular mean of $191.47°$, significantly

290 different from a uniform distribution (Rayleigh's test, r = 197.47, var = 0.46, p < 0.0001).

Does the orientation of canopy gaps affect wood ant nest abundance and volume?

The most parsimonious model of nest abundance included only side, bearing and width (Figure 2;

Table 1). There was a significant interaction between ride bearing, and the side of the ride on which

nests are most abundant (Figure 2), with nests being most abundant on the East sides of rides that

have bearings around the NW-SE axis and most abundant on the West side of rides that have bearings

around the NE-SW axis. For rides that have bearings on the NS axis, no difference in abundance

between sides was seen. Overall, the number of nests per side was compensated by the other (Figure

2): when the east side has few nests (at bearings that cause the east side to be shaded), the west side

has more and vice-versa, so the total sum of nests per ride varied little with bearing. Ride width did

not have a significant effect on nest abundance (Table 1).

Figure 1. Left: The frequency of rides at different bearings was determined by the availability of
suitable rides at the field sites; ride bearings include most possible directions without excessively
favouring one direction. Right: The aspect (direction of the lowest side of the nest) of nests was most
commonly just west of south (180-225 degrees).

Table 1. Generalised additive models to predict several wood ant population measures using ride characteristics. Models 1 and 2 included the side of the ride that nests were on as a 'by' term (fitting the two sides as separate splines) so that we could test the significance of each smooths (difference from line with gradient of 0) for data from the east side of rides and the west, and test the smooths against one another (East:West). Both R^2 and deviance explained are included, as deviance explained is preferred for the non-Gaussian model 1. Expanded table in Supplementary Material (Table S.1.).

Model	Dependant variable	'By' interaction	Smoothing term		Effective degrees of freedom	Test statistic (F unless specified)	P-Value	N	R^2 (Deviance explained)	Notes
1	Nest abundance	Side	Ride Bearing	Side = East	4	17.741 (X^2)	0.001	98	0.054 (31%)	Nest abundance offset by ride length (Figure 2)
				Side = West	4	16.883 (X^2)	0.002			
				East:West	4	17.75 (X^2)	0.001			
			Ride Width		1.142	0.407 (X^2)	0.557			
2	Mean nest volume	Side	Ride Bearing	Side = East	5	3.476	0.008	69	0.267 (40.5%)	(Figure 2)
				Side = West	5	1.059	0.393			
				East:West	5	0.877	0.502			
			Ride Width		1.823	5.487	0.006			
3	Margins nest abundance change per year	None	Ride Bearing		4	4.443	0.082	28	0.163 (41.1%)	(Figure 4)
			Ride Width		4	2.061	0.126			
4	Margins distance change per year	None	Ride Bearing		4	1.934	0.148	28	0.256 (50.8%)	(Figure 4)
			Ride Width		4	1.011	0.428			
			Site		1.154	1.53	0.098			

Figure 2. Left: the total volume of all nests on each side of rides of different bearings. The lines of predicted values from a generalised additive model (GAM; $R^2 = 0.267$, deviance explained = 40.5%) were plotted for values of ride bearing (lines) alongside the original data (points) and 95%CIs (shaded areas). Although the east side smooth was significantly different from a line at the intercept of gradient = 0 (p < 0.01), there was no significant difference between the east and west smooths and no clear pattern is discernible from the plot. Mean width was also significant as a smoothing term (Figure 3; Table 1). Right: the number of nests per side of a ride (E or W) for rides of a range of bearings. The points represent the real data, the lines are predicted values and the shaded areas are 95%CIs. Both east and west smooths were significantly different from 0 (p < 0.005) and significantly different from each other (p = 0.001).

The most parsimonious model of nest volume included site, bearing and width as smoothing terms,

however the inclusion of site only marginally improved explanatory power and did not justify a more

complex model, so was dropped (Table S.1; Figure S.3). In both models, the ride bearing significantly

affected volume only for nests on the East side, and relationship between ride bearing and nest

volume on the East and West sides did not differ significantly from one another (Figure 2; Table 1).

The asymmetry of this result, and the lack of difference between the two sides, indicates that overall,

ride bearing does not effectively predict mean nest volume in this model. In contrast, mean ride width

was significant as a smoothing term ($p < 0.01$; Table 1) and a plot of model predictions (Figure 3)

shows that the mean volume of nests is lower on wider rides but does not continue to decrease as ride

width exceeds 8 m. A relatively small number of large nests on rides narrower 6m may be major

contributors to this effect; however, this small number of nests were distributed across all three sites.

Does the orientation of canopy gaps affect wood ant dispersal rate?

We used two measures of population margins growth: the change in the number of nests at the margin

and the distance that the margins moved per year over the study period. The model that best explained

the change in nest number per year included ride bearing and width (Figure 4; Table 1). Neither term

was below the significance threshold of 0.05 (ride bearing $p = 0.082$, width $p = 0.126$) and the model

explained only approximately 15% of the variation. Overall, we see a similar shaped pattern in the

rate of nest number change (Figure 4) as we saw for nest abundance on each side of rides (Figure 2),

with minima for rides bearing approximately NW-SE and NE-SW (intercardinal), though not reaching

statistical significance. While this model has poor predictive power, it does suggest that bearing may

play some role in the rate of change in nest number.

Figure 3. Generalised additive model (GAM) with the formula volume per nest ≈ s(bearing) + s(width), R^2 = 0.18, deviance explained = 28.7% were plotted for values of width (line) alongside the original data (points) and 95%CIs (shaded area), with the points representing field data. The smooth of mean width was significantly different from the intercept ($p < 0.01$), whereas ride bearing alone (i.e. not split by side) was not.

332
Figure 4. Upper: a generalised additive model (GAM) was fit with the formula nest abundance
change per year ≈ s(bearing) + s(width), R^2 = 0.163, deviance explained = 41.1% and the predictions
for values of ride bearing were plotted (solid line) with 95%CIs (shaded areas). Ride bearing was not
significant as a smoothing term and the model has little explanatory power. Ride width is not
correlated with bearing. Including width as a smoothing term lower predicted values than the
observed values due to the fixing of ride width at the mean value for non-Gaussian data. Lower: A
generalised additive model was fit with the formula margins change ≈ s(bearing) + s(width) + s(site),
R^2 = 0.256, deviance explained = 50.8%. Empirical data (points), model predictions (lines) and
95%CIs (shaded areas) are shown. None of the smoothing terms were significant, but site was a
significant random effect and its inclusion resulted in the most parsimonious model.

The median rate of nest abundance change was greater on rides with cardinal orientations than those
with intercardinal ones (approximately 0.5 nests/year; Figure 5), but this difference did not reach
statistical significance (Welch's *t*-test, $t = 2.02$, df = 21.37, p = 0.056). A further split in the data was
tested to N-S, E-W, NW-SE and NE-SW against each other, but no statistically significant difference
was found between them for either variable (Figures S.4 and S.5).

The distance that the margins moved per year over the study period was also included as a response
variable in a GAM and the best model includes site, ride bearing and ride as smoothing terms (Figure
4; Table 1). This model performed better than the nest number change model ($R^2 = 0.256$) and the
overall pattern was the same (minima on rides near intercardinal directions), but none of the
smoothing terms were significant (p > 0.05; Table 1). Model predictions showed that rides in Cropton
Forest experienced more wood ant population growth than either of the other two sites.

Overall, rides running in intercardinal resulted in lower population margins growth (Figure 5). We
found that the rides that bear in cardinal directions experienced three times the annual margins
expansion than intercardinal ones (15m/year compared to 5m/year; Welch's *t*-test, $t = 2.122$, df =
22.497, p = 0.045; Figure 5; Table 2) and the effect size was medium (Hedges' *g* = 0.72).

Table 2. Dispersal range was measured as rate of change of the number of nests at the margin, and distance of the final nest along a ride from the original (2011) population extent to the final (2022) population extent. We made compared rides of varied orientation relative to north; cardinal (N-S and E-W +/- 22.5°) and intercardinal (NW-SE and NE-SW+/- 22.5°) bearings.

Measure	Bins	Tests	Degrees of freedom	Test statistic	P-value	Effect size (Hedges' g)
Rate of change in nest number at margins (Figure 5)	2	Welch's *t*-test	21.37	t = 2.02	0.056	0.68
Rate of distance change at margins (Figure 5)	2	Welch's *t*-test	22.50	t = 2.12	0.045	0.72

360   **Figure 5.** Ride data for change in nest number (A; Welch's *t*-test, t = 2.02, p = 0.056), and distance of
361   margin change (B; Welch's *t*-test, t = 2.12, p = 0.045), comparing cardinal (22.5° +/- N, S, E or W)
362   and intercardinal (22.5° +/- NE, NW, SE or SW) bearings (Table 2 for statistical details).

Discussion

Summary of results

We aimed to determine the effect of anthropogenic canopy gap characteristics on populations of an edge specialist, the northern hairy wood ant *Formica lugubris*. Our results clearly show that the orientation and width of linear canopy gaps affect *F. lugubris* distribution and dispersal, shaping their population dynamics. Nests overwhelmingly had a south facing aspect, as would be expected if nests are constructed to present a large surface area to the sun and maximise insolation. Nests were more abundant on the northern side of rides for all orientations, as predicted, but orientation did not similarly affect nest size in a consistent way. In general, narrower rides had larger nests, while wider rides were populated by smaller nests, consistent with predictions that nests receiving less insolation are larger (Chen and Robinson 2014). Finally, we found that, as predicted, the orientation of rides affected *F. lugubris* dispersal. Over the preceding 10 years, populations of *F. lugubris* were both more numerous and dispersed more quickly along rides that run in cardinal directions (N, E, S, W) than ones that run in intercardinal directions (NE, SE, SW, NW).

The importance of edge characteristics

We showed that the orientation of edge habitats relative to north is an important factor in edge quality for *Formica lugubris* and is sufficient to have measurable effect on nest abundance and population margin expansion. Ride width is also a factor in edge quality for these wood ants; we found larger nests in narrower rides. In general, larger nests are found in shadier areas because larger nests are better able to thermoregulate (Chen and Robinson, 2014). The strong south-facing bias in nest aspect we found confirms that our other results are likely to driven by insolation, as sunlight is important for many aspects of wood ant biology. Nests lacking direct sunlight, especially small nests, may initiate foraging later (Rosengren *et al.*, 1987), have slower brood maturation (Kadochová and Frouz, 2013) and therefore potentially slower colony growth.

Dispersal along rides

In addition to ride bearing affecting wood ant population density (nest volume and abundance), it also affects the ability of the wood ants to disperse through these plantation forests, with most rapid dispersal along rides oriented along the cardinal bearings (N-S or E-W). Due to the generally slow dispersal of *F. lugubris*, the cardinal ride dispersal distance of 15 m/year is equivalent to three times the rate of dispersal in intercardinal rides (5 m/year). We hypothesise that the slower population margin expansion observed for rides with bearings in intercardinal directions is due to a minimum threshold of insolation for newly established nests of *F. lugubris* to be viable. Above this threshold, a wood ant nest's outcomes continue to improve. We suggest that the south side of a ride with an intercardinal bearing receives insolation below this minimum threshold, due to being shaded for most of the day by the trees on the southern side, while the north side will receive less insolation than the

north side of a ride running east-west. This means nests on these bearings are in a 'worst of both
worlds' environment resulting in fewer individuals and a reduced ability to form new nests in the
polydomous network. Polydomy allows for food and labour sharing between nests that can subsidise a
new nest while it grows big enough to establish foraging (Ellis and Robinson, 2015). Newly
established nests in shadier areas may require extra support from neighbouring nests (Rosengren *et*
al., 1987), so being further from the centre of this network may leave new nests more vulnerable to
abandonment. The combination of lower rates of nest foundation and a higher failure rate for nests
that have been established further from the parental nests, could explain the reduced margins growth
that we measured in shadier, intercardinal rides.

Management implications

Given that wood ants increase biodiversity in plantation forests and may provide additional pest
control benefits (Laine and Niemela, 1980; Karhu and Neuvonen 1998; Zingg *et al.*, 2018), the
results that we present here may be useful in planning management of plantations. Wood ants can
reduce defoliating pest burden (Karhu and Neuvonen 1998), influence tree growth via their effects on
aphids (Kilpeläinen *et al.*, 2009), compete with other insectivores (Haemig, 1992), and change the
spatial distribution of soil resources (Podesta, 2023), so the effects of ride characteristics on their
dispersal have major implications for conservation and forest management. When planning new areas
of forestry adjacent to existing populations of wood ants, or in areas where translocation of wood
ants' nests is being considered (Nielsen *et al.*, 2018), we would recommend cutting firebreaks, access
tracks etc. in cardinal directions, to maximise the ability of wood ants to disperse. The width of the
ride should also be considered when managing forest to promote species richness or dispersal.
Previous work has suggested that canopy gaps should be at least 15 m wide in order to harbour
species that would be shaded out in the forest interior (Oxbrough *et al.,* 2006; Smith *et al.,* 2007), and
our data suggest that wood ants similarly have a minimum threshold of insolation (also impacted by
ride width) that allows them to thrive and disperse.

Wider ecological implications

The work presented here adds to a body of research showing that linear canopy gaps in plantation
forests have important implications for forest dwelling species. The characteristics of rides, such as
width, affect the diversity or abundance of butterflies (Greatorex-Davies *et al.*, 1993), spiders
(Oxbrough *et al.,* 2006; Carter *et al.,* 1991), ground beetles (Carter *et al.,* 1991), vascular plants
(Sparks *et al.,* 1996; Smith *et al.,* 2007), and bryophytes (Smith *et al.,* 2007). We have shown that, in
addition to the effects that ride characteristics can have on the diversity and abundance of forest
species in plantation, the orientation of rides can influence the dispersal ability of an edge specialist,
opening up the question of whether other species with similar habitat requirements might be similarly
affected.

Managing plantations in a manner favourable to wood ant dispersal (by cutting firebreaks and other
necessary linear gaps along cardinal directions) could be beneficial to other species and biodiversity
in economically productive forests in two ways. Firstly, the presence of wood ants themselves can be
beneficial to other species. The nests of wood ants are home to many myrmecophilous species, with
over 100 myrmecophile species known from wood ants' nests (Robinson *et al*., 2016; Parmentier *et*
al., 2014). Additionally, wood ants can play an important role in the dispersal of seed, including some
declining and charismatic wildflowers like the endangered *Melampyrum cristatum* (crested cow-
wheat) which is dispersed by wood ants and is likely range-limited by the availability of dispersers
(Stachnowicz, 2013). Secondly, other species with similar habitat requirements and dispersal ability to
the wood ants would benefit from the same rides as dispersal corridors as the wood ants. Species that
might benefit from the same corridor design as wood ants could include wildflowers that likewise
require direct sunlight but benefit from other conditions provided by the forest edge such as protection
from wind, acidic soil or edge related nutrient influx (Weathers *et al*., 2001).

On the other hand, species movement can be detrimental; there are well documented negative
economic (Cuthbert *et al.*, 2022) and environmental (Pyšek *et al.*, 2020) effects of species invasions.
Edge specialists with the potential to benefit from improved dispersal due to edge orientation may
also be pests, for example, the larvae of *Gilpinia virens,* a pine sawfly and common pest of *Pinus*
sylvestris, are found in higher numbers in stands with relatively open canopies (Gawęda and Grodzki,
2020). As a result, pests such as *G. virens* may also benefit from the presence of canopy gaps cut in
favourable, cardinal directions, although it should be noted that it does not necessarily follow that a
preference for canopy gaps will result in the same response to different gap characteristics as we see
in wood ants.

Conclusions

Our work clearly shows the potential of anthropogenic linear canopy gaps for promoting dispersal in
plantation forests. Using a species that favours woodland edges to allow access to both light and
resources, we demonstrate that the properties of linear canopy gaps influence distribution and
dispersal rates. In practical terms, this means that canopy gaps created during plantation forest
management, such as firebreaks, row-thinning, or access for logging machinery, can have the
secondary function of promoting the dispersal of edge specialist forest species, especially if
orientations near N-S or E-W are predominant. The control that forestry planners have over the
placement of firebreaks and other linear canopy gaps, thus allows the possibility of deliberately
making plantation forests more porous to certain species, to gain the ecosystem services they provide
(for example, the suppression of defoliating pests by wood ants) or to promote biodiversity. The
increase in demand for timber products and forested areas to reach carbon offset goals will likely lead
to an increase in the area and intensity of plantation forests in the future (McEwan et al., 2020). If
these economic and environmental goals to be achieved without detriment to biodiversity or resilience

to pests, care must be taken to ensure the development of forestry practices that allow wildlife to

proliferate in plantation forests, both for the ecosystem services they can provide and for the

biodiversity that they represent.

Acknowledgements

473 We thank Duncan Procter and Megan Holgate for additional data that contributed to the work and

474 Forest Research for site access permission. We also thank Imre Sándor Piross and Mark Hodson for

475 advice with the analysis and manuscript respectively, and Anne Oxbrough for feedback on the

476 original thesis chapter. The project was funded by the NERC ACCE2 DTP.

References

Bennett, A.F. (1990) 'Habitat corridors and the conservation of small mammals in a fragmented forest
environment', *Landscape Ecology*, 4(2–3), pp. 109–122.

Bernaschini, M.L., Valladares, G. and Salvo, A. (2020) 'Edge effects on insect–plant food webs:
assessing the influence of geographical orientation and microclimatic conditions', *Ecological*
Entomology, 45(4), pp. 806-820.

Brito, T.F. *et al.* (2017) 'Forest reserves and riparian corridors help maintain orchid bee
(Hymenoptera: Euglossini) communities in oil palm plantations in Brazil', *Apidologie*, 48(5), pp.
575–587.

Brunzel, S., Elligsen, H. and Frankl, R. (2004) 'Distribution of the Cinnabar moth *Tyria jacobaeae* L.
at landscape scale: Use of linear landscape structures in egg laying on larval hostplant exposures',
Landscape Ecology, 19(1), pp. 21-27.

Burns, D.D.R. *et al.* (2020) 'A longitudinal study of nest occupancy, trail networks and foraging in a
polydomous wood ant population', *Insectes Sociaux*, 67(3), pp. 419–427.

Carter, C. I. (1991). 'Paper 4: Ride Orientation and Invertebrate Activity', *Edge management in*
woodlands, 28, pp. 17-21.

Chen, Y.H. and Robinson, E.J.H. (2013) 'A comparison of mark-release-recapture methods for
estimating colony size in the wood ant *Formica lugubris*', *Insectes Sociaux*, 60(3), pp. 351–359.

Chen, Y.H. and Robinson, E.J.H. (2014) 'The relationship between canopy cover and colony size of
the wood ant *Formica lugubris*-implications for the thermal effects on a keystone ant species', *PLoS*
ONE, 9(12), e116113.

Cuthbert, R.N. *et al.* (2022) 'Are the "100 of the world's worst" invasive species also the costliest?',
Biological Invasions, 24(7), pp. 1895–1904.

Defra (2021) *The England Trees Action Plan 2021-2024*.

Domisch, T., Risch, A. and Robinson, E.J.H. (2016) 'Wood ant foraging and mutualism with aphids',
in J. Stockan and E.J.H. Robinson (eds.), *Wood ant ecology and conservation* Cambridge: Cambridge
University Press, pp. 145-176.

Ellis, S. and Robinson, E.J.H. (2014) 'Polydomy in red wood ants', *Insectes Sociaux*, 61(2), pp. 111–
122.

Ellis, S. and Robinson, E.J.H. (2015) 'The role of non-foraging nests in polydomous wood ant
colonies', *PLoS ONE*, 10(11), e0143901.

Ewers, R. M. and Didham, R. K. (2005) 'Confounding factors in the detection of species responses to
habitat fragmentation', *Biological Reviews*, 81(1), pp. 117-142.

Gawęda, P. and Grodzki, W. (2020) ' The influence of stand and habitat characteristics on the
occurrence of pine sawflies *Diprion pini* L. and *Gilpinia virens* (Klug) (Hymenoptera, Diprionidae) in
selected areas of northern Poland ', *Forest Research Papers*, 81(3), pp. 99-106.

Greatorex-Davies, J. N., Sparks, T. H., Hall, M. L., and Marss, R. H. (1993). 'The influence of shade
on butterflies in rides of coniferised lowland woods in southern England and implications for
conservation management', *Biological Conservation*, 63(1), 31-41.

Haemig, P.D. (1992) 'Competition between Ants and Birds in a Swedish Forest', *Oikos*, 65(3), pp.
479-483.

Holgate, M.K. (2021) 'The effects of land use change on the population expansion of a poor disperser,
the wood ant *Formica lugubris'*, PhD Thesis, University of York, York.

Kadochová, Š. and Frouz, J. (2013) 'Thermoregulation strategies in ants in comparison to other social
insects, with a focus on *Formica rufa'*, *F1000Research*, 2:280.

Karhu, K.J. and Neuvonen, S. (1998) 'Wood ants and a geometrid defoliator of birch: Predation
outweighs beneficial effects through the host plant', *Oecologia*, 113(4), pp. 509–516.

Kilpeläinen, J., et al., Does the mutualism between wood ants (*Formica rufa* group) and Cinara aphids
affect Norway spruce growth? *Forest Ecology and Management*, 2009. 257(1): p. 238-243

Laine, K.J. and Niemela, P. (1980) 'The Influence of Ants on the Survival of Mountain Birches
During an *Oporinia autumnata* (Lep., Geometridae) Outbreak', *Oecologia*, 47, pp. 39–42.

Lamb, E.G. and Mallik, A.U. (2003) 'Plant species traits across a riparian-zone/forest ecotone',
Journal of Vegetation Science, 14(6), pp. 853-858.

Lenoir, J. and Svenning, J.C. (2015) 'Climate-related range shifts - a global multidimensional
synthesis and new research directions', *Ecography*, 38(1), pp. 15-28.

Liira, J. and Paal, T. (2013) 'Do forest-dwelling plant species disperse along landscape corridors?',
Plant Ecology, 214(3), pp. 455-470.

Loyn, R.H. (1987) 'Effects of patch area and habitat on bird abundances, species numbers and tree
health in fragmented Victoria forests.', A. Saunders, G.W. Arnold, A.A. Burbidge, and A.J.M.
Hopkins (eds.) *Nature conservation: the role of remnants of native vegetation* pp. 65-77.

Maeder, A. *et al.* (2016) 'Wood ant reproductive biology and social systems', in J.A. Stockan and
E.J.H.E. Robinson (eds) *Wood Ant Ecology and Conservation*. Cambridge: Cambridge University
Press, pp. 37–50.

Magura, T., Lövei, G.L. and Tóthmérész, B. (2017) 'Edge responses are different in edges under
natural versus anthropogenic influence: a meta-analysis using ground beetles', *Ecology and Evolution*,
7(3), pp. 1009-1017.

Major, R.E., Christie, F.J. and Gowing, G. (2001) 'Influence of remnant and landscape attributes on
Australian woodland bird communities', *Biological Conservation*, 102(1), pp. 47-66.

McEwan, A. *et al.* (2020) 'Past, present and future of industrial plantation forestry and implication on
future timber harvesting technology', *Journal of Forestry Research*. 31, pp. 339–351.

Mendenhall, C.D. *et al.* (2014) 'Predicting biodiversity change and averting collapse in agricultural
landscapes', *Nature*, 509, pp. 213–217 .

Nielsen, J.S. *et al.* (2018) 'Experiences in transplanting wood ants into plantations for integrated pest
management', *Sociobiology*, 65(3), pp. 403–414.

Oxbrough, A. G., Gittings, T., O'Halloran, J., Giller, P. S., and Kelly, T. C. (2006). 'The influence of
open space on ground-dwelling spider assemblages within plantation forests', *Forest Ecology and
Management*, 237(1-3), 404-417.

Parmentier, T., W. Dekoninck and T. Wenseleers (2014) 'A highly diverse microcosm in a hostile
world: a review on the associates of red wood ants (*Formica rufa* group)', *Insectes Sociaux* 61(3):
229-237.

Podesta, J. (2023). Interactions between the wood ant *Formica lugubris* and plantation forests: how
ants affect soil properties and how forest management affects the dispersal of an ecosystem engineer.
PhD thesis, University of York, York.

Procter, D.S. (2016) 'The effects of forest cover change and polydomous colony organisation on the
wood ant *Formica lugubris*', PhD Thesis, University of York, York.

Procter, D.S. *et al.* (2015) 'Do non-native conifer plantations provide benefits for a native forest
specialist, the wood ant *Formica lugubris*?', *Forest Ecology and Management*, 357, pp. 22–32.

Pyšek, P. *et al.* (2020) 'Scientists' warning on invasive alien species', *Biological Reviews*, 95(6), pp.
1511-1534.

Ries, L. and Sisk, T.D. (2010) 'What is an edge species? The implications of sensitivity to habitat
edges', *Oikos*, 119(10), pp. 1636-1642.

Robinson, E.J.H., Stockan, J.A. and Iason, G.R. (2016) 'Wood ants and their interaction with other
organisms', in J.A. Stockan and E.J.H. Robinson (eds.) *Wood Ant Ecology and Conservation*.
Cambridge: Cambridge University Press, pp. 177–199.

Roberts, D., Ciuti, S., Barber, Q. E., Willier, C., and Nielsen, S. E. (2018) 'Accelerated seed dispersal
along linear disturbances in the Canadian oil sands region', *Scientific Reports*, 8(1), 4828.

Rosengren, R. *et al.* (1987) 'Phenology and causation of nest heating and thermoregulation in red
wood ants of the *Formica rufa* group studied in coniferous forest habitats in southern Finland.',
Annales Zoologici Fennici, 24(2), pp. 147–155.

Ruete, A. *et al.* (2017) 'Contrasting long-term effects of transient anthropogenic edges and forest
fragment size on generalist and specialist deadwood-dwelling fungi', *Journal of Applied Ecology*,
54(4), pp. 1142–1151.

Shepard, D.B., Kuhns, A.R., Dreslik, M.J., and Phillips, C.A. (2008). Roads as barriers to animal
movement in fragmented landscapes. *Animal conservation*, 11(4), 288-296.

Smith, G. F., Iremonger, S., Kelly, D. L., O'Donoghue, S., and Mitchell, F. J. (2007). Enhancing
vegetation diversity in glades, rides and roads in plantation forests. *Biological Conservation*, 136(2),
pp. 283-294.

Sparks, T. H., Greatorex-Davies, J. N., Mountford, J. O., Hall, M. L., and Marrs, R. H. (1996) 'The
effects of shade on the plant communities of rides in plantation woodland and implications for
butterfly conservation', *Forest Ecology and Management*, 80(1-3), pp. 197-207.

Stachnowicz, W. (2013) 'Melampyrum Cristatum L. – A Rare River Corridor Plant in Wielkopolska
and Poland', *Biodiversity Research and Conservation*, 32(1) pp. 29-44.

Suárez-Esteban, A. *et al.* (2016) 'Can anthropogenic linear gaps increase plant abundance and
diversity?', *Landscape Ecology*, 31, pp. 721-729.

Sudd, J.H. *et al.* (1977) 'The distribution of wood☐ants (*Formica lugubris* Zetterstedt) in a northern
English forest', *Ecological Entomology*, 2(4), pp. 301-313.

Taylor, R.S., Oldland, J.M. and Clarke, M.F. (2008) 'Edge geometry influences patch-level habitat
use by an edge specialist in south-eastern Australia', *Landscape Ecology*, 23(4), pp. 377–389.

van Schalkwyk, J. *et al.* (2022) 'Corridor width and orientation are complementary design variables
for butterflies in conservation corridors', *Landscape Ecology*, 37, pp. 2535-2549.

Weathers, K.C., Cadenasso, M.L. and Pickett, S.T.A. (2001) 'Forest edges as nutrient and pollutant
concentrators: Potential synergisms between fragmentation, forest canopies, and the atmosphere',
Conservation Biology, 15(6), pp. 1506-1514.

Williamson, S. and Hirth, D. (1985) 'An Evaluation of Edge Use by White-tailed Deer', *Wildlife
Society Bulletin*, 13(3), pp. 252-257.

Wood, S.N. (2006) 'Generalized Additive Models: An Introduction with R' (2nd ed.). Chapman and
Hall/CRC.

Wood, S.N. (2011) 'Fast stable restricted maximum likelihood and marginal likelihood estimation of
semiparametric generalized linear models', *Journal of the Royal Statistical Society (B)*, 73(1), pp. 3–
36.

Zingg, S. *et al.* (2018) 'The negative effect of wood ant presence on tick abundance', *Parasites and
Vectors*, 11(1), 164.