Mechanistic computational modeling of sFLT1 secretion dynamics

Amy Gill[1,2], Karina Kinghorn[3,4], Victoria L. Bautch[3,4,5], Feilim Mac Gabhann[1,2]

[1] Institute for Computational Medicine, Johns Hopkins University, Baltimore, MD, USA
[2] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD, USA
[3] Department of Biology, University of North Carolina, Chapel Hill, NC, USA
[4] Curriculum in Cell Biology and Physiology, University of North Carolina, Chapel Hill, NC, USA
[5] McAllister Heart Institute, University of North Carolina, Chapel Hill, NC, USA

Abstract

Constitutively secreted by endothelial cells, soluble FLT1 (sFLT1 or sVEGFR1) binds and sequesters extracellular vascular endothelial growth factors (VEGF), thereby reducing VEGF binding to VEGF receptor tyrosine kinases and their downstream signaling. In doing so, sFLT1 plays an important role in vascular development and in the patterning of new blood vessels in angiogenesis. Here, we develop multiple mechanistic models of sFLT1 secretion and identify a minimal mechanistic model that recapitulates key qualitative and quantitative features of temporal experimental datasets of sFLT1 secretion from multiple studies. We show that the experimental data on sFLT1 secretion is best represented by a delay differential equation (DDE) system including a maturation term, reflecting the time required between synthesis and secretion. Using optimization to identify appropriate values for the key mechanistic parameters in the model, we show that two model parameters (extracellular degradation rate constant and maturation time) are very strongly constrained by the experimental data, and that the remaining parameters are related by two strongly constrained constants. Thus, only one degree of freedom remains, and measurements of the intracellular levels of sFLT1 would fix the remaining parameters. Comparison between simulation predictions and additional experimental data of the outcomes of chemical inhibitors and genetic perturbations suggest that intermediate values of the secretion rate constant best match the simulation with experiments, which would completely constrain the model. However, some of the inhibitors tested produce results that cannot be reproduced by the model simulations, suggesting that additional mechanisms not included here are required to explain those inhibitors. Overall, the model reproduces most available experimental data and suggests targets for further quantitative investigation of the sFLT1 system.

Introduction

Secreted proteins, which represent 15% of the human genome and 9% of the human proteome, are essential local and systemic signaling factors in multicellular systems.[1] Most proteins destined for the extracellular space, or for intracellular locations other than the cytoplasm and nucleus, mature via the secretory pathway, in which new proteins are cotranslationally translocated into the endoplasmic reticulum (ER), folded, transported to the Golgi apparatus, post-translationally modified, sorted, and trafficked by vesicles to target compartments.[2] Many proteins synthesized through the secretory pathway are released by exocytosis, while others reside and function mainly in intracellular compartments (such as the Golgi or endosomes).[3–5] Many proteins known to function extracellularly or at the plasma membrane also have biologically meaningful interactions in intracellular compartments, including receptor tyrosine kinases.[6–10] Although all secreted proteins begin as intracellular proteins, intracellular roles and dynamics of extracellularly secreted proteins are generally understudied.

Here, we explore the secretion of soluble FLT1 (sFLT1, sVEGFR1), a key negative regulator of the vascular endothelial growth factors (VEGF), a family of cytokines that promote angiogenesis. sFLT1 is an alternatively spliced truncated isoform of VEGF Receptor 1 (VEGFR1 or FLT1) that retains the extracellular domains for binding ligands (including VEGFA, VEGFB, and placental growth factor (PlGF)) and heparan sulfate proteoglycans (HSPGs), but lacks transmembrane or intracellular tyrosine kinase domains, yielding a secreted, non-membrane anchored, and non-signaling receptor.[11,12] Constitutively secreted by endothelial cells,[13] sFLT1 binds and sequesters most extracellular VEGF isoforms, thereby reducing binding of these VEGF isoforms to VEGF receptor tyrosine kinases and the downstream signaling.[11,14–16]

As a VEGF antagonist and antiangiogenic factor, sFLT1 prevents vascular overgrowth and is

essential for corneal avascularity.[17] This inhibitory role is also important during angiogenesis for

proper patterning of new blood vessel networks: sFLT1 from endothelial cells interacts

extracellularly with stromal VEGF to reinforce VEGF gradients extending away from existing

vasculature, guiding nascent vascular sprouts to reduce vascular overgrowth and

disorganization.[18–20]

Dysregulated sFLT1 production and secretion is a pathogenic factor in several disorders

including preeclampsia, a syndrome of endothelial dysfunction during pregnancy characterized

by hypertension, kidney damage, and significant maternal and fetal morbidity and mortality. In

preeclampsia, excessive sFLT1 production and secretion from the placenta leads to elevated

serum sFLT1, reducing free serum levels of VEGF and PlGF, inhibiting VEGF-mediated

vasodilation and angiogenesis, causing hypertension and damaging vasculature.[21] While we do

not focus on pathology here, a quantitative understanding of sFLT1 production and secretion

from endothelial cells could be extended in the future to simulate pathologies.

While the extracellular role of sFLT1 is well characterized, less is known about its

intracellular biology. Recent work suggests unexpectedly high intracellular sFLT1 levels in

endothelial cells and long intracellular residence time,[13,22] motivating a quantitative

characterization of sFLT1 trafficking and elucidation of the intracellular dynamics of sFLT1 and

its possible intracellular roles. Past studies also suggest the full-length membrane-integral FLT1

(mFLT1) protein resides largely intracellularly,[23] much of it in the Golgi, and it traffics to the

plasma membrane in a calcium-dependent manner.[8] sFLT1 may be retained in secretory

organelles by a similar mechanism to the larger mFLT1 isoform.

Mechanistic computational modeling represents biochemical molecular networks as systems of equations based on prior knowledge of regulatory interactions and parameterized with experimental data. These models allow virtual exploration of molecular networks while tracking components that may be difficult to measure experimentally, and are powerful tools for testing mechanistic hypotheses and screening new treatment approaches.[24] While many mechanistic models describe systems as sets of ordinary differential equations (ODEs), time delays between processes are common in biology, and recent work shows that delay differential equations (DDEs) can better represent the behavior of several common network motifs in biochemical reaction networks.[25] This study builds on our recent model of sFLT1 production and secretion in endothelial cells,[13] which we believe is the first mechanistic model to include intracellular sFLT1. Several previous models of the VEGF system have included the impact of sFLT1 secretion on extracellular sFLT1 and VEGF levels, including pharmacological-like distribution across tissues[26–29] and local gradients,[19] but did not explore the intracellular dynamics of sFLT1 secretion. In our recent work characterizing intracellular sFLT1 trafficking from protein synthesis to secretion,[13] we briefly introduced a model linking intracellular and extracellular sFLT1 pools through protein synthesis, secretion, and intracellular and extracellular degradation. We used this model to estimate a basal secretion rate of ~30,000 molecules per cell per hour in human umbilical vein endothelial cells (HUVEC), comparable to other secreted proteins, and to infer that Golgi trafficking is absolutely required for sFLT1 secretion.[13]

Here, we extend our previous work to develop multiple mechanistic models of sFLT1 secretion and to identify a minimal mechanistic model that recapitulates key qualitative and quantitative features of experimental sFLT1 time course datasets. We compare several ODE-based and DDE-based models to capture the dynamics of sFLT1 secretion, using experimental

100   data from multiple studies to parameterize the models and evaluate their relative performance.

101   We then compare simulation predictions of perturbation by secretion pathway inhibitors to the

102   outcomes of equivalent experiments.

₁₀₄ Methods

₁₀₅ **Models of sFLT1 production and secretion.** We constructed several mechanistic models to

₁₀₆ explore sFLT1 secretion dynamics, each of which consists of two coupled nonlinear differential

₁₀₇ equations describing the levels of sFLT1 in two locations: extracellular sFLT1 (*X*) that has been

₁₀₈ secreted out of the cell into conditioned media, and intracellular sFLT1 (*I*), a sum of immature

₁₀₉ and mature protein inside the cell, treating all intracellular locations (ER, Golgi, vesicles, etc.) as

₁₁₀ one well-mixed compartment. Molecular levels of *I* and *X* were tracked in units of number per

₁₁₁ cell (#/cell). We briefly introduced one example of these models in recent work;[13] here we

₁₁₂ provide additional detail on model construction and analysis, compare it to additional model

₁₁₃ formulations, and re-parameterize based on additional experimental data.

₁₁₄ **ODE-based model.** We created an initial model using ordinary differential equations (ODEs)

₁₁₅ (**Figure 1A**) to describe how the mechanistic processes impact the level of intracellular and

₁₁₆ extracellular sFLT1. The equations for intracellular and extracellular sFLT1 include rates for

₁₁₇ four mechanistic processes:

₁₁₈ ● *production*, the synthesis of new intracellular protein ($\emptyset \rightarrow I$) characterized as a zeroth-

₁₁₉ order process with a rate constant α (when on) or 0 (when off), representing translation of

₁₂₀ intracellular protein and assuming no feedback dependence of sFLT1 production on

₁₂₁ current sFLT1 levels;

₁₂₂ ● *secretion*, the transport from the intracellular to extracellular space ($I \rightarrow X$) characterized as

₁₂₃ a first order process with rate constant β, representing exocytosis of mature sFLT1

₁₂₄ protein;

•   *intracellular degradation*, the depletion of intracellular sFLT1 ($I \rightarrow \emptyset$) characterized as a

first order process with rate constant γ, representing lysosomal degradation or other

proteolytic degradation mediated by intracellular factors;

•   *extracellular degradation*, the depletion of extracellular sFLT1 ($X \rightarrow \emptyset$) characterized as a

first order process with rate constant δ, representing proteolytic degradation mediated by

soluble, membrane-bound, or matrix-bound extracellular factors.

Thus, the following equations define the time evolution of the system:

132
$$\frac{dI(t)}{dt} = \alpha - \beta \cdot I(t) - \gamma \cdot I(t) \tag{Eq. 1}$$

133
$$\frac{dX(t)}{dt} = \beta \cdot I(t) - \delta \cdot X(t) \tag{Eq. 2}$$

134   **DDE-based model.** We converted our initial ODE model to a system of delay differential

135   equations (DDEs) (**Figure 2A**) by introducing a fixed time delay τ to the secretion and

136   intracellular degradation processes, representing a required protein maturation period before I

137   can be transported or degraded:

138
$$\frac{dI(t)}{dt} = \alpha - \beta \cdot I(t - \tau) - \gamma \cdot I(t - \tau) \tag{Eq. 3}$$

139
$$\frac{dX(t)}{dt} = \beta \cdot I(t - \tau) - \delta \cdot X(t) \tag{Eq. 4}$$

Candidate mechanistic models. We formulated eight distinct candidate models containing

different combinations of 3 processes:

● *maturation delay*, the fixed time delay τ described above affecting secretion and

intracellular degradation

● *internalization,* a first order transport reaction from the extracellular to intracellular space

($X{\rightarrow}I$) with rate constant ε

● *production decay*, an exponential decay $e^{-\kappa t}$ of production when transitioning from an

expression pulse to chase at *t=0*, with rate constant κ representing first order decay of

labeled amino acids; note this does not affect the constitutive secretion scenario or the

pulse segment of the pulse-chase scenario

Model equations were constructed in piecewise fashion, with different rate terms depending on

which processes were included in a given candidate model (**Supplemental Table S1**). Note that

the model with all processes absent (M1) is equivalent to our original ODE model (**Figure 1A**),

and the model with maturation delay only (M2) is equivalent to the base DDE model (**Figure**

2A), and equivalent to the model formulation used in our previous publication,[13] but here with

updated parameter values based on parameterization with additional experimental data.

Simulation & Analysis Software. We simulated differential equations in MATLAB version

R2022a. We used R version 4.3.1 with RStudio release 2023.09.1 for all further data analysis and

data visualization. See *Supplemental References* for the full list of packages and toolboxes used.

All code is available at https://github.com/gillsignals/sFlt1-paper.

Simulating constitutive secretion experiments. We simulate experiments in which cells secrete

sFLT1 constitutively into the extracellular medium for a defined period of time (**Figure 1B, top**).

First, we run a pre-simulation, representing the time period of cell plating and culture before the

experimental measurements, in which the cells come to a steady state. For this, we initialize

intracellular and extracellular sFLT1 to $(I_0, X_0) = (0,0)$, we set rate parameters to desired

values, and we run the simulation to intracellular steady state I_{SS} (defined as less than 0.5%

change in I over 20h of simulation). These conditions are then used as the initial conditions for

the simulation of the actual experiment, except that to represent media change at $t = 0$, we reset

the extracellular concentration X to 0 without changing the intracellular sFlt1 ($I = I_{SS}$). We then

run the simulation for either 24h or 72h, matching the specific experiments being simulated,

reporting (I, X) at 1-minute intervals.

Simulating pulse-chase experiments. We also simulate experiments in which labeled media is

introduced briefly and then removed, in order to track secretion of newly-synthesized (and thus

labeled) sFLT1 **(Figure 1B, bottom)**. For these pulse-chase experiments, we initialize the

intracellular and extracellular sFLT1 to $(I_0, X_0) = (0,0)$, set the parameters to desired values

(including nonzero sFLT1 production rate constant α), and run the simulation for 20m from $t =$

$(t_{pulse}, 0)$ to mimic synthesis of labeled sFLT1 by cells exposed to pulse media. We used the

resulting system state as the initial condition for the remaining simulation, with X reset to 0 at

$t = 0$ without changing I to represent media change to unlabeled chase media. We then reduced

α to zero and continued the simulation for 10h, reporting (I, X) at 1-minute intervals. For

candidate models with exponential decay of production, the production rate during the chase was

instead αe^{-kt}.

Simulation algorithms. For ODE models, we simulated equations using `ode23s` in MATLAB

with an absolute tolerance of 10^{-12}, relative tolerance of 10^{-8}, initial step size of 10^{-2}, and all

species constrained to non-negative values. For the DDE and other candidate models, we

simulated equations using `dde23` in MATLAB with an absolute tolerance of 10^{-4}, relative

tolerance of 10^{-3}, initial step size of 10^{-2}, and any negative concentrations that emerge due to

numerical approximations (which are always very small) are forced to 0.

Experimental data for optimization. We compiled three independently published datasets of

the time course of sFLT1 secretion by cultured human umbilical vein endothelial cells

(HUVECs) (**Supplemental Table S2**). To distinguish the three datasets, we will refer to them by

the name of the first author in each case. Two of these studies quantified constitutive sFLT1

secretion after media change in the presence of minimal serum and no added perturbation: one

using ELISA measurements of absolute sFLT1 concentration in conditioned media ("Hornig"[30])

and one using quantitative Western blots of sFLT1 in both conditioned media and cell lysate

("Kinghorn"[13]). The third study used a pulse-chase experiment to quantify newly-synthesized

sFLT1 in both conditioned media and cell lysate with quantitative Western blots ("Jung"[22]). For

the Jung and Hornig studies, summary data were extracted from published figures using the

Figure Calibration function in ImageJ.[31] Two data points were omitted: Hornig 3h (assumed

below the limit of quantitation for ELISA) and Jung 10h (trend inconsistent with all other

experimental data and all simulations). The Kinghorn data were available in original tabulated

form; to give all datasets comparable weight in optimization, the mean values at each time point

were used in datasets where multiple measurements per time point were available (mean and

median values were comparable).

Initial guesses for parameter values. We ran optimization on each model starting from $n =$

100 (ODE and candidate DDE models) or $n = 1000$ (final DDE model) sets of initial guesses

for parameter values. These initial values were randomly sampled from ranges (**Supplemental**

Table S3) based on previous rate estimates of protein synthesis, transport, and degradation

processes from experimental proteomic approaches[32–34] and previous mechanistic models of

VEGF receptor tyrosine kinases,[26,27] including parameter values from our previous

implementation of candidate model M2.[13]

Optimization cost function. For each dataset, we first converted the predicted X and I values

(from the simulations) from units of #/cell to the units measured by the experimental assay.

Then, for each observation in that dataset, we calculated the relative error between simulated and

experimental results, (simulated value – experimental value)/(experimental value). Finally, we

calculated the sum of squares of these errors, equally weighting the error of each data point. To

minimize the cost function, we performed nonlinear least squares optimization using

`lsqnonlin` in MATLAB. We bounded the search space for each parameter as shown in

Supplemental Table S3.

Scoring of candidate models. Each candidate model was assigned a quality score using the

corrected Akaike Information Criterion (AIC$_C$):

$$n \, \log\left(\frac{c}{n}\right) + 2k + \frac{2k(k+1)}{n-k-1} \qquad \text{(Eq. 5)}$$

where n is the number of experimental data points, C is the cost of a given model calculated

using the custom cost function, and k is the number of degrees of freedom (number of

parameters + 1) for that model.[35,36] The first two terms represent the uncorrected AIC, and the

last term is a correction factor for small sample size because $\frac{n}{k} < 40$.

Sensitivity analysis. Both local and global sensitivity analyses were performed on the DDE

model starting from a baseline of the median optimized parameter set. Local univariate

sensitivity of key output variables (such as sFlt1 concentrations) to key mechanistic parameters

was calculated by running simulations with one parameter increased by 10%, and the relative

change in the output variable compared to baseline. For global univariate sensitivity analysis, we

ran simulations for a wide span of parameter values, each parameter in turn being scanned from

100x lower to 100x higher than the baseline value.

Simulation of chemical inhibition or genetic downregulation. To represent the response to the

acute addition of a chemical inhibitor of one of the model processes, we simulate constitutive

secretion as defined above, with the pre-simulation running the system to steady state using the

median set of optimized parameters, but when media change is applied at $t = 0$, we also adjust a

single process parameter (α, β, γ, δ, or τ) using $p' = (1 - f) * p$ where p is the initial parameter

value and f is the inhibitor's effect, i.e. the fractional inhibition of that process. To represent

longer-term system perturbation, e.g. due to genetic mutation of a system component or by

expression downregulation of a trafficking protein via RNA interference days before an

experimental time course, we again simulate constitutive secretion, this time running both the

pre-simulation to steady state and the experimental simulation with a single process parameter

adjusted as $p' = (1 - f) * p$, where again p is the initial parameter value and f is the

magnitude of the downregulation, i.e. the fractional inhibition of that process. To compare the

predicted effects of acute and chronic perturbations of trafficking mechanisms, we used

experimental measurements of intracellular and extracellular sFLT1 following inhibitor

treatment, expressed as a fraction of control (i.e. the absence of perturbation).[13] Using our model,

evaluated at a range of inhibition strengths, we estimated the fractional inhibition for each

inhibitor of that inhibitor's target process, i.e. the predicted parameter reduction required for the

model to produce the same relative change in sFLT1 as experimentally observed.

Results

ODE model of sFLT1 secretion. The ODE model (see *Methods*, **Equations 1-2**) includes rate

terms for production, secretion, intracellular degradation, and extracellular degradation,

described by four parameters (**Figure 1A**). A general analysis of the ODE model solution

dynamics (**Figure S1**) is described in more detail in *Supplemental Results*. By setting the time

derivatives to zero, we calculated theoretical steady state solutions:

$$I_{SS} = \frac{\alpha}{\beta+\gamma} \qquad \text{(Eq. 6)}$$

$$X_{SS} = \frac{\beta}{\delta} I_{SS} = \frac{\alpha\beta}{\delta(\beta+\gamma)} \qquad \text{(Eq. 7)}$$

and characteristic times to half-maximal concentrations for intracellular and extracellular sFLT1

assuming first-order kinetics for secretion and degradation:

$$T_{50\_I} = \frac{ln(2)}{\beta+\gamma} \qquad \text{(Eq. 8)}$$

$$T_{50\_X} = \frac{ln(2)}{\delta} \qquad \text{(Eq. 9)}$$

The derivations of Equations 6-9 are given in *Supplemental Methods*.

To calibrate the model parameter values to sFLT1 biology, we used optimization

algorithms to fit simulation predictions of sFLT1 concentrations simultaneously to intracellular

and extracellular sFLT1 concentration datasets from three independent studies of sFLT1

secretion in human umbilical vein endothelial cells (HUVECs): one pulse-chase time course in

relative units ("Jung"[22]) and two constitutive secretion time courses, one in relative units

("Kinghorn"[13]) and one in absolute units ("Hornig"[30]) (**Supplemental Table S2**; **Figure 1B**).

The pulse-chase secretion experiment provided particularly useful data for characterizing early

secretion behavior, while the constitutive secretion experiments allowed for estimation of

biologically relevant production parameter values. The optimization appeared at first to be well-

constrained, but resulted in a set of solutions with identical cost (goodness of fit) that produced

identical extracellular and normalized intracellular time courses (**Figure 1C**, $n = 100$

superimposed lines) and yet differed in absolute intracellular sFLT1 amount over 2 orders of

magnitude (**Figures 1D-E**). These optimized fits reproduced experimentally observed dynamics

of constitutive sFLT1 secretion into culture media, save 2 outlier data points (X_{72h} and I_{24h}) that

were inconsistent with other experimental observations. However, this initial model could not

recapitulate two clearly observed features of early sFLT1 secretion dynamics in the chase phase

of the pulse-chase scenario: (1) maintenance or increase of intracellular sFLT1 level for several

hours after production ends and (2) a multi-hour delay in extracellular sFLT1 accumulation.

Delay differential equation (DDE) model of sFLT1 secretion. To more accurately model the

observed sFLT1 secretion dynamics, we converted our model to a system of two delay

differential equations (DDEs) by introducing a fixed time delay between sFLT1 production and

trafficking (**Figure 2A**; see *Methods*, **Equations 3-4**). This "maturation delay" represents the

time required for newly synthesized intracellular proteins to traverse the secretory pathway,

reach a mature form, and be sorted into vesicles for trafficking. This delay is particularly

important in the pulse-chase scenario because only newly synthesized (and therefore labeled)

proteins are experimentally measured, and so even though unlabeled sFLT1 made before the

media change might be secreted sooner, the labeled sFLT1 will only be extracellularly secreted

and observed after the full delay. The model with the delay would therefore only represent the

labeled sFLT1, for comparison to the experimental data. Under this scenario, the rate terms for

secretion and intracellular degradation are functions of intracellular sFLT1 concentration at time

$t - \tau$, for some time delay τ, rather than at time t as in the initial model. A general analysis of the

DDE model solution dynamics (**Figure S2**) is described in more detail in *Supplemental Results*.

Because at steady state $I(t - \tau) = I(t)$, this model has the same steady state properties as the

ODE model (**Equations 6, 7**).

We optimized the DDE model parameters to the constitutive and pulse-chase secretion

datasets similarly to the ODE model. The optimized DDE model generated pulse-chase

dynamics that fit experimental data better than the ODE model, and fit the constitutive secretion

data as effectively as the ODE model (**Figure 2B**). However, note that in the pulse-chase

experiment, intracellular sFLT1 increased above its $t = 0$ baseline at 2h before decreasing, and

in this DDE model formulation there is no mechanism by which intracellular sFLT1 could

increase after production stops at $t = 0$. This inconsistency could suggest that our model

overlooks one or more processes essential to describing the biology of sFLT1 secretion, or it

could be a consequence of noise in the data, in which case adding parameters to the model risks

overfitting the data.

Testing alternate candidate models. We examined whether we could further improve our

model by including different combinations of three processes: the maturation delay described

above; first-order internalization of extracellular sFLT1; and exponential decay of production at

the end of an expression pulse (see *Methods*). We defined rate terms for each reaction for cases

when each process was present or absent; for example, the internalization process had a rate of

$\epsilon \cdot X$ when present but a rate of 0 when absent. Note that the absence of a process is equivalent to

setting $\varepsilon = 0$ (internalization), $\kappa = 0$ (production decay), or $\tau = 0$ (maturation). Using these

terms, we built an ensemble of eight models by generating each model's equations in a piecewise

fashion (**Figure 3A**, **Supplemental Table S1**). Note that the model with all additional processes

absent (M1) is equivalent to our original ODE model (**Figure 1A**), and the model with

maturation delay only (M2) is equivalent to the base DDE model (**Figure 2A**). We applied our

optimization algorithm to each model with 100 different sets of initial parameters to compare the

'best fit' version of each model.

When considering only the lowest cost fit for each model, we found that all models with

additional processes had a slightly improved fit (lower cost, defined as the sum of squared

differences between simulated and experimental data points) relative to the original ODE model

M1 (**Figure 3B**, middle). While models performed similarly in constitutive scenarios (**Figure**

S3B, D), models that included maturation delay τ were better able to describe extracellular

sFLT1 behavior in the pulse-chase scenario (**Figure S3A**), while inclusion of either maturation

delay τ or production decay κ improved model fit to intracellular sFLT1 during pulse-chase

(**Figure S3C**). Although every parameter addition improved goodness of fit, addition of

maturation delay τ consistently caused the largest cost reduction (**Figure 3C**).

Improvements in fit required introducing additional parameters, so to reduce the risk of

overfitting, we scored the models by balancing cost and complexity (number of parameters)

using the corrected Akaike Information Criterion (AIC$_C$, **Equation 5**).[35,36] While adding

maturation delay τ to any model reduced AIC$_C$, adding production decay κ only reduced AIC$_C$ if

maturation delay τ is not already included, and adding internalization ε never reduced AIC$_C$.

Model M2, which includes maturation delay τ but not internalization ε or production decay κ

and was analyzed in **Figure 2**, had the lowest AIC$_C$ and thus had the best fit to the data given its

number of parameters (**Figure 3B**: bottom). We interpret these results as showing that model M2

performs as well as the more complex models we tested based on the limited experimental data,

so we therefore chose to move on to downstream analysis using this model (**Figure 2A**).

Optimized parameter values. To more thoroughly explore the properties of our selected DDE

model, and to evaluate uncertainty in optimized parameter values for the model, we repeated the

analysis with a larger set of 1000 optimization runs starting from randomly sampled initial

values. While most resulting fits follow the experimental data well, numerous fits failed to

converge to the experimental data (**Figure S4A**). To enrich for high-quality fits, we filtered for

optimization runs where the cost of the resulting fit was within 10% of the minimum cost (n =

594) (**Figure 2B**, **Figure S4B**). Comparing the distribution of initial parameter values for all

1000 runs (**Figure S4C**, yellow; **Supplemental Table S3**) to the distribution of initial values for

the 594 runs that passed the cost filter (**Figure S4C**, black), we can see that the filtered-out runs

generally had extreme initial parameter values, especially large initial values of β .

After filtering, optimized values of δ and τ were very well-constrained ($\delta = 0.057 \pm$

0.0014 1/h, $\tau = 1.958 \pm 0.020$ h), while the other parameters (α, β, γ) spanned 2-3 orders of

magnitude (**Figure 4A**, **Table 1**, **Figure S5A**). Correlation plots show algebraic relationships

between α, β, and γ, including an inverse nonlinear relationship between α and β, and an inverse

linear relationship between β and γ (**Figure 4B, Figure S5B**). Because of these apparent

relationships, we looked for joint constraints among combinations of parameters and found that

both the product $c_1 = \alpha\beta$ and the sum $c_2 = \beta + \gamma$ are well-constrained to single values ($c_1 =$

7252 ± 131 #/cell/h^2, $c_2 = 0.173 \pm 0.00064$ 1/h, mean $\pm$ SD) (**Figure 4C**, **Table 2**). This five-

parameter model thus has four constraints (δ, τ, c_1, c_2) and only one degree of freedom: if any

one of α, β, or γ is specified, then the values of the other two parameters are determined by the

constants c_1 and c_2. When these constraints are present, Equations 6 and 7 can be extended as:

$$I_{SS} = \frac{\alpha}{\beta+\gamma} = \frac{\alpha}{c_2} \tag{Eq. 6a}$$

$$X_{SS} = \frac{\beta}{\delta} I_{SS} = \frac{\alpha\beta}{\delta(\beta+\gamma)} = \frac{c_1}{\delta c_2} \tag{Eq. 7a}$$

Steady state intracellular sFLT1. During constitutive simulation, intracellular sFLT1 remains

at a steady state (I_{SS}) whose absolute amount varies across parameter sets within a range of 2.0 x

10^5 - 2.6 x 10^7 molecules per cell, with a median of 8.2 x 10^5 #/cell (**Figures S6**). The simulated

steady state values match the theoretical definition of $I_{SS} = \frac{\alpha}{\beta+\gamma}$ across all fits (**Figure 4D**,

Equation 6). Relationships between I_{SS} and α, β, and γ also match the theory as follows. α is

directly proportional to I_{SS} (**Figure 4E**). β is inversely proportional to I_{SS} and approaches a

maximum value of c_2 as I_{SS} approaches its minimum value (**Figure 4F**); as β increases, α

decreases (because $c_1 = \alpha\beta$) and thus I_{SS} declines. γ is inversely related to β and approaches a

maximum value of c_2 as β approaches 0; this causes α to increase, and so I_{SS} approaches infinity

(**Figure 4G**). The presence of the c_1 constraint explains why β and γ have opposite relationships

to I_{SS} despite both being in the denominator of $I_{SS} = \frac{\alpha}{\beta+\gamma}$. Of note, each intracellular steady state

is associated with a unique set of (α, β, γ) values, so high-confidence experimental

measurements of intracellular sFLT1 would provide enough data to fully constrain our

mechanistic model.

Process fluxes. We explored the flux of each process, i.e. the rate of molecules traversing each

edge of the reaction network graph (**Figure 5A-B**). During constitutive secretion, extracellular

degradation flux increases hyperbolically over time, while the other three processes have

constant fluxes over time (**Figure 5C-D**). The extracellular degradation flux converges towards a

single time course directly proportional to the time course of extracellular sFLT1 (**Figure 2B**).

The secretion flux converges towards a single value of $\frac{c_1}{c_2} = \frac{\alpha\beta}{\beta+\gamma}$ across all model fits (**Figure 5E,**

Figure S7A). In contrast, production and intracellular degradation flux values differ across fits

and their values are paired such that any increased production flux is offset by intracellular

degradation (**Figure S7B**), and both are algebraically related to the observed intracellular steady

state concentration (**Figure S7C-D,** *Supplemental Methods*). The fit-specific differences in

production and intracellular degradation flux also occur during simulation of pulse-chase

secretion (**Figure 5F-G**), though all models had similar relative fluxes ($\frac{\Phi}{\Phi_{max}}$) over time (**Figure**

S8).

Local univariate sensitivity analysis. We evaluated how uncertainty in each input parameter

affects the uncertainty of selected output variables during constitutive secretion, including

extracellular sFLT1 after 72h (X_{72h}), intracellular sFLT1 (I_{72h}; note that for constitutive

secretion, $I_{72h} = I_{SS}$), and time to half-steady-state for extracellular sFLT1 (T_{50_X} = time to

$X_{SS}/2$), where X_{SS} is calculated theoretically. Starting from the median parameter set (**Table 1**),

we individually increased the value of each parameter by 10%, simulated constitutive secretion,

and calculated the relative change in output variables (**Figure 6A**). Intracellular and extracellular

sFLT1 increase linearly with production rate constant α (sensitivity = 1). Increasing the secretion

rate constant β causes a moderate rise in extracellular sFLT1 but a mild decrease in intracellular

sFLT1, while increasing the intracellular degradation rate constant γ causes a decline in both

intracellular and extracellular levels. The extracellular degradation rate constant δ has a strong

negative effect on extracellular sFLT1 but has no impact on intracellular protein. Notably, δ is

the only parameter which affects the time to half-steady-state (T_{50_X}) for extracellular sFLT1,

with higher δ associated with a faster time to steady state.

Global univariate sensitivity analysis. To explore how individual parameters affect sFLT1

trafficking over a larger range of values, we varied each parameter individually over several

orders of magnitude and simulated constitutive secretion (**Figure 6B, Figure S9**). For all

parameter sets tested, intracellular sFLT1 remains at its relevant steady state throughout a

simulation, while extracellular sFLT1 increases hyperbolically towards a steady state. Consistent

with the local sensitivity analysis, both extracellular and intracellular sFLT1 are most sensitive to

changes in α, with changes in α causing directly proportional changes in both steady state

intracellular sFLT1 and extracellular sFLT1 72 hours after media change, across the range of α

values. β, γ, and δ have narrower ranges of effects than α (**Figure 6B, Figure S9**), because

below or above certain thresholds changing the parameter value has no effect (**Figure S9**). While

β and γ affect extracellular sFLT1 in opposite directions, they influence intracellular sFLT1 in

the same direction but with different magnitudes, with the response to γ being larger and over a

wider range than β. δ does not affect intracellular sFLT1, but decreasing δ increases extracellular

sFLT1 while slowing its rate of convergence to steady state. The constitutive secretion scenario

masks any impact of changing the secretion delay τ, with altered simulation results at high τ only

because long delay interferes with our method of initializing the system to steady state before

media change. While the constraints (c_1, c_2) are fixed for our sFLT1 system, we also explored

the sensitivity of the system to changes in these values (*Supplemental Results*, **Figure S10**).

Chemical perturbation of sFLT1 secretion. We explored how our model of sFLT1 secretion

responds to either chemical inhibition or genetic downregulation of various processes during

constitutive secretion. Chemical inhibition represents the application of a druglike inhibitor at a

specific experimental time point, while genetic downregulation reflects a system where genes

encoding cellular trafficking machinery are mutated or their expression is blocked by RNAi

before the experimental time course begins (**Figure 7A**). Chemical inhibition is acute, genetic

downregulation more chronic; thus, to simulate chemical inhibition, we first ran the simulation to

steady state with baseline parameter values, and then we applied media change ($X \rightarrow 0$) and

inhibitor treatment (parameter value adjustment) at $t = 0$, while to simulate genetic

downregulation, we applied the parameter changes continually from the beginning of the

simulation, before running the simulation to steady state.

Starting from the median parameter set (**Table 1**), we chemically inhibited each

parameter individually by 90% and investigated how inhibition affects intracellular and

extracellular sFLT1 over time (**Figure 7B-C**). Inhibition of the production rate constant α causes

a gradual but large decrease in intracellular sFLT1, which approaches a new lower steady state

within 16h, while extracellular sFLT1 initially increases before declining to a new steady state

within 48h. Inhibition of the secretion rate constant β causes a gradual small increase in

intracellular sFLT1, which approaches a new higher steady state within 16h, while extracellular

sFLT1 hyperbolically approaches steady state with lower concentration but similar dynamics as

the uninhibited case. Inhibiting the intracellular degradation rate constant (γ) leads to a

hyperbolic increase in both intracellular and extracellular sFLT1, both of which take over 48h to

approach steady state. Inhibiting extracellular degradation (δ) does not alter intracellular sFLT1

but causes extracellular sFLT1 to increase continuously throughout the entire 72h simulation.

Adjustment of the secretion delay time (τ) does not influence sFLT1 during constitutive

secretion.

We then evaluated the impact of strength of inhibition on extracellular and intracellular

sFLT1 at two time points: 18 hours, a time point at which inhibitors of sFLT1 secretion have

been experimentally tested,[13] and 72 hours, the standard length of our constitutive simulation.

Consistent with previous findings, inhibiting production (α) decreases both intracellular and

extracellular sFLT1, inhibiting secretion (β) slightly increases intracellular sFLT1 but strongly

decreases extracellular sFLT1, inhibiting intracellular degradation (γ) increases both intracellular

and extracellular sFLT1, and inhibiting extracellular degradation (δ) does not affect intracellular

sFLT1 but increases extracellular sFLT1 (**Figure S11A**). All these effects increase in size as

fraction inhibition increases, although the effect of inhibiting α increases linearly with fraction

inhibition and the effects of inhibiting other parameters increase superlinearly. While the effects

of inhibiting α and β are roughly equivalent at 18h and 72h, the impact of inhibiting γ and δ are

larger at the later time point, consistent with only the α and β-perturbed simulations approaching

their new steady state by 18h.

We compared our model predictions to previous experiments[13] evaluating how chemical

inhibitors of the secretory pathway affect intracellular and extracellular sFLT1 after 18h of

treatment (**Figure 7D**, **Table 3**). Based on observed experimental changes in extracellular

sFLT1, our model suggests the lysosomal degradation inhibitor chloroquine inhibits the sFLT1

intracellular degradation rate constant γ by 98%, while the observed change in intracellular

sFLT1 suggests a lower but still substantial 69% inhibition of γ. Similarly, based on extracellular

sFLT1 experimental measurements, our model suggests the Golgi inhibitor brefeldin inhibits the

sFLT1 secretion rate constant β by 89%, while the vesicular transport inhibitor chlorpromazine

and the exocytosis inhibitor TATNSF700 only inhibit β by 58% and 54% respectively. However,

the experimental intracellular sFLT1 concentrations observed for all 3 drugs exceeded the

predicted concentration expected by our model even if secretion is fully blocked. This suggests

our model may be missing an important component describing intracellular retention of sFLT1,

as inhibiting secretion leads to much larger accumulation of intracellular sFLT1 than our model

can explain. This could also suggest that the effect of secretion-inhibiting drugs is more complex

than simply altering the secretion rate constant, possibly also reducing intracellular degradation,

e.g. by relocating sFLT1 to subcellular compartments from which degradation does not occur.

Genetic perturbation of sFLT1 secretion. Genetic inhibition of individual parameters (**Table**

4) by 90% produced similar results as chemical inhibition at later time points (**Figure 7E**), but

dynamics of early inhibition differed in many cases due to the presence of inhibition during the

pre-simulation stage (**Figure 7F**); for example, intracellular sFLT1 remains at steady state after

media change at $t = 0$, but this steady state level is reduced by inhibiting the production rate

constant α or increased by inhibiting the secretion rate constant β or intracellular degradation rate

constant γ. The steady state intracellular levels induced by 90% genetic inhibition are the same as

the intracellular levels approached over time during 90% chemical inhibition, but because the

genetic inhibitor is applied earlier, these steady state levels have already been reached by $t = 0$.

Extracellular sFLT1 time courses for genetic and chemical inhibition are identical for inhibition

of β, γ, and the extracellular degradation rate constant δ. Genetic inhibition of α causes

extracellular sFLT1 to hyperbolically approach its decreased steady state level, unlike in

chemical inhibition where extracellular sFLT1 initially overshoots its final level before a delayed

decrease to the new steady state.

Genetic parameter inhibition has the same directional effects on extracellular and

intracellular sFLT1 at 18h and 72h as in chemical inhibition (**Figure S11B**). The impact of

inhibiting extracellular degradation (δ) is larger at the later time point, with the discrepancy

increasing in size as fraction inhibition increases. For all other parameters, genetic inhibition led

to the same changes in intracellular and extracellular sFLT1 at both 18h and 72h.

We compared our model predictions to previous experiments[13] exploring how small

interfering RNA (siRNA) inhibition of the secretory pathway affects intracellular and

extracellular sFLT1 (**Figure 7G**; **Table 4**), assuming that the inhibition reached its maximum

effectiveness by the media change at $t = 0$ and remained fully effective throughout the

remaining experimental time. Based on observed experimental changes in extracellular sFLT1,

our model suggests that inhibition of the tested Golgi and vesicular trafficking machinery (STX6,

RAB27a, STX3, VAMP3) decreases the effective secretion rate constant β by 78-98%,

consistent with these factors playing a role in sFLT1 secretion. However, while our model

predicts a minor increase in intracellular sFLT1 with inhibition of the secretion rate constant,

genetic inhibition of STX3 leads to a much larger intracellular sFLT1 accumulation than

expected by our model even if secretion is fully blocked, and genetic inhibition of the other

tested factors caused a paradoxical decrease in intracellular sFLT1. These observations again

suggest our model is an incomplete descriptor of intracellular sFLT1 biology, despite model

agreement with other observed experimental data; for example, the targets of inhibition can

affect the trafficking of many proteins, which may themselves have a role in sFLT1 trafficking.

Evaluating uncertainty: simulating inhibition using different initial parameter sets. Since

there is some remaining uncertainty in parameter values, we explored how this uncertainty

impacts the above insights into chemical and genetic inhibition. We generated five parameter

sets consistent with the observed constraints, by setting the baseline secretion rate constant β to

one of five different values ($0.2*c_2$, $0.4*c_2$, $0.6*c_2$, $0.8*c_2$, c_2), then calculating $\alpha = c_1/\beta$ and

$\gamma = c_2 - \beta$ for each case. To reiterate, these parameter sets all satisfy the underlying temporal

data; so our question was: would comparing simulations for different parameter sets to the

inhibitor data help us to distinguish which parameter set or sets are best? Thus, for each

parameter set (**Figure S12**), we tested the effect of chemical or genetic inhibition by a range of

inhibition fractions at both 18h and 72h (**Figure S13**). In chemical inhibition, the effects of

inhibiting sFLT1 production (α) or extracellular degradation (δ) are independent of the initial

parameter set, but the choice of parameter set affects the size (though not the direction) of the

system's response to inhibiting either sFLT1 secretion (β) or intracellular degradation (γ)

(**Figure S13A**). As β increases, intracellular sFLT1 becomes more sensitive to inhibiting β and

less sensitive to inhibiting γ. Extracellular sFLT1 response to γ inhibition also depends on the

initial parameter set, while the initial parameter set has a smaller impact on extracellular sFLT1

response to β inhibition and effects differ only at intermediate β inhibition fractions. These

differences between parameter sets are more pronounced at 72h than at 18h. For genetic

inhibition, simulated results resemble later time points for chemical inhibition simulations for the

intracellular parameters α, β, and γ (**Figure S13B**). Extracellular sFLT1 is more sensitive to

increasing fraction inhibition of β or γ at 72h than at 18h, but this effect is independent of the

initial parameter set, and for all other parameters the results of genetic inhibition match at these

two time points.

We then compared our inhibition simulations to experimental results[13] (**Figure 8**). Based

on extracellular sFLT1 data from chemical inhibitors, all parameter sets generated plausible

estimates (<100%) for inhibition fraction of secretion (β), with higher fraction inhibition

estimated for higher initial β values (i.e. lower intracellular sFLT1 levels), and with more

uncertainty in the effect size for the moderate inhibitors chlorpromazine and TATNSF700 (55-

75%) than for the strong inhibitor brefeldin (88-95%) (**Figure 8A**). In contrast, for intracellular

sFLT1 following chemical inhibitors, the estimated β inhibition fraction increases as β decreases

(intracellular sFLT1 increases), and a feasible fraction inhibition of β (<100%) can only be

estimated for parameter sets with initial β values above a certain threshold, which differs for each

inhibitor (approximately, β>0.5 for TATNSF700, β>0.7 for brefeldin, and β>1.1 for

chlorpromazine). When initial values of β are low, even full inhibition of β does not generate a

large enough fold change in intracellular sFLT1 to match experimental results. This suggests an

additional constraint on the value of the secretion rate constant β. The intracellular degradation

(γ) inhibitor chloroquine shows the opposite pattern: for both intracellular and extracellular

sFLT1, the estimated γ inhibition fraction increases as β increases (intracellular sFLT1

decreases), and a feasible fraction inhibition of γ (≤100%) can only be estimated at lower initial

values of β (and thus high initial values of γ and intracellular sFLT1), approximately β<0.1.

Collectively, these suggest that an intermediate value of β, similar to the median value of 0.072

(Table 1), can provide a basis for most inhibitor observations; our earlier results showed that a

constrained value of β also constrains α and I_{SS}.

Comparing simulated and experimental genetic inhibitors of β (**Figure 8B**), we find that

all parameter sets except those with the highest initial β values (lowest intracellular sFLT1) led

to reasonable (<100%) estimates of inhibition fraction, again consistent with an intermediate β

value. However, experimental downregulation of the secretion factors VAMP3, RAB27a, and

STX6 decreased intracellular sFLT1, whereas in our simulations, inhibiting secretion increased

intracellular sFLT1 for all inhibition fractions. Only experimental STX3 inhibition changed

intracellular sFLT1 by a magnitude consistent with our simulations, and only for parameter sets

with high initial values of β > 0.11 (low intracellular sFLT1). Our interpretation is that these

inhibitors may have mechanistic effects different to those included in our model – such as effects

on the trafficking on other proteins that themselves impact sFLT1, as mentioned above – and

bear further investigation.

Discussion

Here, we have developed mechanistic computational models of sFLT1 secretion by endothelial cells. sFLT1 is a soluble form of the VEGFR1 protein, whose membrane-associated form is one of three receptor tyrosine kinases that bind VEGF family cytokines, a key family of ligands that induces blood vessel growth in health and disease. Because sFLT1 can bind the ligand but has no tyrosine kinase domain, it does not initiate signaling and therefore acts as a sink for ligand, downregulating its signaling potential. Therefore, sFLT1 is an important negative regulator of blood vessel growth, and understanding its production, secretion, and levels outside the cell are important to understanding VEGF regulation.

There have been a limited number of experimental studies that quantified sFLT1 secretion over time by endothelial cells,[13,22,30,37] and here we have used the data on sFLT1 levels inside and outside the cell from several of those studies to parameterize the models. We also used data from a recent study[13] that tested multiple chemical and genetic inhibitors of proteins known to be involved in protein trafficking and secretion, to determine which were relevant for sFLT1 secretion, and in so doing quantified the level of disruption to sFLT1 extracellular levels over time (and therefore sFLT1 secretion) due to these inhibitors. Since the inhibitors block or reduce the effectiveness of the mechanisms included in our models, we explored model perturbations equivalent to these inhibitors in the simulations also.

A key first step in our study was comparing different mechanistic models for sFLT1 secretion. These models differed in the incorporation or exclusion of mechanisms such as: a maturation time delay between production and availability for secretion; a gradual decay in production rate when production ceases; and the internalization of previously-secreted

extracellular sFLT1. The addition of some of these time-based processes also requires shifting

from a typical ODE-based model to a DDE (delay differential equation)-based model. The

resulting eight candidate models were tested, i.e. the parameters for each were optimized

independently against the three experimental datasets described earlier and the predictions of the

best-fit parameter sets for each model then compared across the eight models. The criterion used

for overall model choice was the Akaike Information Criterion (AIC), which balances goodness

of fit (i.e. how closely the simulation predictions match the experimental data) with the

complexity of the model (i.e. how many parameters are needed to define it, with the

understanding that more parameters would require more data for determination). As a result,

model M2 - a DDE-based model that included maturation delay but not internalization or

production decay - was identified as the best model and was used for the remainder of our study.

This does not mean that production decay and internalization do not happen in the actual cells,

nor indeed that they don't improve goodness of fit of the model when included in the

simulations; rather, it means that their effect is relatively small and the improvement is not worth

the additional model complexity.

Based on this chosen model, and comparing the results of hundreds of optimization runs

attempting to fit to the experimental data, of the five mechanistic parameters ($\alpha, \beta, \gamma, \delta, \tau$), two

of them (δ, extracellular degradation; τ, maturation delay) are very well constrained. These

parameters are likely well-constrained because they are integral to the shape of the concentration

curves over time, something that is relatively invariant to the total amount of sFLT1 in the

system; this is relevant because much of the experimental data is not in absolute concentration

units, and thus the comparisons between simulation and experiment are done after both are

normalized to be on the same scale. In the absence of an absolute intracellular sFLT1

measurement, the other parameters (α, β, γ) are not fully constrained. Thus, we have a family of

solutions - however, we observed that these α, β, and γ values are not all independent, and two

further constraints are apparent when comparing the optimized parameter sets: $c_1 = \alpha\beta$ (i.e.

production rate times secretion rate constant is a fixed value), and $c_2 = \beta + \gamma$ (i.e. the sum of

secretion and intracellular degradation rate constants is a fixed value). As a result, there is only

one degree of freedom remaining, and setting any one of these three parameter values (or

knowing the value of intracellular sFLT1 levels) determines the other parameters. Thus, even in

the absence of a single completely unique parameter set, the resulting model of sFLT1 is still

well defined because (a) the overall dynamics are fairly consistent due to the constraint of δ and

τ, and (b) the additional constraints on α, β, and γ mean that there is a unique solution for any

given intracellular sFLT1 level, i.e. if it were possible to measure an absolute value for

intracellular sFLT1 levels, then the system would be fully determined. Experiments that measure

the dynamic responses to selective perturbations (e.g. of intracellular degradation) may help pin

down the specific rate constants; we also explore the uncertainty in model predictions due to

different parameter sets that are consistent with the experiments and span the range of possible

parameter set values.

We use the parameterized model to simulate chemical inhibitors (e.g. small molecules

that interfere with key trafficking processes) and genetic inhibitors (e.g. small interfering RNAs

(siRNA) that downregulate the expression of key proteins involved in trafficking), in particular a

subset of the inhibitors that were experimentally tested in a recent study.[13] The inhibitors are

simulated in the model by the modification of the rate parameters for the process that the

inhibitors are thought to interfere with; for example, if the inhibitor blocks intracellular

degradation, the value of γ is decreased. Note that there are known to be many sequential steps

to secretion in cells, and the inhibitors tested included those targeting different molecules at

different steps, but in our model we aggregate all secretion steps to one process. We distinguish

between chemical and genetic inhibitors in our model based on when we apply the change in

parameter values: to simulate chemical inhibition, we allow the system to reach steady state with

baseline parameters before adjusting any parameter values, whereas we simulate genetic

inhibition by adjusting the target parameter from the start of the simulation. This timing

difference – longer-term genetic inhibition vs acute chemical inhibition – may give time for

genetic inhibition to result in second-order effects where proteins and processes not included in

the model are affected and can then themselves impact sFLT1 trafficking.

Overall, we see mixed results when comparing our simulations to the experimental

observations. Our model is consistent with the experimental data for all tested chemical

inhibitors of secretion and intracellular degradation for at least some of the possible parameter

sets tested. For example, the secretion inhibitor brefeldin decreases extracellular sFLT1 and

increases intracellular sFLT1, and our simulations predict the drug decreases the secretion rate

constant by over 50% and likely closer to 90%, the precise value depending on which parameter

set is used. However, the experimental data for several genetic inhibitors behave unexpectedly

and change in the opposite direction compared to our model. For example, we simulate inhibiting

the vesicular trafficking factor VAMP3 as decreasing the secretion rate constant while leaving

other parameters unchanged, which is predicted to decrease extracellular sFLT1 while increasing

intracellular sFLT1; but experimental downregulation of VAMP3 instead decreases intracellular

sFLT1. Furthermore, this mismatch occurs regardless of which initial conditions we choose from

the family of optimized parameter sets.

It is not immediately clear how to interpret or resolve this discrepancy between our model

and experiments for these genetic inhibitors. It is difficult to identify a single missing process

that, if included in the model, would explain lower intracellular sFLT1 when secretion is

blocked. Rather, as noted above it is more likely that there is some complex regulation at play for

those specific inhibitors. Importantly, the chemical and genetic inhibitors explored are not

specific to sFLT1, but regulate trafficking for many other proteins, some of which may

themselves modulate sFLT1 synthesis and secretion. Furthermore, nonspecific trafficking

inhibitors like brefeldin can cause massive accumulation of immature protein in the ER and

Golgi, and the resulting cell stress can lead to the unfolded protein response (UPR), a complex

homeostatic response with multifactorial effects on the secretory pathway. Among its many

effects, the UPR increases protein turnover, and while global protein synthesis slows during the

UPR, there is increased transcription and translation of secretory pathway components affecting

protein maturation and trafficking.[38] As such, any inhibitor that triggers the UPR could have

second-order effects altering the rate constants for sFLT1 production, trafficking, and

degradation. Cases like this, where the experimental perturbations affect multiple model

parameters and invoke dynamic feedback regulation, would be difficult to parse in this model.

Focusing on acute inhibition or measuring changes as soon as feasible after inhibition may better

serve to avoid second-order effects. What the model does do, however, is identify precisely those

inhibitors whose observed behavior/outcome does not match what would be expected to happen,

and therefore provides a useful shortlist of inhibitors whose more detailed mechanisms and

effects would be interesting to follow up on.

Returning again to the remaining uncertainty in the model parameter values, we explored

whether the inhibitor experiments could help to pin down the model parameters. Each drug has a

different set of parameter ranges for which modeling inhibition is well-behaved, however these

ranges are not all consistent; for example, some drugs seem to support a high-β-value parameter

set, others a low-β-value parameter set. An intermediate β-value is the most broadly consistent

with experimental data, but further model development, perhaps including more detailed

inhibitor mechanisms, may better constrain our parameter estimates.

Our results provide mechanistic insights about sFLT1 secretion in vitro and suggest

intriguing questions about intracellular behavior of sFLT1. It remains to be seen whether these

insights would also apply in vivo; if so, it suggests that physiological or pathological changes in

sFLT1 concentration could have multiple causes. For example, elevated serum sFLT1

(analogous to extracellular sFLT1 in our model) could result from an increase in sFLT1

production, or increased secretion, or decreased intracellular or extracellular degradation, or any

combination of these changes. Different disorders are likely driven by distinct changes in sFLT1

kinetics, with implications for treatment approaches. In vivo conditions are further complicated

by multi-cellular expression and secretion of sFLT1; for example, in preeclampsia, non-

endothelial cells in the placenta overexpress sFLT1 mRNA[21]. These models could be extended to

understand the effects of such multicellular environments.

Interestingly, our model suggests that less than 20% of mature intracellular sFLT1 is

removed from the intracellular compartment each hour under basal conditions, consistent with a

previous estimate[22] and corresponding to a long intracellular residence time for sFLT1. This

extended residence time raises the question of whether sFLT1 has functional VEGF-binding and

VEGFR-binding roles within the secretory pathway. Interactions can happen because the full-

length receptors accumulate in the same intracellular locations. The full-length mFLT1/VEGFR1

isoform localizes primarily to the Golgi and traffics to the plasma membrane in a calcium-

dependent manner,[8] while VEGFR2 is largely retained in the trans-Golgi and traffics to the cell

surface in response to VEGF stimulation.[39] RTKs are also known to initiate signals from

intracellular organelles[7–10] as well as from their more canonically considered cell surface

locations, and this may also occur within the Golgi and be inhibited by Golgi-resident sFLT1.

Our approach identifies a minimal mechanistic model consistent with experimental data

on sFLT1 secretion, but includes many assumptions that simplify the underlying biology, both

for general features of the secretory pathway and for specific characteristics of sFLT1. The key

assumptions involve homogeneity and the choice of model compartments, species, and reactions,

each described in the following paragraphs. This mass action kinetics modeling approach uses

several assumptions of homogeneity, including that all sFLT1 molecules in a compartment are

identical and equally likely to react, all cells are identical with the same rate constants, and that

system geometry is constant over space and time. However, biological systems are highly

heterogeneous. Tissues consist of numerous cell types of different lineages that vary significantly

in their gene expression and phenotypic profile, typically arranged into higher-order three-

dimensional structures that are not spatially uniform. Individual cell types occupy various states

and can change states over time. Even in monolayer culture of single cell lines, cells are not

identical, as the stochasticity of biochemical reactions leads to natural gradients and fluctuations.

Furthermore, here we assume a constant number of cells over time when simulating or

interpreting experimental results, while the cell culture environment allows cell growth and

proliferation. Since HUVECs proliferate slowly in culture,[40] this is likely not a major confounder

at early time points, but cell number may have increased enough by our 72 hour time point to

impact data interpretation. A hierarchical model that includes both sFLT1 trafficking and cell

population dynamics may add insight and reveal how changing cell number affects interpretation

of the experimental data.

An agent-based modeling (ABM) approach could incorporate cells of different types with

distinct phenotypes and reaction propensities, simulate stochasticity, represent cell proliferation,

and track spatial heterogeneity with partial differential equations. However, ABMs tend to be

computationally expensive, we may not have sufficient data for parameterization, and their

complexity may not be necessary to understand key features of sFLT1 secretion in cell culture.

An intermediate approach that allows modeling of heterogeneous cell communities without the

full computational demand or spatial detail would be to use ODEs to model several separate

intracellular compartments (corresponding to cell subtypes) linked to a common extracellular

compartment. However, since sFLT1 expression varies not only across cell types but across

adjacent endothelial cells,[20] it may be more appropriate to use a spatially-aware modeling

approach like an ABM to explore heterogeneous cell populations. While ABMs have been

extensively applied to the biology of the VEGF family,[41–45] some including extracellular

sFLT1,[46,47] to our knowledge they have not been used to explore intracellular sFLT1 behavior

and secretion.

In our model, all locations for sFLT1 are aggregated into only two compartments for

sFLT1: intracellular and extracellular. In reality, both consist of several subcompartments where

molecules can have different levels and be exposed to different conditions and therefore have

different reaction propensities. For example, the intracellular space represents the entire

endomembrane system of organelles (such as ER, Golgi, vesicles, and endosomes). While

traversing these organelles, intracellular sFLT1 undergoes a series of maturation and trafficking

steps that make a newly synthesized molecule more likely to be secreted over time, meaning

reaction propensities are not constant. Although our final model addresses this maturation

problem by including a maturation delay (τ) in between synthesis and secretion or degradation,

protein maturation is likely more variable and stochastic than can be represented with a fixed

time delay. sFLT1 could also localize to noncanonical compartments; both VEGFR1 and

VEGFR2 have been shown to localize to the nucleus under certain conditions in endothelial and

microvascular cells.[48] Outside the cell, since sFLT1 binds HSPGs in the extracellular matrix

(ECM) or on the cell surface, and also dimerizes with membrane-integral VEGFR isoforms,

extracellular sFLT1 can also be subdivided into cell-surface-associated, ECM-associated, and

freely soluble forms. Explicitly representing these subcompartments or maturation states would

add new molecular species and reactions, increasing model complexity and increasing the range

of possible behaviors in our model. However, they would also have significantly more

parameters, and estimating those parameter values would require quantitative data about sFLT1

organelle distribution or post-translational modification. As a separate but related issue, our

optimization to experimental data assumes that all extracellular sFLT1 is in conditioned media

and that protein detected in cell lysates is intracellular. However, extracellular sFLT1 bound to

ECM may not be aspirated with conditioned media, and extracellular sFLT1 associated with the

cell surface (either dimerized to membrane VEGFR or nonspecifically bound to HSPGs) could

be isolated with the lysate.

Our current model assumes that only sFLT1 affects the system, that we can omit other

molecules and reactions from our model, and that the processes we do model are well-described

by zeroth- and first-order kinetics with fixed rate constants. This framework does not address all

the complexity of cell biology; for example, factors affecting global gene expression, such as

ribosomal subunit expression and nutrient availability, could dynamically alter the sFLT1

production rate. Similarly, other secretory pathway cargo interacts with shared trafficking

machinery, so high production of other secretory proteins could reduce the trafficking factors

available to sFLT1 and reduce the effective secretion rate constant β.

sFLT1 binds other intracellular and extracellular components, which not only mediate its

effects on VEGF signaling but also can affect the dynamics we depict on our model, as bound

sFLT1 may not be available for trafficking reactions. sFLT1 acts as a ligand trap for extracellular

VEGF and PlGF, and also potentially as a dimerization partner for membrane-integral VEGF

receptor isoforms that creates dominant negative receptor complexes.[12] These networks of

biochemical reactions may have emergent properties inconsistent with our assumption of

invariant rate constants. For example, VEGF-A upregulates sFLT1 expression, which then

sequesters VEGF and reduces VEGF signaling.[37,49] This negative feedback loop could generate

pulses of sFLT1 expression or oscillations,[50] either of which would change the sFLT1 production

rate over time. Similar feedback effects could also affect proteins involved in secretion or

degradation. Alternatively, some evidence suggests sFLT1 can be generated by proteolytic

cleavage of membrane-integral VEGFR1,[51] a potential second production route for sFLT1, and a

model incorporating that reaction would also need to track membrane isoforms and potentially

be expanded to the level of gene expression and FLT1/VEGFR1 RNA alternative splicing.

sFLT1 also contains heparin-binding domains that nonspecifically bind glycoproteins on the

ECM and cell surface[52–54] and accumulates in the ECM.[55] sFLT1-ECM binding is important for

establishing and maintaining extracellular VEGF gradients that are directional cues in

angiogenesis. Importantly, protein glycosylation occurs in intracellular secretory pathway

organelles, especially the Golgi,[56] and interaction with intracellular glycoproteins is a potential

mechanism for sFLT1 retention in the Golgi. Adding such extensive additional detail to the

model would require substantial amounts of additional specific experimental data, and would

likely not impact on the core observed secretion dynamics as reproduced by the current model.

793   These caveats and considerations noted above can provide further directions for

794   development of the model of sFLT1 secretion developed here. Even without these additions, the

795   model is already well-behaved, recreates most experimental observations of sFLT1 secretion,

796   and identifies two classes of inhibitors or perturbations: those whose behavior is well represented

797   by the model, and those whose behavior seems contrary to the model and worthy of further

798   investigation.

References

1. Uhlén, M. *et al.* Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).

2. Palade, G. Intracellular aspects of the process of protein synthesis. *Science* **189**, 347–358

(1975).

3. Uhlén, M. *et al.* The human secretome. *Sci. Signal.* **12**, eaaz0274 (2019).

4. Munro, S. Localization of proteins to the Golgi apparatus. *Trends Cell Biol.* **8**, 11–15 (1998).

5. Sun, X., Mahajan, D., Chen, B., Song, Z. & Lu, L. A quantitative study of the Golgi

retention of glycosyltransferases. *J. Cell Sci.* **134**, jcs258564 (2021).

6. Du, Y. *et al.* Syntaxin 6-mediated Golgi translocation plays an important role in nuclear

functions of EGFR through microtubule-dependent trafficking. *Oncogene* **33**, 756–770

(2014).

7. Huang, S. S. & Huang, J. S. Rapid turnover of the platelet-derived growth factor receptor in

sis-transformed cells and reversal by suramin. Implications for the mechanism of autocrine

transformation. *J. Biol. Chem.* **263**, 12608–12618 (1988).

8. Mittar, S. *et al.* VEGFR1 receptor tyrosine kinase localization to the Golgi apparatus is

calcium-dependent. *Exp. Cell Res.* **315**, 877–889 (2009).

9. Sorkin, A. & von Zastrow, M. Endocytosis and signalling: intertwining molecular networks.

Nat. Rev. Mol. Cell Biol. **10**, 609–622 (2009).

10. Sorkin, A. & Goh, L. K. Endocytosis and intracellular trafficking of ErbBs. *Exp. Cell Res.*

314, 3093–3106 (2008).

11. Kendall, R. L. & Thomas, K. A. Inhibition of vascular endothelial cell growth factor activity

by an endogenously encoded soluble receptor. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 10705–

10709 (1993).

12. Kendall, R. L., Wang, G. & Thomas, K. A. Identification of a Natural Soluble Form of the

Vascular Endothelial Growth Factor Receptor, FLT-1, and Its Heterodimerization with KDR.

Biochem. Biophys. Res. Commun. **226**, 324–328 (1996).

13. Kinghorn, K. *et al.* A defined clathrin-mediated trafficking pathway regulates

sFLT1/VEGFR1 secretion from endothelial cells. *Angiogenesis* **27**, 67–89 (2024).

14. Hiratsuka, S., Minowa, O., Kuno, J., Noda, T. & Shibuya, M. Flt-1 lacking the tyrosine

kinase domain is sufficient for normal development and angiogenesis in mice. *Proc. Natl.*

Acad. Sci. U. S. A. **95**, 9349–9354 (1998).

15. Bautch, V. L. VEGF-Directed Blood Vessel Patterning: From Cells to Organism. *Cold*

Spring Harb. Perspect. Med. **2**, a006452 (2012).

16. Wu, F. T. H. *et al.* A systems biology perspective on sVEGFR1: its biological function,

pathogenic role and therapeutic use. *J. Cell. Mol. Med.* **14**, 528–552 (2010).

17. Ambati, B. K. *et al.* Corneal avascularity is due to soluble VEGF receptor-1. *Nature* **443**,

993–997 (2006).

18. Chappell, J. C., Taylor, S. M., Ferrara, N. & Bautch, V. L. Local Guidance of Emerging

Vessel Sprouts Requires Soluble Flt-1. *Dev. Cell* **17**, 377–386 (2009).

19. Hashambhoy, Y. L., Chappell, J. C., Peirce, S. M., Bautch, V. L. & Mac Gabhann, F.

Computational modeling of interacting VEGF and soluble VEGF receptor concentration

gradients. *Front. Physiol.* **2**, 62 (2011).

20. Kappas, N. C. *et al.* The VEGF receptor Flt-1 spatially modulates Flk-1 signaling and blood

vessel branching. *J. Cell Biol.* **181**, 847–858 (2008).

21. Maynard, S. E. *et al.* Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may

844   contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J. Clin.*

845   *Invest.* **111**, 649–658 (2003).

846   22. Jung, J.-J. *et al.* Secretion of Soluble Vascular Endothelial Growth Factor Receptor 1

847   (sVEGFR1/sFlt1) Requires Arf1, Arf6, and Rab11 GTPases. *PLoS ONE* **7**, e44572 (2012).

848   23. Sarabipour, S. *et al.* Trafficking dynamics of VEGFR1, VEGFR2, and NRP1 in human

849   endothelial cells. *PLOS Comput. Biol.* **20**, e1011798 (2024).

850   24. Clegg, L. E. & Mac Gabhann, F. Molecular mechanism matters: Benefits of mechanistic

851   computational models for drug development. *Pharmacol. Res.* **99**, 149–154 (2015).

852   25. Glass, D. S., Jin, X. & Riedel-Kruse, I. H. Nonlinear delay differential equations and their

853   application to modeling biological network motifs. *Nat. Commun.* **12**, 1788 (2021).

854   26. Wu, F. T. H. *et al.* Computational kinetic model of VEGF trapping by soluble VEGF

855   receptor-1: effects of transendothelial and lymphatic macromolecular transport. *Physiol.*

856   *Genomics* **38**, 29–41 (2009).

857   27. Wu, F. T. H. *et al.* VEGF and soluble VEGF receptor-1 (sFlt-1) distributions in peripheral

858   arterial disease: an in silico model. *Am. J. Physiol. Heart Circ. Physiol.* **298**, H2174-2191

859   (2010).

860   28. Clegg, L. E. & Mac Gabhann, F. A computational analysis of in vivo VEGFR activation by

861   multiple co-expressed ligands. *PLoS Comput. Biol.* **13**, e1005445 (2017).

862   29. Clegg, L. E. & Mac Gabhann, F. A computational analysis of pro-angiogenic therapies for

863   peripheral artery disease. *Integr. Biol. Quant. Biosci. Nano Macro* **10**, 18–33 (2018).

864   30. Hornig, C. *et al.* Release and Complex Formation of Soluble VEGFR-1 from Endothelial

865   Cells and Biological Fluids. *Lab. Invest.* **80**, 443–454 (2000).

31. Schindelin, J. *et al.* Fiji - an Open Source platform for biological image analysis. *Nat.*

Methods **9**, 10.1038/nmeth.2019 (2012).

32. Eden, E. *et al.* Proteome half-life dynamics in living human cells. *Science* **331**, 764–768

(2011).

33. Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control.

Nature **473**, 337–342 (2011).

34. Hausser, J., Mayo, A., Keren, L. & Alon, U. Central dogma rates and the trade-off between

precision and economy in gene expression. *Nat. Commun.* **10**, 68 (2019).

35. Akaike, H. Information Theory and an Extension of the Maximum Likelihood Principle. in

Selected Papers of Hirotugu Akaike (eds. Parzen, E., Tanabe, K. & Kitagawa, G.) 199–213

(Springer, New York, NY, 1998). doi:10.1007/978-1-4612-1694-0_15.

36. Bozdogan, H. Model selection and Akaike's Information Criterion (AIC): The general theory

and its analytical extensions. *Psychometrika* **52**, 345–370 (1987).

37. Ahmad, S. *et al.* Autocrine activity of soluble Flt-1 controls endothelial cell function and

angiogenesis. *Vasc. Cell* **3**, 15 (2011).

38. Hetz, C., Zhang, K. & Kaufman, R. J. Mechanism, regulation and functions of the unfolded

protein response. *Nat. Rev. Mol. Cell Biol.* **21**, 421–438 (2020).

39. Manickam, V. *et al.* Regulation of vascular endothelial growth factor receptor 2 trafficking

and angiogenesis by Golgi localized t-SNARE syntaxin 6. *Blood* **117**, 1425–1435 (2011).

40. Jaffe, E. A., Nachman, R. L., Becker, C. G. & Minick, C. R. Culture of Human Endothelial

Cells Derived from Umbilical Veins. IDENTIFICATION BY MORPHOLOGIC AND

IMMUNOLOGIC CRITERIA. *J. Clin. Invest.* **52**, 2745–2756 (1973).

41. Bauer, A. L., Jackson, T. L. & Jiang, Y. A cell-based model exhibiting branching and

anastomosis during tumor-induced angiogenesis. *Biophys. J.* **92**, 3105–3121 (2007).

42. Qutub, A. A. & Popel, A. S. Elongation, proliferation & migration differentiate endothelial

cell phenotypes and determine capillary sprouting. *BMC Syst. Biol.* **3**, 13 (2009).

43. Perfahl, H. *et al.* Multiscale Modelling of Vascular Tumour Growth in 3D: The Roles of

Domain Size and Boundary Conditions. *PLoS ONE* **6**, e14790 (2011).

44. Walpole, J., Mac Gabhann, F., Peirce, S. M. & Chappell, J. C. Agent-based computational

model of retinal angiogenesis simulates microvascular network morphology as a function of

pericyte coverage. *Microcirc. N. Y. N 1994* **24**, (2017).

45. Song, M. & Finley, S. D. Mechanistic characterization of endothelial sprouting mediated by

pro-angiogenic signaling. *Microcirc. N. Y. N 1994* **29**, e12744 (2022).

46. Chappell, J. C. *et al.* Flt-1 (VEGFR-1) coordinates discrete stages of blood vessel formation.

Cardiovasc. Res. **111**, 84–93 (2016).

47. Walpole, J. *et al.* Agent-based model of angiogenesis simulates capillary sprout initiation in

multicellular networks. *Integr. Biol. Quant. Biosci. Nano Macro* **7**, 987–997 (2015).

48. Silva, J. A. F., Qi, X., Grant, M. B. & Boulton, M. E. Spatial and temporal VEGF receptor

intracellular trafficking in microvascular and macrovascular endothelial cells. *Sci. Rep.* **11**,

17400 (2021).

49. Saito, T. *et al.* VEGF-A induces its negative regulator, soluble form of VEGFR-1, by

modulating its alternative splicing. *FEBS Lett.* **587**, 2179–2185 (2013).

50. Alon, U. *An Introduction to Systems Biology: Design Principles of Biological Circuits*.

(Chapman and Hall/CRC, New York, 2006). doi:10.1201/9781420011432.

51. Rahimi, N., Golde, T. E. & Meyer, R. D. Identification of Ligand-Induced Proteolytic

Cleavage and Ectodomain Shedding of VEGFR-1/FLT1 in Leukemic Cancer Cells. *Cancer*

Res. **69**, 2607–2614 (2009).

52. Roeckl, W. *et al.* Differential Binding Characteristics and Cellular Inhibition by Soluble

VEGF Receptors 1 and 2. *Exp. Cell Res.* **241**, 161–170 (1998).

53. Park, M. & Lee, S. T. The fourth immunoglobulin-like loop in the extracellular domain of

FLT-1, a VEGF receptor, includes a major heparin-binding site. *Biochem. Biophys. Res.*

Commun. **264**, 730–734 (1999).

54. Sela, S. *et al.* Local retention versus systemic release of soluble VEGF receptor-1 are

mediated by heparin-binding and regulated by heparanase. *Circ. Res.* **108**, 1063–1070

(2011).

55. Orecchia, A. *et al.* Vascular endothelial growth factor receptor-1 is deposited in the

extracellular matrix by endothelial cells and is a ligand for theα5β1 integrin. *J. Cell Sci.* **116**,

3479–3489 (2003).

56. Reily, C., Stewart, T. J., Renfrow, M. B. & Novak, J. Glycosylation in health and disease.

Nat. Rev. Nephrol. **15**, 346–366 (2019).

929 Figures

930 Figure 1

931
932

Figure 1. Modeling sFLT1 secretion with ordinary differential equations (ODEs). (A) Model of sFLT1

secretion dynamics describing the impact of production (rate α), secretion (rate constant β), intracellular degradation

(γ), and extracellular degradation (δ) on the concentrations of intracellular (I) and extracellular (X) sFLT1 as a

system of ordinary differential equations (ODEs). Created with Biorender.com. **(B)** Comparison of simulation (top

row) and experimental (bottom row) approaches for studying sFLT1 secretion in the constitutive secretion scenario

(left column) and pulse-chase secretion scenario (right column). **(C)** Simulated time courses ($n=100$) of extracellular

(X, top) and intracellular (I, bottom) sFLT1 secretion for pulse-chase (left) and constitutive (right) cases using the

ODE model. X was normalized to its value at 8h (upper left) or 24h (upper right), while I was normalized to its value

at 0h. The highlighted region (P) marks the 20-minute pulse. **(D)** Number of intracellular sFLT1 (I) molecules per

cell over time and **(E)** distribution of steady state intracellular sFLT1 (I_{SS}) during simulations of constitutive

secretion using the ODE model.

Figure 2

946
947

Figure 2. Modeling sFLT1 secretion with delay differential equations (DDEs). (A) Model of sFLT1 secretion
dynamics as a system of delay differential equations (DDEs), extending the ODE model (Figure 1A) with a fixed
maturation delay (τ) affecting secretion and intracellular degradation processes. Created with Biorender.com. **(B)**
Simulated time courses ($n = 594$) of extracellular (X, top) and intracellular (I, bottom) sFLT1 secretion for pulse-
chase (left) and constitutive (right) cases using the base DDE model. X was normalized to its value at 8h (upper left)
or 24h (upper right), while I was normalized to its value at 0h. The highlighted region (P) marks the 20-minute
pulse.

Figure 3

Figure 3. Comparing candidate models of sFLT1 secretion. (A) Structure of candidate model equations and

equation term options for each process when present or absent in a given model. In the production decay term, the

value of *[chase]* is 1 during the chase phase of pulse-chase simulations and 0 otherwise. **(B)** Top: Processes included

(filled) or excluded (empty) in each model. ε: internalization, κ: production decay, τ: maturation delay. Middle: Cost

(goodness of fit) of each model, defined as the sum of squared differences (SSD) between experimental data points

and the lowest cost simulation of each model, with colors indicating the contribution of each experimental dataset.

X: extracellular sFLT1, *I*: intracellular sFLT1. Bottom: Corrected Akaike Information Criterion (AIC$_C$) scores for

each model. **(C)** Impact of additional processes on the best-fit cost of the candidate models. Model nodes (points)

are plotted as minimum model cost versus number of parameters. Edges (arrows) represent transitions between

candidate model structures by adding the labeled process (τ (maturation delay): pink arrows, κ (production decay):

blue arrows, ε (internalization): gray arrows. The vertical height of each arrow represents the decrease in cost from

adding the labeled process to a given model.

Figure 4

Figure 4. Properties of optimized parameter sets from the DDE model of sFLT1 secretion. (A) Distributions of

initial (yellow) and optimized (black) model parameter values for the delay differential equation model after filtering

for low-cost fits ($n=594$). Y axes are normalized such that each distribution has a maximum density of 1. **(B)**

Correlations between α, β, and γ values in optimized parameter sets. Each point represents the observed values of

the listed parameters in a single run of the delay differential equation model after filtering for low-cost fits ($n=594$).

(C) Distributions of initial (yellow) and optimized (black) values of calculated compound parameters $c_1=\alpha\beta$ and

$c_2=\beta+\gamma$ for the delay differential equation model after filtering for low-cost fits ($n=594$). **(D-G)** Correlation between

intracellular steady state sFLT1 (I_{SS}) and **(D)** $\alpha/(\beta+\gamma)$, **(E)** α, **(F)** β, **(G)** γ. Dashed lines indicate theoretical bounds

($\min(I_{SS}) = c_1/c_2^2$, $\min(\alpha) = c_1/c_2$, $\min(\beta) = \min(\gamma) = 0$, $\max(\beta) = \max(\gamma) = 0$; see *Supplemental Results*).

Bounds and highlighted trendlines are calculated using median values of c_1 and c_2 (Table 1).

Figure 5

Figure 5. Analysis of simulated sFLT1 kinetic process fluxes. (A) Reaction network graph showing process

fluxes (Φ). **(B)** Equations for process fluxes. **(C)** Process fluxes over time during simulation of constitutive sFLT1

secretion. **(D)** Rates of change of extracellular sFLT1 (dX/dt = Φ(Secr) − Φ(XDeg)) and intracellular sFLT1 (dX/dt

= Φ(Prod) − Φ(Secr) − Φ(IDeg)) during constitutive secretion. **(E)** Rates of change of extracellular sFLT1 (dX/dt =

Φ(Secr) − Φ(XDeg)) and intracellular sFLT1 (dI/dt = Φ(Prod) − Φ(Secr) − Φ(XDeg)) during constitutive secretion.

(F) Constitutive flux distributions for time-constant processes. **(G)** Rates of change of extracellular and intracellular

sFLT1 during pulse-chase secretion. In C, D, F, and G, each line represents a different optimized parameter set.

Prod = production, Secr = secretion, IDeg = intracellular degradation, XDeg = extracellular degradation.

Figure 6

Figure 6. Local and global univariate sensitivity analysis of the sFLT1 secretion DDE model. (A) Relative

sensitivity of sFLT1 secretion output variables to local changes in input parameters during simulation of constitutive

secretion. The base case is the median parameter set, and all other cells represent the observed fractional increase in

the output variable per fractional increase in the parameter. X_{72h}: extracellular sFLT1 after 72 hours; I_{72h}:

intracellular sFLT1 after 72 hours; T_{50_X}: time to half-steady-state for extracellular sFLT1. **(B)** Global univariate

sensitivity analysis of extracellular and intracellular sFLT1 showing the effect of multiplying each input parameter

by a scaling factor during constitutive secretion. The base case (scaling factor 1) is the median parameter set from

the optimized parameter sets (Table 1).

1005

1006   Figure 7

1007

 Figure 7. Simulated chemical inhibition and genetic downregulation of sFLT1 secretion. **(A)** Simulation

timeline schematic for chemical inhibition (top) and genetic downregulation (bottom) of constitutive secretion. $p =$

p' represents change of parameter values to apply inhibitor effect. **(B)** Effects of chemical inhibition of individual

parameters by 90% on extracellular sFLT1 (X) and intracellular sFLT1 (I) at 18h and 72h, expressed as log2 fold

changes relative to the control (uninhibited) case. **(C)** Time courses for absolute concentrations of extracellular (X)

and intracellular (I) sFLT1 with 90% chemical inhibition of each parameter. The impact of inhibiting each process is

how different each line is from the prediction of the median, unmodified parameter set (black line). Black points,

experimental ELISA data for the uninhibited case (Hornig et al., 2000). **(D)** Annotated curves for chemical

inhibition of β (red) and γ (orange) reproduced from (C). Horizontal lines indicate experimentally observed relative

changes in extracellular (X_{18h}) and intracellular (I_{18h}) sFLT1 18 hours after drug addition. Vertical lines indicate the

fraction inhibition of the target parameter required to reproduce experimental results. **(E)** Effects of genetic

downregulation of individual parameters by 90%, as described in (B). **(F)** Time courses for absolute concentrations

of extracellular (X) and intracellular (I) sFLT1 with 90% genetic downregulation of each parameter, as described in

(C). **(G)** Annotated curves for genetic downregulation of β (red) and γ (orange), as described in (D).

1022

1023

1024

1025   Figure 8

1026

1027

Figure 8. Simulating chemical and genetic inhibition of sFLT1 secretion from different initial parameter sets.

(A) Annotated curves for chemical inhibition of β (secretion, first 3 columns) and γ (intracellular degradation,

rightmost column) at 18 hours. Horizontal lines indicate experimentally observed relative changes in extracellular

(X_{18h}) and intracellular (I_{18h}) sFLT1 18 hours after addition of brefeldin, chlorpromazine, TATNSF700, or

chloroquine. Colored vertical lines mark the fraction inhibition required to match the observed change in sFLT1

when starting from the parameter set with the matching β value. **(B)** Annotated curves for genetic inhibition of β

(secretion rate constant) at 18 hours post media change. Horizontal lines indicate experimentally observed relative

changes in extracellular (X_{18h}) and intracellular (I_{18h}) sFLT1 after 18 hours after media change in cells transfected

with siRNA targeting the listed secretion-linked protein 48 hours previously.[13] Colored vertical lines mark the

fraction inhibition required to match the observed change in sFLT1 when starting from the parameter set with the

matching β value.

Tables

Table 1

1041 **Table 1.** Statistics for the distribution of optimized values of core model parameters (n=594 parameter

1042 sets). SD = standard deviation, CV = coefficient of variation, MAD = median absolute deviation, IQR =

1043 interquartile range.

parameter	α	β	γ	δ	τ
unit	#/cell/h	1/h	1/h	1/h	h
mean	5.177×10^5	7.229×10^{-2}	1.005×10^{-1}	5.725×10^{-2}	1.958
SD	7.692×10^5	6.411×10^{-2}	6.419×10^{-2}	1.444×10^{-3}	2.046×10^{-2}
CV	1.487	8.868×10^{-1}	6.389×10^{-1}	2.523×10^{-2}	1.045×10^{-2}
median	1.419×10^5	5.123×10^{-2}	1.215×10^{-1}	5.743×10^{-2}	1.958
MAD	1.470×10^5	6.639×10^{-2}	6.649×10^{-2}	2.361×10^{-5}	1.570×10^{-2}
IQR	6.310×10^5	1.337×10^{-1}	1.337×10^{-1}	3.149×10^{-5}	2.097×10^{-5}
median/IQR	4.447	2.609	1.1	5.483×10^{-4}	1.071×10^{-5}
min	3.524×10^4	1.592×10^{-3}	1.842×10^{-4}	4.577×10^{-2}	1.695
max	4.563×10^6	1.745×10^{-1}	1.767×10^{-1}	7.296×10^{-2}	2.197

1044

1045 Table 2

1046 **Table 2.** Statistics for the distribution of compound parameters $c_1 = \alpha\beta$ and $c_2 = \beta + \gamma$ resulting from

1047 the optimized parameter values (Table 1). SD = standard deviation, CV = coefficient of variation, MAD =

1048 median absolute deviation, IQR = interquartile range.

1049

parameter	$c_1 = \alpha\beta$	$c_2 = \beta + \gamma$
unit	#/cell/h^2	1/h
mean	7.252×10^3	1.728×10^{-1}
SD	1.316×10^2	6.418×10^{-4}
CV	1.814×10^{-2}	3.715×10^{-2}
median	$\mathbf{7.269 \times 10^3}$	$\mathbf{1.727 \times 10^{-1}}$
MAD	1.207	2.302×10^{-7}
IQR	1.606	3.145×10^{-7}
IQR/median	2.210×10^{-4}	1.821×10^{-6}
min	5.999×10^3	1.685×10^{-1}
max	8.460×10^3	1.835×10^{-1}

1050

1051 Table 3

1052 **Table 3.** Predicted inhibition strengths of chemical inhibitors based on comparing experimental data[13] to

1053 model simulations. *: Simulations match the direction of the experimental perturbation but cannot

1054 achieve the observed strength of impact.

Treatment	Inhibited process	Parameter affected	Percent inhibition in model to match	
			X_{18h}	I_{18h}
brefeldin	Golgi assembly and maturation (Misumi et al., 1986; Fujiwara et al., 1988; Lippincott-Schwartz et al., 1989)	β	89	> 100*
chlorpromazine	clathrin-mediated vesicle trafficking (Wang et al., 1993)	β	58	> 100*
TATNSF700	exocytosis; vesicle fusion (Matsushita et al., 2005)	β	54	> 100*
chloroquine	lysosomal degradation (Mauthe et al., 2018; Lie and Schofield, 1973)	γ	98	69

1055

1056

1057 Table 4

1058 **Table 4.** Predicted inhibition strengths of genetic inhibitors based on comparing experimental data[13] to

1059 model simulations. *: Simulations match the direction of the experimental perturbation but cannot achieve

1060 the observed strength of impact. †: Simulations cannot match experimental data as experimental

1061 perturbation has opposite outcome to simulated perturbation.

Treatment	Target role	Detailed role	Parameter affected	Percent inhibition in model to match	
				X_{18h}	I_{18h}
siRAB27a	vesicle trafficking to plasma membrane	small GTPase coordinating vesicle transport along actin (Wu et al., 2001; Alzahofi et al., 2020)	β	97	$< 0^\dagger$
siSTX3	vesicle trafficking to plasma membrane	t-SNARE at plasma membrane (Sharma et al., 2006)	β	82	$> 100*$
siSTX6	vesicle trafficking from Golgi	t-SNARE in trans-Golgi network (Bock et al., 1997; Wendler and Tooze, 2001)	β	98	$< 0^\dagger$
siVAMP3	vesicle trafficking to plasma membrane	v-SNARE in endosome-derived vesicles (Hu et al, 2007)	β	78	$< 0^\dagger$

1062