Dissection and internal observation of the giant tropical bont tick *Amblyomma variegatum*

Naomie Pature[1,2,3], Nonito Pagès[1,2], Valérie Rodrigues[1,2] and Damien F. Meyer*[1,2]

[1] CIRAD, UMR ASTRE, F-1970 Petit-Bourg – Guadeloupe, France

[2] ASTRE, CIRAD, INRAE, Univ Montpellier – Montpellier, France

[3] Université des Antilles, École doctorale n° 636 DEECA, Campus de Fouillole, Pointe-à-Pitre – Guadeloupe, France

*Corresponding author

Correspondence: damien.meyer@cirad.fr

ABSTRACT

Ticks are ectoparasites that feed on the blood of animals and humans and are among the arthropods responsible for numerous human and animal diseases worldwide. The tropical bont tick, *Amblyomma variegatum*, is a significant natural vector of *Ehrlichia ruminantium*, the causative agent of heartwater. This disease is endemic in many countries, including those in sub-Saharan Africa and certain Caribbean islands. Due to the potential of heartwater and its vector to pose an economic and health threats to non-endemic regions, they have been designated as high-consequence foreign animal diseases and pests by the United States Department of Agriculture's Animal and Plant Health Inspection Service. Understanding the ecology and biology of *A. variegatum* is critical for developing effective control methods to limit its spread to other geographic regions. A key step in this process is the isolation of tick organs and further elucidation of their functions (i.e. vector competence). However, the literature contains limited documentation of the complete internal structure of *Amblyomma* tick species. In this study, we describe the internal structure of unfed nymphs, partially engorged males, and unfed and partially engorged

females of the tick species *A. variegatum*. The objective is to facilitate the identification and the localization organs of interest while highlighting the morphological differences between the tropical bont tick and other tick species. This foundational work will support the development of various applications following the isolation of tick organs. These findings not only deepen our understanding of *Amblyomma variegatum* anatomy but also provide a critical foundation for tackling the broader challenges of tick-borne diseases and advancing vector control strategies in both endemic and non-endemic regions.

Keywords: *Amblyomma variegatum*, heartwater, dissection, tick organs, anatomy, hard-bodied ticks

Introduction

The earliest known study of insects was initiated by the Greek naturalist and philosopher Aristotle, who emphasized their insignificance (Friedrich et al., 2014). The internal anatomy of arthropods became the subject of extensive study for many centuries (Kennedy et al., 2021). Indeed, the study of insects advanced with Galileo's invention of the telescope in the early 17th century, which was adapted for observing insects. This progress was followed by the work of Antonie van Leeuwenhoek, considered the father of microscopy, whose studies included dissection of spiders (Cocquyt, 2022).

The process of dissection under a microscope can be as straightforward as exploring the internal phenome of arthropods (Deans et al., 2012). This was illustrated by Zacaï *et al.*, who utilized microscopy to visualize an arthropod fossil, *Sidneyia inexpectans*, dating back several million years ago (Zacaï et al., 2016). They emphasized the internal digestive system associated with the fossilized gut contents and compared it to that of modern chelicerates. Since then, a variety of dissection techniques have been developed for use with diverse arthropods species. Primarily, the dissection of larger arthropods offers a rapid approach to conducting internal investigations. Identifying the internal organs of hard ticks can be achieved through dissection and subsequent removal of internal organs. The use of simple tools such as fine tweezers and sharpened blades is beneficial during the dissection process. Similarly, the application of strong tape or wax facilitates the dissection process.

Even with the advent of several powerful imaging techniques such as X-ray, microtomography (μCT), or confocal laser scanning microscopy (CLSM), light-optical microscopy remains the best method for analyzing morphological diversity (Sasso Porto et al., 2016). To date, the fragility of the internal structure of arthropods has meant that microscopic dissection involves a risk of tissue damage. Consequently, the manipulator's skill in performing dissections is of paramount importance to prevent the destruction of organs, particularly in morphological studies. To address this limitation, a recent study has focused on the development of spectral-domain optical coherence tomography (SD-OCT), which enables non-invasive imaging while preserving the anatomical features of tick specimens (Lee et al., 2021). However, this method does not permit the collection of arthropod organs of interest for interaction studies with pathogens (Deans et al., 2012).

The tropical bont tick, *Amblyomma variegatum* (Fabricius, 1794), is an obligate hematophagous arthropod belonging to the phylum Arthropoda, subphylum Chelicerata, Class Arachnida, subclass Acari, and order Ixodida (Camargo-Mathias, 2018). This tropical tick is also referred to as the Senegalese tick in the Guadeloupe archipelago and the Antigua gold tick in Antigua (Pegram et al., 2004). It belongs to the family Ixodidae, characterized by a hard body with a dorsal shield (scutum) that may be ornamented in the case of males, or may not be ornamented for females.

The tick is the vector for heartwater, a major disease in the Caribbean region and is frequently associated with dermatophilosis (Prine and Hodges, 2012). Furthermore, this tick contributes significantly to lameness and the formation of deep abscesses, which can ultimately lead to immunosuppression and metabolic disorders in small ruminants (Barré, 1989). Consequently, the presence of *A. variegatum* results in considerable economic and health losses. More than ten species of *Amblyomma* ticks transmit *E. ruminantium* in Africa. Of these, *A. variegatum* and *A. hebraeum* are identified as the main vectors (Bezuidenhout, 1987). The majority of heartwater-endemic areas overlap with the geographic distribution of *Amblyomma* ticks. The vector potential of these ticks depends on factors such as their abundance, activity, and adaptability to their environment (Uilenberg, 1983).

The tropical bont tick, *A. variegatum*, is not only a significant vector of pathogens in endemic regions but also poses a potential threat to non-endemic areas due to livestock movement and climate change (Estrada-Pena et al., 2012). These phenomena may play an indirect role in the abundance and distribution of ticks via changes in hosts. (Gilbert, 2021). Knowing its anatomy is crucial for studying the tick pathogen interactions, which are important part of potential vector control strategies addressing the socioeconomic and health impacts of tick-borne diseases, which disproportionately affect resource-limited regions. While studies on *Rhipicephalus microplus* and *Ixodes scapularis* have advanced our understanding of tick physiology, the anatomical and functional adaptations of *A. variegatum* remain understudied, necessitating detailed investigation. These adaptations include reproductive efficiency, feeding behavior, and ecological resilience, all of which contribute to its status as a major vector.

The morphological identification of the internal organs of the ticks is of significant value in
entomology and parasitology, as it facilitates the development of effective control methods, whether
natural or chemical. To further develop control technologies, it is necessary to anatomize the internal
structure of ticks. The study of the internal anatomy of arthropods is of great interest due to the
remarkable diversity observed, particularly regarding the specific barriers such as target organs they
possess. Given the widespread distribution of *A. variegatum*, particularly in some Caribbean islands,
there is a clear need to study the tick and its role as a pathogen vector. The impetus for composing this
article was the desire to produce a paper identifying the organs of the *A. variegatum* tick and
determining the optimal dissection procedures.

Despite the numerous dissection documents available for accurately identifying the internal organs of
ticks, particularly those in the genus *Rhipicephalus*, there remains a pressing need for greater emphasis
on the internal observations of *Amblyomma* ticks (Cossio-Bayugar et al., 2023; Tidwell et al., 2021). High-
quality histologic images of the ixodid tick reproductive and digestive systems are available online
(Camargo-Mathias, 2018; de la Vega et al., 2012; Mathias, 2013). Additionally, excellent scanning
electron microscopy images of male *Amblyomma* testes and female *Amblyomma* intestines are available
(Anholeto et al., 2015; Caperucci et al., 2009). However, apart from the 2009 article by Edwards *et al.*,
there are few studies based on the dissection of *Amblyomma* ticks (Edwards, 2009). Despite the pictorial
instructions provided in this guide, two different *Amblyomma* species had to be used to visualize the
entirety of the tick's internal organs.

Here, we provide an anatomical observation of the internal cavity of *A. variegatum*, from nymphs to
adult ticks at different stages of engorgement. The dissected organs are described and illustrated in detail
in this document. This report will facilitate the accurate and reproducible sampling of *A. variegatum* tick
tissues, enabling highly precise studies of tick pathogen interactions, and, *in fine*, the development of
new tick control technologies.

Approach

Materials for the dissection
- Forceps with pointed tips
- Petri dish or glass dissection plate
- Disposable scalpel (No. 3 handle with a No. 11 surgical blade)
- Double-sided tape
- Falcon tubes
- Pins (optional)
- Dulbecco's phosphate buffered saline
- 70% Ethanol
- Sterile ultrapure water
- Microcapillary tubes for hemolymph collection
- Pipette tips
All the dissection tools were treated aseptically before use. Specifically, all the forceps were
autoclaved, then immersed in 70% ethanol for 5 minutes and rinsed with sterile ultrapure water. Tools
were allowed to air dry on a clean paper towel. All the petri dishes were wiped or sprayed with 70%
ethanol and allowed to air dry. In certain cases, glass petri dishes could be used to avoid the damaging of
pointed forceps. A LEICA M205FA fluorescence stereomicroscope equipped with Fluocombi (LEICA,
Wetzlar, Germany) was used for image acquisition.
169
Troubleshooting and Practical Notes
During dissections, care must be taken to avoid tissue tearing, especially for fragile organs like ovaries
and synganglion. Using fine, aseptic tools and working under a stereomicroscope significantly reduces the
risk of damage. Adhesive tapes and clean dissecting surfaces are essential to maintain tissue integrity and
minimize contamination. For less experienced researchers, it is advisable to practice dissections on
nymphal or male ticks, which have simpler anatomy, before attempting engorged females. If tissues are
intended for downstream analyses, such as transcriptomics or proteomics, meticulous handling and
immediate storage in appropriate solutions (r.g., PBS or RNAlater) are critical to preserve RNA or protein
quality.
179
Tick general dissection
Tick rearing
For all experiments, batches of unfed *A. variegatum* ticks were reared as a colony for more than 13
years at the CIRAD animal facility in Guadeloupe. Larvae and nymphs were placed on the ear side of a
goat in a Mölnlycke Tubifast green line bandage, secured with Kamar adhesive (Vital Concept, Loudeac,
France). Following the completion of feeding and subsequent detachment, all the ticks were collected
and allowed to molt in a container maintained at room temperature and 70-80% of relative humidity.
After ecdysis, nymphs were used for dissections, while adult females were placed on the side of a goat
until semi-engorgement for some and detachment for others.
189
Preparation of the tick
Prior to the dissection, the ticks were washed twice for 30 seconds in ultrapure water and once in PBS
in three 0.5 mL Eppendorf microtubes to remove impurities on the tick's cuticle. During the washing
process, the tubes containing the ticks were gently inverted by hand. As *A. variegatum* ticks are rapid
crawlers, they could optionally be cooled on ice to slow them down. Following the washing step, the ticks
were completely dried on Kimtech paper towels (Kimberly-Clark, Nanterre, France) before being
immobilize for dissection.
The fixation of the tick was performed by immobilizing a single tick, with its ventral surface facing
down, on a piece of double-sided adhesive tape within a Petri dish (Fig. 1). To ensure fixation of the tick,
all four legs were extended and secured to the double-sided tape with tweezers. For nymphs and unfed
adults, multiple ticks could be fixed in a single Petri dish. Once immobilized, a few drops of sterile, filtered
Dulbecco's Phosphate Buffered Saline (PBS) 1X were applied to cover the entire tick body. This prevented
desiccation of the tick's tissues and facilitated dissection by swelling the internal organs.

Hemolymph and saliva collection

For hemolymph collection, the same procedure was used to wash the external structure of the tick.
The ticks were then fixed on a double-sided tape in the same manner as for the dissection. To collect the
hemolymph, the first two legs of the tick were fully extended. Then, the most distal joint of the tick's
tarsus was cut with a scalpel, and the drop was collected with a micropipette through a microcapillary
tube (Fig. 1B). For larger volumes of hemolymph, several legs may be amputated and the tick's body was
gently squeezed to promote drainage.

To collect saliva, semi-engorged ticks that had been removed from animals or artificially fed were
used to obtain larger amounts of saliva. After washing the tick as described above, it was immobilized on
double-sided tape placed in the upper part of a petri dish, with its ventral surface facing up to expose its
mouthparts, including the rostrum and palps. Another piece of double-sided tape was placed near the
tick's basis capitula on the edge of the petri dish. A capillary tube was pressed between the hypostome
and palps, ensuring the pedipalps remained outside the capillary tube.

A solution of pilocarpine (20mg/ml) was prepared, and 50 µL was gently injected into the tick's
stigmata or spiracles (Sp) (Fig 10A) using a Hamilton syringe. Salivation was observed, and the saliva was
collected using a pipette tip inserted into a micropipette. If the tick did not salivate after 10 minutes, an
additional 20 µL of pilocarpine was injected. The saliva was stored at -80°C for further analysis or use.

Dissecting the external structure: Scutum/alloscutum/conscutum removal

To properly dissect the tropical bont tick, *A. variegatum*, it is fundamental to know the external
appearance of the tick, which is the natural vector of the bacterium that causes heartwater in ruminants.
The overall appearance of the tick body is similar to that of parasitiform acarines, such as mites.

The tick body is divided into two main regions: the gnathosoma (capitulum) and the idiosoma (body
region (Fig. 1). The capitulum, considered as the "head", is located in the anterior part and bears the
mouthparts, including the chelicerae, hypostome, and palps. However, this designation requires caution
because the eyes and brain are situated inside the body region. Additionally, the basis capituli, located in
the capitulum, bears two paired structures known as porose areas.

The idiosoma is divided into the anterior podosoma, containing the genital gonopore and the four
pairs of legs, and the posterior opisthosoma, which includes the spiracular plates and the anal pore
(Sonenshine and Simo, 2021). Additionally, the idiosoma includes the walking legs. Ixodid ticks, often
referred as "hard-bodied ticks", are characterized by their teardrop shape and a dorsal shield or scutum
(Cupp, 1991).

In male ixodid ticks, the scutum, known as the conscutum, covers the entire dorsum, preventing the
ingestion of large quantities of blood during blood feeding. *A. variegatum* males are unique in having an
ornate conscutum with bright orange colors and a dark lateral background (Fig. 1B, 6A).

In contrast, the scutum of adult females, nymphs and larvae, does not cover the entire dorsum. The
female scutum consists of a large subtriangular plate near to the capitulum, while the remaining two-
thirds of the idiosoma is made up of a less sclerotized plate, the alloscutum (Fig 1A, 4A). During feeding,
this part of the female tick expands, allowing it to weigh up to 100 times its initial weight.

Furthermore, the alloscutum of larvae and nymphs tends to enlarge as they ingest blood. Lateral to
the scutum/ conscutum, two hemispherical, dark-colored orbital eyes can be observed (Fig. 1A, 1B, 4A).
The tick's locomotor system consists of four pairs of legs in the adult and nymph stasis, and only three
pairs of legs in larvae. Each walking leg is composed of six podomeres: coxa, trochanter, femur, patella,
tibia, and tarsus, arranged proximally to distally on the podosoma (Voigt and Gorb, 2017). The legs of
ticks are highly flexible, owing to elastically articulated segments with bright colors in *A. variegatum* ticks
(Fig. 1). At the posterior end of the opisthosoma of male and female ticks, as well as nymphs, several
festoons are present, separated by marginal grooves.

Once the tick was firmly immobilized, the scutum/ alloscutum/ conscutum (Sac) was removed. The
Sac was removed with a no. 11 sterile blade. A sterile blade was used to incise the dorsal cuticle at the
distal part of the basis capitulum. A microscalpel was required to dissect ticks, particularly nymphs, to
avoid damaging internal organs. The cutting step continued around the tick's Sac, passing above the
festoons, and reaching the opposite side precisely (Fig. 4A). Special care was taken to keep the internal

cavity intact while cutting the dorsum of the tick by bypassing the spiracular plates, which are connected
to the internal organs.

Following the incision of the Sac, two dissecting forceps (Entomopraxis, Barcelona, Spain) were used
to remove it. Simply put, one forceps held the tick's body on the double-sided tape, while the other
forceps carefully lifted the Sac away from the rest of the body. Once the cuticle was lifted, all of the thin
white tubules and attached muscles were removed from the Sac to gain complete access to the tick's
internal anatomy (Fig. 2). The dorsum was either glued to the double-sided tape or cut off completely
from the tick. At this point, the small white tubes with a spider's web shape are designated as tracheae
(Tr). These structures are part of the respiratory system of ticks and facilitate gas exchanges through the
body and organs of the tick (Woolley, 1972). The tracheal system of the tick is extensive and opens
externally through two lateral spiracular plates (Sp) (Starck et al., 2018). It is noteworthy that there is
considerable variation in the structure of the respiratory system among ticks, particularly with regard to
the morphology of the spiracles. In *A. variegatum*, the spiracular plates of the female specimen are
considerably larger than those of the male specimen. This is due to the fact that females have a larger
body size and distinct physiology. These physiological mechanisms, including the digestion of large blood
meals, excretion, and egg production, require larger spiracular plates to ensure sufficient surface area for
energy exchange via gas exchange (Baker, 1997). Due to their small size, the tracheae of larvae,
compared to nymphs and adults, are significantly less developed. They have the peculiarity to facilitate
gas exchange directly across the integument (Binnington and Obenchain, 1982).
275
In ixodid ticks, the pharynx is an elongated organ situated on the ventral surface of the basis
capitulum. It facilitate the passage of ingested host blood through the food channel to the esophagus and
midgut (Sonenshine and Simo, 2021). The gut extends throughout the entire body region of the tick and
represents its largest organ. The entire tick gut includes the midgut, with a central portion, the stomach
(St), which is connected to several dark brown, spider-shaped diverticula (Div) (Fig. 2, 4B, 6B). During the
tick's engorgement phase, the midgut-diverticula is fill with host blood to be digested. Due to the
considerable size of the midgut, other internal organs may be displaced or compressed (Sonenshine and
Simo, 2021). This phenomenon is particularly salient for *A. variegatum* females, as they possess a larger
body size and consume greater quantities of blood compared to other tick species. The hindgut and the
Malpighian tubules (Mt) are key components of the tick's excretory system, playing vital roles in
maintaining homeostasis and excreting guanine-rich wastes while transporting them to the rectal sac (Rs)
(Šimo and Park, 2014). Two pairs of larger white tubes located laterally to the midgut, represent the
Malpighian tubules, which are clearly visible. In ticks, nitrogenous excreta from the Malpighian tubules
and undigested waste from the midgut are concentrated in the rectal sac and eliminated through the
anus via a short rectum (Fig. 4B) (Šimo and Park, 2014). These connections are situated beneath the
intestinal tissue. In Fig. 6D, the excretory system is faintly visible once it has been extracted from the
cavity.
293
Dissecting the respiratory system: tracheae removal
Once the attached muscles and connective tissue were carefully removed from the ventral part of the
dorsum, the entire tick cavity became visible (Fig. 2, 4B, 6B). To release the internal organs, two
dissecting forceps were gently used to grasp the tracheae at the spiracular plates between the third and
fourth pairs of legs on each side of the tick (Fig. 5).

In some instances, when dissecting feeding adult female ticks, the tracheal trunks at the base of the
spiracular plates were observed denser. If difficulties arose, the spiracular plates could be incised, and
the tracheal trunk carefully extracted with the forceps. After removal of the tracheae, the salivary glands
(Sg) and the synganglion (Sy) became more prominent (Fig. 3A).

The Gené's organ (Go), a specialized structure found only in female ticks, is located in the anterior
region of the cavity, just above the basis capituli. This structure facilitates coating the eggs with a wax-like
substance during oviposition.
306
Dissecting the digestive system: midgut removal
The midgut stomach was withdrawn from the tick body by grasping the junction of the esophagus
and stomach with one dissecting forceps and retracting the midgut with another pair of forceps. The

esophagus is a narrow, elongated tube that runs anteroventrally from the synganglion to the anterior
part of the stomach.
Subsequently, the diverticula of the midgut could be gently pulled away from the rest of the organs
without damaging them, ensuring that their digestive contents are not discharged (Fig. 2B, 3A, 8B). Since
the midgut is elastic, it can be easily removed while ensuring that it remains intact. The midgut was
rinsed in three successive pools of sterile PBS with aseptic dissecting forceps if analysis was required
(Grabowski and Kissinger, 2020). After removal of the midgut, the salivary glands and the synganglion
became more visible (Fig. 3A, 3B, 4C 6E).
Tick salivary glands are complex, multi-lobed organs that serve multiple functions during both the
tick's parasitic and non-parasitic periods on the host (Bowman and Sauer, 2004). The salivary glands are
paired structures located in the anterolateral regions of the tick's body. The salivary glands of female
ixodid ticks consist of clusters of grape-shaped alveoli, called acini, which are categorized into three types
(type I, II and III) (Šimo et al., 2017).
The salivary glands of ticks are innervated by branched peripheral nerves originating from the
synganglion (Sy), which represents the central nervous system of these arthropods. This organ performs
several essential functions in ticks, including movement control and stimulus processing (Rispe et al.,
2022). The synganglion is located in the ventro-anterior region, posterior to the midgut stomach, and is
composed of multiple ganglia (Obenchain, 1974). The synganglion is a translucent, heart-shaped
structure linked to a multitude of peripheral nerves (Pn) that innervate various parts of the tick body
(Sonenshine and Simo, 2021).
330
Dissecting the digestive and the nervous system: salivary glands and synganglion removal
Two dissection forceps were employed to collect the salivary glands. One forceps grasped the
principal salivary duct at the base of the chelicerae at the anterior upper end, while the other gently
pulled out the salivary glands. As with the midgut, the salivary glands were rinsed in three baths of sterile
PBS prior to collection and cleared of any remaining tracheae or connective tissue (Fig. 2C). The same
procedure was applied to remove the salivary gland on the opposite side. The size of the salivary glands
varies with the feeding status of the ticks, becoming larger in half-fed and fully-fed ticks. The glands could
be rinsed in three pools of PBS on either a Petri dish or a sterile glass slide, using aseptic dissecting
forceps.
Once the salivary glands have been removed, the heart became visible especially in nymphs (Fig. 4C).
As part of an open circulatory system, the pumping action of the heart and the arterial vessels facilitates
circulation within the hemocoel (Binnington and Obenchain, 1982). This organ appears pentagonal during
systole and more spherical during diastole (Cossio-Bayugar et al., 2023). The heart (H) lies in the dorsal
portion along the mid-sagittal axis and is connected anteriorly with the aorta, and leads to the
synganglion (Fig. 4C). Given the fragility of the synganglion, its removal requires precision. One forceps
can be used to grasp the whitish neuronal extensions located on its lateral side.
347
Dissecting the reproductive organs: testes and ovaries removal
Following the removal of the synganglion, the tick's reproductive system, particularly in females,
becomes more apparent (Fig. 5). It is important to note that the reproductive system only develops at the
adult stasis. Consequently, the nymph lacks both reproductive organs and a gonopore, which are
essential for the reproduction (Fig. 4C, 4D).
The male reproductive system includes the genital opening (gonopore), testes (T), vasa deferentia
(Vd), ejaculatory duct, seminal vesicle (Sv), and accessory gland complex (Arg). The male gonopore is
located in the ventroanterior region of the body, ventral to the synganglion, and opposite the anal
aperture (Sampieri et al., 2016). Mated male ticks are capable of excreting a sac-like spermatophore that
protrudes from their genital opening and delivering it with their mouthparts to the female's gonopore
(Fig. 10D)(Kiszewski et al., 2001).
The male posterior region is composed of a pair of tubular, elongated testes, positioned
dorsolaterally within the opisthosoma. These are connected to the accessory reproductive glands by the
vasa deferentia (Fig 6E, 6F)(Anholeto et al., 2015; Sampieri et al., 2014). The testes serve as the primary
site of spermatogenesis, which begins during the nymphal stage. The appearance of tick testes of ticks

may vary slightly even within a single species, depending on the nutritional and reproductive status of the
individual (Oliver and Brinton, 1972).

In metastriate ticks, the testes are linked by an extremely thin strand of filamentous tissue and taper
into the vasa deferentia. The testes are positioned along both sides of the male body, dorsally to the
synganglion and laterally to the rectal sac (Oliver, 1982). A comparison of *R. microplus* males and *A.*
variegatum males reveals significant differences in the morphology of their testes. *A. variegatum* males
possess testes that are thinner and less pronounced in the opisthosoma. The vasa deferentia represents
the narrowing of the paired testes at their anterior ends, joining in the anterior region to form a single
duct called the vas deferens or seminal vesicle.

As with the testes, the shape and the size of the vasa deferentia depend on the reproductive state of
the tick and can vary from narrow, coiled tubes to enlarged, straight structures (Oliver, 1982).
Histologically, the vasa deferentia are indistinguishable from the common vas deferens. The seminal
vesicle serves as the point of connection between the vasa derentia and the ejaculatory duct.

The ejaculatory duct is a short, narrow tube located in the ventroanterior region, extending below the
synganglion to the external gonopore (El Shoura, 1987). The male accessory reproductive glands (Arg)
play critical roles in all stages of the reproductive biology of the mated female, from the sperm
deposition to oviposition (Gillott, 2003). For instance, male accessory glands stimulate female
engorgement and activate spermatozoa upon arrival at the female seminal receptacle (Reuben Kaufman,
2007). This multi-lobed complex is located ventrally to the opisthosoma and connects to the ejaculatory
duct on its ventral face and to the vasa deferentia on the dorso-lateral region (Sampieri et al., 2014).

Dissection of males: testes

Since the internal anatomy of *A. variegatum* males is simpler than that of females, isolating the
organs is relatively easier (Fig. 2A, 6B). After removing the trachea and the midgut, the reproductive
system becomes exposed, particularly the testes, which occupy significant space (Fig. 6C, 6E). Using
dissecting tweezers, the ejaculatory duct can be gently pulled out of the tick's body, bringing the entire
reproductive system with it (Fig. 6F, 6G).

During mating, metastriate males insert only their chelicerae into the female's genital pore, leaving
the hypostome and palps outside, barely in contact with the female's surface (Fig. 10A) (Kiszewski et al.,
2001). This mechanism facilitates the transfer of the sac-like spermatophore that protrudes from the
male genital opening (Fig. 10C, 10D). Similar to males, the female genital opening is located ventrally
relative to the synganglion (Fig. 10A).

Dissecting females: Gené's organ and ovaries

As ticks are known to have a complex life cycle in the wild, some ixodid ticks exhibit high reproductive
potential, with an oviposition capacity of up to 20,000 eggs. This strategy is believed to ensure the
survival of their offspring in the wild. The female genital system comprises the gonopore, similar to that
in male, ovaries (O), a common oviduct or uterus, a bipartite vagina divided into a cervical and vestibular
vagina, and a seminal receptacle (Sr). Additionally, females possess two accessory glands: tubular and
lobular.

The ixodid tick ovary is an elongated, U-shaped tube situated in the posterior region of the body,
above the rectal sac (Fig. 3C, 5). The tick ovary is a hollow organ filled with oogonia and primary oocytes
at various early stages of development. A fold or longitudinal groove extends along the antero-dorsal
surface of the ovary, containing primary oocytes internally at their earliest stage of development, and
more advanced oocytes externally (Fig 9C) (Diehl et al., 1982). During feeding, the female's oocytes tend
to protrude into the hemocel, giving the ovary a grape-like shape (Fig. 8A).

At each end of the ovary are whitish, paired, and folded oviducts (Ovi), which converge to form a
common oviduct or uterus (Fig. 3A, 9C). The oviduct tends to stretch considerably during the passage of
the eggs due to peristaltic contractions of the oviduct walls (Fig. 7D). The uterus connects to the cervical
vagina via a short tube.

The vagina consists of a posterior, broad, and short cervical region, as well as an anterior, narrow, and
long vestibular region (El Shoura, 1989). In addition to the uterus, the seminal receptacle, tubular
accessory glands and the vestibular vagina converge in the cervical vagina (Fig. 9B) (Vega et al., 2012).
The vestibular vagina extends from the gonopore to the cervical vagina and opens externally to the

gonopore via the vulva. In the female *A. variegatum*, the vagina situated in the podosoma is more
complex to observe.
The seminal receptacle is a sac-like extension of the cervical region that serves as the storage site for
spermatozoa after mating, facilitating the fertilization of eggs (Oliver, 1989). Final stages of
spermiogenesis occur in the seminal receptacle of copulated females until full engorgement (Fig. 10C)
(Brinton et al., 1974). This structure is positioned above the uterus and opens postero-dorsally into the
cervical vagina (Diehl et al., 1982).
The Gené's organ is unique to female ticks. It secretes a waxy substance that covers the oviposited
eggs, protecting them from the environment. A peculiarity of this organ is its ability to evert itself from
the camerostomal aperture, located between the posterior base of the capitulum and the anterior region
of the scutum. Internally, this organ is situated in the anterior body region, below the capitulum, and
consists of branched tubular glands that are highly developed in the engorged females (Fig. 9A).
The tubular (Tag) and lobular accessory glands are associated with the Gené's organ and also
contribute to the egg waterproofing during the oviposition period of the female. The tubular glands form
the bulk of the Gené's organ and are located anteriorly within the hemocoel, in continuity with the horns
(Fig. 9B). The paired tubular accessory glands open into the vagina at the dorsal boundaries of the
cervical and the vestibular regions (Booth, 1985). During oviposition, the lumen of each gland fills with a
secretion that coats the egg surface as it passes through the vagina (Diehl et al., 1982).
The lobular accessory gland is a distinctive feature of ixodid ticks. This gland results from the
maturation of the vestibular vaginal epithelium as the entire female reproductive system develops. Two
porose areas are located dorsally on the female basis capituli, below the rostrum (Fig. 1A). During
oviposition, these areas are covered by the Gené's organ. The porose areas secrete a substance that is
incorporated into the waterproofed wax of the organ of Gené as it everts from the camerostomal fold.
440
The tracheal system of *A. variegatum* females is much denser than that of the males, making its
removal more complex (Fig. 2A, 6B). During dissection, with the ventral surface of the tick facing
downward, the Gené's organ is located dorsally, above to the synganglion, while the female reproductive
apparatus is positioned ventrally to the synganglion. For better access to Gené's organ, the capitulum can
be removed with a surgical blade or scissors (Fig. 9A).
An incision should be made horizontally beneath the basis capitulum, just above the camerostome, to
avoid damaging the organ. Once the connective tissue and residual tracheae are cleared from the tick's
internal cavity, the Gené's organ becomes visible. Using dissecting forceps, the Gené's organ can be
extracted from the camerostomal opening by firmly grasping it. It should be noted that the tubular
accessory glands of the Gené's organ do not fully develop until the female has completed a blood meal
(Fig. 9A, 9B). Following the removal of the trachea and the midgut, the oviducts leading to the ovary in
the posterior region, and the vagina, located ventrally to the synganglion, should become visible.
The vagina was then grasped with dissecting forceps, and the entire female reproductive apparatus
was carefully removed from the internal cavity (Fig. 9C). This apparatus includes the bipartite vagina,
seminal receptacle, uterus, accessory glands, oviducts, and ovary. It should be noted that these
components have not been differentiated.

457   **Conclusion**

Collecting tick organs is crucial for understanding vector-pathogen interactions *in vivo*. Here, we
describe *A. variegatum* anatomy and how to collect several organs of interest, especially the tick midgut,
salivary glands, and ovaries, which play crucial role in tick vector competence. Dissection methods are a
reliable approach that requires sufficient dissecting skills to enable the efficient observation and
harvesting of organs without tissue degradation and subsequent contamination. Given the intricate
internal structure of ticks, special care must be taken when dissecting tissues, especially if they are
intended for subsequent analysis. The collection of hemolymph, the fluid that fills the tick's hemocoel,
may be particularly relevant because of its hemocyte composition and interest of these cells. These
components are of fundamental importance to understanding how bacteria can adapt within the tick
compartment. As demonstrated in the case of *A. americanum*, the complete depletion of granulocytic

hemocytes resulted in the death of the tick following an *Ehrlichia chaffeensis* infection (Adegoke et al.,
2023).
Further studies will then be conducted on these specific tissues to elucidate the molecular and
cellular processes implemented by the tick in response to the pathogen, at both biological and
transcriptomic levels. Performing sex-specific dissections may facilitate more precise differentiation
among ticks, particularly in the case of engorged females. Given the established role of male salivary
glands in the reproductive process, involving the secretion and subsequent lubrification of the
spermatophore, it can be hypothesized that female and male ticks may employ divergent strategies in
their developmental processes (Tan et al., 2015). Prior research has indicated that tick sex exert a
considerable influence on their microbiome. In ecology, the bacterial richness and species diversity
present within a single sample are collectively referred to as "alpha diversity" (Andermann et al., 2022).
As demonstrated in the case of *Dermacentor variabilis* male ticks, which exhibit higher alpha diversity
measures than females. This phenomenon may be attributed to the reduced quantity of blood ingested
by males during feeding and their capacity to feed multiple times on a host, which establish a more
diverse bacterial community (Duncan et al., 2022). *Amblyomma variegatum*, the tropical bont tick, has
received comparatively less attention in scientific studies than the cattle tick, *Rhipicephalus microplus*.
This study represents a significant contribution to the field, as it provides the first microscopic
observations and a general description of the internal anatomy of *A. variegatum*.
486
487

488   **Figure 1:** Anatomical description of the dorsal surface of unfed adult *A. variegatum* ticks mounted on
489   double-sided adhesive tape. **(A)** Dorsal view of unfed *A. variegatum* female (0.78x). **(B)** Dorsal view of
490   unfed *A. variegatum* male, with its first leg cut to allow hemolymph collection (0.78x).

Figure 2: Dorsal view of an unfed *A. variegatum* female and its organs. **(A)** Scutum dissection: dorsal
view of an unfed *A. variegatum* female with the scutum removed and immersed in PBS (1.6x). **(B)** Midgut
dissection: close up view of the gut of an unfed *A. variegatum* female (1.25x). **(C)** Salivary glands
dissection: close-up view of the salivary glands of an unfed *A. variegatum* female (3.2x). Ac: acini ; Div:
diverticulum ; Sac: <u>scutum</u>/ alloscutum/conscutum (in the female tick) ; Sd: salivary duct ; Sg: salivary
glands ; St: stomach ; Tr: tracheae.
497
Figure 3: Dorsal view of an unfed *A. variegatum* female with the midgut removed and organs. **(A)**
Midgut dissection: dorsal view of unfed an *A. variegatum* female with scutum removed and immersed in
PBS (1.6x). **(B)** Synganglion dissection: close-up view of the nervous system of unfed an *A. variegatum*
female (5x). **(C)** Reproductive system dissection: close-up of the ovary of an unfed *A. variegatum* female
(5x). O: ovary ; Ovi: oviducts ; Pn: peripheral nerves ; Sac: <u>scutum</u>/ alloscutum/conscutum (here, in the
female tick) ; Sy: synganglion ; Sg: salivary glands ; Tr: tracheae.
504
Figure 4: Dorsal view of an unfed *A. variegatum* nymph. **(A)** Dissection of the scutum: dorsal view of
an unfed *A. variegatum* nymph immersed in PBS, showing a clean incision (yellow dotted line) around the
idiosoma (2.5x). **(B)** Removal of the scutum: dorsal view of an unfed *A. variegatum* nymph immersed in
PBS, with the conscutum lifted using forceps (2.5x). **(C)** Dissection of the midgut: *A. variegatum* nymph
with the entire gut removed and the conscutum cut off (3.2x). **(D)** Unfed *A. variegatum* nymph with the
salivary glands, midgut, tracheae and Malpighian tubules removed (4x). Div: diverticulum ; H: heart ; Mt:
Malpighian tubules ; Rs: rectal sac ; Sac: <u>scutum/ alloscutum</u>/conscutum (here, in the nymph tick) ; Sy:
synganglion ; pink circle : hematin granules released after gut tear ; black triangle: anal aperture.
513
514   **Figure 5:** Dorsal view of an unfed *A. variegatum* female with the midgut, and salivary glands removed
515   (2x). Div: diverticulum ; O: ovary ; Ovi: oviducts ; Rs: rectal sac ; Sp: spiracular plates ; Tr: tracheae.
516
517   **Figure 6:** Dorsal view of a fed and mated *A. variegatum* male. **(A)** Dorsal view of a fully repleted *A.*
518   *variegatum* male with a rounded conscutum (1.25x). **(B)** Scutum dissection: dorsal view of a fed *A.*
519   *variegatum* male with folded conscutum, immersed in PBS (1.6x). **(C)** Midgut dissection: *A. variegatum*

male with the entire gut removed (1,6x). **(D)** Close up of the excretory system (2.5x). **(E)** Salivary glands dissection: *A. variegatum* male with the paired salivary glands removed (2.5x). **(F, G)** Reproductive organs dissection: highlighting the organs of the male reproductive system (1.6x, 2.5x). Arg: accessory reproductive glands ; Ct: connective tissue ; Div: diverticulum ; Hin: hindgut ; Mro: male reproductive organs excluding testes ; Mt: Malpighian tubules ; Rs: rectal sac ; Sac: scutum/ alloscutum/<u>conscutum</u> (in the male tick) ; Sg: salivary glands ; St: stomach ; Sv: seminal vesicle ; Sy: synganglion ; T: testes ; Tr: tracheae ; Vd: vasa deferentia.

Figure 7. View of a fully repleted *A. variegatum* female. **(A)** Dorsal view of a fully engorged *A. variegatum* female (0.78x). **(B)** Dorsal view of an ovipositing *A. variegatum* female with scutum removed (0.78x). **(C)** Dorsal view of *A. variegatum* with thousands of eggs expelled from the oviducts (0.78x). **(D)** Close up of oviducts full of eggs (0.78x). Cp: capitulum ; Div: diverticulum ; E: eggs ; Ooc: oocytes ; O: ovary ; Tr: tracheae.

Figure 8: Dorsal view of a partially fed *A. variegatum* female with the midgut removed and organs. **(A)** Dorsal view of a semi-engorged female *A. variegatum* female with the midgut removed (1x). **(B)** Close-up of the midgut of a partially fed female *A. variegatum* female (0.78x). Mt: Malpighian tubules ; O: ovary ; Sg: salivary glands ; Sp: spiracular plates.

Figure 9: Dorsal view of a partially fed *A. variegatum* female with the midgut, and salivary glands removed, showing reproductive organs. **(A)** Dorsal view of a female *A. variegatum* with a close-up of reproductive system (2x). **(B)** Close up of reproductive organs with accessory glands (2x). **(C)** Close-up of ovary with oocytes (1,6x). Ch: chelicerae ; Fro: female reproductive organs excluding ovary and oviducts ; Go: Gené's organ ; Mt: Malpighian tubules ; Ooc : oocytes ; Sy: synganglion ; Tg: tubular glands ; Tr: tracheae.

Figure 10. View of a mated *A. variegatum* female. **(A)** Ventral view of a semi-engorged *A. variegatum* female with an ectospermatophore attached near to its gonopore shortly after copulation (0.78x). **(B)** Dorsal view of a partially fed *A. variegatum* female with a round alloscutum (0.78x). **(C)** Dorsal view of an endospermatophore contained in the seminal receptacle of a semi-engorged *A. variegatum* female (2.5x). **(D)** Close-up of the endospermatophore dissected a few minutes after the female mated with a male (2.5x). Cc: cement cone ; O: ovary ; Ooc: oocytes ; Rs: rectal sac ; Sac: <u>scutum/ alloscutum/</u>conscutum (in the female tick) ; Sr: seminal receptacle ; Sp: spiracular plates ; Tr: tracheae ; black triangle: anal pore ; pink triangle: gonopore ; red triangle: ectospermatophore.

Acknowledgements

The authors want to thank Mélanie Dhune for providing ticks from the CIRAD animal facility as well as Loïc Jacquet-Crétides and Jimmy Dédy for their support with animal care. Naomie Pature is grateful to Rosalie Aprelon for technical assistance with dissection.

Conflict of interest disclosure

The authors declare that they have no financial conflicts of interest in relation to the content of the article.

Funding

This work is supported by the United States Department of Agriculture grant 58-3022-1-018-F (Risk of Arthropod-borne diseases in the Caribbean).

564   **References**

Adegoke, A., Hanson, J., Smith, R.C., Karim, S., 2023. Ehrlichia chaffeensis co-opts phagocytic hemocytes
for systemic dissemination in the Lone Star tick, Amblyomma americanum. J Innate Immun.
https://doi.org/10.1159/000535986

Andermann, T., Antonelli, A., Barrett, R.L., Silvestro, D., 2022. Estimating Alpha, Beta, and Gamma
Diversity   Through   Deep   Learning.   Front   Plant   Sci   13,   839407.
https://doi.org/10.3389/fpls.2022.839407

Anholeto, L.A., Nunes, P.H., Remédio, R.N., Camargo-Mathias, M.I., 2015. Testes of fed and unfed
mblyomma cajennense ticks (Acari: Ixodidae). First morphological data. Acta Zoologica 96, 375–382.
https://doi.org/10.1111/azo.12083

Baker, G.T., 1997. Spiracular Plate of Nymphal and Adult Hard Ticks (Acarina: Ixodidae): Morphology and
Cuticular Ultrastructure. Invertebrate Biology 116, 341. https://doi.org/10.2307/3226866

Barré, N., 1989. Biologie et écologie de la tique Amblyomma variegatum (Acarina : Ixodina) en
Guadeloupe (Antilles Françaises) (thesis). CIRAD-IEMVT.

Bezuidenhout, J.D., 1987. NATURAL TRANSMISSION OF HEARTWATER. Onderstepoort J. vet Res. 54, 349–
351.

Binnington, K.C., Obenchain, F.D., 1982. Structure and Function of the Circulatory, Nervous, and
Neuroendocrine   Systems   of   Ticks,   in: Physiology   of   Ticks.   Elsevier,   pp.   351–398.
https://doi.org/10.1016/B978-0-08-024937-7.50015-9

Booth, T.F., 1985. The structure and function of Gene's organ and its associated glands in ticks (phd).
Thames Polytechnic.

Bowman, A.S., Sauer, J.R., 2004. Tick salivary glands: function, physiology and future. Parasitology 129
Suppl, S67-81. https://doi.org/10.1017/s0031182004006468

Brinton, L.P., Burgdorfer, W., Oliver, J.H., 1974. Histology and fine structure of spermatozoa and egg
passage in the female tract of Dermacentor andersoni stiles (Acari-Ixodidae). Tissue and Cell 6, 109–
125. https://doi.org/10.1016/0040-8166(74)90026-3

Camargo-Mathias, M.I. (Ed.), 2018. Inside ticks: morphophysiology, toxicology and therapeutic
perspectives. Editora UNESP. https://doi.org/10.7476/9788595462861

Caperucci, D., Mathias, M.I.C., Bechara, G.H., 2009. Histopathology and Ultrastructure Features of the
Midgut of Adult Females of the Tick Amblyomma cajennense Fabricius, 1787 (Acari: Ixodidae) in
Various Feeding Stages and Submitted to Three Infestations. Ultrastructural Pathology 33, 249–259.
https://doi.org/10.3109/01913120903296945

Cossio-Bayugar, R., Miranda-Miranda, E., Kumar, S., 2023. A Laboratory Manual on Rhipicephalus
microplus, Cambridge Scholars Publishing. Lady Stephenson Library, Newcastle upon Tyne, NE6 2PA,
UK.

Cupp, E.W., 1991. Biology of Ticks. Veterinary Clinics of North America: Small Animal Practice 21, 1–26.
https://doi.org/10.1016/S0195-5616(91)50001-2

de la Vega, R., Diaz, G., Galán, M., Fernández, C., 2012. Anatomy and hystology of the female
reproductive system of Boophilus microplus (Acari: Ixodidae). Rev. Salud Anim. 34, 1–10.

Deans, A.R., Mikó, I., Wipfler, B., Friedrich, F., 2012. Evolutionary phenomics and the emerging
enlightenment of arthropod systematics. Invert. Systematics 26, 323.
https://doi.org/10.1071/IS12063

Diehl, P.A., Aeschlimann, A., Obenchain, F.D., 1982. CHAPTER 9 - Tick Reproduction: Oogenesis and
Oviposition, in: Obenchain, FREDERICK D., Galun, R. (Eds.), Physiology of Ticks. Pergamon, pp. 277–
350. https://doi.org/10.1016/B978-0-08-024937-7.50014-7

Duncan, K.T., Elshahed, M.S., Sundstrom, K.D., Little, S.E., Youssef, N.H., 2022. Influence of tick sex and
geographic region on the microbiome of *Dermacentor variabilis* collected from dogs and cats across
the United States. Ticks and Tick-borne Diseases 13, 102002.
https://doi.org/10.1016/j.ttbdis.2022.102002

Edwards, K.T., 2009. Examination of the Internal Morphology of the Ixodid Tick, Amblyomma maculatum
Koch, (Acari: Ixodidae); a "How-to" Pictorial Dissection Guide.

El Shoura, S.M., 1989. Effect of blood meal and mating on the genital tract ultrastructure in the female
camel tick Hyalomma (Hyalomma) dromedarii (Ixodoidea: Ixodidae). Exp Appl Acarol 6, 157–175.
https://doi.org/10.1007/BF01201645

El Shoura, S.M., 1987. Fine Structure of the Vasa Deferentia, Seminal Vesicle, Ejaculatory Duct, And
Accessory Gland of Male Ornithodoros (Pavlovskyella) Erraticus (Acari: Ixodoidea: Argasidae). Journal
of Medical Entomology 24, 235–242. https://doi.org/10.1093/jmedent/24.2.235

Estrada-Pena, A., Ayllon, N., De La Fuente, J., 2012. Impact of Climate Trends on Tick-Borne Pathogen
Transmission. Frontiers in Physiology 3.

Friedrich, F., Matsumura, Y., Pohl, H., Bai, M., Hörnschemeyer, T., Beutel, R.G., 2014. Insect morphology
in the age of phylogenomics: innovative techniques and its future role in systematics. Entomological
Science 17, 1–24. https://doi.org/10.1111/ens.12053

Gilbert, L., 2021. The Impacts of Climate Change on Ticks and Tick-Borne Disease Risk. Annu. Rev.
Entomol. 66, 373–388. https://doi.org/10.1146/annurev-ento-052720-094533

Gillott, C., 2003. Male Accessory Gland Secretions: Modulators of Female Reproductive Physiology and
Behavior. Annu. Rev. Entomol. 48, 163–184. https://doi.org/10.1146/annurev.ento.48.091801.112657

Grabowski, J.M., Kissinger, R., 2020. Ixodid Tick Dissection and Tick Ex Vivo Organ Cultures for Tick-Borne
Virus Research. Curr Protoc Microbiol 59, e118. https://doi.org/10.1002/cpmc.118

Kennedy, B., Trim, S.A., Laudier, D., LaDouceur, E.E.B., Cooper, J.E., 2021. Arthropoda: Arachnida, in:
LaDouceur, E.E.B. (Ed.), Invertebrate Histology. Wiley, pp. 221–246.
https://doi.org/10.1002/9781119507697.ch8

Kiszewski, A.E., Matuschka, F.-R., Spielman, A., 2001. M ATING S TRATEGIES AND S PERMIOGENESIS IN I
XODID T ICKS. Annu. Rev. Entomol. 46, 167–182. https://doi.org/10.1146/annurev.ento.46.1.167

Lee, Junsoo, Ryu, J., Han, S., Ravichandran, N.K., Seong, D., Lee, Jaeyul, Wijesinghe, R.E., Kim, P., Lee, S.-
Y., Jung, H.-Y., Jeon, M., Choi, K.S., Kim, J., 2021. Identification of organs inside hard tick body using
spectral-domain optical coherence tomography. Infrared Physics & Technology 114, 103611.
https://doi.org/10.1016/j.infrared.2020.103611

Mathias, M.I.C., 2013. Guia básico de morfologia interna de carrapatos ixodídeos. Editora UNESP.

Obenchain, F.D., 1974. Neurosecretory system of the American dog tick, Dermacentor variabilis (Acari:
Ixodidae).   I.   Diversity   of   cell   types.   J   Morphol   142,   433–445.
https://doi.org/10.1002/jmor.1051420406

Oliver, J.H., 1989. Biology and Systematics of Ticks (Acari:Ixodida). Annual Review of Ecology and
Systematics 20, 397–430.

Oliver, J.H., 1982. Tick Reproduction: Sperm Development and Cytogenetics, in: Physiology of Ticks.
Elsevier, pp. 245–275. https://doi.org/10.1016/B978-0-08-024937-7.50013-5

Oliver, J.H., Brinton, L.P., 1972. Cytogenetics of ticks (Acari: Ixodoidea). 7. Spermatogenesis in the Pacific
Coast tick, dermacentor occidentalis Marx (Ixodidae). J Parasitol 58, 365–379.

Pegram, R., Indar, L., Eddi, C., George, J., 2004. The Caribbean Amblyomma Program: Some Ecologic
Factors Affecting Its Success. Annals of the New York Academy of Sciences 1026, 302–311.
https://doi.org/10.1196/annals.1307.056

Prine, K.C., Hodges, A.C., 2012. Tropical Bont Tick Amblyomma variegatum Fabricius (Arachnida: Acari:
Ixodidae): EENY-518/IN934, 6/2012. EDIS 2012. https://doi.org/10.32473/edis-in934-2012

Reuben Kaufman, W., 2007. Gluttony and sex in female ixodid ticks: How do they compare to other
blood-sucking   arthropods?   Journal   of   Insect   Physiology   53,   264–273.
https://doi.org/10.1016/j.jinsphys.2006.10.004

Rispe, C., Hervet, C., de la Cotte, N., Daveu, R., Labadie, K., Noel, B., Aury, J.-M., Thany, S., Taillebois, E.,
Cartereau, A., Le Mauff, A., Charvet, C.L., Auger, C., Courtot, E., Neveu, C., Plantard, O., 2022.
Transcriptome of the synganglion in the tick Ixodes ricinus and evolution of the cys-loop ligand-gated
ion channel family in ticks. BMC Genomics 23, 463. https://doi.org/10.1186/s12864-022-08669-4

Sampieri, B.R., Labruna, M.B., Bueno, O.C., Camargo-Mathias, M.I., 2014. Dynamics of cell and tissue
genesis in the male reproductive system of ticks (Acari: Ixodidae) Amblyomma cajennese (Fabricius,
1787) and Amblyomma aureolatum (Pallas, 1772): a comparative analysis. Parasitol Res 113, 1511–
1519. https://doi.org/10.1007/s00436-014-3795-y

Sampieri, B.R., Moreira, J.C.S., Páez, F.A.R., Camargo-Mathias, M.I., 2016. Comparative morphology of the
reproductive system and germ cells of Amblyomma ticks (Acari: Ixodidae): A contribution to Ixodidae
systematics. J Microsc Ultrastruct 4, 95–107. https://doi.org/10.1016/j.jmau.2015.11.003

Sasso Porto, D., Melo, G.A.R., Almeida, E.A.B., 2016. Clearing and dissecting insects for internal skeletal
morphological research with particular reference to bees. Revista Brasileira de Entomologia 60, 109–
113. https://doi.org/10.1016/j.rbe.2015.11.007

Šimo, L., Kazimirova, M., Richardson, J., Bonnet, S.I., 2017. The Essential Role of Tick Salivary Glands and
Saliva in Tick Feeding and Pathogen Transmission. Frontiers in Cellular and Infection Microbiology 7.

Šimo, L., Park, Y., 2014. Neuropeptidergic control of the hindgut in the black-legged tick Ixodes scapularis.
Int J Parasitol 44, 819–826. https://doi.org/10.1016/j.ijpara.2014.06.007

Sonenshine, D.E., Simo, L., 2021. Biology and Molecular Biology of Ixodes scapularis, in: Lyme Disease and
Relapsing Fever Spirochetes: Genomics, Molecular Biology, Host Interactions and Disease
Pathogenesis. Caister Academic Press. https://doi.org/10.21775/9781913652616.12

Starck, J.M., Mehnert, L., Biging, A., Bjarsch, J., Franz-Guess, S., Kleeberger, D., Hörnig, M., 2018.
Morphological   responses   to   feeding   in   ticks   (Ixodes   ricinus).   Zoological   Letters   4,   20.
https://doi.org/10.1186/s40851-018-0104-0

Tan, A.W.L., Francischetti, I.M.B., Slovak, M., Manjunatha, K.R., Ribeiro, J.M.C., 2015. Sexual differences
in the sialomes of the zebra tick, Rhipicephalus pulchellus. J Proteomics 117, 120–144.
https://doi.org/10.1016/j.jprot.2014.12.014

Tidwell, J.P., Treviño, D.E., Thomas, D.B., Mitchell, R.D., Heerman, M.C., Pérez de León, A., Lohmeyer,
K.H., 2021. Pictorial dissection guide and internal anatomy of the cattle tick, Rhipicephalus (Boophilus)
microplus (Canestrini). Ticks Tick Borne Dis 12, 101685. https://doi.org/10.1016/j.ttbdis.2021.101685

Uilenberg, G., 1983. Heartwater (Cowdria ruminantium infection): current status. Adv Vet Sci Comp Med
27, 427–480.

Vega, R., Diaz, G., Galan, M., Fernández, C., 2012. ANATOMY AND HISTOLOGY OF THE FEMALE
REPRODUCTIVE SYSTEM OF Boophilus microplus (ACARI: IXODIDAE).

Voigt, D., Gorb, S., 2017. Functional morphology of tarsal adhesive pads and attachment ability in ticks
Ixodes ricinus (Arachnida, Acari, Ixodidae). Journal of Experimental Biology 220, 1984–1996.
https://doi.org/10.1242/jeb.152942

Woolley, T.A., 1972. Scanning Electron Microscopy of the Respiratory Apparatus of Ticks. Transactions of
the American Microscopical Society 91, 348. https://doi.org/10.2307/3224879

Zacaï, A., Vannier, J., Lerosey-Aubril, R., 2016. Reconstructing the diet of a 505-million-year-old
arthropod: Sidneyia inexpectans from the Burgess Shale fauna. Arthropod Structure & Development,
Special Issue: Fossils as Living Beings 45, 200–220. https://doi.org/10.1016/j.asd.2015.09.003
701

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure 9.

Figure 10.