Dissection and internal anatomy of the giant tropical bont tick *Amblyomma variegatum*

Naomie Pature[1,2,3], Nonito Pagès[1,2], Valérie Rodrigues[1,2] and Damien F. Meyer*[1,2]

[1] CIRAD, UMR ASTRE, F-1970 Petit-Bourg – Guadeloupe, France

[2] ASTRE, CIRAD, INRAE, Univ Montpellier – Montpellier, France

[3] Université des Antilles, École doctorale n° 636 DEECA, Campus de Fouillole, Pointe-à-Pitre – Guadeloupe, France

*Corresponding author

Correspondence: damien.meyer@cirad.fr

ABSTRACT

Ticks are parasites arthropods that feed on the blood of animals and humans, serving as vectors responsible for numerous diseases worldwide. The tropical bont tick, *Amblyomma variegatum*, is a primary vector of *Ehrlichia ruminantium*, the causative agent of heartwater disease, endemic in sub-Saharan Africa and several Caribbean islands. Due to their economic and public health impacts, *A. variegatum* and heartwater have been classified as high-consequence foreign animal diseases and pests by the United States Department of Agriculture's Animal and Plant Health Inspection Service. Understanding the ecology and biology of *A. variegatum* is thus crucial to developing effective control methods to limit its geographic spread. A key initial step in this endeavor involves the precise isolation and detailed functional characterization of tick organs to better understand vector competence. However, complete anatomical descriptions of internal structures within the *Amblyomma* genus remain scarce. In this study, we present a thorough characterization of the internal anatomy of unfed nymphs, partially engorged males, and unfed and partially engorged females of the tick species *A. variegatum*. By accurately identifying and mapping organs of interest, we highlight notable morphological differences from other tick species.

31   This foundational work not only enhances our understanding of *A. variegatum* anatomy but
32   also serves as a crucial resource for investigating tick-borne disease transmission.
33   Ultimately, these findings support the development of targeted vector control strategies,
34   applicable to both endemic and non-endemic regions.

36   ***Keywords:*** *Amblyomma variegatum*, heartwater, dissection, tick organs, anatomy, hard-bodied ticks

37
38

Introduction

The earliest known study of insects was undertaken by the Greek naturalist and philosopher Aristotle, who emphasized their limited significance (Friedrich et al., 2014). For centuries, the internal anatomy of arthropods remained a subject of extensive study (Kennedy et al., 2021). This field advanced markedly with Galileo's early 17[th] century invention of the telescope, which was adapted for observing insects. Building on these developments, Antonie van Leeuwenhoek, often regarded as the father of microscopy, conducted pioneering studies, including the dissection of spiders (Cocquyt, 2022).

Microscopic dissection, a fundamental technique in arthropod research, allows detailed exploration of internal structures (Deans et al., 2012). A noteworthy example comes from Zacaï *et al.*, who employed microscopy to study the internal anatomy of the fossilized arthropod *Sidneyia inexpectans*, shedding light on its digestive system and drawing comparisons to modern chelicerates (Zacaï et al., 2016). Over time, various dissection techniques have been developed, facilitating the study of a wide range of arthropods species. Dissecting larger arthropods, such as hard ticks, offers a straightforward approach to examining their internal anatomy. Tools such as fine tweezers and sharp blades, alongside aids like strong tape or wax, enhance the precision and efficiency of these procedures.

Despite advances in imaging techniques - including X-ray, microtomography (μCT), and confocal laser scanning microscopy (CLSM) - light-optical microscopy remains the gold standard for analyzing morphological diversity (Sasso Porto et al., 2016). However, the fragility of arthropod internal structures makes microscopic dissection inherently risky, as tissue damage can compromise the integrity of specimens. Consequently, the manipulator's skill in performing dissections is crucial to preserve the organs, particularly for morphological studies. To address this limitation, a recent study has focused on the development of spectral-domain optical coherence tomography (SD-OCT), which enables non-invasive imaging while preserving the anatomical features of tick specimens (Lee et al., 2021). Nonetheless, this approach does not allow for the collection of arthropod internal organs of interest for further studies of tick-pathogen interactions (Deans et al., 2012).

The tropical bont tick, *Amblyomma variegatum* (Fabricius, 1794), is an obligate hematophagous arthropod belonging to the phylum Arthropoda, subphylum Chelicerata, Class Arachnida, subclass Acari, and order Ixodida (Camargo-Mathias, 2018). This tropical tick is also referred to as the Senegalese tick in the Guadeloupe archipelago and the Antigua gold tick in Antigua (Pegram et al., 2004). *A. variegatum* belongs to the family Ixodidae, characterized by a hard body with a dorsal shield (scutum) that may be ornamented in the case of males, or may not be ornamented for females.

A. variegatum acts as the primary vector for heartwater, a major and often fatal disease of ruminants in the Caribbean region and Subharian Africa caused by the bacterium *Ehrlichia ruminantium*. This tick is frequently associated with dermatophilosis (Prine and Hodges, 2012). Its presence also contributes to lameness and the formation of deep abscesses in small ruminants, often leading to immunosuppression and metabolic disorders (Barré, 1989). These impacts result in considerable economic and health burdens. While more than ten species of *Amblyomma* ticks transmit *E. ruminantium* in Africa, *A. variegatum* and *A. hebraeum* are the main vectors (Bezuidenhout, 1987). The geographic overlap of heartwater-endemic regions with *Amblyomma* distribution underscores the importance of studying these ticks. Factors such as their abundance, activity, and environmental adaptability directly influence their vector potential (Uilenberg, 1983).

Beyond endemic areas, *A. variegatum*, also poses a potential threat to non-endemic regions due to livestock movement and climate change (Estrada-Pena et al., 2012). Alteration in host populations and environmental conditions may indirectly affect the abundance and distribution of ticks. (Gilbert, 2021). Knowing the tick's anatomy is therefore critical for studying tick-pathogen interactions and developing appropriate vector control strategies to mitigate the socioeconomic and health impacts of tick-borne diseases, which disproportionately affect resource-limited regions. While research on *Rhipicephalus microplus* and *Ixodes scapularis* have provided valuable insights into tick physiology, *A. variegatum* remain relatively understudied. Detailed investigations into its reproductive efficiency, feeding behavior, and ecological adaptations are needed, as these traits contribute to its prominence as a vector.

The precise identification of tick internal organs is of significant value in entomology and parasitology, as it facilitates the development of natural and chemical effective control methods. To further develop control technologies, it is necessary to anatomize the internal structure of ticks. The study of the internal

93   anatomy of arthropods is of great interest due to the remarkable diversity observed, particularly
94   regarding the specific barriers such as target organs they possess. Given the widespread distribution of *A.*
95   *variegatum*, particularly in some Caribbean islands, there is a need to study the tick and its role as a
96   pathogen vector. The impetus for composing this article was the desire to produce a paper identifying
97   the organs of the *A. variegatum* tick and determining the optimal dissection procedures.
98
Despite the numerous dissection guides exist for identifying tick organs, especially for *Rhipicephalus*,
species, comparatively few have focused on *Amblyomma* ticks (Cossio-Bayugar et al., 2023; Tidwell et al.,
2021). High-quality histologic images of the reproductive and digestive systems of ixodid ticks are readily
available (Camargo-Mathias, 2018; de la Vega et al., 2012; Mathias, 2013), along with scanning electron
microscopy images of male and female *Amblyomma* reproductive structures (Anholeto et al., 2015;
Caperucci et al., 2009). However, beyond the seminal work by Edwards *et al.* (Edwards, 2009) ,
dissection-based studies on *Amblyomma* ticks remain scarce. Even the pictorial instructions provided in
this guide rely on two different *Amblyomma* species to provide a comprehensive view
of *Amblyomma* internal organs.
In this study, we present a detailed examination of the internal anatomy of *A. variegatum*, from
nymphs to adult ticks at different stages of engorgement. The dissected organs are described and
illustrated, providing a valuable resource for researchers. This report aims to enable consistent and
reproductible tissue sampling of *A. variegatum*, facilitating precise investigations into tick pathogen
interactions, and, <u>ultimately</u> the development of innovative tick control technologies.

                                    Material & Methods

145
Dissection tools
The materials necessary for tick dissections are detailed in the "Tick general dissection" section
below. All the dissection tools were aseptically prepared before use. Specifically, forceps were
autoclaved, then immersed in 70% ethanol for five minutes and rinsed with sterile ultrapure water. Tools
were allowed to air dry on a clean paper towel. Between each tick sample, the tools were immersed in a
25% bleach solution for several minutes, then placed in 70% ethanol, followed by immersion in ultra-pure
water. After this process, the tools were allowed to dry in air dry or were gently wiped with a sterile
paper towel (https://ehs.stanford.edu/reference/comparing-different-disinfectants).
All the petri dishes were wiped or sprayed with 70% ethanol and allowed to air dry. In certain cases,
glass petri dishes could be used to avoid the damaging of pointed forceps. A LEICA M205FA fluorescence
stereomicroscope equipped with Fluocombi (LEICA, Wetzlar, Germany) was used for image acquisition.
157
Approach
Tick rearing
For all experiments, batches of unfed *A. variegatum* ticks (larvae, nymphs and adults) were reared as
a colony for more than 13 years at the CIRAD animal facility in Guadeloupe. Larvae and nymphs were
placed on the ear side of a naïve creole goat in a Mölnlycke Tubifast green line bandage, secured with
Kamar adhesive (Vital Concept, Loudeac, France). Adults, on the other hand, were placed on the goat's
flanks. Larvae were collected and allowed to molt in a container maintained at room temperature and
70-80% of relative humidity. Subsequent to ecdysis, nymphs were used for dissections after a brief pre-
feeding period (approximately 20 days) during which they developed to a sufficient degree of maturity. In
the case of adults, after completing feeding and detachment, repleted females were collected and were
allowed to undergo their pre-oviposition period (approximately 10 days) before being used for dissection.
Semi-engorged females were collected after a period of 7 days after the onset of their feeding, following
several copulations and after approximately 12 days during reproductive physiological processes.
However, in the case of engorged and sexually mature males, they were manually removed from the goat
after approximately 28 days of feeding and were directly dissected.
173
Preparation of the tick
Prior to the dissection, the ticks were washed twice for 30 seconds in ultrapure water and once in PBS
in three 0.5 mL Eppendorf microtubes to remove impurities on the tick's cuticle. During the washing
process, the tubes containing the ticks were gently inverted by hand. As *A. variegatum* ticks are rapid
crawlers, they could optionally be cooled on ice to slow them down. Following the washing step, the ticks
were completely dried on Kimtech paper towels (Kimberly-Clark, Nanterre, France) before being
immobilized for dissection.
The fixation of the tick was performed by immobilizing a single tick, with its ventral side down, on a
piece of double-sided adhesive tape within a Petri dish (Fig. 1). To ensure fixation of the tick, all four legs
were extended and secured to the double-sided tape with tweezers. For nymphs and unfed adults,
multiple ticks could be fixed in a single Petri dish. Once immobilized, a few drops of sterile, filtered
Dulbecco's Phosphate Buffered Saline (PBS) 1X were applied to cover the entire tick body. This prevented
desiccation of the tick's tissues and facilitated dissection by swelling the internal organs.
187
Hemolymph and saliva collection for research purpose
To further study *A. variegatum* saliva components or hemolymph cell populations, two collection
methods can be used. For hemolymph collection, the same procedure was used to wash the external
structure of the tick. The ticks were then fixed on a double-sided tape in the same manner as for the
dissection. To collect the hemolymph, the first two legs of the tick were fully extended. Then, the most
distal joint of the tick's tarsus was cut with a scalpel, and the drop was collected with a micropipette
through a microcapillary tube (Fig. 1B). For larger volumes of hemolymph, several legs may be amputated
and the tick's body was gently squeezed to promote drainage.

To collect saliva, semi-engorged ticks that had been removed from animals or artificially fed were used to obtain larger amounts of saliva. After washing the tick as described above, it was immobilized on double-sided tape placed in the upper part of a petri dish, with its ventral surface facing up to expose its mouthparts, including the rostrum and palps. Another piece of double-sided tape was placed near the tick's basis capitula on the edge of the petri dish. A capillary tube was pressed between the hypostome and palps, ensuring the pedipalps remained outside the capillary tube.

A solution of pilocarpine (20mg/ml) was prepared, and 50 μL was gently injected into the tick's stigmata or spiracles (Sp) (Fig 10A) using a Hamilton syringe. Salivation was observed, and the saliva was collected using a pipette tip inserted into a micropipette. If the tick did not salivate after 10 minutes, an additional 20 μL of pilocarpine was injected. The saliva was stored at -80°C for further analysis or use.

Tick general dissection
Dissecting the external structure: Scutum/alloscutum/conscutum removal

Once the tick was firmly immobilized, the scutum/ alloscutum/ conscutum (Sac) was removed. A sterile no. 11 blade was used to remove the Sac. A sterile blade was used to incise the dorsal cuticle at the distal part of the basis capitulum. A microscalpel was required to dissect ticks, particularly nymphs, to avoid damaging internal organs. The cutting step continued around the tick's Sac, passing above the festoons, and reaching the opposite side precisely (Fig. 4A). Special care was taken to keep the internal cavity intact while cutting the dorsum of the tick by bypassing the spiracular plates, which are connected to the internal organs.

Following the incision of the Sac, two dissecting forceps (Entomopraxis, Barcelona, Spain) were used to remove it. Simply put, one forceps held the tick's body on the double-sided tape, while the other forceps carefully lifted the Sac away from the rest of the body. Once the cuticle was lifted, all of the thin white tubules and attached muscles were removed from the Sac to gain complete access to the tick's internal anatomy (Fig. 2). The dorsum was either glued to the double-sided tape or cut off completely from the tick. At this point, the small white tubes with a spider's web shape are designated as tracheae (Tr).

Dissecting the respiratory system: tracheae removal

Once the attached muscles and connective tissue were carefully removed from the ventral part of the dorsum, the entire tick cavity became visible (Fig. 2, 4B, 6B). To release the internal organs, two dissecting forceps were gently used to grasp the tracheae at the spiracular plates between the third and fourth pairs of legs on each side of the tick (Fig. 5).

In some instances, when dissecting feeding adult female ticks, the tracheal trunks at the base of the spiracular plates (Sp) were observed denser. If difficulties arose, the spiracular plates could be incised, and the tracheal trunk carefully extracted with the forceps. After removal of the tracheae, the salivary glands (Sg) and the synganglion (Sy) became more prominent (Fig. 3A).

Dissecting the digestive system: midgut removal

The midgut stomach (St) was withdrawn from the tick body by grasping the junction of the esophagus and stomach with one dissecting forceps and retracting the midgut with another pair of forceps. The esophagus is a narrow, elongated tube that runs anteroventrally from the synganglion to the anterior part of the stomach.

Subsequently, the diverticula (Div) of the midgut could be gently pulled away from the rest of the organs without damaging them, ensuring that their digestive contents are not discharged (Fig. 2B, 3A, 8B). Since the midgut is elastic, it can be easily removed while ensuring that it remains intact. The midgut was rinsed in three successive pools of sterile PBS with aseptic dissecting forceps if analysis was required (Grabowski and Kissinger, 2020). After removal of the midgut, the salivary glands (Sg) and the synganglion (Sy)became more visible (Fig. 3A, 3B, 4C 6E).

Two pairs of larger white tubes located laterally to the midgut, represent the Malpighian tubules (Mt), which are clearly visible. The Malpighian tubules, which are part of the tick excretory system, are connected to the posterior of the intestines, known as the hindgut (Hin), and the rectal sac (Rs).

Dissecting the digestive and the nervous system: salivary glands and synganglion removal

Two dissection forceps were employed to collect the salivary glands. One forceps grasped the
principal salivary duct at the base of the chelicerae (Ch) at the anterior upper end, while the other gently
pulled out the salivary glands. As with the midgut, the salivary glands were rinsed in three baths of sterile
PBS prior to collection and cleared of any residual tracheae and connective tissue (Ct) (Fig. 2C). The same
procedure was applied to remove the salivary gland on the opposite side. The size of the salivary glands
varies with the feeding status of the ticks, becoming larger in half-fed and fully-fed ticks. The glands could
be rinsed in three pools of PBS on either a Petri dish or a sterile glass slide, using aseptic dissecting
forceps.
Once the salivary glands have been removed, the heart became visible especially in nymphs (Fig. 4C).
The heart (H) lies in the dorsal portion along the mid-sagittal axis and is connected anteriorly with the
aorta, and leads to the synganglion (Fig. 4C). Given the fragility of the synganglion, its removal requires
precision. One forceps can be used to grasp the whitish neuronal extensions located on its lateral side.
262
263   *Dissecting the reproductive organs: testes and ovaries removal*
264   Following the removal of the synganglion, the tick's reproductive system, particularly in males,
265   becomes more apparent (Fig. 6). It is important to note that the reproductive system only develops at the
266   adult stage. Consequently, the nymph lacks both reproductive organs and a gonopore, which are
267   essential for the reproduction (Fig. 4C, 4D).
268
Dissection of males: testes
The male reproductive system includes the genital opening (gonopore), testes (T), vasa deferentia
(Vd), ejaculatory duct, seminal vesicle (Sv), and accessory reproductive gland complex (Arg). Since the
internal anatomy of *A. variegatum* males is simpler than that of females, isolating the organs is relatively
easier (Fig. 2A, 6B). After removing the trachea and the midgut, the reproductive system (Mro) becomes
exposed, particularly the testes, which occupy significant space (Fig. 6C, 6E). Using dissecting forceps, the
ejaculatory duct can be gently pulled out of the tick's body, bringing the entire reproductive system with
it (Fig. 6F, 6G).
277
Dissection of females: ovaries
The Gené's organ (Go), a specialized structure found only in female ticks, is located in the anterior
region of the cavity, just above the basis capituli. This structure facilitates coating the eggs with a wax-like
substance during oviposition.
During dissection, with the ventral surface of the tick facing downward, the Gené's organ is located
dorsally, above the synganglion, while the female reproductive apparatus is positioned ventrally to the
synganglion. For better access to Gené's organ, the capitulum can be removed with a surgical blade or
scissors (Fig. 9A).
An incision should be made horizontally beneath the basis capitulum, just above the camerostome, to
avoid damaging the organ. Once the connective tissue and residual tracheae are cleared from the tick's
internal cavity, the Gené's organ becomes visible. Using dissecting forceps, the Gené's organ can be
extracted from the camerostomal opening by firmly grasping it. It should be noted that the tubular
accessory glands of the Gené's organ do not fully develop until the female has completed a blood meal
(Fig. 9A, 9B). Following the removal of the trachea and the midgut, the oviducts (Ovi) leading to the ovary
(O) in the posterior region, and the vagina, located ventrally to the synganglion, should become visible.
The vagina was then grasped with dissecting forceps, and the entire female reproductive apparatus
was carefully removed from the internal cavity (Fig. 9C). This apparatus includes the bipartite vagina,
seminal receptacle, uterus, accessory glands, oviducts, and ovary. It should be noted that these
components have not been further differentiated. Similar to males, the female genital opening is located
ventrally relative to the synganglion (Fig. 10A).
298
Practical Notes
During dissections, care must be taken to avoid tissue tearing, especially for fragile organs like ovaries
and synganglion. Using fine, aseptic tools and working under a stereomicroscope significantly reduces the
risk of damage. Adhesive tapes and clean dissecting surfaces are essential to maintain tissue integrity and
minimize contamination. For less experienced researchers, it is advisable to practice dissections on

nymphal or male ticks, which have simpler anatomy, before attempting engorged females. If tissues are
intended for downstream analyses, such as transcriptomics or proteomics, meticulous handling and
immediate storage in appropriate solutions (e.g. Phosphate Buffer Saline (PBS) or RNAlater) are critical to
preserve RNA or protein quality.
308

Results and Discussion

Ticks are ectoparasites arthropods that feed on the blood of vertebrates and invertebrates and are
capable of transmitting various pathogens, thus playing a major role in the global epidemiology of
infectious diseases. Their indirect pathogenicity constitutes a significant threat to both animal and human
health, and contributes to economic losses due to their impact on livestock. In tropical and subtropical
regions, the tropical bont tick (*Amblyomma variegatum*) is the principal vector of *Ehrlichia ruminantium,*
an obligate intracellular bacterium of high pathogenic potential. As such, anatomical visualization and
study of this tick species are of particular interest for further research into vector-pathogens interactions.
317

External Morphology and Key Taxonomic Features
A. variegatum *ticks are characterized by a distinctive external scutum ornamentation*
Proper dissection of *A. variegatum* necessitates familiarity with its external morphology. of the tick,
which is the natural vector of the bacterium that causes heartwater in ruminants. The tick body, similar in
structure to that of parasitiform acarines such as mites is divided into two main regions: the gnathosoma
(capitulum) and the idiosoma (body region) (Fig. 1). The capitulum, considered as the "head", is located
in the anterior part and bears the mouthparts - the chelicerae, hypostome, and palps - but does not
house the eyes or brain, which are located within the idiosoma.
The basis capituli of *A. variegatum* is rectangular to oval, as observed in *A. parvitarsum* and *A.*
lepidum (Abouelhassan et al., 2024; Estrada-Peña et al., 2005). Additionally, the basis capituli, located in
the capitulum, bears two paired structures known as porose areas which are nearly circular in shape in *A.*
variegatum females. The porose areas of the female *A. variegatum* tick are observed consistently with
those arranged in *A. varium* or in the *A. parvitarsum* tick, which belongs to the *A. maculatum* group
(Estrada-Peña et al., 2005; Onofrio et al., 2008). In contrast, a greater interdistance between the porose
areas was observed in *A. maculatum, A. tigrinum,* and *A. triste.*
A. variegatum is considered a large tick, ranging from 6-7 mm when unfed to up to 30 mm in
repleted females, similar in size to *A. crissum* and *A. gaeyi* (Martins et al., 2013; Voltzit, 2007). The size of
A. variegatum hypostome is of a particular concern, as it has been observed to cause further damage to
the skin of its host (Kröber and Guerin, 2007). These ticks are classified within the Longirostrata family,
distinguished by their elongated mouthparts, a feature shared by species such as *A. varium* and *A.*
coelebs (Voltzit, 2007). The hypostome of *A. variegatum* ticks is characterized by its elongated cylindrical
form, situated at the same level as the two paired palps, a feature shared by the *A. sculptum* and *A.*
patinoi tick species (Nava et al., 2014).
The idiosoma is further divided into the anterior podosoma, which includes the genital gonopore and
four pairs of legs, and the posterior opisthosoma, which contains spiracular plates and the anal pore
(Sonenshine and Simo, 2021). Additionally, the idiosoma includes the walking legs. As ixodid ticks (hard-
bodied ticks), *A. variegatum* displays a teardrop-shaped body with a dorsal shield, or scutum (Cupp,
1991).
In males, the scutum (conscutum) fully covers the dorsum, restricting the ingestion of large quantities
of blood during blood feeding. *A. variegatum* males exhibit unique ornate pattern in bright orange colors
and dark hues, a characteristic also found in other African species such as *A. lepidum, A. hebraeum, A.*
pomposum, and *A. eburneum* (Fig. 1B, 6A) (Abouelhassan et al., 2024; Smit et al., 2024).
In contrast, the scutum of adult females, nymphs and larvae, does not cover the entire dorsum. The
female scutum consists of a large subtriangular plate near to the capitulum, while the remaining two-
thirds of the idiosoma is made up of a less sclerotized plate, the alloscutum (Fig 1A, 4A). In contrast to the
distinctive ornamentation, shape, and shiny colors exhibited by *A. variegatum* males, *A. variegatum*
females are distinguished by a less specific figure. The scutum of the female exhibits bright colors that
evoke the conscutum of the male. However, given that the scutum is smaller in size compared to the rest

of the body, several species of *Amblyomma* ticks exhibit remarkably similar patterns (Nava et al., 2014).
During feeding, the alloscutum stretches to accommodate a significant increase in volume, allowing
females to reach up to 100 times their unfed weight (Fig. 7A).
Furthermore, the alloscutum of larvae and nymphs tends to enlarge as they ingest blood. *A.*
variegatum nymphs' overall morphology is similar to that observed in *A. hebraeum* nymphs (Arthur,
1973). Lateral to the scutum/ conscutum, two hemispherical, dark-colored orbital eyes can be observed
(Fig. 1A, 1B, 4A). *A. variegatum* possess eyes that are distinctly convex, displaying notable similarities to
those of *H. impressum* or *H. dromedarii* when compared with *R. sanguineus*, where the eyes are not
distinctive (Abdel-Shafy et al., 2011; Abdullah et al., 2016).
The locomotor system includes four pairs of legs in adults and nymphs, and three in larvae. Each leg is
composed of six podomeres—coxa, trochanter, femur, patella, tibia, and tarsus—arranged proximally to
distally on the podosoma (Sonenshine and Roe, 2013). The legs of ticks are highly flexible, owing to
elastically articulated segments with bright colors in *A. variegatum* ticks (Fig. 1). In the *A. variegatum*, the
coloration of the legs exhibits a pattern alternating between brown and pale rings. This trait is also
observed in some *Hyalomma* species (Abdullah et al., 2016). *A. variegatum* legs are slightly slender in
comparison with the size of its body, in contrast to the bulbous legs exhibited by *Margoropus* ticks
(Walker et al., 2003). At the posterior end of the opisthosoma of male and female ticks, as well as
nymphs, several festoons are present, separated by a marginal groove. In certain species of the
Amblyomma genus, males have been observed to exhibit enamel displayed as patches on individual
festoons (Walker et al., 2003). However, *A. variegatum* males do not exhibit enamel on their festoons, in
contrast to *A. hebraeum*, *A. lepidum*, and *A. eburneum* (Abouelhassan et al., 2024; Smit et al., 2024).
377

Internal Anatomy of *A. variegatum*
A. variegatum *females have a well-developed tracheal system*
The tracheae are part of the respiratory system of ticks and facilitate gas exchanges through the
body and organs of the tick (Woolley, 1972). The tracheal system of the tick is extensive and opens
externally through two lateral spiracular plates (Starck et al., 2018). It is noteworthy that there is
considerable variation in the structure of the respiratory system among ticks, particularly with regard to
the morphology of the spiracles. In *A. variegatum*, the spiracular plates of the female specimen are
considerably larger than those of the male specimen. This is due to the fact that females have a larger
body size and distinct physiology. These physiological mechanisms, including the digestion of large blood
meals, excretion, and egg production, require larger spiracular plates to ensure sufficient surface area for
energy exchange via gas exchange (Baker, 1997). Thus, the tracheal system of *A. variegatum* females is
much denser than that of the males, complicating its dissection (Fig. 2A, 6B). In contrast, larval tracheae
are rudimentary and enable passive gas diffusion across the integument (Binnington and Obenchain,
1982).
392

The digestive system of A. variegatum *ticks is of greater proportions than that of other tick species*
In ixodid ticks, the pharynx is an elongated organ situated on the ventral surface of the basis
capitulum. It facilitates the passage of ingested host blood through the food channel to the esophagus
and midgut (Sonenshine and Simo, 2021). The gut extends throughout the entire body region of the tick
and represents its largest organ. The entire tick gut includes the midgut, with a central portion, the
stomach, which is connected to several dark brown, spider-shaped diverticula (Fig. 2, 4B, 6B). During the
tick's engorgement phase, the midgut-diverticula are filled with host blood to be digested. Due to the
considerable size of the midgut, other internal organs may be displaced or compressed (Sonenshine and
Simo, 2021). This phenomenon is particularly salient for *A. variegatum* females, as they possess a larger
body size and consume greater quantities of blood compared to other tick species. The hindgut and the
Malpighian tubules are key components of the tick's excretory system, playing vital roles in maintaining
homeostasis and excreting guanine-rich wastes while transporting them to the rectal sac (Šimo and Park,
2014). In ticks, nitrogenous excreta from the Malpighian tubules and undigested waste from the midgut
are concentrated in the rectal sac and eliminated through the anus via a short rectum (Fig. 4B) (Šimo and
Park, 2014). These connections are situated beneath the intestinal tissue. In Fig. 6D, the excretory system
is faintly visible once it has been extracted from the cavity.

Tick salivary glands are complex, multi-lobed organs that serve multiple functions during both the
tick's parasitic and non-parasitic periods on the host (Bowman and Sauer, 2004). The salivary glands are
paired structures located in the anterolateral regions of the tick's body. The salivary glands of female
ixodid ticks consist of clusters of grape-shaped alveoli, called acini, which are categorized into three
morphotypes: type I, type II, and type III (Šimo et al., 2017). *A. variegatum* tick salivary glands appeared
slightly more opaque compared to those of *R. microplus* (Cossio-Bayugar et al., 2023) and *I. scapularis*
ticks (Sonenshine and Simo, 2021). As observed in partially engorged but interrupted *A. variegatum*
females, the salivary glands exhibited greater translucency compared to those of unfed females (Fig. 2A,
2C, 3A, 8A). This observation is consistent with the findings of a recent study on *R. haemaphysaloides*,
which reported the degeneration of salivary glands in engorged female ticks after 7 days of feeding
(Wang et al., 2021). The hypothesis that the initiation of autolysis of salivary glands was associated with a
peak in hemolymph ecdysteroid concentration was proposed (Mao and Kaufman, 1999). Furthermore,
evidence suggests that the release of ecdysteroids during the post-engorgement period triggers both
salivary gland degeneration and ovarian development (Friesen and Kaufman, 2009). These
findings support the need for careful handling during saliva collection. In this study, tick saliva was
collected subsequently to the injection of pilocarpine through tick spiracles. However, it has been
presumed that pilocarpine is active in certain bioassays (Ribeiro et al., 2004). Consequently, a cautious
interpretation of the results obtained from tick saliva samples treated with pilocarpine is imperative.
427
428
The nervous system of ixodid ticks, including A. *variegatum, appears to be significantly similar*
The salivary glands are innervated by branched peripheral nerves arising from the synganglion, the
central nervous system of ticks. This organ performs several essential functions in ticks, including
movement control and stimulus processing (Rispe et al., 2022). The synganglion is located in the ventro-
anterior region, posterior to the midgut stomach, and is composed of multiple ganglia (Obenchain, 1974).
The synganglion is a translucent, heart-shaped structure linked to a multitude of peripheral nerves (Pn)
that innervate various parts of the tick body (Sonenshine and Simo, 2021). A comparison of the
anatomical structure of *A. variegatum* synganglion and *R. microplus* or *I. scapularis* reveals significant
similarities (Cossio-Bayugar et al., 2023; Szlendak and Oliver Jr., 1992; Tidwell et al., 2021).
As part of an open circulatory system, the pumping action of the heart and the arterial vessels
facilitates circulation within the hemocoel (Fig. 4C)(Binnington and Obenchain, 1982). The heart appears
pentagonal in systole and spherical in diastole (Cossio-Bayugar et al., 2023). In comparison with nymphs,
the internal structure of adult *A. variegatum* ticks is more complex, which hinders the visualization of the
heart.
443
A. variegatum *male testes exhibit a tubular and elongated shape*
The male reproductive system includes the gonopore, testes, vasa deferentia, ejaculatory duct,
seminal vesicle, and accessory gland complex. The male gonopore is located in the ventroanterior region
of the body, ventral to the synganglion, and opposite the anal aperture (B.R. Sampieri et al., 2016).
The male posterior region is composed of a pair of tubular and elongated testes, positioned
dorsolaterally within the opisthosoma. These are connected to the accessory reproductive glands by the
vasa deferentia (Fig 6E, 6F)(Anholeto et al., 2015; Sampieri et al., 2014). The testes serve as the primary
site of spermatogenesis, which begins during the nymphal stage. The appearance of tick testes of ticks
may vary slightly even within a single species, depending on the nutritional and reproductive status of the
individual (Oliver and Brinton, 1972). In metastriate ticks, the testes are linked by an extremely thin
strand of filamentous tissue and taper into the vasa deferentia. The testes are located dorsally to the
synganglion, and laterally to the rectal sac (Oliver, 1982). *A. variegatum* males possess a reproductive
system that is morphologically similar to that of *R. sanguineus* (Bruno Rodrigues Sampieri et al., 2016).
However, a notable distinction emerges in the structure of the seminal vesicles, which exhibit a specific
V-shape in *A. variegatum* male ticks, contrasting with the more elongated appearance observed in *R.*
sanguineus. While both *A. variegatum* and *R. sanguineus* possess two pairs of testes, *O. rostratus*, by
contrast, appears to lack seminal vesicles and presents a single horseshoe-shaped testis. Furthermore, a
comparison of *R. microplus* males and *A. variegatum* males reveals significant differences in the
morphology of their testes. Specifically, *A. variegatum* males possess testes that are thinner and less

pronounced in the opisthosoma compared to *R. microplus* males, which have more rounded and
translucent testes (Tidwell et al., 2021). Furthermore, the testes and the accessory glands complex of fed
male *A. variegatum* are slightly larger than those of *A. hebraeum* (Weiss et al., 2002). The vasa deferentia
represent the narrowing of the paired testes at their anterior ends, joining in the anterior region to form
a single duct called the vas deferens or seminal vesicle. As with the testes, the shape and the size of the
vasa deferentia depend on the reproductive state of the tick and can vary from narrow, coiled tubes to
enlarged, straight structures (Oliver, 1982). In accordance with observations made in other *Amblyomma*
species, including *A. sculptum*, *A. aureolatum* and *A. triste*, the vasa deferentia re short, consistent with
the characteristics observed in *A. variegatum*. As demonstrated by Sampieri *et al* (2016), *A. triste*
possesses an elongated tissue linking the two testes, in contrast to *A. variegatum* which lacks this
isthmus. Histologically, the vasa deferentia are indistinguishable from the common vas deferens. The
seminal vesicle serves as the point of connection between the vasa deferentia and the ejaculatory duct.

The ejaculatory duct is a short, narrow tube located in the ventroanterior region, extending below the
synganglion to the external gonopore (El Shoura, 1987). The male accessory reproductive glands (Arg)
play critical roles in all stages of the reproductive biology of the mated female, from the sperm
deposition to oviposition (Gillott, 2003). For instance, male accessory glands stimulate engorgement in
females and activate spermatozoa upon arrival at the female seminal receptacle (Reuben Kaufman,
2007). This multi-lobed complex is located ventrally to the opisthosoma and connects to the ejaculatory
duct on its ventral face and to the vasa deferentia on the dorso-lateral region (Sampieri et al., 2014).

Mated male ticks are capable of excreting a sac-like spermatophore that protrudes from their genital
opening and delivering it with their mouthparts to the female's gonopore (Fig. 10D)(Kiszewski et al.,
2001). During mating, metastriate males appear to insert only their chelicerae into the female's genital
pore, leaving the hypostome and palps outside, barely in contact with the female's surface (Kiszewski et
al., 2001)(Pature *et al*, 2025, unpublished data). This mechanism ensures the successful transfer of the
sac-like spermatophore protruding from the male genital opening (Fig. 10C, 10D).

*The female tick (*A. variegatum*) reproductive organs are more challenging to observe under the*
microscope

The female genital system comprises the gonopore, similar to that in male, ovaries, a common
oviduct or uterus, a bipartite vagina divided into a cervical and vestibular vagina, and a seminal
receptacle (Sr). Additionally, females possess two accessory glands: tubular and lobular. In *A. variegatum*,
anatomical visualization of the uterus and vagina proves particularly challenging.

As demonstrated in *A. cajennense* (Denardi et al., 2004), *R. sanguineus* (de Oliveira et al., 2005), and
R. microplus (Saito et al., 2005), *A. variegatum* females possess a unique, tubular, horseshoe-shaped, and
continuous ovary located in the posterior part of the opisthosoma (Fig. 3C, 5). The tick ovary is a hollow
organ filled with oogonia and primary oocytes at various early stages of development. The ovary of unfed
A. variegatum females is slightly smaller than average and appears translucent. It seems to be
surrounded by a tunica propria, as observed in *H. longicornis* (Fig. 2C)(Yang et al., 2014). A fold or
longitudinal groove extends along the antero-dorsal surface of the ovary , containing primary oocytes
internally at their earliest stage of development, and more advanced oocytes externally (Fig 9C) (Diehl et
al., 1982). During feeding, the female's oocytes tend to protrude into the hemocel, giving the ovary a
grape-like shape. As demonstrated in *A. rotundatum*, a parthenogenetic tick species, we observed
oocytes of different sizes at various developmental stages after the dissection of a semi-engorged female
(Fig. 8A)(Sanches et al., 2012). As *A. variegatum* ticks are known to have a complex life cycle in the wild,
they exhibit a high reproductive potential, with an oviposition capacity of up to 20,000 eggs. Here, an *A.*
variegatum female was observed after the pre-oviposition period, indicating its imminent oviposition
(Fig. 7). The tick's entire body was found to be filled with mature eggs and oocytes at earlier stages of
development. It is evident that the tick ovary is significantly larger in size in comparison to its unfed state,
which is consistent with the large size of *A. variegatum* females during their blood meal on a host.

At each end of the ovary are whitish, paired, and folded oviducts (Ovi), which converge to form a
common oviduct or uterus (Fig. 9A). The oviduct tends to stretch considerably during the passage of the
eggs due to peristaltic contractions of the oviduct walls (Fig. 7D). The uterus connects to the cervical
vagina via a short tube. As indicated in the extant literature, the fed but unmated *A. variegatum* female
observed here, possessed two thin folded oviducts that lead to the female reproductive organs (Fig. 9A).

These oviducts have a thin diameter, similar to those observed in *R. microplus* females (de la Vega et al.,
2012).

The vagina consists of a posterior, broad, and short cervical region, as well as an anterior, narrow, and
long vestibular region (El Shoura, 1989). In addition to the uterus, the seminal receptacle, tubular
accessory glands and the vestibular vagina converge in the cervical vagina (Fig. 9B) (Vega et al., 2012).
The vestibular vagina extends from the gonopore to the cervical vagina and opens externally to the
gonopore via the vulva. In the female *A. variegatum*, the vagina situated in the podosoma is more
complex to observe.

The seminal receptacle is a sac-like extension of the cervical region that serves as the storage site for
spermatozoa after mating, facilitating the fertilization of eggs (Oliver, 1989). Final stages of
spermiogenesis occur in the seminal receptacle of copulated females until full engorgement (Fig. 10C)
(Brinton et al., 1974). This structure is positioned above the uterus and opens postero-dorsally into the
cervical vagina (Diehl et al., 1982). In the present study, an ectospermatophore was observed to be
adhered near the genital aperture of the *A. variegatum* female (Fig. 10A). The spermatophore of *A.*
variegatum males appears larger than that of *R. microplus* (de la Vega et al., 2012). *A. variegatum*, like
other Argasid ticks, possesses an ectospermatophore with a bulb-shaped capsule associated with a short
neck (Feldman-Muhsam, 1967). Additionally, an endospermatophore was observed within the seminal
receptacle of the female (Fig. 10C). As previously documented, a sheath surrounding the
endospermatophore was observed in *A. variegatum*, likely forming the capsule enclosing the sperm
(Feldman-Muhsam and Borut, 1983). However, the literature also documents that the
endospermatophore of ixodid ticks is monolobed with the formation of a unique capsule (Feldman-
Muhsam, 1967). However, consistent with the observations made by Oliver *et al* (1974), who noted the
presence of an additional small sphere, termed the "endospermatophore neck", in proximity to the
endospermatophore in *D. occidentalis* (Oliver, 1974), we have observed a comparable structure in the
spermatophore of *A. variegatum*.

The Gené's organ is unique to female ticks. It secretes a waxy substance that covers the oviposited
eggs, protecting them from the environment. A peculiarity of this organ is its ability to evert itself from
the camerostomal aperture, located between the posterior base of the capitulum and the anterior region
of the scutum. Internally, this organ is situated in the anterior body region, below the capitulum, and
consists of branched tubular glands that are highly developed in the engorged females (Fig. 9A). In this
study, we observed that the female *A. variegatum* Gené's organ exhibited a structural similarity to that of
R. microplus females, thereby highlighting the conventional form of this organ (Tidwell et al., 2021).

The tubular (Tag) and lobular accessory glands are associated with the Gené's organ and also
contribute to the egg waterproofing during the oviposition period of the female. The tubular glands form
the bulk of the Gené's organ and are located anteriorly within the hemocoel, in continuity with the horns
(Fig. 9B). The paired tubular accessory glands open into the vagina at the dorsal boundaries of the
cervical and the vestibular regions (Booth, 1985). During oviposition, the lumen of each gland fills with a
secretion that coats the egg surface as it passes through the vagina (Diehl et al., 1982). In contrast with *R.*
microplus, the tubular accessory glands of *A. variegatum* females are less visible when fully engorged
(unpublished data). This observation may be attributable to the considerable place of *A. variegatum* eggs
within the tick body (Fig. 7).

The lobular accessory gland is a distinctive feature of ixodid ticks. This gland results from the
maturation of the vestibular vaginal epithelium as the entire female reproductive system develops. Two
porose areas are located dorsally on the female basis capituli, below the rostrum (Fig. 1A). During
oviposition, these areas are covered by the Gené's organ. The porose areas secrete a substance that is
incorporated into the waterproofed wax of the organ of Gené as it everts from the camerostomal fold. In
this study, the complete reproductive system of the *A. variegatum* female has not been observed. This
could be attributed to the absence of the dissection of a more advanced engorged and mated female
(after 15 days of feeding during their rapid engorgement phase, instead of 12 days) or a repleted, newly
detached *A. variegatum* female, which would allow full anatomical observation of the mature
reproductive system.

568
A. variegatum *adults exhibit a blueish hue in its hemolymph*

Prior to hemolymph and saliva collection, it is essential to sterilize the tick cuticle to eliminate
external bacterial contaminants that could compromise the integrity of internal tissues. In this study,
ticks were surface-decontaminated using ultrapure water followed by phosphate-buffered saline (PBS),
as previously described (da Silva Vaz Junior et al., 2024; Kim et al., 2016). Although some reports suggest
that incomplete decontamination does not critically affect internal microbiome profiles, it may introduce
bias in the composition of bacterial communities recovered from internal compartments (Hoffmann et
al., 2020). Recent studies have highlighted that bleach is more effective than ethanol for surface
decontamination (Binetruy et al., 2019); however, it may also disrupt internal microbiota. These
observations emphasize the need for a standardized, reproductible sterilization protocol.

The collection of hemolymph—the circulatory fluid that fills the tick's hemocoel—is particularly
relevant due to its hemocyte composition, which plays a central role in tick immune responses and
interactions with pathogens. Hemolymph components are crucial to understanding bacterial survival and
adaptation within the tick environment. As demonstrated in the case of *A. americanum*, the complete
depletion of granulocytic hemocytes resulted in the death of the tick following an *Ehrlichia chaffeensis*
infection (Adegoke et al., 2023). In this paper, we propose a simple and reproducible method for
hemolymph collection by excising the distal tarsus and collecting the exuding fluid. This method is
especially useful for partially or fully engorged females. Alternative approaches can be used to maximize
hemolymph yield and purity in fed ticks (Aguilar-Díaz et al., 2022). The hemolymph of insects is
characterized by a translucent appearance, ranging from pale amber to green hues, due to the absence
of hemoglobin pigment (Hackman, 1952). Interestingly, in adult *A. variegatum*, we observed a distinctive
bluish hue during the initial incision, a phenomenon rarely reported in other tick species such
as *Haemaphysalis flava, Ixodes scapularis*, or *Rhipicephalus microplus* (Aguilar-Díaz et al., 2022; Liu et al.,
2022; Patton et al., 2012; Tidwell et al., 2021). The underlying mechanism responsible for this
phenomenon remains to be elucidated due to the paucity of research focusing on tick hemolymph
pigment composition. Additionally, a distinctive odor was noted during dissection, reinforcing the
importance of appropriate personal protective equipment, such as a mask, to ensure operator safety.
596

Prior research has also demonstrated that tick sex significantly influences microbial community
composition. In ecological terms, the bacterial richness and diversity within a sample are referred to as
"alpha diversity" (Andermann et al., 2022). For instance, *Dermacentor variabilis* males exhibit higher
alpha diversity than females, a trend likely attributable to their lower blood intake and repeated feeding
behavior, which allow acquisition of a broader bacterial repertoire (Duncan et al., 2022). Performing sex-
specific dissections may therefore improve the resolution of microbiota analyses, particularly in engorged
females. Given the contribution of male salivary glands to mating via spermatophore production and
delivery, it is plausible that female and male ticks employ distinct developmental and physiological
strategies (Tan et al., 2015).
606

Conclusion

The collection of tick organs is crucial for understanding vector-pathogen interactions *in vivo*. In this
study, we provide a comprehensive anatomical description of *A. variegatum,* long with practical guidance
for isolating key internal structures—most notably the midgut, salivary glands, and ovaries—which play
critical roles in vector competence.

The dissection procedures presented here offer a reliable approach to accessing internal tick tissues,
provided that sufficient technical skill is applied to avoid degradation or contamination. Due to the
intricate organization and fragility of internal organs, especially in engorged females, meticulous handling
is required when preparing specimens for downstream applications.

This foundational work opens the way for future molecular and cellular investigations targeting
specific tissues, to better understand the physiological responses of ticks to pathogen colonization—at
both the biological and transcriptomic levels. Despite its major role in transmitting *Ehrlichia*
ruminantium, A. variegatum remains less studied than other species such as *Rhipicephalus microplus*. The
present study fills an important gap by offering the first high-resolution microscopic observations and a
general anatomical framework for this neglected but epidemiologically significant vector.

Figure 1: Anatomical description of the dorsal surface of unfed adult *A. variegatum* ticks mounted on double-sided adhesive tape. **(A)** Dorsal view of unfed *A. variegatum* female (0.78x). **(B)** Dorsal view of unfed *A. variegatum* male, with its first leg cut to allow hemolymph collection (0.78x).

Figure 2: Dorsal view of an unfed *A. variegatum* female and its organs. **(A)** Scutum dissection: dorsal view of an unfed *A. variegatum* female with the scutum removed and immersed in PBS (1.6x). **(B)** Midgut dissection: close up view of the gut of an unfed *A. variegatum* female (1.25x). **(C)** Salivary glands dissection: close-up view of the salivary glands of an unfed *A. variegatum* female (3.2x). Ac: acini ; Div: diverticulum ; Sac: <u>scutum</u>/ alloscutum/conscutum (in the female tick) ; Sd: salivary duct ; Sg: salivary glands ; St: stomach ; Tr: tracheae.

Figure 3: Dorsal view of an unfed *A. variegatum* female with the midgut removed and organs. **(A)** Midgut dissection: dorsal view of unfed an *A. variegatum* female with scutum removed and immersed in PBS (1.6x). **(B)** Synganglion dissection: close-up view of the nervous system of unfed an *A. variegatum* female (5x). **(C)** Reproductive system dissection: close-up of the ovary of an unfed *A. variegatum* female (5x). O: ovary ; Ovi: oviducts ; Pn: peripheral nerves ; Sac: <u>scutum</u>/ alloscutum/conscutum (here, in the female tick) ; Sy: synganglion ; Sg: salivary glands ; Tr: tracheae.

Figure 4: Dorsal view of an unfed *A. variegatum* nymph. **(A)** Dissection of the scutum: dorsal view of an unfed *A. variegatum* nymph immersed in PBS, showing a clean incision (yellow dotted line) around the idiosoma (2.5x). **(B)** Removal of the scutum: dorsal view of an unfed *A. variegatum* nymph immersed in PBS, with the conscutum lifted using forceps (2.5x). **(C)** Dissection of the midgut: *A. variegatum* nymph with the entire gut removed and the conscutum cut off (3.2x). **(D)** Unfed *A. variegatum* nymph with the salivary glands, midgut, tracheae and Malpighian tubules removed (4x). Div: diverticulum ; H: heart ; Mt: Malpighian tubules ; Rs: rectal sac ; Sac: <u>scutum/ alloscutum</u>/conscutum (here, in the nymph tick) ; Sy: synganglion ; pink circle : hematin granules released after gut tear ; black triangle: anal aperture.

Figure 5: Dorsal view of an unfed *A. variegatum* female with the midgut, and salivary glands removed (2x). Ch: chelicerae ; Div: diverticulum ; O: ovary ; Rs: rectal sac ; Sp: spiracular plates ; Tr: tracheae.

Figure 6: Dorsal view of a fed and mated *A. variegatum* male. **(A)** Dorsal view of a fully repleted *A. variegatum* male with a rounded conscutum (1.25x). **(B)** Scutum dissection: dorsal view of a fed *A. variegatum* male with folded conscutum, immersed in PBS (1.6x). **(C)** Midgut dissection: *A. variegatum* male with the entire gut removed (1,6x). **(D)** Close up of the excretory system (2.5x). **(E)** Salivary glands dissection: *A. variegatum* male with the paired salivary glands removed (2.5x). **(F, G)** Reproductive organs dissection: highlighting the organs of the male reproductive system (1.6x, 2.5x). Arg: accessory reproductive glands ; Cc: cement cone ; Ct: connective tissue ; Div: diverticulum ; Hin: hindgut ; Mro: male reproductive organs excluding testes ; Mt: Malpighian tubules ; Rs: rectal sac ; Sac: scutum/ alloscutum/<u>conscutum</u> (in the male tick) ; Sg: salivary glands ; St: stomach ; Sv: seminal vesicle ; Sy: synganglion ; T: testes ; Tr: tracheae ; Vd: vasa deferentia.

Figure 7. View of a fully repleted *A. variegatum* female. **(A)** Dorsal view of a fully engorged *A. variegatum* female (0.78x). **(B)** Dorsal view of an ovipositing *A. variegatum* female with scutum removed (0.78x). **(C)** Dorsal view of *A. variegatum* with thousands of eggs expelled from the oviducts (0.78x). **(D)** Close up of oviducts full of eggs (0.78x). Cp: capitulum ; Div: diverticulum ; E: eggs ; Ooc: oocytes ; O: ovary ; Tr: tracheae.

Figure 8: Dorsal view of a partially fed *A. variegatum* female with the midgut removed and organs. **(A)** Dorsal view of a semi-engorged female *A. variegatum* female with the midgut removed (1x). **(B)** Close-up of the midgut of a partially fed female *A. variegatum* female (0.78x). Mt: Malpighian tubules ; O: ovary ; Sg: salivary glands ; Sp: spiracular plates.

Figure 9: Dorsal view of a partially fed *A. variegatum* female with the midgut, and salivary glands
removed, showing reproductive organs. **(A)** Dorsal view of a female *A. variegatum* with a close-up of
reproductive system (2x). **(B)** Close up of reproductive organs with accessory glands (2x). **(C)** Close-up of
ovary with oocytes (1,6x). Ch: chelicerae ; Fro: female reproductive organs excluding ovary and oviducts ;
Go: Gené's organ ; Mt: Malpighian tubules ; Ooc : oocytes ; Ovi: oviducts ; Sy: synganglion ; Tag: tubular
accessory glands ; Tr: tracheae.
680
Figure 10. View of a mated *A. variegatum* female. **(A)** Ventral view of a semi-engorged *A. variegatum*
female with an ectospermatophore attached near to its gonopore shortly after copulation (0.78x). **(B)**
Dorsal view of a partially fed *A. variegatum* female with a round alloscutum (0.78x). **(C)** Dorsal view of an
endospermatophore contained in the seminal receptacle of a semi-engorged *A. variegatum* female
(2.5x). **(D)** Close-up of the endospermatophore dissected a few minutes after the female mated with a
male (2.5x). Cc: cement cone ; O: ovary ; Ooc: oocytes ; Rs: rectal sac ; Sac: <u>scutum/</u>
<u>alloscutum/</u>conscutum (in the female tick) ; Sr: seminal receptacle ; Sp: spiracular plates ; Tr: tracheae ;
black triangle: anal pore ; pink triangle: gonopore ; red triangle: ectospermatophore.

Acknowledgements

690 The authors want to thank Mélanie Dhune for providing ticks from the CIRAD animal facility as well as
691 Loïc Jacquet-Crétides and Jimmy Dédy for their support with animal care. Naomie Pature is grateful to
692 Rosalie Aprelon for technical assistance with dissection.

Conflict of interest disclosure

694 The authors declare that they have no financial conflicts of interest in relation to the content of the
695 article.

Funding

697 This work is supported by the United States Department of Agriculture grant 58-3022-1-018-F (Risk of
698 Arthropod-borne diseases in the Caribbean).

References

Abdel-Shafy, S., El Namaky, A.H., Khalil, F.H.M., 2011. Scanning electron microscopy and morphometrics
of nymph and larva of the tick Hyalomma impressum (Acari: Ixodidae). Parasitol Res 109, 1509–1518.
https://doi.org/10.1007/s00436-011-2422-4

Abdullah, H.H.A.M., El-Molla, A., Salib, F.A., Allam, N.A.T., Ghazy, A.A., Abdel-Shafy, S., 2016.
Morphological and molecular identification of the brown dog tick Rhipicephalus sanguineus and the
camel tick Hyalomma dromedarii (Acari: Ixodidae) vectors of Rickettsioses in Egypt. Vet World 9, 1087–
1101. https://doi.org/10.14202/vetworld.2016.1087-1101

Abouelhassan, E.M., GadAllah, S., Kamel, M.S., Kamal, M., Elsayed, H.H., Sallam, N.H., Okely, M., 2024.
Molecular identification and morphological variations of Amblyomma lepidum imported to Egypt, with
notes about its potential distribution under climate change. Parasitol Res 123, 276.
https://doi.org/10.1007/s00436-024-08284-0

Adegoke, A., Hanson, J., Smith, R.C., Karim, S., 2023. Ehrlichia chaffeensis co-opts phagocytic hemocytes
for systemic dissemination in the Lone Star tick, Amblyomma americanum. J Innate Immun.
https://doi.org/10.1159/000535986

Aguilar-Díaz, H., Quiroz-Castañeda, R.E., Salazar-Morales, K., Miranda-Miranda, E., 2022. A newly
optimized protocol to extract high-quality hemolymph from the cattle tick Rhipicephalus microplus:
Improving the old conditions. Current Research in Parasitology & Vector-Borne Diseases 2, 100066.
https://doi.org/10.1016/j.crpvbd.2021.100066

Andermann, T., Antonelli, A., Barrett, R.L., Silvestro, D., 2022. Estimating Alpha, Beta, and Gamma
Diversity Through Deep Learning. Front Plant Sci 13, 839407. https://doi.org/10.3389/fpls.2022.839407

Anholeto, L.A., Nunes, P.H., Remédio, R.N., Camargo-Mathias, M.I., 2015. Testes of fed and unfed
mblyomma cajennense ticks (Acari: Ixodidae). First morphological data. Acta Zoologica 96, 375–382.
https://doi.org/10.1111/azo.12083

Arthur, D.R., 1973. Scanning electron microscope studies on the morphology of the immature stages of
Amblyomma hebraeum Koch, 1844. Journal of the Entomological Society of Southern Africa 36, 63–85.
https://doi.org/10.10520/AJA00128789_3081

Baker, G.T., 1997. Spiracular Plate of Nymphal and Adult Hard Ticks (Acarina: Ixodidae): Morphology and
Cuticular Ultrastructure. Invertebrate Biology 116, 341. https://doi.org/10.2307/3226866

Barré, N., 1989. Biologie et écologie de la tique Amblyomma variegatum (Acarina⬚: Ixodina) en
Guadeloupe (Antilles Françaises) (thesis). CIRAD-IEMVT.

Bezuidenhout, J.D., 1987. NATURAL TRANSMISSION OF HEARTWATER. Onderstepoort J. vet Res. 54, 349–
351.

Binetruy, F., Dupraz, M., Buysse, M., Duron, O., 2019. Surface sterilization methods impact measures of
internal microbial diversity in ticks. Parasites & Vectors 12, 268. https://doi.org/10.1186/s13071-019-
3517-5

Binnington, K.C., Obenchain, F.D., 1982. Structure and Function of the Circulatory, Nervous, and
Neuroendocrine Systems of Ticks, in: Physiology of Ticks. Elsevier, pp. 351–398.
https://doi.org/10.1016/B978-0-08-024937-7.50015-9

Booth, T.F., 1985. The structure and function of Gene's organ and its associated glands in ticks (phd).
Thames Polytechnic.

Bowman, A.S., Sauer, J.R., 2004. Tick salivary glands: function, physiology and future. Parasitology 129
Suppl, S67-81. https://doi.org/10.1017/s0031182004006468

Brinton, L.P., Burgdorfer, W., Oliver, J.H., 1974. Histology and fine structure of spermatozoa and egg
passage in the female tract of Dermacentor andersoni stiles (Acari-Ixodidae). Tissue and Cell 6, 109–125.
https://doi.org/10.1016/0040-8166(74)90026-3

Camargo-Mathias, M.I. (Ed.), 2018. Inside ticks: morphophysiology, toxicology and therapeutic
perspectives. Editora UNESP. https://doi.org/10.7476/9788595462861

Caperucci, D., Mathias, M.I.C., Bechara, G.H., 2009. Histopathology and Ultrastructure Features of the
Midgut of Adult Females of the Tick Amblyomma cajennense Fabricius, 1787 (Acari: Ixodidae) in Various
Feeding Stages and Submitted to Three Infestations. Ultrastructural Pathology 33, 249–259.
https://doi.org/10.3109/01913120903296945

Cossio-Bayugar, R., Miranda-Miranda, E., Kumar, S., 2023. A Laboratory Manual on Rhipicephalus
microplus, Cambridge Scholars Publishing. Lady Stephenson Library, Newcastle upon Tyne, NE6 2PA, UK.

Cupp, E.W., 1991. Biology of Ticks. Veterinary Clinics of North America: Small Animal Practice 21, 1–26.
https://doi.org/10.1016/S0195-5616(91)50001-2

da Silva Vaz Junior, I., Lu, S., Pinto, A.F.M., Diedrich, J.K., Yates, J.R., Mulenga, A., Termignoni, C., Ribeiro,
J.M., Tirloni, L., 2024. Changes in saliva protein profile throughout Rhipicephalus microplus blood feeding.
Parasites & Vectors 17, 36. https://doi.org/10.1186/s13071-024-06136-5

de la Vega, R., Diaz, G., Galán, M., Fernández, C., 2012. Anatomy and hystology of the female
reproductive system of Boophilus microplus (Acari: Ixodidae). Rev. Salud Anim. 34, 1–10.

de Oliveira, P.R., Bechara, G.H., Denardi, S.E., Nunes, É.T., Camargo Mathias, M.I., 2005. Morphological
characterization of the ovary and oocytes vitellogenesis of the tick Rhipicephalus sanguineus (Latreille,
1806) (Acari: Ixodidae). Experimental Parasitology 110, 146–156.
https://doi.org/10.1016/j.exppara.2004.12.016

Deans, A.R., Mikó, I., Wipfler, B., Friedrich, F., 2012. Evolutionary phenomics and the emerging
enlightenment of arthropod systematics. Invert. Systematics 26, 323. https://doi.org/10.1071/IS12063

Denardi, S.E., Bechara, G.H., Oliveira, P.R. de, Nunes, É.T., Saito, K.C., Camargo Mathias, M.I., 2004.
Morphological characterization of the ovary and vitellogenesis dynamics in the tick Amblyomma
cajennense (Acari: Ixodidae). Veterinary Parasitology 125, 379–395.
https://doi.org/10.1016/j.vetpar.2004.07.015

Diehl, P.A., Aeschlimann, A., Obenchain, F.D., 1982. CHAPTER 9 - Tick Reproduction: Oogenesis and
Oviposition, in: Obenchain, FREDERICK D., Galun, R. (Eds.), Physiology of Ticks. Pergamon, pp. 277–350.
https://doi.org/10.1016/B978-0-08-024937-7.50014-7

Duncan, K.T., Elshahed, M.S., Sundstrom, K.D., Little, S.E., Youssef, N.H., 2022. Influence of tick sex and
geographic region on the microbiome of Dermacentor variabilis collected from dogs and cats across the
United States. Ticks and Tick-borne Diseases 13, 102002. https://doi.org/10.1016/j.ttbdis.2022.102002

Edwards, K.T., 2009. Examination of the Internal Morphology of the Ixodid Tick, Amblyomma maculatum
Koch, (Acari: Ixodidae); a "How-to" Pictorial Dissection Guide.

El Shoura, S.M., 1989. Effect of blood meal and mating on the genital tract ultrastructure in the female
camel tick Hyalomma (Hyalomma) dromedarii (Ixodoidea: Ixodidae). Exp Appl Acarol 6, 157–175.
https://doi.org/10.1007/BF01201645

El Shoura, S.M., 1987. Fine Structure of the Vasa Deferentia, Seminal Vesicle, Ejaculatory Duct, And
Accessory Gland of Male Ornithodoros (Pavlovskyella) Erraticus (Acari: Ixodoidea: Argasidae). Journal of
Medical Entomology 24, 235–242. https://doi.org/10.1093/jmedent/24.2.235

Estrada-Pena, A., Ayllon, N., De La Fuente, J., 2012. Impact of Climate Trends on Tick-Borne Pathogen
Transmission. Frontiers in Physiology 3.

Estrada-Peña, A., Venzal, J.M., Mangold, A.J., Cafrune, M.M., Guglielmone, A.A., 2005. The Amblyomma
maculatum Koch, 1844 (Acari: Ixodidae: Amblyomminae) tick group: diagnostic characters, description of
the larva of A. parvitarsum Neumann, 1901, 16S rDNA sequences, distribution and hosts. Syst Parasitol
60, 99–112. https://doi.org/10.1007/s11230-004-1382-9

Feldman-Muhsam, B., 1967. Spermatophore Formation and Sperm Transfer in Ornithodoros Ticks.
Science 156, 1252–1253. https://doi.org/10.1126/science.156.3779.1252

Feldman-Muhsam, B., Borut, S., 1983. On the spermatophore of ixodid ticks. Journal of Insect Physiology
29, 449–457. https://doi.org/10.1016/0022-1910(83)90073-2

Friedrich, F., Matsumura, Y., Pohl, H., Bai, M., Hörnschemeyer, T., Beutel, R.G., 2014. Insect morphology
in the age of phylogenomics: innovative techniques and its future role in systematics. Entomological
Science 17, 1–24. https://doi.org/10.1111/ens.12053

Friesen, K.J., Kaufman, W.R., 2009. Salivary gland degeneration and vitellogenesis in the ixodid tick
Amblyomma hebraeum: Surpassing a critical weight is the prerequisite and detachment from the host is
the trigger. Journal of Insect Physiology 55, 936–942. https://doi.org/10.1016/j.jinsphys.2009.06.007

Gilbert, L., 2021. The Impacts of Climate Change on Ticks and Tick-Borne Disease Risk. Annu. Rev.
Entomol. 66, 373–388. https://doi.org/10.1146/annurev-ento-052720-094533

Gillott, C., 2003. Male Accessory Gland Secretions: Modulators of Female Reproductive Physiology and
Behavior. Annu. Rev. Entomol. 48, 163–184. https://doi.org/10.1146/annurev.ento.48.091801.112657

Grabowski, J.M., Kissinger, R., 2020. Ixodid Tick Dissection and Tick Ex Vivo Organ Cultures for Tick-Borne
Virus Research. Curr Protoc Microbiol 59, e118. https://doi.org/10.1002/cpmc.118

Hackman, R.H., 1952. Green pigments of the hemolymph of insects. Arch Biochem Biophys 41, 166–175.
https://doi.org/10.1016/0003-9861(52)90517-1

Hoffmann, A., Fingerle, V., Noll, M., 2020. Analysis of Tick Surface Decontamination Methods.
Microorganisms 8, 987. https://doi.org/10.3390/microorganisms8070987

Kennedy, B., Trim, S.A., Laudier, D., LaDouceur, E.E.B., Cooper, J.E., 2021. Arthropoda: Arachnida, in:
LaDouceur, E.E.B. (Ed.), Invertebrate Histology. Wiley, pp. 221–246.
https://doi.org/10.1002/9781119507697.ch8

Kim, T.K., Tirloni, L., Pinto, A.F.M., Moresco, J., Iii, J.R.Y., Jr, I. da S.V., Mulenga, A., 2016. Ixodes scapularis
Tick Saliva Proteins Sequentially Secreted Every 24 h during Blood Feeding. PLOS Neglected Tropical
Diseases 10, e0004323. https://doi.org/10.1371/journal.pntd.0004323

Kiszewski, A.E., Matuschka, F.-R., Spielman, A., 2001. M ATING S TRATEGIES AND S PERMIOGENESIS IN I
XODID T ICKS. Annu. Rev. Entomol. 46, 167–182. https://doi.org/10.1146/annurev.ento.46.1.167

Kröber, T., Guerin, P.M., 2007. An *in vitro* feeding assay to test acaricides for control of hard ticks. Pest
Management Science 63, 17–22. https://doi.org/10.1002/ps.1293

Lee, Junsoo, Ryu, J., Han, S., Ravichandran, N.K., Seong, D., Lee, Jaeyul, Wijesinghe, R.E., Kim, P., Lee, S.-
Y., Jung, H.-Y., Jeon, M., Choi, K.S., Kim, J., 2021. Identification of organs inside hard tick body using
spectral-domain optical coherence tomography. Infrared Physics & Technology 114, 103611.
https://doi.org/10.1016/j.infrared.2020.103611

Liu, L., Yan, F., Zhang, L., Wu, Z., Duan, D., Cheng, T., 2022. Protein profiling of hemolymph in
Haemaphysalis flava ticks. Parasites & Vectors 15, 179. https://doi.org/10.1186/s13071-022-05287-7

Mao, H., Kaufman, W.R., 1999. Profile of the ecdysteroid hormone and its receptor in the salivary gland
of the adult female tick, Amblyomma hebraeum. Insect Biochem Mol Biol 29, 33–42.
https://doi.org/10.1016/s0965-1748(98)00102-7

Martins, T.F., Scofield, A., Oliveira, W.B.L., Nunes, P.H., Ramirez, D.G., Barros-Battesti, D.M., Sá, L.R.M.,

Ampuero, F., Souza, J.C., Labruna, M.B., 2013. Morphological description of the nymphal stage of
Amblyomma geayi and new nymphal records of *Amblyomma parkeri*. Ticks and Tick-borne Diseases 4,
181–184. https://doi.org/10.1016/j.ttbdis.2012.11.015

Mathias, M.I.C., 2013. Guia básico de morfologia interna de carrapatos ixodídeos. Editora UNESP.

Nava, S., Beati, L., Labruna, M.B., Cáceres, A.G., Mangold, A.J., Guglielmone, A.A., 2014. Reassessment of
the taxonomic status of *Amblyomma cajennense* (Fabricius, 1787) with the description of three new
species, *Amblyomma tonelliae* n. sp., *Amblyomma interandinum* n. sp. and *Amblyomma patinoi* n. sp.,
and reinstatement of *Amblyomma mixtum* Koch, 1844, and *Amblyomma sculptum* Berlese, 1888 (Ixodida:
Ixodidae). Ticks and Tick-borne Diseases 5, 252–276. https://doi.org/10.1016/j.ttbdis.2013.11.004

Obenchain, F.D., 1974. Neurosecretory system of the American dog tick, Dermacentor variabilis (Acari:
Ixodidae). I. Diversity of cell types. J Morphol 142, 433–445. https://doi.org/10.1002/jmor.1051420406

Oliver, J.H., 1989. Biology and Systematics of Ticks (Acari:Ixodida). Annual Review of Ecology and
Systematics 20, 397–430.

Oliver, J.H., 1982. Tick Reproduction: Sperm Development and Cytogenetics, in: Physiology of Ticks.
Elsevier, pp. 245–275. https://doi.org/10.1016/B978-0-08-024937-7.50013-5

Oliver, J.H., Brinton, L.P., 1972. Cytogenetics of ticks (Acari: Ixodoidea). 7. Spermatogenesis in the Pacific
Coast tick, dermacentor occidentalis Marx (Ixodidae). J Parasitol 58, 365–379.

Oliver, J.H., Jr., 1974. Symposium on Reproduction of Arthropods of Medical and Veterinary Importance:
IV. Reproduction in ticks (Ixodoidea)1. Journal of Medical Entomology 11, 26–34.
https://doi.org/10.1093/jmedent/11.1.26

Onofrio, V.C., Barros-Battesti, D.M., Marques, S., Faccini, J.L.H., Labruna, M.B., Beati, L., Guglielmone,
A.A., 2008. Redescription of Amblyomma varium Koch, 1844 (Acari: Ixodidae) based on light and scanning
electron microscopy. Syst Parasitol 69, 137–144. https://doi.org/10.1007/s11230-007-9128-0

Patton, T.G., Dietrich, G., Brandt, K., Dolan, M.C., Piesman, J., Gilmore, R.D., 2012. Saliva, Salivary Gland,
and Hemolymph Collection from Ixodes scapularis Ticks. J Vis Exp 3894. https://doi.org/10.3791/3894

Pegram, R., Indar, L., Eddi, C., George, J., 2004. The Caribbean *Amblyomma* Program: Some Ecologic
Factors Affecting Its Success. Annals of the New York Academy of Sciences 1026, 302–311.
https://doi.org/10.1196/annals.1307.056

Prine, K.C., Hodges, A.C., 2012. Tropical Bont Tick Amblyomma variegatum Fabricius (Arachnida: Acari:
Ixodidae): EENY-518/IN934, 6/2012. EDIS 2012. https://doi.org/10.32473/edis-in934-2012

Reuben Kaufman, W., 2007. Gluttony and sex in female ixodid ticks: How do they compare to other
blood-sucking arthropods? Journal of Insect Physiology 53, 264–273.
https://doi.org/10.1016/j.jinsphys.2006.10.004

Ribeiro, J.M.C., Zeidner, N.S., Ledin, K., Dolan, M.C., Mather, T.N., 2004. How much pilocarpine
contaminates pilocarpine-induced tick saliva? Med Vet Entomol 18, 20–24.
https://doi.org/10.1111/j.0269-283X.2003.0469.x

Rispe, C., Hervet, C., de la Cotte, N., Daveu, R., Labadie, K., Noel, B., Aury, J.-M., Thany, S., Taillebois, E.,
Cartereau, A., Le Mauff, A., Charvet, C.L., Auger, C., Courtot, E., Neveu, C., Plantard, O., 2022.
Transcriptome of the synganglion in the tick Ixodes ricinus and evolution of the cys-loop ligand-gated ion

channel family in ticks. BMC Genomics 23, 463. https://doi.org/10.1186/s12864-022-08669-4

Saito, K.C., Bechara, G.H., Nunes, É.T., de Oliveira, P.R., Denardi, S.E., Mathias, M.I.C., 2005.
Morphological, histological, and ultrastructural studies of the ovary of the cattle-tick *Boophilus microplus*
(Canestrini, 1887) (Acari: Ixodidae). Veterinary Parasitology 129, 299–311.
https://doi.org/10.1016/j.vetpar.2004.09.020

Sampieri, Bruno Rodrigues, Calligaris, I.B., da Silva Matos, R., Páez, F.A.R., Bueno, O.C., Camargo-Mathias,
M.I., 2016. Comparative analysis of spermatids of Rhipicephalus sanguineus sensu lato (Ixodidae) and
Ornithodoros rostratus ticks (Argasidae): morphophysiology aimed at systematics. Parasitol Res 115,
735–743. https://doi.org/10.1007/s00436-015-4797-0

Sampieri, B.R., Labruna, M.B., Bueno, O.C., Camargo-Mathias, M.I., 2014. Dynamics of cell and tissue
genesis in the male reproductive system of ticks (Acari: Ixodidae) Amblyomma cajennese (Fabricius,
1787) and Amblyomma aureolatum (Pallas, 1772): a comparative analysis. Parasitol Res 113, 1511–1519.
https://doi.org/10.1007/s00436-014-3795-y

Sampieri, B.R., Moreira, J.C.S., Páez, F.A.R., Camargo-Mathias, M.I., 2016. Comparative morphology of the
reproductive system and germ cells of Amblyomma ticks (Acari: Ixodidae): A contribution to Ixodidae
systematics. J Microsc Ultrastruct 4, 95–107. https://doi.org/10.1016/j.jmau.2015.11.003

Sanches, G.S., Araujo, A.M., Martins, T.F., Bechara, G.H., Labruna, M.B., Camargo-Mathias, M.I., 2012.
Morphological records of oocyte maturation in the parthenogenetic tick *Amblyomma rotundatum* Koch,
1844 (Acari: Ixodidae). Ticks and Tick-borne Diseases 3, 59–64.
https://doi.org/10.1016/j.ttbdis.2011.11.001

Sasso Porto, D., Melo, G.A.R., Almeida, E.A.B., 2016. Clearing and dissecting insects for internal skeletal
morphological research with particular reference to bees. Revista Brasileira de Entomologia 60, 109–113.
https://doi.org/10.1016/j.rbe.2015.11.007

Šimo, L., Kazimirova, M., Richardson, J., Bonnet, S.I., 2017. The Essential Role of Tick Salivary Glands and
Saliva in Tick Feeding and Pathogen Transmission. Frontiers in Cellular and Infection Microbiology 7.

Šimo, L., Park, Y., 2014. Neuropeptidergic control of the hindgut in the black-legged tick Ixodes scapularis.
Int J Parasitol 44, 819–826. https://doi.org/10.1016/j.ijpara.2014.06.007

Smit, A., Mulandane, F., Labuschagne, M., Wójick, S.H., Malabwa, C., Sili, G., Mandara, S., Dlamkile, Z.,
Ackermann, R., Rose Vineer, H., Stoltsz, W.H., Huber, K., Horak, I.G., Morar-Leather, D., Makepeace, B.L.,
Neves, L., 2024. Intra- and interspecific variation of Amblyomma ticks from southern Africa. Parasites &
Vectors 17, 364. https://doi.org/10.1186/s13071-024-06394-3

Sonenshine, D.E., Simo, L., 2021. Biology and Molecular Biology of Ixodes scapularis, in: Lyme Disease and
Relapsing Fever Spirochetes: Genomics, Molecular Biology, Host Interactions and Disease Pathogenesis.
Caister Academic Press. https://doi.org/10.21775/9781913652616.12

Sonenshine, E. by D.E., Roe, R.M. (Eds.), 2013. Biology of Ticks Volume 2, Second Edition, New to this
Edition:, Second Edition, New to this Edition: ed. Oxford University Press, Oxford, New York.

Starck, J.M., Mehnert, L., Biging, A., Bjarsch, J., Franz-Guess, S., Kleeberger, D., Hörnig, M., 2018.
Morphological responses to feeding in ticks (Ixodes ricinus). Zoological Letters 4, 20.
https://doi.org/10.1186/s40851-018-0104-0

Szlendak, E., Oliver Jr., J.H., 1992. Anatomy of synganglia, including their neurosecretory regions, in

unfed, virgin female Ixodes scapularis say (Acari: Ixodidae). Journal of Morphology 213, 349–364.
https://doi.org/10.1002/jmor.1052130308

Tan, A.W.L., Francischetti, I.M.B., Slovak, M., Manjunatha, K.R., Ribeiro, J.M.C., 2015. Sexual differences
in the sialomes of the zebra tick, Rhipicephalus pulchellus. J Proteomics 117, 120–144.
https://doi.org/10.1016/j.jprot.2014.12.014

Tidwell, J.P., Treviño, D.E., Thomas, D.B., Mitchell, R.D., Heerman, M.C., Pérez de León, A., Lohmeyer,
K.H., 2021. Pictorial dissection guide and internal anatomy of the cattle tick, Rhipicephalus (Boophilus)
microplus (Canestrini). Ticks Tick Borne Dis 12, 101685. https://doi.org/10.1016/j.ttbdis.2021.101685

Uilenberg, G., 1983. Heartwater (Cowdria ruminantium infection): current status. Adv Vet Sci Comp Med
27, 427–480.

University, © Stanford, Stanford, Complaints, C. 94305 C., n.d. Comparing Different Disinfectants –
Stanford Environmental Health & Safety. URL https://ehs.stanford.edu/reference/comparing-different-
disinfectants (accessed 4.7.25).

Vega, R., Diaz, G., Galan, M., Fernández, C., 2012. ANATOMY AND HISTOLOGY OF THE FEMALE
REPRODUCTIVE SYSTEM OF Boophilus microplus (ACARI: IXODIDAE).

Voltzit, O.V., 2007. A review of neotropical Amblyomma species (Acari: Ixodidae).

Walker, A., Bouattour, A., Camicas, J., Estrada-Peña, A., Horak, I., Latif, A., Pegram, R., Preston, P., 2003.
Ticks of domestic animals in Africa: a guide to identification of species.

Wang, Y., Zhang, H., Luo, L., Zhou, Y., Cao, J., Xuan, X., Suzuki, H., Zhou, J., 2021. ATG5 is instrumental in
the transition from autophagy to apoptosis during the degeneration of tick salivary glands. PLOS
Neglected Tropical Diseases 15, e0009074. https://doi.org/10.1371/journal.pntd.0009074

Weiss, B.L., Stepczynski, J.M., Wong, P., Kaufman, W.R., 2002. Identification and characterization of genes
differentially expressed in the testis/vas deferens of the fed male tick, Amblyomma hebraeum. Insect
Biochemistry and Molecular Biology 32, 785–793. https://doi.org/10.1016/S0965-1748(01)00161-8

Woolley, T.A., 1972. Scanning Electron Microscopy of the Respiratory Apparatus of Ticks. Transactions of
the American Microscopical Society 91, 348. https://doi.org/10.2307/3224879

Yang, X.L., Yu, Z.J., Gao, Z.H., Yang, X.H., Liu, J.Z., 2014. Morphological characteristics and developmental
changes of the ovary in the tick Haemaphysalis longicornis Neumann. Medical Vet Entomology 28, 217–
221. https://doi.org/10.1111/mve.12035

Zacaï, A., Vannier, J., Lerosey-Aubril, R., 2016. Reconstructing the diet of a 505-million-year-old
arthropod: Sidneyia inexpectans from the Burgess Shale fauna. Arthropod Structure & Development,
Special Issue: Fossils as Living Beings 45, 200–220. https://doi.org/10.1016/j.asd.2015.09.003

A

Claw

Legs

Pedipalps

Hypostome

Basis capituli

Camerostome

Porose areas

Eyes

Scutum

Alloscutum

Festoons

Idiosoma /
Podosoma-
Opisthosoma

B

Claw

Hemolymph

Capitulum /
Gnathosoma

Eyes

Conscutum

Scutal ornementation

Festoons

Idiosoma /
Podosoma -
Opisthosoma

Figure 1.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure 9.

Figure 10.