Systematic Metaproteomics Mapping Reveals Functional and Ecological

Landscapes of Ex Vivo Human Gut Microbiota Responses to Therapeutic Drugs

Leyuan Li[1,2,*], Caitlin M.A. Simopoulos[2], Janice Mayne[2], Zhibin Ning[2], Xin Zhang[1], Mona Hamada[2],

James Butcher[2], Joeselle M. Serrana[2,3], Luman Wang[4], Kai Cheng[2,5], Hongye Qin[2], Krystal

Walker[2], Xu Zhang[2], Alain Stintzi[2], Daniel Figeys[2,5,6,7,*]

1 State Key Laboratory of Medical Proteomics, Beijing Proteome Research Center, National Center for

Protein Sciences (Beijing), 102206 Beijing, China.

2 School of Pharmaceutical Sciences, Ottawa Institute of Systems Biology and Department of

Biochemistry, Microbiology and Immunology, Faculty of Medicine, University of Ottawa, Ottawa, ON

K1H 8M5, Canada.

3 Department of Environmental Science (ACES), and the Stockholm University Center for Circular and

Sustainable Systems (SUCCeSS), Stockholm University, 106 91 Stockholm, Sweden.

4 Department of Health Informatics and Management, School of Health Humanities, Peking University,

Beijing 100191, China

5 Quadram Institute Bioscience, Norwich Research Park, Norwich, Norfolk, UK.

6 University of East Anglia, Norwich, Norfolk, UK.

7 Lead contact (Requests for further information and resources should be directed to and will be fulfilled

by the lead contact).

* Correspondence: L.L: lileyuan@ncpsb.org.cn, D.F.: Daniel.Figeys@quadram.ac.uk

Abstract

Therapeutic compounds exert impacts on gut microbiota; however, how they affect the

community functional ecology, especially as reflected at the protein level, remains largely

unexplored. In this study, we systematically map metaproteomic responses of *ex vivo* human gut

microbiota to 312 compounds, generating 4.6 million microbial protein responses, available as an

interactive resource (https://shiny.imetalab.ca/MPR_Viz/). Protein-level analyses identify

significant metaproteomic shifts induced by 47 compounds, with neuropharmaceuticals as the

sole drug class significantly enriched among these hits. Further analyses on the community level

reveal a tri-stability pattern in microbial composition and the emergence of three distinct functional

states, based on a functional beta-diversity metric. Notably, neuropharmaceuticals cause

particularly strong effects on the microbiomes, lowering the proteome-level functional redundancy

and raising the level of antimicrobial resistance proteins, ultimately pushing the microbiome into

an alternative functional state. Preliminary validation suggests that enhancing functional

redundancy may contribute to maintaining microbiota resilience against neuropharmaceutical-induced antimicrobial resistance. Overall, this work establishes a comprehensive view of how drugs influence gut microbiome function and ecology at the protein level, proposes a landscape-based framework for interpreting community resilience, and highlights the need to consider protein-level and ecological responses in the evaluation of therapeutic interventions.

INTRODUCTION

The development of human-targeted drugs typically focuses on mechanisms of drug interactions with human protein targets. However, it is often overlooked that therapeutic compounds can have profound impacts on our gut microbiotas, known as off-target effects[1]. Studies have revealed the effects of commonly prescribed drugs on single gut bacterial strains[2], microbiome composition[3], function[3,4] and drug metabolism[5,6]. A more recent study has exemplified these findings by demonstrating that non-antibiotic drugs inhibited *Escherichia coli in vitro* and that the microbial functional pathways they targeted were from those targeted by traditional antibiotics[7]. However, it's important to note that although *E. coli* serves as a frequently investigated model organism, our gut ecosystem is primarily populated by obligate anaerobes belonging to the Bacillota and Bacteroidota phyla, drug effects on which remain largely unexplored. Additionally, ecological interactions between gut microbial species can lead to more complicated drug responses compared to responses of single species. A microbial community can obtain resistance against drug perturbation through exposure protection[8] and metabolic cooperation[9], or conversely, drug-disrupted cross-feeding networks may sensitize the drug response of some microbial species that are otherwise unresponsive to the drug[10,11]. A recent study systematically investigated such community-level behaviors by analyzing species abundance dynamics in a 32-species synthetic community subjected to 30 different drug treatments, and found that 26% of the tested cases exhibited community-level phenomena such as cross-protection and cross-sensitization[12]. Indeed, drugs response involves the complexity of the ecosystem functioning that goes beyond single species survival[13] or overall compositional responses. In other words, a comprehensive analysis of microbiome drug responses which is observed from a functional omics scope will offer a provide deeper insights of the microbial systems ecology.

The drug modes of action that affect functional microbiome ecosystems vary, but they often involve interactions with key microbial proteins or enzymes. During pharmaceutical drug

development, many drugs are developed solely based on their effects through interactions with
specific human proteins, without considering the fact that species within the human microbiota
may contain structurally similar proteins and may thus also be influenced by these drugs. For
example, the serotonin transporter (SERT, encoded by *SLC6A4* in humans), responsible for
serotonin uptake in humans, is commonly targeted for pharmaceutical inhibition by neurological
drugs such as antidepressants. Interestingly, SERT belongs to the neurotransmitter:sodium
symporters (NSS) family, which is conserved from bacteria to humans[14]. Its bacterial homolog,
Leucine transporter (LeuT), has been shown to contain a NSS binding site[14], and has actually
been used as a model to predict the antidepressant drug-binding specificity to SERT[15]. Such
phenomena suggest that comprehensive observation of drug effects on the metaproteome level –
the frontline of microbiota-drug interaction – could offer invaluable insights for in-depth
exploration of phenotypic microbiome responses on the functional and ecological levels.

Moreover, since therapeutic drugs span a broad array of chemical structures, they are expected
to induce a diverse spectrum of microbial ecosystem responses of the gut microbiota. From the
scope of systems ecology, a key question arises: are these responses widely distributed across
the functional landscape of the microbiome (a map that connects all possible combinations of
microbial functional status)? Or do they cluster around distinct stable functional states, as
researchers have suggested regarding community structure[16,17]? Understanding such microbiome
functional landscape and mapping drug responses onto it holds significant promise for identifying
personalized drug treatments. In microbial systems ecology, functional redundancy (FR) refers to
the presence of multiple species that can perform similar functions[18]. It is widely believed that FR
is positively related to community resilience[19]. This raises the question of whether FR allows the
microbiome to maintain resilience by occupying stable functional states within the functional
landscape.

To comprehensively map human gut microbiome's drug response landscape, robust experimental
approaches are fundamentally required to capture a broad range of microbiome functional states
within the landscape under diverse drug treatments. *In vitro* microbiome culturing has emerged as
a potent tool for uncovering microbial responses to drugs and identify means to manipulate these
responses from interference from host factors. We've previously developed the RapidAIM
platform, which maintains the viability and functional individuality of *ex vivo* human gut microbiota
and robustly captures microbiome drug responses using metaproteomics[4] - a technique for the
large-scale identification and quantification of proteins within a microbiome, providing insights into
the functional state and activities of the microbial community. We have continued to scale up the

RapidAIM workflow, including live microbiota biobanking[20], cost-effective metaproteomic TMT-

labeling[21], and automation[22]. As a result, we have established the RapidAIM 2.0[22] platform for

high-throughput metaproteomic analysis of *in vitro* microbiota cultures, which was applied in this

research to reveal the functional landscape of microbiota drug responses.

In this study, we performed high-throughput assays of 312 commonly prescribed therapeutic

compounds against different individual gut microbiotas using the RapidAIM 2.0 platform, resulting

in nearly 2,000 microbiota cultures and tandem mass tag (TMT)-labelled metaproteomic

screenings. Subsequent in-depth label-free metaproteomics analysis of 110 compounds that

passed the initial screening further detailed over 4.6 million bacterial protein-drug responses,

provided as a resource through an interactive Shiny app. We analyzed the functional and

ecological landscapes of gut microbiota responses to therapeutic drugs across three hierarchical

levels: protein-level, revealing metaproteomic shifts in 47 of the tested compounds with

neuropharmaceuticals significantly enriched; taxonomic composition-level, exhibiting a

compositional tri-stability landscape; and systems ecological-level, where our recently developed

PhyloFunc beta-diveristy metric[23] identified three distinct functional community state types in

response to different compounds, demonstrating remarkable systems-level ecological

consistency in drug responses across individuals. Notably, specific human-targeted compounds,

particularly neuropharmaceuticals, stimulated the expression of microbial antibiotic resistance

proteins (ARP), while reducing community-level FR. Building on the functional energy landscape

hypothesis, we were able to reduce the neuropharmaceutical-induced ARM increase by

introducing a prebiotic into the culture to counteract the FR decrease. These findings underscore

the critical need to map the metaproteomic drug response landscape, as it provides a

foundational framework for identifying key leverage points to tailor and optimize personalized

drug response modulation.

RESULTS

The human gut metaproteome is altered by a diverse array of drugs

In this study, we cultured human gut microbiotas from healthy subjects and exposed them to a

collection of therapeutic compounds in our RapidAIM assay. The functional responses were

measured using high-throughput metaproteomics (Figure 1a). Briefly, the compounds were

selected through a multi-step process: we first included 141 drugs with reported impacts on the

microbiome based on our in-house database (Supplementary Data S1). Subsequently, we cross-

referenced this list with the top 200 most prescribed compounds in the US according to

literature[24], resulting in the addition of 119 compounds. To ensure comprehensive coverage of

drug classes and properties, we expanded the list further with 52 additional compounds by

addressing three additional questions: 1) according to the ATC classification system, are drug

classes being adequately presented or further expansion of drug representatives is needed? 2)

does each ATC class contain representative compounds of similar and different chemical

structures? 3) which drugs are stereoisomers and can be used to test the stereospecific

bioactivity and toxicity? Details of the resulting 312 compounds are listed in Supplementary Data

S1. Assessing the compounds using structure-based clustering suggests a good coverage of

drug diversity (Figure 1b).

Experiments were conducted using biobanked live human gut microbiotas from six distinct

individuals, with each microbiota treated with different compounds in separate wells of 96-well

plates, alongside positive and blank controls (2 mg/mL Kestose and the drug solvent DMSO),

resulting in a total of 1,920 microbiota cultures. TMT-11plex^{TM} was used for a first-pass screening

of compounds that potentially induced functional responses in the microbiomes. A total of 91,641

peptides (average of 10,025 ± 1,841 peptides per multiplexed TMT set) and 21,859

proteinGroups (average of 5,496 ± 519 proteinGroups per multiplexed TMT set) were identified

from the 192 multiplexed TMT sets of 1920 samples. Overall Principal Component Analysis (PCA)

based on proteinGroups showed both inter-individual differences in metaproteomics and the

alteration of metaproteomes by a subset of compounds (Figure 1c-d). Due to the presence of

inter-individual differences, we applied a data analysis pipeline to cluster drug responses by

average distances. Briefly, the distance matrix (measured by Pearson's correlation) between

metaproteomic proteome content networks (PCNs)[25] was first calculated separately for each

individual subject; subsequently, the distances between all compounds and controls were

averaged across all subjects. Finally, K-medoids analysis revealed five clusters of drug

responses (Figure 1e). We showed that three of the compound clusters (clusters 1, 4 and 5)

deviated from the controls (Figure 1f). Interestingly, there appeared to be no visible correlation

between the drug structure clusters in Figure 1B and the compound response clusters in Figure

1E. From the TMT-based compound response clustering, we show that across Anatomical

Therapeutic Chemical (ATC) classification system level 1 classes, the proportion of compounds

belonging to each K-medoids clusters are different, we therefore evaluated the enrichment of K-

medoids clusters 1, 4 and 5 compounds in each ATC level 1 class. The analysis revealed no

significant enrichment for any specific ATC level 1 class compared to the background of all

selected compounds. However, it is still notable that Class N (targeting nervous system) showed

a *p*-value of less than 0.1 (Figure 1g).

Figure 1 Screening for compounds influencing the gut metaproteome.

(a) Schematic representation of the experimental workflow. The study follows the RapidAIM 2.0

procedure[22]; fecal slurry samples of each human gut microbiota were cultured anaerobically with

the presence of a compound or a control substance (DMSO or kestose). Automated

metaproteomics sample processing was performed before samples are analyzed using LC-

MS/MS. This figure is adapted from our previously published work[22] under a Creative Commons

Attribution 4.0 International License (CC BY 4.0). (b) Selection, Anatomical Therapeutic Chemical

(ATC) classification and structure-based clustering of compounds in the compound library used in

this study. Colors of compound labels correspond to three of the clusters in Panel E: magenta –

Cluster 1, red – Cluster 4, purple – Cluster 5. Thick branches indicate clusters with high bootstrap

support (approximately unbiased (AU) p-value > 95%). (c)-(d) Principal component analysis (PCA)

of all samples showing metaproteome individuality as well as responses to compounds. (e)

Distance-based K-medoids clustering of all compounds based on the average of six individual

microbiome responses. Each point represents a compound or a control. Ellipses represent 95%

confidence of each cluster. (f) Distance of compounds to DMSO control samples within each

cluster identified in panel (d). The numbers of compound–control pairs were 96, 269, 373, 120,

and 99 for clusters 1, 2, 3, 4, and 5, respectively. Mean values and standard deviations were

calculated across $N = 6$ individual microbiome samples. Letters a, b, c, and d indicate levels of

significance determined by ANOVA with Tukey's post hoc test. The center line of box represents

the median; the box limits indicate the upper and lower quartiles (Q3 and Q1); the whiskers

extend to the most extreme values within 1.5× the interquartile range (IQR) from the quartiles. (g)

Enrichment of various ATC level 1 drug classes in Clusters 1, 4, and 5 compounds

(corresponding to the clusters identified in panel (d)), among all selected compounds. A one-

sided Fisher's exact test was applied to evaluate whether a given drug class was over-

represented in the test set compared to the background, shown are raw *p*-values uncorrected for

multiple hypothesis testing.

Comprehensive profiling of metaproteomics drug responses reveals alterations in
proteins and functional pathways

A total of 110 compounds from the K-medoids clusters 1, 4 and 5 (Figure 1e and Supplementary

Data S1) were selected for further in-depth label-free quantification (LFQ) – based

metaproteomics analysis. We used deep metagenomic sequencing of each individual microbiome

sample to construct a sample-specific database for the label-free database search following the

Metapro-IQ workflow[26] (Supplementary Figure S1a). A total of 9,326,339 LC-MS/MS spectra were

identified as 195,686 unique peptides, corresponding to 36,075 unique proteinGroups. An

average of 9,871 ± 3,679 peptides per sample was identified by MS/MS; in addition,

9,994 ± 2,052 proteinGroups were quantified per sample with match-between-runs (MBR) option

enabled during protein quantification. The MBR approach allows transfer of peptide identifications

across runs based on retention time and mass-to-charge (m/z) alignment, resulting in a higher

number of proteinGroups being quantified than peptides directly identified in individual samples.

Label-free quantification was then performed using directLFQ[27] followed by data harmonization

using HarmonizR[28]. The processed dataset thus includes over 4.6 million bacterial protein-

compound responses. A series of quality-control assessment suggested high reproducibility and

robustness of the whole process (Supplementary Figure S1b-d).

Given the extensive spectrum of metaproteomic responses across various functional pathways

and taxonomic units, microbial drug responses provide valuable insights into resistance

mechanisms, metabolic adaptations, and ecological interactions under therapeutic interventions.

To facilitate data exploration and hypothesis generation in future drug-microbiome research, we

have developed an interactive Shiny app as a resource for exploring functional and taxonomic

responses to these 110 compounds, available at https://shiny.imetalab.ca/MPR_Viz/. This app

allows readers to select specific ATC drug classes and explore the functional and taxonomic

responses of individual compounds within these classes. In more details, the features that can be

visualized involve functional annotations such as NOG categories and NOG accessions

(annotated by eggNOG mapper), KEGG KO annotations (from the KEGG database), as well as

protein abundances across different taxonomic units. By providing this level of detail, the app

facilitates a comprehensive exploration of the drug-microbiome interactions at both the functional

and taxonomic levels. Readers can visualize data and download the figures, making it a highly

informative tool for further exploration and hypothesis generation about microbiome responses to

drugs. Here, we show an example of the Shiny app interface with an analysis result of showing

several significant increases in the NOG annotation of proteins associated with efflux

transmembrane transporter activity following independent treatments of several different drug

from the ATC Level 1 class N (Supplementary Figure S2). This suggests that exposure to these

drugs may have stimulated an increase in drug resistance activity, likely through the upregulation

of efflux transporters.

We next examined the protein-level responses by performing PCA on the label-free quantified

proteinGroup intensities using the nonlinear Iterative Partial Least Squares (NIPALS) algorithm[29].

As shown in Figure 2A, the PCA plot reveals a densely clustered distribution due to the large

number of samples. Nonetheless, a global shift from the control group is evident, indicating

widespread compound-induced alterations in the metaproteomic profiles of the selected

compounds. We further examined the PCA within each distinct ATC class and at various ATC

levels (Supplementary Figures S3-S4). While only a subset of compounds in each of ATC level-1

classes A (alimentary tract and metabolism), B (blood and blood forming organs), D

(dermatologicals), L (antineoplastic and immunomodulating agents), and M (musculo-skeletal

system) showed separation from the controls, all compounds in ATC level-1 class P (antiparasitic

products, insecticides and repellents) separated from the controls on PC 1 which explains 62.7%

of the variance. Classes C (cardiovascular system), G (genito urinary system and sex hormones),

J (antiinfectives for systemic use) and N (nervous system) contained a larger number of

compounds. We performed further explorations of PCAs on the second ATC level and presented

classes that contained discriminative compounds (Supplementary Figures S3). On ATC level-2,

we show that classes G03 (sex hormones and modulators of the genital system), J01

(antibacterials for systemic use), N04 (anti-parkinson drugs), N05 (psycholeptics) and N06

(psychoanaleptics) classes contained greater than three compounds discriminated from the

controls. Among these, the ATC level-4 code N06AB (selective serotonin reuptake inhibitors)

comprised six compounds with noticeable effects (Supplementary Figures S4). By further

examining PCA by individual compounds, we observed a total of 47 compounds that are

discriminative from the control in all individual microbiomes, denoted as Metaproteomics-

response positive (MPR+) drugs (Supplementary Data S2 and Figure 2b). Among which, 39% are

Class N compounds, and 15% are Class C compounds. Enrichment analysis of these 47

compounds against the background of all 312 drugs showed a significant enrichment of Class N

drugs (Figure 2c), suggesting that drugs targeting the nervous system have the most frequent off-

target effects on gut microbes.

We were curious about which human proteins are targeted by these MPR+ compounds and

whether these targets could have bacterial homologs. Therefore, by searching MPR+ compounds

against the molecular drug target database by Santos et al.[30], we extracted a list of 50 human

protein targets (Supplementary Data S2). KEGG enrichment analysis shows that these proteins

are most significantly enriched in neuroactive ligand-receptor interaction and calcium signaling

pathways (Supplementary Figure S5a-b), both are targets of Class N drugs. Using the HMMER

algorithm[31] to map the drug target Pfams through our metaproteomic sample-specific database,

we discovered 2,528 microbial proteins that are homologous to these human targets. EggNOG

annotation was able to map 2,115 of these proteins to reference taxa and functions which were

mainly originated from the Clostridia class, with signal transduction mechanisms being the most

significantly enriched functional category (Figure 2d-e, and Supplementary Data S3). Notably,

270 bacterial proteins homologous to human neurotransmitter sodium symporters, the targets of

selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors

(SNRIs), were identified within the sodium:neurotransmitter symporter or sodium:solute symporter

family.

Figure 2. Discovery of MPR+ compounds and microbial homologs of their human targets

(a) Overall PCA of the label-free dataset. (b) Principal component analysis (PCA) of

metaproteomics response-positive (MPR+) compounds and control samples, grouped by ATC

level-1 classification. For (A)-(B), the first three components of Principal Component Analysis

(PCA) are shown. All six individual gut microbiome are shown as labeled by different colors. Gray

dashed ellipse denotes 95% confidence interval of each group (red: drugs, black: DMSO and

blue: kestose). (c) Enrichment analysis of ATC level 1 class in MPR+ compounds against all 312

selected compounds. A one-sided Fisher's exact test was applied to evaluate whether a given

drug class was over-represented in the test set compared to the background, shown are raw p-

values uncorrected for multiple hypothesis testing. (d) Discovery of 2,528 microbial proteins

homologous to human drug targets in the individual gut microbiomes' metaproteomics database

using the HMMER algorithm. The homologous proteins were annotated by eggNOG mapper, and

their taxon-specific annotations were summarized and visualized in a Circos plot. (e) COG

285 functional enrichment of the identified homologs was then performed using iMetaShiny, in which a
286 one-sided hypergeometric test was used.

Taxonomic responses to drugs exhibit a compositional tri-stability landscape

Taxonomic responses, as it takes account into cumulative abundances of taxon-specific peptides,
provides an alternative view of microbiomes community responses that is different from protein-
level PCA which takes into consideration proteins as features. The compounds selected for LFQ
analysis span a wide spectrum of drug structure and ATC classes. Our initial intuition was that the
microbiome responses to different compounds can be highly diverse, resulting in a diverse
combination of microbiome composition. The equal-volume processing protocol (see Methods >
RapidAIM 2.0 assay) ensures direct comparability of protein-based taxonomic abundances
between samples. To examine the microbiome responses, we selected the microbial genera that
showed at least one significant drug response across all six microbiomes, and visualized their
abundance changes in a genus-level heatmap (Figure 3A). Despite observing highly diverse
patterns in protein-level analysis, we identified an intriguing phenomenon in the taxonomic
(genus-level) response heatmap: there's a coherent transition between three clusters of
responses (Figure 3a). Among which, one cluster (Cluster 2) only included five genus-level
alterations, while the two other clusters (Clusters 1 and 3) exhibited more extensive changes in
44 genera in response to 63 compounds. By color-coding different genera in each cluster with
blue, orange, and green tones, the overall transition becomes apparent. This can be more easily
observed by visualizing the drug response spectrum of each individual microbiome through
stacked column bar plots (Figure 3b).

This provides evidence for the theory of multiple stability points in the microbiome[16,32], our work
marks the first observation of a tri-stability landscape of individual microbiome compositions
within the drug response space. To explain the phenomenon of steady states in the functional
response of the microbiome to drugs, we used the generalized Lotka-Volterra (gLV) to describe
the response dynamics (see Methods). In this model, we classified the genera observed in the
experiment into three distinct clusters based on co-abundance patterns, assuming that
interactions within each cluster are negligible. By adjusting the model parameters to simulate the
effect of drug stimulation, we can reproduce the switches from one stable state to another (Figure
3c), mirroring the actual microbiome responses observed in Figure 3b.

Figure 3. Taxonomic responses reveal multi-stability of individual gut microbiomes driven by drug stimulation

(a) Heatmap and hierarchical clustering showing log$_2$ fold-change of overall protein biomass specific to different genera. Asterisks indicate significance levels from a two-sided Wilcoxon signed-rank test, with $p < 0.05$ and fold-change >1.5. For each genus-compound pair, the test

was performed to evaluate whether the distribution of values significantly deviated from the

reference value of 1, under the null hypothesis that the median of the differences (Value – 1) is

zero. A continuity correction was applied to the test statistic to improve the normal approximation.

(b) Relative abundance of genus-specific proteins presented in stacked column bar plots. The

sample order agrees with panel A, and different genera are also colored according to the

hierarchical clustering in panel A. (c) gLV competition model considering between-cluster

interactions simulates the switches between three different stable states under drug perturbation.

The time axis in panel C is conceptual and does not correspond to the actual order of drug

treatments depicted in panels A and B.

Community state analysis reveals universal drug-specific microbial ecological responses

Examining protein and taxonomic compositions separately fails to capture the full ecological

landscape from a systems perspective. Therefore, it is crucial to adopt a more ecologically

relevant assessment of microbiome responses. We have previously developed a tool named

PhyloFunc, which integrates phylogenetic and functional information to assess microbial

community dissimilarities (i.e. beta diversity) on the ecosystem level[23]. PhyloFunc analysis was

performed on each individual's microbiome including all LFQ samples of drug treatment and

control conditions. The PhyloFunc distance results from different individuals were combined and

visualized using Principal Coordinates Analysis (PCoA). K-means clustering was then applied,

identifying three distinct community-state types (CSTs) (Figure 4a), with the optimal number of

clusters determined as three based on the Gap statistic (Figure 4b). Based on these three

clusters, we used a Sankey diagram to visualize the distribution of drug names and their sources

across different CSTs (Figure 4c). Notably, all negative control (DMSO) individual microbiomes

clustered into CST-3, while 41 of the compounds caused at least 80% of the tested individual

microbiomes to shift into CST-1 and CST-2. These compounds include Amiodarone, Amlodipine,

Amoxapine, Anacardic, Asenapine, Atomoxetine, Berberine, Canagliflozin, Carvedilol, Clemastine,

Clomiphene, Cyclobenzaprine, Diacerein, Dienestrol, Docusate, Doxepin, Drospirenone,

Duloxetine, Entacapone, Escitalopram, Ethoproprazine, Fluoxetine, Fluphenazine, Fluvoxamine,

Ginsenoside, Hydroxychloroquine, Iloperidone, Linezolid, Loperamide, Mefloquine, Metronidazole,

Oxybutynin, Paroxetine, Ridinilazole, Sertraline, Simvastatin, Tamoxifen, Ticagrelor, Tinidazole,

Tolcapone, and Verapamil.

To further illustrate that PhyloFunc's inter-individual variability is not affected by individual-specific

differences in protein composition, we calculated the Mean and standard deviation (SD) of

PhyloFunc distances across individuals (N = 6) (Figure 4d-4e). We can clearly observe the

overall differences in heatmap intensity between the Mean and SD matrices when using the same

color gradient scale. To further quantify these differences, we applied a two-sided Wilcoxon

signed-rank test between the Mean and SD, which demonstrated a highly significant difference

(p-value < 2.2e-16, N = 12,321 compound-compound interactions). Such systems-level ecological

consistency in drug responses across individuals enabled us to summarize the PhyloFunc results

into an averaged distance matrix (Figure 4f), where we identified a distinct cluster of 13 drugs, 8

of which belonged to the N-class category

Figure 4. PhyloFunc analysis reveals impact of drugs on microbial community states.

(a) Principal Coordinates Analysis (PCoA) of PhyloFunc distances, clustering microbiome

responses into three distinct Community-State Types (CSTs). (b) Determination of the optimal

number of CST clusters using the Gap statistic (, represented by centerlines). Error bars

represent the standard error (   ) values across   Monte-Carlo reference simulations at

each   . (c) Sankey diagram illustrating the distribution of drugs across CSTs, showing the

association of compound names and their ATC level-1 classes with different CSTs. (d) Heatmap

of the Mean PhyloFunc distance matrix, representing the averaged PhyloFunc distance for each

373   compound-compound pair across individuals (N = 6). (e) Heatmap of the Standard Deviation (SD)

374   of PhyloFunc distances (N = 6). (f) PCoA based on averaged PhyloFunc distance matrix across

375   all individuals.

A subset of neuropharmaceuticals exerted similar functional effects on the ex vivo human

gut microbiomes

As an example of specific interest in taxonomic and functional responses, we observed that a

subset of neurological drugs exerted nearly identical effects on the human gut microbiomes as

assessed by protein-level PCA, including seven drugs, namely amoxapine (ATC N06AA),

duloxetine (ATC N06AX), fluoxetine (ATC N06AB), fluphenazine (ATC N05AB), fluvoxamine

(ATC N06AB), paroxetine (ATC N06AB), and sertraline (ATC N06AB) (Figure 5a). We show that

within each individual microbiome, NIPALS distances between pairs of these compounds are not

significantly different compared to that between technical replicates of the DMSO and kestose

control groups. In contrast, this group of compounds showed significantly higher distances from

both the controls (Figures 5b and 5c). Interestingly, they did not belong to a same ATC level-3/4

class, while most of these drugs are antidepressants that have been reported to have a common

target of human SERT proteins, fluphenazine belongs to antipsychotics that does not target

SERT. We therefore considered these drugs as a group of potentially common mode of action on

the gut microbiome and visualized differentially expressed proteins of this drug group using

volcano plot (Figure 5d). In comparison to the DMSO control group, we observed 1,447

significantly decreased proteins and 357 significantly increased proteins. By annotating the

proteins to taxon (genus level) and function (COG category), we observed that the functional

responses showed strong taxonomic specificity, characterised by overall decreased growth of

commensals and enriched proteins of opportunistic pathogens in Pseudomonadota (Figure 5e-f).

Taxa having decreased functional protein classes are consistent with previous reports indicating

that these taxa are also suppressed[2,33].

Figure 5. Seven neurological drugs showing near identical metaproteomic effects

(a) NIPALS PCA showing discriminations between the seven selected neurological drugs and the

controls. (b) Between-sample distances were calculated based on NIPALS scores and the

resulted distances were compared between groups. NC, negative control (DMSO); PC, positive

control (kestose); CPD, seven selected compounds. Two-sided Wilcoxon test was applied, ****

indicates p < 0.0001. From left to right, the numbers of between-sample comparisons are N = 16,

18, 51, 119, 126, and 126, respectively. The center line of box represents the median; the box

limits indicate the upper and lower quartiles (Q3 and Q1); the whiskers extend to the most

extreme values within 1.5× the interquartile range (IQR) from the quartiles. (c) The NIPALS

score-based distance matrix was further visualized using heatmap and hierarchical clustering. (d)

Volcano plot visualizing log2 fold-change of proteins versus FDR-adjusted *p* values (-log$_{10}$) using

two-sided Wilcoxon rank sum test. Significantly differential features were determined by log$_2$ fold-

change threshold of 1.5 and FDR-adjusted *p* value of 0.05. Threshold curves[34] were defined

412   using a curvature of 1 as previously described. (e, f) Sankey plots showing taxon-specific

413   functional responses of the significantly decreased (e) and increased (f) proteins.

Antimicrobial resistance negatively correlates to functional redundancy

We have previously performed a proof-of-concept (POC) study of the RapidAIM assay using 43

compounds, and found that drugs not only cause changes in functional and taxonomic

composition, but also induce changes each individual microbiomes' levels of drug resistance

proteins[4]. We further wondered whether these changes in predicted drug resistance and

alteration of taxonomic composition had any relationship with a community's functional ecology

properties (Supplementary Figure S6a). Recently, we have developed an approach to quantify a

microbial community's proteome-level functional redundancy (FR_p) and its normalized version

relative to the community's taxonomic diversity, nFR_p[25] (Supplementary Figure S6b). FR_p

describes the ability of multiple taxonomically distinct organisms to contribute in similar ways to

an ecosystem, providing informative insights into microbial community ecology on the functional

level. Here, we first reanalyzed the RapidAIM POC dataset[4] and compared nFR_p levels with

community composition and levels of antimicrobial resistance proteins (ARP) (Supplementary

Figure S6c). We observed instances where increased prevalence of ARP occurred concurrently

with a decrease in nFR_p under rifaximin (Supplementary Figure S6D), berberine, flucytosine, and

metronidazole treatments. This case study roused our interest, and we were curious whether

such phenomenon would be reproduced in this expanded collection of drug-microbiome cultures.

Hence, we analyzed the LFQ dataset using the full CARD antimicrobial resistance database[35] to

redo database search and sum ARP abundance to estimate community-level drug resistance. We

observed a strong negative correlation between fold-change of nFR_p and ARP ratio (Pearson's

product-moment correlation coefficient r of -0.695 ± 0.154, Figure 6a). Interestingly, nFR_p

response and ARP ratio were both independent of taxonomic diversity (TD) responses (Figures

6b and 6c), suggesting that antimicrobial resistance corresponds to the functional ecology status

of a community, irrespective of taxonomic diversity. We also observed a mixture distribution on

the scatter plot, which becomes particularly evident when performing K-medoids clustering with

each individual microbiome dataset (Figure 6d). This suggests the presence of alternative

functional states of the microbiome, wherein the microbiome can transition from its original

functional ecological state of high   to an alternative state of low   when influenced by

specific groups of compounds. We demonstrated this concept by illustrating the energy

landscape of individual V45's microbiome using the Kernel Density Estimation (KDE) method

(Figure 6e). The upset plot demonstrates a high level of common presence of these compounds

across each individual microbiome's high   clusters (Figure 6f). We calculated the frequency of

compounds in the high   clusters by summing their occurrences across all individual datasets,

N05A (antipsychotics) and N06A (antidepressants) drug classes of ATC level 3 showed a high

frequency of appearance in the high   clusters compared to other N class drugs targeting the

neural system (Figures 6g-h).

Figure 6. ARP abundance negatively correlates with functional redundancy

(a) Scatter plot and Pearson's correlation between functional redundancy log2 fold-change and

ARP ratio. (b) Scatter plot and Pearson's correlation between taxonomic diversity log2 fold-

change and functional redundancy log2 fold-change. (c) Scatter plot and Pearson's correlation

between taxonomic diversity log2 fold-change and ARP ratio. In panels (a)-(c), box plots show

value distributions based on individual microbiome corresponding to each axis. Each box

includes N=115~117 compound-treated or control metaproteomes for each subject's sample. The

thicker black line in the box plot denotes median, the lower and upper hinges correspond to the

first and third quartiles, the box ranges from the 1.5□×□(interquartile range) below the lower

hinge to 1.5□×□IQR above the upper hinge, and whiskers represent the maximum and minimum

values, excluding outliers. (d) K-medoids clustering of ARP-FR response by individual. (e) Energy

landscape of ARP-nFR_p response by KDE method. (f) Set comparison of Cluster 2 compounds

across individuals, with bar heights representing the number of compounds in each intersection.

(g) Counts (represented by bar heights) of Cluster 1 & 2 compound frequencies in different

classes. (h) Tree map showing frequency of occurrence in Cluster 2. Bold capital letters indicate

ATC level 1 classes, and capital letters with numbers indicate ATC level 3 classes.

Enhanced functional redundancy mitigated drug-induced ARP

Considering the ARP-nFR_p drug response energy landscape shown in Figure 6e, and based on

the dynamic mechanisms of energy landscapes, we wondered if functional redundancy could be

leveraged to prevent drug-induced shifts of the microbiome's functional state from the high nFR_p-

low ARP attractor to the high ARP-low nFR_p attractor. Specifically, our hypothesis was that while

a microbiome was stimulated by an ARP-inducing drug (for example, cluster 2 drugs), adding

another compound that can increase a microbiome's functional redundancy may help combat the

functional redundancy decline and suppress ARP. To test this hypothesis, we designed a 2x2

factorial design as shown in Figure 7a.

We selected sertraline as our model for an ARP-inducing drug, as it showed the strongest effect

in reducing nFR_p and increasing ARP levels across all six individual microbiomes (Supplementary

Figure S7a). Sertraline was also shown to be pathogen-favoring in previous research[36].

Fructooligosaccharide (FOS) was selected as a nFR_p enhancing compound because it increased

nFR_p in a subset of previously tested individual microbiomes[4] (an example is shown in

Supplementary Figure S6c). Since the   response to FOS can vary between individuals, we

performed a pretest using three additional individual microbiomes and selected one microbiome

with the highest level of   increase in the presence of 10 mg/mL FOS (Figure 7b) and

performed the 2x2 factorial design experiment. The effects of the different combinations on

and ARP are measured using metaproteomics. Results show that sertraline alone

decreased nFR and increased ARP (first group of comparisons in Figure 7c and d). When

sertraline was combined with FOS, it did not decrease nFR nor increase ARP, as compared to

the microbiome treated by FOS alone (second group of comparisons in Figure 7c and d).

Furthermore, compared to the microbiome treated only with sertraline, those treated with both

sertraline and FOS showed significantly increased   and decreased ARP (third group of

comparison in Figure 7c and d). In addition, sertraline increased the abundance of

Enterobacterales, but this increase did not occur when sertraline was present alongside FOS

(Supplementary Figure S7b). These findings together suggest the possibility that manipulating

the microbiome to increase   may help counteract the increase in ARP and contribute to

preserving microbiome resilience against drug-induced negative disruptions. However, due to

inter-individual variability, future studies in larger populations are needed to assess the

generalizability of this strategy.

Figure 7. Functional redundancy acts as a potential target to combat drug-induced ARP

(a) Experimental design, sertraline was abbreviated as SER on the figure. (b) Pretest of individual gut microbiome responses to FOS show individualized nFR_p alterations. One single value is represented by each bar. (c) Between-sample comparisons of nFR_p and (d) ARP, two-sided t-test was performed.

DISCUSSION

In this study, we systematically profiled how 312 commonly prescribed and structurally diverse drugs and compounds perturb human gut microbiomes ex vivo using the RapidAIM 2.0 platform. Based on an initial TMT-based screening, 110 drugs that induced distinct microbiome perturbations were selected for in-depth label-free quantitative (LFQ) metaproteomic analysis across the microbiomes. The study was adequately powered for exploratory discovery, enabling the detection of key ecological patterns in drug–microbiome interactions. A central finding of this work is that gut microbiome responses to drugs exhibit a reproducible pattern at the ecological systems level, even amid substantial inter-individual variability in baseline microbiome profiles, and that these responses can potentially be modulated by leveraging functional redundancy as an ecological leverage point.

At the functional and taxonomic levels, we found that drugs targeting the nervous system (ATC level-1 class N), cardiovascular system (C), genitourinary system (G), and anti-infectives (J) induced the most pronounced metaproteomic shifts. Among these, nervous system drugs— including SSRIs and SNRIs—consistently elicited strong perturbations in microbial protein expression and community composition. This was consistent with previous findings that SSRI treatment alters gut microbiota composition and function in humans and animal models[37–39]. Our analysis suggests that these effects may involve unintended interactions between drugs and bacterial homologs of human protein targets. Specifically, we observed over 200 protein-coding genes from the sodium:neurotransmitter symporter or sodium:solute symporter family in the tested gut microbiomes. These bacterial transporters are homologous to human neurotransmitter sodium symporters (NSS), which are the targets of commonly used antidepressants such as SSRIs and SNRIs, including the human serotonin transporter (SERT)[30,33] responsible for serotonin reuptake. However, although these proteins were detected during the first search in our two-step metaproteomic workflow[26], they were not consistently quantified in the second search stage, likely reflecting their relatively low abundance and representing a limitation of current

metaproteomic approaches, which constrained our ability to quantify the responses of these
proteins to the neurological drugs. Another limitation of this study is that we did not directly
assess the interactions between drugs and bacterial homologs of human protein targets in the
human gut microbiomes. Establishing such molecular-level evidence will be critical for future
investigations to fully elucidate the mechanisms underlying drug–microbiome interactions.

We next explored the ecological responses at the taxonomic composition level. Counterintuitively,
the microbial community composition did not exhibit a broad continuum of distributions but
instead formed three distinct states, suggesting the presence of intrinsic multi-stability in gut
microbial communities. To further investigate the theoretical principles underlying this tri-stability,
we applied the generalized Lotka-Volterra (gLV) model to describe the response dynamics and
understand the stability of microbial community states. However, we acknowledge that our
current model is theoretical and may oversimplify the actual microbiome dynamics. Moreover,
while our findings highlight interesting patterns in taxonomic responses, taxonomic analyses
alone are insufficient to fully reveal the ecological response landscape of the microbiome, as they
do not adequately capture how microbial functions reorganize in response to drug
perturbations—especially given the extensive functional redundancy within microbiomes.

By contrast, metaproteomic data provide a valuable foundation for systems-level ecological
analysis, as they directly reflect the functional state of the community under perturbation. To
leverage this potential, we applied our previously developed *PhyloFunc* framework, which
integrates phylogenetic relationships and functional annotations to quantify ecological distances
between microbiome states. Unlike protein-level PCA approaches that rely on data harmonization
methods to minimize inter-individual protein variability, PhyloFunc captures ecosystem diversity
directly from the original protein group data without requiring data harmonization across
individuals. It reduces the impact of compositional differences among phylogenetically related
species, thereby offering a more ecologically meaningful perspective on functional community
responses. Our finding suggests that while metagenomics and metaproteomics have traditionally
revealed high inter-individual microbiome diversities in strains, genes, and proteins, a notable
level of consistency emerges at the functional ecosystem level. This suggests that drugs may
induce conserved shifts in the ecological function of gut microbiomes across individuals—an
insight that cannot be fully captured by taxonomic or protein-level analyses alone. Our findings
emphasize that integrating functional and taxonomic dimensions through an ecological framework
is essential for understanding drug–microbiome interactions.

Notably, in all different levels of response landscapes, we observed that several drugs targeting

the nervous system - amoxapine, duloxetine, fluoxetine, fluphenazine, fluvoxamine, paroxetine,

and sertraline —consistently induced strong perturbations in gut microbial proteins and

community ecology. These drug effects were associated with decreased commensal growth and

enrichment of pathogen-associated proteins, raising concerns about their long-term impact on gut

ecosystem health. The complexity of the microbiome makes it unlikely that off-target drug effects

can be mitigated by targeting individual species. Instead, our findings underscore the need to

consider the functional ecological landscape of the microbiome as a framework for understanding

and managing drug-microbiome interactions. Emerging from our analyses was the observation of

bimodal functional states within the microbiome drug response landscape: a high FR_p[25], low ARP

state, and a low FR_p, high ARP state. These patterns resemble ecosystem regime shifts

described in resilience and redundancy theory[40], wherein loss of functional redundancy may

lower the resilience of the microbial community, increasing susceptibility to dominance by

resistant strains. Notably, such non-linear dynamics are not readily captured by taxonomic or

single-protein analyses alone but require a systems-level ecological perspective. We show that

by using prebiotic FOS, the ecosystem can potentially buffer against the functional redundancy

loss in specific individual microbiomes and prevent the system from shifting to an undesirable

state dominated by resistant strains. However, further validation in larger and more diverse

cohorts will be essential to translate these insights into personalized strategies for maintaining gut

ecosystem health during pharmacological treatments.

Several limitations of this study should be acknowledged. First, as an ex vivo microbiota culture

model, the RapidAIM 2.0 platform cannot recapitulate host–microbiome interactions; rather, it

enables the investigation of microbiome-intrinsic responses to drugs, independent of host factors.

Second, as discussed earlier, this study did not directly investigate the molecular mechanisms by

which drugs interact with microbial proteins, though our findings highlight their potential

importance in shaping drug–microbiome dynamics. Looking forward, integrating systems ecology

frameworks with mechanistic and longitudinal approaches will provide deeper insights into drug–

microbiome interactions and help inform strategies to preserve microbiome resilience and

promote recovery from drug-induced perturbations.

METHODS

Sample collection and biobanking

The protocol for human stool sample collection (# 20160585–01 H) was approved by the Ottawa Health Science Network Research Ethics Board at the Ottawa Hospital, Ottawa, Canada. The inclusion criteria of participants were 18-65 years of age and healthy. Exclusion criteria included diagnosis of irritable bowel syndrome, inflammatory bowel disease, or diabetes; antibiotic use or gastroenteritis episode three months preceding collection; use of pro-/pre-biotics, laxatives, or anti-diarrheal drugs in the last month preceding collection; or pregnancy. In this study, we collected samples from nine healthy subjects (23-48 years of age, four females and five males). Sex/gender of participants was self-reported. Written informed consent was obtained from all participants for the collection and use of samples and data analysis for the present work. Sex/gender was not considered as a study variable in the experimental design, as the study focused on overall microbiome responses rather than sex-specific differences.

The gut microbiomes were collected and biobanked according to our previously validated workflow[20]. Briefly, fecal samples were collected off-site from participants and subsequently processed in the laboratory under anaerobic conditions to prepare a 20% (w/v) fecal slurry in a stabilization buffer consisting of 1× phosphate-buffered saline (PBS), 10% (v/v) glycerol, and 1 mg/mL L-cysteine. Samples were aliquoted in 15 mL Falcon tubes and stored at -80 °C until culturing. We have previously validated that the live microbiota biobanking approach can reproduce the metaproteomic responses of fresh stool samples[20].

Key resources

Key resources used in this study, including reagents for microbiota culturing, metaproteomics analyses, experimental consumables, instruments and software are summarized in Supplementary Table S1.

RapidAIM 2.0 assay

Drug plate preparation: For quality control (QC), each plate included triplicate wells for DMSO (negative control) and triplicate wells for kestose (positive control). Blank wells contained only media and were not inoculated; they were designed to be tested for protein concentration to confirm the absence of contamination. Information of the tested compounds is given in Supplementary Data S1. For most compounds, 10 mM stocks in DMSO were made. For

compounds without exact molecular weights (e.g. mixtures of polymers), 1 mg/ml stocks were

prepared. For compounds dissolvable in ddH$_2$O but not in DMSO, the stocks were prepared in

ddH$_2$O. RapidAIM 2.0 assay plates were next prepared by adding 50 µL of each stock solution in

the well designated to the specific compound to achieve 500 µM (or 50 µg/mL for compounds

without exact molecular weights) concentration in the final culture system. Compounds were

distributed randomly across four plates. For negative control wells, 50 µL of DMSO was used. For

compound stocks prepared with ddH$_2$O, 50 µL of DMSO was added to control the DMSO effect.

The plates were stored at -20 °C before the culturing experiment following RapidAIM 2.0 workflow.

Microbiome culturing and drug treatment: Culture media was optimized and validated in our

previous study[41], which includes the following composition: 2.0 g/L peptone water, 2.0 g/L yeast

extract, 0.5 g/L L-cysteine hydrochloride, 2 mL/L Tween 80, 5 mg/L hemin, 10 µL/L vitamin K1,

1.0 g/L NaCl, 0.4 g/L K$_2$HPO$_4$, 0.4 g/L KH$_2$PO$_4$, 0.1 g/L MgSO$_4$·7H$_2$O, 0.1 g/L CaCl$_2$·2H$_2$O, 4.0 g/L

NaHCO$_3$, 4.0 g/L porcine gastric mucin, 0.25 g/L sodium cholate and 0.25 g/L sodium

chenodeoxycholate. RapidAIM 2.0 culturing experiments were performed in an anaerobic

chamber whose gas contained 5% H$_2$, 5% CO$_2$ and 90% N$_2$. The drug plate containing 50 µL

DMSO and a compound in each treatment well was added with 1 mL culture media and 100 µL

fecal slurries as inoculum. Samples were mixed sufficiently using multi-channel pipettes before

covering the wells tightly with a perforated silicone gel mat and sealed around using lab tape to

prevent popping up due to gas production in the culture. The deepwell plates were shaken at

500 rpm on an orbital shaker for 48 hours at 37 °C.

Microbial cell washing: After culturing, culture plates were centrifuged at 3,000 g for 45 minutes

at 4 °C to collect the microbial cells. The cell pellets were then washed three times with 1 mL of

cold PBS. In the final wash cycle, the pellets were resuspended in an additional 1 mL of cold PBS

and briefly centrifuged at 300 g for 5 minutes at 4 °C to clear debris. The resulting supernatant

was carefully transferred to a new deep well plate and subjected to a final centrifugation at

3,000 g for 45 minutes at 4 °C. The purified cell pellets were subsequently stored at -80 °C until

needed for cell lysis.

Microbial cell lysis and protein double-precipitation: Proteins were then extracted from the

cells using ultra-sonication and were purified using a double-precipitation procedure. Briefly,

microbial cell pellets were lysed with 150 µL of lysis buffer (8 M urea and 4% sodium dodecyl

sulfate (SDS) in 100 mM Tris-HCl, pH = 8.0) per well using a 96-channel liquid handler. The

lysates underwent sonication using Qsonica Q700 at 10 kHz, alternating 10 seconds on and 10

seconds off for a total of 10 minutes, with cooling at 8 °C to prevent overheating. The sonicated

lysates were precipitated overnight in ice-cold protein precipitation solution (50%:50%:0.1% (v:v:v)

of acetone:ethanol:acetic acid) at -20 °C. Following the first precipitation, the proteins were

pelleted and resuspended in 150 μL of protein resuspension buffer (6 M urea in 100 mM Tris-HCl,

pH = 8.0) and subjected to a second precipitation under the same condition.

Protein digestion: Protein concentrations of the DMSO control samples were measured using

the DC Protein Assay kit according to the manufacturer's instructions. An equal volume approach

was applied similar to described previously[4], ensuring comparable protein biomass

measurements between samples: DMSO control samples were diluted to 1 μg/μL with the

resuspension buffer (6 M urea in 100 mM Tris-HCl, pH 8), and the same volume of the

resuspension buffer was added to other sample wells. Protein digestion was performed using an

automated liquid handler (Hamilton Nimbus 96) following standard workflows of reduction,

alkylation, and overnight tryptic digestion as described in the RapidAIM 2.0 protocol[22].

Protein desalting and TMT labelling: After digestion, automated desalting of protein lysate was

performed using reverse-phase (RP) desalting columns (IMCS, 04T-H6R05-1-10-96). A 240 μL

aliquot of the 800 μL desalting elute was used for 96-well format TMT 11plex™-labeling following

an economic and robust approach that was previously described[21]. Briefly, 11 different tandem

mass tags (TMT) reagents are first resuspended in acetonitrile, aliquoted into precise 50 μg TMT

per channel, freeze-dried, and stored at -80 °C. For peptide labeling, the peptide aliquots

(dissolved in 100 mM TEAB with 20% acetonitrile) are incubated in the TMT reagent plates at

25 °C for 1-2 hours. Post-incubation, the reaction is quenched, and the pH adjusted to 2-3. Finally,

the labeled peptides are combined, desalted, and freeze-dried for further analysis.

LC-MS/MS analysis

TMT-labelled samples: LC-MS/MS analyses of TMT-labeled samples were performed using an

UltiMate 3000 RSLCnano system coupled to an Orbitrap Exploris 480 mass spectrometer

(ThermoFisher Scientific Inc.). Peptides were separated using a 75 μm × 50 cm analytical column

packed with reverse phase beads (1.9 μm/120-Å ReproSil-Pur C18 resin, Dr. Maisch HPLC

GmbH), applying a two-hour gradient from 5%~35% of buffer B (0.1% (v/v) formic acid with 80%

(v/v) acetonitrile) ratio with buffer A being 0.1% formic acid, at a flow rate of 300 nL/min. Full MS

scans were acquired in the Orbitrap at a resolution of 120,000 for 350-1,200 m/z. Data were

acquired in a data-dependent mode with a cycle time of 2 seconds between master scans.

MS/MS scans were performed using higher-energy collisional dissociation (HCD) at a resolution

of 45,000, with a collision energy of 30%. The dynamic exclusion was set to 60 seconds after a

single count.

Label-free samples: LC-MS/MS analyses of label-free samples were also performed using the
UltiMate 3000 RSLCnano and the Orbitrap Exploris 480. Peptides were separated using a tip
column (75 μm inner diameter × 15 cm) packed with reverse phase beads (3 μm/120 Å ReproSil-
Pur C18 resin, Dr. Maisch HPLC GmbH), applying a one-hour gradient from 5%~ 35% of buffer B
(0.1% (v/v) formic acid with 80% (v/v) acetonitrile) ratio, with buffer A being 0.1% formic acid, at a
flow rate of 300 nL/min. Full MS scans were acquired in the Orbitrap at a resolution of 60,000 for
350-1,200 m/z by data-dependent MS/MS scan of the 15 most intense ions. MS/MS scans were
performed using higher-energy collisional dissociation (HCD) at a resolution of 15,000, with a
collision energy of 30%. The dynamic exclusion repeat count was set to one, and the repeat
exclusion duration was set to 20 seconds.

Metagenomics analysis

Fecal slurries of different individual microbiomes were first subjected to whole metagenomic DNA
extraction using the FastDNA Spin Kit (MP Biomedicals, 116540600). Quantification of DNA
contents was performed using the Qubit High Sensitivity dsDNA Assay Kit, following the
manufacturer's instructions[42]. Metagenomic DNA sequencing was performed with Illumina
NovaSeq 6000 platform at Génome Québec CES using PCR-free library preparation. A total of
2.58B reads were sequenced for six samples with an average of 430M (±70M) per individual
microbiome. The generated fastq sequences were preprocessed using fastp v.0.23.1[43] to remove
adaptors and quality filter the reads with default parameters. Human and phiX reads were then
removed by mapping the reads to a non-redundant version of the Genome Reference
Consortium Human Build 38 (GRCh38hg38; RefSeq: GCF_000001405.39) and phiX reference
genomes using the Kraken2 v.2.1.2 package[44]. The quality-filtered paired-end reads were then
error-corrected using bbcms v38.34 from the BBTools[45] with default parameters. An average of
420M (±69M) reads were obtained after these steps. Metagenome assembly of the error-
corrected and quality-filtered non-human reads was performed using MEGAHIT v1.2.9[46] using
the --presets meta-large --min-contig-len 1000 parameters. A total of 640,099 contigs with
lengths >1,000 bp were individually assembled, with an average of 106,683 (±32,133) contigs per
individual microbiome. Metagenomic binning was performed using SemiBin v1.5.1[47] followed by
quality filtering (completeness >50%) with DAS Tool v1.1.4[48], the high-quality bins obtained
through this process were considered as metagenome-assembled genomes (MAGs)[23]. Species-
and genus-level richness and shared taxa across metagenomes are shown as Supplementary
Figure S8.

The assembled contigs were then annotated using the PROkaryotic Dynamic programming

Gene-finding ALgorithm (Prodigal) v2.6.3[49] to predict open reading frames (ORF). The contigs

were translated into amino acid sequences using the anonymous gene prediction mode (prodigal

-p meta) and default parameters and annotated a total of 3.8M protein-coding genes. Lastly, the

outputs were compiled into a single FASTA file and used as the metagenome-inferred protein

database for the metaproteomic search.

Database search strategy

The initial compound screening was performed using TMT-labeled data analyzed with MetaLab

(Version 2.3.0)[50], which implements an automated spectral clustering algorithm to group similar

MS/MS spectra prior to database searching against the IGC reference database[51]. The samples

selected from TMT-based results were re-analyzed by label-free LC-MS/MS quantification. Since

reference catalog-based databases often lack microbiome-specific peptides[52], we developed a

bioinformatic workflow for sample set specific database generation (Supplementary Figure S1a).

Briefly, metagenomic prodigal sequences (in FASTA format) from each individual microbiome

were used to construct reduced databases of each subset of individual microbiome's LFQ LC-

MS/MS raw files following the MetaPro-IQ workflow[26]. First, databases were reduced from two-

step X!Tandem search to remove non-expressed sequences based on actual MS data of each

subject's raw file subsets. Next, reduced subject-specific databases were merged and subjected

to redundancy removal at 99% using CD-HIT. The resulting FASTA file containing 491,506

sequences was used for database search of LC-MS/MS label-free files using MaxQuant (version

1.5.2.8), as well as revisiting the TMT dataset's database search using MaxQuant (version

2.0.3.0). Both LFQ and TMT searches applied carbamidomethylation as a fixed modification (C)

and variable modifications including acetylation at the protein N-terminus and oxidation of

methionine, with a fragment ion tolerance of 20 ppm. The false discovery rates (FDR) for peptide

spectrum matches (PSM) and proteins were set to 0.01, with a minimum peptide length of 7.

Match between runs were enabled with matching time window of 0.7 minutes and alignment time

window of 20 minutes. To specifically investigate compound effects on protein expressions

related to antimicrobial resistance, we conducted a focused search against the full CARD

database[35] (downloaded December 27th, 2022) using MaxQuant 1.5.2.8, with the same search

parameters as above.

Data preprocessing

For the TMT dataset, proteinGroup.txt was normalized using MSstatsTMT[53] with the "msstats" option selected, and perform estimate size factors on matrix using Deseq2 R package, the processing resulted in 19,318 proteinGroups across all TMT samples. A Q15 criteria was applied to remove highly random sparse features, and SpectroFM was applied for data imputation, resulting in a matrix of 6085 proteinGroups.

For the LFQ dataset, directLFQ[27] (GUI MacOS version 0.2.8) was used for efficient label-free quantification, followed by protein inference using a custom in-house script (available at https://github.com/ningzhibin/protein-Inference). Razor and unique peptides were considered for protein quantification. Dataset was further harmonized by individual microbiome batch using the R package HarmonizR[28], which implements the ComBat algorithm while appropriately handling missing values[28].

For the ARP database search result, protein intensities in the proteinGroups file were filtered to remove potential contaminants and reverse sequences, then all protein intensities in each sample were summed as ARP intensities. The ARP ratio in each sample was then calculated by dividing the ARP intensities by the total protein intensities in the LFQ search result.

Taxonomic and functional annotation

Taxonomic annotation was performed in MetaLab 2.3.0[50] by searching the peptide sequences against the pep2taxa database in MetaLab. Taxonomic annotations with three or more unique peptides were considered confident, following the conservative criteria used in previous studies[50,54]. eggNOG mapper (version 2.1.12)[55] was used for functional annotation of protein sequences, and COG functional enrichment of selected eggNOG matches was performed using the Enrichment Analysis tool of iMetaShiny[56].

Downstream data analysis

Compound structure clustering: Clustering of compounds was performed using the fingerprint, and rcdk R packages following a method developed by Voicu et al.[57] Briefly, The SMILES (Simplified Molecular Input Line Entry System) notation for each compound (excluding those without a defined chemical structure) was parsed using the parse.smiles() function from the rcdk package, then compound fingerprints were obtained by the get.fingerprint() function of the fingerprint package using type "extended" as suggested by Voicu et al.[57] Next, fp.sim.matrix() function was used to calculates the similarity matrix for the set of compound fingerprints by the

Tanimoto coefficient. Finally, the compound structure similarity matrix was passed to the pvclust

function, which performed hierarchical clustering and generated 100 bootstrap samples to

evaluate the stability of the clustering results.

Kernel density estimation (KDE): Kernel density estimation (KDE) of the two-dimensional ARP

versus nFR_p landscape was performed using the kde2d() function of R package MASS with n

resolution set as 100, and 3D surface plot of the KDE results was visualized by R package plotly.

Feature set comparison: Comparison of shared features between different groups were

performed using upset plot wrapped by the "Sets Explorer" in iMetaShiny[56]

(https://shiny2.imetalab.ca/shiny/rstudio/SetExplorer/).

Principal component analysis: For principal component analysis of metaproteomic drug

responses, protein intensity data was first log10 transformed, then normalized using the

estimateSizeFactorsForMatrix() function of the R package DeSeq2. Next, the pca() function of the

pcaMethods package was used to perform PCA analysis using the nonlinear estimation by the

NIPALS algorithm, allowing for the presence of missing values[29].

Finding microbial homologs of human protein drug targets: Human drug targets of our

selected drugs were extracted from the comprehensive map of molecular drug targets

(NIHMS80822-supplement-Supplementary_Information_S2.xlsx table in the referred paper)

summarized by Santos et al.[30] To identify homologs of human protein drug targets in the tested

microbiomes, we employed a profile Hidden Markov Model (HMM)-based approach using the

HMMER suite (v3.3). For each target protein, *hmm* and *seed* files were downloaded from the

Pfam database[58] (https://www.ebi.ac.uk/interpro/), then hmmsearch[31] was used to perform

microbial homolog sequence search by protein alignment/profile-HMM against our metaproteomic

sequence database of this dataset.

Functional redundancy calculation: Within-sample taxonomic diversity (TD) and functional

redundancy (FR_p) of each metaproteome was calculated using the label-free quantified dataset

following our previously developed approach[25]. Briefly, the proteome-content network of each

individual metaproteome data was generated according to pep2taxa-based taxonomic annotation

and eggNOG-based Clusters of Orthologous Genes (COG) annotations. Genus-level biomass

contributions were estimated by genus-specific peptide intensities in pep2taxa-based taxonomic

annotation. Then, taxonomic diversity (TD) and functional redundancy normalized by TD (nFR_p)

was calculated using the python codes provided by our previous research[25]

(https://github.com/yvonnelee1988/Metaproteome_FRp).

Differential protein analysis: Differential protein expression between DMSO and the seven

neurological drugs (amoxapine, duloxetine, fluoxetine, fluphenazine, fluvoxamine, paroxetine and

sertraline) was analyzed using the Differential Protein Analyzer of iMetaLab Suite[56]. The

functional annotation and taxonomic representatives of the resulting proteins were annotated by

eggNOG mapper and visualized using Sanky plot with geom_alluvium() function of ggplot2.

PhyloFunc analysis and community state types (CST) clustering: The phylogenetic tree was

constructed from the deep metagenomics dataset using IQ-TREE2 (http://www.iqtree.org/), the

tree inference included 1000 replicates for the SH-like approximate likelihood ratio test (aLRT)

and 1000 ultrafast bootstrap replicates. PhyloFunc was then computed with the phylogenetic tree

and the original directLFQ-quantified protein group table, with taxonomy information annotated by

referencing the metagenome-assembled genomes (MAGs), and functional annotation performed

by eggNOG mapper[55]. The Python package *phylofunc* [23] was then used to compute functional

distances matrices across metaproteomics samples.

Other analyses and visualizations: Correlation analysis was performed using cor.test() function

in the stats package. Scatter plots with aligned density plots were created by ggscatterhist()

function in the ggpubr package. Partitioning around medoids analyses were performed using

pam() function in the cluster package. Taxonomic heatmap and clustering was performed using

the pheatmap package.

Computation of the generalized Lotka-Volterra (gLV) model

The gLV competition model, considering between-cluster interactions can be written as follows:

836
$$\frac{dx_i}{dt} = f_i(\mathbf{x}) = r_i x_i \left[\frac{K_i - x_i - \left(\sum_{j \neq i}^{3} \alpha_{ij} x_j \right)}{K_i} \right] \qquad (1),$$

where, i and j represent i^{th} and j^{th} microbial cluster, respectively. In our case, $1 \leq i \neq j \leq 3$. $\mathbf{x}=[x_1, x_2,$

$x_3]$ represents the population size of each microbial cluster. The parameters r_i, K_i and α_{ij}

represent the intrinsic growth rate of the i^{th} microbial cluster, environmental carrying capacity of

the i^{th} microbial cluster, and competition coefficient from the j^{th} to the i^{th} microbial cluster,

respectively. We note that the stable state is dependent on model parameters r_i, K_i and α_{ij} rather

than the initial population size of each microbial cluster. Based on the gLV competition model

considering between-cluster interactions given by equation (1), we can calculate population sizes

at the equilibrium state where $\frac{dx_i}{dt} = 0$. Therefore, at equilibrium, we have:

845
$$x_i^* = K_i - \sum_{j \neq i}^{3} \alpha_{ij} x_j \qquad (2)$$

Based on Lyapunov's indirect method, the Jacobian matrix **J** of the gLV model at the equilibrium

point $\mathbf{x}^*=[x^*_1, \ x^*_2, \ x^*_3]$ can be obtained. By linearizing equation (1) around $\mathbf{x}^*$ and its neighborhood,

it is then transformed into a linear dynamic model:

$$\frac{d\mathbf{x}}{dt} = \mathbf{J}\mathbf{x} \tag{3}$$

Where, $$\mathbf{J} = J_{ij} = \frac{\partial f_i(\mathbf{x})}{\partial x_j} \tag{4}$$

If all eigenvalues of the matrix **J** lie in the left half of the complex plane, then we achieve a stable

equilibrium point of the gLV model.

Ordinary equation of the gLV model was solved using scipy.integrate function odeint() in Jupyter

Notebook running Python 3. And visualization of model dynamics was performed using

matplotlib.pyplot.plot().

Functional redundancy enhancement to combat ARP response

An additional RapidAIM test was performed to investigate the impact of enhanced functional

redundancy on mitigating drug-induced ARP within microbial communities. First, to evaluate the

impact of FOS on functional redundancy, three additional individual microbiomes were initially

screened for their response of nFR_p to 10 mg/mL of FOS treatment. The microbiome exhibiting a

high increase in nFR_p was selected for further analysis. It was subjected to 250 µM sertraline,

10 mg/mL FOS, or 250 µM sertraline + 10 mg/mL FOS in 25 µL DMSO, and blank (25 µL DMSO

only) treatments for 48 hours of culturing, followed by metaproteomic analyses to assess changes

in functional redundancy and ARP levels in different groups. Technical triplicates were performed

for each group.

Statistics and Reproducibility

Estimation of sample size was based on practical considerations of large-scale drug screening

(312 compounds + 6 control samples per biological replicate), which necessitated minimizing the

number of biological replicates per condition. Using an effect size of Cohen's d = 1.2 (as

confirmed by post hoc analysis of the DMSO vs Sertraline comparison), with a significance level

of 0.05 and a statistical power threshold of 0.7, the minimum required sample size was estimated

to be six biological replicates per group. All data were retained for analysis unless missingness

was caused by technical issues during sample processing. One DMSO control sample from

subject V47 was lost during the mass spectrometry acquisition step. In addition, sample

V52_P1E3 yielded only 290 quantified values after database searching and was excluded due to

insufficient data quality. The distribution of compounds across 96-well plates was randomized.

Due to the nature of automated proteomics sample processing, the downstream handling steps
were not randomized; however, for each individual, LC–MS/MS sample injection order was
randomized. Although the investigators were not blinded to the allocation during experiments and
outcome assessment, the automated sample processing ensured that experimental and control
groups were handled in a uniform and unbiased manner.

DATA AVAILABILITY

All metaproteomics data sets reported in this study have been deposited in the
ProteomeXchange database through the PRIDE repository under accession codes PXD056930
[https://www.ebi.ac.uk/pride/archive/projects/PXD056930],
PXD056888[https://www.ebi.ac.uk/pride/archive/projects/PXD056888],
PXD056968[https://www.ebi.ac.uk/pride/archive/projects/PXD056968]. The metagenomics data
used in this study are deposited in the NCBI Sequence Read Archive (SRA) with the accession
numbers SRR30894461[https://www.ncbi.nlm.nih.gov/sra/?term=SRR30894461],
SRR30894462[https://www.ncbi.nlm.nih.gov/sra/?term=SRR30894462],
SRR30894463[https://www.ncbi.nlm.nih.gov/sra/?term=SRR30894463],
SRR30894464[https://www.ncbi.nlm.nih.gov/sra/?term=SRR30894464],
SRR30894465[https://www.ncbi.nlm.nih.gov/sra/?term=SRR30894465] and
SRR29021656[https://www.ncbi.nlm.nih.gov/sra/?term=SRR29021656].

CODE AVAILABILITY

Code and related data to reproduce the analyses in this study has been published in figshare
through DOI: 10.6084/m9.figshare.29816780. In addition, the source code of the Shiny app is
made publicly available via GitHub at https://github.com/yvonnelee1988/MPR_Viz and Zendo at
DOI: 10.5281/zenodo.16990829. Any additional information required to reanalyze the data
reported in this paper is available from the lead contact (Daniel Figeys,
Daniel.Figeys@quadram.ac.uk) upon request.

REFERENCES

1. Lam, K.N., Alexander, M., and Turnbaugh, P.J. (2019). Precision Medicine Goes Microscopic: Engineering the Microbiome to Improve Drug Outcomes. Cell Host Microbe *26*, 22–34. https://doi.org/10.1016/j.chom.2019.06.011.

2. Maier, L., Pruteanu, M., Kuhn, M., Zeller, G., Telzerow, A., Anderson, E.E., Brochado, A.R., Fernandez, K.C., Dose, H., Mori, H., et al. (2018). Extensive impact of non-antibiotic drugs on human gut bacteria. Nature *555*, 623–628. https://doi.org/10.1038/nature25979.

3. Vich Vila, A., Collij, V., Sanna, S., Sinha, T., Imhann, F., Bourgonje, A.R., Mujagic, Z., Jonkers, D.M.A.E., Masclee, A.A.M., Fu, J., et al. (2020). Impact of commonly used drugs on the composition and metabolic function of the gut microbiota. Nat Commun *11*, 362. https://doi.org/10.1038/s41467-019-14177-z.

4. Li, L., Ning, Z., Zhang, X., Mayne, J., Cheng, K., Stintzi, A., and Figeys, D. (2020b). RapidAIM: a culture- and metaproteomics-based Rapid Assay of Individual Microbiome responses to drugs. Microbiome *8*, 33. https://doi.org/10.1186/s40168-020-00806-z.

5. Javdan, B., Lopez, J.G., Chankhamjon, P., Lee, Y.-C.J., Hull, R., Wu, Q., Wang, X., Chatterjee, S., and Donia, M.S. (2020). Personalized Mapping of Drug Metabolism by the Human Gut Microbiome. Cell *181*, 1661-1679.e22. https://doi.org/10.1016/j.cell.2020.05.001.

6. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R., and Goodman, A.L. (2019). Mapping human microbiome drug metabolism by gut bacteria and their genes. Nature *570*, 462–467. https://doi.org/10.1038/s41586-019-1291-3.

7. Noto Guillen, M., Li, C., Rosener, B., and Mitchell, A. (2024). Antibacterial activity of nonantibiotics is orthogonal to standard antibiotics. Science *384*, 93–100. https://doi.org/10.1126/science.adk7368.

8. Bottery, M.J., Matthews, J., Wood, A.J., Krogh Johansen, H., Pitchford, J., and Friman, V.-P. (2022). Inter-species interactions alter antibiotic efficacy in bacterial communities. ISME J *16*, 812–821. https://doi.org/10.1038/s41396-021-01130-6.

9. Yu, J.S.L., Correia-Melo, C., Zorrilla, F., Herrera-Dominguez, L., Wu, M.Y., Hartl, J., Campbell, K., Blasche, S., Kreidl, M., Egger, A.-S., et al. (2022). Microbial communities form rich extracellular metabolomes that foster metabolic interactions and promote drug tolerance. Nat Microbiol *7*, 542–555. https://doi.org/10.1038/s41564-022-01072-5.

10. Lindsay, R.J., Jepson, A., Butt, L., Holder, P.J., Smug, B.J., Gudelj, I., and Chase, J. (2021). Would that it were so simple: Interactions between multiple traits undermine classical single-trait-based predictions of microbial community function and evolution. Ecol Lett *24*, 2775–2795. https://doi.org/10.1111/ele.13861.

11. Flynn, J.M., Cameron, L.C., Wiggen, T.D., Dunitz, J.M., Harcombe, W.R., and Hunter, R.C. (2020).
Disruption of cross-feeding inhibits pathogen growth in the sputa of patients with cystic fibrosis.
mSphere *5*, e00343-20. https://doi.org/10.1128/mSphere.00343-20.

12. Garcia-Santamarina, S., Kuhn, M., Devendran, S., Maier, L., Driessen, M., Mateus, A., Mastrorilli, E.,
Brochado, A.R., Savitski, M.M., Patil, K.R., et al. (2023). Emergence of community behaviors in the gut
microbiota upon drug treatment. Preprint at bioRxiv, https://doi.org/10.1101/2023.06.13.544832
https://doi.org/10.1101/2023.06.13.544832.

13. Bottery, M.J., Pitchford, J.W., and Friman, V.-P. (2021). Ecology and evolution of antimicrobial
resistance in bacterial communities. ISME J *15*, 939–948. https://doi.org/10.1038/s41396-020-00832-
7.

14. Gotfryd, K., Boesen, T., Mortensen, J.S., Khelashvili, G., Quick, M., Terry, D.S., Missel, J.W., LeVine,
M.V., Gourdon, P., Blanchard, S.C., et al. (2020). X-ray structure of LeuT in an inward-facing occluded
conformation reveals mechanism of substrate release. Nat Commun *11*, 1005.
https://doi.org/10.1038/s41467-020-14735-w.

15. Zhou, Z., Zhen, J., Karpowich, N., Law, C., Reith, M., and Wang, D. (2009). Antidepressant Specificity of
Serotonin Transporter Suggested by Three LeuT-SSRI Structures. Nat Struct Mol Biol *16*.
https://doi.org/10.1038/nsmb.1602.

16. Fujita, H., Ushio, M., Suzuki, K., Abe, M.S., Yamamichi, M., Iwayama, K., Canarini, A., Hayashi, I.,
Fukushima, K., Fukuda, S., et al. (2023). Alternative stable states, nonlinear behavior, and
predictability of microbiome dynamics. Microbiome *11*, 63. https://doi.org/10.1186/s40168-023-
01474-5.

17. Beisner, B., Haydon, D., and Cuddington, K. (2003). Alternative stable states in ecology. Front Ecol
Environ *1*, 376–382. https://doi.org/10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2.

18. Louca, S., Polz, M.F., Mazel, F., Albright, M.B.N., Huber, J.A., O'Connor, M.I., Ackermann, M., Hahn,
A.S., Srivastava, D.S., Crowe, S.A., et al. (2018). Function and functional redundancy in microbial
systems. Nat Ecol Evol *2*, 936–943. https://doi.org/10.1038/s41559-018-0519-1.

19. Biggs, C.R., Yeager, L.A., Bolser, D.G., Bonsell, C., Dichiera, A.M., Hou, Z., Keyser, S.R., Khursigara, A.J.,
Lu, K., Muth, A.F., et al. (2020). Does functional redundancy affect ecological stability and resilience?
A review and meta-analysis. Ecosphere *11*, e03184. https://doi.org/10.1002/ecs2.3184.

20. Zhang, X., Walker, K., Mayne, J., Li, L., Ning, Z., Stintzi, A., and Figeys, D. (2022). Evaluating live
microbiota biobanking using an *ex vivo* microbiome assay and metaproteomics. Gut Microbes *14*,
2035658. https://doi.org/10.1080/19490976.2022.2035658.

21. Creskey, M., Li, L., Ning, Z., Brown, E.E., Mayne, J., Walker, K., Ampaw, A., Ben, R., Zhang, X., and
Figeys, D. (2022). An economic and robust TMT labeling approach for high throughput proteomic and
metaproteomic analysis. Proteomics *Online ahead of print*, e2200116.

22. Li, L., Mayne, J., Beltran, A., Zhang, X., Ning, Z., and Figeys, D. (2024). RapidAIM 2.0: a high-throughput
assay to study functional response of human gut microbiome to xenobiotics. Microbiome Res Rep *3*,
26. https://doi.org/10.20517/mrr.2023.57.

23. Wang, L., Simopoulos, C.M.A., Serrana, J.M., Ning, Z., Li, Y., Sun, B., Yuan, J., Figeys, D., and Li, L.
(2025). PhyloFunc: phylogeny-informed functional distance as a new ecological metric for
metaproteomic data analysis. Microbiome *13*, 50. https://doi.org/10.1186/s40168-024-02015-4.

24. Fuentes, A., Pineda, M., and Venkata, K. (2018). Comprehension of Top 200 Prescribed Drugs in the
US as a Resource for Pharmacy Teaching, Training and Practice. Pharmacy *6*, 43.
https://doi.org/10.3390/pharmacy6020043.

25. Li, L., Wang, T., Ning, Z., Zhang, X., Butcher, J., Serrana, J.M., Simopoulos, C.M.A., Mayne, J., Stintzi, A.,
Mack, D.R., et al. (2023). Revealing proteome-level functional redundancy in the human gut
microbiome using ultra-deep metaproteomics. Nat Commun *14*, 3428.
https://doi.org/10.1038/s41467-023-39149-2.

26. Zhang, X., Ning, Z., Mayne, J., Moore, J.I., Li, J., Butcher, J., Deeke, S.A., Chen, R., Chiang, C.-K., Wen,
M., et al. (2016). MetaPro-IQ: a universal metaproteomic approach to studying human and mouse gut
microbiota. Microbiome *4*, 31. https://doi.org/10.1186/s40168-016-0176-z.

27. Ammar, C., Schessner, J.P., Willems, S., Michaelis, A.C., and Mann, M. (2023). Accurate Label-Free
Quantification by directLFQ to Compare Unlimited Numbers of Proteomes. Mol Cell Proteomics *22*,
100581. https://doi.org/10.1016/j.mcpro.2023.100581.

28. Voß, H., Schlumbohm, S., Barwikowski, P., Wurlitzer, M., Dottermusch, M., Neumann, P., Schlüter, H.,
Neumann, J.E., and Krisp, C. (2022). HarmonizR enables data harmonization across independent
proteomic datasets with appropriate handling of missing values. Nat Commun *13*, 3523.
https://doi.org/10.1038/s41467-022-31007-x.

29. Wold, H. (1966). Estimation of principal components and related models by iterative least squares. In
Multivariate Analysis, P. R. Krishnajah, ed. (Academic Press), pp. 391–420.

30. Santos, R., Ursu, O., Gaulton, A., Bento, A.P., Donadi, R.S., Bologa, C.G., Karlsson, A., Al-Lazikani, B.,
Hersey, A., Oprea, T.I., et al. (2017). A comprehensive map of molecular drug targets. Nat Rev Drug
Discov *16*, 19–34. https://doi.org/10.1038/nrd.2016.230.

31. Potter, S.C., Luciani, A., Eddy, S.R., Park, Y., Lopez, R., and Finn, R.D. (2018). HMMER web server: 2018
update. Nucleic Acids Res *46*, W200–W204. https://doi.org/10.1093/nar/gky448.

32. Gonze, D., Lahti, L., Raes, J., and Faust, K. (2017). Multi-stability and the origin of microbial
community types. ISME J *11*, 2159–2166. https://doi.org/10.1038/ismej.2017.60.

33. Zhou, Y., Zhang, Y., Zhao, D., Yu, X., Shen, X., Zhou, Y., Wang, S., Qiu, Y., Chen, Y., and Zhu, F. (2024).
TTD: Therapeutic Target Database describing target druggability information. Nucleic Acids Res *52*,
D1465–D1477. https://doi.org/10.1093/nar/gkad751.

34. Keilhauer, E.C., Hein, M.Y., and Mann, M. (2015). Accurate Protein Complex Retrieval by Affinity
Enrichment Mass Spectrometry (AE-MS) Rather than Affinity Purification Mass Spectrometry (AP-MS).
Mol Cell Proteomics *14*, 120–135. https://doi.org/10.1074/mcp.M114.041012.

35. Alcock, B.P., Huynh, W., Chalil, R., Smith, K.W., Raphenya, A.R., Wlodarski, M.A., Edalatmand, A.,
Petkau, A., Syed, S.A., Tsang, K.K., et al. (2023). CARD 2023: expanded curation, support for machine
learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. Nucleic
Acids Research *51*, D690–D699. https://doi.org/10.1093/nar/gkac920.

36. Grießhammer, A., de la Cuesta-Zuluaga, J., Zahir, T., Müller, P., Gekeler, C., Chang, H., Schmitt, K.,
Planker, C., Bohn, E., Nguyen, T.H., et al. (2023). Non-antibiotic drugs break colonization resistance
against pathogenic Gammaproteobacteria. bioRxiv, 2023.11.06.564936.
https://doi.org/10.1101/2023.11.06.564936.

37. Duan, J., Sun, J., Ma, X., Du, P., Dong, P., Xue, J., Lu, Y., and Jiang, T. (2025). Association of
escitalopram-induced shifts in gut microbiota and sphingolipid metabolism with depression-like
behavior in wistar-kyoto rats. Transl Psychiatry *15*, 54. https://doi.org/10.1038/s41398-025-03277-8.

38. Ait Chait, Y., Mottawea, W., Tompkins, T.A., and Hammami, R. (2020). Unravelling the antimicrobial
action of antidepressants on gut commensal microbes. Sci Rep *10*, 17878.
https://doi.org/10.1038/s41598-020-74934-9.

39. Shen, Y., Yang, X., Li, G., Gao, J., and Liang, Y. (2021). The change of gut microbiota in MDD patients
under SSRIs treatment. Sci Rep *11*, 14918. https://doi.org/10.1038/s41598-021-94481-1.

40. Allison, S.D., and Martiny, J.B.H. (2008). Resistance, resilience, and redundancy in microbial
communities. Proc. Natl. Acad. Sci. U.S.A. *105*, 11512–11519.
https://doi.org/10.1073/pnas.0801925105.

41. Li, L., Abou-Samra, E., Ning, Z., Zhang, X., Mayne, J., Wang, J., Cheng, K., Walker, K., Stintzi, A., and
Figeys, D. (2019). An in vitro model maintaining taxon-specific functional activities of the gut
microbiome. Nat Commun *10*, 4146. https://doi.org/10.1038/s41467-019-12087-8.

42. Yan, A., Butcher, J., Schramm, L., Mack, D.R., and Stintzi, A. (2023). Multiomic spatial analysis reveals
a distinct mucosa-associated virome. Gut Microbes *15*, 2177488.
https://doi.org/10.1080/19490976.2023.2177488.

43. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics *34*, i884–i890. https://doi.org/10.1093/bioinformatics/bty560.

44. Wood, D.E., Lu, J., and Langmead, B. (2019). Improved metagenomic analysis with Kraken 2. Genome
Biol *20*, 257. https://doi.org/10.1186/s13059-019-1891-0.

45. Bushnell, B. (2014). BBMap: A Fast, Accurate, Splice-Aware Aligner. In Report Number: LBNL-7065E.

46. Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. (2015). MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics
31, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033.

47. Pan, S., Zhao, X.-M., and Coelho, L.P. (2023). SemiBin2: self-supervised contrastive learning leads to
better MAGs for short- and long-read sequencing. Bioinformatics *39*, i21–i29.
https://doi.org/10.1093/bioinformatics/btad209.

48. Sieber, C.M.K., Probst, A.J., Sharrar, A., Thomas, B.C., Hess, M., Tringe, S.G., and Banfield, J.F. (2018).
Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nat
Microbiol *3*, 836–843. https://doi.org/10.1038/s41564-018-0171-1.

49. Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010). Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics *11*, 119.
https://doi.org/10.1186/1471-2105-11-119.

50. Cheng, K., Ning, Z., Zhang, X., Li, L., Liao, B., Mayne, J., and Figeys, D. (2020). MetaLab 2.0 Enables
Accurate Post-Translational Modifications Profiling in Metaproteomics. J Am Soc Mass Spectrom *31*,
1473–1482. https://doi.org/10.1021/jasms.0c00083.

51. Li, J., Jia, H., Cai, X., Zhong, H., Feng, Q., Sunagawa, S., Arumugam, M., Kultima, J.R., Prifti, E., Nielsen,
T., et al. (2014). An integrated catalog of reference genes in the human gut microbiome. Nature
Biotechnology *32*, 834–841. https://doi.org/10.1038/nbt.2942.

52. Van Den Bossche, T., Armengaud, J., Benndorf, D., Blakeley-Ruiz, J.A., Brauer, M., Cheng, K., Creskey,
M., Figeys, D., Grenga, L., Griffin, T.J., et al. (2025). The microbiologist's guide to metaproteomics.
iMeta *4*. https://doi.org/10.1002/imt2.70031.

53. Huang, T., Choi, M., Tzouros, M., Golling, S., Pandya, N.J., Banfai, B., Dunkley, T., and Vitek, O. (2020).
MSstatsTMT: Statistical Detection of Differentially Abundant Proteins in Experiments with Isobaric
Labeling and Multiple Mixtures. Mol Cell Proteomics *19*, 1706–1723.
https://doi.org/10.1074/mcp.RA120.002105.

54. Jagtap, P.D., Blakely, A., Murray, K., Stewart, S., Kooren, J., Johnson, J.E., Rhodus, N.L., Rudney, J., and
Griffin, T.J. (2015). Metaproteomic analysis using the Galaxy framework. PROTEOMICS *15*, 3553–3565.
https://doi.org/10.1002/pmic.201500074.

55. Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. (2021). eggNOG-
mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the
Metagenomic Scale. Molecular Biology and Evolution *38*, 5825–5829.
https://doi.org/10.1093/molbev/msab293.

56. Li, L., Ning, Z., Cheng, K., Zhang, X., Simopoulos, C.M.A., and Figeys, D. (2022). iMetaLab Suite: A one-
stop toolset for metaproteomics. iMeta *1*, e25. https://doi.org/10.1002/imt2.25.

57. Voicu, A., Duteanu, N., Voicu, M., Vlad, D., and Dumitrascu, V. (2020). The rcdk and cluster R packages
applied to drug candidate selection. J Cheminform *12*, 3. https://doi.org/10.1186/s13321-019-0405-0.

58. Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A., Bileschi, M.L., Bork,
P., Bridge, A., Colwell, L., et al. (2023). InterPro in 2022. Nucleic Acids Res *51*, D418–D427.
https://doi.org/10.1093/nar/gkac993.

59. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann, A.,
Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene Set Knowledge Discovery
with Enrichr. Current Protocols *1*, e90. https://doi.org/10.1002/cpz1.90.

1082

ACKNOWLEDGMENTS

L.L acknowledges funding from National Natural Science Foundation of China (grant 32370050), and the State Key Laboratory of Medical Proteomics (SKLP-Y202402). D.F. acknowledges funding from Genome Canada, the Province of Ontario and NSERC. The authors would like to thank Dr. Dawei Hu for his advice and help on the gLV model, and Dr. Yangyu Liu for his advice on functional redundancy analysis. The authors also acknowledge technical support platforms and staffs at the Faculty of Medicine, University of Ottawa and National Center for Protein Sciences (Beijing) for providing essential resources and assistance.

AUTHOR CONTRIBUTIONS

D.F. and L.L. designed and supervised the study, and wrote the manuscript; M.H., C.M.A.S., L.L., Xu Z. and D.F. selected compounds for this study. J.M., Xu Z., K.W. and H.Q. recruited healthy volunteers, collected individual fecal samples, collected written consent and biobanked the samples. M.H. and L.L. prepared the therapeutic compound library. L.L. established, validated, and performed the RapidAIM in vitro culturing and metaproteomic sample processing workflows, during which process, J.M., K.W., M.H. and H.Q. assisted with microbiome culturing and microbial cell washing. L.L. and Xin Z. performed the functional redundancy analyses. Z.N. optimized LC-MS/MS parameters, both Z.N. and L.L. performed the metaproteomic LC-MS/MS analysis and tracked LC-MS/MS performance. L.L., K.C. and C.M.A.S. performed metaproteomic database search. J.B. and A.S. performed metagenomic sample processing and sent samples for Illumina NovaSeq analysis. J.M.S. performed metagenomic data analysis and created metagenomics-based protein databases. L.L. and L.W. performed the PhyloFunc analysis. L.L. and C.M.A.S curated the data and performed TMT metaproteomics data analysis and compound structure clustering. L.L. performed all label-free metaproteomics down-stream data analysis. All authors contributed to the writing, editing, and proofreading of the manuscript.

1107

COMPETING INTERESTS STATEMENT

D.F. and A.S. are co-founders of MedBiome Inc., a microbiome nutrition and therapeutic company. C.M.A.S. is currently a staff at Recursion Pharmaceuticals, and K.W. is currently a staff at Carleton University, the presented contribution in this manuscript was made during their employment at the University of Ottawa. Other co-authors declare no conflict of interest.