1

Title: Nano-organization of synapses defines synaptic release properties at cortical neuron dendritic spines

Authors: Haani Jafri[1], Samantha J. Thomas[1], Sung Hoon Yang[1], Rachel E. Cain[1], Matthew B Dalva[1,2]

1: Department of Neuroscience and Jefferson Synaptic Biology Center, Sidney Kimmel Medical College at Thomas Jefferson University, 233 South 10[th] Street, Bluemle Life Sciences Building, Room 324, Philadelphia, PA 19107, USA

2: Tulane Brain Institute, Department of Cell and Molecular Biology, Tulane University, 202 Flower Hall, 6823 St. Charles Avenue, New Orleans, LA 70124, USA

*Author correspondence:

Matthew B. Dalva

Email: mdalva@tulane.edu

Abstract (150/150 words): Visualization of the submicron organization of excitatory synapses

has revealed an unexpectedly ordered architecture consisting of nanocolumns of synaptic

proteins that group into nanomodules which scale in number as spine size increases. How these

features are related to synaptic function has remained unclear. Here, using super-resolution

followed by live-cell line-scan imaging, we find that the size of the smallest miniature calcium

and glutamate events are the same, regardless of whether spines have one or two nanopuncta of

PSD-95, and that miniature synaptic response in all spines are best fit by a three term Poisson.

Two nanomodule spines exhibit more large events without a significant change in event

frequency, with the number of the largest events increasing disproportionately. These data

support a model where nanomodules define sites of synaptic release and where the

nanoarchitecture of synaptic proteins specifies subtypes of excitatory synapses, with increasing

numbers of nanomodules increasing coordinated multivesicular release.

Introduction

Synaptic connections underlie neuronal communication and memory in the nervous system and are implicated in many neurological diseases. Synaptic size varies and is linked to the strength of postsynaptic responses, with increasing spine size associated with increased synaptic strength[1,2]. The nanoscale organization of the synapse is also dynamic and linked to the size of the dendritic spine. The functional components of excitatory synapses – including ionotropic and metabotropic glutamate receptors, scaffolding proteins, and active zone proteins– organize into nanoclusters that align across the synaptic cleft forming trans-synaptic nanocolumns which optimize neurotransmission [3-5]. These nanocolumns scale in number, but not size, as spine size increases, suggesting that they are modular structures ('nanomodules') that may relate to the function of the spine [6]. Consistent with this model, following structural plasticity, which enhances synaptic strength and drives spine enlargement, the number of nanomodules increases in spines that undergo sustained increases in size[6,7]. This suggests that the nanoscale features of spines may specify differences in how synapses at larger and smaller spines function. However, it remains unclear whether the number of nanomodules at a spine is related to synaptic function or transmission.

To determine the relationship between synaptic nanomodules and properties of synaptic function, the nanoscale features within individual spines must be related to their synaptic responses. Electrophysiology techniques have a limited ability to record from single spines, because of their small size, and the difficulty of locating the source of synaptic inputs from recordings in the soma or dendrite [8-11]. Additionally, voltage compartments, like the spines themselves, may alter signal propagation within the neuron [12]. To overcome these limitations, we conducted live-cell

STED nanoscopy followed by time-lapse imaging of single spines in neurons transfected with

the calcium or glutamate sensors, GCaMP8f and iGluSnFR3, respectively. We further simplified

the postsynaptic response by measuring amplitude and frequency of spontaneous miniature

postsynaptic calcium or glutamate events under conditions where action potentials are blocked.

Quantal analysis of these data indicates that one and two nanomodule spines both have the same

size single quantal events but larger two nanomodule spines have more high amplitude multi-

quantal events. These data suggest that the nanoscale organization of spines defines distinct

excitatory synapse subtypes with distinct patterns of quantal spontaneous release.

Results:

Observed miniature synaptic Ca^{2+} transients are NMDAR-dependent

Larger spines generate larger postsynaptic responses [1,2]. However, how the nanoscale

organization of the spine relates to single, spontaneous synaptic events has yet to be determined.

To determine this relationship, we imaged the postsynaptic activity of spines with known

nanostructure by co-transfecting neurons with mTurq2-tagged PSD95-FingR and GCaMP8f,

which enabled the identification of dendritic spines with either one or two nanoclusters of PSD-

95 using STED imaging (Fig. 1a, b, Supplementary Fig. 1a-c). The PSD95-FingR nanostructure

of the spine was imaged using live-cell gated-STED and then the same area of dendrite was

imaged with live-cell time-lapse confocal imaging (83 Hz, 64 x 64 pixels, 406 nm x 406nm pixel

size) of single spines. STED imaging was reserved for nanomodule identification to reduce

photodamage to neurons[13]. To visualize the relationship of PSD-95 nanostructure to neuronal

activity, spontaneous NMDAR-dependent calcium transients were analyzed, which ensured that

the post synaptic responses were not saturated. To isolate NMDAR activity, a cocktail of drugs

was used to block non-specific sources of calcium, including activity-dependent potentials,

intracellular calcium stores, AMPA-type glutamate receptors and metabotropic glutamate

receptors (2 μm TTX, 40 μm nifedipine, 30 μm dantrolene, 10 μm LY341495, 20 μm NBQX,

and 500 μm S-MCPG), allowing for NMDAR-dependent miniature synaptic calcium transients

(mSCTs) to be quantified[14-16]. With these blockers in place, spontaneous activity does not

saturate the NMDAR response, permitting quantification of NMDAR responses under

magnesium-free conditions[17-19]. For imaging of glutamate, only action potentials were blocked.

Calcium imaging revealed transient increases in fluorescence limited to the spine head (Fig. 1c,

d, Movies S1, S2). Analysis of these calcium transients revealed a distinctive profile

characterized by a rapid rising phase, a slower decay phase, and a subsecond time course,

indicative of spontaneous NMDAR events[16,20-22] (Fig. 1e, f). To confirm the NMDAR-mediated

dependence of these events, neurons were treated with the selective NMDAR-antagonist (2R)-

amino-5-phosphonovaleric acid (100 μm AP5). AP5 treatment blocked all fluorescence

transients without changing the baseline fluorescence (Fig. 1g-m, Movies S3, S4). These data

indicated that the recorded events were due to the activity-independent activation of NMDARs,

likely via spontaneous vesicle fusion represented as mSCTs.

Miniature NMDAR-dependent synaptic transients exhibit a time course of approximately 200

milliseconds[23]. To determine the sensitivity of single spine imaging of GCaMP, the kinetics of

mSCTs were recorded using different calcium sensors, GCaMP7f and GCaMP8f using 83 Hz 2D

field scanning. (GCaMP6f did not resolve these events well, (data not shown), but see[16].

GCaMP8f event kinetics were also recorded at either 83 Hz 2D field scanning and 4000 Hz 1D

line scanning (Supplementary Fig. 2a-c). Consistent with design of GCaMP8f[24], mSCTs

recorded using GCaMP8f exhibited significantly faster rise times, showed significantly narrower

full width at half maximum amplitude (FWHM), and had similar decay times as those recorded

with GCaMP7f (Supplementary Fig. 2e, f). Moreover, line scans of GCaMP8f mSCTs

demonstrated significantly faster rise times compared to 2D-scanned mSCTs (Supplementary

Fig. 2d) without significant bleaching (Supplementary Fig. 3a), enabling visualization of the

~200ms time course of NMDAR transients. Therefore, we used an approach combining STED

imaging of PSD95-FingR with line scans of single spines filled with GCaMP8f to best capture

spontaneous calcium events from spines with identified nanoscale organization.

Visualization of mSCTs in individual spines reveals quantal event distributions

Larger spines generate larger postsynaptic responses and have more nanoclusters of pre- and

postsynaptic proteins[6,7]. These clusters of pre- and postsynaptic proteins form in trans-

synaptically paired nanomodules, or aligned nanocolumns, suggesting that synaptic nanomodules

might be related to synaptic function[3,4,25]. The excitatory synapses found at dendritic spines

transmit information by releasing glutamate from the presynaptic terminal, activating

postsynaptic glutamate receptors. There are two ways that increasing the number of

nanocolumns or nanomodules might alter the postsynaptic response. If the release of a single

vesicle activates all postsynaptic nanopuncta, the size of the postsynaptic response should

increase as more nanomodules are added. Alternatively, if a single vesicle can only activate the

proximal nanopuncta, then the size of the postsynaptic response would not change with

increasing number of nanomodules. These possibilities are explored below.

To explore the number of vesicles at spines with known nanoarchitecture, the nanoscale

organization of spines was first determined by STED imaging, and then 4000 Hz line scan

imaging of NMDAR-dependent mSCTs in single spines with known nanoscale organization was

conducted (Fig. 2a). Spines expressing FingR were identified, and STED was used to determine

whether they contained one or two PSD-95 nanomodules (Fig. 2b, d, l, n). These same spines

were imaged for 10 or 20 minutes at 4000 Hz, where unitary mSCTs were both clearly visible

(Fig. 2c, e, m, o, Movies S5-S8) and quantifiable (Fig. 2f, i, p, s) from the line scans. Only one

and two nanopuncta spines were selected for 4000 Hz imaging because spines with larger

numbers of nanopuncta are rare[6] (<15% of all spines).

The frequency of spontaneous events varied from spine to spine, but there was no difference in

mean frequency between one and two nanomodule spines (Supplementary Fig. 4a). Therefore,

we asked whether there were differences in events between spines sharing the same PSD-95

nano-organization. The amplitude of mSCTs in all spines, regardless of frequencies, exhibited

positively skewed mSCT amplitude histograms, irrespective of the PSD-95 nanomodule number

(lower frequency: Fig. 2g, j; higher frequency: Fig. 2r, u). Lower frequency spines with fewer

events exhibited event amplitudes of similar sizes, resulting in a narrow distribution with a low

standard deviation, consistent with single quantum events (Fig. 2h, k). Conversely, higher

frequency spines with more events displayed event amplitudes that fell into multiple, distinct,

evenly spaced ranges that increased progressively in size in a pattern reminiscent of multi-

quantal release [26-28] (Fig. 2q, t). There was a weak negative correlation between mean mSCT

frequency and mean amplitude in one nanomodule spines, while this negative correlation was not

exhibited in two nanomodule spines (Supplementary Fig. 4b, c). Therefore, we hypothesized that

spontaneous activity in both one and two nanomodule spines abides by quantal release

characteristics. One limitation of this approach is that the spontaneous presynaptic response is

being integrated using the activation of postsynaptic NMDAR, raising the possibility that the

distribution of NMDAR mSCT amplitudes simply reflects the number of postsynaptic NMDARs

being activated [29]. Therefore, an approach was developed to measure the amount of glutamate

released at synapses with identified nanoarchitecture.

Vesicular glutamate release is quantal in both one and two nanomodule spines

To determine whether the quantal distribution of mSCTs depended on pre- or postsynaptic

release characteristics, cortical neurons were transfected with the postsynaptically-targeted

glutamate sensor, iGluSnFR3-SGZ[30] (iGluSnFR3), along with PSD95-FingR to determine the

relationship between presynaptic glutamate release and the nanoscale organization of PSD-95 at

dendritic spines. One and two nanomodule spines were identified using STED (Fig. 3a, c, k, m).

Miniature glutamate transients (mGluTs) were seen visually (Fig. 3b, d, l, n, Movies S9-S12) and

quantifiably (Fig. 3e, h, o, r) from line scans. Line scan imaging of single spines resulted in a

slightly higher rate of bleaching than GCaMP8f (4.6% mean reduction in raw fluorescence,

Supplementary Fig. 3b), but small events were easily detectable throughout the imaging period.

As seen with GCaMP8f, spines exhibited a range of event frequencies but there was no

difference in mean frequency between one and two nanopuncta spines (Supplementary Fig. 4d).

In spines with lower levels of activity, the distributions of mGluT amplitudes exhibited positive

skew, irrespective of the number of PSD-95 nanomodules present. Notably, events from lower

frequency spines were distributed within a narrow range (Fig. 3f, i), resulting in low standard

deviation amplitude distributions (Fig. 3g, j) consistent with the release of a single quantum. In

higher frequency spines, the mGluT amplitude distributions displayed a positive skew (Fig. 3q,

t), consistent with multiple quantal events. Similar to mSCTs recorded using GCaMP8f, there

was a weak negative correlation between mGluT frequency and amplitude in one nanomodule

spines, which was reduced in two nanomodule spines (Supplementary Fig. 4e, f). Thus, findings

in neurons transfected with iGluSnFR3 were consistent with results using GCaMP8f, suggesting

that visualizing either miniature glutamate or NMDAR-dependent calcium events from single

spines enables us to visualize spontaneous synaptic events at individual spines.

One and two nanomodule spontaneous activity exhibit no kinetic differences

We determined the event kinetics of miniature events, including rise times, decay times, and

FWHM in one and two nanomodule spines. We hypothesized that there would be no differences

in miniature event kinetics between one and two nanomodule spines, consistent with a model

where nanomodules represent unitary sites of synaptic transmission. To test whether event

kinetics were independent of nanomodule number, the rise times, decay times, FWHM time

courses were calculated. There was no difference in these parameters of mSCTs and mGluTs

between one and two nanomodule spines (Supplementary Fig. 5c-e, h-j). Likely due to the

buffering of intracellular calcium, which increases with respect to spine volume[16], when mSCTs

were visualized in two nanomodules spines, their apparent peak amplitudes were less than mSCT

amplitudes in one nanomodule spines, (Supplementary Fig. 5a, b).The event amplitude of

mGluTs was not different in one or two nanomodule spines (Supplementary Fig. 5f, g) due to

iGluSnFR3 being expressed on the postsynaptic surface. All of the recorded mGluTs and mSCTs

were scaled to the same size between nanomodule groups to further visualize differences

between waveforms from one and two nanomodule spines. There was no difference in the

waveforms of scaled events (Supplementary Fig. 6a-j), consistent with a model where the

kinetics of mSCTs or mGluTs events do not depend on nanomodule number. This suggested

nanoclusters of PSD-95 may reflect the number of independent release sites at a synapse.

Quantal analysis of individual spines resolves quantal release characteristics

To test whether the nanoscale organization of spines relates to synaptic function, distributions of

mSCT and mGluT peak amplitudes at single dendritic spines were examined. In spines with

larger numbers of events, the distribution of event size appeared to segregate into discrete,

evenly spaced amplitude bins. These data were consistent with the model that synaptic events

represented the quantal fusion of synaptic vesicles[31]. However, single spines could not be imaged

long enough to visualize a sufficient number of events for quantal analysis. Therefore, we sought

an approach to increase the number of events recorded. Since the kinetics of single events were

the same in all spines, events recorded from all single spines in neurons expressing either

GCaMP8f or iGluSnFR3 were combined into a single mSCT or mGluT amplitude histogram,

respectively. Consistent with previous reports that did not account for the nanoscale organization

of spines[32], combining all events – irrespective of nanomodule number – showed that the event

data for both GCaMP8f and iGluSnFR3 events were well fit by both a skewed Gaussian

distribution or a multiple term Poisson distribution (Fig. 4a, b). However, further separating

spines by nanomodule number allowed for better visualization of quantal peaks in the combined

amplitude histograms (Supplementary Fig. 8a-d). Despite this improvement, differences in

baseline noise and quantal size between individual spines contributed to a smearing effect when

amplitude histograms were aggregated. To address this, quantal analysis was performed to

normalize these differences between spines and enhance visualization of quantal peaks.

Quantal analyses of each individual spine were performed as follows. The single quantal event

size in each spine was determined by defining a quantum as the minimal detectable amplitude of

the spontaneous events relative to the baseline noise[31] (x = 0, Fig. 4c). The size of the first

quantum was determined for each spine by averaging the absolute values between noise and the

first peaks $\bar{d}$, where quantal size m is determined as the $\Delta F/F$ fluorescence F divided by the sum

of highest amplitude of the noise per spine, N, and $\bar{d}$ ($m = F/(N + \bar{d})$, Fig. 4d-g, l-o,

Supplementary Fig. 8). In the resulting quantized mSCT and mGluT event histograms, the first

peak in each spine histogram was at or near 1 (Fig. 4h-k, p-s), indicating, as expected, that the

most frequent event for each synapse was a single quantum. These data are consistent with a

model where spontaneous excitatory synaptic transmission at single spines consists of miniature

synaptic events of a similar minimum size, regardless of the number of nanopuncta of PSD-95.

Nanoscale organization of spines reflects a functional difference in minisynaptic event

properties.

To determine whether spines with different numbers of nanomodules have functional

differences, the mSCT and mGluT events from one and two nanomodules spines were placed

into separate groups. The median mSCT and mGluT event size between one and two

nanomodule was not significantly different, which is consistent with one quantum events being

the most frequent event size (Fig. 5c, i). If nanomodules represent distinct release sites, spines

with increasing numbers of nanomodules should have more events with a larger number of

quanta[33,34]. Consistent with this model, in two nanomodule spines, both mSCT and mGluT event

amplitude cumulative probability distributions demonstrated significantly increased numbers of

larger events than in one nanomodule spines (Fig. 5d, j). These data suggest that increased

nanomodule number increases the number of synaptic release sites, linking the nanoscale

organization of synaptic proteins to synaptic transmission.

To test whether these increases in event size reflected larger quantal events or more quantal

release, Poisson distributions of mSCTs and mGluTs were determined for one or two

nanomodule spines. Events from both one and two nanomodule spines for mSCTs and mGluTs

were best fit by three quanta models (Fig. 5a, b, g, h). The three-term Poisson functions provided

a better fit to the data than skewed Gaussian functions in all cases using Akaike Information

Criterion (AIC) (Supplementary Fig. 9a, b, d, e). Additionally, the mSCT and mGluT amplitude

histograms from all spines were well fit by three-term Poisson functions (Supplementary Fig.

10a, b). The Poisson distribution of two nanomodule spine mSCT events (Fig. 5e) and mGluT

events (Fig. 5k) had fewer one- and two-quanta events and more three-quanta events than single

PSD-95 nanopuncta spines. Spines with two nanomodules had a 2.2-fold increase in three quanta

mSCTs and 25.3-fold increase in three quanta mGluTs when compared to one nanomodule

spines (Fig. 5f, l). The data from two nanomodule spines were best fit by a three-quanta model,

suggesting that the increased number of larger events was not due to the presence of larger

(four+) quanta in two nanomodule spines. The increased number of three quanta events is

consistent with the cooperativity of release, perhaps by reducing the energy barrier for vesicle

fusion[26,35]. Regardless, these data support a model where both one and two nanomodule spines

have multivesicular release, with an increasing number of nanopuncta leading to disproportionate

increases in three quanta multivesicular release. These data support a model whereby synapses

can modulate their pre- and postsynaptic signaling machinery to alter their strength.

Discussion

The majority of spines contain a single nanomodule of synaptic proteins with <35% of spines

containing two or more pre- and postsynaptic nanomodules[6]. The differences in nanoscale

organization suggest that there may be functional differences between spines of different sizes[6,7].

Using a combined super-resolution and live-cell imaging approach, we find that the size of the

smallest miniature calcium and glutamate events are the same, regardless of whether spines have

one or two nanopuncta of PSD-95, suggesting that nanopuncta of PSD-95 mark independent

synaptic release sites. The distributions of mSCT and mGluT amplitudes show similar quantal

patterns, with both single and two nanomodule spines fit by three quanta Poisson distributions.

Given that the fusion of single vesicles activates similar numbers of postsynaptic receptors, these

data support the model that the precise nanoscale feature of spines are likely to be determinative

for their function.

As spines become larger, they exhibit larger postsynaptic responses[1,2]. Previous work has

suggested that these increases in synaptic strength are due largely to postsynaptic changes, for

instance, in receptor number, subtype, and modulation[36-38]. Visualization of the mSCT and

mGluT events in spines with known nanoarchitecture supports an additional mechanism for the

increased size of postsynaptic responses in larger spines: coordination of multivesicular release.

Spines with two nanopuncta of PSD-95 have significantly more three quanta events than spines

with single nanopuncta, a striking increase in multiquanta events. Multivesicular release has been

shown in central synapses both in the cortex and in the hippocampus[26-28,39,40]. Moreover, live-cell

imaging of pre-and postsynaptic structures at the nanoscale reveals a coordinated increase in

both pre-and postsynaptic nanopuncta following structural plasticity[6]. These data support the

model that dendritic spine size is related to synaptic strength and the nanoscale organization of

synaptic proteins. Taken together, these data suggest that each nanopuncta acts independently,

and increases synaptic strength in larger spines may reflect a change in both presynaptic release

machinery and postsynaptic receptors that is reflected in the nanoscale organization of the

synapse.

Central nervous system synapses have long been considered to function differently from those in

the periphery and to consist of only a single release site[41-45]. Our data suggest that nearly 35% of

spines that contain more than one nanomodule of PSD-95 exhibit similarities to peripheral

synapses and may contain multiple release sites. One challenge to the quantal neuromuscular

junction model in the CNS has been the observation of a skewed distribution of miniature

synaptic event amplitudes[32]. By segregating miniature synaptic responses by nanoarchitecture,

the quantal characteristics of CNS synapses become apparent and suggest that nanostructural

features are key functional units of spine synapses. In large spines with higher nanomodule

number, there are more multivesicular events consistent with a model of a synapse that is

functionally similar to peripheral synapses at the neuromuscular junction, where cortical

nanocolumns resemble the multiple discrete active zones within lower motor neuron boutons[46].

Consistent with multivesicular release, EM analysis of CNS synapses has shown multiple

endocytic pits [47]. Our data provides a model of the central synapse; rather than a single release

site that undergoes primarily postsynaptic modifications, we propose that the as the number of

nanomodules of spine synapses increase there is an increase in coordinated vesicle release,

making the classification of synapses based on their nanostructural features a necessary method

to compare functional differences between spine synapses.

Funding: Support by funding from National Institutes of Health grants F30NS130973 (REC),

R01DA022727 (MBD), R01NS111976 (MBD), R01NS115441 (MBD)

Author contributions: Conceptualization: HJ and MBD; Methodology: HJ, SJT, SHY, REC,

MBD; Investigation: HJ; Visualization: HJ, SJT; Funding acquisition: MBD; Supervision: MBD;

Writing: HJ, SJT, MBD.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary

materials. Analysis tools and imaging data available on request.

321
322

 Figures:

 Figure 1: STED and Ca2+ imaging measurement of NMDAR-dependent events.

(**a,b**) Schematic of live-cell STED nanoscopy of exemplary one (left) and two (right)

nanomodule spines with live-cell STED resolved PSD-95 FingR (magenta) and GCaMP8f as a

cell-fill (green). White dashed line shows morphology. Scale bar is 1µm. (**c,d**) The same one and

two nanomodule spines at three different time points of a Ca^{2+}-transient recorded under activity

blockade: baseline before peak at t = 0ms (left panel), peak amplitude t = 612ms (middle panel),

baseline after peak at t = 1500ms (right panel). Spine outlines (white dashed lines) are based on

high pixel format imaging using STED. (**e,f**) Quantification of the same events with the time

points labeled (red dashed lines). (**g-j**) Images and activity traces from the same spines treated

with APV, demonstrating sustained baseline fluorescence. (**k**) Quantification of the frequency of

spontaneous Ca^{2+}-transients in individual dendritic spines before and after APV addition. N = 29

spines (**** $p < 0.0001$, paired t-test). (**l**) $\Delta F/F$ normalized fluorescence intensity trace of a

dendritic spine under non-NMDAR activity blockade showing five minutes of spontaneous Ca^{2+}-

transient recording. (**m**) Same spine as shown in (l) following the addition of APV.

Figure 2: Quantification of mSCTs from individual dendritic spines.

(a) Model depicting the two-step process of identifying nanomodule number in spines using

STED nanoscopy, followed by drawing a line through selected spines and corresponding parent

dendrite to record mSCT activity. One nanomodule **(b, l)** or two nanomodule **(d, n)** spines with

GCaMP8f as a cell-fill (green) imaged with confocal, STED resolved PSD-95-FingR puncta

(magenta, solid white arrow) and the dashed line indicating the position of the line scan used for

high temporal resolution imaging. All scale bars for morphology are 1 μm. **(c, e, m, o)**

Representative line-scans of the respective spines (b, d, l, n) with an mSCT shown (arrowhead).

The spine and parent dendrite are marked (S = spine, D = dendrite). All scale bars for line scans

are x = 1 μm and t = 200 ms. **(f, i, p, s)** Quantification of 30 seconds of recording from each

respective spine with the corresponding event from representative line scans labeled (white

arrow). **(g, j, p, t)** Quantification of every event recorded from each respective spine. **(h, k, r, u)**

Amplitude histograms depicting all the recorded events from one nanomodule (red) and two

nanomodule (blue) spines with 8000 points of baseline noise (black, inset). Models in (a)

generated using BioRender.

Figure 3: Quantification of mGluTs from individual dendritic spines.

One nanomodule **(a, k)** or two nanomodule **(c, m)** spines with iGluSnFR3 as a cell-fill (green)

imaged with confocal, a STED resolved PSD-95 FingR puncta (magenta, solid white arrow), and

a dashed line indicating the position of the line scan used for high temporal resolution imaging.

Scale bar is 1 μm. **(b, d, l, n)** Representative line-scans of the corresponding spines (a, c, k, m)

with an mGluT shown (solid black arrow). The spine and parent dendrite are marked (S = spine,

D = dendrite). Scale bar is x = 1 μm and t = 200 ms. **(e, h, o ,r)** Quantification of 30 seconds of

recording from each respective spine with the corresponding event from the representative line-

scan labeled (white arrow). **(f, i, p, s)** Quantification of every event recorded from each

respective spine. **(g, j, q, t)** Peak amplitude histogram depicting all the recorded events from the

367 respective one nanomodule (green) or two nanomodule (magenta) spines with 8000 points of

368 baseline noise (black, inset).

410
411
Figure 4: Noise characteristics of spines reveals non-Gaussian amplitude distribution.

(a) Raw peak amplitude histogram from all spines recorded using GCaMP8f, with a skewed

Gaussian distribution (red line) and a three quanta Poisson fit model (black line). N = 25 spines,

2164 events; Gaussian R^2 = 0.889, AIC = -509.96, three quanta Poisson R^2 = 0.976, AIC = -

594.67. **(b)** Raw peak amplitude histogram from all spines recorded using iGluSnFR3-SGZ with

a skewed Gaussian distribution (red line) and a three quanta Poisson fit model (black line). N =

23 spines, 799 events; Gaussian R^2 = 0.934, AIC = -374.36, three quanta Poisson R^2 = 0.977,

AIC = -415.71. **(c)** Model depicting the process by which quantal analysis was performed,

accounting for individual noise and quantal size variation of spines. **(d, e)** Examples of one

nanomodule spines expressing GCaMP8f showing the PSD-95 puncta in magenta and spine as

an outline (d) or in green (e). **(f)** Frequency distribution of event sizes from the spine (d, e) with

recorded events (red) and noise (black) shown as histograms. **(g)** Quantized events from spine in

d-e. Difference between noise and quantal size is also shown (red line, d_1, d_2). **(f, g)** Same as in

d, **e** but for two nanomodule spines with recorded events (blue) and noise (black). **(h-k)** As in **d-**

g but showing a spine with two PSD-95 puncta. (**l**, **m**) Examples of a one nanomodule spine as in

d, e but expressing iGluSnFR3. (**n, o**) Histograms are of recorded iGluSnFR3 events (green) and

noise (black) from the spine shown. **(p-s)** Same as in **l-o** but shows a two nanomodule spine.

Models in **c** generated using BioRender. Scale bars = 1 µm.

Figure 5: Comparing event amplitudes between populations of one and two nanomodule

spines reveals quantal differences.

(**a**) Quantized mSCT peak amplitude histogram from a population of one nanomodule GCaMP8f

spines. Three quanta fit model is shown (black line, $R^2 = 0.962$). Individual components of the

function are shown in gray. N = 901 events across 11 spines. (**b**) Same as left (a), but for a

population of two nanomodule spines (black line, three term Poisson $R^2 = 0.927$). N = 1263

events across 14 spines. (**c**) Violin plots of quantized mSCT amplitudes from one nanomodule

(red) and two nanomodule (blue) spines. Black dashed line represents quartiles. (**d**) Cumulative

probability distributions of one nanomodule (red line) and two nanomodule (blue line) spines

with respect to peak mSCT amplitudes (***p = 0.0004, by K-S test). (**e**) Individual quantal terms

from Poisson distributions of one nanomodule (red line) and two nanomodule (blue line)

GCaMP8f spines from above (a,b) overlaid upon each other. (**f**) The integrated area under the

curve (AUC) calculated from each quantal term in (e) with the $\log_2$-fold change between the

AUC of two nanomodule spines divided by one nanomodule spines (inset, fold change = (AUC

two NM spines/AUC one NM spines)) (**g,h**) Quantized mGluT peak amplitude histograms from

populations of one (g) and two (h) nanomodule iGluSnFR3 spines. Three quanta fit model is

shown (black line, one nanomodule $R^2 = 0.848$, two nanomodule $R^2 = 0.923$) with individual

components of the respective Poisson functions (gray). N = 222 events across 8 one nanomodule

spines, N = 577 events across 15 two nanomodule spines. (**i**) Violin plots of quantized mGluT

amplitudes from one nanomodule (green) and two nanomodule (magenta) spines. Black dashed

line represents quartiles. (**j**) Cumulative probability distributions of one nanomodule (green line)

and two nanomodule (magenta line) spines with respect to peak mSCT amplitudes (*p = 0.0308,

by K-S test). (**k**) Individual quantal terms from the Poisson distributions of one nanomodule

(green line) and two nanomodule (magenta line) iGluSnfR3 spines from above (f, g) overlaid

upon each other. **(l)** Same integrated AUC as in (f) except based on mGluT data in (k).

458
459   **References:**

460   1   Segal, M. Dendritic spines and long-term plasticity. *Nat Rev Neurosci* **6**, 277-284 (2005).
461   https://doi.org/10.1038/nrn1649
462   2   Borczyk, M., Śliwińska, M. A., Caly, A., Bernas, T. & Radwanska, K. Neuronal
463   plasticity affects correlation between the size of dendritic spine and its postsynaptic
464   density. *Sci Rep* **9**, 1693 (2019). https://doi.org/10.1038/s41598-018-38412-7
465   3   Tang, A. H. *et al.* A trans-synaptic nanocolumn aligns neurotransmitter release to
466   receptors. *Nature* **536**, 210-214 (2016). https://doi.org/10.1038/nature19058
467   4   Biederer, T., Kaeser, P. S. & Blanpied, T. A. Transcellular Nanoalignment of Synaptic
468   Function. *Neuron* **96**, 680-696 (2017). https://doi.org/10.1016/j.neuron.2017.10.006
469   5   Goncalves, J. *et al.* Nanoscale co-organization and coactivation of AMPAR, NMDAR,
470   and mGluR at excitatory synapses. *Proc Natl Acad Sci U S A* **117**, 14503-14511 (2020).
471   https://doi.org/10.1073/pnas.1922563117
472   6   Hruska, M., Henderson, N., Le Marchand, S. J., Jafri, H. & Dalva, M. B. Synaptic
473   nanomodules underlie the organization and plasticity of spine synapses. *Nat Neurosci* **21**,
474   671-682 (2018). https://doi.org/10.1038/s41593-018-0138-9
475   7   Hruska, M., Cain, R. E. & Dalva, M. B. Nanoscale rules governing the organization of
476   glutamate receptors in spine synapses are subunit specific. *Nat Commun* **13**, 920 (2022).
477   https://doi.org/10.1038/s41467-022-28504-4
478   8   Beaulieu-Laroche, L. *et al.* Enhanced Dendritic Compartmentalization in Human Cortical
479   Neurons. *Cell* **175**, 643-651.e614 (2018). https://doi.org/10.1016/j.cell.2018.08.045
480   9   Spruston, N., Jaffe, D. B., Williams, S. H. & Johnston, D. Voltage- and space-clamp
481   errors associated with the measurement of electrotonically remote synaptic events. *J*
482   *Neurophysiol* **70**, 781-802 (1993). https://doi.org/10.1152/jn.1993.70.2.781
483   10   Williams, S. R. & Mitchell, S. J. Direct measurement of somatic voltage clamp errors in
484   central neurons. *Nat Neurosci* **11**, 790-798 (2008). https://doi.org/10.1038/nn.2137
485   11   Bar-Yehuda, D. & Korngreen, A. Space-clamp problems when voltage clamping neurons
486   expressing voltage-gated conductances. *J Neurophysiol* **99**, 1127-1136 (2008).
487   https://doi.org/10.1152/jn.01232.2007
488   12   Cornejo, V. H., Ofer, N. & Yuste, R. Voltage compartmentalization in dendritic spines in
489   vivo. *Science* **375**, 82-86 (2022). https://doi.org/10.1126/science.abg0501
490   13   Laissue, P. P., Alghamdi, R. A., Tomancak, P., Reynaud, E. G. & Shroff, H. Assessing
491   phototoxicity in live fluorescence imaging. *Nat Methods* **14**, 657-661 (2017).
492   https://doi.org/10.1038/nmeth.4344
493   14   Reese, A. L. & Kavalali, E. T. Spontaneous neurotransmission signals through store-
494   driven Ca(2+) transients to maintain synaptic homeostasis. *Elife* **4** (2015).
495   https://doi.org/10.7554/eLife.09262
496   15   Higley, M. J. & Sabatini, B. L. Calcium signaling in dendritic spines. *Cold Spring Harb*
497   *Perspect Biol* **4**, a005686 (2012). https://doi.org/10.1101/cshperspect.a005686
498   16   Metzbower, S. R., Joo, Y., Benavides, D. R. & Blanpied, T. A. Properties of Individual
499   Hippocampal Synapses Influencing NMDA-Receptor Activation by Spontaneous
500   Neurotransmission. *eNeuro* **6** (2019). https://doi.org/10.1523/ENEURO.0419-18.2019

17   Mainen, Z. F., Malinow, R. & Svoboda, K. Synaptic calcium transients in single spines
indicate that NMDA receptors are not saturated. *Nature* **399**, 151-155 (1999).
https://doi.org/10.1038/20187
18   Higley, M. J. & Sabatini, B. L. Calcium signaling in dendrites and spines: practical and
functional considerations. *Neuron* **59**, 902-913 (2008).
https://doi.org/10.1016/j.neuron.2008.08.020
19   Kerlin, A. *et al.* Functional clustering of dendritic activity during decision-making. *Elife*
8 (2019). https://doi.org/10.7554/eLife.46966
20   Popescu, G., Robert, A., Howe, J. R. & Auerbach, A. Reaction mechanism determines
NMDA receptor response to repetitive stimulation. *Nature* **430**, 790-793 (2004).
https://doi.org/10.1038/nature02775
21   Murthy, V. N., Sejnowski, T. J. & Stevens, C. F. Dynamics of dendritic calcium
transients evoked by quantal release at excitatory hippocampal synapses. *Proc Natl Acad*
Sci U S A **97**, 901-906 (2000). https://doi.org/10.1073/pnas.97.2.901
22   Lester, R. A., Clements, J. D., Westbrook, G. L. & Jahr, C. E. Channel kinetics determine
the time course of NMDA receptor-mediated synaptic currents. *Nature* **346**, 565-567
(1990). https://doi.org/10.1038/346565a0
23   Sabatini, B. L., Oertner, T. G. & Svoboda, K. The life cycle of Ca(2+) ions in dendritic
spines. *Neuron* **33**, 439-452 (2002). https://doi.org/10.1016/s0896-6273(02)00573-1
24   Zhang, Y. *et al.* Fast and sensitive GCaMP calcium indicators for imaging neural
populations. *Nature* **615**, 884-891 (2023). https://doi.org/10.1038/s41586-023-05828-9
25   Guzikowski, N. J. & Kavalali, E. T. Nano-Organization at the Synapse: Segregation of
Distinct Forms of Neurotransmission. *Front Synaptic Neurosci* **13**, 796498 (2021).
https://doi.org/10.3389/fnsyn.2021.796498
26   Dürst, C. D. *et al.* Vesicular release probability sets the strength of individual Schaffer
collateral synapses. *Nat Commun* **13**, 6126 (2022). https://doi.org/10.1038/s41467-022-
33565-6
27   Soares, C., Trotter, D., Longtin, A., Béïque, J. C. & Naud, R. Parsing Out the Variability
of Transmission at Central Synapses Using Optical Quantal Analysis. *Front Synaptic*
Neurosci **11**, 22 (2019). https://doi.org/10.3389/fnsyn.2019.00022
28   Farsi, Z., Walde, M., Klementowicz, A. E., Paraskevopoulou, F. & Woehler, A. Single
synapse glutamate imaging reveals multiple levels of release mode regulation in
mammalian synapses. *iScience* **24**, 101909 (2021).
https://doi.org/10.1016/j.isci.2020.101909
29   Nimchinsky, E. A., Yasuda, R., Oertner, T. G. & Svoboda, K. The number of glutamate
receptors opened by synaptic stimulation in single hippocampal spines. *J Neurosci* **24**,
2054-2064 (2004). https://doi.org/10.1523/jneurosci.5066-03.2004
30   Aggarwal, A. *et al.* Glutamate indicators with improved activation kinetics and
localization for imaging synaptic transmission. *Nat Methods* **20**, 925-934 (2023).
https://doi.org/10.1038/s41592-023-01863-6
31   Del Castillo, J. & Katz, B. Quantal components of the end-plate potential. *J Physiol* **124**,
560-573 (1954). https://doi.org/10.1113/jphysiol.1954.sp005129
32   Wang, C. S., Chanaday, N. L., Monteggia, L. M. & Kavalali, E. T. Probing the
segregation of evoked and spontaneous neurotransmission via photobleaching and
recovery of a fluorescent glutamate sensor. *Elife* **11** (2022).
https://doi.org/10.7554/eLife.76008

547	33	Holderith, N. *et al.* Release probability of hippocampal glutamatergic terminals scales
548		with the size of the active zone. *Nat Neurosci* **15**, 988-997 (2012).
549		https://doi.org/10.1038/nn.3137
550	34	Körber, C. & Kuner, T. Molecular Machines Regulating the Release Probability of
551		Synaptic Vesicles at the Active Zone. *Front Synaptic Neurosci* **8**, 5 (2016).
552		https://doi.org/10.3389/fnsyn.2016.00005
553	35	Schotten, S. *et al.* Additive effects on the energy barrier for synaptic vesicle fusion cause
554		supralinear effects on the vesicle fusion rate. *Elife* **4**, e05531 (2015).
555		https://doi.org/10.7554/eLife.05531
556	36	Kauer, J. A., Malenka, R. C. & Nicoll, R. A. A persistent postsynaptic modification
557		mediates long-term potentiation in the hippocampus. *Neuron* **1**, 911-917 (1988).
558		https://doi.org/10.1016/0896-6273(88)90148-1
559	37	Shi, S., Hayashi, Y., Esteban, J. A. & Malinow, R. Subunit-specific rules governing
560		AMPA receptor trafficking to synapses in hippocampal pyramidal neurons. *Cell* **105**,
561		331-343 (2001). https://doi.org/10.1016/s0092-8674(01)00321-x
562	38	Barria, A., Muller, D., Derkach, V., Griffith, L. C. & Soderling, T. R. Regulatory
563		phosphorylation of AMPA-type glutamate receptors by CaM-KII during long-term
564		potentiation. *Science* **276**, 2042-2045 (1997).
565		https://doi.org/10.1126/science.276.5321.2042
566	39	Mendonça, P. R. F. *et al.* Asynchronous glutamate release is enhanced in low release
567		efficacy synapses and dispersed across the active zone. *Nat Commun* **13**, 3497 (2022).
568		https://doi.org/10.1038/s41467-022-31070-4
569	40	Rudolph, S., Tsai, M. C., von Gersdorff, H. & Wadiche, J. I. The ubiquitous nature of
570		multivesicular release. *Trends Neurosci* **38**, 428-438 (2015).
571		https://doi.org/10.1016/j.tins.2015.05.008
572	41	Silver, R. A., Lubke, J., Sakmann, B. & Feldmeyer, D. High-probability uniquantal
573		transmission at excitatory synapses in barrel cortex. *Science* **302**, 1981-1984 (2003).
574		https://doi.org/10.1126/science.1087160
575	42	Korn, H., Triller, A., Mallet, A. & Faber, D. S. Fluctuating responses at a central synapse:
576		n of binomial fit predicts number of stained presynaptic boutons. *Science* **213**, 898-901
577		(1981). https://doi.org/10.1126/science.6266015
578	43	Gulyás, A. I. *et al.* Hippocampal pyramidal cells excite inhibitory neurons through a
579		single release site. *Nature* **366**, 683-687 (1993). https://doi.org/10.1038/366683a0
580	44	Biró, A. A., Holderith, N. B. & Nusser, Z. Quantal size is independent of the release
581		probability at hippocampal excitatory synapses. *J Neurosci* **25**, 223-232 (2005).
582		https://doi.org/10.1523/jneurosci.3688-04.2005
583	45	Redman, S. Quantal analysis of synaptic potentials in neurons of the central nervous
584		system. *Physiol Rev* **70**, 165-198 (1990). https://doi.org/10.1152/physrev.1990.70.1.165
585	46	Badawi, Y. & Nishimune, H. Presynaptic active zones of mammalian neuromuscular
586		junctions: Nanoarchitecture and selective impairments in aging. *Neurosci Res* **127**, 78-88
587		(2018). https://doi.org/10.1016/j.neures.2017.11.014
588	47	Kusick, G. F. *et al.* Synaptic vesicles transiently dock to refill release sites. *Nat Neurosci*
589		**23**, 1329-1338 (2020). https://doi.org/10.1038/s41593-020-00716-1
590		