Single-cell multiome and spatial profiling reveals pancreas cell type-specific gene regulatory programs driving type 1 diabetes progression

Rebecca Melton[1,*], Sara Jimenez[2,*], Weston Elison[1], Luca Tucciarone[3], Abigail Howell[3], Gaowei Wang[3], Denise Berti[3], Elisha Beebe[3], Michael Miller[4] , Chun Zeng[3], Carolyn McGrail[1], Kennedy VanderStel[3], Katha Korgaonkar[3], Ruth Elgamal[1], Hannah Mummey[5], Joshua Chiou[6], Emily Griffin[3], Irina Kusmartseva[7], Mark Atkinson[7], Sebastian Preissl[4,9], Fabian J. Theis[2,**], Maike Sander[3,8,10,**], Kyle J. Gaulton[3,**]

1. Biomedical sciences program, University of California San Diego, La Jolla CA
2. Department of Computational Health, Institute of Computational Biology, Helmholtz, Munich, Germany
3. Department of Pediatrics, University of California San Diego, La Jolla CA
4. Center for Epigenomics, University of California San Diego, La Jolla CA
5. Bioinformatics and systems biology program, University of California San Diego, La Jolla CA
6. Pfizer Research and Discovery, Pfizer Inc, Cambridge, MA
7. Department of Pathology, Immunology and Laboratory Medicine, University of Florida, Gainesville FL
8. Department of Cellular & Molecular Medicine, University of California San Diego, La Jolla CA
9. Institute of Experimental and Clinical Pharmacology and Toxicology, Faculty of Medicine, University of Freiburg, Freiburg, Germany
10. Max Delbrück Center for Molecular Medicine, Berlin, Germany

* Authors jointly contributed to this work
** Authors jointly supervised this work

Correspondence to: Kyle J Gaulton, kgaulton@ucsd.edu

Abstract

Cell type-specific regulatory programs that drive type 1 diabetes (T1D) in the pancreas are poorly understood. Here we performed single nucleus multiomics and spatial transcriptomics in up to 32 non-diabetic (ND), autoantibody-positive (AAB+), and T1D pancreas donors. Genomic profiles from 853,005 cells mapped to 12 pancreatic cell types, including multiple exocrine sub-types. Beta, acinar, and other cell types, and related cellular niches, had altered abundance and gene activity in T1D progression, including distinct pathways altered in AAB+ compared to T1D. We identified epigenomic drivers of gene activity in T1D and AAB+ which, combined with genetic association, revealed causal pathways of T1D risk including antigen presentation in beta cells. Finally, single cell and spatial profiles together revealed widespread changes in cell-cell signaling in T1D including signals affecting beta cell regulation. Overall, these results revealed drivers of T1D progression in the pancreas, which form the basis for therapeutic targets for disease prevention.

Introduction

Introduction

Type 1 diabetes (T1D) is a complex endocrine disorder characterized by autoimmune
destruction of beta cells in the pancreatic islets, leading to lifelong dependence on insulin
therapy. The destruction of beta cells in T1D is caused by interactions with multiple cell types in
and surrounding the islet microenvironment including infiltrating immune cells, other endocrine
cells, and endothelial cells[1–3]. Cell types in the pancreas outside the local islet environment,
such as exocrine acinar and ductal cells, are also increasingly implicated in T1D
pathogenesis[4,5]. Beta cells themselves likely contribute to the development of T1D as well
through response to environmental factors, external signaling to immune, beta, and other cell
types, and cellular survival[6]. The sequence of events in the pancreas that drives initiation of
beta cell autoimmunity and progression through stages of T1D, however, as well as the role of
each pancreatic cell type in these processes, remains poorly understood.

Seroconversion to autoantibody positivity (AAB+) against islet proteins (i.e. self-antigens)
precedes T1D onset in nearly all cases and is used as a clinical biomarker of T1D
progression[7,8]. Individuals at T1D diagnosis can present with differing number and type of
autoantibodies, which are associated with varying rates of disease incidence; for example, the
presence of a single islet AAB has a relatively low lifetime risk of T1D while individuals with
multiple AAB have disease rates over 90%[9–11]. As clinical presentation of T1D does not occur
until a large fraction of beta cells has been destroyed, there is a window of time between
seroconversion and T1D onset where disease processes can potentially be halted or reversed[7].
Even after onset of T1D, residual beta cell mass could potentially be modulated therapeutically
to restore metabolic function[12]. Defining changes in disease-relevant cell types across the
stages of T1D progression would both improve our understanding of the mechanisms of T1D as
well as reveal potential targets to prevent or reverse disease. Furthermore, an improved
understanding of key changes associated with progression would also help identify novel
biomarkers of T1D, which are particularly needed in the early stages of disease to identify
progressors and candidates for therapeutic intervention[13].

Single cell technology, the focus of this work, enables profiling of individual cells within the
pancreas[5,14]. Previous single cell studies of the pancreas in T1D have been limited in that they
focused primarily on gene expression profiling of dispersed cells[4,15], which does not provide
information on the spatial localization of cellular transcriptomes within the pancreas nor the

genomic elements driving changes in gene activity. Recent developments in spatial
transcriptomics enables profiling cells in their native location[16], which enables understanding cell
type-specific changes in the context of specific cellular neighborhoods and niches in the
pancreas. This is particularly important in the context of T1D which has extensive heterogeneity
in disease processes within the pancreas[17]. In addition, single cell epigenome profiling, for
example using snATAC-seq or single cell multiome (paired snRNA-seq+snATAC-seq), can
reveal transcriptional regulators, *cis*-regulatory elements (cREs), and gene regulatory networks
driving altered gene expression in T1D[5,14]. Critically, gene regulatory networks and cREs can be
intersected with T1D-associated variation to infer cell type-specific regulatory programs that
may play a causal role in driving disease[18,19].

Previous single cell studies have also been limited in the extent to which they have captured key
windows of T1D progression and pathogenesis[4,15]. Specifically, non-diabetic AAB+ donors in
these efforts were largely those with single glutamic acid decarboxylase (GAD) autoantibodies[4],
which have a relatively lower risk of developing T1D compared to multiple AAB+ donors and do
not reflect the full arc of progression to T1D[20]. Furthermore, many of the T1D donors in these
studies had long-standing T1D where disease processes are potentially more dormant, whereas
profiling donors who had more recently developed T1D may give greater insight into active
disease processes. Third, as these studies profiled purified islets, they offer more restricted
insight into genomic changes in cells outside of the islet microenvironment during T1D
progression, including in exocrine cells which are both altered in T1D as well as implicated
causally in the development of T1D[4,18,21].

In this study, we addressed these limitations by performing single cell gene expression and
epigenome profiling in whole pancreas from 32 non-diabetic, non-diabetic single and multiple
AAB+, recent-onset T1D, and long-standing T1D organ donors, as well as spatial
transcriptomics in a subset of non-diabetic and recent-onset T1D donors. We determined
changes in pancreatic cell type abundance, cellular pathways, gene regulatory networks, and
cell-cell signaling across these stages of T1D progression and pathogenesis and, using T1D
association data, identified pathways and gene networks that may play a causal role in the
development of T1D.

136
137 **Results**
138
A comprehensive, multimodal, spatially resolved map of pancreatic cell types

We selected whole pancreas samples from 32 donors in the nPOD biorepository including 11

non-diabetic (ND), 9 non-diabetic autoantibody positive (ND AAB+), and 12 T1D which we

separated into 7 recent onset (<1 year from diagnosis) and 5 longer duration (>5 years from

diagnosis) (**Supplementary Table 1**). Within the ND AAB+ group, most organ donors, by our

study design, had multiple autoantibodies (multiple ND AAB+). For all samples, we performed

single nucleus RNA-seq (snRNA-seq) and single nucleus ATAC-seq (snATAC-seq) assays and,

for eight of the samples, we performed single nucleus multiome (joint snRNA-seq and snATAC-

seq in the same nucleus) assays. In addition, for six of the samples, we performed spatial

transcriptomics assays using the CosMx Spatial Molecular Imager (**Figure 1A**).

After extensive barcode quality control and filtering steps, (**see Methods**), we performed

integration using Harmony[22] and clustered 276,906 gene expression profiles (**Figure 1B,**

Supplementary Figures 1-2). We annotated the resulting 18 clusters based on the expression

of known cell type marker genes which revealed 12 pancreatic cell types including exocrine

(acinar, ductal), endocrine (alpha, beta, delta), immune (T cell, B cell, macrophage, mast),

stellate, endothelial, and Schwann cells (**Figure 1B-C, Supplementary Table 2**). Cell type

clusters had broadly consistent representation across donors and donor characteristics

(**Supplementary Figures 2-3**). We aggregated expression profiles for all cells in the cell type

and derived normalized expression levels of each gene using transcripts-per-million (TPM),

which revealed, on average, 17,885 genes expressed (TPM>1) per cell type (**Supplementary**

Table 3). For each cell type, we further identified genes with expression levels specific

(FDR<.10) to the cell type which revealed both known as well as previously unreported sets of

genes with cell type-specific expression (**Supplementary Table 4**); for example, well-known

genes with expression highly specific to beta cells included *INS, IAPP* and *G6PC2* as well as

others with no currently known role in beta cell function (e.g., *PLCH2, NRG2, RBFOX3,*

MTUS2).

Several cell types displayed multiple sub-clusters representing both known cell sub-types, such

as active and quiescent stellate cells, blood vessel (BVEC) and lymphatic endothelial (LEC)

cells, and MUC5b+ ductal cells, as well as several potential sub-types of acinar cells (**Figure**

1B,C). As the genomic properties of these sub-types have not been completely described in

previous efforts, we derived sets of marker genes for each sub-type (**see Methods,**

Supplementary Table 4). For BVECs and LECs, in addition to reported marker genes *PLVAP*

(BVECs) and *FLT4* (LECs), we observed specific up-regulation of genes in each sub-type such

as *INHBB, BMP6, FCN3,* and *PCAT19* in BVECs and *EFNA5, COLEC12,* and *MYCT1* in LECs.

In *MUCB5+*-ductal cells*,* there was up-regulation of *ERN2, CYP2C18, MYO7B,* and *DMBT1*

compared to the primary sub-type of ductal cells*.* For acinar cells, the primary cluster, which we

annotated as 'basal' acinar cells, was enriched for genes and pathways involved in digestive

enzyme production and secretion. Other clusters included 'high-enzyme' acinar cells with higher

expression of enzymes such as chymotrypsin (*CTRB1/2),* trypsinogen (*PRSS1, PRSS2*), lipase

(*PNLIP*), carboxyl ester lipase (*CEL*), chymotrypsin-like elastase (*CELA3A/B*) and increased

oxidative phosphorylation and translation, 'signaling' acinar cells with increased signaling and

stress-response activity, and 'signaling/differentiation' acinar cells with increased signaling,

metallothionein (*MT1/MT2),* and identity and differentiation genes (*REG1A/B, PTF1A*) (**Figure**

1D).

To next characterize the spatial organization of pancreatic cell types, we performed RNA *in situ*

hybridization (ISH) of 1,010 genes with CosMx from a subset of donors including three ND and

three recent-onset T1D (**Supplementary Table 1, Supplementary Table 5**). We imaged a

total of 82.6M transcripts from 71 fields of view (FOV) in whole pancreas sections from three ND

(32 FOV) and three recent-onset T1D donors (39 FOV) (**Supplementary Figure 4A**) and

assigned transcripts to 392,248 cells overall (80 median genes, 200 median transcripts per cell),

using the CosMx default segmentation. We first performed unsupervised clustering of cellular

gene expression profiles, which revealed nine distinct clusters including exocrine, endocrine,

endothelial, immune and mast cells (**Supplementary Figure 4B**). We next mapped finer-

grained cell type annotations from the snRNA-seq atlas using moscot[23] (**Supplementary Figure**

4B,C), which revealed 14 total cell types and sub-types that were confirmed based on marker

gene expression (**Figure 1E**). Spatial neighborhood enrichment using squidpy[24] revealed

expected cell types clustering together including acinar sub-types, ductal sub-types, endocrine

cells (beta, alpha, delta), and connective cells (e.g., endothelial, immune, stellate) (**Figure 1F**).

Next, we sought to determine whether spatial neighborhoods form recurrent niches across the

pancreas, by defining niches involving a cell type using a gene-gene covariance matrix[25] in a

spatial neighborhood of 30 cells. We recovered six niches in total, characterized by cell type

abundance (**see Methods**), including three exocrine (acinar-enriched, ductal-enriched, and

MUC5b ductal-enriched) niches, one endocrine niche, one niche including both endocrine and

exocrine cells (endo-exo), and one niche consisting of connective cells (**Figure 1G**). To

characterize each niche, we identified spatially variable genes (Moran's I >0.2, p<.05) that
captured gene signatures specific to the niche (**Figure 1G**). In the acinar-enriched niche, marker
genes from the 'basal' and 'high-enzyme' cell types showed strong spatial clustering (*PRSS2,*
REG1A). In comparison, the ductal-enriched niche had more spatial association with 'signaling'
and 'signaling/differentiation' acinar cells (*MT1X, SOD2, MT2A*). Interestingly, in the *MUC5b*
ductal-enriched niche, spatially variable genes were strongly associated with immune
interactions (*HSPA1A, HLA-A,* and *B2M*). In addition, the endocrine niche had highly distinct
patterns which highlighted multiple endocrine-specific genes (e.g. *INS, GCG, SST,* and *IAPP*)
(**Figure 1G**).

Finally, we determined whether there were changes in abundance of cell types and sub-types in
T1D progression based on snRNA-seq data (**see Methods**). There was a significant decrease
(likelihood ratio test [LRT], FDR<.10) in beta cells (**Figure 1H, Supplementary Table 6**)
although we still observed residual beta cell proportion in T1D particularly in recent-onset
(ND=1.5%, recent-onset T1D=0.93%). We also observed significant decrease (FDR<.10) in
delta cells in T1D, and increased abundance of multiple immune populations in ND AAB+ and
recent-onset T1D. There was also more nominal evidence (p<.05) for altered abundance of
specific cell sub-types including 'enzyme-producing' acinar (p=.037) and MUC5b+ ductal cells
(p=.049). We next asked whether there were corresponding changes in the abundance of
specific niches in T1D in spatial profiles. First, we quantified the pairwise similarity between ND
and T1D spatial graphs using Wasserstein distance[26] (**Supplementary Figure 4D**), which
revealed significant changes in the underlying structure of endocrine cells (alpha and beta) in
T1D. We moreover observed significant changes in the abundance of the endocrine niche, as
well as the *MUC5b+* ductal cell niche, in T1D (p<.05) (**Figure 1I**).

Comprehensive map of pancreatic cell type accessible chromatin
To understand how the epigenome may drive changes in cell type-specific gene expression in
T1D, we next created a matched map of accessible chromatin in pancreatic cell types. For 29
nPOD donors we performed snATAC-seq assays, and we also used the snATAC-seq profiles
from single cell multiome assays of eight donors described above. After quality control, filtering
and initial clustering steps (**see Methods**), we annotated cell type identity by label transfer of
the gene expression map using Seurat[27]. After filtering nuclei with low transfer predictions
(<0.5), there were 203,348 chromatin profiles mapping to the same cell types and sub-types
(**Figure 2A, Supplementary Figures 5,6**). We estimated that label transfer was >97% accurate

at the cell type level by comparing the predicted and actual identity of accessible chromatin

profiles in single cell multiome data. We also confirmed that predicted cell types had accessible

chromatin at the promoter regions of key marker genes (**Figure 2B**). The proportions of each

cell type were highly correlated between expression and accessible chromatin maps (r=.98,

P=1.7×10^{-13}; **Supplementary Figure 7**).

We identified transcription factor (TF) binding motifs preferentially enriched in each pancreatic

cell type and sub-type using chromVAR[28]. At the cell type level, enriched sequence motifs

revealed key regulators of each cell type; for example, beta cells and other endocrine cells were

enriched for RFX and FOXA motifs, ductal cells for HNF1, ONECUT and TEAD motifs,

endothelial cells for ETV, FLI and GABPA motifs, and T cells for RUNX, ETV, and ETS motifs,

among others (**Figure 2C, Supplementary Table 7**). Motif enrichments also highlighted

regulators that distinguished accessible chromatin profiles of cell types within specific lineages;

for example, NEUROD1 and NR3C1 had stronger enrichment in beta compared to other

endocrine cells (**Figure 2C)**. Acinar cells showed distinct sets of enriched TF motifs across

different sub-clusters, including 'signaling' acinar cells which were more enriched for FOS/JUN,

ATF and NFE motifs (**Figure 2C,D**, **Supplementary Table 7)**. In 'high-enzyme' acinar cells, the

strongest enrichments were for TFs such as ZEB, SNAI1-3, and TCF3-4, which were also the

most enriched motifs in acinar cells overall compared to other cell types (**Figure 2C,D**,

Supplementary Table 7). As structurally related TFs often have similar motifs, we linked TF

motifs enriched in sub-clusters to specific TFs in the same structural sub-family with concordant

expression patterns. For example, *FOSL2* and *JUNB/D*, as well as *ATF3*, *NFE2L2* and

BACH1/2, were increased in 'signaling' acinar cells, and *TCF3* had increased expression in

'high-enzyme' acinar cells (**Figure 2D**).

For each cell type and sub-type, we next defined candidate *cis*-regulatory elements (cREs). We

derived 'pseudo'-bulk accessible chromatin profiles by aggregating reads from all cells for that

cell type or sub-type and identified cREs by performing peak calling with MACS2. In total, there

were 368,688 cREs across all cell types and an average of 94.3k cCREs per cell type

(**Supplementary Table 8)**. Given expanded profiling of pancreatic cell types in our study we

determined the proportion of cREs not present in previous catalogs. Among cREs in our study,

9.4% and 7.4% were unique compared to a pan-tissue[29] and pancreas-specific[5] cRE catalog,

respectively, such as a T cell-specific cRE directly upstream of *ZNF746* (**Figure 2E**). We

identified cREs with cell type-specific activity by comparing accessible chromatin profiles across

cell types (**Supplementary Table 9, see Methods)**. Cell type-specific cREs were enriched for

sequence motifs of key cell type TFs as well as proximity to genes involved in cell type-specific

function (**Supplementary Table 10,11**). For example, beta cell-specific cREs were significantly

enriched (FDR<.10) for proximity to insulin secretion-related pathways and RFX, FOXA,

NEUROD, and NKX6.1 TF motifs, whereas endothelial-specific cREs were significantly enriched

for proximity to angiogenesis, blood vessel morphogenesis, and vasculature pathways and FLI,

ETS, and ETV TF motifs (**Supplementary Table 10,11**). We also identified cREs specific to

several of the sub-types within acinar cells; for example, 'signaling' acinar-specific cREs were

enriched for JUN, FOS, and ATF motifs.

Due to the scarcity of immune populations in the pancreas, the epigenome of resident and

infiltrating pancreatic immune cells has not been extensively described. In our study, we

identified multiple immune cell types including T cells, macrophages, B cells and mast cells,

although available cell numbers only enabled defining cREs in T cells and macrophages. T cell-

specific cREs were significantly enriched for proximity to genes involved in T cell activation, T

cell receptor complex, and cytokine receptor activity, and motifs for ETS, ETV and RUNX TFs,

and macrophage-specific cREs were enriched for immune-related processes and PU.1 and

SPIB motifs (**Figure 2F**). Compared to a previous study which profiled several whole pancreas

donors, more than double the number of cREs were identified in each cell type (T cells: 58.8k vs

24.5k; Macrophages: 114.3k vs 55.7k). The increased number of cREs improved annotation of

T1D-associated variants at immune-related loci; for example, at the *PRCKQ* locus likely causal

T1D variant rs947474 (PPA=.88) from published fine-mapping data[5] overlapped a pancreatic T

cell and macrophage cRE not identified in these cell types in the pancreas previously, and not

active in other pancreatic cell types (**Figure 2G**).

We next predicted networks of genes regulated by TF activity in each pancreatic cell type (**see**

Methods). We linked cREs to target genes in each cell type using the activity-by-contact (ABC)

method, which revealed an average of 46,474 cRE-target gene links per cell type, as well as

based on promoter proximity (**Supplementary Table 12**). Using ABC and promoter proximity,

genes were linked to, on average, 2.8 cREs per cell type (**Figure 2H**). We identified genes

which had highly cell type-specific cRE links (**see Methods**), and genes with highly cell type-

specific cRE links included key marker genes such as *INS* in beta cells, *GCG* in alpha cells, *IL2,*

IFNGR1 and *GZMA* in T cells, and *MARCOS* in macrophages. In each cell type, we next

constructed gene regulatory networks (GRNs) for 366 TFs by combining (i) cRE-target gene

links, (ii) TF sequence motif predictions in cREs, and (ii) TF and target gene expression levels

(**Figure 2H, see Methods, Supplementary Data 1**). We then annotated likely cellular functions

of TF GRNs by identifying biological pathways with gene sets that significantly overlapped TF

GRNs. There were thousands of significant relationships linking TF GRNs to biological

pathways across all cell types (Fisher's test, FDR<.10) (**Supplementary Table 13**), which

annotated many known regulators of biological pathway activity as well as many putative

functions of TFs.

Finally, we utilized spatial transcriptomics data in combination with cell type-specific TF GRNs to

infer TF activity within cell types and sub-types in the pancreas. Briefly, we used a univariate

linear model to predict the observed gene expression based on TF–gene interaction weights,

from which we scored TFs as active or inactive in each cell type[30]. We identified TFs with

endocrine-specific activity in line with the previously described regulators of endocrine cell

activity, such as *NEUROD1*, as well as high activity of *PAX6* in beta cells, where it is a key

regulator of beta cell identify, function and survival[31] (**Figure 2I,J**). Among other cell types, we

inferred high activity for *BHLHA15*/*MIST1* in acinar cells, where it may play a role in the

maintenance of pancreatic acinar identity[32], and highly specific activity for *MEOX2* in endothelial

cells and *RUNX3* in T cells (**Figure 2I,J**). Integrating GRNs with spatial transcriptomic profiles

thus confirmed the specificity of key TFs regulating pancreatic cell types, including for TFs not

measured on the spatial panel directly.

Pancreatic cell type gene expression in T1D progression

Changes in genome-wide gene activity within each pancreatic cell type during progression to

T1D are poorly understood. We therefore identified genes and biological pathways in each cell

type with altered activity in ND AAB+ and T1D. To increase our power to detect changes in

endocrine cell types, we also utilized single cell RNA-seq from purified islets of 48 non-diabetic,

ND AAB+ (primarily single AAB+), and T1D donors from the HPAP consortium[4,33,34]. For each

cell type and sub-type, we derived gene counts per sample, tested for differential expression in

single and multiple ND AAB+ and recent and long-standing T1D compared to non-diabetes, and

considered genes significant at FDR<.10 (**see Methods**). We further performed gene set

enrichment of differential expression results for each cell type and sub-type and identified

pathways with significant (FDR<.10) changes in activity in each condition (**see Methods**).

Marked gene expression changes were observed in beta cells in T1D (**Figure 3A**). In recent-
onset T1D, 665 genes in beta cells had significant change (FDR<.10) in expression, where the
most up-regulated genes included MHC class I and related (*CD74, B2M*) genes, cytokines and
cytokine-induced genes (*IL15*, *GBP2, IFIT3*), cytokine-responsive TFs (*STAT1/4, IRF1*), and
components of the 20S proteosome (**Figure 3B, Supplementary Figure 8, Supplementary**
Table 14). We also observed up-regulation of MHC class II genes in T1D, particularly *HLA-*
DPB1. At the pathway level, there was up-regulation of antigen processing and presentation,
interferon signaling, interleukin signaling and JAK-STAT signaling, and down-regulation of
oxidative phosphorylation, translation, mitochondrial function, mitosis, mRNA processing,
protein folding and localization, ER-Golgi transport, and autophagy (**Figure 3C, Supplementary**
Table 15). We examined whether specific pathways up-regulated in T1D showed heterogeneity
in expression across beta cells, and several had evidence for bimodal expression patterns most
prominently ECM-related pathways but also antigen presentation, while others such as
interferon and JAK-STAT signaling did not (**Supplementary Figure 9**). Compared to recent-
onset T1D, the largest changes generally differed in long-standing T1D (**Supplementary Figure**
8), where antigen presentation and class I MHC genes were less pronounced, interferon
signaling was less pronounced although specific IRF TFs had higher expression, and class II
MHC genes had stronger up-regulation. There was also stronger down-regulation in long-
standing T1D of insulin secretion and beta cell function and genes such as *GLIS3* and *G6PC2*
(**Supplementary Table 14,15**)*.*

Given marked changes in gene expression in beta cells in recent-onset T1D, we further
characterized whether these pathways had altered activity within specific localizations in the
pancreas. Of the genes with altered expression in beta cells in recent onset T1D and present in
the spatial gene panel, almost all (95%) were up-regulated in T1D in spatial profiles (**Figure 3D,**
Supplementary Figure 10). Furthermore, multiple up-regulated genes in T1D such as MHC
class I genes (e.g. *HLA-A*, *B2M*) showed spatially-dependent expression patterns (Moran's I
>0.2) within endocrine, immune and ductal cells (**Figure 3D**). We further characterized
pathways in recent-onset T1D with expression profiles dependent on specific niches and altered
in T1D progression. We identified pathways in the PROGENy database in LIANA+[35] to predict
pathways preferentially active in a niche using a multivariate linear model. We identified
multiple pathways with niche-dependent expression, including hypoxia in the endocrine niche
(**Figure 3E**). When further assessing T1D-specific changes in pathway expression, pathways

related to hypoxia and inflammation such as TNFa and JAK-STAT were differentially active in
T1D (**Supplementary Figure 10**).

In contrast to T1D, few individual genes had significant changes in expression in beta cells in
either single or multiple ND AAB+ (**Figure 3A**). We determined whether more subtle changes
might be occurring at these stages. Genes altered in recent-onset T1D had significantly
correlated effects in multiple ND AAB+, although not in single ND AAB+ (**Figure 3F**). At the
pathway level, antigen processing and presentation was up-regulated in both single and multiple
ND AAB+, and interferon signaling was up-regulated in multiple ND AAB+ (**Figure 3G,H,**
Supplementary Table 15). Among key genes in these pathways, MHC class I genes (*HLA-A,*
HLA-B, HLA-C) and interferon signaling *IRF* TFs were up-regulated in multiple but not in single
ND AAB+ (**Figure 3I, Supplementary Table 14**). We also identified pathways altered
specifically in single and multiple ND AAB+ and not in T1D; for example, heat stress response
was up-regulated in single and multiple ND AAB+, extracellular matrix organization, cytokine-
cytokine interactions, and GPCR ligand binding were all down-regulated in multiple ND AAB+,
and TGF beta signaling was down-regulated in single ND AAB+ (**Figure 3H, Supplementary**
Table 15). Additionally, class II MHC antigen presentation was strongly up-regulated in multiple
ND AAB+, but not single ND AAB+, including the class II MHC genes *HLA-DBP1* and *HLA-*
DRB1 (**Supplementary Table 14,15**). These results highlight that single and multiple ND AAB+
have both shared and distinct genomic changes in beta cells compared to T1D.

Changes have been reported in the exocrine pancreas in both T1D and at-risk individuals[21] and
in our study, we observed marked changes in exocrine cell gene expression in T1D
progression. In 'basal' acinar cells, there were 276 genes with altered expression in recent-
onset T1D, almost all of which (95%) had decreased expression (**Figure 3A, Supplementary**
Table 14). Basal acinar and other acinar sub-types showed down-regulation of numerous
pathways in recent-onset T1D including those related to signaling, stimulus response,
metabolism, and protein transport (**Figure 3A, Supplementary Table 15**). In multiple ND AAB+,
the 'basal' and 'high-enzyme' acinar sub-types showed higher expression of genes related to
amino acid metabolism, which is necessary for enzyme production, as well as carbohydrate and
glucose metabolism, insulin signaling, immune signaling, transcriptional activity, and respiration
(**Figure 3J, Supplementary Table 15**). We also observed down-regulation of genes in ductal
cells in recent-onset and long-duration T1D associated with small molecule transport, stimulus

response, cytokine signaling and RNA processing, but no evidence for changes in ND AAB+

(**Supplementary Table 15**).

Other cell types in islets and the surrounding micro-environment also had significant changes in

activity across entire pathways during progression to T1D. In alpha cells, while antigen

presentation, interferon signaling, and other pathways were increased in T1D, in contrast to

beta cells there were few changes in single or multiple ND AAB+ (**Figure 3J, Supplementary**

Table 15). Delta cells showed more prominent changes in multiple ND AAB+, including

increased hypoxia and heat stress response and cell cycle-related pathways and decreased

signaling pathways, as well as in single ND AAB+ (**Figure 3J, Supplementary Table 15**). In

endothelial cells there was increased IL2 and JAK-STAT signaling as well as SCF/KIT signaling,

which promotes angiogenesis[36,37], in recent-onset T1D (**Figure 3J, Supplementary Table**

14,15). In activated stellate cells, there was increased expression of genes associated with fibrin

clotting and decreased expression of translation in ND AAB+, and down-regulation of many

pathways in recent-onset T1D (**Supplementary Table 14,15**). While we did not observe

evidence for significant changes in gene or pathway activity in immune (T cell, macrophage)

cells, this could be due to the small number of cells profiled for these cell types.

Together, these results reveal key genes and pathways altered in pancreatic cell types in ND

AAB+ and T1D donors with both shared and distinct changes in ND AAB+ compared to T1D,

which in ND AAB+ included antigen presentation, interferon signaling, ECM-related and stress

response pathways in beta cells and metabolism and immune signaling in acinar cells.

Changes in the pancreatic cell type-specific epigenome in T1D progression

We next examined to what extent altered gene and pathway activity in pancreatic cell types in

T1D progression is driven by changes in the epigenome of ND AAB+ and T1D donors using

snATAC-seq profiles from 29 nPOD donors.

First, we identified cREs in each cell type with altered activity in T1D progression using a linear

mixed model to account for pseudo-replication (**see Methods**). We observed significant

changes (FDR<.10) in cRE activity in ND AAB+ and T1D for most pancreatic cell types

(**Supplementary Table 16**). Beta cell cREs with increased activity in recent-onset T1D were

significantly enriched (FDR<.10) for sequence motifs of steroid hormone receptors (NC3C1,

NR3C2), NF-Y (NFYA, NFYB, NFYC), interferon response factors (IRF2, IRF7), and stress-

response TFs (ATF4, STAT1, CEBPG) among others (**Figure 4A, Supplementary Table 17**).
Conversely, cREs with decreased activity in T1D were significantly enriched for sequence motifs
of TFs involved in core beta cell functions, such as HNF1 and RFX, with many beta cell identity
TFs (NKX6.1, PDX1) and other TF families including FOXA and MEF showing more nominal
enrichment (**Figure 4A, Supplementary Table 17**). We also identified sequence motifs
enriched in beta cell cREs altered in ND AAB+, including IRF, TCF and STAT TF motifs in cREs
with increased activity and MEF, RFX, and NFAT TFs in cREs with decreased activity, although
other T1D-associated motifs such as HNF1 showed no change in ND AAB+ (**Figure 4A,**
Supplementary Table 17). Sequence motifs were also enriched cREs altered in T1D
progression for other pancreatic cell types, such as MEF and RFX TF motifs in alpha cells,
RUNX TF motifs in activated stellate cells, STAT TF motifs in endothelial cells, and FOS/JUN
motifs in ductal cells.
454
We determined next whether TF motifs enriched in T1D-associated cREs in pancreatic cell
types had broader, genome-wide changes in activity in T1D progression by modeling sequence
motif accessibility across individual cells using chromVAR[28] (**see Methods**). In beta cells, we
observed consistent changes in the genome-wide accessibility of specific sequence motifs in
T1D progression, including increasing accessibility of IRF motifs and decreasing accessibility of
RFX, FOXA, and MEF motifs from ND AAB+ to T1D states (**Figure 4B, Supplementary Table**
18). In other cases, sequence motifs had different patterns in ND AAB+ and T1D, such as
decreased accessibility of HNF1 and increased accessibility of PAR-related and hormone
receptor TFs in T1D only and opposed accessibility of SIX TFs in ND AAB+ and T1D. While
alpha cells showed similar increases in accessibility of hormone receptor, stress-response, and
PAR-related TFs in T1D progression as in beta cells, there were also several marked
differences such as increased accessibility of MEF and RFX motifs in ND AAB+ and recent-
onset T1D, respectively (**Figure 4B, Supplementary Table 18).**
468
We used TF GRNs to determine which TFs drive changes in pathway activity in T1D
progression. In beta cells, pathways altered in ND AAB+ and T1D had highly specific links to TF
GRNs, suggesting key regulators of pathway activity in T1D progression (**Figure 4C,**
Supplementary Table 13). For example, pathways up-regulated in beta cells in T1D and ND
AAB+ such as interferon signaling were linked to GRNs for IRF TF motifs and antigen
processing and presentation were linked to NFY, IRF and NFkB TF GRNs, while down-
regulated pathways in T1D such as ER and Golgi-related processes were linked to CREB3L1,

XBP1 and other TF motifs (**Figure 4C, Supplementary Table 13**). We also identified TF GRNs

linked to pathways altered specifically in ND AAB+, such as heat stress related pathways and

HSF TF GRNs, extracellular matrix-related pathways and ETS, ELK and ELF TFs, and GPCR

signaling pathways and RFX and FOXA GRNs (**Figure 4C, Supplementary Table 13**). While

we observed a strong change in HNF1 motif accessibility, as well as HNF1A expression, in beta

cells in T1D (**Figure 4B,D**), no pathways linked to the HNF1 GRN had significant change in

expression in T1D. However, there was a more nominal change in beta cell development and

function pathways linked to the HNF1 GRN in T1D (**Figure 4D, Supplementary Table 13**),

supporting that reduced HNF1 activity likely underlies altered beta cell function in T1D, as has

been shown in the context of type 2 diabetes[38].

Similarly, in other pancreatic cell types, TF GRNs were linked to pathways with altered activity in

ND AAB+ or T1D. For example, in enzyme-high acinar cells, metabolic pathways altered in ND

AAB+ were linked to GRNs for specific TFs such as glucose metabolism and HNF1, amino acid

metabolism and STAT1, and oxidative phosphorylation and MEF and FOS TF GRNs (**Figure**

4E, Supplementary Table 13). In activated stellate cells, fibrin-related pathways up-regulated

in ND AAB+ were significantly linked to ELK, HOX, CEBP and other TF GRNs. In endothelial

cells, IL2 and JAK-STAT signaling pathways up-regulated in T1D were strongly linked to NFkB

(REL, RELA) and IRF TF GRNs, and SCF/KIT signaling was also linked to HOX family TF

GRNs, among others. We further explored changes in TF activity inferred from spatial gene

expression profiles of TF GRNs across cell types, which revealed increased activity of immune

regulation, inflammation and signaling TFs (e.g. STAT3, RBPJ, FOSL2, JUND), and reduced

activity of endocrine-related TFs (e.g. PAX6, GLI3, MAFA, INSM1, NEUROD1), in T1D

compared to non-disease (**Supplementary Figure 11**).

We next annotated specific beta cell cREs altered in T1D progression with putative target genes

and assessed changes in regulatory programs at specific loci. There were 114 beta cell cREs

with altered activity in T1D progression linked to genes with significant changes in expression.

For example, a beta cell cRE on chromosome 14 in the first intron of *TSHR* had increased

accessibility in recent-onset T1D and was linked to *TSHR,* which had among the largest

increases in expression in recent-onset T1D (**Figure 4F**). We identified similar cREs up-

regulated in recent-onset T1D linked to genes with highly up-regulated expression including

HLA-A (**Figure 4F**)*,* as well as *CD74, GAD1, IL15,* and *STAT1/4.* In other cases, we observed

epigenomic changes in beta cells that may precede changes in expression of cognate target

genes. For example, a cRE upstream of *IAPP* had reduced accessibility in early T1D although *IAPP* itself only had a significant decrease in expression in longer-duration T1D.

Given pathways and transcriptional regulators with altered cell type activity in T1D progression, we determined whether any changes prior to T1D onset showed evidence for a role in genetic risk of T1D. We tested for enrichment of cREs linked to genes in each pathway for T1D associated variants genome-wide (excluding the MHC locus) using fgwas[18,39] (**see Methods**). In beta cells, several pathways altered in ND AAB+ were enriched for T1D-associated variants including antigen processing and presentation (log enrich=4.48), class II MHC antigen presentation (log enrich=4.74), and interferon signaling (log enrich=6.00) as well as several extracellular interaction-related processes (focal adhesion, laminin interactions) and GPCR signaling (**Figure 4G**). By comparison, multiple other pathways previously implicated in driving T1D risk in beta cells such as apoptosis, autophagy, mitophagy, and senescence, showed limited to no enrichment (**Figure 4G**). Among other cell types, we found evidence for enrichment of immune, metabolism, and transcription related pathways in 'high-enzyme' as well as 'basal' acinar cells (**Supplementary Figure 12**).

We further identified specific T1D risk loci that may alter regulatory activity of disease-enriched pathways in key cell types such as beta cells, T cells and other immune populations, and exocrine cells during T1D progression. We identified candidate causal variants at known T1D loci by overlapping cREs altered in T1D progression with published fine-mapping data[5]. In beta cells, multiple candidate causal variants at the *IRF1* locus overlapped cREs with increased activity in T1D including at the promoter and downstream of *IRF1* (**Figure 4H, Supplementary Table 19**). There was increased beta cell expression of *IRF1* through stages of T1D progression and *IRF1* is a driver of beta cell interferon responses, which is a pathway broadly enriched for T1D associated variants (**Figure 4G**). Conversely, at the *STAT4* locus we identified cREs with increased activity in beta cells as well as T cells, although candidate causal variants for T1D at the *STAT4* locus only overlapped cREs active in T cells (**Figure 4I**). This finding supports that while increased *STAT4* activity in beta cells is observed in T1D, the *STAT4* locus more likely affects T1D risk through altered T cell function.

Together these results reveal transcriptional regulators and networks altered in T1D progression, including those regulating pathways that likely play a causal role in the

development of T1D such as antigen presentation, interferon signaling, and extracellular
interactions in beta cells.

Changes in pancreatic cell-cell signaling in T1D progression

External signaling between cell types is a key driver of changes in cell type-specific regulation
and function, and therefore we finally identified cell-cell signaling interactions in the pancreas
altered in T1D progression.

We first inferred cell-cell interactions in snRNA-seq data for non-diabetes, ND AAB+ and T1D
using 1,939 ligand-receptor pairs in CellChat[40] (**see Methods**), which revealed 87,650
interactions significant (FDR<.10) in at least one condition (**Supplementary Table 20)**.
Grouping ligands into functional categories revealed classes of outgoing signals preferentially
produced by each cell type; for example, hormones, neuropeptides, and cell adhesion
molecules from endocrine cells, and enzymes from exocrine cells (**Supplementary Figure 13,**
Supplementary Table 21).

We identified cell-cell interactions with changes in activity in T1D progression using a
permutation test and considered changes significant at FDR<.10 (**see Methods**). Overall, there
was a reduction in the number and strength of interactions in recent-onset and long-standing
T1D compared to non-diabetes, which was largely driven by exocrine cells (**Figure 5A**). In both
ND AAB+ and recent-onset T1D there was increased strength of interactions involving
endocrine cells and other cell types, although the total number of interactions was reduced
(**Figure 5A**). We further identified cell-cell interactions among cells in spatial niches and
determined changes in T1D using spatial transcriptome profiles. We identified spatially co-
expressed ligand-receptor (LR) interactions by Moran's bivariant extension in SpatialDM[41] using
LR pairs from CellChat[40]. We compared the number of detected interactions considering each
FOV as technical replicates of a donor and observed significant heterogeneity across donors
and, like dispersed cell data, fewer interactions in T1D compared to ND donors (H-
statistic=19.6, p=.0015) (**Figure 5A**).

Among specific cell types, endocrine cells displayed significant increases in both outgoing and
incoming signaling in recent-onset T1D (**Figure 5B,C**). We also observed significant increases
in incoming signaling to endothelial, ductal, and activated stellate cells, as well as nominal

changes in 'basal' and 'high-enzyme' acinar, immune, and stellate cells, in recent-onset T1D.
Summarizing signaling by functional category revealed broad classes of cell type-specific
signals altered in T1D; for example, beta and other endocrine cells had increased signaling from
cell adhesion molecules, whereas T cells had increased antigen presentation and interleukin
signaling (**Figure 5D**). We further examined changes in signaling between specific pairs of cell
types in T1D progression (**Supplementary Table 22**). Significant changes (FDR<.10) in recent-
onset T1D included increased incoming and outgoing signaling involving beta cells, including
between beta cells themselves (**Figure 5C**), as well as increased signaling for alpha cells,
outgoing signaling from 'high-enzyme' acinar cells and incoming signaling to endothelial cells.
586
Given the importance of external signaling to beta cells in T1D, we focused specifically on
signals involving beta cells. In recent-onset T1D, autocrine/paracrine signals incoming to beta
cells with significant changes in activity included cell adhesion molecules *NRXN1, CADM1,* and
NEGR1 from all endocrine cell types and the secreted factor *BMP5* from beta cells (**Figure 5E)**.
In addition, high-enzyme acinar cells had increased signaling of trypsinogen (**Figure 5E)**, and
stellate cells had increased signaling of ECM and cell adhesion molecules to beta cells. Among
immune cells, signals with significant changes in signaling to beta cells included *GZMA* and
CCL5 from T cells and *VSIR*, *GRN*, and *LGALS9* from macrophages (**Figure 5E**). In return, beta
cells had increased signaling of *IL7* and MHC class I genes *HLA-A* and *HLA-C* to T cells, as well
as increased signaling of *BMP5*, *EFNA5*, *DLK1,* and *ANGPTL2* to macrophages. Notably,
multiple beta-immune cell signals altered in T1D map to T1D risk loci (e.g. *DLK1, HLA-A, HLA-*
C, IL7R)[18].
599
We next identified differential interactions (p<0.05) in spatial profiles by performing a likelihood
ratio test, which provided support for many T1D-associated interactions identified in dispersed
cell data. For example, interactions involving HLA class I (e.g., *HLA-C*)*, APP, SPP1,* and *BMP5,*
as well as ECM-related interactions, were altered in T1D (**Figure 5F,G**). We also identified
additional interactions enriched in T1D donors, for example between migration inhibitory factor
MIF and its transmembrane receptor *CD74*, consistent with previous studies[42], and involving
several chemokines. Next, we identified spatially-co-expressed ligand-receptor pairs using
Morans' I score in Liana+[35]. We obtained the top interactions associated with each niche using
non-negative matrix factorization (**see Methods**). In T1D, an interaction between *APP* and
CD74 was enriched in the endocrine niche, where APP is involved in inflammation and could
promote immune responses in T1D (**Supplementary Figure 14**). Conversely, interactions

involving *INS*, *IGF1R*, and *INSR* and *CALM1*, among others, were enriched in the endocrine
niche from non-disease donors (**Supplementary Figure 14**).

Several ligands altered in T1D progression BMP5 and GRN (granulin) are growth factors that
have not been previously implicated in T1D. We determined the effects of *in vitro* exposure to
these ligands on gene expression using the beta cell model EndoC-BH1. Exposing beta cells to
BMP5 in culture revealed 1,926 genes with significant change (FDR<.10) in expression, where
the most up-regulated genes were *ID1-4* and *SMAD6-7,* known targets of BMP that regulate
proliferation and differentiation, and the beta cell identity gene *MAFA* (**Figure 5H,**
Supplementary Table 23). More broadly, BMP5 exposure up-regulated pathways (FDR<.10)
related to TGF beta signaling, glycolysis, secretion, and lipid metabolism, and down-regulated
pathways such as antigen presentation and chemokine signaling (**Figure 5H)**. Second, granulin
encodes secreted proteins produced by macrophages and ductal cells. Upon exposure to
granulin, 491 genes had significant change (FDR<.10) in expression including up-regulation of
beta cell function and insulin secretion genes *MAFA, ISL1*, *SOX4, CRY2* and down-regulation of
apoptosis related genes *PEA15, PDCD5,* and *CCAR1* (**Figure 5I, Supplementary Table 24**).
More broadly, granulin up-regulated pathways related to cholesterol and glycerolipid metabolism
and down-regulated interleukin signaling and inflammation, transcription and translation, and
cell death.

These results together reveal broad changes in predicted cell-cell signaling in T1D progression
most prominently among endocrine cells and niches but also involving other cell types, including
altered signals in T1D that modulate T1D-relevant regulatory programs in beta cells.

Discussion
Single cell and spatial profiling of human pancreas donors revealed extensive changes in the
abundance, regulation, and signaling of specific cell types in T1D progression, including
processes that play a likely causal role in driving disease. In beta cells, class I and class II MHC
antigen presentation and interferon signaling pathways, TF regulators of these pathways, and
cREs linked to genes in these pathways all had up-regulated activity in recent-onset T1D and
ND AAB+. Antigen presentation was altered as early as single ND AAB+ donors, suggesting
that aberrant antigen presentation in beta cells may be an initial triggering event in T1D. Further,
the heterogenous activity of antigen presentation pathways in beta cells suggests that subsets
of beta cells may initiate the immune responses. Antigen presentation and interferon signaling

pathways in beta cells were broadly enriched for T1D-associated variants and specific risk loci
for T1D were linked to key genes in these pathways such as *IRF1*. In contrast, we found limited
evidence that pathways directly related to apoptosis, as well as other processes implicated in
T1D in beta cells such as autophagy, senescence, and mitophagy, harbor T1D risk, and are
thus more likely consequences of disease. It has been long hypothesized that beta cells affect
genetic risk of T1D through increased cell death[43–49]. Our results support that beta cells may
primarily contribute to T1D risk via the initiation or exacerbation of immune responses, which
necessitates different cellular models and phenotypic readouts to understand their role in
disease.

In addition to shared pathways, gene activity in beta cells and other pancreatic cell types had
distinct changes in ND AAB+ compared to recent-onset T1D, revealing that genomic profiles
prior to T1D onset are only partially intermediate to those in T1D. In addition, the lack of
individual genes with highly significant changes expression in ND AAB+ suggests that changes
at these stages are likely more subtle, in contrast to previous reports[4]. Several pathways in
beta cells were altered specifically in multiple and single ND AAB+ such as heat shock
response and ECM organization. Heat shock responses are activated by a variety of stressors,
promote antigen presentation in beta cells, and can act as chaperones for antigens and thus
may plausibly contribute to the initiation of autoimmunity[50,51]. The breakdown of ECM is also an
important process in T1D, as both a precursor to immune invasion as well as by affecting
intrinsic beta cell function[52]. We observed a similar pattern of both shared and distinct changes
in the epigenome of beta cells in ND AAB+ compared to T1D, including increased NEUROD1
activity and decreased SIX TF activity. There were also shared and distinct features in T1D
based on the duration of disease; for example, a more pronounced reduction in beta cell
function in long-standing T1D.

In contrast to beta cells, changes in gene activity in alpha cells were largely restricted to after
the onset of T1D, including for antigen presentation and interferon response pathways and
transcriptional regulators of these pathways. This supports that immune responses are more
pronounced within beta cells compared to alpha cells prior to T1D onset, which may reflect
differences in immune targeting as well as the intrinsic properties of each cell type. The latter is
supported by multiple *in vitro* studies showing pronounced responses of beta cells to external
T1D-relevant stressors[19]. A previous study revealed changes in alpha cell function and gene
expression in single ND AAB+ donors using data from the HPAP consortium[53], although there

were overall few genes with altered expression in this study which supports our findings that
genomic changes in alpha cells prior to T1D onset are likely subtle. In addition, several TF
families such as RFX and MEF2 had different patterns of accessibility between alpha cells and
beta cells in T1D progression, further highlighting the unique responses of each cell type to
disease progression. Conversely to alpha cells, delta cells had altered activity of multiple
pathways related to stress and inflammatory responses in single and multiple ND AAB+, as well
as decreased abundance in T1D, suggesting they may play an as-of-yet unappreciated role in
T1D progression.

Given that we profiled whole pancreas donors, our study was uniquely placed to reveal changes
in the exocrine pancreas compared to previous single cell studies which used purified islets[4,15].
We identified multiple clusters of acinar and ductal cell types which had distinct genomic profiles
and may represent heterogeneous sub-types of these cell types. In acinar cells, sub-clusters
were broadly related to enzyme production and signaling responses, and previous reports
highlighted similar heterogeneity in secretory and idling acinar cells[54]. Similar hormone
producing and signaling states have been reported in endocrine cells[55], and thus may represent
a common property of secretory cells. Resolving exocrine sub-clusters revealed genomic
changes within specific exocrine sub-types in T1D. Enzyme-producing acinar and *MUC5B*+
ductal cells were more abundant in ND AAB+ donors, and multiple acinar sub-types had altered
metabolism, immune, and transcriptional pathways, as well as increased signaling to beta cells,
in T1D progression. Specific pathways within acinar cells altered in T1D progression also
harbored T1D-associated variants, further supporting a role for exocrine pancreas in T1D risk[4,18]
and providing new in-roads to determine how cellular processes in acinar cells contribute
causally to T1D.

Signaling relationships between pancreatic cell types revealed incoming and outgoing external
signals during progression to T1D. Cell-cell signaling between immune and beta cells
highlighted known signals in T1D, such as granzyme B incoming to beta cells and class I MHC
presented by beta cells[56,57], as well as potential mechanisms of genes implicated in T1D genetic
risk such as *DLK1* and *IL7* signaling from beta cells to immune cells[18]. Additional signals
incoming to beta cells in T1D such as BMP5 and granulin have no prior known role in disease.
BMP5 has increased autocrine/paracrine signaling in T1D and *in vitro* suppressed antigen
presentation- and chemokine-related genes and enhanced expression of several genes linked
to beta cell proliferation and function. Other BMP proteins have been shown to both enhance

and inhibit beta cell function, maturity, and proliferation[58,58–60], where the direction of effect may

depend on the level of BMP signaling. Granulin suppresses class I MHC expression and T cell

infiltration of ductal adenocarcinoma cells in the context of pancreatic cancer and has been

shown to promote proliferation in mouse models of beta cells[61,62]. Signaling pathways altered in

T1D, particularly those involved in T1D genetic risk, may represent therapeutic areas for

preserving beta cell function to prevent or reverse T1D.

There were multiple limitations of our study that could be used to inform directions for future

studies. For example, while we grouped non-diabetic donors by number of autoantibodies,

there is additional granularity in stages of T1D stages; for example, stage 2 of T1D is marked by

both autoantibody positivity and reduced beta cell function[63]. Future studies may therefore

utilize islet functional measures to help refine characterization of T1D stages for genomic

analyses. In addition, heterogeneity in T1D pathogenesis has been defined based on criteria

such as first-developed autoantibody, HLA background, age and other factors[64–67], and

continued collection of larger sample numbers will enable understanding genomic changes

within donors mapping to disease sub-groups. As immune subsets infiltrating the pancreas

arise from pancreatic lymph nodes (pLNs)[68], studies combining pLN data with pancreas data

from matched donors will be valuable in understanding the role of immune cells in driving T1D

progression in the pancreas. Finally, expanded spatially-resolved profiling of cells will continue

to help reveal cell type-specific changes within disease-related cellular niches and

neighborhoods.

In summary, our study revealed gene regulatory changes in pancreatic cell types in T1D

progression and highlighted pathways, regulatory networks, and signals that may play a causal

role in T1D; efforts that inform both new directions for mechanistic studies and novel targets for

therapies to prevent or reverse T1D. We provide these data and maps in visualization tools

(available at http://t1d-pancreas.isletgenomics.org) to further enhance their utility to the research

community. More broadly, our study highlights the utility of single cell multiomics and spatial

analysis to reveal insight into cellular processes underlying progression to complex disease.

Methods

Sample selection

Whole pancreas tissue was obtained from the Network for Pancreatic Organ Donors with

Diabetes (nPOD) biorepository according to federal guidelines with informed consent obtained

from each donor's legal representative. Studies were considered exempt and approved by the

Institutional Review Board (IRB) of the University of California San Diego. We selected 7 T1D

donors with more recent onset (<1 year from diagnosis) and 5 T1D donors with longer duration

(>5 years from diagnosis), along with 11 age- and sex-matched non-diabetic (ND) individuals.

We also selected 9 non-diabetic donors with T1D autoantibodies (ND TD AAB+), the majority of

which had multiple antibodies although one donor was single GAD+. In total, 32 donors were

obtained for genomic profiling (**Supplementary Table 1**).

Single cell assays

Tissue homogenization

Flash-frozen pancreas tissue was homogenized using mortar and pestle on liquid nitrogen. ~40

mg of ground tissue was used as input for the different single nucleus assays.

Generation of single nucleus ATAC-seq data

Roughly 40 mg of ground pancreas tissue was resuspended in 1 ml of nuclei permeabilization

buffer (10 mM Tris-HCl (pH 7.5), 10 mM NaCl, 3 mM $MgCl_2$, 0.1% Tween-20 (Sigma), 0.1%

IGEPAL-CA630 (Sigma), 0.01% digitonin (Promega) and 1% fatty acid-free BSA (Proliant,

68700) in molecular biology-grade water). Nuclei suspension was filtered with a 30-µm filter

(CellTrics, Sysmex) and then incubated for 5 min at 4 °C on a rotator. Nuclei were pelleted

with a swinging-bucket centrifuge (500 × *g*, 5 min, 4 °C; Eppendorf, 5920 R) and washed

with 1 ml wash buffer (10 mM Tris-HCl (pH 7.5), 10 mM NaCl, 3 mM $MgCl_2$, 0.1% Tween-20,

1% BSA (Proliant, 68700) in molecular biology-grade water). Nuclei were pelleted and

resuspended in 10 µl of 1x Nuclei Buffer (10x Genomics). Nuclei were counted using a

hemocytometer, and 15,360 nuclei were used for tagmentation. snATAC-seq libraries were

generated using the Chromium Single Cell ATAC Library & Gel Bead Kit v1.1 (10x Genomics,

1000175), Chromium Chip H Single Cell ATAC Kit (10x Genomics, 1000161) and indexes

(Single Index Kit N Set A, 1000212) following manufacturer instructions. Final libraries were

quantified using a Qubit fluorometer (Life Technologies), and the nucleosomal pattern was

verified using a TapeStation (High Sensitivity D1000, Agilent). Libraries were sequenced on

NextSeq 500, HiSeq 4000 and NovaSeq 6000 sequencers (Illumina) with the following read

lengths (Read1 + Index1 + Index2 + Read2): 50 + 8 + 16 + 50.

Generation of single nucleus RNA-seq data

Roughly 40 mg of ground pancreas tissue was suspended in 500 µL of nuclei buffer: 0.1%

Triton-X-100 (Sigma-Aldrich, T8787), 1× EDTA free protease inhibitor (Roche or Pierce), 1 mM

DTT, and 0.2 U/µL RNase inhibitor (Promega, N211B), 2% BSA (Sigma-Aldrich, SRE0036) in

PBS. Sample was incubated on a rotator for 5 min at 4 °C and then pelleted with a swinging

bucket centrifuge (500× g, 5 min, 4 °C; 5920 R, Eppendorf). Supernatant was removed and

pellet was resuspended in 400 µl of sort buffer [1 mM EDTA and RNase inhibitor (0.2 U/µl) in

2% BSA (Sigma-Aldrich, SRE0036) in PBS] and stained with DRAQ7 (1:100; Cell Signaling

Technology, 7406). 75,000 nuclei were sorted using an SH800 sorter (Sony) into 50 µl of

collection buffer [RNase inhibitor (1 U/µl) and 5% BSA (Sigma-Aldrich, SRE0036) in PBS].

Sorted nuclei were then centrifuged at 1000g for 15 min (Eppendorf, 5920R; 4°C, ramp speed of

3/3), and supernatant was removed. Nuclei were resuspended in 18 to 25 µl of reaction buffer

[RNase inhibitor (0.2 U/µl) and 1% BSA (Sigma-Aldrich, SRE0036) in PBS] and counted using a

hemocytometer. 16,500 nuclei were loaded onto a Chromium controller (10x Genomics).

Libraries were generated using the 10x Genomics, Chromium Next GEM Single Cell 3' GEM,

Library & Gel Bead Kit v3.1 (10x Genomics, 1000121), Chromium Next GEM Chip G Single Cell

Kit (10x Genomics, 1000120) and indexes (Single Index Kit T Set A, 10x Genomics, 1000213 or

Dual Index Kit TT Set A, 10x Genomics, 1000215) according to the manufacturer specifications.

Complementary DNA was amplified for 12 PCR cycles. SPRISelect reagent (Beckman Coulter)

was used for size selection and cleanup steps. Final library concentration was assessed by the

Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific), and fragment size was checked using a

TapeStation (High Sensitivity D1000, Agilent). Libraries were sequenced using a NextSeq 500

and   a   Novaseq6000   (Illumina)   with   the   following   read   lengths

(Read1 + Index1 + Index2 + Read2):   28   +   8   +   0   +   90   (single   index)   or

28 + 10 + 10 + 90 (dual index).

Generation of joint single-nucleus RNA and ATAC-seq data (Multiome)

~40 mg ground tissue was resuspended in 1 ml of wash buffer (10 mM Tris-HCl (pH 7.4),

10 mM NaCl, 3 mM MgCl$_2$, 0.1% Tween-20 (Sigma), 1% fatty acid-free BSA (Proliant, 68700),

1 mM DTT (Sigma), 1x protease inhibitors (Thermo Fisher Scientific, PIA32965),

1 U µl^{-1} RNasin (Promega, N2515) in molecular biology-grade water). Nuclei suspension was

filtered with a 30-µm filter (CellTrics, Sysmex) and pelleted with a swinging-bucket centrifuge

(500 × g, 5 min, 4 °C; Eppendorf, 5920 R). Nuclei were resuspended in 400 µl of sort

buffer (1% fatty acid-free BSA, 1x protease inhibitors (Thermo Fisher Scientific, PIA32965),

1 U µl⁻¹ RNasin (Promega, N2515) in PBS) and stained with 7-aminoactinomycin D (7-AAD;

1 µM; Thermo Fisher Scientific, A1310). A total of 120,000 nuclei were sorted using an SH800

sorter (Sony) into 87.5 µl of collection buffer (1 U µl⁻¹ RNasin (Promega, N2515), 5% fatty

acid-free BSA (Proliant, 68700) in PBS). Nuclei suspension was mixed in a ratio of 4:1 with 5x

permeabilization buffer (50 mM Tris-HCl (pH 7.4), 50 mM NaCl, 15 mM MgCl₂, 0.5% Tween-

20 (Sigma), 0.5% IGEPAL-CA630 (Sigma), 0.05% digitonin (Promega), 5% fatty acid-free BSA

(Proliant, 68700), 5 mM DTT (Sigma), 5x protease inhibitors (Thermo Fisher Scientific,

PIA32965), 1 U µl⁻¹ RNasin (Promega, N2515) in molecular biology-grade water) and

incubated on ice for 1 min before pelleting with a swinging-bucket centrifuge (500 × g,

5 min, 4 °C; Eppendorf, 5920 R). Supernatant was gently removed, and ~10 µl were left

behind to increase nuclei recovery. A total of 650 µl of wash buffer (10 mM Tris-HCl (pH 7.4),

10 mM NaCl, 3 mM MgCl₂, 0.1% Tween-20 (Sigma), 1% fatty acid-free BSA (Proliant, 68700),

1 mM DTT (Sigma), 1x protease inhibitors (Thermo Fisher Scientific, PIA32965),

1 U µl⁻¹ RNasin (Promega, N2515) in molecular biology-grade water) was added with minimal

disturbance of the pellet, and samples were centrifuged again with a swinging-bucket centrifuge

(500 × g, 5 min, 4 °C; Eppendorf, 5920 R). Supernatant was gently removed without

disturbing the pellet, leaving ~2–3 µl behind. Approximately 7–10 µl of 1x Nuclei Buffer (10x

Genomics) were added, and nuclei were gently resuspended. Nuclei were counted using a

hemocytometer, and 18,300 nuclei were used as input for tagmentation. Single-cell multiome

ATAC and gene expression libraries were generated following manufacturer instructions

(Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle, 10x

Genomics, 1000283; Chromium Next GEM Chip J Single Cell Kit, 10x Genomics, 1000234;

Dual Index Kit TT Set A, 10x Genomics, 1000215; Single Index Kit N Set A, 10x Genomics,

1000212) with the following PCR cycles: 7 cycles for pre-amplification, 8 cycles for ATAC index

PCR, 7 cycles for complementary DNA (cDNA) amplification, and 12 cycles for RNA index PCR.

Final libraries were quantified using a Qubit fluorometer (Life Technologies), and the size

distribution was checked using a TapeStation (High Sensitivity D1000, Agilent). Libraries were

sequenced on NextSeq 500 and NovaSeq 6000 sequencers (Illumina) with the following read

lengths   (Read1 + Index1 + Index2 + Read2):   ATAC   (NovaSeq   6000),

50 + 8 + 24 + 50; ATAC (NextSeq 500 with custom recipe), 50 + 8 + 16 + 50; RNA

(NextSeq 500, NovaSeq 6000), 28 + 10 + 10 + 90

Quality control and filtering

Single nuclei ATAC data was processed and aligned to reference genome hg38, and duplicate

reads were removed using Cellranger ATAC (version 1.1.0). Chromatin accessibility for each

sample was quantified in 5kb genome windows as previously described[69]. Nuclei with less than

1000 unique ATAC-seq fragments were removed. Initial quality control was performed to retain

cells in each sample using the following metrics unique usable reads > 5000, fraction promoters

used > 0.01, TSS enrichment (TSSe) > 0.3 using scanPy v1.8.0. Doublets were removed using

Amulet v1.0 per sample[70]. Single nucleus ATAC-seq datasets were processed and aligned to

reference genome hg38, and duplicate reads were removed using Cellranger ATAC v1.1.0.

Chromatin accessibility for each sample was quantified in 5kb genome windows as previously

described[69]. Nuclei with less than 1000 unique ATAC-seq fragments were removed. Initial

quality control was performed to retain cells in each sample using the following metrics unique

usable reads > 5000, fraction promoters used > 0.01, TSS enrichment (TSSe) > 0.3. Doublets

were removed using Amulet v1.0 per sample[70].

Single nuclei RNA samples were processed using Cellranger (version 6.0.1) with reference

genome hg38[71]. Individual samples were processed for quality initially by removing nuclei with

less than 500 expressed genes. Doublets were detected for each sample using DoubletFinder

(version 2.0.3) using an expected doublet rate of 4% for all samples[72]. In effort to reduce

ambient RNA contamination largely driven by acinar cells, we utilized SoupX (version 1.6.2) and

selected acinar marker genes, REG1A and PRSS1, to estimate contamination rates[73]. Gene

expression count matrices were then corrected for this predicted contamination, these correct

counts were used for both clustering and downstream analysis. Single nucleus RNA-seq

datasets were processed using Cellranger v6.0.1 with reference genome hg38[71]. Individual

samples were processed for quality initially by removing nuclei with less than 500 expressed

genes. Doublets were detected for each sample using DoubletFinder using an expected doublet

rate of 4% for all samples[72]. In an effort to reduce ambient RNA contamination largely driven by

acinar cells, we utilized SoupX and selected acinar marker genes, REG1A and PRSS1, to

estimate contamination rates[73]. Gene expression count matrices were then corrected for this

predicted contamination, these correct counts were used for both clustering and downstream

analysis.

Paired multiome data was processed, aligned, and multiplet reads were removed using

cellranger arc (version 2.0.0) with the reference genome hg38. Individual sample quality control

was done using both modalities to remove low quality nuclei without a minimum of 500

expressed genes and 1000 ATAC-seq fragments. Ambient RNA contamination was removed

using SoupX (version 1.6.2) using the same parameters as previously described. Doublets were

detected and removed for both modalities using DoubletFinder (version 2.0.3) and Amulet

(version 1.0), with the same parameters as above for single modality data[70,72]. Paired multiome

datasets were processed, aligned, and multiplet reads were removed using cellranger arc

(version 2.0.0) with the reference genome hg38. Individual sample quality control was done

using both modalities to remove low quality nuclei without a minimum of 500 expressed genes

and 1000 ATAC-seq fragments. Doublets were detected and removed for both modalities using

DoubletFinder and Amulet, with the same parameters as above for single modality data[70,72].

Clustering

Gene expression

After individual sample quality control, high quality barcodes from single modality snRNA-seq

and the RNA modality of our multiome data were clustered for 40 samples (32 snRNA and 8

snRNA multiome) using Seurat (version 4.3)[74]. Quality control metrics such as high

mitochondrial percentage (>1%), high number of genes detected (>4,000 genes), and high

number of RNA counts (>7,500) were used to remove low quality barcodes. A combined

clustering was created using principal components (PCs) from PCA of gene expression. We

used Harmony[8] (version 1.0.3) to correct the PCs for batch effects across samples, sex, and

sequencing technology. Clusters were removed with low number of cells (<10 cells) and with

quality metrics such as number of detected genes and RNA counts lower than other clusters.

Additional doublet cells were removed based on the expression of 2+ canonical markers from

unrelated cell types.

We leveraged gene expression profiles specific to the wide array of pancreatic cells from

previous work to broadly label each snRNA-seq cluster as one of the following types: alpha

(*GCG*), beta (*INS*), endothelial (*PLVAP*), lymphatic endothelial (*FLT4*), ductal (*CFTR*), acinar

(*REG1A*), stellate (*PDGFRB*), and variety of immune cells including T-cells (*CD3D*),

macrophages (*C1QC*), and mast cells (*KIT*) (**Supplementary Table 2**). Using cell type markers

previously used to annotate cell type and sub-type populations such as activated stellate

(*COL6A3*) and quiescent stellate (*SPARCL1*) we were able to annotate these clusters. We

identified previously characterized ductal subtype MUC5b ductal cells from the presence of

known marker genes such as *MUC5B*, *TIFF3*, and *CRISP3*[54].

Marker genes of acinar sub-clusters were identified using DESeq2[76] (version 1.34), followed by

gene set enrichment of sub-cluster marker genes in KEGG[77-79] and REACTOME[80] pathways

using fGSEA[81] (version 1.20). In brief this was done by first creating two sets of sample pseudo-

bulk count matrices of SoupX corrected gene expression for each cell type, one set which has

the summation of count per sample per gene for that cell type and another with the summation

of counts per sample per gene for all other cell types. We then performed DESeq for each cell

type by concatenating these two matrices as our input and using cell type as outcome variable

with sample ID as a covariate.

Accessible chromatin

We first merged 40 samples (32 snATAC samples, 8 multiome snATAC samples) from 29

donors using read counts in 5kb windows using Signac[82](version 1.9.0). We then performed

latent semantic indexing (LSI) of the combined snATAC data using Signac[82]. Harmony (version

1.0.3) was used to correct for batch effects using the covariates sample, sex, and sequencing

technology[75]. Clustering was performed on the batch-corrected PCs using graph-based Leiden

clustering. We removed nuclei with a TSS enrichment (TSSe) score <2, and removed clusters

with less than 10 cells or with overall lower quality metrics, such as fraction of read in peaks,

number of ATAC fragments per barcode, and fraction of reads in promoters compared to other

clusters. After an initial window-based clustering, we called peaks using MACS2[83] (version

2.2.7.1) (parameters: -q 0.05 --nomodel --keep-dup all) on each cluster and then repeated the

entire clustering process using a consensus set of peaks merged across clusters. Additional

doublets were manually removed based off the presence of promoter accessibility of other cell

type marker genes. This was done using 9 known marker genes (*INS, GCG, REG1A,REG2B,*

CTRB2, PRSS1,PRSS2, CFTR, C1QC); promoter region was considered 2kb upstream of the

TSS. Data was clustered again after the removal of doublets. To identify cell types, we first

assigned gene names to peaks that overlapped 2kb upstream of TSS and gene body using the

gene activity function in Signac and then determined gene activity in established marker genes

for each cell type and sub-type.

We next performed label transfer on the snATAC object using our gene expression map as

reference and the peak-based chromatin data as query in Signac. Due to the size of the

chromatin data, prior to label transfer we randomly split the barcodes in the object into smaller

subsets. We used the 2k most highly variable features from the gene expression map to derive

transfer anchors using canonical correlation analysis (CCA). These anchors were then used to

transfer to our chromatin map using the TransferData function in Seurat (version 4.3). After
each subset object was done with label transfer, we merged the objects and re-clustered all the
chromatin data together using the same methods described above. Finally, we removed cells
with low prediction scores (max.predicted.score<0.5), and all cells passing this threshold were
labelled with the predicted cell type annotation. For acinar cells, we summed the prediction
scores of all acinar subtypes then filtered by a combined acinar max.predicted.score <0.5.

To determine the accuracy of label transfer, we utilized single cell multiome data where the
identity of the accessible chromatin profile is already known from the paired gene expression
profile. Since the gene expression and chromatin profiles for these nuclei were analyzed
separately, we could use them as an independent check to determine how many barcodes were
correctly labeled.   We identified multiome barcodes present both chromatin and gene
expression maps, and then calculated the percentage of accessible chromatin barcodes with
matching cell type assignments in label transfer and from the paired gene expression profile.
Due to the limited transferring of subtypes in the chromatin modality, we calculated a
percentage at both the sub-type and primary cell type levels.

Generation of spatial transcriptomics data
Pancreatic tissue from six nPOD organ donors - three with T1D (6228, 6247, 6456) and three
without diabetes (6431, 6339, 6229), matched by age and sex - was selected for spatial
transcriptomic profiling on the CosMx platform (NanoString, Seattle, WA). For each donor, five
consecutive FFPE tissue sections from the pancreatic body region were cut at a thickness of 4
microns. Sections #1, #2, #4, and #5 were mounted on the back of VWR Superfrost Plus Micro
Slides, centered within the scanning area. After sectioning, the slides were air-dried overnight at
room temperature, sealed, and immediately shipped with desiccant and ice packs to the
NanoString facility (Seattle, Washington), where they were processed within two weeks of
receipt. Section #3 was triple-stained for CD3, insulin, and glucagon using chromogen-based
immunohistochemical staining using the Mach2 Double Stain 1/Mach2 Double Stain 2 HRP-AP
Polymer Detection Kit according to the manufacturer's instructions (Biocare Medical, Pacheco,
CA) and chromogens used included Betazoid DAB (CD3), Warp Red (insulin), and Ferangi Blue
(glucagon; all from Biocare Medical). Slides were then counterstained with hematoxylin. After
staining, the slide was digitized at 20X magnification using an Aperio CS2 slide scanner (Leica
Biosystems, Inc., Wetzlar, Germany), and this image served as a reference for field-of-view
(FOV) selection during CosMx data processing. The FOVs were selected by prioritizing specific

features such as insulitic islets, islets with few insulin-positive cells, insulin-negative islets, and
areas of inflammation in acinar tissue. The gene panel used for spatial imaging included 1010
total genes, including a fixed panel of 1000 genes on the Human Universal Cell Characterization
RNA Panel and 10 additional custom genes selected for this project. The imaging experiments
using the CosMX platform were performed at NanoString (Seattle, WA). Cell segmentation was
performed by NanoString using Giotto[84], which included using immunofluorescence for
glucagon to mark islets, CD3 or CD45 to mark immune cells, and PanCK for ductal cells +
DAPI.
990
Quality control and transcriptomic clustering of segmented cells
For downstream analysis of spatial transcriptomes, we used the python toolkits Scanpy[85] and
Squidpy[24]. For each slide, we imported matrices containing the gene expression, metadata and
positions of segmented cells. We defined a unique cell name and created a merged anndata
object with data from all the slides. We adopted a standard filtering strategy, removing cell with
less than 10 detected genes and removing genes detected in less than 300 cells. We then
normalized the counts per cell, such that every cell has the same total count after normalization
(1e6), and we log-transformed the counts.
999
Clustering of segmented cells and cell type annotation in spatial data
To cluster the segmented cells we first integrated the samples using scVI v1.1.2[86]. We
performed integration by condition using the slide as a categorical covariate. We then used the
latent representation to create a shared nearest neighbor graph and compute UMAP for two-
dimensional visualization. We performed hierarchical clustering on the scVI latent space at
resolutions of 0.5 and 0.7 and we identified 15 and 16 transcriptomic clusters for ND and T1D
respectively. To annotate cell types, we identified marker genes enriched in each cluster for
knowledge-based cell type annotation. We detected endocrine cells by hormone expression,
beta (*INS*, *IAPP*) and alpha (*GCG*, *TTR*); we also identified exocrine cells positively expressing
epithelial marker *EPCAM*, ductal (*SOX9*, *KRT19*) and acinar (*EGF*, *DLL1*, *JAG1*); we further
annotated endothelial cells (*PECAM1*, *VWF*), fibroblasts (*VIM*, *COL1A1*), immune cells (*CD4*,
CD8A), and mast cells (*CPA3*, *TPSAB1*).
1012
Cell type label transfer from reference snRNA-seq data
To achieve a finer annotation on the spatial context, we transferred the cell type labels from the
dissociated reference to the spatial data using spatial mapping function from moscot v0.3.5[87].

First, we performed pseudo-bulking of dissociated data using decoupler v1.6.0[30]. We found the

optimal combination of parameters for the spatial mapping task by hyperparameter tunning per

FOV and we used cosine distance between the modalities. For the annotation mapping, we

selected the label of the annotated cell with the highest matching probability.

1020

1021   *Identification of spatial cellular neighborhoods*

1022   Cellular neighborhoods in the spatial context were computed per FOV utilizing the squidpy[24]

1023   function spatial_neighbors, where we utilized generic coordinates and considered 30 nearest

1024   neighbors.

1025

Identification and annotation of multicellular spatial niches

To identify multicellular niches, we computed the covet representation implemented in envi

v0.3.0[25] per FOV. We used the default parameters, which included 64 genes to represent the

covariance matrix. We then created a shared nearest neighbor graph using the covet

representation and performed unsupervised Leiden clustering with a resolution of 0.2. To

annotate the clusters, we evaluated the relative cell type abundance in each group per fov and

performed hierarchical clustering. We aggregated 'Acinar basal', 'Acinar High Enz', 'Acinar

signal' and 'Acinar sig/diff' subtypes in the acinar niche, 'Ductal' and 'MUC5b ductal'subtypes in

the ductal niche, 'Alpha', 'Beta' and 'Delta' subtypes in the endocrine niche and 'Act stellate', 'Q.

stellate', 'Endothelial', 'Macrophage' and 'T cells' in the connective tissue niche.

1036

Downstream analysis:

Final peak calling and signal tracks

Cell type specific set of chromatin peaks were derived using MACS2[83] v2.2.7.1 on the final cell

type annotations of our chromatin map using the following parameters -q 0.05 --nomodel --keep-

dup all. These peak calls were used to accessible chromatin signal tracks in UCSC genome

browser[88].

1043

Marker CREs

Cell type-specific cREs were derived for each cell type and subtype. We first created a set of

union peaks across the whole dataset. This was achieved by limiting peak size for all called

peaks to 300bp by centering any peaks larger than 300bp at their summit and extending

coordinates 150bp in either direction. We then grouped peaks based on overlap to create

clusters of peaks. Within each cluster, the peak with the highest read count at its summit was

identified as the reference peak for the region. We then generated a list of peaks that did not
overlap any of the reference peaks and iteratively identified additional reference peaks again
until no peaks remained.

1053

We used this set of union peaks to calculate two sets of sample level pseudo-bulk matrices per
cell type as follows: first, we aggregated the number of ATAC fragments within peaks per donor
per cell type, then for each cell type created a second matrix with the summation of fragments
from all other cell types. Normalized counts matrices were generated by dividing number of
fragments within a peak by total number of fragments for that sample in that cell type then
multiplying by scaling factor (1e6). Cell type specific regulatory elements were then determined
for each cell type by comparing the normalized counts matrix for a given cell type with the
normalized counts matrix of all other cell types summed together. To test enrichment of a given
peak for each cell type, we performed a logistic regression model using sample id as a covariate
and corrected for multiple tests using the Benjamini-Hochberg correction method (FDR<0.1).
We limited the marker cREs per cell type to the top 5,000 cREs ranked by fold-change.   We
performed sequence motif enrichment of marker cREs for each cell type compared to a
background of all cREs in the cell type using HOMER[89] v5.0.1 and retained enriched motifs at
FDR<.01. We also tested for gene set enrichment in marker cREs using GREAT[90](version
4.0.4).

1069

1070 *Calculation of TPM*
1071 We derived gene expression profiles for each cell type by creating aggregate count matrices by
1072 donor per cell type. Using GENCODE v38 [91] GRCh38.p13 gene size annotations we calculated
1073 transcript per million (TPM) to normalize for gene size.

1074

Cell type proportion changes
We first scaled the counts for each cell type in a sample to 10,000 total cells per sample. For
several cell types we excluded samples with abnormally high counts (sample 6278 for beta and
delta; sample 6393 for T cells and B cells; sample 6375 for MUC5b+ ductal cells). We then
created a linear model of the log transformed counts as a function of disease status (ND, ND
AAB+, recent-onset T1D, long-duration T1D), age, sex, and BMI, as well as a linear model
without the disease status variable. We performed comparison of the nested models using a
likelihood ratio test in package lmtest[92] in R and considered p-values from the test significant at
.05.

1084

Differential gene expression

1085

To determine disease-related changes in gene expression, we performed differential analysis
using DESeq2[76] v1.34. Using the snRNA- seq data, we derived pseudo-bulk count matrices for
each cell type by aggregating all barcodes of a donor for each gene on a per cell type basis. We
created the count matrices from the SoupX[73] corrected expression counts, and then rounded
counts in the matrix to the nearest integer. We included sex, age, and BMI, as well as proportion
of beta cells, as covariates in the model. For endocrine cell types, we included expression
counts from scRNA-seq of 48 donors from the HPAP consortium[93] derived from a previously
created single cell map[34], and included an additional covariate in the model for cohort. For a
given cell type, we only used samples with at least 20 cells, except for long-duration T1D beta
cells where we included all samples. In addition, genes were only tested for a cell type if
detected in at least 2 samples per tested condition and if there was total of at least 10 counts
across all tested conditions. We further excluded genes for each cell type that are established
marker genes for a different cell type. Multiple test correction was performed using Benjamini-
Hochberg correction and we considered genes significant at FDR<.10.

1100

Differential cRE accessibility

1101

Using cell type specific peak calls from MACS2[83] v2.2.7.1 per cell type we created peak by
barcode fragment count matrices all snATAC-seq donors for each disease condition. Lowly
accessible peaks were removed from analysis, as determined by the average accessibility of
peak across all samples less than median accessibility of all peaks across all samples. In
addition, for each cell type samples were removed with less than 10 barcodes in that cell type.
Lastly, cell types with less than 10 cells were not used in this analysis. We tested each disease
condition against non-diabetic using glmer[94] in R using the logistic regression model [Peak
accessibility ~ Disease + scale(FRiP) + scale(count) + (1|Sample)] using a binary peak count
matrix. We used the fixed covariates of fraction reads in peak (FRiP) and ATAC fragment count
(count) to account for sequencing depth variation and used sample ID as a random effect to
adjust for sample variation. We used sample as random effect to mediate the pseudo-replication
side effect of barcode level analysis. Cell types with more than 30k cells were subsampled down
to 10k for this analysis. Disease related fold change was calculated by the following formula:
(mean(disease peak accessibility) /mean(non-diabetic peak accessibility)). Multiple test
correction was performed using the Benjamini-Hochberg method and we considered cREs
significant at FDR<.10.

1118

Pathway enrichment during T1D using gene expression input

1120 To test for pathways enriched by disease, we performed gene set enrichment analysis
1121 (GSEA)[95,96]. Using the results from our differential expression analysis input genes were ranked
1122 using the following formula (-log10(pvalue)*log2FoldChanges), and fGSEA[81] v1.20 was run
1123 using both KEGG[77–79] and REACTOME[97–103] databases [parameters: eps=0.0, minSize = 0,
1124 maxSize = 1000]. Enriched pathways were filtered down using an FDR cutoff of 10%.

1125

Motif Enrichment

We used chromVAR[20] to measure z-scored motif accessibility in snATAC-seq data. To do so,
we prepared peak count data for input to chromVAR by converting the fixed peak sparse count
matrix into a SummarizedExperiment and estimated GC content bias using chromVAR's built in
method[20,21]. Human TF motifs from JASPAR 2022[22] were accessed using the JASPAR2022
Bioconductor package[23] and motifs were annotated to peaks using motifmatchr[24]. The
SummarizedExperiment and motif annotations were used as inputs into chromVAR's
computeDeviations function to derive GC bias corrected motif accessibility z-scores.

1134

Motifs Enriched in Cell Types

1136 TF motifs were filtered for those with an accessibility >1.2 based on chromVAR's built in
1137 computeVariability function. Cell types with fewer than 50 cells were excluded. Cell type motif
1138 accessibility z-scores were averaged and plotted with pheatmap[25] and RColorBrewer[26].

1139

Motifs Enriched in Acinar Subtypes

After sub-setting the motif matrix to barcodes from acinar cells, we averaged motif accessibility
of each acinar subtype per sample then tested each motif using a two-way ANOVA across
acinar subtypes also including a donor variable. We then calculated FDR on the p-values using
qvalue[104]. To identify which specific subtype a significant motif was most enriched in, motifs
were further tested using a two-way ANOVA comparing motif accessibility within the subtype to
the average motif accessibility for the other acinar subtypes together also including a donor
variable. P-values for each motif were corrected by Holm's method. Motifs were annotated to
sub-clusters based on being significant in the pan-subtype ANOVA, significant in the post-hoc
ANOVA with Holm's correction, and having the highest average deviation score in the given
cluster.

1151

Motif Differential Accessibility

To identify motifs with differential accessibility across disease states we used a linear mixed

model using the lmerTest package[105]. We identified motifs in a cell type enriched in cREs with

altered activity in ND AAB+ or T1D. For these motifs, accessibility was modeled by barcode

using encoded variables to contrast autoantibody, recent-onset and long-duration T1D

independently against non-diabetic controls. Scaled fractions of reads in peaks and scaled

number of counts were used as fixed effect covariates and a random effect for sample was used

to control for pseudo-replication. Samples with less than 10 cells in the given cell type were

excluded and cell types with fewer than 50 cells, or disease states with fewer than 20 cells and

3 samples were not tested. We obtained p-values from the resulting models. Motif accessibility

was averaged by sample and disease state to make boxplots. Average motif accessibility per

condition was generated by averaging sample average motif accessibility and volcano plots

were generated by comparing difference in motif accessibility vs negative log10 q values, with a

difference threshold of 0.25 and q-value cutoff of 0.05 (5% FDR) for dashed lines and coloring

and labeling of samples.

1167

Motif enrichment in differential accessible CREs

To identify TF motifs enriched in cREs differential accessibility in each cell type, we used

HOMER[106] (version 5.0.1). For each cell type, we identified cREs with nominal association

(uncorrected p<.05) and split cREs by fold change as input, and user HOMER function

findMotifsGenome with a background of all cREs for the cell type with a size parameter of 200

and a masked version of the human genome hg38. Multiple test correction was done using the

Benjamini-Hochberg method, and significant motifs were considered at FDR<.10.

1175

ABC analysis

To link cREs to target genes we used Activity-by-contact (ABC)[107] v0.2. This was done by first

creating .bam files for each cell containing only barcodes from the accessible chromatin map.

Since the HiC reference panel used was in hg19 genome build, cell type bams and peaks were

coverted to hg19 using CrossMap[108] v0.6.3, and we called peaks for each cell type with MACS2

v2.2.7.1 using this genome build. To further improve enhancer activity prediction, we used

publicly available H3K27ac ChIP-seq data for acinar, ductal, alpha, beta, and delta cells[109]. We

predicted candidate regions and enhancer activity for each cell type using the following flags: --

peakExtendFromSummit 250, --nStrongestPeaks 150000, and all genes with a TPM greater

than 1 as ubiquitously expressed genes. After ABC analysis, links were converted back to hg38

using CrossMap. We identified genes with cell type-specific cRE link profiles by calculating the

proportion of the total number of ABC links for that gene by cell type and calculating Shannon

entropy based on the proportion.

1189

Constructing TF gene regulatory networks

To determine gene regulatory networks (GRNs), we constructed units of transcription factors

linked to cCREs linked to genes. We first used a position frequency matrix (PFMatrixList object)

of TF DNA-binding preferences from the JASPAR 2022 database[110] and width-fixed peaks[111] as

input to perform TF binding motif analysis. We used the 'matchMotifs' function in the R package

motifmatchr[112] v.1.21.0 to infer cell type specific cREs bound by each TF. We linked cREs

bound by each TF to target genes based on proximity to the gene promoter (±5 kb of a TSS in

GENCODE V19 or through Activity-by-contact (ABC) links [107,113] at a score cutoff of .015. TF

GRNs were retained for analysis if the network includes fewer genes then the 90[th] percentile of

number of genes linked to a given TF. In addition to ensure TF GRNs were active in the

associated cell type, we removed any TF GRNs with an average TF expression (TPM) less than

5.

1202

Identification of cell type specific TF-modules and pathways enrichment

For each pancreatic cell population, we identified pathways and TF modules enriched using our

identified marker CREs. In brief this was done for each cell type by deriving CREs associated

with KEGG and REACTOME paths using the bedtools intersection TF-module gene linked

CREs with union peaks accessible in that cell type. These union peak based pathways were

tested for enrichment using fGSEA. We used the logistic regression marker CRE results to rank

peaks using the following formula (-log10(pvalue)*log2FoldChanges). Similarly, we tested for TF

modules enriched in each cell type by defining union peaks associated with a TF either

proximally or through ABC; then using the logistic regression marker CRE results to rank peaks

and test for enrichment using fGSEA. For both analyses, we used the Benjamini-Hochberg

method for multiple test correction and retained results with an FDR < .10.

1214

Identification of TF GRNs linked to biological pathways altered in T1D

To identify regulators of enriched pathways for each cell type, we next tested enrichment of

each TF-module in pathways identified in our fGSEA analysis. We performed Fisher's exact test

to test for overlap in genes in each TF GRN and genes in each biological pathway in KEGG and

REACTOME for each cell type. We performed multiple test correction using FDR and

considered TF GRNs linked to a pathway at FDR<.10. Next, we filtered results to biological

pathways with significantly altered expression in T1D and TF motifs belonging to TF sub-

families with differentially accessibility in T1D from chromVAR[114] results.

1223

Genetic association enrichment

We tested for enrichment of T1D associated variants using summary statistics from a published

genome-wide association study[5]. We defined groups of cREs in multiple ways; first, by

identifying all cREs in each cell type linked to genes in each biological pathway in KEGG and

REACTOME using ABC and promoter proximity links and, second, by identifying cREs in each

cell type in GRNs for each TF. We calculated Bayes Factors (BFs) for each variant with minor

allele frequency (MAF)>.05 genome-wide, excluding all variants at the MHC locus, using the

method of Wakefield[115]. We then tested for enrichment of T1D associated variants in groups of

cREs genome-wide using fgwas v0.3.6[39] with a block size (-k) of 2,500.

1233

We also tested for enrichment of fine-mapped T1D risk variants using finrich[116], which compares

the cumulative posterior probability of a set of variants in an annotation to a null distribution

drawn from permutations of a background set of annotations. For each enrichment analyses

using subsets of cREs in a cell type, we used the posterior probabilities in credible sets from a

previously published GWAS[5], the full set of cREs for the cell type as background, and 10,000

permutations.

1240

1241 We overlapped cREs in each cell type with credible sets of variants at known T1D signals from

1242 a published fine-mapping study. We further determined which cREs had at least nominal

1243 evidence (uncorrected p<.05) for differential accessibility in ND AAB+ or T1D.

1244

1245 *Cell-cell interactions*

1246 The gene expression data was pre-filtered prior to running CellChat[40] v1.1.3. First, any cell type

1247 represented by fewer than 20 cells for a sample was excluded. Next, cell types that appeared in

1248 fewer than two samples within a control or disease group were excluded from that group.

1249

We considered a ligand expressed in a specific cell type if the average expression of the ligand

in the cell type was greater than half the standard deviation (SD) of its average expression

across all cells in at least two samples. After applying these filters, we ran CellChat using the

RNA data slot of the Seurat object across the entire CCdb with default parameters except for

'trim = 0' in the "computeCommunProb" command and 'thresh=1' in the "subsetCommunication"
command[40]. Each control and disease group were processed independently. Ligands from the
CellChat database (CCdb) were grouped into high level categories by manual curation using
UniProt[117] and GeneCard[118,119] (listed in **Supplementary Table 21**). Gene families were
downloaded using biomart[120] in R.

1259

Results from different conditions were consolidated and subjected to FDR correction using the
Benjamini-Hochberg method with the q-value[104] package. Predicted interactions were
considered with an FDR<0.1 and an IS above the second quartile were considered for
downstream interpretation. To remove residual background contamination due to highly
expressed genes, the following interactions were blacklisted in all cell sources except the ones
listed: *INS* in beta cells, *GCG* in alpha cells, *SST* in delta cells, *PRSS1/2/3* in acinar cells,
CD8A, CD8B, CD8B2 in T cells, and *CD4* in T cells and macrophages.

1267

To assess the significance of differences between conditions, we randomly permuted sample
IDs among conditions and re-performed the CellChat analysis 100 times and comparing these
outcomes with the observed CellChat results. The permutations were produced and filtered
using the identical parameters as those for the observed data. Next, we aggregated the
Interaction Strength (IS) across different "units" by summing all Ligand-Receptor (LR) pairs
within a unit and normalizing this sum by the number of significant interactions for each
condition (for example, the total of all incoming ligands to Beta cells in non-diabetic samples
divided by the number of significant interactions identified in that condition). We then quantified
the difference in effect size (IS-effect size) across contrasts: ND AAB+ vs. ND, recent-onset
T1D vs. ND, and long-duration T1D vs. ND. A p-value was calculated by comparing the
observational results against the simulations using the formula: the number of instances where
simulation IS-effect size exceeded observational IS-effect size, divided by the number of
permutations. Subsequently, p-values were corrected for multiple tests using the Benjamini-
Hochberg method. We considered only interactions with an FDR less than 0.10 as significant.

1282

Functional Analysis of spatial genomics profiles
We inferred TF and pathway activities utilizing the package Liana v1.1.0[35]. For TF activity
inference, we use the cell type-specific GRNs derived from single cell multiome. We then fit a
univariate linear model to infer the interaction wights. To identify cell type-specific TFs we
performed a t-test overestimating the variance of each group and filtered TFs according to an

adjusted p-value <0.05. We inferred pathway activities using the PROGENy model[35]. We used

weights of the top 500 responsive genes ranked by p-value. We then fit a multivariate linear

model to obtain the weights corresponding to pathway interactions. As with the TF activity

analyses, we identified cell type specific pathways by performing a t-test overestimating the

variance of each group.

1293

Cell-Cell communication

We analyzed cell-cell communication in spatial transcriptomic data using SpatialDM v0.2.0[41].

We performed the analysis per condition, and per donor, having each FOV as technical

replicate. For this study, the parameters l and cutoff were set to 100, and 0.2 to represent the

spatial context. Additionally, we computed the weight matrix using the single-cell mode and we

extracted the ligand-receptor interactions from the CellChat database[40]. To compute the global

Morans' I score and the local spot detection, we used the z-score method.

1301

EndoC-Bh1 stimulation experiments and RNA-sequencing

A total of 25,000 EndoC-BH1 cells were seeded in media composed of DMEM (Corning,

10014CV), 2% BSA (Sigma, A1470), 3.5×10^{-4}% 2-mercaptoethanol (Gibco, 21985023),

0.12% Nicotinamide (MilliporeSigma, 481907), 5.5 ng/mL transferrin (MilliporeSigma, T8158),

6.7 pg/mL Sodium Selenite (Sigma, 214485), and 1% Penicillin-Streptomycin (Gibco,

15140122) on a 96-well (CellTreat Scientific Products, 229105) plate coated with ECM (Sigma,

E1270) and Fibronectin (Sigma, F1141). The recombinant protein concentrations used were:

1ug/ml PGRN and 50ng/ml BMP5. EndoC-βH1 cells were obtained from Human Cell Design.

RNA was isolated using the RNeasy Mini Kit (Qiagen) from EndoC-Bh1 cells either stimulated

or unstimulated with each ligand. Samples included three replicates each for PGRN and its

untreated controls, and six replicates each for BMP5 and its untreated controls. RNA integrity

was assessed using a 2200 TapeStation (Agilent Technologies), and all samples achieved an

RNA Integrity Number (RIN) greater than 7. Ribodepleted total RNA libraries were prepared

using the TruSeq Stranded Total RNA Library Prep Gold kit (Illumina, Catalog #20020599) and

sequenced at the UCSD Institute for Genomic Medicine on an Illumina NovaSeq S4 platform.

1317

Bulk RNA-seq analysis

Quality control of the sequencing data was assessed using FastQC[121]. Transcript quantification

was performed using Salmon[122] with default parameters and the hg38 reference indexes.

Counts were imported into R using the tximport[123] package, and genes with fewer than 10 reads

were excluded. Differential gene expression analysis was conducted using DESeq2[76], applying

a false discovery rate (FDR) threshold of less than 0.1. For pathway enrichment analysis, the

fGSEA package was employed using the "stat" column from DESeq2 results. fGSEA analysis

was restricted to gene sets containing more than 10 and fewer than 500 terms. Pathways were

corrected for multiple testing using FDR with a threshold of 0.1, and only pathways belonging to

the KEGG[77,79] or REACTOME[80] databases were considered.

Figure legends

1329

Figure 1. Cell type-specific map of gene expression in the pancreas. (A) Design of study

profiling human pancreas from ND, ND AAB+ and T1D donors using single cell assays. (B)

Uniform manifold approximation and projection (UMAP) plot showing clustering of 276,906

nuclei from single nuclear RNA-seq of 32 whole pancreas donors from the nPOD biorepository.

Clusters are labeled based on cell type and sub-type annotations. (C) Dot plot showing the

normalized expression levels of selected known marker genes for pancreatic cell types and sub-

types. (D) Dot plot of genes with preferential expression across different sub-types of acinar

cells (top left), and normalized enrichment score (NES) of pathways enriched in each subtype

using fGSEA (top right). Donor transcripts per million (TPM) of selected genes with preferential

expression in different sub-types of acinar cells. (E) Representative FOV per condition (ND: top,

T1D: bottom) showing (from left to right) immunofluorescence, coarse cell type annotation with

the spatial gene panel directly, and finer-grained cell type annotation transferred from the

snRNA-seq data. (F) Matrix plots showing the neighborhood enrichment of cell types based on

spatial neighbors. (G) Stacked barplot illustrating the relative abundance of each cell type in

each multicellular niche (left). Dot plot showing the normalized gene expression levels of

spatially variable genes across multi-cellular pancreatic niches (right). (H) Normalized cell

counts for selected pancreatic cell types and sub-types organized by donor T1D and ND AAB+

status. ** FDR<.10, * uncorrected p<.05. (I) Stacked barplot showing the relative abundance of

each multi-cellular niche per condition. Niches with * have altered abundance in ND samples

(p<0.05).

1350

Figure 2. Cell type-specific map of accessible chromatin in the pancreas. (A) Uniform

manifold approximation and projection (UMAP) plot showing clustering of 203,348 nuclei from

single nuclear ATAC-seq of 29 whole pancreas donors from the nPOD biorepository. Clusters

are labeled with cell type and sub-type identity based on label transfer from the gene expression

map. (B) Genome browser showing cell type-specific accessible chromatin signal at the

promoter regions of known marker genes for pancreatic cell types. (C) Heatmap showing

genome-wide accessibility from chromVAR of sequence motifs for selected transcription factors

(TF) across cell types (left), and boxplots showing donor-level accessibility of selected TF

sequence motifs across cell types (right). (D) Genome-wide accessibility of sequence motifs for

TFs with preferential enrichment in specific sub-types of acinar cells (left), and log fold-change

in expression for genes in structural sub-families for the enriched TF motifs (right). (E) Number

of cREs identified across all cell types and the percentage of cREs that do not overlap previous

catalogs of cREs from Zhang et al and Chiou et al (top). Example of a pancreatic T cell-specific

cRE novel to this study compared to previous catalogs at the *ZNF746* locus. (F) Sequence

motifs for TFs enriched in cREs with activity specific to each cell type (left) and barplots showing

-log10 p-values of gene sets enriched for proximity to cell type-specific cREs using GREAT. (G)

Example of a cRE active in pancreatic T cells and macrophages that overlaps a candidate

causal T1D risk variant at the *PRCKQ* locus. (H) Number of gene-CRE links per gene per cell

type (top) and schematic of TF gene regulatory network (GRN) creation (bottom). (I) Matrix plot

showing the scaled Z-score of TF activities for top TFs identified for each cell type using a t-test

with overestimated variance. (J) Spatial plot of selected TFs showing the TF activity profile (top),

and cell type distribution for the respective cell type (bottom).

1373

Figure 3. Cell type-specific changes in gene expression in T1D progression. (A) Number

of genes in each pancreatic cell type with significant (FDR<.10) changes in expression in ND

AAB+ or T1D status compared to non-diabetes. Endocrine cell types include scRNA-seq data

from HPAP donors (top). Number of biological pathways enriched in genes with up- and down-

regulated expression in each cell type in ND AAB+ or T1D (bottom). (B) Volcano plot showing

differential expression results in beta cells in recent-onset T1D compared to ND. (C) Bar plot

showing normalized enrichment score (NES) of biological pathways enriched in up- and down-

regulated genes in beta cells in recent-onset T1D (bottom). (D) Scaled expression in spatial

profiles of genes with up-regulated expression in T1D in beta cells (left). Spatially-dependent

expression of selected genes (HLA-A, B2M) up-regulated in T1D in each cell type (right). (E)

Biological pathways with differential expression within spatial niches in T1D compared to ND in

spatial profiles. (F) Scatterplot of log fold-change in expression of genes in beta cells in single

or multiple ND AAB+ compared to recent-onset T1D (top) and longer-duration T1D (bottom).

Line shown in each plot is from a linear model of log fold-change values, and p-values are from

Spearman correlation. (G) Normalized enrichment score (NES) of biological pathways enriched

in differential expression results of recent-onset T1D and multiple ND AAB+. Pathways are

colored based on significant enrichment (FDR<.10) in either, or both, disease states. (H)
Normalized enrichment score of biological pathways enriched in differential expression results in
beta cells across each T1D state (single ND AAB+, multiple ND AAB+, recent-onset T1D, and
long-duration T1D) compared to non-diabetes. Red stars are for pathways with significant
enrichment (FDR<.10) in each state. (I) Log fold-change of expression of selected MHC and
interferon related genes in beta cells in each state compared to ND. Red stars indicate genes
with significant change in expression (FDR<.10) and red dots indicate more nominal change in
expression (un-corrected p<.05). (J) Normalized enrichment score (NES) of biological pathways
enriched in genes with up- and down-regulated expression in ND AAB+ and T1D in other
pancreatic cell types. Red stars indicate pathways with significant enrichment (FDR<.10).

1400

Figure 4. Epigenomic changes in pancreatic cell types in T1D progression. (A) Fold
enrichment of sequence motifs for transcription factors (TFs) enriched in beta cell cREs with up-
regulated or down-regulated activity in recent-onset T1D (top) or ND AAB+ (bottom). (B) Box
plots showing donor-level genome-wide accessibility of selected TF motifs in beta cells (left) and
alpha cells (right) from chromVAR across non-diabetes (ND), ND AAB+ (AAB) and recent-onset
T1D (T1D). (C) TF GRNs enriched for overlap with genes in biological pathways in beta cells
altered in T1D progression. (D) Biological pathways in beta cells enriched for overlap with the
HNF1A GRN (top). Beta cell expression of HNF1A in T1D progression (middle). Beta cell
activity of biological pathways linked to the HNF1A GRN in T1D progression (bottom). (E)
Genome browser views of the *TSHR* (top) and *HLA-A* (bottom) loci where beta cell cREs with
altered activity in T1D were linked to genes with concordant changes in expression in T1D. (F)
Genome-wide enrichment of T1D-associated variants in beta cell cREs linked to pathways with
altered expression in ND AAB+. (G) Genome browser view of T1D associated variants and
beta cell accessible chromatin in non-diabetes, ND AAB+ and T1D at the *IRF1* locus, where
candidate T1D variant overlaps a beta cell cRE with altered activity in T1D progression. (H)
Genome browser view of T1D associated variants and both beta cell and T cell accessible
chromatin in non-diabetes, ND AAB+, and T1D at the *STAT4* locus. Candidate T1D variants at
this locus overlap a T cell cRE but not a beta cell cRE.

1419

1420 **Figure 5. Cell-cell signaling networks altered in T1D progression.** (A) Summary of total
1421 interaction strength (top) and number of interactions (middle) for each pancreatic cell type
1422 lineage in non-diabetes, ND AAB+, recent-onset T1D and long-duration T1D. Bar plot showing
1423 the number of ligand receptor interactions identified per donor in spatial slides (bottom). (B)

Heatmap showing normalized interaction strength of outgoing and incoming signals for each cell type among donors which were non-diabetes, ND AAB+, recent-onset T1D and long-duration T1D. Stars represent the significance of the difference in interaction strength in each disease state compared to non-diabetes using permutations. **FDR<.01, *FDR<.05. (C) Difference in strength of interactions between beta cells and other pancreatic cell types and sub-types in ND AAB+, recent-onset T1D and long-duration T1D. **FDR<.01, *FDR<.05. (D) Interaction strength of outgoing and incoming signals for each cell type summarized by broad functional categories. **FDR<.01, *FDR<.05. (E) Normalized interaction strength in recent-onset T1D and non-diabetes for ligands with significant change in incoming or outgoing signal involving beta cells. (F) Heatmap per donor showing the interaction score of the top ligand-receptor interactions from a likelihood ratio test comparing ND and T1D donors. (G) Spatial plots of a representative FOV per condition (T1D: top, ND: bottom) highlighting, from left to right, spots where the interaction between HLA-C and CD8A presented a significant spatial pattern and the cell types where this interaction occurs. (H) Volcano plot showing genes with up- and down-regulated expression in EndoC-BH1 cells after treatment with BMP5 compared to no treatment (left). Biological pathways enriched in genes with up- and down-regulated expression in BMP5 exposure (right). (I) Volcano plot showing genes with up- and down-regulated expression in EndoC-BH1 cells after treatment with granulin (GRN) compared to no treatment (left). Biological pathways enriched in genes with up- and down-regulated expression after GRN exposure (right).

Code availability

All code used for this article has been made available at https://github.com/Gaulton-Lab/nPOD and https://github.com/theislab/spatial_pancreas

Data availability

Raw data can be accessed from GEO under the accession numbers GSE273594, GSE273597, GSE273598. Single cell maps can be visualized at http://t1d-pancreas.isletgenomics.org.

Author contributions

R.L.M. performed single cell analyses and wrote the manuscript. S.J. performed spatial analyses and wrote the manuscript. W.E., L.T., E.B., C.M., K.K, R.E., H.M., J.C., E.G., and A.H. performed single cell analyses. C.Z. and D.B. performed sample collection and processing. G.W. performed single cell and spatial analyses. M.M. generated single cell genomics data. K.V. performed in vitro treatment experiments. I.K. and M.A. performed sample collection and

processing. S.P. supervised single cell data analysis and contributed to the design of the study. F.J.T. supervised the spatial data analyses. M.S. designed and supervised the study. K.J.G. designed and supervised the study, performed analyses, and wrote the manuscript.

Acknowledgements

The work in this study was funded by DK120429 and DK122607 to K.J.G. and M.S. and T32 GM00866 to R.L.M. This research was performed with the support of the Network for Pancreatic Organ donors with Diabetes (nPOD; RRID:SCR_014641), a collaborative type 1 diabetes research project supported by JDRF (nPOD: 5-SRA-2018-557-Q-R) and The Leona M. & Harry B. Helmsley Charitable Trust (Grant#2018PG-T1D053). The content and views expressed are the responsibility of the authors and do not necessarily reflect the official view of nPOD. Organ Procurement Organizations (OPO) partnering with nPOD to provide research resources are listed at https://npod.org/for-partners/npod-partners/. This manuscript used data acquired from the Human Pancreas Analysis Program (HPAP-RRID:SCR_016202) Database (https://hpap.pmacs.upenn.edu), a Human Islet Research Network (RRID:SCR_014393) consortium (UC4-DK-112217, U01-DK-123594, UC4-DK-112232, and U01-DK-123716).

Declaration of interests

The following conflicts of interest are reported for several authors. K.J.G has done consulting for Genentech, received honoraria from Pfizer, holds stock in Neurocrine biosciences, and his spouse is employed by Altos Labs, Inc. J.C. and R.M.E hold stock in and are employed by Pfizer Inc. F.J.T. consults for Immunai, Singularity Bio, CytoReason, Cellarity and Omniscope, and has ownership interest in Dermagnostix and Cellarity.

References:

1. Lehuen, A., Diana, J., Zaccone, P. & Cooke, A. Immune cell crosstalk in type 1 diabetes. *Nat. Rev. Immunol.* **10**, 501–513 (2010).

2. Eguchi, K. & Nagai, R. Islet inflammation in type 2 diabetes and physiology Find the latest version□: Islet inflammation in type 2 diabetes and physiology. *J. Clin. Invest.* **127**, 14–23 (2017).

3. Boldison, J. & Wong, F. S. Immune and Pancreatic β Cell Interactions in Type 1 Diabetes. *Trends Endocrinol. Metab.* **27**, 856–867 (2016).

4. Fasolino, M. *et al.* Single-cell multi-omics analysis of human pancreatic islets reveals novel cellular states in type 1 diabetes. *Nat. Metab.* **4**, 284–299 (2022).

5. Chiou, J. *et al.* Interpreting type 1 diabetes risk with genetics and single-cell epigenomics. *Nature* **594**, 398–402 (2021).

6. Mallone, R., Halliez, C., Rui, J. & Herold, K. C. The β-Cell in Type 1 Diabetes Pathogenesis: A Victim of Circumstances or an Instigator of Tragic Events? *Diabetes* **71**, 1603–1610 (2022).

7. Atkinson, M. A. The pathogenesis and natural history of type 1 diabetes. *Cold Spring Harb. Perspect. Med.* **2**, 1–18 (2012).

8. Pihoker, C., Gilliam, L. K., Hampe, C. S. & Lernmark, Å. Autoantibodies in Diabetes. **54**, (2005).

9. Bingley, P. J. *et al.* Combined analysis of autoantibodies improves prediction of IDDM in islet cell antibody-positive relatives. *Diabetes* **43**, 1304–1310 (1994).

10. Bingley, P. J. *et al.* Prediction of IDDM in the general population: strategies based on combinations of autoantibody markers. *Diabetes* **46**, 1701–1710 (1997).

11. Fousteri, G., Ippolito, E., Ahmed, R. & Hamad, A. R. A. Beta-cell specific autoantibodies: Are they just an indicator of type 1 diabetes? *Curr. Diabetes Rev.* **13**, 322–329 (2017).

12. Oram, R. A., Sims, E. K. & Evans-Molina, C. Beta cells in type 1 diabetes: mass and function; sleeping or dead? *Diabetologia* **62**, 567–577 (2019).

13. Mănescu, M., Mănescu, I. B. & Grama, A. A Review of Stage 0 Biomarkers in Type 1 Diabetes: The Holy Grail of Early Detection and Prevention? *J. Pers. Med.* **14**, 878 (2024).

14. Preissl, S., Gaulton, K. J. & Ren, B. Characterizing cis-regulatory elements using single-cell epigenomics. *Nat. Rev. Genet.* **24**, 21–43 (2023).

15. Elgamal, R. M. *et al.* An Integrated Map of Cell Type–Specific Gene Expression in Pancreatic Islets. *Diabetes* **72**, 1719–1728 (2023).

16. Bressan, D., Battistoni, G. & Hannon, G. J. The dawn of spatial omics. *Science* **381**,

eabq4964 (2023).

17. Pugliese, A. Insulitis in the pathogenesis of type 1 diabetes. *Pediatr. Diabetes* **17 Suppl 22**,

31–36 (2016).

18. Chiou, J. *et al.* Interpreting type 1 diabetes risk with genetics and single-cell epigenomics.

Nature **594**, 398–402 (2021).

19. Benaglio, P. *et al.* Type 1 diabetes risk genes mediate pancreatic beta cell survival in

response to proinflammatory cytokines. *Cell Genomics* **2**, 100214 (2022).

20. Bingley, P. J., Boulware, D. C. & Krischer, J. P. The implications of autoantibodies to a

single islet antigen in relatives with normal glucose tolerance: development of other

autoantibodies and progression to type 1 diabetes. *Diabetologia* **59**, 542–549 (2016).

21. Foster, T. P. *et al.* Exocrine Pancreas Dysfunction in Type 1 Diabetes. *Endocr. Pract. Off. J.*

Am. Coll. Endocrinol. Am. Assoc. Clin. Endocrinol. **26**, 1505–1513 (2020).

22. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony.

Nat. Methods **16**, 1289–1296 (2019).

23. Klein, D. *et al.* Mapping cells through time and space with moscot. *Nature* (2025)

doi:10.1038/s41586-024-08453-2.

24. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat. Methods* **19**,

171–178 (2022).

25. Haviv, D., Gatie, M., Hadjantonakis, A.-K., Nawy, T. & Pe'er, D. The covariance

environment defines cellular niches for spatial inference. *BioRxiv Prepr. Serv. Biol.*

2023.04.18.537375 (2023) doi:10.1101/2023.04.18.537375.

26. Ali, M. *et al.* GraphCompass: Spatial metrics for differential analyses of cell organization

across conditions. Preprint at https://doi.org/10.1101/2024.02.02.578605 (2024).

27. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

28. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat. Methods* **14**, 975–978 (2017).

29. Zhang, K. *et al.* A single-cell atlas of chromatin accessibility in the human genome. *Cell* **184**, 5985-6001.e19 (2021).

30. Badia-i-Mompel, P. *et al.* decoupleR: ensemble of computational methods to infer biological activities from omics data. *Bioinforma. Adv.* **2**, vbac016 (2022).

31. Mitchell, R. K. *et al.* The transcription factor Pax6 is required for pancreatic β cell identity, glucose-regulated ATP synthesis, and Ca2+ dynamics in adult mice. *J. Biol. Chem.* **292**, 8892–8906 (2017).

32. Jiang, M. *et al.* MIST1 and PTF1 Collaborate in Feed-Forward Regulatory Loops That Maintain the Pancreatic Acinar Phenotype in Adult Mice. *Mol. Cell. Biol.* **36**, 2945–2955 (2016).

33. Elgamal, R. M. *et al.* An Integrated Map of Cell Type–Specific Gene Expression in Pancreatic Islets. *Diabetes* **72**, 1719–1728 (2023).

34. Kaestner, K. H., Powers, A. C., Naji, A., HPAP Consortium & Atkinson, M. A. NIH Initiative to Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The Human Pancreas Analysis Program (HPAP). *Diabetes* **68**, 1394–1402 (2019).

35. Dimitrov, D. *et al.* LIANA+ provides an all-in-one framework for cell–cell communication inference. *Nat. Cell Biol.* **26**, 1613–1622 (2024).

36. Matsui, J., Wakabayashi, T., Asada, M., Yoshimatsu, K. & Okada, M. Stem cell factor/c-kit signaling promotes the survival, migration, and capillary tube formation of human umbilical vein endothelial cells. *J. Biol. Chem.* **279**, 18600–18607 (2004).

37. Wang, C.-H. *et al.* Stem cell factor attenuates vascular smooth muscle apoptosis and increases intimal hyperplasia after vascular injury. *Arterioscler. Thromb. Vasc. Biol.* **27**, 540–547 (2007).

38. Wang, G. *et al.* Integrating genetics with single-cell multiomic measurements across disease

states identifies mechanisms of beta cell dysfunction in type 2 diabetes. *Nat. Genet.* **55**,

984–994 (2023).

39. Pickrell, J. K. Joint analysis of functional genomic data and genome-wide association

studies of 18 human traits. *Am. J. Hum. Genet.* **94**, 559–573 (2014).

40. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat.*

Commun. **12**, 1088 (2021).

41. Li, Z., Wang, T., Liu, P. & Huang, Y. SpatialDM for rapid identification of spatially co-

expressed ligand–receptor and revealing cell–cell communication patterns. *Nat. Commun.*

14, 3995 (2023).

42. Korf, H. *et al.* MIF inhibition interferes with the inflammatory and T cell-stimulatory capacity

of NOD macrophages and delays autoimmune diabetes onset. *PloS One* **12**, e0187455

(2017).

43. Dooley, J. *et al.* Genetic predisposition for beta cell fragility underlies type 1 and type 2

diabetes. *Nat. Genet.* **48**, 519–527 (2016).

44. Liston, A., Todd, J. A. & Lagou, V. Beta-Cell Fragility As a Common Underlying Risk Factor

in Type 1 and Type 2 Diabetes. *Trends Mol. Med.* **23**, 181–194 (2017).

45. Marroquí, L. *et al.* BACH2, a candidate risk gene for type 1 diabetes, regulates apoptosis in

pancreatic β-cells via JNK1 modulation and crosstalk with the candidate gene PTPN2.

Diabetes **63**, 2516–2527 (2014).

46. Santin, I. & Eizirik, D. L. Candidate genes for type 1 diabetes modulate pancreatic islet

inflammation and β-cell apoptosis. *Diabetes Obes. Metab.* **15 Suppl 3**, 71–81 (2013).

47. Santin, I. *et al.* PTPN2, a candidate gene for type 1 diabetes, modulates pancreatic β-cell

apoptosis via regulation of the BH3-only protein Bim. *Diabetes* **60**, 3279–3288 (2011).

48. Størling, J. & Pociot, F. Type 1 Diabetes Candidate Genes Linked to Pancreatic Islet Cell

Inflammation and Beta-Cell Apoptosis. *Genes* **8**, 72 (2017).

49. Colli, M. L., Moore, F., Gurzov, E. N., Ortis, F. & Eizirik, D. L. MDA5 and PTPN2, two

candidate genes for type 1 diabetes, modify pancreatic beta-cell responses to the viral by-

product double-stranded RNA. *Hum. Mol. Genet.* **19**, 135–146 (2010).

50. Tsan, M.-F. & Gao, B. Heat shock proteins and immune system. *J. Leukoc. Biol.* **85**, 905–

910 (2009).

51. Kregel, K. C. Invited Review: Heat shock proteins: modifying factors in physiological stress

responses and acquired thermotolerance. *J. Appl. Physiol.* **92**, 2177–2186 (2002).

52. Bogdani, M. *et al.* Extracellular matrix components in the pathogenesis of type 1 diabetes.

Curr. Diab. Rep. **14**, 552 (2014).

53. Doliba, N. M. *et al.* α Cell dysfunction in islets from nondiabetic, glutamic acid

decarboxylase autoantibody-positive individuals. *J. Clin. Invest.* **132**, e156243 (2022).

54. Tosti, L. *et al.* Single-Nucleus and In Situ RNA–Sequencing Reveal Cell Topographies in the

Human Pancreas. *Gastroenterology* **160**, 1330-1344.e11 (2021).

55. Chiou, J. *et al.* Single-cell chromatin accessibility identifies pancreatic islet cell type- and

state-specific regulatory programs of diabetes risk. *Nat. Genet.* **53**, 455–466 (2021).

56. Thomas, H. E., Trapani, J. A. & Kay, T. W. H. The role of perforin and granzymes in

diabetes. *Cell Death Differ.* **17**, 577–585 (2010).

57. Notkins, A. L. Immunologic and Genetic Factors in Type 1 Diabetes. *J. Biol. Chem.* **277**,

43545–43548 (2002).

58. Goulley, J., Dahl, U., Baeza, N., Mishina, Y. & Edlund, H. BMP4-BMPR1A signaling in beta

cells is required for and augments glucose-stimulated insulin secretion. *Cell Metab.* **5**, 207–

219 (2007).

59. Chmielowiec, J. *et al.* Human pancreatic microenvironment promotes β-cell differentiation

via non-canonical WNT5A/JNK and BMP signaling. *Nat. Commun.* **13**, 1952 (2022).

60. Urizar, A. I. *et al.* Beta cell dysfunction induced by bone morphogenetic protein (BMP)-2 is

associated with histone modifications and decreased NeuroD1 chromatin binding. *Cell*

Death Dis. **14**, 399 (2023).

61. Barbu, A., Lejonklou, M. H. & Skogseid, B. Progranulin Stimulates Proliferation of Mouse

Pancreatic Islet Cells and Is Overexpressed in the Endocrine Pancreatic Tissue of an MEN1

Mouse Model. *Pancreas* **45**, 533–540 (2016).

62. Cheung, P. F. *et al.* Progranulin mediates immune evasion of pancreatic ductal

adenocarcinoma through regulation of MHCI expression. *Nat. Commun.* **13**, 156 (2022).

63. Insel, R. A. *et al.* Staging Presymptomatic Type 1 Diabetes: A Scientific Statement of JDRF,

the Endocrine Society, and the American Diabetes Association. *Diabetes Care* **38**, 1964–

1974 (2015).

64. Battaglia, M. *et al.* Introducing the Endotype Concept to Address the Challenge of Disease

Heterogeneity in Type 1 Diabetes. *Diabetes Care* **43**, 5–12 (2020).

65. Redondo, M. J. & Morgan, N. G. Heterogeneity and endotypes in type 1 diabetes mellitus.

Nat. Rev. Endocrinol. **19**, 542–554 (2023).

66. McGrail, C. *et al.* Genetic discovery and risk prediction for type 1 diabetes in individuals

without high-risk HLA-DR3/DR4 haplotypes. Preprint at

https://doi.org/10.1101/2023.11.11.23298405 (2023).

67. McGrail, C., Sears, T. J., Kudtarkar, P., Carter, H. & Gaulton, K. Genetic association and

machine learning improves discovery and prediction of type 1 diabetes. *MedRxiv Prepr.*

Serv. Health Sci. 2024.07.31.24311310 (2024) doi:10.1101/2024.07.31.24311310.

68. Abedi, M. *et al.* Aberrant TNF signaling in pancreatic lymph nodes of patients with Type 1

Diabetes. Preprint at https://doi.org/10.1101/2024.05.31.596885 (2024).

69. Chiou, J. *et al.* Single-cell chromatin accessibility identifies pancreatic islet cell type– and

state-specific regulatory programs of diabetes risk. *Nat. Genet.* **53**, 455–466 (2021).

70. Thibodeau, A. *et al.* AMULET: a novel read count-based method for effective multiplet

detection from single nucleus ATAC-seq data. *Genome Biol.* **22**, 252 (2021).

71. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat.*

Commun. **8**, 14049 (2017).

72. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-

Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* **8**, 329-337.e4

(2019).

73. Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-based

single-cell RNA sequencing data. *GigaScience* **9**, giaa151 (2020).

74. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

75. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony.

Nat. Methods **16**, 1289–1296 (2019).

76. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

77. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids*

Res. **28**, 27–30 (2000).

78. Kanehisa, M. Toward understanding the origin and evolution of cellular organisms. *Protein*

Sci. Publ. Protein Soc. **28**, 1947–1951 (2019).

79. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for

taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res.* **51**, D587–D592

(2022).

80. Milacic, M. *et al.* The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Res.* **52**,

D672–D678 (2024).

81. Korotkevich, G. *et al.* Fast gene set enrichment analysis. 060012 Preprint at

https://doi.org/10.1101/060012 (2021).

82. Stuart, T., Srivastava, A., Lareau, C. & Satija, R. Multimodal single-cell chromatin analysis

with Signac. *bioRxiv* 1–17 (2020) doi:10.1101/2020.11.09.373613.

83. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

84. Dries, R. *et al.* Giotto: a toolbox for integrative analysis and visualization of spatial

expression data. *Genome Biol.* **22**, 78 (2021).

85. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

86. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for

single-cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).

87. Klein, D. *et al.* Mapping cells through time and space with moscot. Preprint at

https://doi.org/10.1101/2023.05.11.540374 (2023).

88. Nassar, L. R. *et al.* The UCSC Genome Browser database: 2023 update. *Nucleic Acids*

Res. **51**, D1188–D1195 (2023).

89. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589

(2010).

90. McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions.

Nat. Biotechnol. **28**, 495–501 (2010).

91. Frankish, A. *et al.* GENCODE: reference annotation for the human and mouse genomes in

2023. *Nucleic Acids Res.* **51**, D942–D949 (2023).

92. Hothorn, T., Zeileis, A., Farebrother, R. W. & Cummins, C. lmtest: Testing Linear

Regression Models. 0.9-40 https://doi.org/10.32614/CRAN.package.lmtest (1999).

93. Kaestner, K. H., Powers, A. C., Naji, A., HPAP Consortium & Atkinson, M. A. NIH Initiative

to Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The

Human Pancreas Analysis Program (HPAP). *Diabetes* **68**, 1394–1402 (2019).

94. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, (2015).

95. Mootha, V. K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are
coordinately downregulated in human diabetes. *Nat. Genet.* **34**, 267–273 (2003).

96. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**, 15545–15550
(2005).

97. Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res.* **48**, D498–D503
(2020).

98. Griss, J. *et al.* ReactomeGSA - Efficient Multi-Omics Comparative Pathway Analysis. *Mol.*
Cell. Proteomics MCP **19**, 2115–2125 (2020).

99. Fabregat, A. *et al.* Reactome graph database: Efficient access to complex pathway data.
PLoS Comput. Biol. **14**, e1005968 (2018).

100.   Fabregat, A. *et al.* Reactome diagram viewer: data structures and strategies to boost
performance. *Bioinforma. Oxf. Engl.* **34**, 1208–1214 (2018).

101.   Sidiropoulos, K. *et al.* Reactome enhanced pathway visualization. *Bioinforma. Oxf. Engl.*
33, 3461–3467 (2017).

102.   Wu, G. & Haw, R. Functional Interaction Network Construction and Analysis for Disease
Discovery. *Methods Mol. Biol. Clifton NJ* **1558**, 235–253 (2017).

103.   Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res.* **50**,
D687–D692 (2022).

104.   John D. Storey With Contributions From Andrew J. Bass, A. qvalue. Bioconductor
https://doi.org/10.18129/B9.BIOC.QVALUE (2017).

105.   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. **lmerTest** Package: Tests in
Linear Mixed Effects Models. *J. Stat. Softw.* **82**, (2017).

106.   Duttke, S. H., Chang, M. W., Heinz, S. & Benner, C. Identification and dynamic

1720   quantification of regulatory elements using total RNA. *Genome Res.* **29**, 1836–1846 (2019).

1721   107.   Fulco, C. P. *et al.* Activity-by-contact model of enhancer–promoter regulation from

1722   thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664–1669 (2019).

1723   108.   Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between genome

1724   assemblies. *Bioinformatics* **30**, 1006–1007 (2014).

1725   109.   Arda, H. E. *et al.* A Chromatin Basis for Cell Lineage and Disease Risk in the Human

1726   Pancreas. *Cell Syst.* **7**, 310-322.e4 (2018).

1727   110.   Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access

1728   database of transcription factor binding profiles. *Nucleic Acids Res.* **50**, D165–D173 (2022).

1729   111.   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human

1730   immune cell development and intratumoral T cell exhaustion. *Nat. Biotechnol.* **37**, 925–936

1731   (2019).

1732   112.   Schep, A. motifmatchr: Fast Motif Matching in R.

1733   113.   Nasser, J. *et al.* Genome-wide enhancer maps link risk variants to disease genes.

1734   *Nature* **593**, 238–243 (2021).

1735   114.   Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring

1736   transcription-factor-associated accessibility from single-cell epigenomic data. *Nat. Methods*

1737   **14**, 975–978 (2017).

1738   115.   Wakefield, J. Bayes factors for genome-wide association studies: comparison with P-

1739   values. *Genet. Epidemiol.* **33**, 79–86 (2009).

1740   116.   Aylward, A. *et al.* Glucocorticoid signaling in pancreatic islets modulates gene regulatory

1741   programs and genetic risk of type 2 diabetes. *PLoS Genet.* **17**, e1009531 (2021).

1742   117.   UniProt: the Universal Protein Knowledgebase in 2023 | Nucleic Acids Research |

1743   Oxford Academic. https://academic.oup.com/nar/article/51/D1/D523/6835362?login=true.

118. Stelzer, G. *et al.* The GeneCards Suite: From Gene Data Mining to Disease Genome

Sequence Analyses. *Curr. Protoc. Bioinforma.* **54**, 1.30.1-1.30.33 (2016).

119. Safran, M. *et al.* The GeneCards Suite. in *Practical Guide to Life Science Databases*

(eds. Abugessaisa, I. & Kasukawa, T.) 27–56 (Springer Nature, Singapore, 2021).

doi:10.1007/978-981-16-5812-9_2.

120. Steffen Durinck <Biomartdev@Gmail. Com>, W. H. biomaRt. Bioconductor

https://doi.org/10.18129/B9.BIOC.BIOMART (2017).

121. Andrews, S. No Title. *FastQC: a quality control tool for high throughput sequence data.*

http://www.bioinformatics.babraham.ac.uk/projects/fastqc (2010).

122. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast

and bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

123. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-

level estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2015).

1757
1758
1759