Long-term evaluation of the association between dominance, bib size,

sex and age

Running title: Evaluating a possible melanic badge of status

Authors: Nicolas J. Silva[a], Liliana R. Silva[b,c], Pietro B. D'Amelio[a,d], André C.

Ferreira[a,b,c,e], Annick Lucas[a], Rita Covas[b,c,f], Claire Doutrelant[a,f]

Corresponding author: Nicolas J. Silva, silva.nicolas.j@gmail.com

Affiliations:

[a]CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France;

[b]CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos,

Campus de Vairão, Universidade do Porto, 4485-66, Vairão, Portugal;

[c]BIOPOLIS Program in genomics, Biodiversity and Land Planning, CIBIO, Campus

de Vairão, 4485-661, Vairão, Portugal;

[d]Université Paris-Saclay, CNRS, Institut des Neurosciences Paris-Saclay, 91400,

Saclay, France;

[e]Department of Evolutionary Biology and Environmental Studies, University of

Zurich, Zurich, Switzerland;

[f]FitzPatrick Institute of African Ornithology, DST-NRF Centre of Excellence,

University of Cape Town, Rondebosch 7701, South Africa

Acknowledgments

We thank all involved in the long-term data collection and pre-processing over the years, specifically for help with the yearly captures, dominance videos and bib size picture collection. We thank Franck Theron for his management of the sociable weaver long-term database. We thank Margaux Rat, Inês Duarte and Bruna Fonseca for help scoring dominance videos. Additionally, we thank Margaux Rat for the initial development of the dominance and bib size collection and annotation protocols and Samuel Perret for bib size pictures and scoring. De Beers Mining Corporation provided access to Benfontein Nature Reserve and logistical assistance. The sexing and genotyping analyses were conducted at the CTM lab, CIBIO, and we thank Susana Lopes and all the technical staff that conduct the work. This study was funded by ANR to C.D. (France, grants 15-CE32-0012-02 and 19-CE02-0014-01), FCT (Portugal grants IF/01411/2014/CP1256/CT0007 and PTDC/BIA-EVF/5249/2014) ERC to R.C. (EU, Consolidator grant 866489) and by the DST-NRF Centre of Excellence at the Fitzpatrick Institute of African Ornithology University of Cape Town. N.J.S. was supported by the ANR 19-CE02-0014-01. L.R.S. was funded by ERC 866489 and ANR 19-CE02-0014-01. P.B.D.A. was supported by the south African Claude Leon Postdoctoral Fellowship 2019-20 and by the Marie Curie Individual Fellowship "SOCIAL MATCH" N. 896475 and then by the ANR 19CE02-0014-01. A.C.F was funded by University of Zurich Forschungskredit postdoc grant (K-74312-01-01 University of Zurich), Swiss Federal Commission for Scholarships and by ERC grants 866489 to R.C and. 850859 awarded to Damien Farine). R.C. was funded by FCT (IF/01411/2014/CP1256/CT0007 and CEECIND/03451/2018). The sociable weaver project is supported by the French OSU OREME, European Union's Horizon 2020 Research and Innovation Programme under the Grant Agreement

Conflict of interest statement

50   The authors declare no conflict of interest.

Authors' contribution

The study was initiated by C.D. and R.C., and N.J.S., L.R.S., P.B.D.A. and A.C.F.

further contributed to its conceptualisation and execution. All authors contributed to

the general monitoring data. L.R.S. and A.C.F. collected and processed the dominance

score of the birds. N.J.S., L.R.S., A.L., R.C., and C.D. collected and processed bib

data. N.J.S. developed the deep learning model. N.J.S. and P.B.D.A. produced the

statistical analyses. N.J.S. led the writing of the first draft of the manuscript with

contribution from all co-authors, and all authors contributed critically to the drafts and

gave final approval for publication. P.B.D.A., R.C. and C.D. supervised N.J.S. for this

work. R.C. and C.D. administrated the research project and acquired the funding for

this work.

Data availability statement

65   Analyses reported in this article can be reproduced using the data provided by Silva et

66   al. (2025).

Abstract

The association between dominance and phenotypic traits ('badges of status') has been recently questioned, since it has been found to show more variation across populations and time than expected. Using an eleven-year dataset, encompassing more than 1,800 individuals, we studied the association between the size of a melanin-based plumage trait (a black patch under the beak or 'bib'), individual attributes (age and sex) and dominance. Our study model, the sociable weaver (*Philetairus socius*), is a small passerine, which lives year-round in colonies of varying size. Bib sizes were obtained from pictures and automatically pre-processed with an object segmentation deep learning network. We recorded 36,346 dominance-related interactions to estimate dominance hierarchies. We then investigated the effect of sex and age on bib size, and the overall and between-year association of bib size and age with dominance for each sex. We found that bib size increased with age within individuals of both sexes. Age was a strong and consistent predictor of dominance in males, but not in females. Nevertheless, dominance was not correlated with bib size in males and in females. Finally, these relationships did not strongly vary between years. Our results suggest that males and females may use different factors to regulate dominance interactions. Furthermore, our longitudinal dataset allowed us to show that bib size is not a badge of status in our population even though its growth encodes information on age. These results highlight the importance of considering multiple traits over time when studying morphological signalling of dominance hierarchy.

Keywords: Badge of status; Deep learning; Long-term studies; Melanin-based trait; Social species; Status signalling hypothesis

92 **Data availability statement**

93 Analyses reported in this article can be reproduced using the data provided by Silva et

94 al. (2025).

1. Introduction

Competition over resources (e.g., food, territories, mates) often lead to aggressive interactions between individuals. These agonistic events can be very costly, sometimes leading to severe injuries or death (Clutton-Brock et al., 1979; Neat et al., 1996; Riechert, 1998; Sneddon et al., 2002). To limit these costs, many species establish dominance hierarchies (Tibbetts et al., 2022). Several traits or individual attributes, most commonly sex or age, have been found to be linked to the position of individuals in the dominance hierarchy (Tibbetts et al., 2022).

Dominance signalling can rely on intrinsic competitive abilities of individuals and can be displayed through 'badges of status'. These traits can be visual, olfactory or acoustic and should correlate with the individuals' dominance status (Rohwer, 1975, 1982; Senar, 2006; Smith & Harper 2003; Whiting et al., 2003). They allow to resolve conflicts without direct confrontation, because individuals can assess the competitive ability of others before interacting physically with them. These traits are supposed to be honest and several non-mutually exclusive mechanisms have been shown to be part of the evolutionary stability of the honesty of the signal, among which condition-dependence, social punishment, physiological costs, predation costs, or familiarity (Chaine et al., 2018; Searcy & Nowicki, 2010; Senar, 2006). The signalling function of badges of status has been demonstrated both by correlational and experimental studies in a wide range of taxa (e.g., mammals (Burgener et al., 2009; Clutton-Brock & Albon, 1978), birds (Senar, 2006), reptiles (Stapley & Whiting, 2006), and insects (Tibbetts & Lindsay, 2008)).

In birds, a meta-analysis using 47 studies encompassing 23 species found that plumage was linked to dominance across species and plumage types (e.g., carotenoids,

melanic), suggesting that there is overall statistical support for the status signalling

hypothesis across birds (Santos et al., 2011); however, heterogeneity was high,

indicating low generality of that finding across studies. Regarding melanin-based

plumage patches in birds, the textbook example has been the bib size of male house

sparrows *Passer domesticus*, which was found to be correlated with the fighting

abilities of individuals (Nakagawa et al., 2007). However, this example was

challenged in a meta-analysis conducted on 19 studies (Sánchez-Tójar, Nakagawa et

al., 2018), which showed that when including unpublished studies, the overall effect

size of the correlation between dominance and bib size was not different from zero,

and that the overall evidence for the hypothesis in house sparrows has been declining

over time. This meta-analysis revealed a strong publication bias that calls for

reconsideration of the bib size-based status signalling in this species, and for further

investigation of within-species variation in the relationships between badges of status

and dominance rank.

In addition to dominance signalling through phenotypic traits, dominance

hierarchies have also been shown to be associated with individual attributes, most

notably age. The hierarchies associated with individual attributes are called

'convention-based' hierarchies, and in these, age influences various phenotypic traits

and regulates dominance interactions in many taxa (e.g., in birds (Sergio et al., 2009;

Stahl et al., 2001), insects (Taylor et al., 2020), and mammals (Šárová et al., 2013)).

For instance, in chimpanzees (*Pan troglodytes*), female social status is influenced by

age, while males fight to gain status in the male hierarchy, which ultimately increases

their reproductive success (Foerster et al., 2016).

The mechanisms regulating dominance may differ between the sexes because

the benefits (and costs) of dominance are unlikely to be equivalent, and sex-biased

dispersal may lead to stronger social bonds within one of the sexes. In birds, status

traits have been studied mostly in males and less in females (Doutrelant et al., 2020),

but some studies, with relatively small sample sizes, on mutually ornamented species

found an association between dominance and a given ornament in females (e.g.,

Chaine et al., 2011; Crowhurst et al., 2012; Jawor et al., 2004; Murphy et al., 2014;

Watt, 1986). Hence, the selective pressures affecting these ornaments and their

signalling function may be sex-specific. Finally, studies have also found evidence for

a sex-dependent relationship between dominance and age (Foerster et al., 2016).

However, overall sex differences in the mechanism regulating dominance remains to

be explored in depth.

Trait association can also vary in time and according to environmental

conditions (Kvarnemo & Ahnesjo, 1996; Lucchesi et al., 2019), and dominance is no

exception. Therefore, being able to disentangle the effect of confounding factors, such

as sex, age, or phenotypic traits on dominance and the variations through time is

crucial to understand the evolution and the stability of dominance in wild populations.

In a recent review, Tibbetts et al. (2022) argued that long-term field research is

needed to deduce the factors influencing the stability of dominance hierarchies.

However, collecting such data represents a challenge as it requires continuous funding

to cover both logistical and human resources (Cockburn, 2014). Another challenge is

the processing of the large volumes of data collected, which is very time consuming

and costly when extracting information from videos or images (Weinstein, 2018).

Recent approaches based on deep learning methods (LeCun et al., 2015) can help with

image processing, as they allow to conduct time-consuming and repetitive tasks much

faster than humans (e.g., few milliseconds per images classification; Jocher et al.,

2022) and sometimes outperforming the repeatability and performance of humans

(Ditria et al., 2020). In ecology, deep learning methods are increasingly used for

multiple tasks such as identification, or density estimations, among others (Christin et

al. 2019). However, these methods are still rarely used for fine scale measurements of

morphological traits, as it requires accurate detection and precise segmentation of the

phenotypic traits of interest.

In the present study, we examine the relationship between a putative badge of

status (a black plumage 'bib') and dominance, age, and sex in a highly social colonial

and cooperative bird, the sociable weaver (*Philetairus socius*). To this end, we used a

deep learning approach, based on object-segmentation, to automatically extract the

bib size from approximately 13,300 pictures. We then calculated dominance from

more than 36,000 dominance interactions collected over seven years. In this species,

both male and female adults display a black melanin-based bib, with males only

presenting very slightly larger bibs (+0.032 cm²; Acker et al., 2015). Sociable weavers

establish steep dominance hierarchies within their colonies with males being

dominant over females (Rat et al., 2015; Silva et al., 2018) and older males have been

found to be dominant over younger males (Silva master thesis, 2017). Previous short-

term studies on this species found conflicting results regarding the association

between dominance and bib size, with either a positive association in male-male

interactions only (over two years; Rat et al., 2015), or females with larger bibs being

dominant when interactions were computed in both sexes (Silva master thesis, 2017).

Here, we first evaluated how bib size varies in relation to sex and age, and

performed a longitudinal analysis to understand the within individual variation in bib

size, using 11 years of data. Then, over seven years of data, we tested how dominance

related with bib size, age and sex, and whether these associations varied between

years. Since dominance hierarchies in this species are structured by sex (with most

males being dominant over most females), we then explored whether the traits

associated with dominance hierarchies were different when estimating all individual

compared to when estimating same-sex dominance hierarchies. Longitudinal studies

on badges of status are rare and we expected that by using such a large dataset

spanning across time, we could reliably assess whether bib size functions as a badge

of status, as well as capture potential variation over time of the relationship between

dominance, age and bib size.

2. Materials and Methods

2.1. Species and individual information

Sociable weavers are passerines endemic to the semi-arid savannahs of the

Kalahari region of southern Africa. They are small sexually monomorphic birds of

approximately 28g and are colonial facultative cooperative breeders (Maclean, 1973).

The study was conducted at Benfontein Nature Reserve, in South Africa (-28°818S,

24°816E, 1,190m asl), an area of ~15km² of open savannah where up to 15 colonies

are simultaneously monitored as part of a long-term study since 2010 (some colonies

collapsed during the course of the study and new ones were added to the long-term

monitoring).

The study was conducted between 2013 and 2024. Dominance trials were

conducted on seven of the regularly monitored colonies (from 2013 to 2019), while

bib pictures were taken from all colonies (from 2013 to 2024). We define a year as the

period between 1st of September and the 31st of August, as the birds in the study

colonies are captured at the end of the Southern Hemisphere winter, generally in

August-September, and the breeding season usually starts in September-November

(D'Amelio et al., 2024b). Sociable weavers can form very large colonies up to

approximately 500 individuals (Maclean 1973), but colonies in our study area over

the years of this study varied between 7 and 82 birds (mean±SD, 38±17 individuals).

Individuals are ringed with three coloured rings and a numbered metal ring, allowing

individual visual identification. We also take morphological measurements (mass,

tarsus+intertarsal joint's length) and blood samples to sex genetically all individuals

(D'Amelio et al., 2024a; Griffiths et al., 1998).

Exact age is known for all individuals that were born in the studied colonies as

a result of a routine breeding monitoring (these individuals constitute 77% of the

males and 65% of the females in the dataset). For immigrants, which were first

captured as adults, we computed the age by adding the "average minimum age" of

first dispersal to the first date of identification. This average minimum age of first

dispersal in days was quantified for both males and females based on the long-term

capture-recapture database (mean±SD, males: 690±46 days, N=13; females: 727±291

days, N=184). In sociable weavers, females disperse away from their natal colonies to

breed (Covas et al., 2006; van Dijk et al., 2015). Once they establish themselves in a

colony, they generally settle for several years and no longer change colonies

((mean±SD) 2.9±1.6 of years of presence in the study site after moving in from the

outside and starting to breed, N=562 females; Silva et al., in prep). Note that our

estimation of age is more robust for females (N=184) than for males (N=13). All

protocols were approved by the Northern Cape Nature Conservation (permit FAUNA

1638/2015, 0825/2016, 0212/2017, 0684/2019 and 0059/2021) and the Ethics

Committee of the University of Cape Town (permit 2014/V1/RC and 2018/V20/RC).

2.2. Dominance status

Dominance hierarchies were determined within seven colonies each year

(2013-2019) by analysing and scoring agonistic interactions between individuals

feeding on an artificial food source, a plate with mixture of bird seed, placed beneath

each colony. Behavioural observations were performed using a video camera (Sony

Handycam HD) on a tripod 2-3m from the feeder, to record all interactions seen

approximately up to 1m around the feeder. All encounters were quantified as either

aggression, displacement, threat or avoidance (see Table 1 in Silva et al., 2018 and

Rat et al. 2014), and for each we determined the winner and the looser of the

interaction.

Dominance observations were performed most of the time in September at the

end of winter, before the onset of the breeding season (see Table S15 for precise

dates). From 2014 to 2017 and in 2019 observations were conducted in September; in

2013, from September to November; and in 2018, in December. Over these seven

years, we conducted approximately 4h of observations each day, usually starting

between 06:45 and 12:50; and repeated later in the day, between 13:15 and 16:50, and

the food provided was removed between recordings. We obtained a total of 550 hours

of video and each colony was recorded approximately (mean±SD) 15±12 hours per

year (mean±SD, over approximately 6±4 different days per colony). All the videos

per colony per year were recorded within a period length of (mean±SD) 9±9 days per

year (Table S15). A total of 36,346 agonistic interactions (6,869 acts of aggression,

14,856 displacements, 3,459 threats and 11,162 avoidances) were recorded (mean±SD,

approximately 1,010±1,240 interactions per colony per season). In addition, we

scored 682 agonistic interactions for which we were not able to determine the winner

or loser, as both individuals, after initiating a strength assessment, move away

simultaneously from the confrontation.

To assess individual rank and to obtain dominance hierarchies, we used

David's score (David, 1987), which is considered as one of the most appropriate and

reliable methods available (Sánchez-Tójar, Schroeder et al., 2018), as it allows to

assess overall individual success in agonistic interactions (Gammell et al., 2003).

David's score allows to estimate dominance of an individual by considering the

proportion of wins or losses across all the agonistic interactions. This index is also

adjusted by the total number of observations of the individuals and the proportion of

wins for each of them based on the relative number of wins and losses against all the

other individuals in the group. Additionally, this method allows to compare results to

previous studies on dominance behaviour in this species (Rat et al., 2015; Silva

master thesis, 2017). This score is calculated using the "steepness" R package (De

Vries et al., 2006) and standardized per year and colony between zero and one to

allow comparison between individuals of different groups (see Fig. S1 for the

distribution of the dominance scores in the population).

The hierarchies produced here included the majority of the birds captured in

winter for the studied colonies ((mean±SD) 85%±23% of individual within colony

and year observed during dominance data collection; Table S15). The 15% missing

correspond to birds never observed interacting at the feeders or that were not

alive/present at the colony at the recording time. Individuals seen in several colonies

(N=29 individuals) and those seen interacting less than three times in a given colony

and year (N=45 individuals) were excluded from the analyses to avoid prospecting

individuals (i.e., individuals briefly visiting colonies without being part of it).

In addition to these general dominance hierarchies (both-sexes dominance

hierarchies) including all individuals and interactions, we also extracted dominance

hierarchies by considering i) only male-male dominance interactions (N=16,261

interactions between 256 males, the between-males hierarchies) and ii) only female-

female dominance interactions (N=2,651 interactions between 267 females, the

between-females hierarchies).

To obtain reliable and representative dominance hierarchies for steep

dominance hierarchies, like the sociable weavers' (Silva et al., 2018), Sánchez-Tójar,

Schroeder et al. (2018), proposed that the threshold ratio of total number of observed

interactions to total number of individuals should have a minimum of 10. In our study

site, many colonies did not fulfil this criterium. We therefore decided to run our

models both with and without the colonies with a ratio inferior to 10 (except for

females as only one colony fulfilled this criterium). We found that the estimates were

similar between the models with and without the colonies with a ratio inferior to 10

(Table S3 to S12), and therefore decided to include the colonies with a ratio inferior to

10 in the results.

2.3. Bib size and YOLO: annotations, training bib segmentation and model

validation

For all the birds captured, three top view pictures of the bib per individual

were taken in the field (Rat et al., 2015). We collected pictures of the bib of the birds

for 11 years, between 2013 and 2024, between mid-August and mid-September each

year. The birds were handled and held in an apparatus where they were photographed

in a standardized position. Pictures of the bib and videos for dominance estimation of

the birds were collected with less than a month from each other for all years, except

for three colonies of the year 2013 (approximately 1.5 month between data collection)

and four colonies of the year 2018 (approximately 3.5 months between data

collection).

In previous works, bib size was manually analysed using image graphic editor

(The GIMP Development Team, 2019 or Photoshop CS5.1 (Version 12.1), hereafter

referred as "GIMP method"), which is a time-consuming task when dealing with large

datasets. We improved this method by implementing a deep learning approach using

YOLO (You Only Look Once; Redmon et al., 2016) to automatically detect and

extract bib size from the pictures (Supplementary 2 for detailed description of the

method).

We tested the repeatability of the GIMP and YOLO methods by selecting 20

individuals (300 pictures) on five different years for which bib size was measured

with both methods (rptR package version 0.9.22 (Nakagawa & Schielzeth, 2010)).

Results showed high repeatability and high correlations between the two methods

(Table S1), indicating that the bib size measurements obtained with the deep learning

model were robust. Bib size, extracted in pixels from pictures, was transformed to cm².

2.4. Statistical analysis

All statistical analyses were conducted in R version 4.3.1, using mixed models

ran with the lme4 package version 1.1-33 (Bates et al., 2014). We verified whether

model assumptions were met, specifically that a) the residuals followed normality

using Kolmogorov-Smirnov tests, b) the simulated dispersion of the residuals was equal to the observed dispersion, c) there were no more simulated outliers than expected, and d) there was homoscedasticity of the variance at three quantiles (0.25, 0.5, 0.75) between the residuals and predicted values. These diagnostics were performed using the Dharma package version 0.4.6 (Hartig & Lohse, 2022), and, although not marked, some heteroscedasticity was observed and it is detailed in Supplementary 7 and 8. The plots were created with ggplot2 package version 3.4.2 (Wickham, 2014) and ggeffects version 1.3.2 (Lüdecke, 2023).

2.4.1. Factors associated with bib size

To design and interpret the most appropriate structure for the models testing the association between bib size and dominance, we first needed to understand the relationship of bib size with age within sexes (Fig. S2). These variables were found to be associated in a previous study conducted on this species (using distinct datasets: Acker et al., 2015). In addition, we included tarsus length (as a proxy of body size), to correct for a potential correlative effect between bib size and body size (as recommended by Wysocki et al., 2022). We designed a linear mixed model (LMM) with bib size (in cm²) as the response variable, and tarsus length (in mm), age (in days), sex and the interaction between age and sex as fixed factors. Age is often used as a non-linear factor in many model to test for a senescence effect on the association between age and a trait (here bib size). However, here we did not test for such effect due to the low number of old individuals in our dataset. The oldest birds alive recorded on our study site was approximately 16 years old (D'Amelio et al., 2024a), but here our individual are on average three years old ((mean±sd): males - 1,212±865 days old and 20 individuals older than 10 years only; females - 1,057±739 days old and 10 individuals older than 10 years only). Therefore, we decided that including age,

as a non-linear factor, would not be relevant and would not allow to test for floor or ceiling effect. As random factors, we controlled for year, colony and individual identity. Because photographers did not always follow the same protocol and because we changed cameras over the years, we included photographer ID in our initial model, the observer who took the photos (7 observers) and the camera model used (3 cameras). We also added random slopes to the random intercepts to estimate the link between age and bib size for every individual. We estimated random slopes uncorrelated with random intercepts, as we do not have predictions about individuals with larger bib size also having a greater bib size growth when getting older. Using the estimated slopes for each individual, we assessed the percentage of individuals showing positive growth in bib size over the years. Additionally, to evaluate the proportion of variance explained by the random intercepts and random slopes in the model, we computed two alternative models: (i) one excluding the random slopes and (ii) one excluding both the random intercepts and random slopes. The conditional R^2 (i.e., the proportion of variance explained by both the fixed and random effects) was calculated to estimate the proportion of variance explained by the random slopes and random intercepts when removed from the models (Johnson, 2014).

Tarsus length was computed as the average tarsus length measured during the adult life of the individuals (by the same scorer, R.C., which measured the majority of birds, or by others if no measures by R.C. were found for that individual). Three individuals with extreme tarsus length observations were removed.

We used a dataset composed of 1,863 individuals from 19 colonies over eleven years (from 2014 to 2024, N=3,831 observations). We estimated the influence of the number of repeated measures per individuals on the random slopes variance estimation by computing the same model on a subset of individuals with at least three

measures through their life and we obtained similar conclusions regarding the random

slopes analysis (Table S3). In addition, we used 1,000 bootstraps and 1,000

permutations and estimated the between individual adjusted repeatability of bib size

across years, using the 'rptR' package (Stoffel et al., 2017), on 959 individuals seen at

least twice over the study period (2,927 observations). To estimate the repeatability,

we only corrected for the sex of the individuals.

The residuals of the model using the 3,831 observations did not respect the

normality assumption (Kolmogorov–Smirnov test, P<0.001). This non-normality of

the residuals of the model were mainly due to one year photographed by one scorer.

By removing this scorer from the dataset (N=647 observations), the model fitted the

necessary assumptions. Estimates and effects of the different factors ranged from

highly similar to identical between the two models (Table 1 and Table S2). Therefore,

we decided to use the model containing the full dataset.

2.4.2. Overall association between dominance, sex, bib size and age

First, we used 3,000 bootstraps and 3,000 permutations and estimated the

between individual adjusted repeatability of dominance rank within the colonies

across years, using the 'rptR' package (Stoffel et al., 2017), on 184 individuals

observed at least twice over the study period (498 observations). To estimate the

repeatability, we corrected for the sex of the individuals and used a Poisson log-link

model.

Then, we performed a LMM using the dominance score of the individuals as

the response variable and, as fixed factors, bib size, age, sex, tarsus length, body index,

as well as the interactions between bib size and sex, and age and sex. Gaussian

distribution model was used as they fitted particularly well (see Supplementray 6 Fig.

S4). Body index was based on a scaled mass index (Peig & Green, 2009) that uses mass (here collected at the capture events, approximately when dominance data collection was performed) and a body size metric of an individual (here tarsus length, T) with its arithmetic mean value for the studied population (T_0). The method allows to compute a scaled mass as the predicted body mass when $T=T_0$. To visually describe the structure of our model, we designed a Directed Acyclic Graph (DAG) (Fig. S2). DAGs are visualisation tools that help to represent the links between the variables that we intend to test in a model (Wysocki et al., 2022). In this model, bib size was normalized within colony and year when analysing both-sexes dominance hierarchies, and within colony, year and sex when dealing with sex-specific dominance hierarchies. The normalisation allows us to compare relative bib size of the individuals within colonies and does not allow us to interpret bib size at the population level. Two individuals with a David score of one in different colonies might not have the same bib size but potentially be at the same bib size position in the bib size hierarchy within their colonies. Therefore, they would be assigned the same normalized bib size. Normalisation was performed as $\frac{BS_i - BS_{min}}{BS_{max} - BS_{min}}$, where i represents the individual's bib size within the colony, and BS_{min} and BS_{max} respectively the minimum and maximum bib size in cm^2 within a colony one year. Normalized bib size, body index, tarsus length and age were scaled and mean-centered over the whole dataset. Body index and tarsus length were included because they are often related to dominance in other bird species (Jonart et al., 2007; Tibbetts et al., 2022), and including them allowed to estimate the influence of bib size, while accounting for the influence of the individual's condition and its size on dominance score (Fig. S2 for visual representation of the links between dominance and the predictors). As random factors, we controlled for colony (N=7), individual identity

435   (N=409) and year of data collection (N=7, from 2013 to 2019). We included year as a

436   random factor since we expected potential differences in dominance between years

437   (e.g., arising from differences in social composition, resource availability, breeding

438   onset).

439   2.4.3. Between-years association between dominance, sex, bib size and age

We used the same LMM model structure as when testing the overall

association between dominance and bib size, but instead of having "year" as a random

categorical factor, we used it as a fixed categorical factor. Additionally, we included

the interactions between year, normalized bib size and sex, as well as between year,

age and sex. We then performed a post-hoc test with the emmeans package version

1.8.6 (Lenth et al., 2023) to determine if the slopes of the relationship between

dominance, bib size and age were significantly different from zero between years

(Table S13).

3. Results

3.1. Factors associated with bib size

We found that age and bib size were significantly positively associated

(0.02 ± 0.01 (EST±SE, used throughout the Results section except if specified

otherwise), T=2.58, P=0.010; Table 1), and we estimated that between individuals

that differ by one year, the older birds have approximately 2% larger bibs. Within

individuals, we found a positive association between bib size and age for 99.8% of the

birds (Fig. 1), and this relationship was not affected by sex, with bib size increasing

by 1.37% per year within females and 1.21% per year within males. We found a

conditional R² of 0.67 in the model including both random intercepts and random

slopes, and estimated that the individual random intercepts for bib size accounted for

more variance in the model (Conditional R² = 0.64 in the model without random

slopes) than the random slopes of age's effect on bib size per individual (Conditional

R² = 0.26 in the model without random intercepts and slopes). This suggesting that

individuals vary in their average bib size but have a similar pattern of growth through

time. Moreover, we found a small positive significant effect of sex (0.02±0.01,

T=2.58, P=0.010; Table 1) with males having on average 1.29% larger bib size than

females (mean±SD, 1.55±0.03 cm² for females and 1.57±0.03 cm² for males). Finally,

we found a significant positive association between tarsus length and bib size

(0.02±0.00, T=3.48, P<0.001; Fig. S3; Table 1), hence larger individuals have larger

bibs independently of sex. Finally, bib size was found repeatable between individuals

and across years ((estimate [CI]) 0.41 [0.37; 0.44], P = 0.001).

Figure 1: Linear regression of bib size as a function of age. The random slopes and
intercepts of the 1,863 individuals (995 females (red) and 868 males (blue)) were used
to simulate bib size development every 200 days between one to 5,201 days old (i.e.,
more than 14 years).

Table 1: Estimates of the model parameters to test the influence of the size and the age on the bib size using the whole dataset. Bib size was taken once a year at the end of winter and some individuals were collected multiples years. The model was computing using 3,831 observations (995 females and 868 males). Here, 'Age' is either exact age for individuals born on our studied colonies or minimum age for individuals first caught as adults. Conditional R² was of 0.67.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	1.55	0.03	45.91	<0.001
Age	0.05	0.00	9.78	<0.001
Sex - *Male*	0.02	0.01	2.58	0.010
Tarsus	0.02	0.00	4.48	<0.001
Age : **Sex** - *Male*	-0.01	0.01	-0.86	0.392

Random effect	Variance			
Individual ID - *Intercepts* (N=1,863)	0.013			
Individual ID - *Slopes* (N=1,863)	0.001			
Colony (N=19)	0.001			
Year (N=10)	0.002			
Photographer (N=7)	0.004			
Camera (N=3)	0.001			
Residuals	0.012			

3.2. Overall association between dominance, bib size, age and sex

We found a between individual repeatability of dominance rank of 0.38 ([CI] [0.28; 0.49], P=0.001).

Dominance in relation to bib size and sex. We found no significant association between normalized bib size and dominance in the datasets considering both-sexes dominance hierarchies (0±0.01, T=0.19, P=0.852; Table 2; Fig. 2.A) and between-males dominance hierarchies (-0.02±0.01, T=-1.69, P=0.092; Table S5; Fig. 2.C). However, we found a small positive significant correlation between bib size and dominance in between-females dominance hierarchies (0.04±0.02, T=2.34, P=0.020; Table S7; Fig. 2.E). Females being ranked 25% higher in the normalized bib size

ranks have an estimated 8% higher dominance score. However, the between-years

analysis below shows that the relationship is mainly influenced by two years with

very low sample size (Table S13) and when removing these two years the correlation

is weak and non-significant (0.02±0.02, T=1.06, P=0.290, Table S14).

Table 2: Estimates of the model testing the overall influence of bib size, sex and age
on dominance in both-sexes dominance hierarchies, with colonies with a ratio inferior
to 10 of interactions to the number of individuals. The model was computed with 723
observations. Association of dominance with age, bib size and sex is visually
represented in Fig. 2.A and 2.B.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.26	0.03	9.00	<0.001
Bib size	0.00	0.01	0.19	0.852
Sex - *Male*	0.30	0.02	18.61	<0.001
Age	0.01	0.01	0.65	0.515
Body condition	0.02	0.01	1.89	0.059
Tarsus	0.01	0.01	1.13	0.259
Bib size : **Sex** - *Male*	-0.01	0.01	-0.42	0.674
Age : **Sex** - *Male*	0.11	0.02	7.20	<0.001
Random effect	**Variance**			
Individual ID (N=409)	0.010			
Year (N=7)	0.002			
Colony (N=7)	0.003			
Residuals	0.022			

<u>Dominance in relation to age and sex.</u> In both-sexes dominance hierarchies,

we found a strong significant link between sex and dominance, indicating that males

are approximately two times more dominant than females (0.30±0.02, T=18.61,

P<0.001; Table 2). We also found a strong association between dominance and the

interaction of age and sex (0.11±0.02, T=7.2, P<0.001; Table 2), indicating a different

relationship between age and dominance for males and females (Fig. 2.B), specifically

a much steeper slope for males. Dominance score is 1.36% higher in females and

9.39% higher in males when comparing two different individuals that differ by one

year in age (Table 2). This was found also in the sex-specific interactions' datasets,

where we observed a different association between age and dominance, with a

positive, strong and significant link in between-males dominance hierarchies

(0.16±0.01, T=11.77, P<0.001; Table S5; Fig. 2.D), while there was no statistically

significant relation in between-females dominance hierarchies (-0.01±0.02, T=-0.35,

P=0.725; Table S7; Fig. 2.F).

Figure 2: Association between: (A,C,E) dominance and normalized bib size; and (B,D,F) dominance and age. Results were obtained using both-sexes dominance hierarchies (A,B), between-females dominance hierarchies (C,D) and between-males dominance hierarchies (E,F). Dominance score and bib size were normalized within dominance hierarchies (both-sexes, between-females, and between-males), colony and year of data collection. Females are represented in red (square dots and dashed line) and males are represented in blue (circle dots and solid line).

3.3. Between-years association between dominance, bib size, age and sex

Dominance in relation to bib size, sex and year. We found no evidence that the relationship between dominance and normalized bib size varied between years, with a few exceptions. Out of the seven years used in each analysis, the only statistically significant associations we found were a slightly negative relationship for one year in males in both-sexes dominance hierarchies (-0.06±0.03, T=-2.18, P=0.030; Table S13; Fig. S5.A in year 2016), a slightly negative relationship for one year in between-males dominance hierarchies (-0.06±0.03, T=-2.04, P=0.042 in year 2014; Table S13; Fig. S5.E), and positive relationships for two years in between-females dominance hierarchies (0.14±0.06, T=2.34, P=0.020 in year 2013, and 0.17±0.08, T=2.25, P=0.026 in year 2016; Table S13; Fig. S5.C). However, the sample size was limited for these latter years, with respectively 19 and 12 individuals only (the other years presenting between 17 and 84 individuals, Table S13), and so these relationships remain unclear.

Dominance in relation to age, sex and year. We found no between-years variation in the association between dominance and age in both-sexes dominance hierarchies and in between-males dominance hierarchies, with dominance being positively associated with age in all years for males and never associated for females (Table S13; Fig. S5.B and S5.F). In between-females dominance hierarchies, dominance was negatively associated with age in one year only (-0.07±0.03, T=-2.15, P=0.033; Table S13; Fig. S5.D in year 2017).

4. Discussion

We examined whether bib size and age relate to dominance at artificial feeders in sociable weavers and how these relationships vary between sexes and years. We found that, in both sexes, bib grows with age, and that larger birds have larger bibs. We also confirmed that males were dominant over females. However, normalized bib size was not found to be associated with dominance in any of the hierarchies (except in between-females dominance hierarchies for two years, but based on very low sample size). In addition, we confirmed that age was strongly associated with dominance in males a pattern that was consistent between years. In contrast, age was not found to be associated with female dominance. Our study is one of the largest long-term studies investigating correlates of dominance with a potential badge of status, providing robust results to understand the association between dominance and bib size, and the variation in its associated dominance correlates, given the low uncertainty in the variation obtained between years (see also Parker (2013); Sánchez-Tójar, Nakagawa et al. (2018); Tibbetts et al. (2022)). While previous studies found positive associations between dominance and bib size with only one year (included in this study (season 2013/2014; Silva master thesis, 2017) and two years of data (not included in this study; Rat et al. 2015), our analyses, on a much larger dataset, led to different conclusions. Finally, this study showed a sexual difference in the factor used to regulate dominance interactions.

4.1. Association between dominance, bib size and sex

Relative bib size was clearly not associated with dominance, whether when considering all birds or only males, and this did not change between years. We found

an association within females, but the association was weak and unclear, with small effect sizes, and we had a smaller number of female-female interactions in the hierarchies (32 out of 33 colonies were below the recommended threshold of ratio of 10 of interactions to number of individuals in the group; Sánchez-Tójar, Schroeder et al. 2018). In addition, this association was only present in two years, which had among the lowest sample sizes of all years tested here (respectively N=19 and N=12 females). When these two years are removed there is no longer evidence of an association between relative bib size and dominance in females.

Several factors might explain why bib size does not function as a badge of status in this population. First, the sampling period could influence our results, as the association between bib size, age and dominance could be used by birds only in some parts of the season or some seasons depending on the social and environmental contexts. For example, a density-dependent effect in the association between badge of status and dominance has been found in the black-crested titmouse (*Baeolophus atricristatus*), where males with larger black crests had more access to feeding stations when competition was high but did not display such advantages when population density decreased and competition was low (Queller & Murphy, 2017). Here, the dominance scores were always computed at the end of winter, usually before the onset of the breeding season. Therefore, the association between dominance and bib size should be investigated at another moment of the year to investigate the possibility that bib size is a seasonal badge of status. Second, here we only tested the size of the bib, but other bib features (e.g., saturation, intensity) could signal the dominance abilities (Santos et al., 2011). For instance, in black-capped chickadees (*Poecile atricapilla*), high-ranking males exhibit darker plumages than subordinates (Poston et al., 2005). Finally, when facing known opponents, animals are

expected to use information on the opponents' previous behaviour (i.e., their

'reputation') to decide how to interact (Senar, 2006; Tibbetts & Dale 2007). For

instance, Chaine et al. (2018) experimentally manipulated plumage patch size of

golden-crowned sparrows (*Zonotrichia atricapilla*) and showed that manipulation

only affected interactions between unknown individuals. Sociable weaver groups live

together all year long, interacting on a daily basis, and this familiarity between

individuals could remove the need for badges of status. In addition, male sociable

weavers are philopatric, therefore knowing each other since birth. A previous study

showed that the number of aggressive interactions between two individuals was

negatively correlated with relatedness (Rat et al., 2015). Therefore, the social system

of sociable weavers might not promote the use of badges of status in males. However,

badges of status might still be important for individuals that move between colonies,

especially females, which disperse to reproduce in other colonies (Covas et al., 2006;

van Dijk et al., 2015). Badges of status could also play a more important role among

the residents of very large colonies where individuals might not be able to keep track

of all residents. Specifically, some sociable weaver colonies house several hundreds

of birds (Maclean, 1973), while our study was conducted in colonies that varied in

size between 7 and 82 residents.

Our results also contrast with previous results found in the same species but on

smaller datasets, in which positive association between dominance and bib size was

found in males (Rat et al., 2015) and in females (Silva master thesis, 2017).

Accumulating evidence shows that phenotypic traits and their signalling function may

vary depending on external forces (e.g., availability of food, mates), which vary over

time and can ultimately influence competition intensity (Kvarnemo & Ahnesjo, 1996;

Lucchesi et al., 2019). Long-term studies integrate the wide ecological and social

variation that characterise the environments experienced by individuals (Clutton-Brock & Sheldon, 2010a, 2010b; Cockburn, 2014) and have provided several examples where limited temporal sampling could have resulted in inaccurate generalisation of association patterns (Clutton-Brock & Sheldon, 2010a; Cockburn, 2014; Cornwallis & Uller, 2010; Doutrelant et al., 2020). This study is an additional example illustrating the power of long-term studies in drawing general patterns of traits association.

4.2. Association between dominance, age and sex

We found a strong sexual difference in the association between age and dominance. Evidence of age-based dominance hierarchies (also referred to as social queuing) is common in many taxa and is particularly common in species that form long-term associations with relatives (Tibbetts et al., 2022), like sociable weavers. In ungulates, for instance, individuals, that are young and more inexperienced learn to be subordinates to older and stronger individuals, and do not challenge the hierarchy when getting older, resulting in age-based hierarchies (Favre et al., 2008; Thouless & Guiness, 1986). In sociable weavers, males know each other since birth and a mechanism similar to the ungulates could explain their age-based hierarchy. In contrast, females disperse to reproduce and therefore may not be able to use previous information about the status of dominant or subordinate of their opponents. Females may resort to alternative cues to infer fighting abilities of other females to decide how to interact with them, or may not need to enter in competition with them. It aligns with the low number of female-female interactions observed in this study, that is particularly low compared to the sex ratio observed in the population (Doutrelant et

al., 2004). However, it is important to note that because of this natal dispersal of the

females, our estimation of their age is less accurate than for males.

Social queuing has been suggested to also arise when the costs of reaching a

higher position in the hierarchy are higher than the benefits of reaching that position

(Broom et al., 2009). For instance, in chimpanzees, dominant males benefit from

higher mating and reproductive success, therefore males fight for higher ranks

(Foerster et al., 2016). However, females do not obtain fitness benefits from higher

dominance rank in the hierarchy and intense fights would only bring fitness costs,

resulting in age-dominance hierarchies in females (Foerster et al., 2016). In sociable

weavers, males of the same colony are more related to each other than females and,

within males' fights could either bring direct fitness costs (if the fight is lost), but also

indirect fitness costs (if the fight is won against a related individual). This could

suggest that, for males, the fitness costs of reaching a higher dominance rank might

not be compensated by the fitness benefits of reaching this rank, leading to age-based

dominance hierarchies within males. In male sociable weavers, dominant individuals

are more likely to access food and reproduce (Rat, 2015; Silva et al., 2018), but

further information of the exact fitness costs and benefits on the long-term are needed

to explore the origin of the age-based dominance hierarchies in males.

4.3. Association between bib size, age and sex

In agreement with a previous study (Acker et al. 2015), we found that bib size

increases with age in both sexes. Age-related changes in adult plumage have been

described in several species (e.g., Dreiss & Roulin, 2010; Wiebe & Vitousek, 2015).

However, the bib size growth between years is small, only slightly over 1% for both

sexes, and the perception by the individuals of such difference is questionable. Interestingly, we found that the growth rate of the bib is very similar between individuals, but individuals vary in bib size, indicating that the mechanisms behind bib size first growth varied between individuals. This last result is in line with the strong determination that genetics and development are expected to have on bird melanic patches (Roulin & Ducrest, 2013). Finally, we detected an effect of sex, with a difference of approximately +0.03cm² (approximately +2%) for the bib size of males compared to females. This sexual difference is in line with the one found previously in this species (Acker et al., 2015), but again, the perception of this difference by the birds would need to be demonstrated.

4.4. The growing importance of artificial intelligence in ecology

Finally, our methodology for measuring automatically thousands of bibs from pictures adds to the growing examples that illustrate the effectiveness and accuracy of deep learning-based methods to detect and quantify morphometrical traits, and how this can facilitate the process of extracting large amounts of data from images (i.e., here the mode processed 124 times faster than a human's; based on 30 pictures). This approach can be used in different taxa, from museum specimens (Miele et al., 2020), to live ones (He et al., 2023), but it is still rarely used for such tasks in evolutionary ecology. While in many cases developing deep learning methods would be too demanding technically or too time consuming, data extraction from existing long-term datasets, as was the case here, is particularly well suited to be processed automatically (e.g., because of already laballed large amount of data).

 5. Conclusion

In this long-term study capitalising on machine learning to assess the badge of

status hypothesis in sociable weavers we found no convincing evidence of it, although

for female-female interactions further investigation is needed. These results align with

the previous recommendation of re-evaluating the status signalling hypothesis of

melanic patches as well as evaluating the temporal variation of the correlates with

dominance. By using multiple years of data, we were able to reach strong conclusions

regarding the stability of bib size and age with dominance in our species. The sexual

differences in the association between dominance and age for the two sexes suggest

that male and female dominance interactions are not regulated by the same factors.

These results show the importance of conducting analyses on both sexes when trying

to understand the role of phenotypic traits on dominance, and suggest that males and

females may be subjected to different selective pressures.

References

Acker, P., Grégoire, A., Rat, M., Spottiswoode, C. N., van Dijk, R. E., Paquet, M.,
Kaden, J. C., Pradel, R., Hatchwell, B. J., Covas, R., & Doutrelant, C. (2015).
Disruptive viability selection on a black plumage trait associated with dominance.
Journal of Evolutionary Biology, *28*(11), 2027–2041.
https://doi.org/10.1111/jeb.12717

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2014). Fitting linear mixed-
effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.
https://doi.org/10.18637/jss.v067.i01

Broom, M., Koenig, A., & Borries, C. (2009). Variation in dominance hierarchies
among group-living animals: Modeling stability and the likelihood of coalitions.
Behavioral Ecology, *20*(4), 844–855. https://doi.org/10.1093/beheco/arp069

Burgener, N., Dehnhard, M., Hofer, H., & East, M. L. (2009). Does anal gland scent
signal identity in the spotted hyaena? *Animal Behaviour*, *77*(3), 707–715.
https://doi.org/10.1016/j.anbehav.2008.11.022

Chaine, A. S., Shizuka, D., Block, T. A., Zhang, L., & Lyon, B. E. (2018). Manipulating badges of status only fools strangers. *Ecology Letters*, *21*(10), 1477–1485. https://doi.org/10.1111/ele.13128

Chaine, A. S., Tjernell, K. A., Shizuka, D., & Lyon, B. E. (2011). Sparrows use multiple status signals in winter social flocks. *Animal Behaviour*, *81*(2), 447–453. https://doi.org/10.1016/j.anbehav.2010.11.016

Christin, S., Hervet, É., & Lecomte, N. (2019). Applications for deep learning in ecology. *Methods in Ecology and Evolution*, *10*(10), 1632–1644. https://doi.org/10.1111/2041-210X.13256

Clutton-Brock, T. H., & Albon, S. D. (1978). The roaring of red deer and the evolution of honest advertisement. *Behaviour*, *69*(3–4), 145–170. https://doi.org/10.1163/156853979X00449

Clutton-Brock, T. H., Albon, S. D., Gibson, R. M., & Guinness, F. E. (1979). The logical stag: adaptive aspects of fighting in red deer (Cervus elaphus L.). *Animal Behaviour*, *27*, 211–225. https://doi.org/10.2307/3492065

Clutton-Brock, T. H., & Sheldon, B. C. (2010a). Individuals and populations: The role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends in Ecology and Evolution*, *25*(10), 562–573. https://doi.org/10.1016/j.tree.2010.08.002

Clutton-Brock, T. H., & Sheldon, B. C. (2010b). The seven ages of pan. *Science*, *327*(5970), 1207–1208. https://doi.org/10.1126/science.1187796

Cockburn, A. (2014). Behavioral ecology as big science: 25 years of asking the same questions. *Behavioral Ecology*, *25*(6), 1283–1286. https://doi.org/10.1093/beheco/aru177

Cornwallis, C. K., & Uller, T. (2010). Towards an evolutionary ecology of sexual traits. *Trends in Ecology and Evolution*, *25*(3), 145–152. https://doi.org/10.1016/j.tree.2009.09.008

Covas, R., Dalecky, A., Caizergues, A., & Doutrelant, C. (2006). Kin associations and direct vs indirect fitness benefits in colonial cooperatively breeding sociable weavers Philetairus socius. *Behavioral Ecology and Sociobiology*, *60*, 323–331. https://doi.org/10.1007/s00265-006-0168-2

Crowhurst, C. J., Zanollo, V., Griggio, M., Robertson, J., & Kleindorfer, S. (2012). White flank spots signal feeding dominance in female diamond firetails, Stagonopleura guttata. *Ethology*, *118*(1), 63–75. https://doi.org/10.1111/j.1439-0310.2011.01986.x

D'Amelio, P. B., Covas, R., Ferreira, A. C., Fortuna, R., Silva, L. R., Theron, F., Rybak, F., & Doutrelant, C. (2024a). Benefits of pair-bond duration on reproduction in a lifelong monogamous cooperative passerine. *The American Naturalist*, *203*(5), 576–589. https://doi.org/10.1086/729436

D'Amelio, P. B., Silva, N. J., Silva, L. R., Ferreira, A. C., Theron, F., Doutrelant, C., & Covas, R. (2024b). Highly variable breeding phenology in an arid region colonial cooperative breeder, the Sociable Weaver Philetairus socius. *Ostrich*, *95*, 1–7. https://doi.org/10.2989/00306525.2024.2328654

David, H. A. (1987). Ranking from unbalanced paired-comparison data. *Biometrika*, *74*(2), 432–436. https://doi.org/10.1093/biomet/74.2.432

De Vries, H., Stevens, J. M. G., & Vervaecke, H. (2006). Measuring and testing the steepness of dominance hierarchies. *Animal Behaviour*, *71*(3), 585–592. https://doi.org/10.1016/j.anbehav.2005.05.015

Ditria, E. M., Lopez-Marcano, S., Sievers, M., Jinks, E. L., Brown, C. J., & Connolly, R. M. (2020). Automating the analysis of fish abundance using object detection: optimizing animal ecology with deep learning. *Frontiers in Marine Science*, *7*, 1–9. https://doi.org/10.3389/fmars.2020.00429

Doutrelant, C., Covas, R., Caizergues, A., & Du Plessis, M. (2004). Unexpected sex ratio adjustment in a colonial cooperative bird: pairs with helpers produce more of the helping sex whereas pairs without helpers do not. *Behavioral Ecology and Sociobiology*, *56*(2), 149-154. https://doi.org/10.1007/s00265-004-0756-y

Doutrelant, C., Fargevieille, A., & Grégoire, A. (2020). Evolution of female coloration: What have we learned from birds in general and blue tits in particular. *Advances in the Study of Behavior*, *52*, 123–202. https://doi.org/10.1016/bs.asb.2020.03.001

Dreiss, A. N., & Roulin, A. (2010). Age-related change in melanin-based coloration of Barn owls (Tyto alba): Females that become more female-like and males that become more male-like perform better. *Biological Journal of the Linnean Society*, *101*(3), 689–704. https://doi.org/10.1111/j.1095-8312.2010.01503.x

Favre, M., Martin, J. G. A., & Festa-Bianchet, M. (2008). Determinants and life-history consequences of social dominance in bighorn ewes. *Animal Behaviour*, *76*(4), 1373–1380. https://doi.org/10.1016/j.anbehav.2008.07.003

Foerster, S., Franz, M., Murray, C. M., Gilby, I. C., Feldblum, J. T., Walker, K. K., & Pusey, A. E. (2016). Chimpanzee females queue but males compete for social status. *Scientific Reports*, *6*, 1–11. https://doi.org/10.1038/srep35404

Gammell, M. P., De Vries, H., Jennings, D. J., Carlin, C. M., & Hayden, T. J. (2003). David's score: a more appropriate dominance ranking method than Clutton-Brock et al.'s index. *Animal Behaviour*, *66*(3), 601–605. https://doi.org/10.1006/anbe.2003.2226

Griffiths, R., Double C., M., Orr, K., & Dawson J. G., R. (1998). A DNA test to sex most birds. *Molecular Ecology*, *7*, 1071–1075. https://doi.org/10.1046/j.1365-294x.1998.00389.x

Hartig, F., & Lohse, L. (2022). *R package DHARMa. Version 0.4.6* (pp. 1–65). https://doi.org/10.32614/CRAN.package.DHARMa

He, Y., Cooney, C. R., Maddock, S., & Thomas, G. H. (2023). Using pose estimation to identify regions and points on natural history specimens. *PLoS Computational Biology*, *19*(2), 1–23. https://doi.org/10.1371/journal.pcbi.1010933

Jawor, J. M., Gray, N., Beall, S. M., & Breitwisch, R. (2004). Multiple ornaments correlate with aspects of condition and behaviour in female northern cardinals, Cardinalis cardinalis. *Animal Behaviour*, *67*(5), 875–882. https://doi.org/10.1016/j.anbehav.2003.05.015

Jocher, G., Chaurasia, A., Stoken, A., Borovec, J., Kwon, Y., Michael, K., Fang, J., Yifu, Z., Wong, C., & Montes, D. (2022). ultralytics/yolov5: v7.0 - YOLOv5 SOTA Realtime Instance Segmentation. In *Zenodo*. https://doi.org/10.5281/zenodo.7347926

Johnson, P. C. (2014). Extension of Nakagawa & Schielzeth's R2GLMM to random slopes models. *Methods in ecology and evolution*, *5*(9), 944-946. https://doi.org/10.1111/2041-210X.12225

Jonart, L. M., Hill, G. E., & Badyaev, A. V. (2007). Fighting ability and motivation: determinants of dominance and contest strategies in females of a passerine bird. *Animal Behaviour*, *74*(6), 1675–1681. https://doi.org/10.1016/j.anbehav.2007.03.012

Kvarnemo, C., & Ahnesjö, I. (1996). The dynamics of operational sex ratios and competition for mates. *Trends in Ecology and Evolution*, *11*(10), 404–408. https://doi.org/10.1016/0169-5347(96)10056-2

LeCun, Y., Bengio, Y., & Hinton, G. (2015). Deep learning. *Nature, 521*(7553), 436-444. https://doi.org/10.1038/nature14539

Lenth, R. L., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., & Singmann, H. (2023). *R package emmeans*. https://doi.org/10.32614/CRAN.package.emmeans

Lucchesi, S., Cheng, L., Janmaat, K., Mundry, R., Pisor, A., & Surbeck, M. (2020). Beyond the group: how food, mates, and group size influence intergroup encounters in wild bonobos. *Behavioral Ecology*, *31*(2), 519–532. https://doi.org/10.1093/BEHECO/ARZ214

Lüdecke, D. (2023). *R package ggeffects*. https://doi.org/10.32614/CRAN.package.ggeffects

Maclean, G. L. (1973). The sociable weaver, part 1: description, distribution, dispersion and populations. *Ostrich*, *44*(3–4), 176–190. https://doi.org/10.1080/00306525.1973.9639158

Miele, V., Dussert, G., Cucchi, T., & Renaud, S. (2020). Deep learning for species identification of modern and fossil rodent molars. *BioRxiv*, 2020.08.20.259176. https://doi.org/10.1101/2020.08.20.259176

Murphy, T. G., West, J. A., Pham, T. T., Cevallos, L. M., Simpson, R. K., & Tarvin, K. A. (2014). Same trait, different receiver response: Unlike females, male American goldfinches do not signal status with bill colour. *Animal Behaviour*, *93*, 121–127. https://doi.org/10.1016/j.anbehav.2014.04.034

Nakagawa, S., Ockendon, N., Gillespie, D. O., Hatchwell, B. J., & Burke, T. (2007). Assessing the function of house sparrows' bib size using a flexible meta-analysis method. *Behavioral Ecology*, *18*(5), 831-840. https://doi.org/10.1093/beheco/arm050

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical guide for biologists. *Biological Reviews*, *85*(4), 935–956. https://doi.org/10.1111/j.1469-185X.2010.00141.x

Neat, F. C., Taylor, A. C., & Huntingford, F. A. (1998). Proximate costs of fighting in male cichlid fish: The role of injuries and energy metabolism. *Animal Behaviour*, *55*(4), 875–882. https://doi.org/10.1006/anbe.1997.0668

Parker, T. H. (2013). What do we really know about the signalling role of plumage colour in blue tits? A case study of impediments to progress in evolutionary biology. *Biological Reviews*, *88*(3), 511–536.https://doi.org/10.1111/brv.12013

Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from mass/length data: The scaled mass index as an alternative method. *Oikos*, *118*(12), 1883–1891. https://doi.org/10.1111/j.1600-0706.2009.17643.x

Poston, J. P., Hasselquist, D., Stewart, I. R. K., & Westneat, D. F. (2005). Dietary amino acids influence plumage traits and immune responses of male house sparrows, Passer domesticus, but not as expected. *Animal Behaviour, 70*(5), 1171–1181. https://doi.org/10.1016/j.anbehav.2005.02.015

Queller, P. S., & Murphy, T. G. (2017). Seasonal variation in the utility of a status signaling system: Plumage ornament predicts foraging success only during periods of high competition. *PLoS ONE, 12*(10), 1–14. https://doi.org/10.1371/journal.pone.0185584

Rat, M. (2015). *Dominance, social organisation and cooperation in the sociable weavers (Philetarius socius)* (Issue June 2015). http://hdl.handle.net/11427/16714

Rat, M., van Dijk, R. E., Covas, R., & Doutrelant, C. (2015). Dominance hierarchies and associated signalling in a cooperative passerine. *Behavioral Ecology and Sociobiology*, *69*, 437–448. https://doi.org/10.1007/s00265-014-1856-y

Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You Only Look Once: Unified, Real-Time Object Detection. *Proceedings of the IEEE Conference on*

Computer Vision and Pattern Recognition (CVPR), 779–788.
https://doi.org/10.48550/arXiv.1506.02640

Rohwer, S. (1975). The Social Significance of Avian Winter Plumage Variability.
Evolution, *29*(4), 593–610. https://doi.org/10.2307/2407071

Rohwer, S. (1982). The evolution of reliable and unreliable badges of fighting ability.
American Zoologist, *22*(3), 531–546. https://doi.org/10.1093/icb/22.3.531

Roulin, A., & Ducrest, A. L. (2013). Genetics of colouration in birds. *Seminars in*
Cell and Developmental Biology, *24*(6–7), 594–608.
https://doi.org/10.1016/j.semcdb.2013.05.005

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S.,
Girndt, A., Bókony, V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T.,
& Schroeder, J. (2018). Meta-analysis challenges a textbook example of status
signalling and demonstrates publication bias. *ELife*, *7*, 1–26.
https://doi.org/10.7554/eLife.37385

Sánchez-Tójar, A., Schroeder, J., & Farine, D. R. (2018). A practical guide for
inferring reliable dominance hierarchies and estimating their uncertainty. *Journal*
of Animal Ecology, *87*(3), 594–608. https://doi.org/10.1111/1365-2656.12776

Santos, E. S. A., Scheck, D., & Nakagawa, S. (2011). Dominance and plumage traits:
Meta-analysis and metaregression analysis. *Animal Behaviour*, *82*(1), 3–19.
https://doi.org/10.1016/j.anbehav.2011.03.022

Šárová, R., Špinka, M., Stěhulová, I., Ceacero, F., Šimečková, M., & Kotrba, R.
(2013). Pay respect to the elders: Age, more than body mass, determines
dominance in female beef cattle. *Animal Behaviour*, *86*(6), 1315–1323.
https://doi.org/10.1016/j.anbehav.2013.10.002

Searcy, W. A., & Nowicki, S. (2010). The evolution of animal communication:
reliability and deception in signaling systems. *In The Evolution of Animal*
Communication. Princeton University Press

Senar, J. C. (2006). Bird colors as intrasexual signals of aggression and dominance.
Bird Coloration, *2*, 125–193. https://doi.org/10.2307/j.ctv22jnr8k.6

Sergio, F., Blas, J., & Hiraldo, F. (2009). Predictors of floater status in a long-lived
bird: A cross-sectional and longitudinal test of hypotheses. *Journal of Animal*
Ecology, *78*(1), 109–118. https://doi.org/10.1111/j.1365-2656.2008.01484.x

Silk, J. B. (2019). Hyena politics: The dynamics of dynasties. *Proceedings of the*
National Academy of Sciences of the United States of America, *116*(18), 8644–
8645. https://doi.org/10.1073/pnas.1903407116

Silva, L. R. (2017). Assessing the physiological costs of dominance in a highly social
bird. *Master Thesis*. https://repositorio-
aberto.up.pt/bitstream/10216/90951/2/176196.pdf

Silva, L. R., Lardy, S., Ferreira, A. C., Rey, B., Doutrelant, C., & Covas, R. (2018).
Females pay the oxidative cost of dominance in a highly social bird. *Animal*
Behaviour, 144, 135–146. https://doi.org/10.1016/j.anbehav.2018.08.006

Silva, Nicolas João; Silva, Liliana R.; D'Amelio, Pietro B., Ferreira, A. C., Covas, R.,
& Doutrelant, C. (2025). Data from: Long-term evaluation of the association
between dominance, bib size, sex and age. Behavioral Ecology.
https://doi.org/10.5061/dryad.v15dv4290

Smith, J. M., & Harper, D. (2003). *Animal signals*. Oxford University Press.

Sneddon, L. U., Huntingford, F. A., & Taylor, A. C. (1998). Impact of an ecological
factor on the costs of resource acquisition: Fighting and metabolic physiology of
crabs. *Functional Ecology, 12*(5), 808–815. https://doi.org/10.1046/j.1365-
2435.1998.00249.x

Stahl, J., Tolsma, P. H., Loonen, M. J. J. E., & Drent, R. H. (2001). Subordinates
explore but dominants profit: Resource competition in high arctic barnacle goose
flocks. *Animal Behaviour, 61*(1), 257–264.
https://doi.org/10.1006/anbe.2000.1564

Stapley, J., & Whiting, M. J. (2006). Ultraviolet signals fighting ability in a lizard.
Biology Letters, 2(2), 169–172. https://doi.org/10.1098/rsbl.2005.0419

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation
and variance decomposition by generalized linear mixed-effects models.
Methods in Ecology and Evolution, 8(11), 1639–1644.
https://doi.org/10.1111/2041-210X.12797

Taylor, B. A., Cini, A., Cervo, R., Reuter, M., & Sumner, S. (2020). Queen
succession conflict in the paper wasp Polistes dominula is mitigated by age-
based convention. *Behavioral Ecology, 31*(4), 992–1002.
https://doi.org/10.1093/beheco/araa045

Thouless, C. R., & Guinness, F. E. (1986). Conflict between red deer hinds: the
winner always wins. *Animal Behaviour, 34*(4), 1166–1171.
https://doi.org/10.1016/S0003-3472(86)80176-2

Tibbetts, E. A., & Dale, J. (2007). Individual recognition: it is good to be different.
Trends in Ecology and Evolution, 22(10), 529–537.
https://doi.org/10.1016/j.tree.2007.09.001

Tibbetts, E. A., & Lindsay, R. (2008). Visual signals of status and rival assessment in
Polistes dominulus paper wasps. *Biology Letters, 4*(3), 237–239.
https://doi.org/10.1098/rsbl.2008.0048

Tibbetts, E. A., Pardo-Sanchez, J., & Weise, C. (2022). The establishment and
maintenance of dominance hierarchies. *Philosophical Transactions of the Royal*
Society B: Biological Sciences, 377(1845), 20200450.
https://doi.org/10.1098/rstb.2020.0450

van Dijk, R. E., Covas, R., Doutrelant, C., Spottiswoode, C. N., & Hatchwell, B. J. (2015). Fine-scale genetic structure reflects sex-specific dispersal strategies in a population of sociable weavers (Philetairus socius). *Molecular Ecology, 24*(16), 4296–4311. https://doi.org/10.1111/mec.13308

Watt, D. J. (1986). Relationship of plumage variability, size and sex to social dominance in Harris' sparrows. *Animal Behaviour*, *34*, 16–27. https://doi.org/10.1016/0003-3472(86)90002-3

Weinstein, B. G. (2018). A computer vision for animal ecology. *Journal of Animal Ecology, 87*(3), 533–545. https://doi.org/10.1111/1365-2656.12780

Whiting, M. J., Nagy, K. A., & Bateman, P. W. (2003). Evolution and maintenance of social status-signaling badges. *Lizard Social Behavior*, 47–82. https://researchers.mq.edu.au/en/publications/evolution-and-maintenance-of-social-status-signalling-badges-expe

Wickham, M. H. (2014). Package 'ggplot2.' In *Create elegant data visualisations using the grammar of graphics. Version, 2(1)* (pp. 1–189). https://ggplot2.tidyverse.org/

Wiebe, K. L., & Vitousek, M. N. (2015). Melanin plumage ornaments in both sexes of Northern Flicker are associated with body condition and predict reproductive output independent of age. *Auk, 132*(2), 507–517. https://doi.org/10.1642/AUK-14-281.1

1   <u>**Supplementary materials:**</u> **Long-term evaluation of the association**
2   **between dominance, bib size, sex and age**
3

Authors: Nicolas J. Silva[a], Liliana R. Silva[b,c], Pietro B. D'Amelio[a,d], André C.

Ferreira[a,b,c,e], Annick Lucas[a], Rita Covas[b,c,f], Claire Doutrelant[a,f]

Corresponding author: Nicolas J. Silva, silva.nicolas.j@gmail.com

Affiliations:

[a]CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France;

[b]CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos,

Campus de Vairão, Universidade do Porto, 4485-66, Vairão, Portugal;

[c]BIOPOLIS Program in genomics, Biodiversity and Land Planning, CIBIO, Campus

de Vairão, 4485-661, Vairão, Portugal;

[d]Université Paris-Saclay, CNRS, Institut des Neurosciences Paris-Saclay, 91400,

Saclay, France;

[e]Department of Evolutionary Biology and Environmental Studies, University of

Zurich, Zurich, Switzerland;

[f]FitzPatrick Institute of African Ornithology, DST-NRF Centre of Excellence,

University of Cape Town, Rondebosch 7701, South Africa

Acknowledgments

We thank all involved in the long-term data collection and pre-processing over
the years, specifically for help with the yearly captures, dominance videos and bib

size picture collection. We thank Franck Theron for his management of the sociable weaver long-term database. We thank Margaux Rat, Inês Duarte and Bruna Fonseca for help scoring dominance videos. Additionally, we thank Margaux Rat for the initial development of the dominance and bib size collection and annotation protocols and Samuel Perret for bib size pictures and scoring. De Beers Mining Corporation provided access to Benfontein Nature Reserve and logistical assistance. The sexing and genotyping analyses were conducted at the CTM lab, CIBIO, and we thank Susana Lopes and all the technical staff that conduct the work. This study was funded by ANR to C.D. (France, grants 15-CE32-0012-02 and 19-CE02-0014-01), FCT (Portugal grants IF/01411/2014/CP1256/CT0007 and PTDC/BIA-EVF/5249/2014) ERC to R.C. (EU, Consolidator grant 866489) and by the DST-NRF Centre of Excellence at the Fitzpatrick Institute of African Ornithology University of Cape Town. N.J.S. was supported by the ANR 19-CE02-0014-01. L.R.S. was funded by ERC 866489 and ANR 19-CE02-0014-01. P.B.D.A. was supported by the south African Claude Leon Postdoctoral Fellowship 2019-20 and by the Marie Curie Individual Fellowship "SOCIAL MATCH" N. 896475 and then by the ANR 19CE02-0014-01. A.C.F was funded by University of Zurich Forschungskredit postdoc grant (K-74312-01-01 University of Zurich), Swiss Federal Commission for Scholarships and by ERC grants 866489 to R.C and. 850859 awarded to Damien Farine). R.C. was funded by FCT (IF/01411/2014/CP1256/CT0007 and CEECIND/03451/2018). The sociable weaver project is supported by the French OSU OREME, European Union's Horizon 2020 Research and Innovation Programme under the Grant Agreement Number 857251, and the CNRS programme on Long-Term Studies in Ecology and Evolution (SEE-Life).

Data availability statement

Analyses reported in this article can be reproduced using the data provided by Silva et al. (2025).

Silva, Nicolas J.; Silva, Liliana R.; D'Amelio, Pietro B., Ferreira, André C., Covas, Rita, & Doutrelant, Claire (2025). Data from: Long-term evaluation of the association between dominance, bib size, sex and age. Behavioral Ecology. https://doi.org/10.5061/dryad.v15dv4290

This supplementary materials include:

(i) Distribution of the normalised dominance score in the dataset (Fig. S1)

(ii) The description and evaluation of the deep learning method used for automatic bib

segmentation on pictures of the birds (Table S1).

(iii) The evaluation of the relationship between bib size, age, and sex. Two models are

presented: one with all the individuals; and one with a subsample due to non-

validation of the mixed model assumptions in the first model (Table S2).

(iv) The Directed Acyclic Graph of the model testing the association of dominance

with bib size and age (Fig. S2).

(v) The association between bib size and tarsus length (Fig. S3).

(vi) Justification of the use of Gaussian distribution models to test for the association

between dominance and bib size and age (Fig. S4).

(vii) The summary of the mixed models testing the overall association of dominance

with bib size and age in relation to sex. The models were computed using both-sexes

dominance hierarchies (Table S3 and S4), between-males dominance hierarchies

(Table S5 and S6) and between females dominance hierarchies (Table S7). For each

dominance hierarchies, the models were computed both with and without the

recommended threshold of ratio of interactions to number of individuals in the group

superior to 10 (Sánchez-Tójar et al., 2018) (Table S4 and S6), except for female

dominance hierarchies for which not enough colonies matched the criterion.

(viii) The summary of the mixed model testing the between-years association of

dominance with bib size and age in relation to sex. The models were computed using

both-sexes dominance hierarchies (Table S8 and S9), between-males dominance

hierarchies (Table S10 and S11) and between females dominance hierarchies (Table

S12). For each dominance hierarchies, the models were computed both with and

without the recommended threshold of ratio of interactions to number of individuals

in the group superior to 10 (Sánchez-Tójar et al., 2018) (Table S9 and S11), except for

female dominance hierarchies for which not enough colonies matched the criterion.

We also tested, using the package "emmeans" (Lenth et al., 2023), the slopes of the

relationship between dominance with bib size and the relationship between

dominance and age for each years (Table S13; Figure S5).

(ix) The influence of year 2013 and 2016 on the relationship between dominance and

bib size in between-females dominance hierarchies (Table S14).

(x) Timing of data collection for bib size and dominance and proportion of individuals

per colony per year observed during dominance video collection (Table S15).

Supplementary 1: Distribution of the normalised dominance score in the dataset.

Figure S1: Distribution of the dominance score, normalised by colony by year
between zero and one for the 723 individuals present in the dataset. Zero represent the
least dominant birds and one the more dominant birds. In the dataset, 13 individuals
had a score of exactly zero and 27 had a score of exactly one.

Supplementary 2: Evaluation of the automatic YOLO method and comparison with previous manual annotations.

YOLO is a deep learning model initially specialised in object detection (Redmon et al., 2016), whereas YOLOv8 allows to perform image classification, object segmentation, and pose estimation (Jocher et al., 2022). We used YOLOv8l, a sub-model of YOLOv8, to perform bib segmentation on the birds, due to a good trade-off between number of parameters and processing speed (Jocher et al., 2022; Terven & Cordova-Esparza, 2023).

To train our fine-tuned YOLOv8l model, 1037 pictures were manually annotated using VGG Annotator version 1.0.6 (Dutta & Zisserman, 2019) and we transformed the annotations into the YOLO input format. We separated the dataset into 934 train pictures and 103 validation pictures. Since the pictures were highly standardised, we did not used a test dataset as the validation was enough in these conditions. The model was trained using batches of 14 pictures, an initial learning rate of 0.01 and a final learning rate of 0.001, and data augmentation (flip left/right with a probability of 0.5 on each picture). We used a different image size input from the original YOLOv8 model because we needed accurate segmentation on high quality pictures. Therefore, we defined an image size input of 960x960 pixels.

To evaluate the model we used an Average Precision (AP) metric. The AP computes the number of True Positives divided by the number of True Positives + False Negatives. However, evaluating a segmentation is more complex than a common classification as we need to determine when the segmentation is good enough. For that we use the Intersection over Union (IoU), which is the ratio of the intersection between the true location of the object and the predicted location divided by the union of both the true location and the predicted one. The higher this ratio, the

higher is the quality of the detection. An IoU threshold can be set up depending on

how accurate the model needs to be. Here, we set up the threshold to 0.9, as we need a

very precise overlap between true locations and predicted locations (threshold is

usually set to 0.5 when using object detection). At a IoU of 0.9, we obtained an AP of

0.98, which indicates that nearly 100% of the bibs in the 103 validation pictures were

at least segmented with 90% of their real location.

The model's segmentation with YOLO presented similar high repeatability

between consecutive pictures of a same individual with the image graphic editor

GIMP: YOLO, (mean±SE) 0.90±0.016; CI: [0.866, 0.93]; P<0.001; N=300 versus

GIMP manual observer: (mean±SE) 0.92±0.014; CI: [0.885, 0.941]; P<0.001; N=300.

The speed to process 30 pictures was also compared, and the deep learning model

showed much faster performance than the GIMP method: (mean±SD) 10.2±0.04

seconds to process 30 pictures using the YOLOv8l model (N=10 repetitions) versus

21min07s seconds for the GIMP method (N=1 repetition). Additionally, the accuracy

of detection between real bib sizes (VGG annotations), the model's segmentation, and

the previous method (GIMP measures) were compared. Paired T-tests and

correlations tests were performed on 82 pictures and the GIMP method seems to be

closer to the VGG annotations than the YOLO method, but the YOLO model was

slightly more correlated with the VGG annotations (Table S1).

These tests allow us to validate the method using deep learning object

segmentation for fine-scale measurement of phenotypic traits. In the dataset used for

the analysis, only six years out of seven have the bib size computed using the YOLO

method. The GIMP method was used to obtain the bib sizes of the season 2013-2014

due to a loss of the pictures this year and the impossibility to compute the pictures

again with the YOLO method. However, bib sizes obtained using the GIMP and the

YOLO methods are highly correlated (Table S1), and as bib sizes are normalized

within year by colony, results still be comparable.

We also evaluated the reliability of the bib sizes obtained using pictures. We

blindly selected 27 individuals for which we computed the repeatability between two

to three set of three pictures taken a few dozens of minutes apart. We used the average

bib size, extracted with the YOLO method, on the consecutive three pictures per

individuals and we compared the replicates, for which we obtained high repeatability

(0.838±0.06 (mean±SE); CI: [0.723, 0.93]; P<0.001, N=58).

Table S1: Paired T-test and correlation test between the VGG annotations, the GIMP
data processing and the YOLOv8l fine-tuned model.

Method 1	Method 2	Estimate	Confidence intervals	p-value	Correlation
GIMP	VGG	-3328	[-4064, -2592]	<0.001	0.96
GIMP	YOLO	-14477	[-15342, -13611]	<0.001	0.95
VGG	YOLO	-11149	[-11762, -10536]	<0.001	0.97

Supplementary 3: Evaluation of the link between bib size, age and tarsus length

Table S2: Estimates of the model parameters to test the influence of the size and the
age on the bib size without one Photographer that was responsible for variance
deviation. Bib size was taken once a year at the end of winter and some individuals
were collected multiples years. The model was computedon a reduced dataset without
one of the scorers (N=3,373 observations, 814 males and 930 females). The model did
meet the residuals normality assumption.   Here, 'Age' is either exact age for
individuals born on our studied colonies or minimum age for individuals first caught
as adults.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	1.52	0.02	63.43	<0.001
Age	0.04	0.00	9.01	<0.001
Sex - *Male*	0.02	0.01	2.51	0.012
Tarsus	0.02	0.00	4.83	<0.001
Age : **Sex** - *Male*	-0.00	0.01	-0.44	0.657

Random effect	Variance			
Individual ID - *Intercept* (N=1,744)	0.013			
Individual ID - *Slopes* (N=1,744)	0.001			
Colony (N=19)	0.001			
Year (N=9)	0.002			
Photographer (N=6)	0.002			
Camera (N=3)	0.000			
Residuals	0.010			

Table S3: Estimates of the model parameters to test the influence of the size and the
age on the bib size with a subset of individual with at least three measures per
individuals ((mean ±SD) 3.88±1.15 measures per individuals). The table is the
summary of the effects using 2,081 observations (313 males and 223 females). Here,
'Age' is either exact age for individuals born on our studied colonies or minimum age
for individuals first caught as adults.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	1.56	0.03	50.33	<0.001
Sex - *Male*	0.02	0.01	1.66	0.097
Age	0.04	0.01	5.22	<0.001
Tarsus	0.02	0.01	2.96	0.003
Age : **Sex** - *Male*	-0.00	0.01	-0.61	0.543

Random effect	Variance
Individual ID - *Intercept* (N=536)	0.012
Individual ID - *Slopes* (N=536)	0.001
Colony (N=19)	0.001
Year (N=10)	0.002
Photographer (N=7)	0.004
Camera (N=3)	0.000
Residuals	0.012

Supplementary 4: Directed Acyclic Graph of the model testing the effect of the

bib on dominance.

Figure S2: DAG of the model testing the factors impacting dominance. Arrows
represent the direction of the effect. In the model structure, interactions between age
and sex, and bib size and sex were added as fixed factors. Colony ID, individual
identity and year of the data collection were added as random factors.

Supplementary 5: Association between bib size and tarsus length

Figure S3: Association between bib size (cm²) and tarsus (mm). Dots correspond to
raw data and lines represent the estimated linear fit and its 95% confidence intervals
from the mixed model in Table 1.

Supplementary 6: Justification of the use of Gaussian distribution models to test

for the association between dominance and bib size and age.

For the model testing the association between dominance and the different factors, we used a

Linear Mixed Model which implies Gaussian distribution. Initially, we aimed at a Beta

distribution model, adapted to estimate distribution between 0 and 1, and because Beta

distribution excludes exact 0 and 1, we added and subtracted 10-10 to respectively 0 and 1.

However, the model did not fit well (see analyses of the residuals; Fig. S4). The distribution

of the dependent variable is not what determines the link function to be used in the model,

and it is only an indication of the model type that we could use but it is not what it determines

the link function chosen. It is true that the estimated distribution will not be bounded by 0 and

1 as our data is, but since these data point are relatively rare (13 points with a score of zero

and 27 with a score of one), lower and higher estimates will be rare. Analyses of the residuals

of the model (Fig. S4) show that the Linear Mixed Model using a Gaussian link fitted better

than the Beta distribution model and fitted particularly well with no statistical deviation from

normality, no statistical over-dispersion of the residuals and no abnormal presence of outliers.

Figure S4: QQ-plot analyses of the residuals of the mixed model testing the association
between dominance with bib size and age using the both-sexes dominance hierarchies. The

model was initially tested using a Beta distribution model that did not fit well our data (see
QQ-plot on the right). The Gaussian distribution model fitted better and fitted well (see QQ-
plot on the left). The Gaussian distribution model was therefore kept for the analyses.

Supplementary 7: Evaluation of the effect of the ratio of dyads per individuals on

the overall association between dominance and bib size.

Using only the colonies with a ration of number of interactions to individuals

over 10, no major differences were observed in the values and directions of the

estimates of the significant effects in both-sexes dominance hierarchies and in

between-males dominance hierarchies (Table S4 to S6). No model was computed for

between-females dominance hierarchies as only one colony one year respected the

minimum ratio of interactions to individuals superior to 10. Both the models using

both-sexes dominance hierarchies and between-males dominance hierarchies

presented an improvement of the models' assumptions for validation with less

heteroscedasticity at the first and third quantile of the residuals when being computed

with the colonies with a ration of number of interactions to individuals over 10 (Table

S4 to S6). In these models, Age was either exact age for individuals born on our

studied colonies or minimum age for individuals first caught as adults.

Table S4: Estimates of the model testing the overall influence of bib size, sex and age
on dominance in both-sexes dominance hierarchies, without colonies with a ratio
inferior to 10 of interactions to the number of individuals. The model was computed
with 612 observations. The model's residuals followed normality, but
heteroscedasticity was observed near the residuals' quantiles 0.25 and 0.75.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.26	0.04	6.87	<0.001
Bib size	0.01	0.01	0.80	0.423
Sex - *Male*	0.30	0.02	19.05	<0.001
Age	0.00	0.01	0.38	0.702
Body condition	0.01	0.01	1.01	0.311
Tarsus	0.00	0.01	0.31	0.755
Bib size : **Sex** - *Male*	0.00	0.01	-0.04	0.968

Fixed effect	Estimate	SE	t-value	p-value
Age : **Sex** - *Male*	0.11	0.02	7.00	<0.001

Random effect	Variance			
Individual ID (N=377)	0.008			
Year (N=7)	0.004			
Colony (N=6)	0.004			
Residuals	0.020			

Table S5: Estimates of the model testing the overall influence of bib size, sex and age
on dominance in between-males dominance hierarchies, with colonies with a ratio
inferior to 10 of interactions to the number of individuals. The model was computed
with 419 observations. The model's residuals did not follow normality, and
heteroscedasticity was observed near the residuals' quantiles 0.25, 0.5 and 0.75.
Association of dominance with bib size and sex is visually represented in Fig. 2.E and
2.F.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.49	0.04	13.08	<0.001
Bib size	-0.02	0.01	-1.69	0.092
Age	0.16	0.01	11.77	<0.001
Body condition	0.03	0.02	1.53	0.128
Tarsus	0.01	0.02	0.40	0.690

Random effect	Variance			
Individual ID (N=207)	0.018			
Year (N=7)	0.001			
Colony (N=7)	0.006			
Residuals	0.034			

237 **Table S6:** Estimates of the model testing the overall influence of bib size, sex and age
238 on dominance in between-males dominance hierarchies, without colonies with a ratio
239 inferior to 10 of interactions to the number of individuals. The model was computed
240 with 188 observations. The model's residuals followed normality, but
241 heteroscedasticity was observed near the residuals' quantiles 0.25 and 0.75.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.45	0.03	13.67	<0.001
Bib size	-0.01	0.02	-0.47	0.640
Age	0.18	0.02	9.51	<0.001
Body condition	-0.01	0.03	-0.42	0.673
Tarsus	-0.05	0.03	-1.84	0.067

Random effect	Variance
Individual ID (N=130)	0.021
Year (N=7)	0.005
Colony (N=4)	0.000
Residuals	0.026

Table S7: Estimates of the model testing the overall influence of bib size, sex and age
on dominance in between-females dominance hierarchies, with colonies with a ratio
inferior to 10 of interactions to the number of individuals. The model was computed
with 249 observations. The model's residuals followed normality, and small
heteroscedasticity was observed near the residuals' quantiles 0.5. Association of
dominance with bib size and sex is visually represented in Fig. 2.C and 2.D.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.50	0.03	19.63	<0.001
Bib size	0.04	0.02	2.34	0.020
Age	-0.01	0.02	-0.35	0.725
Body condition	0.00	0.03	0.12	0.907
Tarsus	-0.01	0.03	-0.53	0.598

Random effect	Variance			
Individual ID (N=180)	0.023			
Year (N=7)	0.001			
Colony (N=6)	0.001			
Residuals	0.048			

Supplementary 8: Evaluation of the effect of the ratio of dyads per individuals on the between years association between dominance and bib size.

In both-sexes dominance hierarchies, using only the colonies with a ratio of number of interactions to individuals over 10, the sign and estimates of the significant effects did not change (Table S8 and S9), but the interaction between bib size and sex became significant for one season (Table S9). However, (i) the significance of this interactions this year is still in the same direction than the reference year in the model, making the correlation between bib and sex for male being different than the relation between bib and sex for female this year; (ii) the model presented a non-normality of the residuals, potentially affecting the reliability of this additional significant factor. In between-males dominance hierarchies, the signs of the significant effects remained the same between the models using the full dataset and the one using the dataset without the colonies with a ratio of number of interactions to individuals over 10, but the estimates of one of the season levels differed (0.14±0.04 (EST±SE), P=0.002; Table S10; and 0.34±0.09 (EST±SE), P=<0.001; Table S11). No model was computed for the between-females interactions dataset only as only one colony one year respected the minimum ratio of interactions to individuals superior to 10. In these models, Age was either exact age for individuals born on our studied colonies or minimum age for individuals first caught as adults.

Table S8: Estimates of the model testing the between years influence of bib size, sex and age on dominance in both-sexes dominance hierarchies, with colonies with a ratio inferior to 10 of interactions to the number of individuals. The model was computed with 723 observations. The model's residuals followed normality, but heteroscedasticity was observed near the residuals' quantile 0.75. Association of dominance with bib size and sex is visually represented in Fig. 3.A and 3.B.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.23	0.05	4.98	<0.001
Bib size	0.02	0.04	0.60	0.551

Sex - *Male*	0.30	0.05	5.96	<0.001
Year - *2014*	0.05	0.05	0.83	0.409
Year - *2015*	0.06	0.05	1.18	0.238
Year - *2016*	-0.03	0.06	-0.42	0.676
Year - *2017*	0.02	0.05	0.51	0.612
Year - *2018*	-0.01	0.05	-0.16	0.876
Year - *2019*	0.04	0.05	0.83	0.409
Age	0.02	0.04	0.50	0.620
Body condition	0.02	0.01	1.72	0.086
Tarsus	0.01	0.01	1.02	0.309
Bib size : **Sex** - *Male*	-0.04	0.04	-0.97	0.334
Bib size : **Year** - *2014*	0.00	0.05	-0.10	0.921
Bib size : **Year** - *2015*	0.03	0.05	0.57	0.571
Bib size : **Year** - *2016*	0.01	0.05	0.27	0.788
Bib size : **Year** - *2017*	-0.03	0.04	-0.67	0.502
Bib size : **Year** - *2018*	-0.05	0.05	-1.06	0.288
Bib size : **Year** - *2019*	-0.05	0.05	-1.10	0.271
Sex - *Male* : **Year** - *2014*	0.05	0.06	0.74	0.458
Sex - *Male* : **Year** - *2015*	0.05	0.06	0.86	0.390
Sex - *Male* : **Year** - *2016*	-0.03	0.07	-0.43	0.665
Sex - *Male* : **Year** - *2017*	-0.04	0.05	-0.79	0.429
Sex - *Male* : **Year** - *2018*	0.04	0.06	0.65	0.514
Sex - *Male* : **Year** - *2019*	0.00	0.06	0.07	0.947
Sex - *Male* : **Age**	0.09	0.04	2.32	0.021
Age :**Year** - *2014*	-0.02	0.04	-0.35	0.728
Age :**Year** - *2015*	-0.04	0.04	-0.84	0.400
Age :**Year** - *2016*	0.03	0.06	0.52	0.605
Age :**Year** - *2017*	-0.01	0.04	-0.15	0.878
Age :**Year** - *2018*	-0.03	0.05	-0.60	0.547

Fixed effect	Estimate	SE	t-value	p-value
Age :**Year** - *2019*	0.00	0.04	0.00	0.997
Year - *2014* : **Sex** - *Male* : **Bib size**	0.00	0.06	-0.06	0.950
Year - *2015* : **Sex** - *Male* : **Bib size**	-0.01	0.06	-0.12	0.902
Year - *2016* : **Sex** - *Male* : **Bib size**	-0.05	0.06	-0.88	0.380
Year - *2017* : **Sex** - *Male* : **Bib size**	0.07	0.05	1.40	0.162
Year - *2018* : **Sex** - *Male* : **Bib size**	0.08	0.06	1.51	0.131
Year - *2019* : **Sex** - *Male* : **Bib size**	0.07	0.06	1.23	0.218
Year - *2014* : **Sex** - *Male* : **Age**	0.00	0.05	0.02	0.980
Year - *2015* : **Sex** - *Male* : **Age**	0.08	0.05	1.44	0.152
Year - *2016* : **Sex** - *Male* : **Age**	0.06	0.07	0.94	0.346
Year - *2017* : **Sex** - *Male* : **Age**	-0.02	0.05	-0.35	0.726
Year - *2018* : **Sex** - *Male* : **Age**	0.06	0.06	1.07	0.287
Year - *2019* : **Sex** - *Male* : **Age**	0.00	0.05	-0.09	0.928

Random effect	Variance			
Individual ID (N=409)	0.009			
Colony (N=7)	0.002			
Residuals	0.022			

276 **Table S9:** Estimates of the model testing the between years influence of bib size, sex
277 and age on dominance in both-sexes dominance hierarchies, without colonies with a
278 ratio inferior to 10 of interactions to the number of individuals. The model was
279 computed with 612 observations. The model's residuals did not follow normality, and
280 heteroscedasticity was observed near the residuals' quantiles 0.25 and 0.75.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.24	0.05	4.73	<0.001
Bib size	0.01	0.04	0.25	0.799
Sex - *Male*	0.30	0.05	6.14	<0.001
Year - *2014*	0.06	0.06	0.99	0.321
Year - *2015*	0.05	0.05	1.04	0.297
Year - *2016*	-0.04	0.07	-0.57	0.572
Year - *2017*	0.01	0.05	0.32	0.750
Year - *2018*	-0.08	0.05	-1.48	0.139

Year - *2019*	0.11	0.06	1.95	0.052
Age	0.02	0.03	0.68	0.495
Body condition	0.01	0.01	0.58	0.563
Tarsus	0.00	0.01	-0.02	0.982
Bib size : **Sex** - *Male*	-0.03	0.04	-0.64	0.521
Bib size : **Year** - *2014*	0.02	0.05	0.35	0.727
Bib size : **Year** - *2015*	0.04	0.05	0.78	0.434
Bib size : **Year** - *2016*	0.02	0.06	0.37	0.709
Bib size : **Year** - *2017*	-0.01	0.04	-0.21	0.833
Bib size : **Year** - *2018*	0.00	0.05	0.01	0.990
Bib size : **Year** - *2019*	-0.03	0.05	-0.59	0.556
Sex - *Male* : **Year** - *2014*	0.03	0.07	0.50	0.614
Sex - *Male* : **Year** - *2015*	0.05	0.06	0.87	0.384
Sex - *Male* : **Year** - *2016*	-0.03	0.08	-0.37	0.711
Sex - *Male* : **Year** - *2017*	-0.04	0.05	-0.83	0.408
Sex - *Male* : **Year** - *2018*	0.03	0.06	0.47	0.640
Sex - *Male* : **Year** - *2019*	-0.02	0.07	-0.30	0.765
Sex - *Male* : **Age**	0.08	0.04	2.14	0.033
Age : **Year** - *2014*	-0.02	0.04	-0.36	0.722
Age : **Year** - *2015*	-0.05	0.04	-1.07	0.285
Age : **Year** - *2016*	-0.05	0.07	-0.75	0.451
Age : **Year** - *2017*	-0.01	0.04	-0.13	0.897
Age : **Year** - *2018*	-0.01	0.05	-0.22	0.826
Age : **Year** - *2019*	-0.03	0.04	-0.74	0.462
Year - *2014* : **Sex** - *Male* : **Bib size**	0.02	0.06	0.26	0.799
Year - *2015* : **Sex** - *Male* : **Bib size**	-0.01	0.06	-0.24	0.808
Year - *2016* : **Sex** - *Male* : **Bib size**	-0.06	0.07	-0.9	0.371
Year - *2017* : **Sex** - *Male* : **Bib size**	0.06	0.05	1.09	0.277
Year - *2018* : **Sex** - *Male* : **Bib size**	0.05	0.06	0.80	0.424

Fixed effect	Estimate	SE	t-value	p-value
Year - *2019* : **Sex** - *Male* : **Bib size**	0.05	0.06	0.95	0.341
Year - *2014* : **Sex** - *Male* : **Age**	-0.02	0.05	-0.37	0.712
Year - *2015* : **Sex** - *Male* : **Age**	0.09	0.05	1.76	0.078
Year - *2016* : **Sex** - *Male* : **Age**	0.18	0.09	1.99	0.047
Year - *2017* : **Sex** - *Male* : **Age**	0.00	0.05	-0.10	0.918
Year - *2018* : **Sex** - *Male* : **Age**	0.06	0.06	0.97	0.334
Year - *2019* : **Sex** - *Male* : **Age**	0.03	0.06	0.50	0.615

Random effect	Variance
Individual ID (N=377)	0.008
Colony (N=6)	0.004
Residuals	0.020

Table S10: Estimates of the model testing the between years influence of bib size, sex
and age on dominance in between-males dominance hierarchies, with colonies with a
ratio inferior to 10 of interactions to the number of individuals. The model was
computed with 419 observations. The model's residuals followed normality, but
heteroscedasticity was observed near the residuals' quantiles 0.25, 0.5 and 0.75.
Association of dominance with bib size and sex is visually represented in Fig. 3.E and
3.F.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.42	0.05	8.89	<0.001
Bib size	-0.02	0.03	-0.64	0.519
Year - *2014*	0.14	0.04	3.24	0.001
Year - *2015*	0.12	0.04	2.73	0.007
Year - *2016*	0.06	0.05	1.32	0.188
Year - *2017*	0.03	0.04	0.80	0.427
Year - *2018*	0.08	0.05	1.61	0.108
Year - *2019*	0.07	0.05	1.43	0.154
Age	0.16	0.02	6.39	<0.001
Body condition	0.03	0.02	1.43	0.155
Tarsus	0.01	0.02	0.45	0.656
Bib size : **Year** - *2014*	-0.04	0.04	-1.02	0.310
Bib size : **Year** - *2015*	0.01	0.04	0.15	0.879

Bib size : **Year** - *2016*	-0.03	0.04	-0.7	0.487
Bib size : **Year** - *2017*	0.02	0.04	0.51	0.609
Bib size : **Year** - *2018*	0.02	0.04	0.39	0.695
Bib size : **Year** - *2019*	0.01	0.04	0.23	0.816
Age :**Year** - *2014*	-0.01	0.04	-0.31	0.760
Age :**Year** - *2015*	0.03	0.04	0.79	0.428
Age :**Year** - *2016*	0.05	0.05	0.98	0.326
Age :**Year** - *2017*	-0.02	0.03	-0.66	0.510
Age :**Year** - *2018*	0.00	0.04	-0.08	0.937
Age :**Year** - *2019*	-0.03	0.04	-0.62	0.538

Random effect	**Variance**			
Individual ID (N=207)	0.018			
Colony (N=7)	0.006			
Residuals	0.035			

292 **Table S11:** Estimates of the model testing the between years influence of bib size, sex
293 and age on dominance in between-males dominance hierarchies, without colonies
294 with a ratio inferior to 10 of interactions to the number of individuals. The model was
295 computed with 188 observations. The model's residuals followed normality, but
296 heteroscedasticity was observed near the residuals' quantiles 0.25, 0.5 and 0.75.

Fixed effect	**Estimate**	**SE**	**t-value**	**p-value**
Intercept	0.34	0.06	5.34	<0.001
Bib size	-0.03	0.06	-0.47	0.639
Year - *2014*	0.34	0.09	3.73	<0.001
Year - *2015*	0.14	0.07	2.15	0.033
Year - *2016*	0.10	0.07	1.34	0.181
Year - *2017*	0.12	0.07	1.68	0.094
Year - *2018*	0.04	0.08	0.47	0.635
Year - *2019*	0.10	0.08	1.25	0.212
Age	0.13	0.05	2.88	0.005
Body condition	-0.02	0.03	-0.59	0.558
Tarsus	-0.05	0.03	-1.76	0.080

Fixed effect	Estimate	SE	t-value	p-value
Bib size : Year - *2014*	-0.10	0.08	-1.22	0.225
Bib size : Year - *2015*	0.01	0.06	0.21	0.832
Bib size : Year - *2016*	-0.02	0.08	-0.29	0.771
Bib size : Year - *2017*	0.05	0.07	0.81	0.417
Bib size : Year - *2018*	0.09	0.07	1.27	0.205
Bib size : Year - *2019*	0.03	0.07	0.46	0.646
Age :Year - *2014*	0.09	0.08	1.13	0.261
Age :Year - *2015*	0.09	0.06	1.63	0.106
Age :Year - *2016*	0.06	0.09	0.70	0.486
Age :Year - *2017*	0.08	0.06	1.27	0.207
Age :Year - *2018*	0.11	0.06	1.97	0.051
Age :Year - *2019*	0.00	0.06	0.00	0.999

Random effect	Variance			
Individual ID (N=130)	0.024			
Colony (N=4)	0.000			
Residuals	0.022			

Table S12: Estimates of the model testing the between years influence of bib size, sex
and age on dominance in between-females dominance hierarchies, with colonies with
a ratio inferior to 10 of interactions to the number of individuals. The model was
computed with 249 observations. The model's residuals followed normality, but
heteroscedasticity was observed near the residuals' quantile 0.75. Association of
dominance with bib size and sex is visually represented in Fig. 3.C and 3.D.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.47	0.06	7.25	<0.001
Bib size	0.14	0.06	2.36	0.019
Year - *2014*	-0.05	0.09	-0.53	0.593
Year - *2015*	-0.01	0.07	-0.20	0.843
Year - *2016*	0.03	0.10	0.33	0.739
Year - *2017*	0.07	0.07	1.07	0.287
Year - *2018*	0.04	0.08	0.45	0.650
Year - *2019*	0.03	0.08	0.43	0.665
Age	0.00	0.05	-0.05	0.958

Body condition	-0.01	0.03	-0.27	0.787
Tarsus	-0.02	0.03	-0.83	0.405
Bib size : Year - *2014*	-0.07	0.08	-0.83	0.407
Bib size : Year - *2015*	-0.08	0.07	-1.08	0.280
Bib size : Year - *2016*	0.04	0.10	0.39	0.696
Bib size : Year - *2017*	-0.10	0.07	-1.50	0.135
Bib size : Year - *2018*	-0.13	0.08	-1.67	0.096
Bib size : Year - *2019*	-0.13	0.07	-1.89	0.060
Age :Year - *2014*	0.09	0.06	1.47	0.142
Age :Year - *2015*	-0.03	0.06	-0.48	0.631
Age :Year - *2016*	0.11	0.10	1.05	0.297
Age :Year - *2017*	-0.07	0.06	-1.21	0.228
Age :Year - *2018*	0.11	0.09	1.13	0.260
Age :Year - *2019*	0.00	0.06	-0.07	0.946

Random effect	**Variance**
Individual ID (N=180)	0.025
Colony (N=6)	0.002
Residuals	0.046

305 **Table S13:** Post-hoc tests analysing bib size's and age's slopes of the between-years
306 variation GLMM for both-sexes dominance hierarchies, between-males dominance
307 hierarchies, and between-females dominance hierarchies. Sample size corresponds to
308 the number of individuals present in the dataset for a specific year. The tests were
309 performed with the package "emmeans".

Both-sexes dominance hierarchies

Variable	Year	Sample size	Slope	SE	df	t-value	p-value
Bib size Females	2013	17	0.02	0.04	669	0.60	0.552
	2014	22	0.02	0.03	659	0.53	0.594
	2015	47	0.05	0.03	669	1.67	0.095
	2016	16	0.04	0.03	672	1.05	0.292
	2017	92	-0.01	0.02	667	-0.30	0.765
	2018	45	-0.03	0.03	675	-0.98	0.326
	2019	41	-0.03	0.03	634	-1.04	0.301
Bib size	2013	51	-0.02	0.02	625	-0.86	0.392

Variable	Year	Sample size	Slope	SE	df	t-value	p-value
Males	2014	49	-0.03	0.02	591	-1.25	0.211
	2015	63	0.00	0.02	557	-0.03	0.978
	2016	42	-0.06	0.03	486	-2.18	0.030
	2017	117	0.02	0.02	647	1.25	0.213
	2018	58	0.02	0.02	563	0.70	0.488
	2019	63	0.00	0.02	615	-0.11	0.915
Age Females	2013	17	0.02	0.04	656	0.50	0.621
	2014	22	0.00	0.03	673	0.09	0.932
	2015	47	-0.02	0.03	669	-0.73	0.469
	2016	16	0.05	0.04	676	1.09	0.276
	2017	92	0.01	0.02	664	0.53	0.600
	2018	45	-0.01	0.03	605	-0.35	0.728
	2019	41	0.02	0.03	640	0.66	0.512
Age Males	2013	51	0.11	0.02	674	6.20	<0.001
	2014	49	0.10	0.03	628	3.85	<0.001
	2015	63	0.15	0.03	596	5.59	<0.001
	2016	42	0.20	0.03	587	6.36	<0.001
	2017	117	0.09	0.02	633	4.50	<0.001
	2018	58	0.14	0.02	585	6.11	<0.001
	2019	63	0.10	0.03	605	4.02	<0.001

Between-males dominance hierarchies

Variable	Year	Sample size	Slope	SE	df	t-value	p-value
Bib size	2013	45	-0.02	0.03	377	-0.64	0.521
	2014	48	-0.06	0.03	384	-2.04	0.042
	2015	59	-0.01	0.03	345	-0.39	0.695
	2016	41	-0.05	0.03	309	-1.55	0.122
	2017	112	0.00	0.03	388	0.02	0.981
	2018	52	0.00	0.03	337	-0.09	0.925
	2019	62	-0.01	0.03	374	-0.39	0.700
Age	2013	45	0.16	0.03	390	6.37	<0.001
	2014	48	0.15	0.03	384	4.59	<0.001

	Year	Sample size	Slope	SE	df	t-value	p-value
	2015	59	0.19	0.04	368	5.47	<0.001
	2016	41	0.21	0.04	343	4.90	<0.001
	2017	112	0.14	0.03	387	5.41	<0.001
	2018	52	0.16	0.03	362	5.21	<0.001
	2019	62	0.13	0.03	372	4.00	<0.001

Between-females dominance hierarchies

Variable	Year	Sample size	Slope	SE	df	t-value	p-value
	2013	19	0.14	0.06	223	2.34	0.020
	2014	17	0.07	0.06	216	1.25	0.214
	2015	45	0.06	0.04	198	1.42	0.157
Bib size	2016	12	0.17	0.08	221	2.25	0.026
	2017	84	0.04	0.04	225	1.04	0.298
	2018	35	0.01	0.05	213	0.19	0.850
	2019	37	0.01	0.04	169	0.14	0.892
	2013	19	0.00	0.05	223	-0.05	0.958
	2014	17	0.09	0.05	220	1.83	0.069
	2015	45	-0.03	0.04	205	-0.78	0.436
Age	2016	12	0.11	0.09	223	1.16	0.249
	2017	84	-0.07	0.03	176	-2.15	0.033
	2018	35	0.10	0.08	156	1.28	0.201
	2019	37	-0.01	0.04	181	-0.17	0.868

Figure S5: Slopes of the association between: (A,C,E) dominance and normalized bib size; and (B,D,F) dominance and age. Results were obtained from the models using both-sexes dominance hierarchies (A,B), between-female dominance hierarchies (C,D) and between-male dominance hierarchies (E,F). Dots represent the estimated means of the relationships and bar are the confidence intervals.

Supplementary 9: Influence of year 2013 and 2016 on the relationship between dominance and bib size in between-females dominance hierarchies.

Table S14: Estimates of the model testing the overall influence of bib size, sex and age on dominance in between-females dominance hierarchies, with colonies with a ratio inferior to 10 of interactions to the number of individuals but without year 2013 and 2016. These two seasons were removed to test their influence on the relationship between dominance and bib size found in Table S7. The model was computed with 219 observations. The model's residuals followed normality, and small heteroscedasticity was observed near the residuals' quantiles 0.75.

Fixed effect	Estimate	SE	t-value	p-value
Intercept	0.50	0.03	17.22	<0.001
Bib size	0.02	0.02	1.06	0.290
Age	-0.01	0.02	-0.38	0.704
Body condition	0.01	0.03	0.46	0.646
Tarsus	-0.02	0.03	-0.67	0.503

Random effect	Variance			
Individual ID (N=163)	0.019			
Year (N=5)	0.001			
Colony (N=6)	0.001			
Residuals	0.047			

Supplementary 10: Timing of data collection for bib size and dominance and proportion of individuals per colony per year observed during dominance video collection.

Table S15: Timing of data collection for bib size and dominance and proportion of individuals per colony per year observed during dominance video collection. Date for bib size data collection are given as month.day. Dominance data collection timing are given as the number of days after bib size collection. The total number of individuals used to compute the proportion of the colony observed is the total number of birds capture at a colony a year during the day of bib size collection. Proportion higher to 1 can happen when more individuals are observed on the videos than caught before (due to dispersal for instance). Stars corresponds to colonies where dominance was sampled more than 40 days after bib size collection date. In total, (mean±SD) 0.85%±0.23% were observed on average per colony per year, and the dominance data were collected on average on a period of (mean±SD) 9±9 days.

Season	Colony	Bib size date	Days after bib size collection dominance was collected	Number of days recording	Proportion of the colony observed
2013	11	08.31	4 to 22	16	0.80
	20*	09.04	59 to 60 and 81 to 92	14	0.74
	21*	09.09	48 to 59	12	1.00
	27	09.10	23 to 35	10	0.82
	71	09.07	18 to 30	11	1.55
	81	09.13	5 and 27 to 40	13	0.43
2014	11	08.25	12 to 19	5	0.66
	20	08.28	15 to 18	4	0.76
	21	09.02	7 to 20	8	1.14
	27	08.30	18 to 21	4	0.76
	71	09.01	5 to 12	5	0.81
	81	08.23	15 to 34	10	0.63
2015	11	09.04	-7 to 5	9	1.16
	20	08.31	-1 to 10	11	0.90
	21	08.27	1 to 13	11	1.25
	27	08.29	0 to 12	10	1.07
	71	09.03	-2 to 7	9	0.98
2016	11	08.30	20 to 25	4	0.49
	20	08.25	15 to 24	4	0.67
	27	08.31	-1 to 20	7	0.78
	43	08.22	10 to 35	6	0.62

	11	08.25	13 to 19	5	1.00
	20	08.29	9	1	0.96
2017	27	08.30	9	1	0.94
	43	08.23	10 to 12	3	0.93
	71	08.26	12	1	0.93
	11*	09.04	102	1	0.54
	20*	08.28	108	1	0.43
2018	27*	08.23	108 to 115	4	0.78
	43*	08.26	105 to 106	2	0.73
	71*	09.05	96 to 103	4	0.86
	11	09.08	10 to 12	3	0.98
	20	08.31	18 to 20	3	0.96
2019	27	09.06	12 to 14	3	0.81
	43	09.02	16 to 22	3	0.70
	71	09.10	9 to 11	2	0.88

References

Dutta, A., & Zisserman, A. (2019). The VIA annotation software for images, audio and video. *MM 2019 - Proceedings of the 27th ACM International Conference on Multimedia*, 2276–2279. doi:10.1145/3343031.3350535

Lenth, R. L., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., & Singmann, H. (2023). Estimated Marginal Means, aka Least-Squares Means. 1–101. https://github.com/rvlenth/emmeans

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S., Girndt, A., Bókony, V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T., & Schroeder, J. (2018). Meta-analysis challenges a textbook example of status signalling and demonstrates publication bias. *ELife, 7*, 1–26. doi:10.7554/eLife.37385

Terven, J., & Cordova-Esparza, D. (2023). A Comprehensive Review of YOLO: From YOLOv1 and Beyond. *ArXiv Preprint ArXiv:2304.00501*, 1–34. http://arxiv.org/abs/2304.00501