Long-term evaluation of the association between dominance, bib size, sex and age

Authors: Nicolas J. Silva[a], Liliana R. Silva[b,c], Pietro B. D'Amelio[a,d], André C. Ferreira[a,b,c,e], Annick Lucas[a], Rita Covas[b,c,f], Claire Doutrelant[a,f]

Corresponding author: Nicolas J. Silva, silva.nicolas.j@gmail.com

Affiliations:

[a]CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France;

[b]CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Campus de Vairão, Universidade do Porto, 4485-66, Vairão, Portugal;

[c]BIOPOLIS Program in genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661, Vairão, Portugal;

[d]Université Paris-Saclay, CNRS, Institut des Neurosciences Paris-Saclay, 91400, Saclay, France;

[e]Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland;

[f]FitzPatrick Institute of African Ornithology, DST-NRF Centre of Excellence, University of Cape Town, Rondebosch 7701, South Africa

Acknowledgments

We thank all involved in the long-term data collection and pre-processing over the years, specifically for help with the yearly captures, dominance videos and bib size picture collection. We thank Franck Theron for his management of the sociable weaver long-term database. We thank Margaux Rat, Inês Duarte and Bruna Fonseca for help scoring dominance videos. Additionally, we thank Margaux Rat for the initial development of the dominance and bib size collection and annotation protocols and Samuel Perret for bib size pictures and scoring. De Beers Mining Corporation provided access to Benfontein Nature Reserve and logistical assistance. The sexing and genotyping analyses were conducted at the CTM lab, CIBIO, and we thank Susana Lopes and all the technical staff that conduct the work. This study was funded by ANR to C.D. (France, grants 15-CE32-0012-02 and 19-CE02-0014-01), FCT (Portugal grants IF/01411/2014/CP1256/CT0007 and PTDC/BIA-EVF/5249/2014) ERC to R.C. (EU, Consolidator grant 866489) and by the DST-NRF Centre of Excellence at the Fitzpatrick Institute of African Ornithology University of Cape Town. N.J.S. was supported by the ANR 19-CE02-0014-01. L.R.S. was funded by ERC 866489 and ANR 19-CE02-0014-01. P.B.D.A. was supported by the South African Claude Leon Postdoctoral Fellowship 2019-20 and by the Marie Curie Individual Fellowship "SOCIAL MATCH" N. 896475 and then by the ANR 19CE02-0014-01. A.C.F was funded by University of Zurich Forschungskredit postdoc grant (K-74312-01-01 University of Zurich), Swiss Federal Commission for Scholarships and by ERC grants 866489 to R.C and. 850859 awarded to Damien Farine). R.C. was funded by FCT (IF/01411/2014/CP1256/CT0007 and CEECIND/03451/2018). The sociable weaver project is supported by the French OSU OREME, European Union's Horizon 2020 Research and Innovation Programme under the Grant Agreement

Conflict of interest statement

48 The authors declare no conflict of interest.

Authors' contribution

The study was initiated by C.D. and R.C., and N.J.S., L.R.S., P.B.D.A. and A.C.F.

further contributed to its conceptualisation and execution. All authors contributed to

the general monitoring data. L.R.S. and A.C.F. collected and processed the dominance

score of the birds. N.J.S., L.R.S., A.L., R.C., and C.D. collected and processed bib

data. N.J.S. developed the deep learning model. N.J.S. and P.B.D.A. produced the

statistical analyses. N.J.S. led the writing of the first draft of the manuscript with

contribution from all co-authors, and all authors contributed critically to the drafts and

gave final approval for publication. P.B.D.A., R.C. and C.D. supervised N.J.S. for this

work. R.C. and C.D. administrated the research project and acquired the funding for

this work.

Data availability statement

63 Analyses reported in this article can be reproduced using the data provided here:

64 https://figshare.com/s/cafa846429419e5a31bf

Abstract

Individual-based long-term datasets are needed to investigate the relationships between drivers of labile individual traits. A case in point is dominance structures, which has been shown to vary greatly across years and populations. Using an eight-year dataset, encompassing more than 700 individuals, we studied the association between the size of a melanin-based plumage trait (a black patch under the beak or 'bib'), individual attributes (age and sex) and dominance. Our study model, the sociable weaver (*Philetairus socius*), is a small passerine, which lives year-round in colonies of varying size. Bib sizes were obtained from pictures and automatically pre-processed with an object segmentation deep learning network based on YOLO. We recorded 36,346 dominance-related interactions to estimate dominance hierarchies. We then investigated the effect of sex and age on bib size, and the overall and between-year association of bib size and age with dominance for each sex. We found that bib size increased with age within individuals of both sexes. Age was a strong and consistent predictor of dominance in males, but not in females. Nevertheless, dominance was not correlated with bib size in males and not in females. Finally, these relationships did not strongly vary between years. Our results suggest that males and females may use different factors to regulate dominance interactions. Furthermore, our longitudinal dataset allowed us to show that bib size is not a badge of status under our population conditions even though its growth encodes age information, highlighting the importance of considering multiple traits over time when studying morphological signalling of hierarchy positions.

Keywords: Badge of status; Dominance; Deep learning; Long-term studies; Object segmentation; *Philetairus socius*; Social species; Status signalling hypothesis

1. Introduction

Competition over resources (e.g., food, territories, mates) often lead to aggressive interactions between individuals. These agonistic events can be very costly, sometimes leading to severe injuries or death (Clutton-Brock et al., 1979; Neat et al., 1996; Riechert, 1998; Sneddon et al., 2002). To limit these costs, many species establish dominance hierarchies (Tibbetts et al., 2022). Several traits or individual attributes, most commonly sex or age, have been found to be linked to the position of individuals in the dominance hierarchy (Tibbetts et al., 2022). However, dominance relations are likely to vary according to individual and environmental conditions and in a recent review, Tibbetts et al. (2022) argued that long-term field research is needed to deduce the factors influencing the stability of dominance hierarchies. In parallel, accumulating evidence shows that phenotypic traits and their signalling function may vary depending on external forces (e.g., availability of food, mates), which vary over time and can ultimately influence competition intensity (Kvarnemo & Ahnesjo, 1996; Lucchesi et al., 2019). Long-term studies integrate the wide ecological and social variation that characterise the environments experienced by individuals (Clutton-Brock & Sheldon, 2010a, 2010b; Cockburn, 2014) and have provided several examples where limited temporal sampling could have resulted in inaccurate generalisation of association patterns (Clutton-Brock & Sheldon, 2010a; Cockburn, 2014; Cornwallis & Uller, 2010; Doutrelant et al., 2020).

Dominance signalling relies on intrinsic competitive abilities of individuals and can be displayed through 'badges of status'. These traits can be visual, olfactory or acoustic and correlate with the individuals' dominance status (Rohwer, 1975, 1982; Senar, 2006; Smith & Harper 2003; Whiting et al., 2003). They therefore allow to

resolve conflicts without direct confrontation, as individuals can assess the
competitive ability of others and decide how to interact with them. The signalling
function of badges of status has been demonstrated both by correlational and
experimental studies in a wide range of taxa (e.g., mammals (Burgener et al., 2009;
Clutton-Brock & Albon, 1978), birds (Senar, 2006), reptiles (Stapley & Whiting,
2006), and insects (Tibbetts & Lindsay, 2008)).

In birds, a meta-analysis using 47 studies encompassing 23 species found that
plumage was linked to dominance across species and plumage types (e.g., carotenoids,
melanic), suggesting that there is overall statistical support for the status signalling
hypothesis across birds (Santos et al., 2011); however, heterogeneity was high,
indicating low generality of that finding across studies. Regarding melanin-based
plumage patches in birds, the textbook example has been the bib size of male house
sparrows *Passer domesticus*, which was found to be correlated with the fighting
abilities of individuals (Nakagawa et al., 2007). However, this example was
challenged in a meta-analysis conducted on 19 studies (Sánchez-Tójar, Nakagawa et
al., 2018), which showed that when including unpublished studies, the overall effect
size of the correlation between dominance and bib size was not different from zero,
and that the overall evidence for the hypothesis in house sparrows has been declining
over time. This meta-analysis revealed a strong publication bias that calls for
reconsideration of the bib size-based status signalling in this species, and for further
investigation of within-species variation in the relationships between badges of status
and dominance rank.

In addition to dominance signalling through phenotypic traits, dominance
hierarchies have also been shown to be associated with individual attributes, most

notably age. The hierarchies associated with individual attributes are called 'convention-based' hierarchies, and in these, age influences various phenotypic traits and regulates dominance interactions in many taxa (e.g., in birds (Sergio et al., 2009; Stahl et al., 2001), insects (Taylor et al., 2020), and mammals (Šárová et al., 2013)). For instance, in chimpanzees (*Pan troglodytes*), female social status is influenced by age, while males fight to gain status in the male hierarchy, which ultimately increases their reproductive success (Foerster et al., 2016).

The mechanisms regulating dominance may differ between the sexes because the benefits (and costs) of dominance are unlikely to be equivalent, and sex-biased dispersal may lead to stronger social bonds within one of the sexes. In birds, status traits have been studied mostly in males and less in females (Doutrelant et al., 2020), but some studies on mutually ornamented species found an association between dominance and a given ornament only in females (e.g., Chaine et al., 2011; Crowhurst et al., 2012; Jawor et al., 2004; Murphy et al., 2014; Watt, 1986). Hence, the selective pressures affecting these ornaments and their signalling function may be sex-specific. Finally, studies have also found evidence for a sex-dependent relationship between dominance and age (Foerster et al., 2016). However, overall sex differences in the mechanism regulating dominance remains to be explored in depth.

Long-term datasets are important because they allow to account for environmental variability. However, collecting such data can represent a challenge as it requires continuous funding to cover both logistical and human resources (Cockburn, 2014). Another challenge is the processing of the large volumes of data collected, which is very time consuming and costly when extracting information from videos or images (Weinstein, 2018). Recent approaches based on deep learning

methods (LeCun et al., 2015) can help with image processing, as they allow to

conduct time-consuming and repetitive tasks much faster than humans (e.g., few

milliseconds per images classification; Jocher et al., 2022) and sometimes

outperforming the repeatability and performance of humans (Ditria et al., 2020). In

ecology, deep learning methods are increasingly used for multiple tasks such as

identification, or density estimations, among others (Christin et al. 2019). However,

these methods are still rarely used for fine scale measurements of morphological traits,

as it requires accurate detection and precise segmentation of the phenotypic traits of

interest.

In the present study, we examine the relationship between a putative badge of

status (a black plumage 'bib') and dominance, age, and sex in a highly social colonial

and cooperative bird, the sociable weaver (*Philetairus socius*). To this end, we used a

deep learning approach, based on object-segmentation, to automatically extract the

bib size from approximately 10,000 pictures. We then calculated dominance from

more than 36,346 dominance interactions collected over seven years. In this species,

both male and female adults display a black melanin-based bib, with males only

presenting very slightly larger bibs (+0.032 cm²; Acker et al., 2015). Sociable weavers

establish steep dominance hierarchies within their colonies with males being

dominant over females (Rat et al., 2015; Silva et al., 2018) and older males have been

found to be dominant over younger males (Silva master thesis, 2017). Previous short-

term studies on this species found conflicting results regarding the association

between dominance and bib size, with either a positive association in male-male

interactions only (over two years; Rat et al., 2015), or females with larger bibs being

dominant when interacting with both sexes (Silva master thesis, 2017).

Here, we first evaluated how bib size varies in relation to sex and age, and performed a longitudinal analysis to understand the within individual variation in bib size. Then, we tested how dominance related with bib size, age and sex, and whether these associations varied between years. Since dominance hierarchies in this species are structured by sex (with most males being dominant over most females), we then explored whether the traits associated with dominance were different when estimating all individual hierarchies compared to when estimating same-sex dominance hierarchies. Long-term studies on badges of status are rare and we expected that by using such a large dataset spanning across time, we could reliably assess whether bib size functions as a badge of status, as well as capture potential variation over time of the relationship between dominance, age and bib size.

2. Materials and Methods

2.1. Species and individual information

Sociable weavers are passerines endemic to the semi-arid savannahs of the Kalahari region of southern Africa. They are small sexually monomorphic birds of approximately 28g and are colonial facultative cooperative breeders (Maclean, 1973). The study was conducted at Benfontein Nature Reserve, in South Africa (-28°818S, 24°816E, 1190m asl), an area of ~15km² of open savannah where up to 15 colonies are simultaneously monitored as part of a long-term study since 2010 (some colonies collapsed during the course of the study and new ones were added to the long-term monitoring).

The study was conducted between 2013 and 2021. Dominance trials were conducted on 5 of the regularly monitored colonies, while bib pictures were taken

from all colonies. We define a year as the period between 1st of September and the 31st of August, as the birds in the study colonies are captured at the end of the Southern Hemisphere winter, generally in August-September, and the breeding season usually starts in September-November (D'Amelio et al., 2024b). Sociable weavers can form very large colonies up to approximately 500 individuals (Maclean 1973), but colonies in our study area over the years of this study varied between 7 and 82 birds (mean±SD, 38±17 individuals). Individuals are ringed with three colored rings and a numbered metal ring, allowing individual visual identification. We also take morphological measurements (mass, tarsus+intertarsal joint's length) and blood samples to sex genetically all individuals (D'Amelio et al., 2024a; Griffiths et al., 1998). Exact age is known for all individuals that were born in the studied colonies as a result of a routine breeding monitoring (these individuals constitute 77% of the males and 65% of the females in the dataset). For immigrants, which were first captured as adults, we computed the age by adding the "average minimum age" of first dispersion to the first date of identification. This average minimum age of first dispersion in days was quantified for both males and females based on the long-term capture-recapture database (mean±SD, males: 690±46 days, N=13; females: 727±291 days, N=184). All protocols were approved by the Northern Cape Nature Conservation (permit FAUNA 1638/2015, 0825/2016, 0212/2017, 0684/2019 and 0059/2021) and the Ethics Committee of the University of Cape Town (permit 2014/V1/RC and 2018/V20/RC).

2.2. Dominance status

Using an established protocol (Rat et al., 2015; Silva et al., 2018), dominance hierarchies were determined within seven colonies each year (2013-2019) by

analysing and scoring agonistic interactions between individuals feeding on an artificial food source, a plate with mixture of bird seed, placed beneath each colony. Behavioural observations were performed using a video camera (Sony Handycam HD) on a tripod 2-3m from the feeder, to record all interactions seen approximately up to 1m around the feeder. From 2014 to 2017 and in 2019 observations were conducted in September; in 2013, from September to November; and in 2018, in December. Over these seven years, we conducted approximately 4h of observations each day, usually starting between 06:45 and 12:50; and repeated later in the day, between 13:15 and 16:50, and the food provided was removed between recordings. We obtained a total of 550 hours of video and each colony was recorded approximately (mean±SD) 15±12 hours per year (mean±SD, over approximately 6±4 days per colony). Both the type and direction of agonistic interactions observed were scored (as aggressions, displacements, threats and avoidances (Table 1 in Silva et al., 2018)). A total of 36,346 agonistic interactions (6,869 acts of aggression, 14,856 displacements, 3,459 threats and 11,162 avoidances) were recorded (mean±SD, approximately 1,010±1,240 interactions per colony per season).

To assess individual rank and to obtain dominance hierarchies, we used David's score (David, 1987), which is considered as one of the most appropriate and reliable methods available (Sánchez-Tójar, Schroeder et al., 2018), as it allows to assess overall individual success in agonistic interactions (Gammell et al., 2003). Additionally, this method allows to compare results to previous studies on dominance behaviour in this species. This score is calculated using the "steepness" R package (De Vries et al., 2006) and standardized per year and colony to allow comparison between individuals of different groups.

The hierarchies produced here included the majority of the birds captured in

winter for the studied colonies (85%±22% (mean±SD)). The 15% missing correspond

to birds never observed interacting at the feeders or that were not alive/present at the

colony at the recording time. Individuals seen in several colonies (N=29 individuals)

and those seen interacting less than three times in a given colony and year (N=45

individuals) were excluded from the analyses to avoid prospecting individuals (i.e.,

individuals briefly visiting colonies without being part of it).

In addition to these general dominance hierarchies (both-sexes dominance

hierarchies) including all individuals and interactions, we also extracted dominance

hierarchies by considering i) only male-male dominance interactions (N=16,261

interactions between 256 males, the between-males hierarchies) and ii) only female-

female dominance interactions (N=2,651 interactions between 267 females, the

between-females hierarchies).

To obtain reliable and representative dominance hierarchies for steep

dominance hierarchies, like the sociable weavers' (Silva et al., 2018), Sánchez-Tójar,

Schroeder et al. (2018), proposed that the threshold ratio of total number of observed

interactions to total number of individuals should have a minimum of 10. In our study

site, many colonies did not fulfil this criterium. We therefore decided to run our

models both with and without the colonies with a ratio inferior to 10 (except for

females as one colony was not enough to run the model with colonies with ratio

inferior to 10). We found that the estimates were similar between the models with and

without the colonies with a ratio inferior to 10 (Table S3 to S12), and therefore

decided to include the colonies with a ratio inferior to 10 in the results.

2.3. Bib size and YOLO: annotations, training bib segmentation and model validation

For all the birds captured, three top view pictures of the bib per individual were taken in the field (Rat et al., 2015). In previous works, bib size was manually analysed using image graphic editor (The GIMP Development Team, 2019 or Photoshop CS5.1 (Version 12.1), hereafter referred as "GIMP method"), which is a time-consuming task when dealing with large datasets. We improved this method by implementing a deep learning approach using YOLO (You Only Look Once; Redmon et al., 2016) to automatically detect and extract bib size from the pictures (Supplementary 1 for detailed description of the method).

We tested the repeatability of the GIMP and YOLO methods by selecting 20 individuals (300 pictures) on five different years for which bib size was measured with both methods (rptR package version 0.9.22 (Nakagawa & Schielzeth, 2010)). Results showed high repeatability and high correlations between the two methods (Table S1), indicating that the bib size measurements obtained with the deep learning model were robust. Bib size, extracted in pixels from pictures, was measured in cm².

2.4. Statistical analysis

All statistical analyses were conducted on R version 4.3.1, using mixed models ran with the lme4 package version 1.1-33 (Bates et al., 2014). We verified whether model assumptions were met, specifically that a) the residuals followed normality using Kolmogorov-Smirnov tests, b) the simulated dispersion of the residuals was equal to the observed dispersion, c) there were no more simulated outliers than expected, and d) there was homoscedasticity of the variance at three

quantiles (0.25, 0.5, 0.75) between the residuals and predicted values. These diagnostics were performed using the Dharma package version 0.4.6 (Hartig & Lohse, 2022), and, although not marked, some heteroscedasticity was observed and it is detailed in Supplementary 5 and 6. The plots were created with ggplot2 package version 3.4.2 (Wickham, 2014) and ggeffects version 1.3.2 (Lüdecke, 2023).

2.4.1. Factors associated with bib size

To design and interpret the most appropriate structure for the models testing the association between bib size and dominance, we first needed to understand the relationship of bib size with age within sexes (Fig S1). These variables were found to be associated in a previous study conducted on this species (using distinct datasets: Acker et al., 2015). In addition, we included tarsus length (as a proxy of body size), to correct for a potential correlative effect between bib size and body size. We designed a linear mixed model (LMM) with bib size (in cm²) as the response variable, and tarsus (in mm), age (in days), sex and the interaction between age and sex as fixed factors. As random factors, we controlled for year, colony and individual identity, the observer who took the photos (7 observers) and the camera model used (3 cameras). We also added random slopes to the random intercepts to estimate the link between age and bib size for every individual. We estimated random slopes uncorrelated with random intercepts, as we do not predict that individuals with larger bib size also have a higher bib size growth when getting older. We used a dataset composed of 743 individuals from 17 colonies over eight years (from 2014 to 2021, N=2,209 observations). Tarsus length was computed as the average tarsus length measured during the adult life of the individuals (by the same scorer, R.C., which measured the majority of birds, or by others if no measures by R.C. were found for that individual). Three individuals with extreme tarsus size observations were removed. The residuals

of the model using the 2,209 observations did not respect the normality assumption (Kolmogorov–Smirnov test, P<0.001). This non-normality of the residuals of the model were mainly due to one year photographed by one scorer. By removing this scorer from the dataset (N=647 observations), the model fitted the necessary assumptions. Estimates and effects of the different factors ranged from highly similar to identical between the two models (Table S2). Therefore, we decided to use the model containing the full dataset.

2.4.2. Overall association between dominance, sex, bib size and age

We performed a LMM using the dominance score of the individuals as the response variable and, as fixed factors, bib size, age, sex, tarsus, body index, as well as the interactions between bib size and sex, and age and sex. Body index was based on a scaled mass index (Peig & Green, 2009) that uses mass (here collected at the capture events, approximately when dominance data collection was performed) and a body size metric of an individual (here tarsus length, T) with its arithmetic mean value for the studied population (T_0). The method allows to compute a scaled mass as the predicted body mass when $T=T_0$. To visually describe the structure of our model, we designed a Directed Acyclic Graph (DAG) (Fig. S1). DAGs are visualisation tools that help to represent the links between the variables that we intend to test in a model (Wysocki et al., 2022). In this model, bib size was normalized within colony and year when analysing both-sexes dominance hierarchies, and within colony, year and sex when dealing with sex-specific dominance hierarchies. The normalisation allows us to compare relative bib size of the individuals within colonies and does not allow us to interpret bib size at the population level. Two individuals with a David score of one in different colonies might not have the same bib size but potentially be at the same bib size position in the bib size hierarchy within their colonies. Therefore, they would be

assigned the same normalized bib size. normalized bib size, body index, tarsus and

age were scaled and mean-centered over the whole dataset. Body index and tarsus

length were included because they are often related to dominance in other bird species

(Jonart et al., 2007; Tibbetts et al., 2022). As random factors, we controlled for colony

(N=7), individual identity (N=409) and year of data collection (N=7, from 2013 to

2019). We included year as a random factor since we expected potential differences in

dominance between years (e.g., arising from differences in social composition,

resource availability, breeding onset).

<u>2.4.3. Between-years association between dominance, sex, bib size and age</u>

We used the same LMM model structure as when testing the overall

association between dominance and bib size, but instead of having "year" as a random

categorical factor, we used it as a fixed categorical factor. Additionally, we included

the interactions between year, normalized bib size and sex, as well as between year,

age and sex. We then performed a post-hoc test with the emmeans package version

1.8.6 (Lenth et al., 2023) to determine if the slopes of the relationship between

dominance, bib size and age were significantly different from zero between years

(Table S13).

3. Results

3.1. Factors associated with bib size

We found that age and bib size were significantly positively associated

(0.03±0.01 (EST±SE, used throughout the Results section except if specified

otherwise), T=4.98, P<0.001; Table S2), and we estimated that between individuals

that differ by one year, the older birds have approximately 1% larger bibs. Within

individuals, we found a positive association between bib size and age for 99% of the

birds (Fig. 1), and this relationship was affected by sex, with bib size increasing by

0.97% per year within females and 1.98% per year within males. The variance

explained by random intercepts (Table S2, V=0.012) was higher than the variance

explained by random slopes (Table S2, V=0.001) suggesting that individuals vary in

their average bib size but have a similar pattern of growth through time. Moreover, we

found a small positive significant effect of sex (0.03±0.01, T=2.66, P=0.008; Table S2)

with males having on average 1.89% larger bib size than females (mean±SD,

1.56±0.03 cm² for females and 1.59±0.03 cm² for males). Finally, we found a

significant positive association between tarsus and bib size (0.02±0.00, T=3.41,

P=0.001; Fig. S2; Table S2), hence larger individuals have larger bibs independently

of sex.

Figure 1: Linear regression of bib size as a function of age. The random slopes and

intercepts of the 743 individuals (344 females (red) and 399 males (blue)) were used

to simulate bib size development every 200 days between one to 5,201 days old (i.e.,

more than 14 years).

3.2. Overall association between dominance, bib size, age and sex

<u>Dominance in relation to bib size and sex.</u> We found no significant association

between normalized bib size and dominance in the datasets considering both-sexes

dominance hierarchies (0±0.01, T=0.19, P=0.852; Table S3; Fig. 2.A) and between-

males dominance hierarchies (-0.02±0.01, T=-1.69, P=0.092; Table S5; Fig. 2.C).

However, we found a small positive significant correlation between bib size and

dominance in between-females dominance hierarchies (0.04±0.02, T=2.34, P=0.020;

Table S7; Fig. 2.E). Females being ranked 25% higher in the normalized bib size

ranks have an estimated 8% higher dominance score. However, the between-years

analysis bellow shows that the relationship is mainly influenced by two years with

very low sample size (Table S13) and when removing these two years the correlation

is weak and non-significant (0.02±0.02, T=1.06, P=0.290, Table S14).

<u>Dominance in relation to age and sex.</u> In both-sexes dominance hierarchies,

we found a strong significant link between sex and dominance, indicating that males

are approximately two times more dominant than females (0.30±0.02, T=18.61,

P<0.001; Table S3). We also found a strong association between dominance and the

interaction of age and sex (0.11±0.02, T=7.2, P<0.001; Table S3), indicating a

different relationship between age and dominance for males and females (Fig. 2.B),

specifically a much steeper slope for males. Dominance score is 1.36% higher in

females and 9.39% higher in males when comparing individuals that differ by one

year (Table S3). This was found also in the sex-specific interactions' datasets, where

we observed a different correlation between age and dominance, with a positive,

strong and significant association in between-males dominance hierarchies (0.16±0.01,

T=11.77, P<0.001; Table S5; Fig. 2.D), while there was no statistically significant

relation in between-females dominance hierarchies (-0.01±0.02, T=-0.35, P=0.725;

Table S7; Fig. 2.F).

Figure 2: Association between: (A,C,E) dominance and normalized bib size; and (B,D,F) dominance and age. Results were obtained using both-sexes dominance hierarchies (A,B), between-females dominance hierarchies (C,D) and between-males dominance hierarchies (E,F). Dominance score was standardized within colony and

year of data collection, and bib size has been normalized within colony and year of

data collection. Females are represented in red (square dots and dashed line) and

males are represented in blue (circle dots and solid line).

3.3. Between-years association between dominance, bib size, age and sex

Dominance in relation to bib size, sex and year. We found no evidence that the

relationship between dominance and normalized bib size varied between years, with a

few exceptions. Out of the seven years used in each analysis, the only significant

associations we found were a slightly negative relationship for one year in males in

both-sexes dominance hierarchies (-0.06±0.03, T=-2.18, P=0.030; Table S13; Fig. 3.A

in year 2016), a slightly negative relationship for one year in between-males

dominance hierarchies (-0.06±0.03, T=-2.04, P=0.042 in year 2014; Table S13; Fig.

3.E), and positive relationships for two years in between-females dominance

hierarchies (0.14±0.06, T=2.34, P=0.020 in year 2013, and 0.17±0.08, T=2.25,

P=0.026 in year 2016; Table S13; Fig. 3.C). However, the sample size was limited for

these latter years, with respectively 19 and 12 individuals only (the other years

presenting between 17 and 84 individuals, Table S13), and so these relationships

remain unclear.

Dominance in relation to age, sex and year. We found no between-years

variation in the association between dominance and age in both-sexes dominance

hierarchies and in between-males dominance hierarchies, with dominance being

positively associated with age in all years for males and never associated for females

(Table S13; Fig. 3.B and 3.F). In between-females dominance hierarchies, dominance

was negatively associated with age in one year only (-0.07±0.03, T=-2.15, P=0.033;

Table S13; Fig. 3.D in year 2017).

Figure 3: Slopes of the association between: (A,C,E) dominance and normalized bib

size; and (B,D,F) dominance and age. Results were obtained from the models using

both-sexes dominance hierarchies (A,B), between-female dominance hierarchies (C,D) and between-male dominance hierarchies (E,F). Dots represent the estimated means of the relationships and bar are the confidence intervals.

4. Discussion

We examined whether bib size and age relate to dominance at artificial feeders in sociable weavers and how these relationships vary between sexes and years. We found that, in both sexes, bib grows with age, and that larger birds have larger bibs. We also confirmed that males were dominant over females. However, normalized bib size was not found to be associated with dominance in any of the hierarchies (except in between-females dominance hierarchies for two years, but based on very low sample size). In addition, we confirmed that age was strongly associated with dominance in males a pattern that was consistent between years. In contrast, age was not found to be associated with female dominance. Our study is one of the largest long-term studies investigating correlates of dominance with a potential badge of status, providing robust results to understand the association between dominance and bib size, and the variation in its associated dominance correlates, given the low uncertainty in the variation obtained between years (see also Parker (2013); Sánchez-Tójar, Nakagawa et al. (2018); Tibbetts et al. (2022)). While previous studies found positive associations between dominance and bib size with only one year (included in this study (season /2014); Silva master thesis, 2017) and two years of data (not included in this study; Rat et al. 2015), our analyses, on a much larger dataset, led to very different conclusions. Finally, this study showed a strong sexual difference in the factor used to regulate dominance interactions

4.1. Association between dominance, bib size and sex

Relative bib size was clearly not associated with dominance, whether when

considering all birds or only males, and this did not change between years. We found

an association within females, but the association was weak and unclear, with small

effect sizes, and we had a smaller number of female-female interactions in the

hierarchies (32 out of 33 colonies were below the recommended threshold of ratio of

10 of interactions to number of individuals in the group; Sánchez-Tójar, Schroeder et

al. 2018). In addition, this association was only present in two years, which had

among the lowest sample sizes of all years tested here (respectively N=19 and N=12

females). When these two years are removed there is no longer evidence of an

association between relative bib size and dominance in females. If confirmed, these

results would add to the growing number of studies showing that badges of status

might function only in females but not in males (e.g., Chaine et al., 2011; Crowhurst

et al., 2012; Jawor et al., 2004; Murphy et al., 2014; Watt, 1986), but more studies on

both males and females are needed to better understand the possible functions of

badges of statues in both sexes (Doutrelant et al., 2020).

Several factors might explain why bib size does not function as a badge of

status in this population. First, the sampling period could influence our results, as the

association between bib size, age and dominance being potentially used by the birds

only in some parts of the season or some seasons only depending on the social and

environmental contexts. A density-dependent effect in the association between badge

of status and dominance has been found in the black-crested titmouse (*Baeolophus*

atricristatus), where males with larger black crests had more access to feeding

stations when competition was high but did not display such advantages when population density decreased and competition was low (Queller & Murphy, 2017). Here, the dominance scores were always computed at the end of winter, most of the time before the onset of the breeding season. Therefore, the association between dominance and bib size should be investigated at another moment of the year and over a longer time scale to confirm the results obtained here. Second, here we only tested the size of the bib, while other bib features could signal the dominance abilities of the individuals (e.g., Santos et al., 2011). For instance, in black-capped chickadees (*Poecile atricapilla*), high-ranking males exhibit darker plumages than subordinates (Poston et al., 2005). Finally, when facing known opponents, animals are expected to use information on the opponents' previous behaviour (i.e., their 'reputation') to decide how to interact (Senar, 2006; Tibbetts & Dale 2007). For instance, Chaine et al. (2018) experimentally manipulated plumage patch size of golden-crowned sparrows (*Zonotrichia atricapilla*) and showed that manipulation only affected interactions between unknown individuals. Sociable weaver groups live together all year long, interacting on a daily basis, and this increased familiarity between individuals, could remove the need badges of status. In addition, male sociable weavers are philopatric, therefore knowing each other since birth. A previous study showed that the number of aggressive interactions between two individuals was negatively correlated with relatedness (Rat et al., 2015). Therefore, the social system of sociable weavers might not promote the use of badges of status in males. However, badges of status might still be important for individuals that move between colonies, especially females, which disperse to reproduce in other colonies (Covas et al., 2006; van Dijk et al., 2015). Badges of status could also play a more important role among the residents of very large colonies where individuals might not be able to keep track of all residents.

Specifically, some sociable weaver colonies house several hundreds of birds (Maclean,

1973), while our study was conducted in colonies that varied in size between 7 and 82

residents.

4.2. Association between dominance, age and sex

We found a strong sexual difference in the association between age and

dominance. Evidence of age-based dominance hierarchies (also referred to as social

queuing) is common in many taxa and has particularly been found in species that form

long-term associations with relatives (Tibbetts et al., 2022), like sociable weavers. In

ungulates, for instance, individuals, that are young and more inexperienced learn to be

subordinates to older and stronger individuals, and do not challenge the hierarchy

when getting older, resulting in age-based hierarchies (Favre et al., 2008; Thouless &

Guiness, 1986). In sociable weavers, males know each other since birth and a

mechanism similar to the ungulates could explain their age-based hierarchy. In

contrast, females disperse to reproduce and therefore may not be able to use previous

information about the status of dominant or subordinate of their opponents. Females

may resort to alternative cues to infer fighting abilities of other females to decide how

to interact with them, or may not need to enter in competition with them. However, it

is important to note that because of this natal dispersal of the females, our estimation

of age for this sex is less accurate than for the males.

Social queuing has been suggested to also arise when the costs of reaching a

higher position in the hierarchy are higher than the benefits of reaching that position

(Broom et al., 2009). For instance, in chimpanzees, dominant males benefit from

higher mating and reproductive success, therefore males fight for higher ranks

(Foerster et al., 2016). However, females do not obtain fitness benefits from higher

dominance rank in the hierarchy and intense fights would only bring fitness costs,

resulting in age-dominance hierarchies in females (Foerster et al., 2016). In sociable

weavers, males of the same colony are more related to each other than females and,

within males' fights could either bring direct fitness costs (if the fight is lost), but also

indirect fitness costs (if the fight is won against a related individual). This could

suggest that, for males, the fitness costs of reaching a higher dominance rank might

not be compensated by the fitness benefits of reaching this rank, leading to age-based

dominance hierarchies within males. In male sociable weavers, dominant individuals

are more likely to access food and reproduce (Rat, 2015; Silva et al., 2018), but

further information of the exact fitness costs and benefits on the long-term for the

individuals are needed to explore the origin of the age-based dominance hierarchies in

males.

4.3. Association between bib size, age and sex

In agreement with a previous study (Acker et al. 2015), we found that bib size

increases with age in both sexes. Age-related changes in adult plumage have been

described in several species (e.g., Dreiss & Roulin, 2010; Wiebe & Vitousek, 2015).

However, the bib size growth between years is small, only 1% and 2% growth per

year for females and males respectively, and the perception by the individuals of such

difference is questionable. Interestingly, we found that the growth rate of the bib is

very similar between individuals, but individuals vary in bib size, indicating that the

mechanisms behind bib size first growth varied between individuals. This last result is

in line with the strong determination that genetics and development are expected to

have on bird melanic patches (Roulin & Ducrest, 2013). Finally, we detected an effect

of sex, with a difference of approximately +0.03cm² (approximately +2%) for the bib

size of males compared to females. This sexual difference is in line with the one

found previously in this species (Acker et al., 2015), but again, the perception of this

difference by the birds would need to be demonstrated.

4.4. The growing importance of artificial intelligence in ecology

Finally, our methodology for automatically measure thousands of bibs from

pictures adds to the growing examples that illustrate the effectiveness and accuracy of

deep learning-based methods to detect and quantify morphometrical traits, and how

this can facilitate the process of extracting large amounts of data from images. This

approach can be used in different taxa, from museum specimens (Miele et al., 2020),

to live ones (He et al., 2023), but it is still rarely used for such tasks in evolutionary

ecology. While in many cases developing deep learning methods would be too

demanding technically or too time consuming, data extraction from existing long-term

datasets, as was the case here, is particularly well suited to be processed automatically

(i.e., here the mode processed 124 times faster than a human's; based on 30 pictures).

5. Conclusion

In this study we found no convincing evidence that bib size functions as a

badge of status in sociable weavers, although for female-female interactions further

investigation is needed. These results align with the previous recommendation of re-

evaluating the status signalling hypothesis of melanic patches as well as evaluating

the temporal variation of the correlates with dominance. By using multiple years of

data, we were able to reach stronger conclusions regarding the stability of the

correlates (i.e., bib size and age) with dominance in our species. The sexual

differences in the association between dominance and age for the two sexes suggest

that male and female dominance interactions are potentially not regulated by the same

factors. These results show the importance of conducting analyses on both sexes when

trying to understand the role of phenotypic traits on dominance, and suggest that

males and females may be subjected to different selective pressures.

References

Acker, P., Grégoire, A., Rat, M., Spottiswoode, C. N., van Dijk, R. E., Paquet, M.,
Kaden, J. C., Pradel, R., Hatchwell, B. J., Covas, R., & Doutrelant, C. (2015).
Disruptive viability selection on a black plumage trait associated with dominance.
Journal of Evolutionary Biology, *28*(11), 2027–2041.
https://doi.org/10.1111/jeb.12717

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2014). Fitting linear mixed-
effects models using lme4. *Journal of Statistical Software, 67*(1), 1–48.
https://doi.org/10.18637/jss.v067.i01

Broom, M., Koenig, A., & Borries, C. (2009). Variation in dominance hierarchies
among group-living animals: Modeling stability and the likelihood of coalitions.
Behavioral Ecology, 20(4), 844–855. https://doi.org/10.1093/beheco/arp069

Burgener, N., Dehnhard, M., Hofer, H., & East, M. L. (2009). Does anal gland scent
signal identity in the spotted hyaena? *Animal Behaviour, 77*(3), 707–715.
https://doi.org/10.1016/j.anbehav.2008.11.022

Chaine, A. S., Shizuka, D., Block, T. A., Zhang, L., & Lyon, B. E. (2018).
Manipulating badges of status only fools strangers. *Ecology Letters, 21*(10),
1477–1485. https://doi.org/10.1111/ele.13128

Chaine, A. S., Tjernell, K. A., Shizuka, D., & Lyon, B. E. (2011). Sparrows use
multiple status signals in winter social flocks. *Animal Behaviour, 81*(2), 447–453.
https://doi.org/10.1016/j.anbehav.2010.11.016

Christin, S., Hervet, É., & Lecomte, N. (2019). Applications for deep learning in
ecology. *Methods in Ecology and Evolution, 10*(10), 1632–1644.
https://doi.org/10.1111/2041-210X.13256

Clutton-Brock, T. H., & Albon, S. D. (1978). The roaring of red deer and the evolution of honest advertisement. *Behaviour, 69*(3–4), 145–170. https://doi.org/10.1163/156853979X00449

Clutton-Brock, T. H., Albon, S. D., Gibson, R. M., & Guinness, F. E. (1979). The logical stag: adaptive aspects of fighting in red deer (Cervus elaphus L.). *Animal Behaviour, 27*, 211–225. https://doi.org/10.2307/3492065

Clutton-Brock, T. H., & Sheldon, B. C. (2010a). Individuals and populations: The role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends in Ecology and Evolution, 25*(10), 562–573. https://doi.org/10.1016/j.tree.2010.08.002

Clutton-Brock, T. H., & Sheldon, B. C. (2010b). The seven ages of pan. *Science, 327*(5970), 1207–1208. https://doi.org/10.1126/science.1187796

Cockburn, A. (2014). Behavioral ecology as big science: 25 years of asking the same questions. *Behavioral Ecology, 25*(6), 1283–1286. https://doi.org/10.1093/beheco/aru177

Cornwallis, C. K., & Uller, T. (2010). Towards an evolutionary ecology of sexual traits. *Trends in Ecology and Evolution, 25*(3), 145–152. https://doi.org/10.1016/j.tree.2009.09.008

Covas, R., Dalecky, A., Caizergues, A., & Doutrelant, C. (2006). Kin associations and direct vs indirect fitness benefits in colonial cooperatively breeding sociable weavers Philetairus socius. *Behavioral Ecology and Sociobiology, 60*, 323–331. https://doi.org/10.1007/s00265-006-0168-2

Crowhurst, C. J., Zanollo, V., Griggio, M., Robertson, J., & Kleindorfer, S. (2012). White flank spots signal feeding dominance in female diamond firetails, Stagonopleura guttata. *Ethology, 118*(1), 63–75. https://doi.org/10.1111/j.1439-0310.2011.01986.x

D'Amelio, P. B., Covas, R., Ferreira, A. C., Fortuna, R., Silva, L. R., Theron, F., Rybak, F., & Doutrelant, C. (2024a). Benefits of pair-bond duration on reproduction in a lifelong monogamous cooperative passerine. *The American Naturalist, 203*(5), 576–589. https://doi.org/10.1086/729436

D'Amelio, P. B., Silva, N. J., Silva, L. R., Ferreira, A. C., Theron, F., Doutrelant, C., & Covas, R. (2024b). Highly variable breeding phenology in an arid region colonial cooperative breeder, the Sociable Weaver Philetairus socius. *Ostrich, 95*, 1–7. https://doi.org/10.2989/00306525.2024.2328654

David, H. A. (1987). Ranking from unbalanced paired-comparison data. *Biometrika, 74*(2), 432–436. https://doi.org/10.1093/biomet/74.2.432

De Vries, H., Stevens, J. M. G., & Vervaecke, H. (2006). Measuring and testing the steepness of dominance hierarchies. *Animal Behaviour, 71*(3), 585–592. https://doi.org/10.1016/j.anbehav.2005.05.015

Ditria, E. M., Lopez-Marcano, S., Sievers, M., Jinks, E. L., Brown, C. J., & Connolly, R. M. (2020). Automating the analysis of fish abundance using object detection: optimizing animal ecology with deep learning. *Frontiers in Marine Science*, *7*, 1–9. https://doi.org/10.3389/fmars.2020.00429

Doutrelant, C., Fargevieille, A., & Grégoire, A. (2020). Evolution of female coloration: What have we learned from birds in general and blue tits in particular. *Advances in the Study of Behavior*, *52*, 123–202. https://doi.org/10.1016/bs.asb.2020.03.001

Dreiss, A. N., & Roulin, A. (2010). Age-related change in melanin-based coloration of Barn owls (Tyto alba): Females that become more female-like and males that become more male-like perform better. *Biological Journal of the Linnean Society*, *101*(3), 689–704. https://doi.org/10.1111/j.1095-8312.2010.01503.x

Favre, M., Martin, J. G. A., & Festa-Bianchet, M. (2008). Determinants and life-history consequences of social dominance in bighorn ewes. *Animal Behaviour*, *76*(4), 1373–1380. https://doi.org/10.1016/j.anbehav.2008.07.003

Foerster, S., Franz, M., Murray, C. M., Gilby, I. C., Feldblum, J. T., Walker, K. K., & Pusey, A. E. (2016). Chimpanzee females queue but males compete for social status. *Scientific Reports*, *6*, 1–11. https://doi.org/10.1038/srep35404

Gammell, M. P., De Vries, H., Jennings, D. J., Carlin, C. M., & Hayden, T. J. (2003). David's score: a more appropriate dominance ranking method than Clutton-Brock et al.'s index. *Animal Behaviour*, *66*(3), 601–605. https://doi.org/10.1006/anbe.2003.2226

Griffiths, R., Double C., M., Orr, K., & Dawson J. G., R. (1998). A DNA test to sex most birds. *Molecular Ecology*, *7*, 1071–1075. https://doi.org/10.1046/j.1365-294x.1998.00389.x

Hartig, F., & Lohse, L. (2022). *R package DHARMa. Version 0.4.6* (pp. 1–65). https://doi.org/10.32614/CRAN.package.DHARMa

He, Y., Cooney, C. R., Maddock, S., & Thomas, G. H. (2023). Using pose estimation to identify regions and points on natural history specimens. *PLoS Computational Biology*, *19*(2), 1–23. https://doi.org/10.1371/journal.pcbi.1010933

Jawor, J. M., Gray, N., Beall, S. M., & Breitwisch, R. (2004). Multiple ornaments correlate with aspects of condition and behaviour in female northern cardinals, Cardinalis cardinalis. *Animal Behaviour*, *67*(5), 875–882. https://doi.org/10.1016/j.anbehav.2003.05.015

Jocher, G., Chaurasia, A., Stoken, A., Borovec, J., Kwon, Y., Michael, K., Fang, J., Yifu, Z., Wong, C., & Montes, D. (2022). ultralytics/yolov5: v7.0 - YOLOv5 SOTA Realtime Instance Segmentation. In *Zenodo*. https://doi.org/10.5281/zenodo.7347926

Jonart, L. M., Hill, G. E., & Badyaev, A. V. (2007). Fighting ability and motivation:
determinants of dominance and contest strategies in females of a passerine bird.
Animal Behaviour, *74*(6), 1675–1681.
https://doi.org/10.1016/j.anbehav.2007.03.012

Kvarnemo, C., & Ahnesjö, I. (1996). The dynamics of operational sex ratios and
competition for mates. *Trends in Ecology and Evolution*, *11*(10), 404–408.
https://doi.org/10.1016/0169-5347(96)10056-2

LeCun, Y., Bengio, Y., & Hinton, G. (2015). Deep learning. *Nature, 521*(7553), 436-
444. https://doi.org/10.1038/nature14539

Lenth, R. L., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love,
J., Miguez, F., Riebl, H., & Singmann, H. (2023). *R package emmeans*.
https://doi.org/10.32614/CRAN.package.emmeans

Lucchesi, S., Cheng, L., Janmaat, K., Mundry, R., Pisor, A., & Surbeck, M. (2020).
Beyond the group: how food, mates, and group size influence intergroup
encounters in wild bonobos. *Behavioral Ecology*, *31*(2), 519–532.
https://doi.org/10.1093/BEHECO/ARZ214

Lüdecke, D. (2023). *R package ggeffects*.
https://doi.org/10.32614/CRAN.package.ggeffects

Maclean, G. L. (1973). The sociable weaver, part 1: description, distribution,
dispersion and populations. *Ostrich*, *44*(3–4), 176–190.
https://doi.org/10.1080/00306525.1973.9639158

Miele, V., Dussert, G., Cucchi, T., & Renaud, S. (2020). Deep learning for species
identification of modern and fossil rodent molars. *BioRxiv*, 2020.08.20.259176.
https://doi.org/10.1101/2020.08.20.259176

Murphy, T. G., West, J. A., Pham, T. T., Cevallos, L. M., Simpson, R. K., & Tarvin,
K. A. (2014). Same trait, different receiver response: Unlike females, male
American goldfinches do not signal status with bill colour. *Animal Behaviour, 93*,
121–127. https://doi.org/10.1016/j.anbehav.2014.04.034

Nakagawa, S., Ockendon, N., Gillespie, D. O., Hatchwell, B. J., & Burke, T. (2007).
Assessing the function of house sparrows' bib size using a flexible meta-analysis
method. *Behavioral Ecology*, *18*(5), 831-840.
https://doi.org/10.1093/beheco/arm050

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian
data: A practical guide for biologists. *Biological Reviews, 85*(4), 935–956.
https://doi.org/10.1111/j.1469-185X.2010.00141.x

Neat, F. C., Taylor, A. C., & Huntingford, F. A. (1998). Proximate costs of fighting in
male cichlid fish: The role of injuries and energy metabolism. *Animal Behaviour,
55*(4), 875–882. https://doi.org/10.1006/anbe.1997.0668

Parker, T. H. (2013). What do we really know about the signalling role of plumage colour in blue tits? A case study of impediments to progress in evolutionary biology. *Biological Reviews, 88*(3), 511–536.https://doi.org/10.1111/brv.12013

Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from mass/length data: The scaled mass index as an alternative method. *Oikos, 118*(12), 1883–1891. https://doi.org/10.1111/j.1600-0706.2009.17643.x

Poston, J. P., Hasselquist, D., Stewart, I. R. K., & Westneat, D. F. (2005). Dietary amino acids influence plumage traits and immune responses of male house sparrows, Passer domesticus, but not as expected. *Animal Behaviour, 70*(5), 1171–1181. https://doi.org/10.1016/j.anbehav.2005.02.015

Queller, P. S., & Murphy, T. G. (2017). Seasonal variation in the utility of a status signaling system: Plumage ornament predicts foraging success only during periods of high competition. *PLoS ONE, 12*(10), 1–14. https://doi.org/10.1371/journal.pone.0185584

Rat, M. (2015). *Dominance, social organisation and cooperation in the sociable weavers (Philetarius socius)* (Issue June 2015). http://hdl.handle.net/11427/16714

Rat, M., van Dijk, R. E., Covas, R., & Doutrelant, C. (2015). Dominance hierarchies and associated signalling in a cooperative passerine. *Behavioral Ecology and Sociobiology, 69*, 437–448. https://doi.org/10.1007/s00265-014-1856-y

Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You Only Look Once: Unified, Real-Time Object Detection. *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR),* 779–788. https://doi.org/10.48550/arXiv.1506.02640

Rohwer, S. (1975). The Social Significance of Avian Winter Plumage Variability. *Evolution, 29*(4), 593–610. https://doi.org/10.2307/2407071

Rohwer, S. (1982). The evolution of reliable and unreliable badges of fighting ability. *American Zoologist, 22*(3), 531–546. https://doi.org/10.1093/icb/22.3.531

Roulin, A., & Ducrest, A. L. (2013). Genetics of colouration in birds. *Seminars in Cell and Developmental Biology, 24*(6–7), 594–608. https://doi.org/10.1016/j.semcdb.2013.05.005

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S., Girndt, A., Bókony, V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T., & Schroeder, J. (2018). Meta-analysis challenges a textbook example of status signalling and demonstrates publication bias. *ELife, 7,* 1–26. https://doi.org/10.7554/eLife.37385

Sánchez-Tójar, A., Schroeder, J., & Farine, D. R. (2018). A practical guide for inferring reliable dominance hierarchies and estimating their uncertainty. *Journal of Animal Ecology, 87*(3), 594–608. https://doi.org/10.1111/1365-2656.12776

Santos, E. S. A., Scheck, D., & Nakagawa, S. (2011). Dominance and plumage traits:
Meta-analysis and metaregression analysis. *Animal Behaviour*, *82*(1), 3–19.
https://doi.org/10.1016/j.anbehav.2011.03.022

Šárová, R., Špinka, M., Stěhulová, I., Ceacero, F., Šimečková, M., & Kotrba, R.
(2013). Pay respect to the elders: Age, more than body mass, determines
dominance in female beef cattle. *Animal Behaviour*, *86*(6), 1315–1323.
https://doi.org/10.1016/j.anbehav.2013.10.002

Senar, J. C. (2006). Bird colors as intrasexual signals of aggression and dominance.
Bird Coloration, *2*, 125–193. https://doi.org/10.2307/j.ctv22jnr8k.6

Sergio, F., Blas, J., & Hiraldo, F. (2009). Predictors of floater status in a long-lived
bird: A cross-sectional and longitudinal test of hypotheses. *Journal of Animal
Ecology*, *78*(1), 109–118. https://doi.org/10.1111/j.1365-2656.2008.01484.x

Silk, J. B. (2019). Hyena politics: The dynamics of dynasties. *Proceedings of the
National Academy of Sciences of the United States of America*, *116*(18), 8644–
8645. https://doi.org/10.1073/pnas.1903407116

Silva, L. R. (2017). Assessing the physiological costs of dominance in a highly social
bird. *Master Thesis*. https://repositorio-
aberto.up.pt/bitstream/10216/90951/2/176196.pdf

Silva, L. R., Lardy, S., Ferreira, A. C., Rey, B., Doutrelant, C., & Covas, R. (2018).
Females pay the oxidative cost of dominance in a highly social bird. *Animal
Behaviour*, *144*, 135–146. https://doi.org/10.1016/j.anbehav.2018.08.006

Smith, J. M., & Harper, D. (2003). *Animal signals*. Oxford University Press.

Sneddon, L. U., Huntingford, F. A., & Taylor, A. C. (1998). Impact of an ecological
factor on the costs of resource acquisition: Fighting and metabolic physiology of
crabs. *Functional Ecology*, *12*(5), 808–815. https://doi.org/10.1046/j.1365-
2435.1998.00249.x

Stahl, J., Tolsma, P. H., Loonen, M. J. J. E., & Drent, R. H. (2001). Subordinates
explore but dominants profit: Resource competition in high arctic barnacle goose
flocks. *Animal Behaviour*, *61*(1), 257–264.
https://doi.org/10.1006/anbe.2000.1564

Stapley, J., & Whiting, M. J. (2006). Ultraviolet signals fighting ability in a lizard.
Biology Letters, *2*(2), 169–172. https://doi.org/10.1098/rsbl.2005.0419

Taylor, B. A., Cini, A., Cervo, R., Reuter, M., & Sumner, S. (2020). Queen
succession conflict in the paper wasp Polistes dominula is mitigated by age-
based convention. *Behavioral Ecology*, *31*(4), 992–1002.
https://doi.org/10.1093/beheco/araa045

Thouless, C. R., & Guinness, F. E. (1986). Conflict between red deer hinds: the
winner always wins. *Animal Behaviour, 34*(4), 1166–1171.
https://doi.org/10.1016/S0003-3472(86)80176-2

Tibbetts, E. A., & Dale, J. (2007). Individual recognition: it is good to be different.
Trends in Ecology and Evolution, 22(10), 529–537.
https://doi.org/10.1016/j.tree.2007.09.001

Tibbetts, E. A., & Lindsay, R. (2008). Visual signals of status and rival assessment in
Polistes dominulus paper wasps. *Biology Letters, 4*(3), 237–239.
https://doi.org/10.1098/rsbl.2008.0048

Tibbetts, E. A., Pardo-Sanchez, J., & Weise, C. (2022). The establishment and
maintenance of dominance hierarchies. *Philosophical Transactions of the Royal
Society B: Biological Sciences, 377*(1845), 20200450.
https://doi.org/10.1098/rstb.2020.0450

van Dijk, R. E., Covas, R., Doutrelant, C., Spottiswoode, C. N., & Hatchwell, B. J.
(2015). Fine-scale genetic structure reflects sex-specific dispersal strategies in a
population of sociable weavers (Philetairus socius). *Molecular Ecology, 24*(16),
4296–4311. https://doi.org/10.1111/mec.13308

Watt, D. J. (1986). Relationship of plumage variability, size and sex to social
dominance in Harris' sparrows. *Animal Behaviour, 34*, 16–27.
https://doi.org/10.1016/0003-3472(86)90002-3

Weinstein, B. G. (2018). A computer vision for animal ecology. *Journal of Animal
Ecology, 87*(3), 533–545. https://doi.org/10.1111/1365-2656.12780

Whiting, M. J., Nagy, K. A., & Bateman, P. W. (2003). Evolution and maintenance of
social status-signaling badges. *Lizard Social Behavior*, 47–82.
https://researchers.mq.edu.au/en/publications/evolution-and-maintenance-of-
social-status-signalling-badges-expe

Wickham, M. H. (2014). Package 'ggplot2.' In *Create elegant data visualisations
using the grammar of graphics. Version, 2(1)* (pp. 1–189).
https://ggplot2.tidyverse.org/

Wiebe, K. L., & Vitousek, M. N. (2015). Melanin plumage ornaments in both sexes of
Northern Flicker are associated with body condition and predict reproductive
output independent of age. *Auk, 132*(2), 507–517. https://doi.org/10.1642/AUK-
14-281.1