1
2
3 # Host-microbe mutualism drives urea carbon salvage and acetogenesis during hibernation

9 Matthew D. Regan*[1,2], Edna Chiang[3], Michael Grahn[1], Marco Tonelli[4], Fariba M. Assadi-Porter[5],
10 Garret Suen[3], Hannah V. Carey[1]
11
12

**Affiliations**

[1]Department of Comparative Biosciences, University of Wisconsin-Madison, Madison, WI, USA.

[2]Département de Sciences Biologiques, Université de Montréal, Montréal, QC, Canada.

[3]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA.

[4]National Magnetic Resonance Facility at Madison (NMRFAM), University of Wisconsin-
Madison, Madison, WI, USA.

[5]Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA.

*Corresponding author.

**Keywords**

**Abstract**

Hibernation is a wintertime survival strategy employed by certain mammals that, through torpor use, reduces overall energy expenditure and permits long-term fasting. Although fasting resolves the challenge of winter food scarcity, it also removes dietary carbon, a critical biomolecular building block. Here, we demonstrate a process of urea carbon salvage (UCS) in the 13-lined ground squirrel, whereby urea carbon is reclaimed through gut microbial ureolysis and used in reductive acetogenesis, producing acetate—a short-chain fatty acid (SCFA) of major value to the host and its microbial community. We find that urea carbon incorporation into acetate is more efficient during hibernation than the summer active season, and that while both host and gut microbes oxidize acetate for energy throughout the year, the host's ability to absorb and oxidize acetate is highest during hibernation. Metagenomic data of the gut microbiome indicate retention during hibernation of genes involved in the degradation of gut mucin, an abundant endogenous nutrient. The hydrogen disposal associated with reductive acetogenesis from urea carbon provides a luminal environment that sustains fermentation of mucin oligosaccharides, thereby generating SCFAs and other metabolites usable by both the host and its gut microbes. Our findings introduce UCS as a novel mechanism that enables hibernating squirrels and their gut microbes to exploit two key endogenous nutrient sources – urea and mucins – in the resource-limited hibernation season.

**Introduction**

Winter is a time of resource limitation for hibernating mammals and their gut microbes. This is especially true for species like the 13-lined ground squirrel (*Ictidomys tridecemlineatus*), which, thanks to their use of torpor, stop eating for approximately six months each year [1]. This effectively solves the animal's winter food scarcity problem, but it simultaneously deprives both the host and its gut microbes of exogenous nutrients. This can in turn jeopardize the dietary supply of valuable resources that are needed to sustain life throughout hibernation. However, mutualistic relationships exist between hosts and their microbial symbionts that allow the re-use of existing resources to meet this challenge.

One example of resource acquisition during fasting is urea nitrogen salvage (UNS), a gut microbe-dependent process that is used by ground squirrels to recycle urea nitrogen back into their tissue protein pools during hibernation [2], when they lack the typical dietary supply of nitrogen. This process, which peaks during late winter after months of fasting and just prior to the squirrel's emergence into the spring mating season, recycles nitrogen to support tissue protein synthesis throughout winter, and likely contributes to these hibernators' renowned

resistance to muscle atrophy [2,3]. Urea nitrogen is also incorporated into the microbial protein

pool, thus confirming that UNS benefits both symbiotic partners [2]. Thus, urea, which is

excreted as a waste product during summer, serves as an endogenous nitrogen source during

hibernation when the animals and their gut microbes lack an exogenous supply of this important

biomolecular building block.

In addition to nitrogen, urea contains carbon, another essential nutrient that is limited during

hibernation due to long-term fasting. Theoretically, the ability to capture this carbon via urea

carbon salvage (UCS) would also provide nutritional benefits to the microbiota and potentially to

the hibernator host. The existence of UCS has, to our knowledge, never been proposed in the

literature, perhaps because most animals do not experience extended periods with no carbon

intake where UCS could be beneficial. This is not the case for fasting hibernators, and

preliminary results from our previous study [2] suggested that urea carbon is indeed liberated by

microbial urea hydrolysis, biologically fixed, and incorporated into organic molecules. Thus, it is

possible that both host and microbes benefit from UCS during resource-limited hibernation.

In this study, we tested more rigorously the hypothesis that carbon liberated by urea

hydrolysis is incorporated into organic molecules, and that these molecules provide benefits for

ground squirrels and/or their microbiota during winter fasting. To accomplish this, we treated

squirrels with $^{13}C,^{15}N$-labeled urea and measured urea carbon (i.e., $^{13}C$) incorporation into the

microbiota, tissue metabolome and protein. We performed real-time in vivo $\delta^{13}C$ breath

experiments to estimate rates of microbial urea hydrolysis and absorption/metabolism of orally

gavaged acetate, as well as metagenomic analyses of bacterial genes involved in biological

carbon fixation and acetogenesis. In addition, we expanded the notion that fasted hibernators

and their gut microbes participate in mutualistic interactions that enable the use of host-secreted

compounds as nutrients by examining whether hibernator microbiomes are enriched in genes

coding for enzymes that degrade intestinal mucins, which are the most abundant endogenous

nutrients in the squirrel gut ecosystem during winter fasting.

**Materials & methods**

*Experimental animals*

From a total of 65 13-lined ground squirrels (39 females, 26 males; hereafter called squirrels),

39 were used for the doubly-labelled urea treatment experiments described in [2], 15 were

pretreated with oral gavages of $^{13}C$- acetate, and gut contents from 11 others were used for

mucin degradation metagenomics analyses. Squirrels were born in a vivarium on the University

of Wisconsin-Madison (UW-Madison) campus to wild-caught female squirrels that were

collected around Madison, Wisconsin in May 2018 and May 2019. The pregnant squirrels were

housed individually at 22°C with a 12:12 h light-dark cycle with ad libitum access to water and

rodent chow (Teklad no. 2020X, Envigo, Indianapolis, IN, USA) supplemented with fruit (apples)

and sunflower seeds once per week. After birth, pups remained with their mothers for 5 weeks

and then moved to individual cages. Following 2 weeks of ad libitum chow and fruit, food was

restricted to 12 g chow per day (supplemented with 1 g of sunflower seeds once per week) to

prevent excessive weight gain that occurs in captive-born squirrel pups. The consistent diet

minimized diet-induced effects on the gut microbiome and the molar ratio of $^{13}C$ to $^{12}C$ in

exhaled $CO_2$. Squirrels were held under these conditions until the day of experiment (Summer

group) or transferred to the cold room at the beginning of the hibernation season (Early and

Late winter groups).

In mid-September, squirrels in the Winter groups were moved to a 4°C environmental

chamber in the UW-Madison Biotron. The chamber was held in constant darkness except for ~5

min/day of dim light to enable activity checks. Food and water were removed after squirrels

began regular torpor-arousal cycles, which was 24–72 h after being moved into the chamber.

The 39 urea treatment squirrels were studied at in one of three time points: Summer (14

squirrels), Early Winter (12 squirrels), and Late Winter (13 squirrels). Each seasonal group was

divided into three treatment groups: $^{13}C,^{15}N$-urea with antibiotics, $^{13}C,^{15}N$-urea without

antibiotics, and unlabeled urea without antibiotics. Mean days in hibernation for the Early Winter

and Late Winter groups were 36 d (range 31–41 d) and 127 d (range 109–135 d), respectively.

Mean body masses on experiment days for Summer, Early Winter, and Late Winter groups

were 187.1 g (range 165.0–224.0 g), 155.6 g (range 123.4–187.2 g), and 134.1 g (range 105.5–

167.0 g), respectively.

The 15 acetate treatment squirrels were divided into four groups: Summer with antibiotics (4

squirrels), Summer without antibiotics (3 squirrels), Late Winter with antibiotics (3 squirrels), and

Late Winter without antibiotics (5 squirrels). Mean days in hibernation for the combined Late

Winter groups combined was 92 d (range 68–145 d). Mean body masses on experiment days

for Summer and Late Winter were 206.1 g (range 172.6–241.2 g) and 151.2 g (range 128.3–

174.0 g), respectively.

The 11 squirrels used for metagenomic analysis of mucin degradation genes were divided

into two groups: Summer without antibiotics (6 squirrels) and Late Winter without antibiotics (5

squirrels). Mean days in hibernation for the Late Winter group were 135 d (range 127–144 d).

Mean body masses on experiment days for Summer and Late Winter were 183.0 g (range

144.0–225.0 g) and 132.5 g (range 116.5–144.6 g)

The UW-Madison Institutional Animal Care and Use Committee approved all procedures.

*Animal treatments and stable isotope breath-testing*

For the 39 squirrels treated with doubly-labelled urea, the antibiotic treatments to deplete the gut

microbiota, the $^{13}C,^{15}N$-labeled urea treatments (200 mg urea kg$^{-1}$ body mass), the breath

testing experiments via cavity ring-down spectrometry (CRDS), and the euthanasia protocol

were all carried out as described in [2].

The 15 squirrels used to estimate host and microbe acetate oxidation capacity were treated

with $^{13}C$-labeled acetate (500 mg kg$^{-1}$ body mass) via oral gavage following light anesthesia (4%

isoflurane-$O_2$ mixture). Acetate oxidation capacity was measured using the CRDS-based breath

analysis, after which squirrels were euthanized. Antibiotics were administered as described in

[2].

The 11 squirrels used for metagenomic-based estimates of mucin glycan degradation were

administered saline via oral gavage following light anesthesia (4% isoflurane-$O_2$ mixture). On

the day of sampling, CRDS-based breath analysis was performed, after which they were

euthanized. There were no pretreatments for these squirrels.

In all cases, Early and Late Winter experiments were performed on euthermic squirrels

following forced arousal.

*Tissue harvesting and preparation*

147    Following euthanasia, we collected host tissues (liver, quadriceps) and cecal contents (a

148    proxy for the cecal microbiota) and prepared them for isotope ratio mass spectrometry (IRMS)

149    and nuclear magnetic resonance (NMR)-based metabolomics analyses as described in [2]. The

150    IRMS and NMR techniques were executed as described in [2]. All $^{1}H$ and $^{13}C$ spectra were

151    analyzed using the Chenomx software package (Version 8.5; Edmonton, AB, Canada).

*Metagenomics*

Generation of metagenomic data from cecal contents were performed as previously described

[2]. Briefly, genomic DNA was extracted using the Qiagen DNeasy PowerSoil Kit (Qiagen,

Hilden, Germany) according to manufacturer's specifications and quantified using a Qubit

Fluorometer (Invitrogen, San Diego, CA, USA). Sequencing was performed on an Illumina

NovaSeq6000 (Illumina, San Diego, CA, USA) using an S4 Reagent Kit v1.5 at 2x150bp PE.

Trimmomatic v0.38 [4] was used to remove sequencing adapters and low-quality reads, while

host DNA was filtered using bowtie2 v2.2.2 [5] against the 13-lined ground squirrel genome

(GenBank and RefSeq assembly accession = GCA_000236235.1). The remaining reads were

assembled using metaSPAdes v3.13.0 [6] and open reading frames (ORFs) were predicted

using prodigal v2.6.3 [7]

For the carbon fixation-related analyses, cecal content samples from three microbiota-

intact individuals were randomly chosen for Summer and Late Winter. Translated proteins from

the ORF predictions were compared against the Pfam [7] profiles for proteins involved in carbon

fixation and mucin degradation, as defined by KEGG using HMMER [8]. Identified sequences

were then mapped back to their contig of origin and taxonomically classified using centrifuge [9].

The percentage of target genes (*i.e.,* genes for carbon fixation or mucin degradation) in the

metagenomes were calculated by dividing the number of identified target genes by the total

number of predicted genes. The percentage of target genes classified to different taxa was

calculated by dividing the number of target genes classified to a specific taxon by the total

number of detected target genes.

For the mucin degradation analysis, we first identified putative enzymes thought to be

involved in mucin degradation from the Carbohydrate-Active enZYmes (CAZy) database [10],

including glycoside hydrolase (GH) 20 (exo-acting *β-N*-acetylglucosaminidases, *β-N*-

acetylgalactosamindase and *β*-6-SO$_3$-*N*-acetylglucosaminidases), GH33 (sialidase), GH84 ( *β-*

*N*-acetylglucosaminidases), GH85 (Endo-β-N-acetylglucosaminidases), and GH109 (α-*N*-

acetylgalactosaminidase). We then compared the translated proteins from the predicted ORFs

of the metagenomes against the CAZy database using dbCAN [11] and HMMER v3.3 [8].

To determine differences in mucin degradation genes between summer and late winter

metagenomes, we first normalized CAZyme mucin degradation genes by calculating reads per

kb per genome equivalent (RPKG) [12]. For each ORF, ORF length (kb) and the number of

mapped short reads were determined using bowtie2. Next, the number of mapped short reads

for each ORF was divided by ORF length and summed in each sample. This sum was then

divided by genome equivalents to calculate RPKG. RPKG seasonal differences were tested

using Kruskal-Wallis in R [13] (*kruskal.test*) and average genome size and genome equivalents

were tested using ANOVA (*aov*, stats package in R) to compare between summer and late

winter. Overrepresented CAZymes were identified with DESeq2 [14] and normalized counts

were transformed using the Hellinger method (*hellinger*, labdsv package in R).

*Statistical analyses*

Multivariate [13]C-metabolomic data for cecal contents, liver and muscle were analyzed by

combining principal components analysis (PCA) for visualization and PERMANOVA analysis for

quantitation (MetaboAnalyst 6.0, [www.metaboanalyst.ca](www.metaboanalyst.ca)). Effects of season and microbiota

presence on $^{13}$C-incorporation into identifiable metabolites (via NMR) and protein fraction of

tissue (via IRMS) were analyzed using 2-way ANOVA (GraphPad Prism v10.3). Breath $\delta^{13}$C

was plotted as a function of time following labeled urea or acetate treatment (2.5 to 3 h

depending on individual), and then area-under-the-curve (AUC) was calculated and expressed

relative to each individual's total breath analysis time. The data for urea-treated squirrels

originally reported in [2] were expressed here exponentially to eliminate negative values and

enable urea-to-acetate conversion efficiency calculations, which were expressed as the $^{13}$C-

acetate abundance relative to the exponentially converted breath $\delta^{13}$C AUC for the urea-treated

squirrels. Time course traces for breath responses to doubly-labelled urea-treated squirrels can

be found in [2]. Seasonal differences in breath $\delta^{13}$C AUC for urea-treated squirrels and relative

abundances of gut bacterial genes involved in biological carbon fixation (via Wood-Ljungdahl

reaction) and mucin degradation were analyzed using 1-way ANOVA with Tukey multiple

comparisons test (carbon fixation) and t-test (mucin degradation). Effects of season and

microbiota presence on breath $\delta^{13}$C AUC for acetate-treated squirrels were analyzed using 2-

way ANOVA. Non-normally distributed data via Shapiro-Wilk test were log-transformed prior to

analysis. In cases where data remained non-normal, non-parametric tests were used. For data

presented in whisker plots, the upper and lower bounds of the box represent 75$^{th}$ and 25$^{th}$

percentiles, respectively; the horizontal line within the box represents the median value, and the

upper and lower whiskers represent the maximum and minimum values within the data set,

respectively. Statistical significance was set at $P < 0.05$ for all comparisons.

**Results**

*Urea carbon ($^{13}$C) incorporation into host and microbe metabolome and protein*

We observed a total of 10, 12 and 15 metabolites enriched in urea-derived $^{13}$C in cecal

contents, liver and muscle, respectively (Figs. 1, 2).  The paired PCA-PERMANOVA multivariate

analysis revealed that, among the six treatment groups within each compartment, $^{13}$C-enriched

metabolite pools varied with season in cecal contents ($F=3.0992$, $R^2=0.4025$, $P=0.007$) and liver

($F=2.7784$, $R^2=0.3766$, $P=0.009$), but not in muscle ($F=1.2785$, $R^2=0.2175$, $P=0.288$) (Fig. 1).

However, among all observable metabolites, only cecal content $^{13}$C-acetate was significantly

affected by the presence of a microbiota, whereby its concentrations in microbiota-intact

squirrels were 2- to 7-fold higher than those of microbiota-depleted squirrels (Fig. 2A; ANOVA

$P < 0.0001$; original data were reported in Fig. S6 in [2] and expressed here in different units).

This microbiota effect confirmed that the incorporation of urea carbon into acetate molecules

was dependent upon microbial activity, most likely reflecting reductive acetogenesis that utilized

$^{13}CO_2$ generated by ureolytic bacteria after administration of $^{13}C,^{15}N$-urea. Microbial involvement

in $^{13}C$-acetate formation was not observed in liver despite $^{13}C$-acetate concentrations being

slightly higher in Late Winter than Summer or Early Winter (Fig. 2B; ANOVA P>0.05), or in

muscle where concentrations of $^{13}C$-acetate were below detectable limits (Fig. 2C).

In contrast to our previous observations [2] that urea-derived $^{15}N$ is incorporated into host

(liver, muscle) and microbiota (cecal contents) protein pools, we found no evidence of

microbiota-dependent incorporation of $^{13}C$ into protein fractions of the liver, muscle, or

microbiota after $^{13}C,^{15}N$-urea administration (Fig. S1). The small sample volumes of cecal

contents in the microbiota-intact squirrels required pooling all samples into a single aliquot for

IRMS analysis, which ultimately precluded statistical testing. Sample volumes for the

microbiota-depleted squirrels were also too small for analysis, likely because the protein fraction

of the cecal contentss was largely bacterial and thus absent from the microbiota-depleted

squirrels. However, we observed a trend for greater incorporation of $^{13}C$ into protein fractions of

Summer cecal contents compared with Early or Late Winter fractions (Fig. S1).

*Seasonal effect on ureolytic activity and urea-to-acetate efficiency*

The CRDS-based measurements of breath $\delta^{13}C$ reported in [2] for estimation of in vivo gut

microbial ureolytic activity were transformed here to express the $\delta^{13}C$ values as positive

numbers, thus enabling efficiency measurements (see Materials and Methods). This analysis

revealed that, as reported in [2], microbial ureolytic activity was higher in Summer than in Early

or Late Winter squirrels, which were not different (Fig. 3A; ANOVA P=0.0017).

Each squirrel's urea-to-acetate efficiency was expressed as the ratio of its cecal urea-

derived $^{13}C$-acetate concentration to its total $^{13}CO_2$ abundance, represented as its transformed

breath $\delta^{13}C$ value shown in Fig. 3A. This efficiency measure reflects the amount of liberated

urea carbon that is directed towards acetogenesis. We found that urea-to-acetate efficiency is

generally higher during hibernation than summer, with peak efficiency in Early Winter (Fig. 3B;

ANOVA P=0.0135; t-test P=0.0112 when grouping Early and Late Winter into a single

hibernation group).

*Seasonal effect on microbial biological carbon fixation capacity*

To estimate the capacity of the squirrel gut microbiome to biologically fix carbon and enable

acetogenesis, we analyzed the microbial metagenome for genes coding for enzymes involved in

the Wood-Ljungdahl (W-L) pathway of $CO_2$ fixation. The relative abundances of W-L-related

genes did not change from Summer to Late Winter (Fig. 4A; t-test P=0.2882) despite the ~90%

reduction in bacterial density by Late Winter [15]. Analysis of the top ten bacterial taxa from

taxonomic classification of W-L genes, as determined by sequence abundance, revealed a

seasonal shift from Summer to Late Winter in the bacterial potential for reductive acetogenesis

that could include use of urea carbon (Fig. 4B). Most notable was the finding that the greatest

percentage of W-L genes in Summer metagenomes (1.74%) belonged to the *Bacteroides*,

whereas over twice that amount (3.63%) was classified to *Blautia* in Late Winter. The

percentage of W-L gene sequences represented in *Blautia* spp. in Late Winter metagenomes

was over 5-fold than in Summer metagenomes.

*Capacities for acetate oxidation*

An increase in $\delta^{13}C$ breath after administration of $^{13}C,^{15}N$-urea indicates that oxidation of one or

more bioenergetic metabolites by the microbiota, by squirrel cells, or both is a consequence of

ureolysis in the gut lumen. Our identification of acetate molecules in the microbial metabolome

that contain urea-derived carbon [2] implicated acetate as one of those metabolites. To explore

the site of acetate oxidation in the squirrel holobiont, and whether there is a seasonal difference

in capacity for acetate oxidation by the microbiota and/or host, we orally gavaged microbiota-

intact and -depleted squirrels from Summer and Late Winter with a body mass-specific dose of

$^{13}C$-labeled acetate, and then measured their breath $\delta^{13}C$ using CRDS. None had been treated

with labeled urea. We found that squirrels from all groups exhibited rapid and large increases in

breath $\delta^{13}C$ (Fig. 5A), indicative of acetate oxidation and subsequent release of $^{13}CO_2$ in breath.

Squirrels in Late Winter generally exhibited greater breath $\delta^{13}C$ responses than squirrels in

Summer (Fig. 5B; ANOVA P=0.0003), but there were seasonal differences in how the presence

of the gut microbiota influenced the responses. In Summer, breath $\delta^{13}C$ as measured by AUC

was highest in squirrels with intact microbiotas, whereas in Late Winter, breath $\delta^{13}C$ was highest

in squirrels with depleted microbiotas (Fig. 5B). Consequently, microbiota-depleted squirrels

exhibited both the lowest (Summer) and highest (Late Winter) responses, with a >3-fold

difference between the two (Fig. 5B; ANOVA P=0.001). Season also influenced the time

required to reach peak breath $\delta^{13}C$, with Summer squirrels reaching peak values early (~30 min

post-gavage) relative to Late Winter squirrels (~75 min post-gavage) (Fig. 5A).

*Mucin degrading capacity in the squirrel microbiome*

Like urea, mucin glycoproteins are host-derived compounds that provide benefits to gut

microbes and their hosts. Mucins are secreted from goblet cells in the gut epithelial layer and

are either bound to epithelial cell membranes or secreted directly into the lumen. The glycan

component of these molecules are complex carbohydrates whose degradation releases a

variety of smaller oligosaccharides that are fermented anaerobically, producing energy-rich

metabolites including SCFAs that can be oxidized for ATP generation, used as signaling

molecules, or other purposes.

We initially planned to examine seasonal variation in recycling of mucin carbon from the

microbiota back into squirrel tissues using isotope labeling methods similar to those used for

urea [2], but were unable to do so due to a lack of commercially available $^{13}$C-labeled mucin

oligosaccharides that are resistant to metabolism by mammalian enzymes and could be

gavaged into squirrel guts. Therefore, we used metagenomics to test the hypothesis that the

seasonal shifts in gut microbiota composition we previously identified [16,17] is associated with

a change in the metagenome from Summer to Winter that favors retention of genes capable of

degrading mucins. We focused on gene families that encode CAZymes with the capacity to

degrade dietary polysaccharides, mucin glycans, or both, and identified 17 that were

overrepresented in either Summer or Winter metagenomes. Of the 17, we found five CAZymes

that were enriched in Winter vs. Summer, all of which are capable of degrading intestinal

mucins (Fig. 6).

**Discussion**

This study was designed to provide functional context for our observation that, in 13-lined

ground squirrels, carbon contained in urea molecules that enter the gut lumen is incorporated

into acetate within the microbial metabolome [2]. Here we show that this effect is restricted to

acetate among a suite of metabolites we examined and is more efficient during winter fasting

than the summer fed season. These and other findings discussed below support the idea that

through the symbiotic relationship between hibernators and their gut microbes, urea is

repurposed as a dual-use endogenous nutrient source during hibernation, with both its carbon

and nitrogen salvaged and used for synthesis of important biomolecules. Moreover, our findings

reveal a previously undescribed pathway for acetate production in the squirrel gut that uses $CO_2$

released by ureolytic microbes for hydrogen disposal via reductive acetogenesis. This helps to

maintain an efficient fermentative environment for the degradation of mucin glycoproteins, the

most abundant endogenous nutrients in the gut in fasted hibernators.

*Urea carbon salvage*

Like UNS, UCS begins with the movement of urea molecules from the blood into the gut lumen,

mediated by urea transporters (UT-B), where it is hydrolyzed by ureolytic gut microbes. We

recently showed that the abundance of UT-B in the squirrel cecum increases from Summer to

Late Winter, suggesting a greater capacity to transport urea into the lumen during the

hibernation season [2]. We also showed that ureolytic bacteria are found in the squirrel

microbiota and that rates of ureolysis are higher in Summer than in Early or Late Winter, likely

due to the reduction in total bacterial abundance that occurs in ground squirrels from summer to

winter [15,16].

Following urea hydrolysis, the next step in UCS involves urea carbon incorporation into

organic compounds (metabolites), a process carried out by gut microbes. These metabolites

would then be available for use by microbes and/or absorption and processing by the host. To

investigate this, we screened the metabolomes of the luminal cecal contents (a proxy for the

microbiota) and the squirrels' liver and skeletal muscle specifically for $^{13}$C-enriched metabolites

which, because our squirrels were treated with $^{13}$C,$^{15}$N-labeled urea, would indicate urea

carbon-containing metabolites. Looking broadly across the multivariate data via paired PCA and

PERMANOVA analyses, we found no significant effect of gut microbiota presence/absence on

the $^{13}$C-metabolomes of these three compartments (Fig. 1), suggesting that at this level of

analysis there is minimal incorporation of urea carbon into organic compounds. However, after

analyzing individual metabolites, we observed a significant gut microbe-dependent incorporation

of urea carbon into acetate (Fig. 2).

Our finding that urea carbon-derived acetogenesis occurs in the gut microbiota of

squirrels was notable for several reasons. First, although cecal acetogenesis by the reduction of

$CO_2$ with $H_2$ was first reported in 1977 [18] and replicated subsequently in the hindgut of many

monogastric species, the carbon source in those studies was not identified and was generally

assumed to be oligosaccharides produced by fermentation of dietary carbohydrates. Thus, our

study is the first to report incorporation of urea carbon in acetogenic reactions in the hindgut.

Also notable is that the cecal $^{13}$C-acetate concentrations are similar throughout the year

despite the decrease in bacterial abundance during hibernation [15,16]. This is coupled with the

sustained percentage from summer to winter of bacterial genes that code for enzymes in the

Wood-Ljungdahl pathway (Fig. 4). In this pathway, $CO_2$ is reduced by $H_2$ to form acetate and

water, thus linking the $CO_2$ produced by urea hydrolysis with acetogenesis. Taken together,

these findings suggest that the chronic lack of dietary substrates during hibernation may induce

a compensatory response in the squirrel microbiota that enhances metabolic functions and

survival in those bacteria that persist during winter fasting. This would include those genes that

code for carbon fixation and acetogenesis, facilitating the production of bioenergetic substrates

such as acetate. This is consistent with the stringent response, whereby bacteria under nutrient

stress such as carbon starvation reallocate resources from growth towards the upregulation of

survival-related genes until conditions improve [19–21].

Another example of molecular shifts activated by changing nutrients is the canonical

glucose-acetate transition in *Escherichia coli*, whereby cells excrete acetate while growing on

glucose and later absorb and catabolize it for energy when glucose is depleted. This is

facilitated by rapid gene expression modifications that suppress biosynthetic pathways and

upregulate catabolic ones [22,23]. Here, we posit that a similar situation may exist for the

microbiomes of fasting hibernators. When typical carbon sources in the gut such as glucose

derived from dietary polysaccharides are depleted over the winter fast, bacteria likely scavenge

for other sources [22]. Urea-derived acetogenesis via biological carbon fixation appears to be

one such pathway that persists throughout hibernation, possibly enabled by altered expression

of carbon fixation and acetogenesis genes in certain bacterial species, despite an overall

reduction of bacterial density due to winter fasting [15,16]. In the case of fasted hibernators, the

contributions of pathways that recycle endogenous host-derived compounds such as urea and

mucins may play significant roles in energy-sparing, thus conserving stored lipids that are the

major fuel source during torpor and arousals [24]. Although we detected no seasonal changes in

the proportion of genes for carbon fixation (Fig. 4A), transcriptomic or proteomic analyses might

reveal that ground squirrel microbiomes upregulate these enzymatic pathways from summer

feeding to winter fasting.

Interestingly, more of the urea carbon liberated by microbial ureolysis appears to be

used for acetogenesis during hibernation than during the summer active season. Specifically,

the ratio of $^{13}$C-acetate to total exhaled $^{13}CO_2$ was higher for hibernating squirrels than for

Summer squirrels (Fig. 3B). This suggests that less urea carbon is lost as $CO_2$ in breath during

winter, when exogenous carbon intake has ceased and when urea carbon, via carbon fixation

and subsequent acetogenesis, would most benefit the hibernator holobiont.

*Potential benefits of urea carbon-derived acetate for hibernator host and microbiota*

We perceive at least four benefits that UCS-mediated acetogenesis may offer hibernating

species during winter fasting. First, acetate can serve as a direct energy source for host cells,

with intestinal epithelial cells (IECs) being most likely to benefit given their dependence on

luminal SCFAs such as butyrate to meet energy needs [25]. Under normal fed conditions,

butyrate is the preferred energetic substrate for IEC; however, when butyrate availability is

limited, acetate oxidation can supply ATP [26]. A similar shift in the nutritional landscape of the

cecal lumen occurs during hibernation, where concentrations of all SCFAs, including butyrate

concentrations, fall to 3% of Summer levels. Importantly, acetate concentrations remain at 30-

40% [15,16] and it is therefore possible that UCS-derived acetogenesis contributes to this

energy supply for the host, particularly at the level of IECs.

To lend empirical support for acetate's role as an energy source for hibernators and their

gut microbes, we orally gavaged $^{13}C$-acetate into Summer and Winter squirrels. We then used

CRDS to measure breath $\delta^{13}C$, whereby elevated values above baseline indicate oxidation of

$^{13}C$-acetate to $^{13}CO_2$. Breath $\delta^{13}C$ rapidly increased following $^{13}C$-acetate gavage in microbiota-

intact Summer and Late Winter squirrels, resulting in similar AUC values for the 3 h post-gavage

data collection period (Fig. 5). To distinguish host acetate oxidation from potential microbial

acetate oxidation, we repeated the experiments on microbiota-depleted squirrels in both

seasons. In Summer squirrels, depleting the microbiota halved the breath $\delta^{13}C$ AUC (suggesting

a microbial contribution to acetate oxidation), whereas in Late Winter squirrels it had no effect

on breath $\delta^{13}C$ AUC but produced a higher peak in $\delta^{13}C$ that was more quickly attained. Overall,

the Late Winter microbiota-depleted squirrels had AUC values 3-fold higher than Summer

microbiota-depleted squirrels. These results suggest two things about acetate use during

hibernation. First, during summer, gut microbes oxidize much of the acetate they produce,

whereas during hibernation, a greater proportion of the acetate is absorbed and oxidized by the

host. Thus, the host appears to benefit most from the bioenergetic contributions of acetate

during hibernation. We suspect that the primary site of oxidation for microbially generated

acetate in squirrels, especially during hibernation, is within IECs. This is consistent with the lack

of observable $^{13}C$-acetate in liver and muscle, and with the generally deficient energy state of

IECs during long periods of food deprivation [27]. Second, the 3-fold higher breath $\delta^{13}C$

responses after $^{13}C$-acetate gavage in Late Winter compared with Summer microbiota-depleted

squirrels indicate that the capacity to absorb and oxidize acetate increases during hibernation,

when resources are limited and organic compounds such as acetate are especially valuable.

Although we have not directly measured rates of intestinal acetate absorption, these results are

consistent with our previous work showing higher normalized rates of $Na^+$, glucose, and amino

acid absorption by the small intestinal epithelium in hibernating than summer squirrels [28,29].

Similar results were observed for nutrient uptake into brush border membrane vesicles [30], and

it is plausible that similar seasonal effects on intestinal acetate absorption via monocarboxylate

transporters underlie the results seen here. A second, indirect way that urea carbon-derived

acetate may influence IEC metabolism during winter fasting is through its conversion to their

preferred fuel source by acetate-consuming, butyrate-producing bacteria within the microbiome
[31]. Together, these results suggest that the hibernator's physiology is seasonally optimized to
benefit from UCS-derived acetogenesis, which may suggest positive selection on this
mechanism.

A second potential role for acetate produced from urea carbon for the hibernating host is
its role as a ketogenic substrate. The ketone β-hydroxybutyrate is the preferred fuel source for
the heart and brain of ground squirrels during hibernation [32].  Ketogenesis, which takes place
primarily in the liver but can occur in other sites including the gut [33,34], is regulated by the
rate-limiting enzyme (3-hydroxy-3-methylglutaryl-CoA synthetase 2; HMGCS2). We showed
previously that HMGCS2 is expressed in 13-lined ground squirrel intestine and its protein
abundance is 2-fold higher during hibernation than summer [35]. This finding is consistent with
those from mice and rats whereby short-term fasting elevates both intestinal gene expression of
HMGCS2 [36] and the conversion of acetate to ketone bodies in isolated colonocytes [26].
Furthermore, a role for the microbiota in stimulating hepatic ketogenesis in mice under nutrient
deprivation was shown when mice with intact gut microbiotas elevated hepatic ketogenesis
following a 24 h fast, but the response was reduced in mice lacking a microbiota [37]. Given that
levels of fatty acids, which are the most common substrates for ketogenesis, are likely reduced
in hibernator IECs due to the lack of dietary lipid intake, acetate may be an effective ketogenic
precursor for the gut during hibernation.

A third potential way that UCS-derived acetate benefits the hibernating host is through
its effects on secretory IgA (sIgA), the most abundant immunoglobulin in mammals, and one
that provides the first line of defense at the epithelial surface. We showed previously that small
intestinal expression of sIgA is greater in torpor phases of hibernation compared with summer
levels, part of a suite of changes in the intestinal immune system that appears to strengthen
anti-microbial defenses and thereby homeostasis of the host with its gut microbiota [38]. More
recent studies have implicated microbial-derived acetate, but not butyrate, in playing an
essential role in regulation of sIgA production that helps shape commensal microbiota
composition and minimize microbial pathogen colonization [39,40]. Thus, maintenance of host-
microbiota mutualism during winter fasting may require alternative means to generate acetate
for immune protection such as UCS-derived acetogenesis.

Finally, gut microbial acetogenesis plays a key role in maintaining an environment
conducive to ongoing fermentation. The fermentative breakdown of complex carbohydrates
produces $H_2$ which, if not removed or metabolized, will eventually inhibit fermentative reactions
by inhibiting the regeneration of the coenzyme $NAD^+$ from NADH, thus slowing the

thermodynamic favourability of SCFA production and hindering bacterial growth [41].

Acetogenic bacteria help to keep $H_2$ partial pressure at an optimal level by reducing $CO_2$ with

$H_2$, producing acetate and $H_2O$. Methanogenesis plays a similar role in many animals (but

without formation of a bioenergetic compound like acetate), though there is no evidence for

hydrogenotrophic methanogens (Archaea) in the microbiotas of hibernating ground squirrels

[15–17] or bats [42]. Thus, acetogenesis is likely the primary $H_2$ disposal mechanism used in the

hindgut of mammalian hibernators. This is useful for the holobiont during hibernation because,

while there is a lack of exogenous complex carbohydrates during the winter fast, there is a

continuous supply of mucin glycoproteins in the intestinal lumen that, as described in the

following section, can serve as endogenous source of fermentable carbohydrates.

*Seasonal changes in mucin degradation capacity in the ground squirrel microbiota*

Mucins are complex glycoproteins that are secreted from goblet cells in the gut epithelium that,

upon hydration in the lumen, form mucus, a complex viscoelastic matrix that protects the

epithelium and serves as a habitat for a diverse assemblage of microbes [43]. Mucins also

provide nutritional benefits to gut microbes and their hosts [43], which are particularly beneficial

during winter fasting. In the absence of dietary (typically plant) glycans, host-derived mucins

become the major substrates that support microbial growth and proliferation, and contribute to

host energetic needs by producing SCFAs that fuel IECs [44]. Coupled with our previous

observation that mucin (MUC2) secretion continues throughout the hibernation season [17],

these molecules, like urea, may serve as valuable sources of carbon, nitrogen and energy

during the winter fast when exogenous nutrients are absent. Intact mucins are not digestible by

mammalian enzymes, but bacteria that express certain glycosyl hydrolases (GHs) can cleave

and metabolize glycan linkages, releasing oligosaccharides that are further fermented by the

same or other bacteria for energy, and in so doing produce beneficial metabolites for the host

[45]. There are two features of the gut ecosystem that must be in place for mucin glycan

degradation to proceed during hibernation: a thermally permissive environment for reactions

catalyzed by GHs and other enzymes, and a mechanism to dispose of $H_2$ whose accumulation

in the lumen slows fermentation. The thermal requirement is met by the ~12-24 h IBA periods

when host body temperature is above 30 ºC. The requirement for $H_2$ disposal, which is likely

compromised in hibernators by reduced luminal $CO_2$ generation due to the absence of

fermentable dietary substrates, can be met by the dual actions of $CO_2$ production by ureolytic

bacteria and its subsequent use in hydrogen disposal by acetogens.

We initially planned to examine seasonal variation in recycling of mucin-derived carbon

from the microbiota into squirrel tissues using isotope labeling methods similar to those used for

urea [2], but were unable to do so due to a lack of commercially available $^{13}$C-labeled mucin

oligosaccharides that are resistant to breakdown by mammalian enzymes. We therefore

investigated whether the capacity for mucin degradation in squirrel microbiotas changes over

the annual hibernation cycle. Our metagenomic analysis suggests an important role for bacteria

with mucin degrading capacity during winter fasting, as the five CAZymes gene families that are

enriched in Winter compared with Summer microbiomes all have mucin degradation capacity,

whereas those enriched in Summer microbiomes include a mix of mucin and non-mucin

degraders. This is consistent with our previous findings that several of the bacteria whose

relative abundances increase in Winter are known mucin degraders [17,28]. Moreover, all five

genes enriched in the Winter microbiomes were classified to at least one member of the phylum

Bacteroidota, whose relative abundance in the squirrel microbiota increases over 2-fold from

Summer to Winter [16,17]. Noteworthy genera to which winter-enriched gene families are

classified include *Akkermansia* and *Alistipes spp*. *Akkermansia muciniphila*, which is the sole

member of the phylum Verrucomicrobiota present in the ground squirrel microbiota [15,16], can

subsist on mucin alone as its sole carbon and nitrogen source [46]. This property may be

responsible for the over 4-fold increase in relative abundance of Verrucomicrobiota during

hibernation [16,17]. The genus *Alistipes*, a member of Bacteroidota and another prominent

mucin degrader [45], also increases during hibernation [16,17]. Interestingly, beyond its role as

a mucin degrader, *Alistipes* has the highest percentage of urease genes in the ground squirrel

microbiome during hibernation, increasing nearly two-fold in abundance from Summer to Winter

[2]. It therefore plays dual roles in processing of host endogenous nutrients with potential

benefits to the host and its microbiota.

*Conclusion*

Collectively, these findings introduce UCS as a novel, mutually beneficial mechanism through

which hibernating mammals and their gut microbes meet metabolic demands during prolonged

fasting. We propose a process whereby carbon limitation in the gut of fasted hibernators is

ameliorated by a sequential set of host-microbe interactions involving first ureolytic and then

acetogenic bacteria. This two-step mutualism, initiated by urea hydrolysis, first generates

acetate that is avidly absorbed and oxidized by the hibernator host at levels that exceed

summer values and may participate in other host functions including immunity and intestinal

barrier function. Next, via hydrogen disposal, it helps sustain the gut ecosystem as an efficient

environment for microbial fermentation of host compounds such as mucin glycoproteins, thus

unlocking the value of endogenous nutrient sources for both the microbiota and host. In

conjunction with UNS — which enables sustained protein synthesis in host tissues during

hibernation [2] — UCS establishes urea as a valuable endogenous nutrient source throughout

the winter fast and underscores the integral role that gut microbiota play in hibernation's

metabolic phenotype. Consistent shifts in gut microbial composition observed across various

hibernating mammals [47,48] and the finding that UCS also takes place in summer squirrels

suggest that urea's dual role in enabling its nitrogen and carbon to be recycled into molecules

with nutritional value may be widespread among mammals. Taken together, these insights

deepen our understanding of the host–microbe interactions that underlie metabolic resilience

and highlight the vital roles gut microbiota play as animals endure extended periods without

external nutrient sources.

**References**

1.  Staples JF. 2016 Metabolic Flexibility: Hibernation, Torpor, and Estivation. In
*Comprehensive Physiology* (ed R Terjung), pp. 737–771. Hoboken, NJ, USA: John Wiley &
Sons, Inc. (doi:10.1002/cphy.c140064)

2.  Regan MD, Chiang E, Liu Y, Tonelli M, Verdoorn KM, Gugel SR, Suen G, Carey HV,
Assadi-Porter FM. 2022 Nitrogen recycling via gut symbionts increases in ground squirrels
over the hibernation season. *Science* **375**, 460–463. (doi:10.1126/science.abh2950)

3.  Hindle AG, Otis JP, Epperson LE, Hornberger TA, Goodman CA, Carey HV, Martin SL.
2015 Prioritization of skeletal muscle growth for emergence from hibernation. *J. Exp. Biol.*
**218**, 276–284. (doi:10.1242/jeb.109512)

4.  Langmead B, Salzberg SL. 2012 Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
**9**, 357–359. (doi:10.1038/nmeth.1923)

5.  Li H *et al.* 2009 The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079. (doi:10.1093/bioinformatics/btp352)

6.  Mikheenko A, Saveliev V, Gurevich A. 2016 MetaQUAST: evaluation of metagenome
assemblies. *Bioinformatics* **32**, 1088–1090. (doi:10.1093/bioinformatics/btv697)

7.  Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010 Prodigal:
prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*
**11**. (doi:10.1186/1471-2105-11-119)

8.  El-Gebali S *et al.* 2019 The Pfam protein families database in 2019. *Nucleic Acids Res.* **47**,
D427–D432. (doi:10.1093/nar/gky995)

9.   Kim D, Song L, Breitwieser FP, Salzberg SL. 2016 Centrifuge: rapid and sensitive
classification of metagenomic sequences. *Genome Res.* **26**, 1721–1729.
(doi:10.1101/gr.210641.116)

10. Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. 2022 The
carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Res.* **50**,
D571–D577. (doi:10.1093/nar/gkab1045)

11. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012 dbCAN: a web resource for automated
carbohydrate-active enzyme annotation. *Nucleic Acids Res.* **40**, W445-451.
(doi:10.1093/nar/gks479)

12. Nayfach S, Pollard KS. 2015 Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome.
*Genome Biol.* **16**, 51. (doi:10.1186/s13059-015-0611-7)

13. Team RC. 2020 R A language and environment for statistical computing, R Foundation for
Statistical. *Computing*.

14. Love MI, Huber W, Anders S. 2014 Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550. (doi:10.1186/s13059-014-0550-8)

15. Stevenson TJ, Duddleston KN, Buck CL. 2014 Effects of Season and Host Physiological
State on the Diversity, Density, and Activity of the Arctic Ground Squirrel Cecal Microbiota.
*Appl. Environ. Microbiol.* **80**, 5611–5622. (doi:10.1128/AEM.01537-14)

16. Carey HV, Walters WA, Knight R. 2013 Seasonal restructuring of the ground squirrel gut
microbiota over the annual hibernation cycle. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.*
**304**, R33–R42. (doi:10.1152/ajpregu.00387.2012)

17. Dill-McFarland KA, Neil KL, Zeng A, Sprenger RJ, Kurtz CC, Suen G, Carey HV. 2014
Hibernation alters the diversity and composition of mucosa-associated bacteria while
enhancing antimicrobial defence in the gut of 13-lined ground squirrels. *Mol. Ecol.* **23**,
4658–4669. (doi:10.1111/mec.12884)

18. Prins RA, Lankhorst A. 1977 Synthesis of acetate from $CO_2$ in the cecum of some rodents.
*FEMS Microbiol. Lett.* **1**, 255–258. (doi:10.1111/j.1574-6968.1977.tb00627.x)

19. Groisman EA, Han W, Krypotou E. 2023 Advancing the fitness of gut commensal bacteria.
*Science* **382**, 766–768. (doi:10.1126/science.adh9165)

20. Schofield WB, Zimmermann-Kogadeeva M, Zimmermann M, Barry NA, Goodman AL. 2018
The Stringent Response Determines the Ability of a Commensal Bacterium to Survive
Starvation and to Persist in the Gut. *Cell Host Microbe* **24**, 120-132.e6.
(doi:10.1016/j.chom.2018.06.002)

21. Zhu M, Dai X. 2023 Stringent response ensures the timely adaptation of bacterial growth to
nutrient downshift. *Nat. Commun.* **14**, 467. (doi:10.1038/s41467-023-36254-0)

22. Wolfe AJ. 2005 The Acetate Switch. *Microbiol. Mol. Biol. Rev.* **69**, 12–50.
(doi:10.1128/mmbr.69.1.12-50.2005)

23. Enjalbert B, Letisse F, Portais J-C. 2013 Physiological and Molecular Timing of the Glucose
to Acetate Transition in Escherichia coli. *Metabolites* **3**, 820–837.
(doi:10.3390/metabo3030820)

24. Regan MD, Chiang E, Martin SL, Porter WP, Assadi-Porter FM, Carey HV. 2019 Shifts in
metabolic fuel use coincide with maximal rates of ventilation and body surface rewarming in
an arousing hibernator. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **316**, R764–R775.
(doi:10.1152/ajpregu.00379.2018)

25. Fagundes RR, Belt SC, Bakker BM, Dijkstra G, Harmsen HJM, Faber KN. 2024 Beyond
butyrate: microbial fiber metabolism supporting colonic epithelial homeostasis. *Trends
Microbiol.* **32**, 178–189. (doi:10.1016/j.tim.2023.07.014)

26. Clausen MR, Mortensen PB. 1994 Kinetic studies on the metabolism of short-chain fatty
acids and glucose by isolated rat colonocytes. *Gastroenterology* **106**, 423–432.
(doi:10.1016/0016-5085(94)90601-7)

27. Bekebrede AF, de Boer VCJ, Gerrits WJJ, Keijer J. 2023 Functional and molecular profiling
of fasted piglets reveals decreased energy metabolic function and cell proliferation in the
small intestine. *Am. J. Physiol.-Gastrointest. Liver Physiol.* **325**, G539–G555.
(doi:10.1152/ajpgi.00240.2022)

28. Carey HV. 1990 Seasonal changes in mucosal structure and function in ground squirrel
intestine. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **259**, R385–R392.
(doi:10.1152/ajpregu.1990.259.2.R385)

29. Carey HV, Sills NS. 1992 Maintenance of intestinal nutrient transport during hibernation.
*Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **263**, R517–R523.
(doi:10.1152/ajpregu.1992.263.3.R517)

30. Carey HV, Sills NS. 1996 Hibernation enhances D-glucose uptake by intestinal brush border
membrane vesicles in ground squirrels. *J. Comp. Physiol. [B]* **166**, 254–261.
(doi:10.1007/BF00262869)

31. Wolin MJ, Miller TL, Yerry S, Zhang Y, Bank S, Weaver GA. 1999 Changes of Fermentation
Pathways of Fecal Microbial Communities Associated with a Drug Treatment That Increases
Dietary Starch in the Human Colon. *Appl. Environ. Microbiol.* **65**, 2807–2812.
(doi:10.1128/AEM.65.7.2807-2812.1999)

32. Andrews MT, Russeth KP, Drewes LR, Henry P-G. 2009 Adaptive mechanisms regulate
preferred utilization of ketones in the heart and brain of a hibernating mammal during
arousal from torpor. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **296**, R383–R393.
(doi:10.1152/ajpregu.90795.2008)

33. Cheng C-W *et al.* 2019 Ketone Body Signaling Mediates Intestinal Stem Cell Homeostasis
and Adaptation to Diet. *Cell* **178**, 1115-1131.e15. (doi:10.1016/j.cell.2019.07.048)

34. Wang Q, Zhou Y, Rychahou P, Fan TW-M, Lane AN, Weiss HL, Evers BM. 2017
Ketogenesis contributes to intestinal cell differentiation. *Cell Death Differ.* **24**, 458–468.
(doi:10.1038/cdd.2016.142)

35. Martin SL, Epperson LE, Rose JC, Kurtz CC, Ané C, Carey HV. 2008 Proteomic analysis of
the winter-protected phenotype of hibernating ground squirrel intestine. *Am. J. Physiol.-*
*Regul. Integr. Comp. Physiol.* **295**, R316–R328. (doi:10.1152/ajpregu.00418.2007)

36. Sokolović M *et al.* 2007 Fasting induces a biphasic adaptive metabolic response in murine
small intestine. *BMC Genomics* **8**, 361. (doi:10.1186/1471-2164-8-361)

37. Crawford PA, Crowley JR, Sambandam N, Muegge BD, Costello EK, Hamady M, Knight R,
Gordon JI. 2009 Regulation of myocardial ketone body metabolism by the gut microbiota
during nutrient deprivation. *Proc. Natl. Acad. Sci.* **106**, 11276–11281.
(doi:10.1073/pnas.0902366106)

38. Kurtz CC, Carey HV. 2007 Seasonal changes in the intestinal immune system of hibernating
ground squirrels. *Dev. Comp. Immunol.* **31**, 415–428. (doi:10.1016/j.dci.2006.07.003)

39. Takeuchi T *et al.* 2021 Acetate differentially regulates IgA reactivity to commensal bacteria.
*Nature* **595**, 560–564. (doi:10.1038/s41586-021-03727-5)

40. Wu W *et al.* 2017 Microbiota metabolite short-chain fatty acid acetate promotes intestinal
IgA response to microbiota which is mediated by GPR43. *Mucosal Immunol.* **10**, 946–956.
(doi:10.1038/mi.2016.114)

41. Smith NW, Shorten PR, Altermann EH, Roy NC, McNabb WC. 2019 Hydrogen cross-
feeders of the human gastrointestinal tract. *Gut Microbes* **10**, 270–288.
(doi:10.1080/19490976.2018.1546522)

42. Xiao G, Liu S, Xiao Y, Zhu Y, Zhao H, Li A, Li Z, Feng J. 2019 Seasonal Changes in Gut
Microbiota Diversity and Composition in the Greater Horseshoe Bat. *Front. Microbiol.* **10**.
(doi:10.3389/fmicb.2019.02247)

43. Wang BX, Wu CM, Ribbeck K. 2021 Home, sweet home: how mucus accommodates our
microbiota. *FEBS J.* **288**, 1789–1799. (doi:10.1111/febs.15504)

44. Berkhout MD, Plugge CM, Belzer C. 2022 How microbial glycosyl hydrolase activity in the
gut mucosa initiates microbial cross-feeding. *Glycobiology* **32**, 182–200.
(doi:10.1093/glycob/cwab105)

45. Glover JS, Ticer TD, Engevik MA. 2022 Characterizing the mucin-degrading capacity of the
human gut microbiota. *Sci. Rep.* **12**, 8456. (doi:10.1038/s41598-022-11819-z)

46. Derrien M, Vaughan EE, Plugge CM, de Vos WM. 2004 Akkermansia muciniphila gen. nov.,
sp. nov., a human intestinal mucin-degrading bacterium. *Int. J. Syst. Evol. Microbiol.* **54**,
1469–1476. (doi:10.1099/ijs.0.02873-0)

47. Carey HV, Assadi-Porter FM. 2017 The Hibernator Microbiome: Host-Bacterial Interactions
in an Extreme Nutritional Symbiosis. *Annu. Rev. Nutr.* **37**, 477–500. (doi:10.1146/annurev-
nutr-071816-064740)

48. Kurtz CC, Otis JP, Regan MD, Carey HV. 2021 How the gut and liver hibernate. *Comp.*
*Biochem. Physiol. A. Mol. Integr. Physiol.* **253**, 110875. (doi:10.1016/j.cbpa.2020.110875)

**Figure captions**

**Figure 1.** Principal components analyses (PCA) for $^{13}$C-enriched metabolomes from 13-lined ground squirrel cecal contents (A), liver (B), and quadriceps muscle (C). Treatment groups include gut microbiota-intact and -depleted groups from Summer (sum), early winter (EW), and late winter (LW) squirrels. PERMANOVA results for each tissue are presented next to its respective score plot.

**Figure 2**. $^{13}$C-urea carbon incorporation into metabolomes for cecal content (A), liver (B), and quadriceps muscle (C) of microbiome-intact (filled bars) and microbiome-depleted (open bars) 13-lined ground squirrels. For each panel, the plots portray $^{13}$C-incorporation into the metabolite named above the plot. Asterisks indicate a significant effect of season (Season) or gut microbiota presence/absence (ABX), where * means $P<0.05$, ** means $P<0.01$, and *** means $P<0.001$. n = 5 for all data.

**Figure 3.** (A) Breath $\delta^{13}$C values for 13-lined ground squirrels treated with $^{13}$C-labelled urea. The data represent total area under the curve (AUC) for the 3 h period following urea treatment, as first reported in [2]. The data are expressed here exponentially to eliminate negative values and thus enable the urea-to-acetate efficiency calculations. (B) Urea-to-acetate efficiency, which is the ratio of an individual squirrel's $^{13}$C-enriched cecal acetate concentration to its breath $\delta^{13}$C AUC value.

**Figure 4.** Seasonal effect on (A) the percentage of gut bacterial genes in the 13-lined ground squirrel gut microbiome related to the Wood-Ljungdahl pathway for biological carbon fixation, as well as (B) the top 10 bacterial taxa from taxonomic classification of Wood-Ljungdahl genes as determined by sequence abundance.

**Figure 5.** (A) Mean breath $\delta^{13}$C for Summer (green) and Late Winter (blue) 13-lined ground squirrels treated with $^{13}$C-labeled acetate via oral gavage at time 0. Horizontal and vertical error bars represent SEM, and closed and open symbols represent squirrels with intact and depleted microbiotas. (B) Mean area under the curve (AUC) values for data in (A), with 2-way ANOVA results presented on the graph and asterisks indicating significant differences as determined by post-hoc test, where * indicates $P<0.05$, ** indicates $P<0.01$, and *** indicates $P<0.001$.

**Figure 6.** Glycoside hydrolase (GH) carbohydrate-active enzymes (CAZymes) involved in mucin

degradation that are seasonally enriched in Late Winter compared with Summer squirrels.

Displayed counts represent DESeq2-normalized counts that were Hellinger-transformed. Darker

colors indicate higher counts and lighter colors indicate lower counts. T-test results: GH20

$P=0.0043$; GH33 $P<0.0001$; GH84 $P=0.0048$; GH85 $P=0.0027$; GH109 $P=0.0051$. N=6 for

Summer, N=5 for Later Winter.

**A  Cecum content**

PC 2 (21.1 %)

- ○ EW-Depleted
- ○ EW-Intact
- ○ LW-Depleted
- ○ LW-Intact
- ○ Sum-Depleted
- ○ Sum-Intact

PERMANOVA
F-value: 3.0992
R$^2$: 0.40254
P-value: 0.007

PC 1 ( 48.7 %)

**B  Liver**

PC 2 (17.1 %)

- ○ EW-Depleted
- ○ EW-Intact
- ○ LW-Depleted
- ○ LW-Intact
- ○ Sum-Depleted
- ○ Sum-Intact

PERMANOVA
F-value: 2.7784
R$^2$: 0.37656
P-value: 0.009

PC 1 ( 24.2 %)

**C  Muscle**

PC 2 (15 %)

- ○ EW-Depleted
- ○ EW-Intact
- ○ LW-Depleted
- ○ LW-Intact
- ○ Sum-Depleted
- ○ Sum-Intact

PERMANOVA
F-value: 1.2785
R$^2$: 0.21749
P-value: 0.288

PC 1 ( 30.1 %)

**Figure 1.** Principal components analyses (PCA) for 13C-enriched metabolomes from 13-lined ground squirrel cecal contents (A), liver (B), and quadriceps muscle (C). Treatment groups include gut microbiota-intact and -depleted groups from Summer (sum), early winter (EW), and late winter (LW) squirrels. PERMANOVA results for each tissue are presented next to its respective score plot.

**Figure 2**. [13]C-urea carbon incorporation into metabolomes for cecal content (A), liver (B), and quadriceps muscle (C) of microbiome-intact (filled bars) and microbiome-depleted (open bars) 13-lined ground squirrels. For each panel, the plots portray [13]C-incorporation into the metabolite named above the plot. Asterisks indicate a significant effect of season (Season) or gut microbiota presence/absence (ABX), where * means $P<0.05$, ** means $P<0.01$, and *** means $P<0.001$. n = 5 for all data.

**Figure 3**. (A) Breath $\delta^{13}C$ values for 13-lined ground squirrels treated with $^{13}C$-labelled urea. The data represent total area under the curve (AUC) for the 3 h period following urea treatment, as first reported in [2]. The data are expressed here exponentially to eliminate negative values and thus enable the urea-to-acetate efficiency calculations. (B) Urea-to-acetate efficiency, which is the ratio of an individual squirrel's 13C-enriched cecal acetate concentration to its breath $\delta^{13}C$ AUC value.

**A**

**B**

| | Summer | | Late Winter | |
|---|---|---|---|---|
| **Taxon** | **% W-L Genes** | **Taxon** | | **% W-L Genes** |
| *Bacteroides* | 1.74% | *Blautia* | | 3.63% |
| *Clostridium* | 0.96% | *Clostridium* | | 1.67% |
| *Desulfallas* | 0.92% | *Bacteroides* | | 1.60% |
| *Alistipes* | 0.78% | *Lachnoclostri* | | 0.73% |
| *Terrabacteria* | 0.78% | *Terrabacteria* | | 0.58% |
| *Treponema* | 0.74% | *Eggerthella* | | 0.58% |
| *Blautia* | 0.67% | *Treponema* | | 0.51% |
| *Lactobacillus* | 0.63% | *Oscillibacter* | | 0.51% |
| *Neisseria* | 0.55% | *Olsenella* | | 0.44% |
| *Corynebacter* | 0.37% | *Desulfallas* | | 0.44% |

**Figure 4.** Seasonal effect on (A) the percentage of gut bacterial genes in the 13-lined ground squirrel gut microbiome related to the Wood-Ljungdahl pathway for biological carbon fixation, as well as (B) the top 10 bacterial taxa from taxonomic classification of Wood-Ljungdahl genes as determined by sequence abundance.

**Figure 5.** (A) Mean breath δ¹³C for Summer (green) and Late Winter (blue) 13-lined ground squirrels treated with 13C-labeled acetate via oral gavage at time 0. Horizontal and vertical error bars represent SEM, and closed and open symbols represent squirrels with intact and depleted microbiotas. (B) Mean area under the curve (AUC) values for data in (A), with 2-way ANOVA results presented on the graph and asterisks indicating significant differences as determined by post-hoc test, where * indicates P<0.05, ** indicates P<0.01, and *** indicates P<0.001.

**Figure 6.** Glycoside hydrolase (GH) carbohydrate-active enzymes (CAZymes) involved in mucin degradation that are seasonally enriched in Late Winter compared with Summer squirrels. Displayed counts represent DESeq2-normalized counts that were Hellinger-transformed. Darker colors indicate higher counts and lighter colors indicate lower counts. T-test results: GH20 P=0.0043; GH33 P<0.0001; GH84 P=0.0048; GH85 P=0.0027; GH109 P=0.0051. N=6 for Summer, N=5 for Later Winter.

**Figure S1.** Isotope ratio mass spectrometry-based measurements of [13]C-urea carbon incorporation into the protein fractions of (A) cecum content, (B) liver and (C) quadriceps muscle. [13]C-urea carbon incorporation into "total" cecum content is also shown (D), which is an analyses run on the total cecal content milieu, not just the protein fraction. Cecal content panels (A, D) lack microbiota-depleted data because ABX-treated animals had insufficient content for analyses. Asterisks indicate a significant effect of season (Season) or gut microbiota presence/absence (ABX), where ** indicates $P<0.01$ and *** indicates $P<0.001$. n = 5 for all data except for cecum content (A, D; n=1, pooled samples).