Dehalobacter dechlorinates dichloroanilines and contributes to the natural attenuation of dichloronitrobenzenes at a complex industrial site

Sofia P. Araújo,[a] Line Lomheim,[a] Suzana P. Q. Kraus,[b] E. Erin Mack,[c] Jim C. Spain,[d] Sávia Gavazza,[e] Elizabeth A. Edwards[a]*

[a]Department of Chemical Engineering and Applied Chemistry, University of Toronto, Toronto, Ontario, M5S 3E5, Canada.

[b]Geosyntec Consultants International Inc, Etobicoke, Ontario, M8X 1Y9, Canada

[c]Corteva Remediation Group, Wilmington, Delaware, 19805, United States

[d]Center for Environmental Diagnostics and Bioremediation, University of West Florida, Pensacola, Florida, 32514-5751, United States

[e]Departamento de Engenharia Civil e Ambiental, Universidade Federal de Pernambuco, Recife, Pernambuco, 50740-530, Brazil

*Corresponding author: elizabeth.edwards@utoronto.ca

ABSTRACT

The potential for bioremediation of dichloroanilines (2,3- and 3,4-DCA) and

dichloronitrobenzenes (3,4-, 2,5- and 2,3-DCNB) was investigated using inocula from an industrial

site in northeast of Brazil. Anaerobic biotransformation of these chlorinated compounds was

observed in microcosms simulating site conditions, particularly when electron donor was added.

To disentangle specific transformation reactions, sub-cultures from active anaerobic microcosms

were enriched with individual DCA or DCNB isomers. In these single-compound enrichment

cultures, DCNB isomers were completely reduced to the corresponding DCA isomers. Each DCA

isomer was stoichiometrically dechlorinated to the corresponding monochloroaniline (CA)

isomers. Further dechlorination of CA to aniline was noted. The reduction of DCNB to DCA was

sustained by known fermenters and possibly *Desulfitobacterium*, while *Dehalobacter* was clearly

responsible for dechlorination of DCAs. These findings helped to attribute function to microbial

community data retrieved from field samples to inform the conceptual site model and enable

identification of bioactive zones and rate-limiting conditions at a complex site. More generally,

these data broaden the understanding of the metabolic repertoire of *Dehalobacter* and provide a

model for incorporating microbial community data into site remediation efforts.

Graphical Abstract

Keywords: Dichloronitrobenzene;   Dichloroaniline;   Monochloroaniline;   anaerobic

biotransformation; bioremediation.

Short synopsis statement

A diverse microbial community contributes to natural attenuation at a complex contaminated

site. Notably, dichloroanilines produced by biologically-mediated nitroreduction of

dichloronitrobenzenes were dechlorinated by *Dehalobacter*.

1   INTRODUCTION

Many historically-contaminated sites worldwide are a challenge to remediate owing to complex

mixtures of toxic contaminants and heterogeneous stratigraphy. For example, an industrial facility

located in Bahia in NE Brazil has been the subject of several remedial investigations.[1,2] As a former

manufacturing facility of the herbicide diuron, it is impacted by high concentrations of multiple

compounds including dichloronitrobenzenes (DCNB; ~3,000 mg/kg) and dichloroanilines (DCA;

~700 mg/kg) among many others.[1,2] Anoxic conditions typically prevail in regions with low

permeability, high organic matter, or high contaminant concentrations, where the rate of oxygen

depletion exceeds rates of replenishment by diffusion or groundwater flow. Therefore, we sought

to understand the fate of DCNB and DCA isomers under anoxic conditions with particular

attention to associated microbial populations to elucidate microbial pathways and thus create a

better conceptual model of the site and a more effective remediation strategy.

In subsurface anoxic environments, nitroaromatic compounds tend to become reduced to their

corresponding aromatic amines. As early as 1975, the pesticide pentachloronitrobenzene was

shown to be reduced to pentachloroaniline.[3] Nitro group reduction to the corresponding amine is

now understood to involve broad specificity and oxygen-independent nitroreductases and related

enzymes that are widely distributed in bacteria and archaea.[4] These reactions produce toxic

transient nitroso and hydroxylamino intermediates, which are highly reactive and unstable.[4–6]

Nitroaromatic reduction may also be a respiratory process akin to nitrate reduction.[7–9] A 1983

study reported the reduction of 3,4-DCNB to a variety of poorly quantified products including 3,4-

DCA, and condensation products.[5] Despite their environmental relevance, few studies have

investigated the anaerobic biodegradation of DCNB isomers.

Literature on the anaerobic biodegradation of chloroanilines is also limited. The reductive

dechlorination of 3,4-DCA to a mixture of monochloroaniline isomers (3-CA and 4-CA) under

anaerobic/methanogenic conditions was first reported in 1989 and 1990, but the organisms

responsible were not identified.[10–12] Under nitrate-reducing conditions, a novel reductive

deamination reaction was observed in a *Rhodococcus* strain where 3,4-DCA was deaminated

primarily to dichlorobenzene,[13] however, additional reports of this process could not be found.

More recently, *Dehalococcoides mccartyi* strain CBDB1 and *Dehalobacter sp*. strain 14DCB1,

both obligate organohalide-respiring-bacteria (OHRB) that dechlorinate chlorinated benzenes,

were shown to dechlorinate 2,3-DCA to 2- and 3-CA and 3,4-DCA to 3-CA with no further

transformation when acetate was provided as carbon source and hydrogen as electron donor.[14]

Reports of further dechlorination of monochloroanilines to aniline are sparse and inconclusive.

Early reports (1997-98) described dechlorination of DCA to CA and then to small amounts of

aniline under sulfate-reducing conditions.[15,16] These studies did not characterize the microbial

community. A 2019 study showed that *Geobacter* sp. KT5 could transform dichloroanilines under

iron-reducing conditions; mineralization of the products via aniline and 4-aminobenzoate was

proposed.[17] Two years later, a mixed culture composed of *Geobacter* sp., *Paracoccus*

denitrificans, *Pseudomonas* sp., and *Rhodococcus* sp. strains was described to transform propanil

to 3,4-DCA, and further to 3-CA, chlorobenzene, and aniline, prior to mineralization.[18] Another

study found that 2-aminoanthraquinone-graphene oxide enhanced dechlorination in an enriched

anaerobic sludge community that was able to dechlorinate 2-CA. In this work, some aniline and

hexanoic acid were detected and organisms from *Oscillospira*, *Lactobacillales*, *Veillonellaceae*,

and *Ruminococcus* groups were positively correlated with 2-CA transformation.[19]

Given the scarce literature on the biodegradation of dichloronitrobenzenes and dichloroanilines,

the goal of this study was to identify organisms responsible for transformation reactions of such

compounds and to use the information to design biomarkers to track biodegradation and delineate

bioactive zones across a contaminated site. We determined that *Dehalobacter* plays a very active

and critical role in the dechlorination of dichloroanilines at a complex site in Brazil. This

knowledge is crucial for developing cost-effective *in situ* remediation strategies for many other

complex contaminated sites.

2   MATERIALS AND METHODS

2.1   Chemicals and stock solutions

Substrates added to enrichment cultures in this study were 3,4-dichloronitrobenzene (3,4-DCNB), 2,5-dichloronitrobenzene (2,5-DCNB), 2,3-dichloronitrobenzene (2,3-DCNB), 3,4-dichloroaniline (3,4-DCA), and 2,3-dichloroaniline (2,3-DCA) (Sigma Aldrich, purity > 99%, (Figure S1). Stock solutions of each compound were prepared in acetone (Sigma Aldrich) at a concentration of 5 g/L. Ethanol (100%: Sigma Aldrich) and 60% sodium lactate (TCI America) were used as electron donors. The mentioned DCNB and DCA isomers, aniline, 2-, 3-, and 4-chloroaniline (CA), 2,5-DCA, DCB isomers (Sigma Aldrich, purity > 99%), and methane (Scott) were also used as analytical standards. Methanol (Sigma Aldrich) was used as the solvent for the chlorinated standards and acetonitrile (Fisher Scientific) was used for the mobile phase for liquid chromatography (described below).

2.2   Preparation of original microcosm study and enrichment cultures

The original microcosms were constructed in 2015 with solid subsurface material and groundwater from the industrial facility in Brazil, incubated under anaerobic conditions at site pH (4.6 to 6.5), and amended with various electron donors and acceptors. The microcosms were amended with various mixtures of chloroanilines, dichloroanilines, dichloronitrobenzenes and chlorobenzenes at concentrations typical of the site, and biodegradation was monitored for over one year. The microcosm setup is detailed in Supporting Information Text S1 and in a Master's Thesis.[20] Results are summarized in Table S1 and Figure S2.

Five microcosms established with electron donors to stimulate methanogenic conditions

(Cam16, Cam17, Cam64, Cam88, and Cam89) showed the most activity and were therefore

selected for enrichment (Figure S1). All transfer cultures as well as controls were prepared in

autoclaved 250-mL Boston round bottles sealed with Mininert caps, containing 100 mL of

anaerobic mineral medium (pH 7) reduced with iron sulfide (FeS).[21] Only medium supernatant

was used to avoid adding precipitated FeS. To simulate site groundwater conditions and to promote

mineralization of nitrogen-containing compounds, ammonium was omitted from the medium

recipe. All cultures were incubated statically in the dark at room temperature in an anaerobic

glovebox (Coy) supplied with a gas mix of $N_2:CO_2:H_2$ (80:10:10%).

The first transfer (T1) cultures were inoculated in March 2020 with the cell pellet from the

respective parent microcosm. The pellet was obtained from a 40 ml sample from a shaken

microcosm by centrifuging at 12,000 g for 15 min. These T1 enrichment cultures were amended

with individual target compounds and ethanol and lactate as electron donors (Figure S1). An

autoclaved killed control (KC) was also prepared with a cell pellet from an original microcosm

(Cam 65). This control was autoclaved three times on three consecutive days prior to being

amended with a mix of six compounds (3,4-DCNB, 2,5-DCNB, 3,4-DCA, 2,3-DCA, 2-CA, and

aniline). An abiotic control (AC1) was also established containing only reduced mineral medium

amended with a slightly different set of six compounds (3,4-DCNB, 2,3-DCNB, 3,4-DCA, 2,3-

DCA, 2-CA, and aniline).

The 5 active T1 transfer cultures, labelled 34DCNB-T1, 25DCNB-T1, 23DCNB-T1, 34DCA-

T1 and 23DCA-T1, were repeatedly amended with individual chlorinated substrates (Figure S1).

These semi-volatile substrates (electron acceptors) were added as evaporated acetone stocks as

described in Text S2. The target concentration was initially 10 mg/L, which was increased to ~27

mg/L over time. Electron donors were also added to the active cultures and the two controls (AC1

and KC) initially and were re-amended to active cultures when degradation rates slowed. Electron

donor was added in ~17-fold excess in electron equivalents (eeq) considering both nitroreduction

and complete dechlorination of DCNB to aniline (Text S2). The initial pH was between 6.6 and

7.1 in all active enrichments.

A second set of transfer cultures (T2) were prepared from T1 cultures after ~ 4 months (Figure

S1). The DCNB T2 cultures were inoculated with a 50% transfer from T1 cultures (50 mL into 50

mL fresh medium) while DCA T2 cultures were inoculated with a 10% transfer from T1 cultures

(10 mL of T1 into 90 mL). Additional mineral medium supernatant was added to a culture bottle

when its volume was depleted to below 50 mL due to sampling. Occasionally, a little more medium

was added to gradually increase the volume of culture. A second abiotic control (AC2) was

prepared containing 3,4-DCNB, 2,3-DCNB, 3,4-DCA, and 2,3-DCA, and mineral medium devoid

of FeS. The data presented herein focus on the second transfer (T2) enrichment cultures.

2.3   Analytical procedures

Concentrations of DCNB, DCA and CA isomers and aniline were measured using a Hewlett-

Packard 1050 series high-performance liquid chromatography (HPLC) system (Acclaim™ 120

C18 column, Thermo Scientific) equipped with a UV detector (set at 254 nm). The mobile phase

was an isocratic 50:50 mixture of Milli-Q water and acetonitrile (Millipore Sigma) at 1 mL/min

with a runtime of 25 min. Samples (2 mL) were taken from cultures using a 2 mL gastight glass

syringe and filtered through a 0.22 μm (13 mm) Chromspec PTFE syringe filter (pre-wetted with

methanol) prior to being injected into the HPLC. Lactate, acetate, propionate, formate, butyrate,

pyruvate, chloride, nitrate, nitrite, sulfate, and phosphate were measured by ion chromatography

(IC) using a Dionex™ Integrion™ IC system (Thermo Fisher Scientific). Methane was measured

in headspace samples by gas chromatography (GC) using an Agilent 7890A gas chromatograph

with headspace autosampler G1888 (Agilent Technologies). Method details are provided in Text

S3.

2.4   Microbial community analysis and qPCR

Well-mixed samples (4 mL) from enrichment cultures were collected periodically, centrifuged

at 10,000 rpm for 20 minutes, the supernatant removed, and cell pellets stored at -80ºC. DNA was

extracted using the DNeasy PowerSoil Kit (Qiagen) and quantified using a Qubit® 3.0

Fluorometer (Thermo Fisher Scientific). DNA samples were sent for Illumina MiSeq PE300

amplicon sequencing at an external laboratory (Genome Quebec Innovation Centre, Montreal,

Canada), using modified "staggered end"[22] versions of primers 926f-modified (5'-

AAACTYAAAKGAATWGRCGG-3') and 1392r-modified (5'-ACGGGCGGTGWGTRC-3'),

which target the V6−V8 region of the 16S rRNA gene from both Bacteria and Archaea, as well as

the 18S rRNA gene in Eukarya.[23,24] Sequencing data was processed using QIIME 2 version

2021.2.[25] After trimming the primer region with the cutadapt plug-in, amplicon sequence variants

(ASVs) were generated using the DADA2 plugin,[26] and taxonomy was assigned to the ASVs

through the feature-classifier plugin using the SSU SILVA v.138 database.[27] Data were further

processed in Excel. Sequences <400 bp in length were removed and ASVs were ranked by

maximum sequence abundance across all DNA samples. ASVs with abundance greater than 1%

in at least one sample were kept for graphing. The Basic Local Alignment Search Tool (BLAST)

available at the National Center for Biotechnology Information (NCBI) was used to compare ASV

sequences of interest. Raw DNA sequences were uploaded to the NCBI (Accession:

PRJNA1040917; Table S2).

DNA samples were also analyzed by quantitative polymerase chain reaction (qPCR) using a

CFX96TM real-time PCR detection system (Bio-Rad Laboratories Inc.). Primers GenBac1055f

(5'-ATGGYTGTCGTCAGCT-3') and GenBac1392r (5'-ACGGGCGGTGTGTAC-3') targeting

most bacteria[24,28,29] and primers GenArch787f (5'-ATTAGATACCCGBGTAGTCC-3') and

GenArch1059r (5'-GCCATGCACCWCCTCT-3') targeting most archaea[30] were used. A no-

template control was also included in each run. All qPCR samples were run in triplicate. Details

on the qPCR method are provided in Text S4. Absolute total bacterial and archaeal abundance

obtained by qPCR (copies 16S rRNA gene per ml) was used to convert the relative abundances

obtained from amplicon sequencing to absolute abundance for each ASV.

A total of forty-eight DNA samples were sequenced and analyzed by qPCR, including samples

from the original subsurface solid material, microcosms, and enrichments. On average 66,000 16S

rRNA gene amplicon reads were retrieved per sample (Table S3a). Reads were grouped by bacteria

and archaea, then sequences with relative abundance >1% in any sample (Table S3b for bacteria

and Table S3c for archaea) were used to calculate absolute abundances (copies/mL) of each ASV

using total abundances of bacteria and archaea determined by qPCR (Table S4). Final absolute

abundances for each ASV (copies/mL) are reported in Tables S5a to S5f for bacteria; Tables S6a

to S6f for archaea. Finally, Non-metric Multi-dimensional Scaling (NMDS) and bubble plots were

constructed as explained in Text S5.

3   RESULTS AND DISCUSSION

3.1   Abiotic transformation of DCNB was not sustained

Contaminant profiles in three controls (AC1, AC2 and KC) were assessed for possible abiotic

reactions. Chloroanilines and aniline were not transformed in any of these controls.

Dichloronitrobenzenes were not transformed in medium control (AC2) without FeS (Figure S3;

Table S7). In the medium control (AC1) with FeS, 3,4-DCNB was not transformed, but 2,3-DCNB

was reduced to 2,3-DCA (Figure S4; Table S7); however, this activity was not sustained upon

reamending with 2,3-DCNB. These data are consistent with reports that reduced minerals, such as

FeS or perhaps reduced resazurin, can sometimes mediate reduction of the nitro group. [31–38]

Curiously, DCNB isomers were repeatedly transformed to DCA in the autoclaved control KC

(Figure S5; Table S7); however later analyses (to be discussed) revealed that KC was in fact not

sterile.

3.2   Biologically-mediated nitroreduction and dechlorination

Each transfer culture established on single substrates remained active, sustained by refeeding

chlorinated substrate and electron donors as needed for over 500 days. Historical data are provided

in Table S8 (T1) and Table S9 (T2). Focusing on T2 transfers from day 280 to 520 and on the three

DCNB-amended cultures first, we observed consistent and complete transformation of each

DCNB isomer to the corresponding dichloroaniline (Figure 1a-c). In the culture amended with 3,4-

DCNB the product, 3,4-DCA, was further dechlorinated to 3- and 4-CA (Figure 1a), while in the

cultures amended with 2,3- and 2,5-DCNB, the transformation stopped at the corresponding

dichloroaniline (Figures 1b and 1c). The sum of measured chlorinated products agreed well with

the cumulative DCNB added. Reduction of 2,3-DCNB (<~1 µM per day) was ~5 times slower

than the reduction of either 3,4- or 2,5-DCNB (5-6 µM per day).

Figure 1. Degradation profiles (upper panels a), b) and c); µmole/bottle) and corresponding major

microbial taxa (lower panels d), e) and f); copies/mL) in DCNB-amended T2 transfer cultures from

day 280 to 520. Cumulative DCNB added (———) is shown for each amendment (step) of 70 to

140 µM (10 to 20 µmol per bottle). The sum of products (– –○– –), including DCA (———■———), 4-

CA (———▲———), 3-CA (———▲———), 2-CA and aniline (not shown, mostly below detection), illustrates a

reasonable mole balance. Measured DCNB concentrations are not shown for clarity but can be

found in Table S9. In lower panels amplicon sequence variants (ASVs) from the same taxonomic

designation are assigned the same colour.

Focusing next on the two DCA-amended transfers, 3,4-DCA was dechlorinated to a mix of 3- and 4-CA (Figure 2a), while 2,3-DCA was dechlorinated primarily to 3-CA, with traces of 2-CA (Figure 2b). These cultures dechlorinated about ~5 to 6 µM DCA to CA per day.

Figure 2. Degradation profiles (upper panels a), b) and c); µmole/bottle) and corresponding major microbial taxa (lower panels, d) and e); copies/mL) in two DCA-amended T2 transfer cultures from day 280 to 520 (panels a), b), d), and e)) and from day 560 to 625 (panel c)). In panels a) and b), cumulative DCA added (———) is shown for each amendment (step) of ~80 µM (~12 µmol per bottle). The sum of products (– –o– –), including DCA (—■—), 4-CA (—▲—), 3-CA (—▲—), 2-CA (—▲—), and aniline (not shown, mostly below detection), illustrates a reasonable mole balance. Measured DCA concentrations are not shown for clarity but can be found in Table S9. Panel c) shows degradation profile after 2,3-DCA-T2 was fed with 2,3-DCNB (—●—) instead of

2,3-DCA on Day 568. The open circle shows the amount of 2,3-DCNB fed to 23DCA-T2 (─O─

). In lower panels amplicon sequence variants (ASVs) from the same taxonomic designation are

assigned the same color.

Given the very slow transformation rate in 23DCNB-T2, we wondered if the enrichment culture

dechlorinating 2,3-DCA (23DCA-T2; Figure 2b) could also reduce 2,3-DCNB, even though this

culture - and the microcosm (Cam16) from which it was derived - had never been exposed to any

DCNB isomer. Surprisingly, 2,3-DCNB was reduced without a lag to 2,3-DCA which was further

dechlorinated to 3-CA (Figure 2c). We then tested all active DCA-degrading cultures with a mix

of the three DCNB isomers; all the cultures were able to reduce DCNBs to their respective DCA

isomer (data not shown).

3.3   Microbial community composition reflects electron balance

To relate the microbial community to activity, we first identified abundant bacterial and archaeal

ASVs (>1%) and then calculated their absolute abundance in active cultures focusing on 2-time

points for each sample. Figure 3 provides a summary of the most relevant taxa identified. Most of

the enriched bacterial ASVs belong to the phylum Bacillota (Firmicutes) including *Sporomusa*,

Anaerospora, and unassigned members of the Veillonellales-Selenomonadales. These organisms

clearly grew on the ample electron donor amendments to the cultures. ASVs for

Anaeromyxobacter, *Desulfitobacterium* and *Desulfovibrio* were also retrieved. These genera are

also often implicated in nitrate or sulfate reduction though some members may be facultative

dechlorinators. ASVs for obligate dechlorinators of the genus *Dehalobacter* were also identified

(Figure 3). A complete overview of all major taxa (>1%) are shown in Figure S6, for bacteria, and

Figure S7, for archaea.

Figure 3. Absolute abundance of major organisms (copies/mL) and proportion of electron equivalents from amended donors channeled to each electron-accepting processes (% electron equivalents) for days 297 and 517. Veillonellales-Selenomonadales order is shortened to "Veill"; *Sporomusa* genus is shortened to "*Sporo*". Calculated absolute abundances for all ASVs are available in Table S5 (bacteria) and Table S6 (archaea).

To confirm the extent of dechlorination, an electron balance of possible donors and acceptors was completed using concentrations of chlorinated organics (Table S9) and anions (Table S10). As expected from stoichiometry (e.g., $C_6H_5Cl_2N + H_2 \rightarrow C_6H_6ClN + HCl$), chloride concentrations increased proportionally in cultures that exhibited dechlorination (34DCNB-T2, 34DCA-T2, and 23DCA-T2). The measured increase in chloride was slightly higher than expected in almost all cases (Table S11), likely because of the error associated with measuring small changes against a high background of ~1900 μM chloride.

Potential electron-accepting processes include acetogenesis, methanogenesis, nitroreduction, dechlorination, and sulfate reduction. For T2 cultures between days 280 and 520, we calculated the proportion of electron equivalents from added donors channeled to each of the different

electron-accepting processes (Figure 3, bottom table). In DCNB-amended cultures, acetogenesis

from lactate and ethanol was the dominant process, accounting for over 80% of electrons from

donors. Methanogenesis was severely limited. By contrast, in DCA-amended cultures,

methanogenesis accounted for 60-70% of electron equivalents from donors, which combined with

acetogenesis accounted for over 98% of electron equivalents. A buildup of acetate in all cultures

(Tables S10 and S12) suggested possible inhibition of acetoclastic methanogens specifically,

particularly by DCNBs. Culture 34DCNB-T2 produced the most methane among DCNB-amended

cultures but also exhibited the most rapid and extensive nitroreduction and dechlorination, thus

relieving inhibition associated with DCNBs and DCAs.

Other electron accepting processes accounted for a much lower proportion of activity (Figure

3). Nitroreduction consumed 5% of electron equivalents in cultures fed with 3,4- and 2,5-DCNB

isomers and up to 20% of electron equivalents in culture 23DCNB-T2, although activity in this

culture was very low and no lactate was consumed (Tables S10 and S12). In all cases,

dechlorination accounted for less than 3% of all electron equivalents provided. The small amount

of sulfate (~30 µM) present in the medium was rapidly depleted in the 4 highly active cultures

(34DCNB-T2, 25DCNB-T2, 34DCA-T2, and 23DCA-T2). However, less than 1% of electron

equivalents were channeled to sulfate reduction. In culture 23DCNB-T2, sulfate was not

consumed, consistent with low activity observed in this enrichment (Table S11).

We compared the proportions of electrons consumed by each electron-accepting process with

corresponding microbial proportions (Figure 3) to infer function (Table S12). Methane production

was greatest in cultures 23DCA-T2 and 34DCA-T2, lower in 34DCNB-T2, very low in 25DCNB-

T2 and absent in 23DCNB-T2. Methanogens were correspondingly most abundant in 23DCA-T2,

34DCA-T2, and 34DCNB-T2. ASVs corresponding to hydrogenotrophic *Methanoregula*

dominated the archaeal population, with a much lower abundance of *Methanocella* and other

hydrogenotrophic genera (Figure S7). The presence of only hydrogenotrophic methanogens

combined with substantial accumulation of acetate strongly indicates that DCNB isomers and to a

lesser extent DCA isomers inhibited acetoclastic methanogens.

3.4   Nitro group reduction can be mediated by many different microbes

We had hoped to observe enrichment of specific microbes in DCNB-amended cultures to enable

functional attribution. Rather, it seems that many different genera can likely reduce the nitro group.

Biologically regenerated FeS is unlikely to play a role as a reductant, as there was no evidence of

sulfate or iron cycling in the cultures, and the rates of DCNB reduction were similar regardless of

FeS concentrations. In all cultures, there was significant representation of fermenting or acetogenic

microbes, particularly Veillonellales-Selenomonadales, *Sporomusaceae*, *Sporomusa*, and

Anaerospora (Figure S6). This is consistent with the ample supply of ethanol and lactate to all

cultures.

We wondered if these fermenting and acetogenic organisms could also be reducing DCNB to

DCA as a growth-associated electron-accepting process akin to nitrate reduction. In 25DCNB-T2,

where only nitroreduction was observed without dechlorination, 6 ASVs (ASV1 to 5 and 8) (>90%

identity) classified only to the order level as Veillonellales-Selenomonadales were predominant

(Table S5b). They represented more than 80% of the bacterial community in this culture on day

517 (Figure 1e). The closest matches to these sequences are affiliated with the genera *Pelosinus*,

Anaerospora, *Sporomusa*, *Dendrosporobacter*, among other organisms; interestingly, one of the

closely matching sequences (KF460373.1) comes from a study on the impact of nitrate on

pentachlorophenol dechlorination[39] (Figure S8).

In fact, ASVs from Veillonellales-Selenomonadales were abundant in all 5 active cultures

(Figures 1d, e, and f, Figures 2d and e and Figure 3), whether fed DCNB or DCA, making

functional attribution difficult. Finding that DCNB isomers could be reduced in all cultures

regardless of enrichment substrate is perhaps then not so surprizing. In the autoclaved "killed"

control (KC) that was amended with a mixture of compounds (3,4-DCNB, 2,5-DCNB, 3,4-DCA,

2,3-DCA, 2-CA, aniline) both 3,4-DCNB and 2,5-DCNB were reduced to the corresponding DCA

isomers (Figure S5) but we suspected that it was not sterile. DNA was readily extracted from this

control, and although the total microbial abundance was an order of magnitude lower than other

cultures, the KC culture was dominated by a unique Butyricoccaceaea ASV as well as other spore-

forming and presumably heat-resistant microbes. Among the microbes detected in KC after

refeeding 3,4- and 2,5-DCNB 4 times, *Desulfitobacterium* and *Anaerospora* were also detected in

other active nitroreducing cultures (i.e., 23DCNB-T2 and 25DCNB-T2) (Figure S6), suggesting

these genera are also candidates for nitroreduction. Ultimately it became clear that nitroreduction

is a widespread microbial activity that was easy to stimulate with electron donor in the cultures.

3.5   Dechlorination of 2,3- and 3,4-DCA supports growth of *Dehalobacter*

In actively dechlorinating cultures, *Dehalobacter* was the only microbe to systematically show

a substantial increase in abundance. *Dehalobacter* was not detected in cultures that were only

catalyzing nitroreduction. Given that it is an obligate organohalide respiring bacterium, we

conclude that *Dehalobacter* was responsible for the DCA-dechlorinating activity in enrichment

cultures amended with 3,4-DCNB, 3,4-DCA, and 2,3-DCA (Figures 1d and 2d and e).

Two distinct *Dehalobacter* ASVs were identified: ASV1 in 34DCNB-T2 and ASV2 in 34DCA-

T2 and 23DCA-T2 (Text S6), with maximum bacterial abundances of 14, 25, and 31% in each

culture, respectively (Table S3b). ASV1 (468 bp) is 100% identical to several previously published

16S rRNA gene sequences of *Dehalobacter restrictus* strains, including strain SAD (CP148032.1),

strain DSM 9455 (CP007033.1), and strain 12DCA (CP046996.1), as well as to three

Dehalobacter strains classified only to the genus level, including strain DCA (CP003869.1), strain

DAD (CP148031.1), and strain CF (1CP003870.1). The known substrates for these *Dehalobacter*

strains include tetra- and trichloroethene,[40] 1,2-dichloroethane,[41] tri- and dichloromethane,[42] and

1,1,1-trichloroethane and 1,1-dichloroethane.[43,44]

Dehalobacter ASV2 (468 bp) was not a perfect match to any 16S rRNA gene sequence in the

NCBI database. It shares only 97.01% identity to ASV1 and 98.93% to *Dehalobacter* strain DCM

(CP092282.1) described as "dichloromethane-degrading bacterium from Xi river" (Direct

Submission by Yang, Y. *et al*., 13-Jan-2022). Neither *Dehalobacter* ASV1 nor ASV2 sequences

matched (<95% identity) to the 16S rRNA gene fragment (937 bp) from *Dehalobacter* sp. strain

14DCB1 (JN051267.1), a strain enriched on dichlorobenzene[45] but reported to dechlorinate

dichloroaniline.[14]

To further confirm the role of *Dehalobacter* in chloroaniline dechlorination, we estimated

growth yields from qPCR and concentration data to compare with previously published yields.

The net growth yields ranged from 1.4×10^{13} to 6.5×10^{13} cells per mol of chloride released

assuming 3 copies of the 16S rRNA gene per cell (Table S13). The yields calculated in this study

are consistent with many previously reported yields for other *Dehalobacter* strains using various

chloroethanes, chlorobenzenes, and chloromethanes as electron acceptors for growth (Table S14).

In fact the yields align with the highest yields previously reported, solidifying chloroaniline

dechlorination as a growth-related process.

Additional evidence for *Dehalobacter* being responsible for DCA dichlorination was obtained

when non-dechlorinating culture 25DCNB-T2 was inoculated with 3 mL of dechlorinating culture

34DCNB-T2 (on day 653). After inoculation, residual 2,5-DCA previously formed from 2,5-

DCNB was dechlorinated to 3-CA and the relative abundance of *Dehalobacter*_ASV1 increased

from zero on day 517 to ~9% on day 653 (Figure S9).

A shift in activity in culture 23DCA-T2 was noted as monitoring continued for over 1000 days

(Figure S10). As time progressed, 2-CA rather than 3-CA became the major product from 2,3-

DCA. Thereafter, 2-CA was slowly yet consistently transformed to aniline. On day 881,

*Dehalobacter*_ASV2 constituted 16% of the bacterial community (Table S5e). 2-CA was

dechlorinated to aniline in subsequent transfers from this culture as well (data not shown). A

subsample of 23DCA-T2 was taken on day 653 for additional growth experiments to ultimately

build up enough culture to measure compound specific isotopic enrichment factors. This ongoing

research implicates *Dehalobacter* in aniline formation as described in a companion publication.[46]

3.7   Other organisms identified in enriched cultures

Several other ASVs belonging to known facultative OHRB were identified, including

Anaeromyxobacter, *Desulfovibrio*, and *Desulfitobacterium* (Figure 3). While some members of

these genera can dechlorinate, most are known to be quite versatile and may reduce sulfate, nitrate

or even grow fermentatively in the absence of a terminal electron acceptor. An *Anaeromyxobacter*

ASV found in culture 34DCNB-T2 is 99.79% similar over ~450 bp to the 16S rRNA gene of

Anaeromyxobacter sp. strain RF71 (MZ254878.1), 99.57% similar to *A. dehalogenans* 2CP-C

(CP000251.1) and 99.36% similar to *A. dehalogenans* 2CP-1 (CP001359.1). Both strains 2CP-C

and 2CP-1 can gain energy from the reductive dechlorination of chlorophenol and contain putative

dehalogenases.[47–49] However, the abundance of *Anaeromyxobacter* in the culture is far less than

Dehalobacter and is therefore unlikely to be primarily responsible for dechlorination.

A single ASV of *Desulfovibrio* increased in abundance over time in the three cultures

dechlorinating DCA isomers (Figure 3). *Desulfovibrio* are sulfate-reducing bacteria that can also

ferment substrates such as lactate; some can respire organohalides.[50,51] In cultures 23DCA-T2 and

34DCA-T2, *Desulfovibrio* increased between days 280 and 520 to similar extent as *Dehalobacter*

(Figures 2d and 2e). However, a similar increase was not seen in dechlorinating culture 34DCNB-

T2 (Figure 1d). The amount of sulfate in the medium (around 30 μM) is too small to explain this

growth, therefore *Desulfovibrio* is more likely metabolizing lactate to acetate, possibly

participating in dechlorination alongside *Dehalobacter*. The *Desulfovibrio* ASV in our cultures is

100% identical to a non-dechlorinating *Desulfovibrio* sp. (JQ599691.1) detected in a

dichloroethane dechlorinating enrichment.[52] The sequence was also 100% identical to an

uncultured *Desulfovibrio* clone (AB854345.1) found in a non-dechlorinating environment.[53]

Therefore, the more likely role of this organism is fermentation and acetogenesis, in support of

dechlorinating *Dehalobacter*.

Several ASVs (sharing > 94% identity) were assigned to *Desulfitobacterium*. Interestingly, these

ASVs were only found in the cultures reducing DCNBs and in the KC control that exhibited

nitroreduction; they were not detected in the DCA-amended cultures (Figure 3). This seems to

implicate *Desulfitobacterium* in nitroreduction, however the ASVs were in low abundance.

Furthermore, DCA-amended cultures could also rapidly reduce the nitro group, therefore

Desulfitobacterium is not solely responsible for nitroreduction, if at all. Moreover, the 8 most

abundant *Desulfitobacterium* ASV sequences are > 95% similar to a *Desulfitobacterium* sequence

found in *Dehalococcoide*s-containing dechlorinating cultures (AB596883.1) (Figure S11), where

they were not implicated in dechlorination,[54] a likely scenario in our study as well.

3.8   Data integration and application to improving remediation outcomes

Non-metric multidimensional scaling (NMDS) was used to evaluate relationships among

cultures as they were enriched from subsurface material to initial microcosms and finally to

enrichment cultures (Figure 4). First, there is clearly an impact of the original source material,

shown in red, blue and green for three original sediment sources used to construct the microcosms.

Second, there was a convergence of community depending on substrate. Enrichment cultures that

dechlorinate DCAs are distinct from those that only carry out nitroreduction. Samples from

34DCNB-T2 culture, in which both nitroreduction and dehalogenation are observed, are located

between the dechlorinating-exclusive culture samples (34DCA-T2, and 23DCA-T2) and the

nitroreducing-exclusive cultures (25DCNB-T2, 23DCNB-T2 and KC). *Dehalobacter* in the top

groupings and *Desulfitobacterium* in the bottom two cultures (23DCNB-T2 and KC) are major

genera driving the vertical spread of samples in the NMDS plot.

Figure 4. Non-metric multidimensional scaling plot based on relative abundance of 16S rRNA amplicons in 45 samples from the original site material used for microcosms, microcosms and enrichment cultures.

In a survey of over 60 groundwater samples from around the site in Brazil that were analyzed by amplicon sequencing, approximately 14% returned *Dehalobacter*. Of the ~400 samples that have been sequenced from various remediation trials – including samples from pilot wetlands and pre and post biosparging units – many different *Dehalobacter* ASVs (including those enriched in this study) were detected in 78 samples (~20%) clustered around more impacted zones with lower redox potential. Relative abundances of *Dehalobacter* reached 2% of the bacterial population in some locations, which is high considering the complex mixture of

chemicals and conditions at the site.[1,2] These abundances are being used to delineate biologically

active zones and to predict attenuation rates to most effectively remediate the site.

Other ASVs detected in the enrichment cultures are detected in site samples, including the

ones affiliated with *Anaeromyxobacter* and *Desulfitobacterium*, which were detected in ~53%

and ~18% of the total ~460 samples, respectively. The ASVs classified as Veillonellales-

Selenomonadales were detected in ~11% of the same group of samples, while the ones classified

further to *Sporomusa* and *Anaerospora* genera in ~3% and ~6%. The high relative abundance of

the Veillonellales-Selenomonadales order in our enriched cultures indicates those organisms

were favoured by the ample availability of donor (ethanol and lactate) added to the cultures,

which is not the case in the field.

The fact that the microbes identified in enrichment cultures are also present at the site attests to

the utility of such enrichments in site characterization and provides very strong evidence that these

processes are contributing to active natural attenuation in anaerobic zones. At the site,

concentrations of dichloroanilines are generally higher than dichloronitrobenzenes, consistent with

our finding that biologically-mediated nitroreduction occurs readily and that there are many genera

capable of catalyzing this reaction. It is highly likely that some of the other contaminants at the

site could be electron donors, including petroleum hydrocarbons, solvents and plasticizers. In

addition, abiotic reactions may also occur driven by reduced minerals or by reduced natural organic

matter, which can donate electrons for nitroreduction.[37,55–58] Further studies will seek to better

understand the players and nitroreduction mechanisms in DCNB cultures.

Two distinct *Dehalobacter* ASVs were unequivocally linked to dechlorination, and their

presence along with other ASVs in site samples is indicative of dechlorinating activity in situ. The

two ASVs originated from geographically distinct locations at the site. Traces of aniline were

found and continue to accumulate in 23DCA-T2, suggesting further dechlorination of CA is

possible in situ. Inhibition of acetoclastic methanogens was noted, which is also consistent with

site data and provides a means to evaluate site recovery: as DCNB and DCA concentrations

decrease with active remediation efforts, methanogenic populations in anoxic zones should

rebound. Biomarkers based on 16S rRNA genes are already being deployed at the site to track

population changes as remediation alternatives are evaluated. The robust enrichment cultures

generated herein will be invaluable for future studies of biotransformation mechanisms,

development of isotopic enrichment factors and of additional biomarkers, and possibly even for

use as bioaugmentation cultures.

SUPPORTING INFORMATION

The following files are available free of charge:

Additional details on experiments, materials and methods, and data obtained, including texts and

figures mentioned in this paper (PDF); Tables including data obtained and calculations mentioned

in this paper (.xls).

ACKNOWLEDGEMENTS

For the financial support for Sofia Araújo, we thank FACEPE (Science and Technology

Foundation of the State of Pernambuco, Brazil – process number BFD-0013-3.07/21), CNPq

(National Council for Scientific and Technological Development, Brazil - process number

165130/2017-2), and CAPES (Coordination for the Improvement of Higher Education Personnel)

for the Graduate Program support (Proap) and Institutional Internationalization Program (PrInt)

(process number 88887.363316/2019-00). This work was also supported by Mitacs through the

Mitacs Accelerate and Mitacs Elevate programs (Application ref. IT23315 and IT34875, Canada).

The authors also acknowledge the grant from CNPq (process number 304862/2018-5) and

FACEPE (process number APQ 0456-3.07/20), and NSERC (Natural Sciences and Engineering

Research Council of Canada) Collaborative Research and Development (CRD) project, which has

Geosyntec Consultants and Corteva Agriscience™ as the two main industrial partners.

The authors do not have a conflict of interest to declare. This publication is in accordance with

the Brazilian biodiversity law n.º 13.123, from May 20th, 2015. This is non-profit research and has

no commercial intention between the parties. This is a collaboration project between the parties

involved. Access Registration No. A3D9F11.

We would also like to thank Elodie Passeport, Shuping Wang, and Luiz Pereira for their helpful

discussions and collaborative work on this project.

REFERENCES

(1)   Melo, N.; Araújo, S. P.; de Paula Queiroz Kraus, S.; Lomheim, L.; Quintero, P. B.; Mack, E. E.; Edwards, E. A.; Spain, J.; Gavazza, S. Strategies for Bioremediation of Soil from an Industrial Site Exposed to Chlorinated and Nitroaromatic Compounds. *Groundwater Monit. Rem.* **2023**, *43* (3), 108–120. https://doi.org/10.1111/gwmr.12596.

(2)   Barreto, P.; Lemes, M.; Jimenez, J.; Mack, E. E.; Henderson, J.; Freedman, D. L. Evaluation of Strategies to Remediate Mixed Wastes at an Industrial Site in Brazil. *Groundwater Monit. Rem.* **2023**, *43* (3), 93–107. https://doi.org/10.1111/gwmr.12607.

(3)   Buser, H.-R.; Bosshardt, H.-P. Studies on the Possible Formation of Polychloroazobenzenes in Quintozene Treated Soil. *Pestic Sci* **1975**, *6*, 35–41.

(4)   Roldán, M. D.; Pérez-Reinado, E.; Castillo, F.; Moreno-Vivián, C. Reduction of Polynitroaromatic Compounds: The Bacterial Nitroreductases. *FEMS Microbiol. Rev.* **2008**, pp 474–500. https://doi.org/10.1111/j.1574-6976.2008.00107.x.

(5)   Bunce, N. J.; Merrick, R. L.; Corke, C. T. Reductive Transformations of Nitrate with 3,4-Dichloroaniline and Related Compounds by *Escherichia Coli*. *J Agric Food Chem* **1983**, *31* (5), 299–326.

(6)   Spain, J. C. Biodegradation of Nitroaromatic Compounds. *Annu Rev Microbiol* **1995**, *49*, 523–555.

(7)   Esteve-Nunez, A.; Lucchesi, G.; Philipp, B.; Schink, B.; Ramos, J. L. Respiration of 2,4,6-Trinitrotoluene by *Pseudomonas* sp. strain JLR11. *J Bacteriol* **2000**, *182* (5), 1352–1355.

(8)   Huang, J.; Ning, G.; Li, F.; Sheng, G. D. Biotransformation of 2,4-Dinitrotoluene by Obligate Marine *Shewanella marisflavi* EP1 under Anaerobic Conditions. *Bioresour Technol* **2015**, *180*, 200–206. https://doi.org/10.1016/j.biortech.2014.12.108.

(9)   Madeira, C. L.; Menezes, O.; Park, D.; Jog, K. V.; Hatt, J. K.; Gavazza, S.; Krzmarzick, M. J.; Sierra-Alvarez, R.; Spain, J. C.; Konstantinidis, K. T.; Field, J. A. Bacteria Make a Living Breathing the Nitroheterocyclic Insensitive Munitions Compound 3-Nitro-1,2,4-Triazol-5-One (NTO). *Environ Sci Technol* **2021**, *55* (9), 5806–5814. https://doi.org/10.1021/acs.est.0c07161.

(10)   Struijs, J.; Rogers, J. E. Reductive Dehalogenation of Dichloroanilines by Anaerobic Microorganisms in Fresh and Dichlorophenol-Acclimated Pond Sediment. *Appl Environ Microbiol* **1989**, *55* (10), 2527–2531.

(11) Kuhn, E. P.; Suflita, J. M. Sequential Reductive Dehalogenation of Chloroanilines by Microorganisms from a Methanogenic Aquifer. *Environ Sci Technol* **1989**, *23* (7), 848–852. https://doi.org/10.1021/es00065a014.

(12) Kuhn, E. P.; Townsend, G. T.; Suflita, J. M. Effect of Sulfate and Organic Carbon Supplements on Reductive Dehalogenation of Chloroanilines in Anaerobic Aquifer Slurries. *Appl Environ Microbiol* **1990**, *56* (9), 2630–2637. https://doi.org/10.1128/aem.56.9.2630-2637.1990.

(13) Travkin, V.; Baskunov, B. P.; Golovlev, E. L.; Boersma, M. G.; Boeren, S.; Vervoort, J.; Berkel, W. J. H.; Rietjens, I. M. C. M.; Golovleva, L. A. Reductive Deamination as a New Step in the Anaerobic Microbial Degradation of Halogenated Anilines. *FEMS Microbiol Lett* **2002**, *209* (2), 307–312. https://doi.org/10.1111/j.1574-6968.2002.tb11149.x..

(14) Zhang, S.; Wondrousch, D.; Cooper, M.; Zinder, S. H.; Schüürmann, G.; Adrian, L. Anaerobic Dehalogenation of Chloroanilines by *Dehalococcoides* m*ccartyi* strain CBDB1 and *Dehalobacter* strain 14DCB1 via Different Pathways as Related to Molecular Electronic Structure. *Environ Sci Technol* **2017**, *51* (7), 3714–3724. https://doi.org/10.1021/acs.est.6b05730.

(15) Susarla, S.; Yonezawa, Y.; Masunaga, S. Reductive Dehalogenation of Chloroanilines in Anaerobic Estuarine Sediment. *Environ. Technol.* **1997**, *18* (1), 75–83. https://doi.org/10.1080/09593331808616514.

(16) Susarla, S.; Yonezawa, Y.; Masunaga, S. Reductive Transformations of Halogenated Aromatics in Anaerobic Estuarine Sediment: Kinetics, Products and Pathways. *Water Res* **1998**, *32* (3), 369–348.

(17) Duc, H. D.; Oanh, N. T. Anaerobic Degradation of Chloroanilines by *Geobacter* sp. KT5. *Curr Microbiol* **2019**, *76* (2), 248–257. https://doi.org/10.1007/s00284-018-1617-7.

(18) Oanh, N. T.; Duc, H. D. Anaerobic Degradation of Propanil in Soil and Sediment Using Mixed Bacterial Culture. *Curr Microbiol* **2021**, *78* (4), 1499–1508. https://doi.org/10.1007/s00284-021-02419-7.

(19) Lu, H.; Zhang, T.; Zhou, Y.; Zhou, J.; Wang, J.; Wang, X. Enhanced Dechlorination and Biodegradation of 2-Chloroaniline by a 2-Aminoanthraquinone-Graphene Oxide Composite under Anaerobic Conditions. *Sci Rep* **2019**, *9* (1). https://doi.org/10.1038/s41598-019-48904-9.

(20) Kraus, S. Aerobic and Anaerobic Biotransformation of Chloroanilines, Chlorobenzenes, and Dichlonitrobenzenes at a Complex Industrial Site in Brazil and Analysis of Associated Microbial Communities. Master Thesis, University of Toronto, Toronto, 2018. https://hdl.handle.net/1807/91484 (accessed 2024-06-11).

(21) Edwards, E. A.; Grbic-Galic, D. Anaerobic Degradation of Toluene and O-Xylene by a Methanogenic Consortium. *Appl Environ Microbiol* **1994**, 313–322.

(22) Toth, C. R. A.; Luo, F.; Bawa, N.; Webb, J.; Guo, S.; Dworatzek, S.; Edwards, E. A. Anaerobic Benzene Biodegradation Linked to the Growth of Highly Specific Bacterial Clades. *Environ Sci Technol* **2021**, *55* (12), 7970–7980. https://doi.org/10.1021/acs.est.1c00508.

(23) Engelbrektson, A.; Kunin, V.; Wrighton, K. C.; Zvenigorodsky, N.; Chen, F.; Ochman, H.; Hugenholtz, P. Experimental Factors Affecting PCR-Based Estimates of Microbial Species Richness and Evenness. *ISME Journal* **2010**, *4* (5), 642–647. https://doi.org/10.1038/ismej.2009.153.

(24) Ferris, M. J.; Muyzer, G.; Ward, D. M. Denaturing Gradient Gel Electrophoresis Profiles of 16S rRNA-Defined Populations Inhabiting a Hot Spring Microbial Mat Community. *Appl Environ Microbiol* **1996**, *62* (2), 340–346.

(25) Bolyen, E.; Rideout, J. R.; Dillon, M. R. Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2. *Nat. Biotechnol*, **2019**, pp 852–857. https://doi.org/10.1038/s41587-019-0190-3.

(26) Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S. P. DADA2: High-Resolution Sample Inference from Illumina Amplicon Data. *Nat Methods* **2016**, *13* (7), 581–583. https://doi.org/10.1038/nmeth.3869.

(27) Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F. O. The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing and Web-Based Tools. *Nucleic Acids Res* **2013**, *41* (D1). https://doi.org/10.1093/nar/gks1219.

(28) Ritalahti, K. M.; Amos, B. K.; Sung, Y.; Wu, Q.; Koenigsberg, S. S.; Löffler, F. E. Quantitative PCR Targeting 16S rRNA and Reductive Dehalogenase Genes Simultaneously Monitors Multiple *Dehalococcoides* strains. *Appl Environ Microbiol* **2006**, *72* (4), 2765–2774. https://doi.org/10.1128/AEM.72.4.2765-2774.2006.

(29) Lane, D. J. 16S/23S rRNA Sequencing. In *Nucleic acid techniques in bacterial systematics*; Stackebrandt, E., Goodfellow, M., Eds.; John Wiley and Sons: Chichester, United Kingdom, 1991; pp 115–175.

(30) Yu, Y.; Lee, C.; Kim, J.; Hwang, S. Group-Specific Primer and Probe Sets to Detect Methanogenic Communities Using Quantitative Real-Time Polymerase Chain Reaction. *Biotechnol Bioeng* **2005**, *89* (6), 670–679. https://doi.org/10.1002/bit.20347.

(31) Cheng, J.; Suidan, M. T.; Venosa, A. D. Abiotic Reduction of 2,4-Dinitrotoluene in the Presence of Sulfide Minerals under Anoxic Conditions. *Wat. Sci. Tech.* **1996**, *34* (10), 25–33. https://doi.org/10.1016/S0273-1223(96)00693-2.

(32) Elsner, M.; Schwarzenbach, R. P.; Haderlein, S. B. Reactivity of Fe(II)-Bearing Minerals toward Reductive Transformation of Organic Contaminants. *Environ Sci Technol* **2004**, *38* (3), 799–807. https://doi.org/10.1021/es0345569.

(33) Klupinski, T. P. Abiotic Degradation of Pentachloronitrobenzene by Fe (II): Reactions on Goethite and Iron Oxide Nanoparticles. *Environ. Sci. Technol* **2004**, *38*, 4353–4360. https://doi.org/10.1021/es035434j.

(34) Niedźwiecka, J. B.; Drew, S. R.; Schlautman, M. A.; Millerick, K. A.; Grubbs, E.; Tharayil, N.; Finneran, K. T. Iron and Electron Shuttle Mediated (Bio)Degradation of 2,4-Dinitroanisole (DNAN). *Environ Sci Technol* **2017**, *51* (18), 10729–10735. https://doi.org/10.1021/acs.est.7b02433.

(35) Khatiwada, R.; Root, R. A.; Abrell, L.; Sierra-Alvarez, R.; Field, J. A.; Chorover, J. Abiotic Reduction of Insensitive Munition Compounds by Sulfate Green Rust. *Environ. Chem.* **2018**, *15* (5), 259–266. https://doi.org/10.1071/EN17221.

(36) Cárdenas-Hernández, P. A.; Anderson, K. A.; Murillo-Gelvez, J.; Di Toro, D. M.; Allen, H. E.; Carbonaro, R. F.; Chiu, P. C. Reduction of 3-Nitro-1,2,4-Triazol-5-One (NTO) by the Hematite-Aqueous Fe(II) Redox Couple. *Environ Sci Technol* **2020**, *54* (19), 12191–12201. https://doi.org/10.1021/acs.est.0c03872.

(37) Niedźwiecka, J. B.; McGee, K.; Finneran, K. T. Combined Biotic-Abiotic 2,4-Dinitroanisole Degradation in the Presence of Hexahydro-1,3,5-Trinitro-1,3,5-Triazine. *Environ Sci Technol* **2020**, *54* (17), 10638–10645. https://doi.org/10.1021/acs.est.0c02363.

(38) Menezes, O.; Yu, Y.; Root, R. A.; Gavazza, S.; Chorover, J.; Sierra-Alvarez, R.; Field, J. A. Iron(II) Monosulfide (FeS) Minerals Reductively Transform the Insensitive Munitions Compounds 2,4-Dinitroanisole (DNAN) and 3-Nitro-1,2,4-Triazol-5-One (NTO). *Chemosphere* **2021**, *285*. https://doi.org/10.1016/j.chemosphere.2021.131409.

(39) Yu, H. Y.; Wang, Y. kui; Chen, P. cheng; Li, F. bai; Chen, M. jia; Hu, M.; Ouyang, X. Effect of Nitrate Addition on Reductive Transformation of Pentachlorophenol in Paddy Soil in Relation to Iron(III) Reduction. *J Environ Manage* **2014**, *132*, 42–48. https://doi.org/10.1016/j.jenvman.2013.10.020.

(40) Kruse, T.; Maillard, J.; Goodwin, L.; Woyke, T.; Teshima, H.; Bruce, D.; Detter, C.; Tapia, R.; Han, C.; Huntemann, M.; Wei, C. L.; Han, J.; Chen, A.; Kyrpides, N.; Szeto, E.; Markowitz, V.; Ivanova, N.; Pagani, I.; Pati, A.; Pitluck, S.; Nolan, M.; Holliger, C.; Smidt, H. Complete Genome Sequence of *Dehalobacter restrictus* PER-K23T. *Stand Genomic Sci* **2013**, *8* (3). https://doi.org/10.4056/sigs.3787426.

(41) Molenda, O.; Puentes Jácome, L. A.; Cao, X.; Nesbø, C. L.; Tang, S.; Morson, N.; Patron, J.; Lomheim, L.; Wishart, D. S.; Edwards, E. A. Insights into Origins and Function of the Unexplored Majority of the Reductive Dehalogenase Gene Family as a Result of Genome Assembly and Ortholog Group Classification. *Environ Sci Process Impacts* **2020**, *22* (3), 663–678. https://doi.org/10.1039/c9em00605b.

(42) Bulka, O.; Webb, J.; Dworatzek, S.; Mahadevan, R.; Edwards, E. A. A Multifunctional *Dehalobacter*? Tandem Chloroform and Dichloromethane Degradation in a Mixed Microbial Culture. *Environ Sci Technol* **2023**, *57* (48), 19912–19920. https://doi.org/10.1021/acs.est.3c06686.

(43) Tang, S.; Gong, Y.; Edwards, E. A. Semi-Automatic in Silico Gap Closure Enabled de Novo Assembly of Two *Dehalobacter* Genomes from Metagenomic Data. *PLoS One* **2012**, *7* (12). https://doi.org/10.1371/journal.pone.0052038.

(44) Tang, S.; Wang, P. H.; Higgins, S. A.; Löffler, F. E.; Edwards, E. A. Sister *Dehalobacter* Genomes Reveal Specialization in Organohalide Respiration and Recent Strain Differentiation Likely Driven by Chlorinated Substrates. *Front Microbiol* **2016**, *7* (FEB). https://doi.org/10.3389/fmicb.2016.00100.

(45) Nelson, J. L.; Fung, J. M.; Cadillo-Quiroz, H.; Cheng, X.; Zinder, S. H. A Role for *Dehalobacter* spp. in the Reductive Dehalogenation of Dichlorobenzenes and Monochlorobenzene. *Environ Sci Technol* **2011**, *45* (16), 6806–6813. https://doi.org/10.1021/es200480k.

(46) Wang, S.; Araujo, S. P.; Lomheim, L.; Mack, E. E.; Edwards, E. A.; Passport, E. *Dehalobacter converts 2,3-dichloroaniline to aniline via 2-chloroaniline in a methanogenic enrichment culture*; 2025. Manuscript in preparation (preprint on *BioRxiv*).

(47) Cole, J. R.; Cascarelli, A. L.; Mohn, W. W.; Tiedjel, J. M. Isolation and Characterization of a Novel Bacterium Growing via Reductive Dehalogenation of 2-Chlorophenol. *Appl Environ Microbiol* **1994**, *60* (10), 3536–3542.

(48) Sanford, R. A.; Cole, J. R.; Tiedje, J. M. Characterization and Description of *Anaeromyxobacter dehalogenans* gen. nov., sp. nov., an Aryl-Halorespiring Facultative Anaerobic Myxobacterium. *Appl Environ Microbiol* **2002**, *68* (2), 893–900. https://doi.org/10.1128/AEM.68.2.893-900.2002.

(49) Thomas, S. H.; Wagner, R. D.; Arakaki, A. K.; Skolnick, J.; Kirby, J. R.; Shimkets, L. J.; Sanford, R. A.; Löffler, F. E. The Mosaic Genome of *Anaeromyxobacter dehalogenans* strain 2CP-C Suggests an Aerobic Common Ancestor to the Delta-Proteobacteria. *PLoS One* **2008**, *3* (5). https://doi.org/10.1371/journal.pone.0002103.

(50) Fincker, M.; Spormann, A. M. Biochemistry of Catabolic Reductive Dehalogenation. *Annu Rev Biochem* **2017**, *86*, 357–386. https://doi.org/10.1146/annurev-biochem.

(51) Qian, Y.; Xu, M.; Deng, T.; Hu, W.; He, Z.; Yang, X.; Wang, B.; Song, D.; Chen, L.; Huang, Y.; Sun, G. Synergistic Interactions of *Desulfovibrio* and *Petrimonas* for Sulfate-Reduction Coupling Polycyclic Aromatic Hydrocarbon Degradation. *J Hazard Mater* **2021**, *407*. https://doi.org/10.1016/j.jhazmat.2020.124385.

(52) Manchester, M. J.; Hug, L. A.; Zarek, M.; Zila, A.; Edwards, E. A. Discovery of a Trans-Dichloroethene-Respiring *Dehalogenimonas* Species in the 1,1,2,2-Tetrachloroethane-

Dechlorinating WBC-2 Consortium. *Appl Environ Microbiol* **2012**, *78* (15), 5280–5287. https://doi.org/10.1128/AEM.00384-12.

(53) Kato, S.; Yumoto, I.; Kamagata, Y. Isolation of Acetogenic Bacteria That Induce Biocorrosion by Utilizing Metallic Iron as the Sole Electron Donor. *Appl Environ Microbiol* **2015**, *81* (1), 67–73. https://doi.org/10.1128/AEM.02767-14.

(54) Ise, K.; Suto, K.; Inoue, C. Microbial Diversity and Changes in the Distribution of Dehalogenase Genes during Dechlorination with Different Concentrations of Cis -DCE. *Environ Sci Technol* **2011**, *45* (12), 5339–5345. https://doi.org/10.1021/es104199y.

(55) Menezes, O.; Kocaman, K.; Wong, S.; Rios-Valenciana, E. E.; Baker, E. J.; Hatt, J. K.; Zhao, J.; Madeira, C. L.; Krzmarzick, M. J.; Spain, J. C.; Sierra-Alvarez, R.; Konstantinidis, K. T.; Field, J. A. Quinone Moieties Link the Microbial Respiration of Natural Organic Matter to the Chemical Reduction of Diverse Nitroaromatic Compounds. *Environ Sci Technol* **2022**, *56* (13), 9387–9397. https://doi.org/10.1021/acs.est.2c01329.

(56) Murillo-Gelvez, J.; Di Toro, D. M.; Allen, H. E.; Carbonaro, R. F.; Chiu, P. C. Reductive Transformation of 3-Nitro-1,2,4-Triazol-5-One (NTO) by Leonardite Humic Acid and Anthraquinone-2,6-Disulfonate (AQDS). *Environ Sci Technol* **2021**, *55* (19), 12973–12983. https://doi.org/10.1021/acs.est.1c03333.

(57) Ratasuk, N.; Nanny, M. A. Characterization and Quantification of Reversible Redox Sites in Humic Substances. *Environ Sci Technol* **2007**, *41* (22), 7844–7850. https://doi.org/10.1021/es071389u.

(58) Uchimiya, M.; Stone, A. T. Reversible Redox Chemistry of Quinones: Impact on Biogeochemical Cycles. *Chemosphere* **2009**, *77* (4), 451–458. https://doi.org/10.1016/j.chemosphere.2009.07.025.