Dehalobacter dechlorinates dichloroanilines and

contributes to the natural attenuation of

dichloronitrobenzenes at a complex industrial site

Sofia P. Araújo,[a] Line Lomheim,[a] Suzana P. Q. Kraus,[b] E. Erin Mack,[c] Jim C. Spain,[d] Sávia

Gavazza,[e] Elizabeth A. Edwards[a]*

[a]Department of Chemical Engineering and Applied Chemistry, University of Toronto,

Toronto, Ontario, M5S 3E5, Canada.

[b]Geosyntec Consultants International Inc, Etobicoke, Ontario, M8X 1Y9, Canada

[c]Corteva Remediation Group, Wilmington, Delaware, 19805, United States

[d]Center for Environmental Diagnostics and Bioremediation, University of West Florida,

Pensacola, Florida, 32514-5751, United States

[e]Departamento de Engenharia Civil e Ambiental, Universidade Federal de Pernambuco,

Recife, Pernambuco, 50740-530, Brazil

*Corresponding author: elizabeth.edwards@utoronto.ca

ABSTRACT

The potential for bioremediation of dichloroanilines (2,3- and 3,4-DCA) and

dichloronitrobenzenes (3,4-, 2,5- and 2,3-DCNB) was investigated using inocula from an

industrial site in northeast of Brazil. Anaerobic biotransformation of these chlorinated

compounds was observed in microcosms simulating site conditions, particularly when electron

donor was added. To disentangle specific transformation reactions, sub-cultures from active

anaerobic microcosms were enriched with individual DCA or DCNB isomers. In these single-

compound enrichment cultures, DCNB isomers were reduced to the corresponding DCA. Each

DCA isomer was stoichiometrically dechlorinated to the corresponding monochloroaniline

(CA) isomers. Further dechlorination of CA to aniline was noted. The reduction of DCNB to

DCA was sustained by known fermenters and possibly *Desulfitobacterium*, while

Dehalobacter was clearly responsible for dechlorination of DCAs. These findings helped to

attribute function to microbial community data retrieved from field samples to inform the

conceptual site model and enable identification of bioactive zones and rate-limiting conditions

at a complex site. More generally, these data broaden the understanding of the metabolic

repertoire of *Dehalobacter* and provide a model for incorporating microbial community data

into site remediation efforts.

Graphical Abstract

40 Keywords: Dichloronitrobenzene; Dichloroaniline; Monochloroaniline; anaerobic

41 biotransformation; bioremediation.

Short synopsis statement

43 A diverse microbial community contributes to natural attenuation at a complex

44 contaminated site. Notably, dichloroanilines produced by biologically-mediated

45 nitroreduction of dichloronitrobenzenes were dechlorinated by *Dehalobacter*.

1 INTRODUCTION

Many historically-contaminated sites worldwide are a challenge to remediate owing to complex mixtures of toxic contaminants and heterogeneous stratigraphy. For example, an industrial facility located in Bahia in NE Brazil has been the subject of several remedial investigations.[1,2] As a former manufacturing facility of the herbicide diuron, it is impacted by high concentrations of multiple compounds including dichloronitrobenzenes (DCNB; ~3,000 mg/kg) and dichloroanilines (DCA; ~700 mg/kg) among many others.[1,2] Anoxic conditions typically prevail in regions with low permeability, high organic matter, or high contaminant concentrations, where the rate of oxygen depletion exceeds rates of replenishment by diffusion or groundwater flow. Therefore, we sought to understand the fate of DCNB and DCA isomers under anoxic conditions with particular attention to associated microbial populations to elucidate microbial pathways and thus create a better conceptual model of the site and a more effective remediation strategy.

In subsurface anoxic environments, nitroaromatic compounds tend to become reduced to their corresponding aromatic amines. As early as 1975, the pesticide pentachloronitrobenzene was shown to be reduced to pentachloroaniline.[3] Nitro group reduction to the corresponding amine is now understood to involve broad specificity and oxygen-independent nitroreductases and related enzymes that are widely distributed in bacteria and archaea.[4] These reactions produce toxic transient nitroso and hydroxylamino intermediates, which are highly reactive and unstable.[4–6] Nitroaromatic reduction may also be a respiratory process akin to nitrate

reduction.[7–9] A 1983 study reported the reduction of 3,4-DCNB to a variety of poorly quantified

products including 3,4-DCA, and condensation products.[5] Despite their environmental

relevance, few studies have investigated the anaerobic biodegradation of DCNB isomers.

Literature on the anaerobic biodegradation of chloroanilines is also limited. The reductive

dechlorination of 3,4-DCA to a mixture of monochloroaniline isomers (3-CA and 4-CA) under

anaerobic/methanogenic conditions was first reported in 1989 and 1990, but the organisms

responsible were not identified.[10–12] Under nitrate-reducing conditions, a novel reductive

deamination reaction was observed in a *Rhodococcus* strain where 3,4-DCA was deaminated

primarily to dichlorobenzene,[13] however, additional reports of this process could not be found.

More recently, *Dehalococcoides mccartyi* strain CBDB1 and *Dehalobacter sp.* strain 14DCB1,

both obligate organohalide-respiring-bacteria (OHRB) that dechlorinate chlorinated

benzenes, were shown to dechlorinate 2,3-DCA to 2- and 3-CA and 3,4-DCA to 3-CA with no

further transformation when acetate was provided as carbon source and hydrogen as electron

donor.[14]

Reports of further dechlorination of monochloroanilines to aniline are sparse and

inconclusive. Early reports (1997-98) described dechlorination of DCA to CA and then to small

amounts of aniline under sulfate-reducing conditions.[15,16] These studies did not characterize

the microbial community. A 2019 study showed that *Geobacter* sp. KT5 could transform

dichloroanilines under iron-reducing conditions; mineralization of the products via aniline

and 4-aminobenzoate was proposed.[17] Two years later, a mixed culture composed of *Geobacter*

sp., *Paracoccus denitrificans*, *Pseudomonas* sp., and *Rhodococcus* sp. strains was described to

transform propanil to 3,4-DCA, and further to 3-CA, chlorobenzene, and aniline, prior to

mineralization.[18] Another study found that 2-aminoanthraquinone-graphene oxide enhanced

dechlorination in an enriched anaerobic sludge community that was able to dechlorinate 2-

CA. In this work, some aniline and hexanoic acid were detected and organisms from

Oscillospira, *Lactobacillales*, *Veillonellaceae*, and *Ruminococcus* groups were positively

correlated with 2-CA transformation.[19]

Given the scarce literature on the biodegradation of dichloronitrobenzenes and

dichloroanilines, the goal of this study was to identify organisms responsible for

transformation reactions of such compounds and to use the information to design biomarkers

to track biodegradation and delineate bioactive zones across a contaminated site. We

determined that *Dehalobacter* plays a very active and critical role in the dechlorination of

dichloroanilines at a complex site in Brazil. This knowledge is crucial for developing cost-

effective *in situ* remediation strategies for many other complex contaminated sites.

2   MATERIALS AND METHODS

2.1   Chemicals and stock solutions

Substrates added to enrichment cultures in this study were 3,4-dichloronitrobenzene (3,4-

DCNB), 2,5-dichloronitrobenzene (2,5-DCNB), 2,3-dichloronitrobenzene (2,3-DCNB), 3,4-

dichloroaniline (3,4-DCA), and 2,3-dichloroaniline (2,3-DCA) (Sigma Aldrich, purity > 99%,

(Figure S1). Stock solutions of each compound were prepared in acetone (Sigma Aldrich) at a

concentration of 5 g/L. Ethanol (100%: Sigma Aldrich) and 60% sodium lactate (TCI America)

were used as electron donors. The mentioned DCNB and DCA isomers, aniline, 2-, 3-, and 4-

chloroaniline (CA), 2,5-DCA, DCB isomers (Sigma Aldrich, purity > 99%), and methane (Scott)

were also used as analytical standards. Methanol (Sigma Aldrich) was used as the solvent for

the chlorinated standards and acetonitrile (Fisher Scientific) was used for the mobile phase for

liquid chromatography (described below).

2.2   Preparation of original microcosm study and enrichment cultures

Cultures were enriched from microcosms originating from a large laboratory treatability

study done to evaluate the potential for in situ bioremediation of chlorinated and

nitroaromatic compounds at an industrial site in Brazil. The original microcosms were

constructed in 2015 with solid subsurface material and groundwater from the industrial site,

incubated under anaerobic conditions at site pH (4.6 to 6.5), and amended with various

electron   donors   and   acceptors.   The   microcosms   had   mixtures   of   chloroanilines,

dichloroanilines, dichloronitrobenzenes and chlorobenzenes at concentrations typical of the

site, and biodegradation was monitored closely over a period of one year. After this, some of

the most active microcosms continued to be monitored and re-amended with contaminant

mixtures and electron donors (ethanol and lactate), and topped up with groundwater, as

needed, until March 2020. At this time, six of these most active microcosms (microcosms

Cam16, Cam17, Cam64, Cam65, Cam88, and Cam89), all established with electron donor to

stimulate methanogenic conditions, were selected as inoculum for the bottles described in this

study (Figure S1). The microcosm setup is detailed in Supporting Information Text S1 and in a

Master's Thesis.[20] Results are summarized in Table S1 and Figure S2.

The first set of transfer cultures (34DCNB-T1, 25DCNB-T1, 23DCNB-T1, 34DCA-T1 and

23DCA-T1) and an autoclaved killed control (KC) were inoculated with cell pellets from one

of the microcosms mentioned above (Figure S1). The pellet used as inoculum was obtained

from a 40 ml sample from a shaken microcosm by centrifuging at 12,000 g for 15 min. These

transfer cultures (group T1) were repeatedly amended with single individual chlorinated

substrates: 3,4-DCNB, 2,5-DCNB, 2,3-DCNB, 3,4-DCA, and 2,3-DCA (Figure S1). The killed

control (KC) was autoclaved three times on three consecutive days prior to being amended

with a mix of six compounds: 3,4-DCNB, 2,5-DCNB, 3,4-DCA, 2,3-DCA, 2-CA, and aniline.

An abiotic control (AC1) was also prepared, containing only reduced mineral medium

supplemented with a slightly modified set of six compounds: 3,4-DCNB, 2,3-DCNB, 3,4-DCA,

2,3-DCA, 2-CA, and aniline. This modification was necessary because 2,5-DCNB and 2,3-

DCNB co-elute under the chromatographic conditions used to measure compound

concentrations.

A second generation of transfer cultures (34DCNB-T2, 25DCNB-T2, 23DCNB-T2, 34DCA-

T2, and 23DCA-T2) was prepared from each T1 culture after ~4 months. These were amended

with the same individual compounds as the first generation transfers (3,4-DCNB, 2,5-DCNB,

2,3-DCNB, 3,4-DCA, and 2,3-DCA, respectively) (Figure S1). The DCNB-T2 cultures were

inoculated with a 50% transfer from the corresponding T1 culture (50 mL into 50 mL fresh

medium) while DCA-T2 cultures were inoculated with a 10% transfer from T1 cultures (10

mL of T1 into 90 mL). At the time we initiated the T2 cultures, the mechanism of DCNB

reduction was unclear (discussed later). Therefore, we opted for a more conservative transfer

approach, using a 50% inoculum instead of the 10% used for DCA cultures. Additional mineral

medium supernatant was added to a T2 culture bottle when its volume was depleted to below

50 mL due to sampling, and the cultures were topped up to at least 80 mL. Occasionally, a little

more medium was added to increase the volume of culture up to 250 mL. A second abiotic

control (AC2) was prepared with mineral medium devoid of FeS and amended with 3,4-DCNB,

2,3-DCNB, 3,4-DCA, and 2,3-DCA. The data presented herein focus on the second transfer

(T2) enrichment cultures.

All transfer cultures as well as controls were prepared in autoclaved 250-mL Boston round

bottles sealed with Mininert caps, containing 100 mL of anaerobic mineral medium (pH 7)

reduced with iron sulfide (FeS).[21] Only medium supernatant was used to avoid adding

precipitated FeS. To simulate site groundwater conditions and to promote mineralization of

nitrogen-containing compounds, ammonium was omitted from the medium recipe. All bottles

were incubated statically in the dark at room temperature in an anaerobic glovebox (Coy)

supplied with a gas mix of N_2:CO_2:H_2 (80:10:10%). No replicates were conducted.

The semi-volatile substrates (electron acceptors) were added as evaporated acetone stocks as

described in Text S2. The target concentration was 10 mg/L (52 µM [DCNB] and 62 µM [DCA].

Second transfer enrichments had the concentration increased up to ˜27 mg/L (140 µM [DCNB]

and 170 µM [DCA]) over time, to expose the cultures to a higher concentration of the

corresponding compound for selection purposes. As electron donors, ethanol and lactate were

added to the active cultures and the two controls (AC1 and KC) initially and were re-amended

to active cultures when degradation rates slowed. Electron donors were added in ˜17-fold

excess in electron equivalents (eeq) considering both nitroreduction and complete

dechlorination of DCNB to aniline (Text S2). The initial pH was between 6.6 and 7.1 in all

active enrichments.

2.3   Analytical procedures

Concentrations of DCNB, DCA and CA isomers and aniline were measured using a Hewlett-

Packard 1050 series high-performance liquid chromatography (HPLC) system (Acclaim™ 120

C18 column, Thermo Scientific) equipped with a UV detector (set at 254 nm). The mobile

phase was an isocratic 50:50 mixture of Milli-Q water and acetonitrile (Millipore Sigma) at 1

mL/min with a runtime of 25 min. Samples (2 mL) were taken from cultures using a 2 mL

gastight glass syringe and filtered through a 0.22 µm (13 mm) Chromspec PTFE syringe filter

(pre-wetted with methanol) prior to being injected into the HPLC. Lactate, acetate,

propionate, formate, butyrate, pyruvate, chloride, nitrate, nitrite, sulfate, and phosphate were

measured by ion chromatography (IC) using a Dionex™ Integrion™ IC system (Thermo Fisher

Scientific). Methane was measured in headspace samples by gas chromatography (GC) using

an Agilent 7890A gas chromatograph with headspace autosampler G1888 (Agilent

Technologies). Method details are provided in Text S3.

2.4   Microbial community analysis and qPCR

Well-mixed samples (4 mL) from enrichment cultures were collected periodically and

centrifuged at 10,000 rpm for 20 minutes. The supernatant was removed and cell pellets stored

at -80°C. DNA was extracted using the DNeasy PowerSoil Kit (Qiagen) and quantified using a

Qubit® 3.0 Fluorometer (Thermo Fisher Scientific). DNA samples were sent for Illumina

MiSeq PE300 amplicon sequencing at an external laboratory (Genome Quebec Innovation

Centre, Montreal, Canada), using modified "staggered end"[22] versions of primers 926f-modified

(5'-AAACTYAAAKGAATWGRCGG-3') and 1392r-modified (5'-ACGGGCGGTGWGTRC-

3'), which target the V6–V8 region of the 16S rRNA gene from both Bacteria and Archaea, as

well as the 18S rRNA gene in Eukarya.[23,24] Sequencing data was processed using QIIME 2

version 2021.2.[25] After trimming the primer region with the cutadapt plug-in, amplicon sequence

variants (ASVs) were generated using the DADA2 plugin,[26] and taxonomy was assigned to the

ASVs through the feature-classifier plugin using the SSU SILVA v.138 database.[27] Data were

further processed in Excel. Sequences <400 bp in length were removed and ASVs were ranked

by maximum sequence abundance across all DNA samples. ASVs with abundance greater than

1% in at least one sample were kept for graphing. The Basic Local Alignment Search Tool

(BLAST) available at the National Center for Biotechnology Information (NCBI) was used to

compare ASV sequences of interest. To investigate the presence of the organisms of interest

identified in the cultures, key genera were subsequently searched in a database containing 16S

rRNA gene sequences retrieved from field samples collected by our collaborating partner

during various remediation pilot projects conducted at the site. This database also contains all

raw DNA sequences from microcosms and enrichments and was uploaded to the NCBI

(Accession PRJNA1040917 as described in Table S2).

DNA samples were also analyzed by quantitative polymerase chain reaction (qPCR) using a

CFX96TM real-time PCR detection system (Bio-Rad Laboratories Inc.). Primers GenBac1055f

(5'-ATGGYTGTCGTCAGCT-3') and GenBac1392r (5'-ACGGGCGGTGTGTAC-3') targeting

most bacteria[24,28,29] and primers GenArch787f (5'-ATTAGATACCCGBGTAGTCC-3') and

GenArch1059r (5'-GCCATGCACCWCCTCT-3') targeting most archaea[30] were used. A no-

template control was also included in each run. All qPCR samples were run in triplicate.

Details on the qPCR method are provided in Text S4. Absolute total bacterial and archaeal

abundance obtained by qPCR (copies 16S rRNA gene per ml) was used to convert the relative

abundances obtained from amplicon sequencing to absolute abundance for each ASV.

A total of forty-eight DNA samples were sequenced and analyzed by qPCR, including

samples from the original subsurface solid material, microcosms, and enrichments. On average

66,000 16S rRNA gene amplicon reads were retrieved per sample (Table S3a). Reads were

grouped by bacteria and archaea, then sequences with relative abundance >1% in any sample

(Table S3b for bacteria and Table S3c for archaea) were used to calculate absolute abundances

(copies/mL) of each ASV using total abundances of bacteria and archaea determined by qPCR

(Table S4). Final absolute abundances for each ASV (copies/mL) are reported in Tables S5a to

S5f for bacteria; Tables S6a to S6f for archaea. Finally, Non-metric Multi-dimensional Scaling

(NMDS) and bubble plots were constructed as explained in Text S5.

3   RESULTS AND DISCUSSION

3.1   Abiotic transformation of DCNB was not sustained

Contaminant profiles in three controls (AC1, AC2 and KC) were assessed for possible abiotic

reactions. Chloroanilines and aniline were not transformed in any of these controls.

Dichloronitrobenzenes were not transformed in medium control (AC2) without FeS (Figure

S3; Table S7). In the medium control (AC1) with FeS, 3,4-DCNB was not transformed, but 2,3-

DCNB was reduced to 2,3-DCA (Figure S4; Table S7); however, this activity was not sustained

upon reamending with 2,3-DCNB. These data are consistent with reports that reduced

minerals, such as FeS or perhaps reduced resazurin, can sometimes mediate reduction of the

nitro group. [31–38] Curiously, DCNB isomers were repeatedly transformed to DCA in the

autoclaved control KC (Figure S5; Table S7); however later analyses (to be discussed) revealed

that KC was in fact not sterile.

3.2 Biologically-mediated nitroreduction and dechlorination

Each transfer culture established on single substrates remained active, sustained by refeeding

chlorinated substrates (DCNB or DCA) and electron donors as needed for over 500 days.

Historical data are provided in Table S8 (T1) and Table S9 (T2). Focusing on T2 transfers from

day 280 to 520 and on the three DCNB-amended cultures first, we observed consistent and

complete transformation of each DCNB isomer to the corresponding dichloroaniline (Figure

1a-c). In the culture amended with 3,4-DCNB the product, 3,4-DCA, was further

dechlorinated to 3- and 4-CA (Figure 1a), while in the cultures amended with 2,3- and 2,5-

DCNB, the transformation stopped at the corresponding dichloroaniline (Figures 1b and 1c).

The sum of measured chlorinated products agreed well with the cumulative DCNB added.

Reduction of 2,3-DCNB (<~1 µM per day) was ~5 times slower than the reduction of either

3,4- or 2,5-DCNB (5-6 µM per day).

Figure 1. Degradation profiles (upper panels a), b) and c); μmole/bottle) and corresponding major microbial taxa (lower panels d), e) and f); copies/mL) in DCNB-amended T2 transfer cultures from day 280 to 520. Cumulative DCNB added (———) is shown for each amendment (step) of 70 to 140 μM (10 to 20 μmol per bottle). The sum of products (- -○- -), including DCA (—■—), 4-CA (—▲—), 3-CA (—▲—), 2-CA and aniline (not shown, mostly below detection), illustrates a reasonable mole balance. Measured DCNB concentrations are not shown for clarity but can be found in Table S9. In lower panels amplicon sequence variants (ASVs) from the same taxonomic designation are assigned the same colour.

Focusing next on the two DCA-amended transfers, 3,4-DCA was dechlorinated to a mix of

3- and 4-CA (Figure 2a), while 2,3-DCA was dechlorinated primarily to 3-CA, with traces of

2-CA (Figure 2b). These cultures dechlorinated about ~5 to 6 µM DCA to CA per day.

Figure 2. Degradation profiles (upper panels a), b) and c); µmole/bottle) and corresponding

major microbial taxa (lower panels, d) and e); copies/mL) in two DCA-amended T2 transfer

cultures from day 280 to 520 (panels a), b), d), and e)) and from day 560 to 625 (panel c)). In

panels a) and b), cumulative DCA added (————) is shown for each amendment (step) of ~80

µM (~12 µmol per bottle). The sum of products (- -o- -), including DCA (——■——), 4-CA (

——▲——), 3-CA (——▲——), 2-CA (——▲——), and aniline (not shown, mostly below detection),

illustrates a reasonable mole balance. Measured DCA concentrations are not shown for clarity

but can be found in Table S9. Panel c) shows degradation profile after 2,3-DCA-T2 was fed

with 2,3-DCNB (—●—) instead of 2,3-DCA on Day 568. The open circle shows the amount of

2,3-DCNB fed to 23DCA-T2 (—○—). In lower panels amplicon sequence variants (ASVs) from

the same taxonomic designation are assigned the same color.

Given the very slow transformation rate in 23DCNB-T2, we wondered if the enrichment

culture dechlorinating 2,3-DCA (23DCA-T2; Figure 2b) could also reduce 2,3-DCNB, even

though this culture - and the microcosm (Cam16) from which it was derived - had never been

exposed to any DCNB isomer. Surprisingly, 2,3-DCNB was reduced without a lag to 2,3-DCA

which was further dechlorinated to 3-CA (Figure 2c). We then tested all active DCA-

degrading cultures with a mix of the three DCNB isomers; all the cultures were able to reduce

DCNBs to their respective DCA isomer (data not shown).

3.3 Microbial community composition reflects electron balance

To relate the microbial community to activity, we first identified abundant bacterial and

archaeal ASVs (>1%) and then calculated their absolute abundance in active cultures focusing

on 2-time points for each sample. Figure 3 provides a summary of the most relevant taxa

identified. Most of the enriched bacterial ASVs belong to the phylum Bacillota (Firmicutes)

including *Sporomusa*, *Anaerospora*, and unassigned members of the Veillonellales-

Selenomonadales. These organisms clearly grew on the ample electron donor amendments to

the cultures. ASVs for *Anaeromyxobacter*, *Desulfitobacterium* and *Desulfovibrio* were also

retrieved. These genera are also often implicated in nitrate or sulfate reduction though some

members may be facultative dechlorinators. ASVs for obligate dechlorinators of the genus

Dehalobacter were also identified (Figure 3). A complete overview of all major taxa (>1%) are

shown in Figure S6, for bacteria, and Figure S7, for archaea.

Figure 3. Absolute abundance of major organisms (copies/mL) and proportion of electron

equivalents from amended donors channeled to each electron-accepting processes (% electron

equivalents) for days 297 and 517. Veillonellales-Selenomonadales order is shortened to

"Veill"; *Sporomusa* genus is shortened to "*Sporo*". Calculated absolute abundances for all ASVs

are available in Table S5 (bacteria) and Table S6 (archaea).

To confirm the extent of dechlorination, an electron balance of possible donors and acceptors

was completed using measured concentrations of chlorinated organics, methane (Table S9) and

anions (Table S10). As expected from stoichiometry (e.g., $C_6H_5Cl_2N + H_2 \rightarrow C_6H_6ClN + HCl$),

chloride concentrations increased proportionally in cultures that exhibited dechlorination

(34DCNB-T2, 34DCA-T2, and 23DCA-T2). The measured increase in chloride was slightly

higher than expected in almost all cases (Table S11), likely because of the error associated with

measuring small changes against a high background of ˜1900 µM chloride.

Potential electron-accepting processes include acetogenesis, methanogenesis,

nitroreduction, dechlorination, and sulfate reduction. For T2 cultures between days 280 and

520, we calculated the proportion of electron equivalents from added donors channeled to

each of the different electron-accepting processes (Figure 3, bottom table). In DCNB-amended

cultures, acetogenesis from lactate and ethanol was the dominant process, accounting for over

80% of electrons from donors. Methanogenesis was severely limited. By contrast, in DCA-

amended cultures, methanogenesis accounted for 60-70% of electron equivalents from donors,

which combined with acetogenesis accounted for over 98% of electron equivalents. A buildup

of acetate in all cultures (Tables S10 and S12) suggested possible inhibition of acetoclastic

methanogens specifically, particularly by DCNBs. Culture 34DCNB-T2 produced the most

methane among DCNB-amended cultures but also exhibited the most rapid and extensive

nitroreduction and dechlorination, thus relieving inhibition associated with DCNBs and

DCAs.

Other electron accepting processes accounted for a much lower proportion of activity

(Figure 3). Nitroreduction consumed 5% of electron equivalents in cultures fed with 3,4- and

2,5-DCNB isomers and up to 20% of electron equivalents in culture 23DCNB-T2, although

activity in this culture was very low and no lactate was consumed (Tables S10 and S12). In all

cases, dechlorination accounted for less than 3% of all electron equivalents provided. The small

amount of sulfate (~30 µM) present in the medium was rapidly depleted in the 4 highly active

cultures (34DCNB-T2, 25DCNB-T2, 34DCA-T2, and 23DCA-T2). However, less than 1% of

electron equivalents were channeled to sulfate reduction. In culture 23DCNB-T2, sulfate was

not consumed, consistent with low activity observed in this enrichment (Table S11).

We compared the proportions of electrons consumed by each electron-accepting process

with corresponding microbial proportions (Figure 3) to infer function (Table S12). Methane

production was greatest in cultures 23DCA-T2 and 34DCA-T2, lower in 34DCNB-T2, very

low in 25DCNB-T2 and absent in 23DCNB-T2. Methanogens were correspondingly most

abundant in 23DCA-T2, 34DCA-T2, and 34DCNB-T2. ASVs corresponding to

hydrogenotrophic *Methanoregula* dominated the archaeal population, with a much lower

abundance of *Methanocella* and other hydrogenotrophic genera (Figure S7). The presence of

only hydrogenotrophic methanogens combined with substantial accumulation of acetate

strongly indicates that DCNB isomers and to a lesser extent DCA isomers inhibited acetoclastic

methanogens.

3.4   Nitro group reduction can be mediated by many different microbes

We had hoped to observe enrichment of specific microbes in DCNB-amended cultures to

enable functional attribution. Rather, it seems that many different genera can likely reduce

the nitro group. Biologically regenerated FeS is unlikely to play a role as a reductant, as there

was no evidence of sulfate or iron cycling in the cultures, and the rates of DCNB reduction

were similar regardless of FeS concentrations. In all cultures, there was significant

representation of fermenting or acetogenic microbes, particularly Veillonellales-

Selenomonadales, *Sporomusaceae*, *Sporomusa*, and *Anaerospora* (Figure S6). This is consistent

with the ample supply of ethanol and lactate to all cultures.

We wondered if these fermenting and acetogenic organisms could also be reducing DCNB

to DCA as a growth-associated electron-accepting process akin to nitrate reduction. In

25DCNB-T2, where only nitroreduction was observed without dechlorination, 6 ASVs (ASV1

to 5 and 8) (>90% identity) classified only to the order level as Veillonellales-Selenomonadales

were predominant (Table S5b). They represented more than 80% of the bacterial community

in this culture on day 517 (Figure 1e). The closest matches to these sequences are affiliated

with the genera *Pelosinus*, *Anaerospora*, *Sporomusa*, *Dendrosporobacter*, among other

organisms; interestingly, one of the closely matching sequences (KF460373.1) comes from a

study on the impact of nitrate on pentachlorophenol dechlorination[39] (Figure S8).

In fact, ASVs from Veillonellales-Selenomonadales were abundant in all 5 active cultures

(Figures 1d, e, and f, Figures 2d and e and Figure 3), whether fed DCNB or DCA, making

functional attribution difficult. Finding that DCNB isomers could be reduced in all cultures

regardless of enrichment substrate is perhaps then not so surprising. An autoclaved control

(KC) amended with a mixture of compounds (3,4-DCNB, 2,5-DCNB, 3,4-DCA, 2,3-DCA, 2-

CA, aniline) was prepared to try to distinguish between biotic and abiotic reactions.

Unfortunately, the bottle did not achieve sterility, despite autoclaving it on three consecutive

days. In this "killed" control, both 3,4-DCNB and 2,5-DCNB were reduced to the

corresponding DCA isomers (Figure S5). DNA was readily extracted from this control, and

although the total microbial abundance was an order of magnitude lower than other cultures,

the KC culture was dominated by a unique Butyricoccaceaea ASV as well as other spore-

forming and presumably heat-resistant microbes. Among the microbes detected in KC after

refeeding 3,4- and 2,5-DCNB 4 times, *Desulfitobacterium* and *Anaerospora* were also detected

in other active nitroreducing cultures (i.e., 23DCNB-T2 and 25DCNB-T2) (Figure S6),

suggesting these genera are also candidates for nitroreduction. Ultimately it became clear that

nitroreduction is a widespread microbial activity that was easy to stimulate with electron

donor in the cultures, however, the contribution of biotic nitroreduction to microbial growth

is yet to be investigated.

3.5 Dechlorination of 2,3- and 3,4-DCA supports growth of *Dehalobacter*

In actively dechlorinating cultures, *Dehalobacter* was the only microbe to systematically

show a substantial increase in abundance. *Dehalobacter* was not detected in cultures that were

only catalyzing nitroreduction (25DCNB-T2 and 23DCNB-T2). Given that it is an obligate

organohalide respiring bacterium, we conclude that *Dehalobacter* was responsible for the

DCA-dechlorinating activity in enrichment cultures amended with 3,4-DCNB, 3,4-DCA, and

2,3-DCA (Figures 1d and 2d and e).

Two distinct *Dehalobacter* ASVs were identified: ASV1 in 34DCNB-T2 and ASV2 in 34DCA-

T2 and 23DCA-T2 (Text S6), with maximum bacterial abundances of 14, 25, and 31% in each

culture, respectively (Table S3b). ASV1 (468 bp) is 100% identical to several previously

published 16S rRNA gene sequences of *Dehalobacter restrictus* strains, including strain SAD

(CP148032.1), strain DSM 9455 (CP007033.1), and strain 12DCA (CP046996.1), as well as to

three *Dehalobacter* strains classified only to the genus level, including strain DCA

(CP003869.1), strain DAD (CP148031.1), and strain CF (1CP003870.1). The known substrates

for these *Dehalobacter* strains include tetra- and trichloroethene,[40] 1,2-dichloroethane,[41] tri-

and dichloromethane,[42] and 1,1,1-trichloroethane and 1,1-dichloroethane.[43,44]

Dehalobacter ASV2 (468 bp) was not a perfect match to any 16S rRNA gene sequence in the

NCBI database. It shares only 97.01% identity to ASV1 and 98.93% to *Dehalobacter* strain

DCM (CP092282.1) described as "dichloromethane-degrading bacterium from Xi river" (Direct

Submission by Yang, Y. *et al.*, 13-Jan-2022). Neither *Dehalobacter* ASV1 nor ASV2 sequences

matched (<95% identity) to the 16S rRNA gene fragment (937 bp) from *Dehalobacter* sp. strain

14DCB1 (JN051267.1), a strain enriched on dichlorobenzene[45] but reported to dechlorinate

dichloroaniline.[14]

To further confirm the role of *Dehalobacter* in chloroaniline dechlorination, we estimated

growth yields from qPCR and concentration data to compare with previously published yields.

The net growth yields ranged from 1.4×10^{13} to 6.5×10^{13} cells per mol of chloride released

assuming 3 copies of the 16S rRNA gene per cell (Table S13). The yields calculated in this study

are consistent with many previously reported yields for other *Dehalobacter* strains using

various chloroethanes, chlorobenzenes, and chloromethanes as electron acceptors for growth

(Table S14). In fact the yields align with the highest yields previously reported, solidifying

chloroaniline dechlorination as a growth-related process.

Additional evidence for *Dehalobacter* being responsible for DCA dechlorination was

obtained when non-dechlorinating culture 25DCNB-T2 was inoculated with 3 mL of

dechlorinating culture 34DCNB-T2 (on day 653). After inoculation, residual 2,5-DCA

previously formed from 2,5-DCNB was dechlorinated to 3-CA and the relative abundance of

*Dehalobacter*_ASV1 increased from zero on day 517 to ~9% on day 653 (Figure S9).

A shift in activity in culture 23DCA-T2 was noted as monitoring continued for over 1000

days (Figure S10). As time progressed, 2-CA rather than 3-CA became the major product from

2,3-DCA. Thereafter, 2-CA was slowly yet consistently transformed to aniline. On day 881,

*Dehalobacter*_ASV2 constituted 16% of the bacterial community (Table S5e). 2-CA was

dechlorinated to aniline in subsequent transfers from this culture as well (data not shown). A

subsample of 23DCA-T2 was taken on day 653 for additional growth experiments to ultimately

build up enough culture to measure compound specific isotopic enrichment factors. This

ongoing research implicates *Dehalobacter* in aniline formation as described in a companion

publication.[46]

3.7   Other organisms identified in enriched cultures

Several other ASVs belonging to known facultative OHRB were identified, including

Anaeromyxobacter, *Desulfovibrio*, and *Desulfitobacterium* (Figure 3). While some members

of these genera can dechlorinate, most are known to be quite versatile and may reduce sulfate,

nitrate or even grow fermentatively in the absence of a terminal electron acceptor. An

Anaeromyxobacter ASV found in culture 34DCNB-T2 is 99.79% similar over ~450 bp to the

16S rRNA gene of *Anaeromyxobacter* sp. strain RF71 (MZ254878.1), 99.57% similar to *A.*

dehalogenans 2CP-C (CP000251.1) and 99.36% similar to *A. dehalogenans* 2CP-1

(CP001359.1). Both strains 2CP-C and 2CP-1 can gain energy from the reductive

dechlorination of chlorophenol and contain putative dehalogenases.[47–49] However, the

abundance of *Anaeromyxobacter* in the culture was far less than *Dehalobacter* and is therefore

unlikely to be primarily responsible for dechlorination.

A single ASV of *Desulfovibrio* increased in abundance over time in the three cultures

dechlorinating DCA isomers (Figure 3). *Desulfovibrio* are sulfate-reducing bacteria that can

also ferment substrates such as lactate; some can respire organohalides.[50,51] In cultures 23DCA-

T2 and 34DCA-T2, *Desulfovibrio* increased between days 280 and 520 to similar extent as

Dehalobacter (Figures 2d and 2e). However, a similar increase was not seen in dechlorinating

culture 34DCNB-T2 (Figure 1d). The amount of sulfate in the medium (around 30 μM) is too

small to explain this growth, therefore *Desulfovibrio* was more likely metabolizing lactate to

acetate, possibly participating in dechlorination alongside *Dehalobacter*. The *Desulfovibrio*

ASV in our cultures is 100% identical to a non-dechlorinating *Desulfovibrio* sp. (JQ599691.1)

detected in a dichloroethane dechlorinating enrichment.[52] The sequence was also 100%

identical to an uncultured *Desulfovibrio* clone (AB854345.1) found in a non-dechlorinating

environment.[53] Therefore, the more likely role of this organism is fermentation and

acetogenesis, in support of dechlorinating *Dehalobacter*.

Several ASVs (sharing > 94% identity) were assigned to *Desulfitobacterium*. Interestingly,

these ASVs were only found in the cultures reducing DCNBs and in the KC control that

exhibited nitroreduction; they were not detected in the DCA-amended cultures (Figure 3).

This seems to implicate *Desulfitobacterium* in nitroreduction, however the ASVs were in low

abundance. Furthermore, DCA-amended cultures could also rapidly reduce the nitro group,

therefore *Desulfitobacterium* was not solely responsible for nitroreduction, if at all. Moreover,

the 8 most abundant *Desulfitobacterium* ASV sequences are > 95% similar to a

Desulfitobacterium sequence found in *Dehalococcoide*s-containing dechlorinating cultures

(AB596883.1) (Figure S11), where they were not implicated in dechlorination,[54] a likely

scenario in our study as well.

3.8   Data integration and application to improving remediation outcomes

Non-metric multidimensional scaling (NMDS) was used to evaluate relationships among

cultures as they were enriched from subsurface material to initial microcosms and finally to

enrichment cultures (Figure 4). First, there is clearly an impact of the original source material,

shown in red, blue and green for three original sediment sources used to construct the

microcosms. Second, there was a convergence of community depending on substrate.

Enrichment cultures that dechlorinate DCAs are distinct from those that only carry out

nitroreduction. Samples from 34DCNB-T2 culture, in which both nitroreduction and

dehalogenation were observed, are located between the dechlorinating-exclusive culture

samples (34DCA-T2, and 23DCA-T2) and the nitroreducing-exclusive cultures (25DCNB-T2,

23DCNB-T2 and KC). *Dehalobacter* in the top groupings and *Desulfitobacterium* in the bottom

two cultures (23DCNB-T2 and KC) are major genera driving the vertical spread of samples in

the NMDS plot. Relationships between original microcosms and transfers are also illustrated

in Figure S1.

Figure 4. Non-metric multidimensional scaling plot based on relative abundance of 16S rRNA amplicons in 45 samples from the original site material used for microcosms and enrichment cultures.

In a survey of over 60 groundwater samples from around the site in Brazil that were analyzed by amplicon sequencing, approximately 14% returned *Dehalobacter*. Of the ~400 samples that have been sequenced from various remediation trials implemented at the industrial site – including samples from pilot wetlands and pre and post biosparging units – many different *Dehalobacter* ASVs (including those enriched in this study) were detected in 78 samples

(~20%) clustered around more impacted zones with lower redox potential. Relative

abundances of *Dehalobacter* reached 2% of the bacterial population in some locations, which

is high considering the complex mixture of chemicals and conditions at the site.[1,2] These

abundances are being used to delineate biologically active zones and to predict attenuation

rates to most effectively remediate the site.

Other ASVs detected in the enrichment cultures were detected in site samples, including the

ones affiliated with *Anaeromyxobacter* and *Desulfitobacterium*, which were detected in ~53%

and ~18% of the total ~460 samples collected by the partner company, respectively. The ASVs

classified as Veillonellales-Selenomonadales were detected in ~11% of the same group of

samples, while the ones classified further to *Sporomusa* and *Anaerospora* genera in ~3% and

~6%. The high relative abundance of the Veillonellales-Selenomonadales order in our

enriched cultures indicates those organisms were favoured by the ample availability of donor

(ethanol and lactate) added to the cultures, which is not the case in the field.

The fact that the microbes identified in enrichment cultures are also present at the site attests

to the utility of such enrichments in site characterization and provides very strong evidence

that these processes are contributing to active natural attenuation in anaerobic zones. At the

site, concentrations of dichloroanilines are generally higher than dichloronitrobenzenes,

consistent with our finding that biologically-mediated nitroreduction occurs readily and that

there are many genera capable of catalyzing this reaction. It is highly likely that some of the

other contaminants at the site could be electron donors, including petroleum hydrocarbons,

solvents and plasticizers. In addition, abiotic reactions may also occur driven by reduced

minerals or by reduced natural organic matter, which can donate electrons for

nitroreduction.[37,55–58] Further studies will seek to better understand the players and

nitroreduction mechanisms in DCNB cultures.

Two distinct *Dehalobacter* ASVs were unequivocally linked to dechlorination, and their

presence along with other ASVs in site samples is indicative of dechlorinating activity in situ.

The two ASVs originated from geographically distinct locations at the site. Traces of aniline

were found and continue to accumulate in 23DCA-T2, suggesting further dechlorination of

CA is possible in situ. Inhibition of acetoclastic methanogens was noted, which is also

consistent with site data and provides a means to evaluate site recovery: as DCNB and DCA

concentrations decrease with active remediation efforts, methanogenic populations in anoxic

zones should rebound. Biomarkers based on 16S rRNA genes are already being deployed at the

site to track population changes as remediation alternatives are evaluated. The robust

enrichment cultures generated herein will be invaluable for future studies of

biotransformation mechanisms, development of isotopic enrichment factors and of additional

biomarkers, and possibly even for use as bioaugmentation cultures.

SUPPORTING INFORMATION

The following files are available free of charge:

Additional details on experiments, materials and methods, and data obtained, including texts

and figures mentioned in this paper (PDF); Tables including data obtained and calculations

mentioned in this paper (.xls).

ACKNOWLEDGEMENTS

For the financial support for Sofia Araújo, we thank FACEPE (Science and Technology

Foundation of the State of Pernambuco, Brazil – process number BFD-0013-3.07/21), CNPq

(National Council for Scientific and Technological Development, Brazil - process number

165130/2017-2), and CAPES (Coordination for the Improvement of Higher Education

Personnel) for the Graduate Program support (Proap) and Institutional Internationalization

Program (PrInt) (process number 88887.363316/2019-00). This work was also supported by

Mitacs through the Mitacs Accelerate and Mitacs Elevate programs (Application ref. IT23315

and IT34875, Canada).

The authors also acknowledge the grant from CNPq (process number 304862/2018-5) and

FACEPE (process number APQ 0456-3.07/20), and NSERC (Natural Sciences and Engineering

Research Council of Canada) Alliance project (ALLRP 575243-22), which has Geosyntec

Consultants and Corteva Agriscience™ as the two main industrial partners.

The authors do not have a conflict of interest to declare. This publication is in accordance

with the Brazilian biodiversity law n.º 13.123, from May 20[th], 2015. This is non-profit research

and has no commercial intention between the parties. This is a collaboration project between

the parties involved. Access Registration No. A3D9F11.

We would also like to thank Elodie Passeport, Shuping Wang, and Luiz Pereira for their

helpful discussions and collaborative work on this project.

REFERENCES

(1)   Melo, N.; Araújo, S. P.; de Paula Queiroz Kraus, S.; Lomheim, L.; Quintero, P. B.; Mack, E. E.; Edwards, E. A.; Spain, J.; Gavazza, S. Strategies for Bioremediation of Soil from an Industrial Site Exposed to Chlorinated and Nitroaromatic Compounds. *Groundwater Monit. Rem.* **2023**, *43* (3), 108–120. https://doi.org/10.1111/gwmr.12596.

(2)   Barreto, P.; Lemes, M.; Jimenez, J.; Mack, E. E.; Henderson, J.; Freedman, D. L. Evaluation of Strategies to Remediate Mixed Wastes at an Industrial Site in Brazil. *Groundwater Monit. Rem.* **2023**, *43* (3), 93–107. https://doi.org/10.1111/gwmr.12607.

(3)   Buser, H.-R.; Bosshardt, H.-P. Studies on the Possible Formation of Polychloroazobenzenes in Quintozene Treated Soil. *Pestic Sci* **1975**, *6*, 35–41.

(4)   Roldán, M. D.; Pérez-Reinado, E.; Castillo, F.; Moreno-Vivián, C. Reduction of Polynitroaromatic Compounds: The Bacterial Nitroreductases. *FEMS Microbiol. Rev.* **2008**, pp 474–500. https://doi.org/10.1111/j.1574-6976.2008.00107.x.

(5)   Bunce, N. J.; Merrick, R. L.; Corke, C. T. Reductive Transformations of Nitrate with 3,4-Dichloroaniline and Related Compounds by *Escherichia Coli*. *J Agric Food Chem* **1983**, *31* (5), 299–326.

(6)   Spain, J. C. Biodegradation of Nitroaromatic Compounds. *Annu Rev Microbiol* **1995**, *49*, 523–555.

(7)   Esteve-Nunez, A.; Lucchesi, G.; Philipp, B.; Schink, B.; Ramos, J. L. Respiration of 2,4,6-Trinitrotoluene by *Pseudomonas* sp. strain JLR11. *J Bacteriol* **2000**, *182* (5), 1352–1355.

(8)   Huang, J.; Ning, G.; Li, F.; Sheng, G. D. Biotransformation of 2,4-Dinitrotoluene by Obligate Marine *Shewanella marisflavi* EP1 under Anaerobic Conditions. *Bioresour Technol* **2015**, *180*, 200–206. https://doi.org/10.1016/j.biortech.2014.12.108.

(9)   Madeira, C. L.; Menezes, O.; Park, D.; Jog, K. V.; Hatt, J. K.; Gavazza, S.; Krzmarzick, M. J.; Sierra-Alvarez, R.; Spain, J. C.; Konstantinidis, K. T.; Field, J. A. Bacteria Make a Living Breathing the Nitroheterocyclic Insensitive Munitions Compound 3-Nitro-1,2,4-Triazol-5-One (NTO). *Environ Sci Technol* **2021**, *55* (9), 5806–5814. https://doi.org/10.1021/acs.est.0c07161.

(10) Struijs, J.; Rogers, J. E. Reductive Dehalogenation of Dichloroanilines by Anaerobic Microorganisms in Fresh and Dichlorophenol-Acclimated Pond Sediment. *Appl Environ Microbiol* **1989**, *55* (10), 2527–2531.

(11) Kuhn, E. P.; Suflita, J. M. Sequential Reductive Dehalogenation of Chloroanilines by Microorganisms from a Methanogenic Aquifer. *Environ Sci Technol* **1989**, *23* (7), 848–852. https://doi.org/10.1021/es00065a014.

(12) Kuhn, E. P.; Townsend, G. T.; Suflita, J. M. Effect of Sulfate and Organic Carbon Supplements on Reductive Dehalogenation of Chloroanilines in Anaerobic Aquifer Slurries. *Appl Environ Microbiol* **1990**, *56* (9), 2630–2637. https://doi.org/10.1128/aem.56.9.2630-2637.1990.

(13) Travkin, V.; Baskunov, B. P.; Golovlev, E. L.; Boersma, M. G.; Boeren, S.; Vervoort, J.; Berkel, W. J. H.; Rietjens, I. M. C. M.; Golovleva, L. A. Reductive Deamination as a New Step in the Anaerobic Microbial Degradation of Halogenated Anilines. *FEMS Microbiol Lett* **2002**, *209* (2), 307–312. https://doi.org/10.1111/j.1574-6968.2002.tb11149.x..

(14) Zhang, S.; Wondrousch, D.; Cooper, M.; Zinder, S. H.; Schüürmann, G.; Adrian, L. Anaerobic Dehalogenation of Chloroanilines by *Dehalococcoides* m*ccartyi* strain CBDB1 and *Dehalobacter* strain 14DCB1 via Different Pathways as Related to Molecular Electronic Structure. *Environ Sci Technol* **2017**, *51* (7), 3714–3724. https://doi.org/10.1021/acs.est.6b05730.

(15) Susarla, S.; Yonezawa, Y.; Masunaga, S. Reductive Dehalogenation of Chloroanilines in Anaerobic Estuarine Sediment. *Environ. Technol.* **1997**, *18* (1), 75–83. https://doi.org/10.1080/09593331808616514.

(16) Susarla, S.; Yonezawa, Y.; Masunaga, S. Reductive Transformations of Halogenated Aromatics in Anaerobic Estuarine Sediment: Kinetics, Products and Pathways. *Water Res* **1998**, *32* (3), 369–348.

(17) Duc, H. D.; Oanh, N. T. Anaerobic Degradation of Chloroanilines by *Geobacter* sp. KT5. *Curr Microbiol* **2019**, *76* (2), 248–257. https://doi.org/10.1007/s00284-018-1617-7.

(18) Oanh, N. T.; Duc, H. D. Anaerobic Degradation of Propanil in Soil and Sediment Using Mixed Bacterial Culture. *Curr Microbiol* **2021**, *78* (4), 1499–1508. https://doi.org/10.1007/s00284-021-02419-7.

(19) Lu, H.; Zhang, T.; Zhou, Y.; Zhou, J.; Wang, J.; Wang, X. Enhanced Dechlorination and Biodegradation of 2-Chloroaniline by a 2-Aminoanthraquinone-Graphene Oxide Composite

under Anaerobic Conditions. *Sci Rep* **2019**, *9* (1), 12376. https://doi.org/10.1038/s41598-019-48904-9.

(20) Kraus, S. Aerobic and Anaerobic Biotransformation of Chloroanilines, Chlorobenzenes, and Dichlonitrobenzenes at a Complex Industrial Site in Brazil and Analysis of Associated Microbial Communities. Master Thesis, University of Toronto, Toronto, 2018. https://hdl.handle.net/1807/91484 (accessed 2024-06-11).

(21) Edwards, E. A.; Grbic-Galic, D. Anaerobic Degradation of Toluene and O-Xylene by a Methanogenic Consortium. *Appl Environ Microbiol* **1994**, 313–322.

(22) Toth, C. R. A.; Luo, F.; Bawa, N.; Webb, J.; Guo, S.; Dworatzek, S.; Edwards, E. A. Anaerobic Benzene Biodegradation Linked to the Growth of Highly Specific Bacterial Clades. *Environ Sci Technol* **2021**, *55* (12), 7970–7980. https://doi.org/10.1021/acs.est.1c00508.

(23) Engelbrektson, A.; Kunin, V.; Wrighton, K. C.; Zvenigorodsky, N.; Chen, F.; Ochman, H.; Hugenholtz, P. Experimental Factors Affecting PCR-Based Estimates of Microbial Species Richness and Evenness. *ISME Journal* **2010**, *4* (5), 642–647. https://doi.org/10.1038/ismej.2009.153.

(24) Ferris, M. J.; Muyzer, G.; Ward, D. M. Denaturing Gradient Gel Electrophoresis Profiles of 16S rRNA-Defined Populations Inhabiting a Hot Spring Microbial Mat Community. *Appl Environ Microbiol* **1996**, *62* (2), 340–346.

(25) Bolyen, E.; Rideout, J. R.; Dillon, M. R. Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2. *Nat. Biotechnol*, **2019**, pp 852–857. https://doi.org/10.1038/s41587-019-0190-3.

(26) Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S. P. DADA2: High-Resolution Sample Inference from Illumina Amplicon Data. *Nat Methods* **2016**, *13* (7), 581–583. https://doi.org/10.1038/nmeth.3869.

(27) Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F. O. The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing and Web-Based Tools. *Nucleic Acids Res* **2013**, *41*, D590-6. https://doi.org/10.1093/nar/gks1219.

(28) Ritalahti, K. M.; Amos, B. K.; Sung, Y.; Wu, Q.; Koenigsberg, S. S.; Löffler, F. E. Quantitative PCR Targeting 16S rRNA and Reductive Dehalogenase Genes Simultaneously Monitors Multiple *Dehalococcoides* strains. *Appl Environ Microbiol* **2006**, *72* (4), 2765–2774. https://doi.org/10.1128/AEM.72.4.2765-2774.2006.

(29) Lane, D. J. 16S/23S rRNA Sequencing. In *Nucleic acid techniques in bacterial systematics*; Stackebrandt, E., Goodfellow, M., Eds.; John Wiley and Sons: Chichester, United Kingdom, 1991; pp 115–175.

(30) Yu, Y.; Lee, C.; Kim, J.; Hwang, S. Group-Specific Primer and Probe Sets to Detect Methanogenic Communities Using Quantitative Real-Time Polymerase Chain Reaction. *Biotechnol Bioeng* **2005**, *89* (6), 670–679. https://doi.org/10.1002/bit.20347.

(31) Cheng, J.; Suidan, M. T.; Venosa, A. D. Abiotic Reduction of 2,4-Dinitrotoluene in the Presence of Sulfide Minerals under Anoxic Conditions. *Wat. Sci. Tech.* **1996**, *34* (10), 25–33. https://doi.org/10.1016/S0273-1223(96)00693-2.

(32) Elsner, M.; Schwarzenbach, R. P.; Haderlein, S. B. Reactivity of Fe(II)-Bearing Minerals toward Reductive Transformation of Organic Contaminants. *Environ Sci Technol* **2004**, *38* (3), 799–807. https://doi.org/10.1021/es0345569.

(33) Klupinski, T. P. Abiotic Degradation of Pentachloronitrobenzene by Fe (II): Reactions on Goethite and Iron Oxide Nanoparticles. *Environ. Sci. Technol* **2004**, *38*, 4353–4360. https://doi.org/10.1021/es035434j.

(34) Niedźwiecka, J. B.; Drew, S. R.; Schlautman, M. A.; Millerick, K. A.; Grubbs, E.; Tharayil, N.; Finneran, K. T. Iron and Electron Shuttle Mediated (Bio)Degradation of 2,4-Dinitroanisole (DNAN). *Environ Sci Technol* **2017**, *51* (18), 10729–10735. https://doi.org/10.1021/acs.est.7b02433.

(35) Khatiwada, R.; Root, R. A.; Abrell, L.; Sierra-Alvarez, R.; Field, J. A.; Chorover, J. Abiotic Reduction of Insensitive Munition Compounds by Sulfate Green Rust. *Environ. Chem.* **2018**, *15* (5), 259–266. https://doi.org/10.1071/EN17221.

(36) Cárdenas-Hernández, P. A.; Anderson, K. A.; Murillo-Gelvez, J.; Di Toro, D. M.; Allen, H. E.; Carbonaro, R. F.; Chiu, P. C. Reduction of 3-Nitro-1,2,4-Triazol-5-One (NTO) by the Hematite-Aqueous Fe(II) Redox Couple. *Environ Sci Technol* **2020**, *54* (19), 12191–12201. https://doi.org/10.1021/acs.est.0c03872.

(37) Niedźwiecka, J. B.; McGee, K.; Finneran, K. T. Combined Biotic-Abiotic 2,4-Dinitroanisole Degradation in the Presence of Hexahydro-1,3,5-Trinitro-1,3,5-Triazine. *Environ Sci Technol* **2020**, *54* (17), 10638–10645. https://doi.org/10.1021/acs.est.0c02363.

(38) Menezes, O.; Yu, Y.; Root, R. A.; Gavazza, S.; Chorover, J.; Sierra-Alvarez, R.; Field, J. A. Iron(II) Monosulfide (FeS) Minerals Reductively Transform the Insensitive Munitions

Compounds 2,4-Dinitroanisole (DNAN) and 3-Nitro-1,2,4-Triazol-5-One (NTO). *Chemosphere* **2021**, *285*, 131409. https://doi.org/10.1016/j.chemosphere.2021.131409.

(39) Yu, H. Y.; Wang, Y. kui; Chen, P. cheng; Li, F. bai; Chen, M. jia; Hu, M.; Ouyang, X. Effect of Nitrate Addition on Reductive Transformation of Pentachlorophenol in Paddy Soil in Relation to Iron(III) Reduction. *J Environ Manage* **2014**, *132*, 42–48. https://doi.org/10.1016/j.jenvman.2013.10.020.

(40) Kruse, T.; Maillard, J.; Goodwin, L.; Woyke, T.; Teshima, H.; Bruce, D.; Detter, C.; Tapia, R.; Han, C.; Huntemann, M.; Wei, C. L.; Han, J.; Chen, A.; Kyrpides, N.; Szeto, E.; Markowitz, V.; Ivanova, N.; Pagani, I.; Pati, A.; Pitluck, S.; Nolan, M.; Holliger, C.; Smidt, H. Complete Genome Sequence of *Dehalobacter restrictus* PER-K23T. *Stand Genomic Sci* **2013**, *8* (3), 375-388. https://doi.org/10.4056/sigs.3787426.

(41) Molenda, O.; Puentes Jácome, L. A.; Cao, X.; Nesbø, C. L.; Tang, S.; Morson, N.; Patron, J.; Lomheim, L.; Wishart, D. S.; Edwards, E. A. Insights into Origins and Function of the Unexplored Majority of the Reductive Dehalogenase Gene Family as a Result of Genome Assembly and Ortholog Group Classification. *Environ Sci Process Impacts* **2020**, *22* (3), 663–678. https://doi.org/10.1039/c9em00605b.

(42) Bulka, O.; Webb, J.; Dworatzek, S.; Mahadevan, R.; Edwards, E. A. A Multifunctional *Dehalobacter*? Tandem Chloroform and Dichloromethane Degradation in a Mixed Microbial Culture. *Environ Sci Technol* **2023**, *57* (48), 19912–19920. https://doi.org/10.1021/acs.est.3c06686.

(43) Tang, S.; Gong, Y.; Edwards, E. A. Semi-Automatic in Silico Gap Closure Enabled de Novo Assembly of Two *Dehalobacter* Genomes from Metagenomic Data. *PLoS One* **2012**, *7* (12), e52038. https://doi.org/10.1371/journal.pone.0052038.

(44) Tang, S.; Wang, P. H.; Higgins, S. A.; Löffler, F. E.; Edwards, E. A. Sister *Dehalobacter* Genomes Reveal Specialization in Organohalide Respiration and Recent Strain Differentiation Likely Driven by Chlorinated Substrates. *Front Microbiol* **2016**, *7*, 100. https://doi.org/10.3389/fmicb.2016.00100.

(45) Nelson, J. L.; Fung, J. M.; Cadillo-Quiroz, H.; Cheng, X.; Zinder, S. H. A Role for *Dehalobacter* spp. in the Reductive Dehalogenation of Dichlorobenzenes and Monochlorobenzene. *Environ Sci Technol* **2011**, *45* (16), 6806–6813. https://doi.org/10.1021/es200480k.

(46) Wang, S.; Araujo, S. P.; Lomheim, L.; Mack, E. E.; Edwards, E. A.; Passport, E. *Dehalobacter converts 2,3-dichloroaniline to aniline via 2-chloroaniline in a methanogenic enrichment culture*; 2025. Manuscript in preparation (preprint on *BioRxiv*).

(47) Cole, J. R.; Cascarelli, A. L.; Mohn, W. W.; Tiedjel, J. M. Isolation and Characterization of a Novel Bacterium Growing via Reductive Dehalogenation of 2-Chlorophenol. *Appl Environ Microbiol* **1994**, *60* (10), 3536–3542.

(48) Sanford, R. A.; Cole, J. R.; Tiedje, J. M. Characterization and Description of *Anaeromyxobacter dehalogenans* gen. nov., sp. nov., an Aryl-Halorespiring Facultative Anaerobic Myxobacterium. *Appl Environ Microbiol* **2002**, *68* (2), 893–900. https://doi.org/10.1128/AEM.68.2.893-900.2002.

(49) Thomas, S. H.; Wagner, R. D.; Arakaki, A. K.; Skolnick, J.; Kirby, J. R.; Shimkets, L. J.; Sanford, R. A.; Löffler, F. E. The Mosaic Genome of *Anaeromyxobacter dehalogenans* strain 2CP-C Suggests an Aerobic Common Ancestor to the Delta-Proteobacteria. *PLoS One* **2008**, *3* (5), e2103. https://doi.org/10.1371/journal.pone.0002103.

(50) Fincker, M.; Spormann, A. M. Biochemistry of Catabolic Reductive Dehalogenation. *Annu Rev Biochem* **2017**, *86*, 357–386. https://doi.org/10.1146/annurev-biochem.

(51) Qian, Y.; Xu, M.; Deng, T.; Hu, W.; He, Z.; Yang, X.; Wang, B.; Song, D.; Chen, L.; Huang, Y.; Sun, G. Synergistic Interactions of *Desulfovibrio* and *Petrimonas* for Sulfate-Reduction Coupling Polycyclic Aromatic Hydrocarbon Degradation. *J Hazard Mater* **2021**, *407*, 124385. https://doi.org/10.1016/j.jhazmat.2020.124385.

(52) Manchester, M. J.; Hug, L. A.; Zarek, M.; Zila, A.; Edwards, E. A. Discovery of a Trans-Dichloroethene-Respiring *Dehalogenimonas* Species in the 1,1,2,2-Tetrachloroethane-Dechlorinating WBC-2 Consortium. *Appl Environ Microbiol* **2012**, *78* (15), 5280–5287. https://doi.org/10.1128/AEM.00384-12.

(53) Kato, S.; Yumoto, I.; Kamagata, Y. Isolation of Acetogenic Bacteria That Induce Biocorrosion by Utilizing Metallic Iron as the Sole Electron Donor. *Appl Environ Microbiol* **2015**, *81* (1), 67–73. https://doi.org/10.1128/AEM.02767-14.

(54) Ise, K.; Suto, K.; Inoue, C. Microbial Diversity and Changes in the Distribution of Dehalogenase Genes during Dechlorination with Different Concentrations of Cis -DCE. *Environ Sci Technol* **2011**, *45* (12), 5339–5345. https://doi.org/10.1021/es104199y.

(55) Menezes, O.; Kocaman, K.; Wong, S.; Rios-Valenciana, E. E.; Baker, E. J.; Hatt, J. K.; Zhao, J.; Madeira, C. L.; Krzmarzick, M. J.; Spain, J. C.; Sierra-Alvarez, R.; Konstantinidis, K. T.; Field, J. A. Quinone Moieties Link the Microbial Respiration of Natural Organic Matter to the Chemical Reduction of Diverse Nitroaromatic Compounds. *Environ Sci Technol* **2022**, *56* (13), 9387–9397. https://doi.org/10.1021/acs.est.2c01329.

(56) Murillo-Gelvez, J.; Di Toro, D. M.; Allen, H. E.; Carbonaro, R. F.; Chiu, P. C. Reductive Transformation of 3-Nitro-1,2,4-Triazol-5-One (NTO) by Leonardite Humic Acid and Anthraquinone-2,6-Disulfonate (AQDS). *Environ Sci Technol* **2021**, *55* (19), 12973–12983. https://doi.org/10.1021/acs.est.1c03333.

(57) Ratasuk, N.; Nanny, M. A. Characterization and Quantification of Reversible Redox Sites in Humic Substances. *Environ Sci Technol* **2007**, *41* (22), 7844–7850. https://doi.org/10.1021/es071389u.

(58) Uchimiya, M.; Stone, A. T. Reversible Redox Chemistry of Quinones: Impact on Biogeochemical Cycles. *Chemosphere* **2009**, *77* (4), 451–458. https://doi.org/10.1016/j.chemosphere.2009.07.025.