Transposons accelerate chromosomal speciation by centromere expansion

and chromosome fission

Jui-Hung Tai [1,2,3,†], Tzi-Yuan Wang [3,†], Gwo-Chin Ma[4], Yu-Wei Wu[5,6,7], Tsung-Han Yu[8],

Chen-Fan Wang[9], Te-Yu Liao[10], Shih-Pin Huang[3], Feng-Yu Wang[11], Yohey Terai[12], Tsung-

Han Wu[1], Chun-Hua Hsu[1,13], Hon-Tsen Yu[14], Kwang-Tsao Shao[3,15*], Shu-Miaw Chaw [3*],

Hurng-Yi Wang [2,8,16,17*]

[1] Genome and Systems Biology Degree Program, National Taiwan University and Academia

Sinica, Taipei, Taiwan

[2] Graduate Institute of Clinical Medicine, College of Medicine, National Taiwan University,

Taipei, Taiwan

[3] Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

[4] Department of Genomic Medicine, Changhua Christian Hospital, Changhua, Taiwan.

[5] Graduate Institute of Biomedical Informatics, College of Medical Science and Technology,

Taipei Medical University, Taipei, Taiwan.

[6] Clinical Big Data Research Center, Taipei Medical University Hospital, Taipei, Taiwan

[7] TMU Research Center for Digestive Medicine, Taipei Medical University, Taipei, Taiwan

[8] Institute of Ecology and Evolutionary Biology, National Taiwan University, Taipei,

Taiwan

[9] Ming Der Municipal High School, New Taipei City, Taiwan

[10] Department of Oceanography, National Sun Yat-sen University, Kaohsiung, Taiwan

[11] Taiwan Ocean Research Institute, National Applied Research Laboratories, Kaohsiung,

Taiwan.

[12] Research Center for Integrative Evolutionary Science, The Graduate University for

Advanced Studies, SOKENDAI, Hayama, Kanagawa, Japan

[13] Department of Agricultural Chemistry, National Taiwan University, Taipei, Taiwan

[14] Department of Life Science, National Taiwan University, Taipei, Taiwan

[15] Institute of Marine Biology, National Taiwan Ocean University, Keelung, Taiwan

[16] Graduate Institute of Medical Genomics and Proteomics, College of Medicine, National

Taiwan University, Taipei, Taiwan

[17] Department of Entomology, National Taiwan University, Taipei, Taiwan

[†] These authors contribute equally to this work.

[*] Corresponding authors. Email: KTS, zoskt@gate.sinica.edu.tw; SMC,

smchaw@sinica.edu.tw; HYW, hurngyi@ntu.edu.tw

Abstract

Chromosomal speciation poses a paradox: if rearrangements cause postmating isolation,

how do they persist and spread? We propose that centromere remodeling, driven by

transposable element accumulation, provides a mechanistic solution. In *Opsariichthys*, a

hyper-diverse lineage of Asian river chub with 2N = 76–78 chromosomes, we identify ~15

fission events coinciding with TE-fueled centromeric DNA expansion, potentially linked to a

mutation in the PIWI1 protein. These expansions enable neokinetochore formation, allowing

fissioned chromosomes to segregate faithfully, while subsequent rearrangements disrupt

pairing and reduce gene flow. Comparative genomics reveals that fissioned chromosomes

exhibit lower migration and greater divergence, acting as reproductive barriers. Our findings

show that TE-mediated kinetochore duplication can facilitate extensive chromosome fission

without meiotic disruption, providing a viable and previously underappreciated path to rapid

speciation.

Main Text

Chromosomal speciation model posits that fissions generate reproductive barriers that drive speciation (*1, 2*). In monocentric species, however, fission can produce acentric fragments or centromere imbalance, leading to meiotic failure (*3*). It remains unclear how chromosome fission occurs and contributes to reproductive isolation, despite the occasional occurrence of massive fission events in vertebrate evolution (*4*). One proposed solution is neokinetochore formation (*5, 6*): during gametogenesis, it can generate dicentric chromatids, and if these achieve bipolar attachment in meiosis, fission may yield two chromosomes with functional kinetochores and minimal phenotypic effect. Subsequent pericentric inversion may complicate pairing between fissioned and intact homologs, thus creating barriers to gene flow and driving speciation. While theoretically compelling, this model lacks strong empirical support (*7, 8*), and the mechanism triggering neokinetochore formation remains unknown.

Transposable elements (TEs) have recurrently shaped genomic variation and host diversity throughout evolution (*9*). Their activity drives chromosomal rearrangements—including fusions, fissions, translocations, and inversions—that can fuel evolutionary innovation and speciation (*10, 11*). We propose that active TE accumulation triggers mutagenic processes such as unequal exchange and replication slippage in the centromere (*12*), promoting the expansion of centromeric DNA and initiating neokinetochore formation.

The Opsariichthyini tribe (Asian river chub) (Cypriniformes: Xenocyprididae) includes small riverine minnows broadly distributed across East Asia, from Russia to northern Vietnam (*13*). This monophyletic group contains multiple genera, ranging from early-diverging lineages such as *Parazacco*, *Candidia*, and *Nipponocypris* to more recently diverged genera such as *Zacco* and *Opsariichthys*. Most species in the tribe have 2N = 48 chromosomes, except *Opsariichthys*, which exhibits a strikingly elevated count of 2N = 74 - 78 (*14, 15*). Interestingly, hybrids between *Zacco* and *Opsariichthys,* as well as between different *Opsariichthys* species, have occasionally been seen in natural environments where they are sympatric (*16, 17*). Despite the increase in chromosome number, overall DNA content remains conserved, implicating extensive chromosome fission rather than genome expansion as the primary driver of karyotypic change (*18*). Notably, *Opsariichthys* contains over 15 recognized species (*19*), markedly more than the typical three to four species found in other genera in the tribe (*20*). This exceptional diversity, alongside its tolerance to widespread fission, makes *Opsariichthys* a compelling system for testing chromosomal speciation models.

To investigate the relationship between chromosome fission and TE activity in
Opsariichthys, and to assess whether fission contributes to accelerated speciation, we
sequenced and annotated the genomes of *Candidia barbatus* (Cb), *Zacco platypus* (Zp), *O.*
pachycephalus (Op), and *O. evolans* (Oe), each sampled from multiple populations. In
parallel, we analyzed the karyotypes of these species to enable integrated analysis of
chromosomal variation and fission patterns. For comparison, the genome and karyotype of *O.*
bidens (Ob) were included in subsequent analyses based on published data (*21*). These five
species capture the genomic and ecological diversity within the Opsariichthyini tribe. Our
results suggest that chromosome fissions were initiated by lineage-specific TE activity, likely
beginning in the common ancestor of Opsariichthyini. Fissioned chromosomes showed
reduced migration and increased divergence relative to intact chromosomes, indicating their
role in impeding gene flow and facilitating speciation.

Karyotyping, genome assembly and annotation
Karyotypes were determined by examining multiple individuals from each species. No
heteromorphic elements were detected between males and females. Cb and Zp both possess
2N = 48 chromosomes, while Op and Oe have 2N = 76 and 78, respectively (Fig. 1, fig. S1
and table S1). Despite the wide variation in chromosome number, the fundamental numbers
are comparable across species. Genome assembly details are given in Supplementary Text
(Genome assembly statistics; see also fig. S2, table S2-S6). Chromosome-level assemblies
are consistent with karyotyping results and exhibit high completeness, with BUSCO scores
exceeding 96% for all species (table S4). Transposable elements (TEs) account for 51% to
55% of the opsariichthyin genomes. Notably, the Opsariichthyini show a higher proportion of
long interspersed nuclear elements (LINEs), particularly L2, and a significant increase in
Gypsy within their long terminal repeats (LTRs) content compared to other cyprinids (table
S5).

Chromosomal evolution is associated with recent TE activities
Synteny analysis revealed that chromosome numbers in *Opsariichthys* increased
dramatically from N = 24 to 38 or 39, driven by 14 to 15 chromosome fission events (Fig. 2).
We identified 'breaking regions' in Cb and Zp, where segments of ancestral chromosome no
longer align with their fissioned homologs (fig. S3), likely due to structural changes or
sequence loss during fission. These breaking regions are enriched with Gypsy and L2, with

Zp showing a higher Gypsy density than Cb (fig. S4A). This enrichment extends to the

homologous regions in the fissioned chromosomes of Op, Oe, and Ob (fig. S3).

Repeat landscapes revealed recent expansions of L2 and Gypsy (k-distance < 5%; young

TEs) across all examined opsariichthyin genomes (fig. S4B). In Cb and Zp, 56% and 43% of

L2 insertions, respectively, are recent, increasing to 75% and 72% within breaking regions

(fig. S4C). Recent Gypsy expansion is more pronounced in Zp, increasing from 17%

genome-wide to 31% in breaking regions, whereas in Cb, only 10% of Gypsy are recently

inserted in both genomic and breaking regions. Furthermore, 14 of the 15 breaking regions in

Cb show recent L2 expansion and 8 show recent Gypsy expansion (table S7; fig. S5), while

all breaking regions in Zp exhibit recent expansion of both elements (table S8; fig. S6). These

observations suggest that L2 and Gypsy expansion occurred asynchronously, with L2

expanding before Gypsy.

Chromosome fission within centromere regions

To investigate the mechanism of chromosome fission, we identified putative centromere

regions by searching for regions containing centromeric DNA, low recombination rates, and

enriched with L2 and Gypsy (Supplementary Text: Centromere identification; fig. S5-S10).

In general, karyotyping results derived *in silico* analysis and microscopic observations are

largely consistent (Fig. 1 and table S1), demonstrating that the putative centromeres identified

are authentic. In Cb, 10 of 15 breaking regions overlapped with putative centromeres (fig. S5;

table S7), while in Zp, all 15 coincided (fig. S6; table S8). The difference is primarily due to

chromosomal rearrangements, especially pericentric inversions, that occurred during

evolution (fig. S11). Recent L2 expansions were observed in centromeres across all species.

Gypsy expansions were prevalent in nearly all centromeres of Zp, Op, Oe, and Ob, but

occurred in only 14 of Cb's 24 centromeres (table S7-S11). These patterns, consistent with

breaking regions, support earlier L2 expansion than Gypsy.

Notably, in chromosomes that later underwent fission in *Opsariichthys*, the

corresponding regions in Zp already exhibit significantly greater centromeric DNA content

and broader centromeric spans than in Cb (Fig. 3A). While the number of L2 in the

centromere regions is similar among species, that of Gypsy is significantly greater in Zp (Fig.

3B). These findings suggest that recent Gypsy expansion may have promoted both

centromeric DNA duplication and centromere expansion through unequal exchange and

replication slippage. Together, these processes could facilitate neokinetochore formation,

leading to dicentric chromosomes (*6*), in which each fissioned product retains a functional
kinetochore for proper segregation.

Insertion preferences and interaction between L2 and Gypsy

The expansion of young L2 in putative centromeres suggests that the reduced
recombination rates in these regions limit transposon removal, enabling their accumulation
over time (*22*). This is evidenced by a negative correlation between young L2 density and
recombination rates in Zp (r = -0.44) and Cb (r = -0.25) (fig. S12A). Because centromeres
can shift over time (fig. S11), ancient L2 (k-distance > 5%) show reduced or no correlation
with recombination rate (Zp: r = –0.11; Cb: n.s.) (fig. S12B).

The expansion of L2 preceding that of Gypsy suggests a potential interdependence (fig.
S13), with expansion of Gypsy possibly relying on a genomic context shaped by L2. LTR
insertion is typically marked by a 5-bp target site duplication (TSD) flanked by palindromic
motifs generated during integration (*23*). Using LTR_finder and LTR_retriever, we identified
a conserved palindromic motif flanking the TSD in opsariichthyin species: TNACA-TSD-
TGTNA (fig. S14A). Similar motifs are present in L2 (fig. S14B), suggesting that each L2
replication and transposition event creates a new target site for subsequent Gypsy integration.

Chromosome fission decrease gene flow and increase genetic divergence

According to the chromosome speciation model, chromosomal rearrangements can lead
to the formation of a reproductive barrier that promotes speciation. Divergence at loci on
rearranged chromosomes is also expected to exceed that on non-rearranged chromosomes. To
test this, we used gIMble (*24*) to calculate the blockwise site frequency spectrum (bSFS)
between Zp and Oe, as well as between Zp and Op. The isolation with migration (IM) model
was the best fit (table S12). Migration near the breaking regions was notably reduced (Fig.
4A and fig. S15-S16), and fission chromosomes exhibited significantly lower migration rates
than non-fission chromosomes in both comparisons ($p < 0.05$) (Fig. 4B).

We next calculated the genetic divergence using single-copy genes. Chromosomes that
underwent fission in *Opsariichthys* showed significantly greater genetic distances than non-
fission chromosomes in all comparisons with Zp ($p < 10^{-6}$) (Fig. 4C). By contrast, when
comparing Cb and Zp, where both sets of chromosomes retained the ancestral, unfissioned
state, no difference in divergence was detected between the two chromosome sets.

It is generally believed that fission does not reduce recombination. Indeed, fissioned
chromosomes in both Op and Oe exhibited higher average recombination rates than non-

fissioned chromosomes (fig. S17). In Zp, the homologous chromosomes showed comparable

recombination rates across both groups.

Mutations in PIWI1 protein associates with L2 expansion and species diversification

While the recent Gypsy expansion occurs in multiple cyprinids (fig. S18A), recent L2

expansion appears unique to the Opsariichthyini, suggesting possible shifts in transposon

silencing. Given the critical role of PIWI1 in transposon repression (*25*), we compared PIWI1

sequences between opsariichthyins and other cyprinids (Supplementary Text: Comparative

Analysis of Piwi1 Protein). All opsariichthyins share two unique amino acid substitutions,

G348S and V380I, in the PAZ domain, a region essential for piRNA binding and TE

silencing (fig. S18C). V380I is particularly notable, as it lies adjacent to the nucleic acid-

binding interface at residue 379 (*26*).

Repeat landscape analysis shows that cyprinids with the ancestral V380 lack L2

expansion (fig. S18A), while *Triplophysa* (Nemacheilidae), a cyprinid outgroup carrying

V380I, has undergone recent L2 expansion (fig. S18B). Structural modeling using AlphaFold

3 revealed that residue 380 lies within the piRNA binding pocket of the PAZ domain (fig.

S19A). Although it does not directly contact the binding interface (fig. S19B, upper panel),

substitution of valine with the bulkier and more hydrophobic isoleucine (V380I) may subtly

alter local conformation (fig. S19B, lower panel), potentially impairing piRNA loading or

destabilizing the complex (*27, 28*). Given the phylogenetic distance between Opsariichthyini

and *Triplophysa*, the parallel emergence of both the V380I mutation and L2 expansion

suggests convergent evolution. These observations point to a shared mechanism, whereby

V380I interferes with PIWI1 function, diminishing transposon silencing and enabling L2

proliferation.

In *Triplophysa*, chromosome numbers remain stable (2N = 48–50), yet karyotypes and

fundamental numbers vary substantially (*29*), coinciding with L2 expansion (fig. S18).

Notably, *Triplophysa* comprises 147 described species, far exceeding the family average of

16, suggesting a possible link between TE activity and elevated species diversification.

Together, these findings implicate the V380I mutation in reshaping genome architecture and

possibly contributing to lineage-specific diversification.

Discussion

Transposon-fueled centromere remodeling sets the stage for chromosome fission

Through comparative analysis of opsariichthyin genomes and interspecific gene flow,
we provide direct evidence linking TE activity to chromosomal evolution and species
diversification (Fig. 5A). L2 expansion is specific to Opsariichthyini within Cyprinoidei (Fig.
3B and fig. S18A) and is enriched in putative centromeres (Fig. 5B), where reduced
recombination likely facilitates accumulation over time (22). L2 elements also harbor
sequence motifs preferentially targeted by Gypsy, guiding its insertion into centromeric
regions (fig. S14B). Moreover, LTRs frequently insert into the 3' UTRs of other LTRs,
especially Gypsy (30) , creating a positive feedback loop that concentrates both elements in
centromeres.

Active TE accumulation may trigger mutagenic processes, such as unequal exchange
and replication slippage in centromere(12), leading to further expansion of repetitive
sequences including centromeric DNA. This is evident in Zp, where chromosomes that
underwent fission after *Opsariichthys* diverged from *Zacco* contain significantly more
centromeric DNA and exhibit centromere expansion compared to homologs in Cb (Fig. 5C),
likely due to a recent Gypsy expansion (Fig. 3). According to kinetochore reproduction
theory, expanded centromeric DNA increases the likelihood of neokinetochore formation (5,
6) (Fig. 5D). In the presence of kinetochores, dicentric chromosomes may arise. During
replication, chromatids with monopolar spindle attachment segregate normally, while those
with opposing bipolar attachment undergo fission (Fig. 5D and Fig 1B of ref 5).

Of the 15 identified breakpoints, 10 coincide with putative centromeres in Cb (fig. S5;
table S7), while the remaining five, initially non-centromeric in Cb, became centromeres in
Zp via pericentric inversions (fig. S11), resulting in all 15 regions perfectly match
centromeres in Zp (fig. S6; table S8). Notably, similar associations between active TEs and
chromosomal breakpoints have been observed in primate evolution (Supplementary Text:
TEs in the breaking region of macaque chromosome 7).

247 How fissioned chromosomes escape meiotic disruption and drive speciation
Following fission, accurate pairing of homologous chromosomes is required for proper
segregation during meiosis. Centromeres have traditionally been considered essential not
only for segregation but also for homologous pairing (3), raising concerns that new
centromeres might disrupt this process. However, studies in wheat and *Arabidopsis* suggest
centromere may not play a direct role in meiotic pairing (31, 32). Instead, chromosome
pairing initiates at both ends and extends toward the center during early meiosis, indicating
that pairing is end-driven rather than centromere-dependent (33). This provides a mechanistic

explanation for how massive chromosome fission, as observed in *Opsariichthys*, can occur
without disrupting meiotic pairing or segregation (Fig. 5E, left). Furthermore, selection may
favor the retention of fissioned over intact homologs, especially if smaller chromosomes are
more likely than larger ones to segregate without error during cell division (*7, 34*). As a result,
fission karyotypes can rapidly increase in frequency within local populations without
deleterious effects. For further evidence that frequent fission can proceed without deleterious
effects in Opsariichthys, see Supplementary Text (Frequent fission and fusion in
Opsariichthys genomes).

At later stages, additional rearrangement, such as pericentric inversion, common in the
Opsariichthini (fig. S11), can relocate centromeres (Fig. 5E, right). Such relocation may
interfere with pairing between fissioned and unfissioned homologs, thereby impeding gene
flow, promoting genetic isolation, and ultimately facilitating speciation (Fig. 5F). Supporting
this, gene flow between *Zacco* and *Opsariichthys* is reduced at chromosomal breakpoints,
with fissioned chromosomes showing significantly lower migration estimates and greater
genetic divergence than un-fissioned ones (Fig. 4).

Taken together, these findings invite a fresh perspective on the chromosomal speciation
model, which traditionally posits that rearrangements such as fissions promote speciation by
impeding gene flow, yet at the cost of meiotic dysfunction due to centromere imbalance.
However, perhaps the paradox lies in the premise itself: the longstanding notion that
chromosome rearrangements are inherently deleterious may overlook mechanisms that allow
them to persist. Our results suggest that kinetochore duplication and end-driven pairing
provide a viable path for fissioned chromosomes to segregate correctly and spread within
populations, quietly laying the groundwork for reproductive isolation and speciation.

Reference and Notes

1. M. J. D. White, *Modes of speciation*. A series of books in biology (W.H. Freeman and Company) (W.H. Freeman, San Francisco, 1978).

2. A. Navarro, N. H. Barton, Accumulating postzygotic isolation genes in parapatry: a new twist on chromosomal speciation. *Evolution* **57**, 447-459 (2003).

3. O. Da Ines, C. I. White, Centromere Associations in Meiotic Chromosome Pairing. *Annu. Rev. Genet.* **49**, 95-114 (2015).

4. M. A. Ferguson-Smith, V. Trifonov, Mammalian karyotype evolution. *Nat. Rev. Genet.* **8**, 950-962 (2007).

5. N. B. Todd, Karyotypic fissioning and Canid phylogeny. *J. Theor. Biol.* **26**, 445-480 (1970).

6. R. L. Kolnicki, Kinetochore reproduction in animal evolution: cell biological explanation of karyotypic fission theory. *Proc Natl Acad Sci U S A* **97**, 9493-9497 (2000).

7. L. R. Godfrey, J. C. Masters, Kinetochore reproduction theory may explain rapid chromosome evolution. *Proc Natl Acad Sci U S A* **97**, 9821-9823 (2000).

8. H. T. Imai, Y. Satta, N. Takahata, Integrative Study on Chromosome Evolution of Mammals, Ants and Wasps Based on the Minimum Interaction Theory. *J. Theor. Biol.* **210**, 475-497 (2001).

9. A. Böhne, F. Brunet, D. Galiana-Arnoux, C. Schultheis, J.-N. Volff, Transposable elements as drivers of genomic and biological diversity in vertebrates. *Chromosome Research* **16**, 203-215 (2008).

10. S. J. Klein, R. J. O'Neill, Transposable elements: genome innovation, chromosome diversity, and centromere conflict. *Chromosome Res* **26**, 5-23 (2018).

11. F. J. Ayala, M. Coluzzi, Chromosome speciation: humans, Drosophila, and mosquitoes. *Proc Natl Acad Sci U S A* **102 Suppl 1**, 6535-6542 (2005).

12. E. Balzano, S. Giunta, Centromeres under Pressure: Evolutionary Innovation in Conflict with Conserved Function. *Genes (Basel)* **11**, (2020).

13. H.-Y. Wang, C.-F. Wang, S.-Y. Du, S.-C. Lee, New insights on molecular systematics of opsariichthines based on cytochrome b sequencing. *J. Fish Biol.* **71**, 18-32 (2007).

14. X. Yu, T. Zhou, K. Li, Y. Li, M. Zhou, On the karyosystematics of cyprinid fishes and a summary of fish chromosome studies in China. *Genetica* **72**, 225-235 (1987).

15. Y. Ojima, M. Hayashi, K. Ueno, Cytogenetic studies in lower vertebrates. X. Karyotype and DNA studies in 15 species of Japanese Cyprinidae. *The Japanese journal of genetics* **47**, 431-440 (1972).

16. N. L. Liao, S. P. Huang, T. Y. Wang, Interspecific Mating Behavior Between Introduced Zacco platypus and Native Opsariichthys evolans in Taiwan. *Zool Stud* **59**, e6 (2020).

17. H.-Y. Wang, S.-C. Lee, M.-J. Yu, Genetic evidence to clarify the systematic status of
the genera Zacco and Candidia. *Zool. Stud.* **36**, 170-177 (1997).

18. A. Takai, Karyotype and Chromosomal Distribution of C-band-Positive Constitutive
Heterochromatin in Opsariichthys uncirostris (Cyprinidae, Pisces). *Memoirs of Osaka*
Shin-ai College **53**, 1-5 (2019).

19. X. Peng, J.-J. Zhou, H.-D. Gao, J.-Q. Yang, A new species of Opsariichthys
(Teleostei, Cypriniformes, Xenocyprididae) from Southeast China. *ZooKeys* **1214**,
(2024).

20. R. Froese, D. Pauly, *FishBase*. (WorldFish, 2024), vol. 1594.

21. X. Xu *et al.*, Chromosome-Level Assembly of the Chinese Hooksnout Carp
(Opsariichthys bidens) Genome Using PacBio Sequencing and Hi-C Technology.
Frontiers in Genetics **12**, (2022).

22. P. B. Talbert, S. Henikoff, What makes a centromere? *Exp. Cell Res.* **389**, 111895
(2020).

23. R. S. Linheiro, C. M. Bergman, Whole genome resequencing reveals natural target
site preferences of transposable elements in Drosophila melanogaster. *PLoS One* **7**,
e30008 (2012).

24. D. R. Laetsch *et al.*, Demographically explicit scans for barriers to gene flow using
gIMble. *PLOS Genetics* **19**, e1010999 (2023).

25. S. Houwing *et al.*, A role for Piwi and piRNAs in germ cell maintenance and
transposon silencing in Zebrafish. *Cell* **129**, 69-82 (2007).

26. J. B. Ma, K. Ye, D. J. Patel, Structural basis for overhang-specific small interfering
RNA recognition by the PAZ domain. *Nature* **429**, 318-322 (2004).

27. Y. Tian, D. K. Simanshu, J.-B. Ma, D. J. Patel, Structural basis for piRNA 2'-O-
methylated 3'-end recognition by Piwi PAZ (Piwi/Argonaute/Zwille) domains.
Proceedings of the National Academy of Sciences **108**, 903-910 (2011).

28. L.-p. Dang *et al.*, Investigating structural impact of a valine to isoleucine substitution
on anti-Müllerian hormone in silico and genetic association of the variant and AMH
expression with egg production in chickens. *Journal of Integrative Agriculture* **19**,
1635-1643 (2020).

29. Y. Niu, Q. Zhao, H. Zhao, A. Ludwig, Z. Peng, Karyotype and genome size of an
endangered cavefish (Triplophysa rosa Chen & Yang, 2005). *J. Appl. Ichthyol.* **33**,
124-126 (2017).

30. P. Jedlicka, M. Lexa, I. Vanat, R. Hobza, E. Kejnovsky, Nested plant LTR
retrotransposons target specific regions of other elements, while all LTR
retrotransposons often target palindromes and nucleosome-occupied regions: in silico
study. *Mob DNA* **10**, 50 (2019).

31. O. Da Ines, K. Abe, C. Goubely, M. E. Gallego, C. I. White, Differing requirements
for RAD51 and DMC1 in meiotic pairing of centromeres and chromosome arms in
Arabidopsis thaliana. *PLoS Genet.* **8**, e1002636 (2012).
32. E. Corredor, A. J. Lukaszewski, P. Pachón, D. C. Allen, T. s. Naranjo, Terminal
Regions of Wheat Chromosomes Select Their Pairing Partners in Meiosis. *Genetics*
177, 699-706 (2007).
33. H. You *et al.*, Chromosome ends initiate homologous chromosome pairing during rice
meiosis. *Plant Physiol.* **195**, 2617-2634 (2024).
34. I. Schubert, J. L. Oud, There Is an Upper Limit of Chromosome Size for Normal
Development of an Organism. *Cell* **88**, 515-520 (1997).
364
365

Acknowledgements

The authors wish to express their gratitude to Professor Chi-Chang Liu for providing valuable samples. Additionally, we thank the NGS High Throughput Genomics Core at the Biodiversity Research Center, Academia Sinica, for their support in library preparation and Hi-C sequencing. We are deeply grateful to Professor Shau-Ping Lin and Professor Isheng Jason Tsai for their invaluable advice throughout this research.

Funding

This study was supported by grants from the National Science and Technology Council (NSTC), Taiwan (113-2327-B-002-003-, MOST 109-2311-B-002-023-MY3, MOST 105-2311-B-001-064, MOST 106-2311-B-001-022, and MOST 107-2311-B-001-007), and National Taiwan University (113L7223). This study was also facilitated by the National Key Area International Cooperation Alliance: University Academic Alliance in Taiwan (UAAT) - Kyushu-Okinawa Open University (KOOU) - Medicine and Life Sciences Integrative Program. Supported by the Ministry of Education, Taiwan, the program fosters international collaboration in cutting-edge research.

Author contributions

Conceived the idea: TYW, KTS, SMC, HYW

Collected samples: JHT, TYW, GCM, THY, CFW, TYL, SPH, FYW, YT

Performed the experiment: TYW, GCM, THW, CHH, HTY, YT

Performed the analysis: JHT, TYW, CFW, YWW, THY

Supervised the project and secured the fundings: KTS, SMC, HYW

Wrote the manuscript: JHT, TYW, THY, KTS, SMC, HYW

Competing interests

The authors declare no competing interests.

Data and materials availability

All data in this study has been deposited to NCBI bioproject under accession PRJNA1159521 (Cb), PRJNA1160456 (Zp), PRJNA1192024 (Op), PRJNA1192752 (Oe) and PRJNA1203011 (Hybrids). The details were listed in Supplementary Tables and Supplementary Information.

Supplementary Materials

Materials and Methods

Supplementary Text

402   Figs. S1 to S21

403   Tables S1 to S13

404   References (35–85)

Fig. 1. Karyotypes of opsariichthyins examined.

A, Cb (*Candidia barbatus*) **B,** Zp (*Zacco platypus*) **C,** Op (*Opsariichthys pachycephalus*) and **D,** Oe (*O. evolans*). m: metacentric; sm: submetacentric; st: sub-telocentric; t: telocentric.

Fig. 2. The phylogeny (left panel) and syntenic relationships (right panel) among Opsariichthyini tribe species and the outgroups.
Dark black branches represent lineages of the Opsariichthyini tribe. The numbers represent the chromosome number (N). Drer: *Danio rerio*; Mamb: *Megalobrama amblycephala*; Cb: *Candidia barbatus*; Zp: *Zacco platypus*; Oe: *Opsariichthys evolans*; Op: *O. pachycephalus*; Ob: *O. bidens*

Fig. 3. The distribution of centromeric DNA, L2, and Gypsy within putative

centromeres of Cb and Zp in fissioned chromosomes.

A, The left panel displays the regions spanned by centromere markers, and the right panel

shows their counts in fissioned chromosomes. **B,** The distribution of L2 (left panel) and

Gypsy (right panel) around centromeres in fissioned chromosomes. X-axis represents

distance to the center of putative centromere regions.

** $p < 0.01$

Fig. 4. Gene flow was reduced in fissioned chromosomes.

A, Migration rate (Me) between Zp (*Z. platypus*) and Oe (*Opsariichthys evolans*) across various chromosomes, exemplified here by chromosome 10 of Zp, with additional data for other chromosomes presented in fig. S8-S9. Red dash line indicates breaking regions. **B,** The average Me estimates for fission and non-fission chromosomes between Zp and Oe (left panel) and between Zp and Op (right panel). **C,** Average genetic distance of single copy genes on fission and un-fissioned chromosomes between Zp and each of Cb, Oe, Ob, and Op. * p < 0.05; *** p < 10-6, NS: not significant.

Fig. 5. Transposable element expansion facilitates extensive chromosome fissions leading to species diversification

A, Under normal conditions (inset at the top right), the PIWI1 protein suppresses TE activity. However, mutations in the PAZ domain of the *piwi1* gene in the ancestor of the Opsariichthyini enabled the expansion of L2 within the genome. **B,** Reduced recombination within centromeres likely contributed to the accumulation of L2. **C,** L2 elements harbor sequence motifs preferentially for Gypsy, guiding its insertion into these motifs within centromeric regions and resulting in centromere expansion. **D,** Centromere expansion may facilitate neokinetochore formation, resulting in dicentric chromosomes. During replication, dicentric chromosomes with opposing bipolar attachment undergo chromosome fission. **E,** Following fission, the distal ends of the daughter chromosomes remain intact, ensuring proper pairing, while each of the two resulting chromosomes is equipped with a kinetochore, which enables accurate segregation (left). At later stage, however, chromosome rearrangement such as pericentric inversion may contribute to centromere relocation (right). **F.** Such rearrangements can act as reproductive barriers, reducing gene flow and driving speciation. Consequently, the species diversity of *Opsariichthys* increased.