1 **The prevalence of two known pathogens of Atlantic Blue Crab *Callinectes sapidus***
2 **(Rathbun, 1896) in coastal Adriatic, Aegean sea, and Atlantic Iberian coast**
3
4

5 **Authors:**
6 Eric Schott[1]* Marina Brailo Šćepanović[2], Jasna Maršić-Lučić[3], Olivia Pares[1], Mingli Zhao[4], Jerko
7 Hrabar[3], Luka Glamuzina[2], Marijana Pećarević[2], Sanja Grđan[2], João Pedro da Silva Encarnação[5],
8 Pedro Morais[6] Ana Pešić[7], Ilija Ćetković[7], Branko Glamuzina[2]
9
10 [submitted Nov 10, 2024; reviews received March 23, 2025; revised manuscript submitted May
11 23, 2025]
12
Affiliations
[1]University of Maryland Center for Environmental Science, Institute of Marine and Environmental
Technology, 701 E Pratt St. Baltimore MD 21202, USA
[2] University of Dubrovnik, Department of Applied Ecology, Dubrovnik, Croatia
[3] Institute of Oceanography and Fisheries, Laboratory for Aquaculture, Split, Croatia
[4]Royal Veterinary College, University of London, United Kingdom
[5]Centre of Marine Sciences (CCMAR) University of Algarve, Faro, Portugal
[6]Department of Environmental Science, Policy, and Management, University of California,
Berkeley, California, USA
[7] University of Montenegro, Institute of Marine Biology, Kotor, Montenegro
* Corresponding author: Eric Schott schott@umces.edu

24
25
26 **Abstract**
27

In the Mediterranean Sea, the abundance of the invasive portunid crab, *Callinectes sapidus*, has
dramatically increased in recent years. This raises concerns about damage to ecosystems, but
also offers opportunities for exploitation of a new fishery. Newly invasive species may escape
from pathogens in their native range, may introduce new pathogens, or can become host to
endemic pathogens. Understanding these factors is important for predicting or managing natural
resources in the invaded range. This study investigated the prevalence of two pathogens common
in *C. sapidus* in its home range of North America: the reovirus CsRV1 and the protozoan
parasite *Hematodinium perezi*. In crabs collected from Aegean, Adriatic, and Atlantic waters, the
CsRV1 virus was not detected. In contrast, the parasite *H. perezi* was found in crabs from all
areas except the Aegean Sea. Sequence analysis of the *H. perezi* ITS1 gene indicated that the
strains observed are most related to genotypes already described in Europe and the

Mediterranean, and not to strains from the Americas or Asia. The arrival of new species and new
potential pathogens is ongoing through transfer of ballast water to the Atlantic and
Mediterranean. Although systems are in place to exchange or inactivate ballast water, it is
advisable to continue and expand surveillance for pathogens in introduced species, to inform
management of movement of these species between regions.
44
45 **Introduction**
46

The Atlantic blue crab *Callinectes sapidus* Rathbun,1896 (Brachyura: Portunidae) is a swimming
crab native to the western Atlantic in coastal and estuarine waters of New England (USA) to
Uruguay, including the Gulf of Mexico and the Caribbean Sea (Mancinelli *et al.,* 2021, 2017). In
recent years it has also emerged as one of the most concerning invasive species in the
Mediterranean Sea. It was first recorded (as *Neptunus pelagicus*) in Europe in 1900 (Bouvier
1901), and appears to be established on the Atlantic coast of France and England (Pezy *et al.,*
2019). Other records show the species has also moved south to the upper Iberian coast (Ribeiro
and Veríssimo 2014). In the Mediterranean Sea it was first recorded in 1949 in the Gulf of
Venice, Italy, in the northern Adriatic Sea (Giordani Soika 1951), and possibly as early as 1935
near Turkey (Enzenroß *et al.,* 1997). It has become invasive in the eastern-most and western
basins of the Mediterranean and the Black Sea (Vella *et al.,* 2023). Recently, the species was
reported in Malta and Sicily (Vella *et al.,* 2023), Sardinia (Culurgioni *et al.,* 2020), Spain
(Clavero *et al.,* 2022), the Adriatic and Aegean seas (Glamuzina *et al.,* 2023, 2021; Cilenti *et al.*,
2015; Kevrekidis *et al.,* 2023), and the Mediterranean coast of Africa (Kara and Chaoui, 2021;
Taybi and Mabrouki, 2020). Also recently, *C. sapidus* has appeared in abundance on the
southern Atlantic Coast of Portugal, and has been reported on the Atlantic coast of Morocco,
possibly reflecting an expansion through the straits of Gibraltar (Morais *et al.,* 2019; González-
Ortegón *et al.,* 2022; Lamkhalkhal *et al.,* 2024).

65
The life history of *C. sapidus* includes a 28-35 day oceanic planktonic phase, before
metamorphosis to benthic postlarvae (megalopa stage), that contributes to dispersal. Blue crabs
are relatively short-lived, with an estimated longevity of 2-3 years in some parts of the US native
range (Kahn and Hestler 2005), and a recent estimate of 4.8 years in an Italian lagoon
(Mancinelli *et al*., 2024). Therefore, in locations with consistently high crab abundance, there
must be recurring sources of larvae, either through local self recruitment or from broodstock
sources upstream in oceanic currents. For example, in the south-eastern Adriatic Sea, a strong
anti-clockwise gyre connects the coast of Italy and Croatia, while in the Aegean Sea, where blue
crabs have been abundant for decades (Holthuis 1961) currents are weak and the many islands
ensure promote connectivity and self recruitment (Estournel *et al.,* 2021). The local-recruitment
hypothesis is supported by Schubart *et al.,* (2022) who found low genetic connectivity between

different populations of *C. sapidus* in the invaded range, consistent with multiple founder
introductions.
79
The success of the blue crab in new regions is likely enhanced by its omnivory and ability to
adapt to a wide range of salinity, temperature and dissolved oxygen (Glandon *et al.,* 2019).
Success of non-native species in a new range is often also attributed to an escape from predators
and infectious diseases that suppressed their numbers in the home range (sometimes referred to
as the enemy release hypothesis) (Blakeslee *et al.,* 2009; Torchin *et al.,* 2007; Schultz *et al.,*
2019). While a lot has been reported on occurrence and the invasiveness of *C. sapidus* in the
Mediterranean, there is still little knowledge of the infections that it may acquire in this new
range. The enemy release (pathogen escape) concept assumes that newly arrived species have not
carried their home-range pathogens with them.
89
In its North American range, *C. sapidus* is infected by a number of pathogenic parasites and
viruses (Shields and Overstreet 2007). The most studied is the parasitic dinoflagellate,
Hematodinium perezi, which is known to cause mortality in *C. sapidus*, and is a host generalist
that infects a wide range of crustacean species. The type species is described in *Liocarcinus*
depurator from the Atlantic coast of Europe (Chatton and Poisson, 1931; Small et al. 2102).
Three different molecular *H. perezi* clades are recognized, each characteristic of different global
regions, and distinguished by genomic changes in the ribosomal RNA gene ITS1 region (Small
et al., 2012, Pagenkopp-Lohan *et al.,* 2012). The genotype infecting crabs in the Mediterranean
and Atlantic coast of Europe is referred at Type I, while those in Asia, often associated with
mortality in aquaculture, are type II, whereas *H. perezi* found in *C. sapidus* and other hosts on
the Atlantic estuaries of north America are type III (Small *et al*., 2012).
101
In the United States, regional population declines of *C. sapidus* have coincided with elevated
prevalence *Hematodinium* sp. (Lee and Frischer 2004). A related *Hematodinium* sp. that infects
Norway lobster *(Nephrops norvegicus)* is a known driver of mortality in this fishery on the west
coast of Scotland (Stentiford and Neil 2011; Molto-Martin *et al*., 2024). In the Mediterranean,
there are few reports of *H. perezi* (or other *Hematodinium* sp.) in any host, however it has been
reported in *C. sapidus* and other crabs in the south of Italy (Pagliara and Mancinelli 2018;
Mancinelli *et al.,* 2013).
109
The life cycle of *H. perezi* is complex, with a progression of several morphological cell types
within the host, culminating in the release of dinospores that emerge from mouth or gills
(Stentiford and Shields, 2005). Transmission of the parasite is thought to be via dinospores,
which remain infective for hours to days in the water column (Coffey *et al*., 2012). Prevalence
and infectivity of *H. perezi* are responsive to salinity, with higher prevalence in crabs captured in
water above 18 psu, and dinospores becoming inactive at salinity below 10 psu (Messick and
Shields 2000; Coffey *et al*., 2012).

The blue crab is also infected across its native range by a virus known as CsRV1, which was first

observed in electron microscopy as "reo-like virus" (RLV) in the 1970s (Johnson 1977). This

virus is associated with the majority of mortality in captive blue crabs in the United States

(Bowers *et al.,* 2010; Spitznagel *et al.,* 2019). Surprisingly, the CsRV1 genome has a region of

high genetic similarity to a reovirus, named P virus, described in the portunid crab *Liocarcinus*

depurator, on the Mediterranean coast of France (Flowers *et al.,* 2016; Montanie *et al.,* 1993;

Walton *et al.,* 1999). There has been no publication on evidence of CsRV1 or P virus in the

Mediterranean in the intervening 25 years. CsRV1 has been studied thoroughly within the crab's

native range (Zhao *et al.,* 2020). CsRV1 is present from Uruguay to New England, but has not

been reported in wild capture of any other species than *C. sapidus* (Zhao *et al.,* 2020). The

genotype of CsRV1 varies in a manner consistent with geographic origins (Zhao *et al.,* 2023).

As *C. sapidus* continues to expand and increase in abundance across the Mediterranean and

Atlantic coast of Europe, several authors point out that there are still considerable gaps in our

knowledge of the ecology of this crustacean in the new range, even as the species is being

exploited as a fishery resource in some locations (e.g., Mancinelli *et al.,* 2017; Vella *et al.,* 2023;

Azzurro *et al.,* 2024; Galil 2024). Given the importance of *H. perezi* and CsRV1 in the ecology

of *C. sapidus* in its native range, it is relevant to investigate how these pathogens are distributed

in *C. sapidus* in the invaded range. Given the global distribution of *H. perezi* and other

Hematodinium sp., the bi-hemispheric distribution of CsRV1 and molecular evidence that

CsRV1 may have a close relative in Europe, we investigated the prevalence of these two known

C. sapidus pathogens in the Adriatic and Aegean Seas, where *C. sapidus* has been reported for

over a decade, and on the newly-invaded coast of Portugal in the Guadiana estuary.

Materials and methods

Crab sampling and environmental data

Thirty adult *Callinectes sapidus* were collected from each of 6 estuaries in the Mediterranean and

Atlantic Iberian coast between 2021 and 2023 (Figure 1), for a total of 180 crabs.

Wire traps were used in Mediterranean sampling locations of the South Adriatic Sea in Croatia

(Parila lagoon in Neretva river delta, Ston saltworks, Mlini in Župa Bay) and Montenegro (Ada

Bojana). Blue crabs originating from the Aegean sea (North Euboean Gulf) in Greece were also

captured using commercial wire traps and obtained from the fish market. Mediterranean crab

samples were transported to laboratory on ice and processed immediately upon arriving.

Temperature and salinity were measured at the time of the sampling in Parila lagoon and Ston

saltworks by YSI Pro30 probe, and in Ada Bojana by HANNA HI 98194 multiparameter probe.

Environmental data for Mlini were acquired from a year round study of the Župa bay (M.

Pećarević, personal communication) while the temperature of the Aegean sea was obtained by

daily satellite readings provided by NOAA, some 70 km southeast from the sampling site.

Crabs from Atlantic coast were captured by fishermen using gill and fyke nets in the tidal Guadiana River in Portugal (bordering Spain). Salinity and temperature were collected as a contribution to the International Group for Marine Ecology Time Series (IGMETS) monitoring program, in the Guadiana lower estuary station, southwest Iberian Peninsula. These crabs were transported to shore within 2 hours, and frozen at -20 ˚C until processing in the laboratory.

Table 1. *Callinectes sapidus* sampling sites, dates and environmental conditions

Location	Map ID	Collect date (month-year)	Latitude (°N)	Longitude (°E)	Temp (°C)	Sal (psu)
Parila lagoon, South Adriatic Sea	1	February and March 2021	43.0307	17.4512	15	18
Mlini, South Adriatic Sea	2	February and May 2021	42.8301	17.6975	21.7	38
Ston saltworks, South Adriatic Sea	3	May, June and October 2021	42.6178	18.2031	18.8	38
Ada Bojana, South Adriatic Sea	4	September 2022	41.8538	19.3572	22.5	32
North Euboean Gulf, Aegean Sea	5	April 2023	38.7107	23.2851	15.0	n.a.*
Foz de Odeleite, Guadiana estuary	6	June 2023	37.3534	-7.4409	18.3	13

*n.a.   not available

167   **Figure 1.** Sampling locations on the Adriatic, Aegean and Iberian Atlantic coasts. Numbers at
168   each location correspond to numbers in Table 1.
169

Dissection, nucleic acid extraction and polymerase chain reactions

Crab legs (fresh or partially thawed on ice) were externally cleaned with RNaseZap (Thermo
Fisher Scientific) and dissected with new razor blades. Two samples of 50-100 mg muscle tissue
were dissected from a walking leg of each crab and stored at 4°C in 96% ethanol (Gram-Mol,
Croatia) for DNA analysis, as well as in TriReagent solution (Thermo Fisher Scientific) for RNA
analysis. All samples were transported to the Laboratory for Aquaculture of the Institute of
Oceanography and Fisheries in Split, Croatia for nucleic acid extraction and histological analysis
see Methods 2.3.
178

DNA extraction

Genomic DNA was extracted following a simplified phenol-chloroform-isoamyl DNA isolation
procedure (Laird *et al.,* 1991). Resulting DNA pellets were suspended in 30 µl of TE buffer,
checked for purity and concentration by NanoPhotometer (Implen) and stored at -20°C until
shipment. DNA was then re-precipitated with ethanol and shipped as precipitates to the Institute
Marine and Environmental Technology (IMET) in Baltimore, USA for PCR analyses.
185

RNA extraction

For the RNA extraction the tissue was homogenized with 1 mL TriReagent solution (Thermo
Fisher Scientific) with ceramic beads using MagNa Lyser (Roche) and then the procedure
followed the protocol used by Spitznagel *et al.* (2019). Obtained RNA pellets were suspended in
50 µl of nuclease-free water and checked for purity and concentration by NanoPhotometer
(Implen). RNA was re-precipitated in ethanol and pelleted nucleic acid samples were shipped to
the Institute of Marine and Environmental Technology (IMET) in Baltimore, USA where they
were stored at -20°C until resuspension in nuclease-free water for RT-qPCR analysis.
194

RTt-qPCR detection of CsRV1

The presence of CsRV1 RNA was investigated using quantitative reverse transcription PCR (RT-
qPCR) methods adapted from Flowers *et al.,* (2016), using a primer pair designed to amplify a
158-bp region of the ninth genome segment of CsRV1: 5'-
TGCGTTGGATGCGAAGTGACAAAG-3' (RLVset1F) and 5'-
GCGCCATACCGAGCAAGTTCAAAT-3' (RLVset1R). Positive controls and standard curves
consisted of a 10-fold dilution series of purified double-stranded RNA (dsRNA) purified from
crabs infected with CsRV1, and containing 10 to 10^6 CsRV1 genome copies per µl, serially
diluted in 25 ng µl^{-1} yeast tRNA carrier. The RT-qPCR cycling conditions and reagents were as
described by Spitznagel *et al.* (2019), using qScript XLT One-Step RT-PCR Kit (Quantabio,

Beverly MA) in 10 µl reactions containing 0.5 µM of each primer, on a QuantStudio 3 Real Time
Thermocycler. Gene target copies were calculated as copies mg^{-1} of crab muscle, and samples
with >500 copies mg^{-1} were recorded as CsRV1-positive according to Flowers *et al.* (2016),
which is based on empirical measurement of background contamination from field and
laboratory procedures. RNA amplification quality was assessed by using endpoint RT-PCR to
amplify a fragment the blue crab small subunit ribosomal RNA transcript using primers
CrabSSUfor 5'-CATGTCTAAGTACAAGCCGAA-3' and CrabSSURev 5'-
GGGTAATTTGCGTGCCT-3', which produce a 405 nucleotide amplicon (GenBank
JAHFWG010003967; Bachvaroff *et al.,* 2021). PCR products were assessed by gel
electrophoresis. RNA samples that failed to amplify with this rRNA primer set were deemed low
quality and were not included in the CsRV1 analyses.

PCR detection of Hematodinium sp.

The presence and amount of *Hematodinium* sp. DNA was assessed by the qPCR method of
Nagle *et al.,* (2009), which amplifies a 121-nucleotide region of the small subunit ribosomal
RNA gene. Analyses were performed in duplicate on a QuantStudio 3 Real Time Thermocycler
using PerfeCTa SYBR Green FastMix (Quantabio) with the following thermocycling conditions;
50°C for 2 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min. The
qPCR reactions included negative (no template) and positive (cloned target gene) controls. The
number of *Hematodinium* sp. target copies was quantified by comparison to a standard curve (10
to 10^6 copies per µl) of cloned *Hematodinium* sp. DNA (pES103, see Hanif *et al.,* 2013) that was
run on each plate. Based on the fraction of total DNA used in the qPCR reaction and the weight
of the tissue from which it was extracted, the *Hematodinium* sp. infection intensity was
expressed as the number of gene copies mg^{-1} of muscle.

Sequencing and analysis of H. perezi ITS1 gene

A 783 bp region of the 18S rRNA gene, from the SSU across the first internal transcribed spacer
(ITS1) region to the 5.8S region, was amplified and sequenced for phylogenetic analysis.
Amplification of this region was performed using the primers PlaceF (5'-
GGGTAATCTTCTGAAAACGCATCGT-3') from Hanif *et al.*, 2013 and a reverse primer (5'-
CTGGTCTCAGCGTCTGTTCA -3') from the 5.8S region of the *H. perezi* rRNA gene cluster.
The PCR was conducted on a BioRad C1000 thermocycler with AccuStart II PCR ToughMix
under the following conditions: initial denaturation at 94°C for 5 minutes, followed by 31 cycles
of 94°C for 25 seconds, 58°C for 25 seconds, and 72°C for 1 minute. Amplified products were
visualized by 2% (w/v) agarose gel electrophoresis, stained with SYBR Safe DNA Gel Stain, and
examined under ultraviolet light. Aliquots of each reaction were sequenced using an ABI 3130
genetic analyzer. Quality trimming and removal of primer sequences and low quality sequence
was carried out using CLC Genomics Workbench 9.5.2 (Qiagen, Hilden, Germany). Assembled
forward plus reverse assemblies were manually inspected, and consensus sequences were
exported for further analysis.

Comparison of sequences derived from crabs in this study was conducted by alignment tools in
CLC Workbench. A BLAST search of the 650 nt region identified candidate *H. perezi* sequence
accessions in GenBank for comparison to experimental data. Selected candidates were chosen
that represented genotypes I, II, and III (Small *et al.*, 2012) and geographic origins of Europe,
Asia and North America. The reulsting alignment was trimed to the sequence area in common,
visually inspected, and visualized using the tree functions using https://ngphylogeny.fr/. Default
parameters were used to create a Neighbor Joining tree with bootstrap values determined using
100 iterations.
253
Histological analysis

At the time of dissection for DNA and RNA extractions, pieces of hepatopancreas were excised
from the crab samples and fixed for 24 hours in modified Davidson's fixative, prepared with
filtered seawater to maintain isotonicity. After fixation, the samples were transferred to 70%
ethanol and stored at 4°C. Selected samples of fixed tissue were dehydrated in graded ethanol
series, cleared in xylene and paraffin-embedded. Tissue sections were cut at 3-5 μm thickness,
stained with Mayer's hematoxylin and eosin (MHE) and coverslipped in NeoMount. Stained
sections were inspected under a Leica DM3000 LED microscope. Images were captured with a
Leica DMC 4500 camera and assembled and annotated in PhotoShop CS5 (Adobe Systems).
263
264
265 **Results**
266
There was no evidence found for CsRV1 virus RNA in the samples of *C. sapidus* collected in
this study, from the Aegean Sea, Adriatic Sea or Guadiana estuary. This represented a total of
166 individual samples from 6 locations. CsRV1 RT-qPCR assays were conducted on all RNA
samples, but of these, several RNA samples (10 from Portugal and 4 from Greece), did not
produce RT-PCR amplicons with the positive control SSU primers, and so were excluded from
analysis.

DNA of the parasite *H. perezi* was detected in crabs from the Adriatic Sea and Guadiana estuary,
but not from the Aegean Sea. The proportion of infected crabs and the mean intensity (copies of
H. perezi target DNA mg^{-1} muscle tissue) are listed in Table 2. Individual crabs had *H. perezi*
DNA intensity ranging from 30 copies to over 100,000 copies of target DNA mg^{-1} of tissue
(supplemental material available).
278

279 **Table 2.** Prevalence and mean copy number of *H. perezi* and CsRV1 as determined by qPCR.
280 Copy number of cloned target genes per mg tissue.

Location	*Hematodinium perezi*			CsRV1	
	Sample	Prevalence	Mean	Sample	Prevalence

	size	(%)	intensity (copies mg^{-1})	size*	(%)
Parila lagoon, South Adriatic Sea	30	23	1.28 x 10^3	30	0
Ston saltworks, South Adriatic Sea	30	40	1.54 x 10^5	30	0
Mlini, South Adriatic Sea	30	67	3.00 x 10^5	30	0
Ada Bojana, South Adriatic Sea	30	10	3.72 x 10^3	30	0
North Euboean Gulf, Aegean Sea	30	0	n.a.	26	0
Foz de Odeleite, Guadiana estuary	30	87	3.25 x 10^4	20	0

*some samples were excluded because they failed the rRNA amplification quality check.

Histological analysis of the hepatopancreas of a crab from Ada Bojana Montenegro that had 1.8
x10e3 gene copies mg^{-1} of fresh weight is shown in **Figure 2**. The interstitial spaces between
tubules is distended and heavily invaded by parasite cells. Occasionally, cellular aggregates,
possibly fixed phagocytes, are seen. Some of the tubules (T) are depleted of lipid inclusions
(arrows).

289
290
Figure 2. Representative micrographs of a hepatopancreas of the *C. sapidus* from Ada Bojana
Montenegro infected with *H. perezi* at high infection intensity. **(A)** The intertubular space is
distended and the interstitial tissue is almost completely replaced by vegetative stages of the
parasite (asterisks). Occasionally, cellular aggregates, presumably fixed phagocytes, can be seen
(arrowheads). Some of the tubules (T) show a depletion of lipid inclusions (arrows). **(B)** High

magnification (400x) showing details of the developmental stages of the parasite with fixed
phagocytes (arrowhead) and parasites adhering to the tubule epithelium (curved arrows). H&E
staining. Scale bar: (A) 200 μm, (B) 50 μm
299
H. perezi infections (qPCR signal) in three locations were intense enough to enable
amplification of a 888 nt region of the ITS and gene. Each of these sequences is deposited in
GenBank, under the accession numbers PP854984 (Portugal), PP854699 (Montenegro),
PP854700 (Zupa). As can be seen in **Figure 3**, a Maximum Likelihood phylogenetic analysis of
aligned ITS1 regions of these sequences with corresponding regions of 256 nucleotides of the *H.*
perezi ITS1 from other regions of the eastern Atlantic, North America, and Asia shows that the
two *H. perezi* sequences from the Adriatic are identical, and slightly less similar to the sequence
from Portugal (98% identity). All of them are more similar to *H. perezi* sequences from
European waters than from Asia or North America. Bootstrap values are below 75 for all nodes
except those at the branchpoints between continental regions.
310

311
312
Figure 3. Neighbor joining tree from alignment (261 nt) of *H. perezi* ITS1 sequences
from this study (no shading) as compared to *H. perezi* sequences in hosts from the
Atlantic coast of Europe and North Africa, Mediterranean, Asia and North America.
Numbers at branches are bootstrap values for 100 replicates. Genbank accession of *H.*
perezi representing genotypes I, II and III are shaded with different colors as indicated
in the legend.
319
320 **Discussion**

The spread of *C. sapidus* throughout the Mediterranean and eastern Atlantic is highly consequential for the ecology and economy of the region. Measuring, predicting and adapting to the specific impacts of this invasive species is important for practical reasons, and presents a rare opportunity to evaluate ecological and evolutionary theories about invasive species escaping predators and disease. There has been some indication that blue crabs can dramatically decrease the population of aquatic prey (Céspedes *et al.,* 2024; Nardelli *et al.,* 2024). In Italy and Spain, bivalve farmers have abandoned the practice because of damage caused by *C. sapidus* predation (Chiesa *et al*., 2025). Often, invasive species exhibit a surge in abundance, followed by a decline and leveling of abundance in a new region after some years (Simberloff and Gibbons, 2004). In some cases this could be a consequence of the introduced species disrupting its own prey or habitat resources. Additionally, within the "enemy escape" framework/concept, this could be a result of both external predators and internal pathogens adapting to and exploiting the presence of the new species.

The current study gives insight to the presence of two known infectious agents of the Atlantic blue crab *C. sapidus* in selected locations of western Mediterranean and Atlantic Iberian estuaries. The opportunity to analyze crabs from the Aegean Sea, Adriatic, and Guadiana estuary represents an east-to-west range that corresponds to the possible expansion trajectory of the species over time (Mancinelli *et al.,* 2021). The two infectious agents studied represent very different life history strategies for pathogens. The CsRV1 virus is described only in the *C. sapidus* native range, and has not been yet been reported in any other species of host (Zhao *et al.,* 2022). In contrast, the protozoan parasite (parasitic dinoflagellate) *H. perezi* is a globally-distributed crustacean host generalist (Small and Li 2024).

There are relatively few studies related to the health status of *C. sapidus* in the Mediterranean and adjacent waters. At the time of sampling, ours was the first study to investigate the virus CsRV1 in the Mediterranean or eastern Atlantic. Although we did not detect CsRV1 in any of the *C. sapidus* tested, our limited study cannot be taken as evidence that the virus is entirely absent from the Mediterranean or eastern Atlantic. Perhaps the most intriguing suggestion that CsRV1 could be present in the Mediterranean is the report by Walton *et al.,* (1999) on the sequence of a segment of the P virus genome in the portunid crab *Liocarcinus depurator*. The segment of P virus has over 95% identity to segment 4 of CsRV1 (Flowers *et al.,* 2016). This level of identity is in the range of genome difference between CsRV1 ecotypes in north America versus south America (Zhao *et al.,* 2020). Future investigations of CsRV1 in the Mediterranean should include assessments of portunid species such as *Liocarcinus* sp. and *Carcinus* sp.

On the North American Atlantic coast, studies have shown that CsRV1 prevalence is highly variable across time and space, often as low as 0% or as high as 75% prevalence (Flowers *et al.,* 2018). The only consistent trend seen for CsRV1 in the native range is the near absence in the

tropics (Zhao *et al.,* 2020). Flowers *et al.,* (2018) noted that high prevalence of CsRV1was
coincident with higher *C. sapidus* populations in Chesapeake Bay, which is consistent with the
principles of host-dependent prevalence (Cortez and Duffy 2021). All of the locations surveyed
in the current study are temperate, but the overall blue crab population numbers are also
unknown in each of the areas. In the coming 5 – 10 years, it will be informative to re-survey for
the presence of CsRV1 virus in areas with particularly high *C. sapidus* populations or where the
species has been recorded for a long time.
368
The presence of *H. perezi* in Mediterranean *C. sapidus* was first reported by Mancinelli *et al.,*
(2013) in ecosystems of the Apulia coastline. In that study, over 70% of the crabs examined (*C.*
sapidus, Eriphia verrucosa, Pachygrapsus marmoratus and *Carcinus mediterraneus*) in the
Torre Colimena (Ionean Sea) basin and Acquatina lagoon (West Adriatic Sea) were infected by
Hematodinium sp. (Mancinelli *et al.,* 2013). Our study showed that just across the southern
Adriatic, prevalence of *H. perezi* was moderate to high at all locations. These two coasts are
highly connected by an anticlockwise gyre in the southern (Estournel *et al.,* 2021), which would
have the potential to rapidly convey both crab larvae and parasite propagules between them.
377
In the Aegean Sea, Lattos *et al.,* (2024) recently reported 85% prevalence of *H. perezi* in *C.*
sapidus that were collected in November following 2020 mortality events. They suggest this high
prevalence is responsible for recent declines in *C. sapidus* abundance in that region of Greece.
These authors pointed out that there were no external signs of disease observed during the
sampling of *C. sapidus* but that infected crabs had a characteristic "milky" coloration of
hemolymph. In our study of Aegean Sea crabs harvested in April of 2021, we detected no *H.*
perezi. Our collection was from a local fish market, so lacked fine scale specificity of geographic
origin. We are aware that the use of commercially harvested crabs may bias against heavily
infected or moribund crabs, the work of Lattos *et al.* (2024) reported that both "moribund and
healthy individuals" were found with *H. perezi*. The association of crab declines with *H. perezi* is
still relevant to our current study; there are anecdotal reports from fishermen in Mlini, which had
the 2[nd] highest prevalence (76%), that they didn't encounter blue crabs after the summer of 2021
for almost a year (M. Pećarević, personal communication). While there are many possible
reasons for this, one is that infections by *H. perezi* reduced the abundance of blue crabs, similar
to what has been reported in the *C. sapidus* native range (Lee & Frischer 2004).
393
On the Atlantic coast of north Morroco, Lamkhalkhal *et al.,* (2024) also recently reported *H.*
perezi in *C. sapidus* at 25-60% prevalence. Moreover, they show that the sequence of the ITS1
gene of *H. perezi* from Morocco is similar to that in strains from Europe and the Mediterranean.
Our study is the first in our knowledge to document *H. perezi* on the Atlantic coast of Portugal.
We are aware that a 2021 metagenomic investigation of *C. sapidus* stomach contents from the
Rio Formosa estuary near Faro, Portugal also revealed evidence of *H. perezi* DNA. In this case,
the sequence was a match to the mitochondrial COX1 gene (J. Encarnacao and P. Morais,

personal communication), so no direct comparison can be made to the ITS1 sequences.
Nevertheless, this suggests that the parasite has been in Atlantic coastal estuaries of Portugal
since at least 2021.
404
In the *C. sapidus* native range, the prevalence of *H. perezi* is typically associated with high
salinity (Messick and Shields 2000; Gandy *et al.,* 2015). In our study, there is no strict
correspondence between salinity at the sites sampled and the prevalence of *Hematodinium* sp. In
fact, the highest prevalence was found in the Guadiana River, which had locally low salinity but
is less than 20 km away from the ocean. In the Eastern Adriatic Sea, salinity was above 30 psu
except in Parila lagoon (18 psu), which also had the lowest *Hematodinium* sp. prevalence in that
region. The only sample without detection of *H. perezi* was in the Aegean Sea, where crabs were
obtained from a fish market. Therefore precise location and salinity information was not
available. This is in contrast to the recent report by Lattos *et al.,* (2024) of *H. perezi* prevalence
over 80% across the Aegean Sea, which they note was coincident with a decline in the
abundance of *C. sapidus* as reported by fishermen. The contrasting prevalence found in our study
and that of Lattos *et al.,* (2024) is not unusual for marine disease studies, where diverse
environmental factors interact with shifting population size and connectivity to produce highly
variable prevalence.
419
The relationship of *H. perezi* ITS1 sequences in our study with corresponding regions of *H.*
perezi strains across the globe is consistent with prior observations that *H. perezi* within different
continents are similar to one another, and distinct from other regions, regardless of host. The
high identity of ITS1 sequences between Greece and Portugal strains stands out, however, and
may suggest a rapid movement from one location to another. One possible mechanism for such
transport of a parasite strain is ballast water (Nehring, 2011; Cantrell *et al*., 2020). To comply
with the International Convention for the Control and Management of Ships' Ballast Water and
Sediments (IMO, 2004), prior to entering the Mediterranean, cargo ships exchange ballast water
in the Atlantic. Therefore, it is possible that a strain of *H. perezi* or a host infected with it was
transferred in ballast from the Atlantic to the Aegean Sea. Additional comparisons of *H. perezi*
genetics will be needed to lend support or refute the idea that that the parasite has recently
moved.
432
It is important to understand how endemic infectious agents find or adapt to newly arrived
invasive species for at least two reasons. One is the effect that these agents may have on the
health of native species; will they proliferate in a new reservoir host, or perhaps the new host will
constitute a dead end that lowers transmission of the infectious agent. Evidence is that the
genotype I of *H. perezi* is well- adapted to *C. sapidus*. A second reason to understand infectious
agents is the impact on the invasive species over time. In many regions of Europe, there is still a
policy to reduce the number of *C. sapidus* in the invaded range. As income from *C. sapidus*
fisheries grow, however, there may be concern about disease-related mortality if it affects

harvests. In North America, the CsRV1 virus is associated with stress and high populations of *C.*
sapidus (Spitznagel *et al.,* 2019; Flowers *et al.,* 2018). Elsewhere in the Americas, other viruses
have been recently identified in *C. sapidus* (Tavares *et al*., 2022; Zhao *et al*., 2022), illustrating
that there is a potential for various pathogens to be transferred to or within European waters by
ballast water transfer or movement of live crustaceans. Surveillance of *C. sapidus* health is
therefore advisable in all regions of Europe.

Funding and Acknowledgements

OP received support from the Smithsonian Environmental Research Center. ES and OP were
partially funded by the National Oceanic and Atmospheric Administration, Office of Education
Educational Partnership Program award number (NA21SEC4810005). Its contents are solely the
responsibility of the award recipient and do not necessarily represent the official views of the
U.S. Department of Commerce, National Oceanic and Atmospheric Administration.

Statements and Declarations

The authors have no financial or non-financial interests that are directly or indirectly related to
the work submitted for publication.

References

Azzurro, E., Bonanomi, S., Chiappi, M., De Marco, R., Luna, G.M., *et al.,* 2024. Uncovering
unmet demand and key insights for the invasive blue crab (*Callinectes sapidus*) market before
and after the Italian outbreak: Implications for policymakers and industry stakeholders. *Marine*
Policy, 167, 106295.

Bachvaroff, T.R., McDonald, R.C., Plough, L.V., Chung, J.S., 2021. Chromosome-level genome
assembly of the blue crab, *Callinectes sapidus. G3 (Bethesda)* 11 (9), jkab212.

Blakeslee, A.M.H., Keogh, C.L., Byers, J.E., Kuris, A.M., Lafferty, K.D., *et al.,* 2009.
Differential escape from parasites by two competing introduced crabs. *Marine Ecology Progress*
Series, 393, 83-96.

Bouvier, E. L., 1901. Sur quelques Crustacés du Japon, offerts au Muséum par M. le Dr.
Harmand. *Bulletin du Muséum national d'histoire naturelle,* 7, 332-334.

Bowers, H.A., Messick, G.A., Hanif, A., Jagus, R., Carrion, L., *et al.,* 2010. Physicochemical
properties of double-stranded RNA used to discover a reo-like virus from blue crab *Callinectes*
sapidus. Diseases of Aquatic Organisms, 93, 17−29.

Cantrell, D.L., Groner, M.L., Ben-Horin, T., Grant, J. and Revie, C.W., 2020. Modeling
pathogen dispersal in marine fish and shellfish. *Trends in Parasitology*, *36*(3), pp.239-249.
485

486 Céspedes, V., Bernardo-Madrid, R., Picazo, F., Vilà, M., Rubio, C., *et al.,* 2024. Massive decline
487 of invasive apple snail populations after blue crab invasion in the Ebro River, Spain. *Biological*
488 *Invasions*, 26, 2387-2395.
489

490 Chatton, E. & Poisson, R. 1931. Sur l'existence, dans le sang des Crabes, de Pe ridiniens
491 parasites: *Hematodinium perezi* n. g., n. sp. (Syndinidae). *C.R. Se´ances Soc. Biol. Paris,* 105,
492 553–557
493

494 Chiesa, S., Petochi, T., Brusà, R.B., Raicevich, S., Cacciatore, F., Franceschini, G., Antonini, C.,
495 Vallini, C., Bernarello, V., Oselladore, F. and Ciani, M., 2025. Impacts of the blue crab invasion
496 on Manila clam aquaculture in Po Delta coastal lagoons (Northern Adriatic Sea, Italy).
497 *Estuarine, Coastal and Shelf Science*, *312*, 109037.
498

499 Cilenti, L., Pazienza, G., Scirocco, T., Fabbrocini, A., D'Adamo, R., 2015. First record of
500 ovigerous *Callinectes sapidus* (Rathbun, 1896) in the Gargano Lagoons (south-west Adriatic
501 Sea). *BioInvasions Record*, 4(4), 281-287.
502

503 Clavero, M., Franch, N., Bernardo-Madrid, R., López, V., Abelló, P., *et al.,* 2022. Severe, rapid
504 and widespread impacts of an Atlantic blue crab invasion. *Marine Pollution Bulletin*, 176,
505 113479.
506

507 Cortez, M.H., Duffy, M.A., 2021. The context-dependent effects of host competence,
508 competition, and pathogen transmission mode on disease prevalence. *The American Naturalist*,
509 *198*(2), pp.179-194.
510

511 Culurgioni, J., Diciotti, R., Satta, C. T., Camedda, A., de Lucia, G. A., Pulina, S. *et al.,* 2020.
512 Distribution of the alien species *Callinectes sapidus* (Rathbun, 1896) in Sardinian waters
513 (western Mediterranean). *BioInvasions Record*, 9(1), 65-73.
514

515 Estournel, C., Marsaleix, P., Ulses, C., 2021. A new assessment of the circulation of Atlantic and
516 Intermediate Waters in the Eastern Mediterranean. *Progress in Oceanography*, 198, 102673.
517

518 Enzenroß, R., Enzenroß, L., Bingel, F., 1997. Occurrence of blue crab, *Callinectes sapidus*
519 (Rathbun, 1896) (Crustacea, Brachyura) on the Turkish Mediterranean and the adjacent Aegean
520 coast and its size distribution in the bay of Iskenderun. *Turkish Journal of Zoology*, 21, 113-122.
521

Flowers, E.M., Simmonds, K., Messick, G.A., Sullivan, L., Schott, E.J., 2016. PCR-based prevalence of a fatal reovirus of the blue crab, *Callinectes sapidus* (Rathbun) along the northern Atlantic coast of the USA. *Journal of Fish Diseases*, 39, 705-714.

Flowers, E.M., Johnson, A.F., Aguilar, R., Schott, E.J. 2018. Prevalence of a pathogenic crab virus *Callinectes sapidus* reovirus 1 in near flow-through aquaculture in Chesapeake Bay, USA. *Diseases of Aquatic Organisms*, 129(2), 135-144.

Galil, B. 2024. *Callinectes sapidus* (blue crab). CABI Digital Library datasheet. https://doi.org/10.1079/cabicompendium.90126

Gandy, R., Schott, E.J., Crowley, C., Leone, E.H. 2015. Temperature correlates with annual changes in *Hematodinium perezi* prevalence in blue crab *Callinectes sapidus* in Florida, USA. *Diseases of Aquatic Organisms* 113, 235-243.

Giordani Soika A (1951) Il Neptunus pelagicus (L.) nell'altoAdriatico. *Natura* 42, 18–20

Glamuzina, L., Conides, A., Mancinelli, G., Glamuzina, B., 2021. A comparison of traditional and locally novel fishing gear for the exploitation of the invasive Atlantic blue crab in the Eastern Adriatic Sea. *Journal of Marine Science and Engineering*, 9(9), 1019.

Glamuzina, L., Pešić, A., Marković, O., Tomanić, J., Pećarević, M., *et al.,* 2023. Population structure of the invasive Atlantic blue crab, *Callinectes sapidus* on the Eastern Adriatic coast (Croatia, Montenegro). *Naše more*, 70, 153-159.

Glandon, H.L., Paynter, K.T., Rowe, C.L., Miller, T.J., 2019. Resilience of oxygen consumption rates in the juvenile blue crab *Callinectes sapidus* to future predicted increases in environmental temperature and pCO2 in the mesohaline Chesapeake Bay. *Journal of Shellfish Research*, 38(3), 711-723.

González-Ortegón, E., Berger, S., Encarnação, J., Chairi, H., Morais, P., *et al.,* 2022. Free pass through the pillars of Hercules? Genetic and historical insights into the recent expansion of the Atlantic Blue Crab *Callinectes sapidus* to the West and the East of the Strait of Gibraltar. *Frontiers in Marine Science*, 9, 918026.

Holthuis, L.B. 1961. Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey and the Balkans. *Zoologische Verhandelingen*, 47, 1-67.

Johnson, P.T., 1977. A viral disease of the blue crab, *Callinectes sapidus*: histopathology and differential diagnosis. *Journal of Invertebrate Pathology*, 29, 201-209.

563   Kahn, D.M., Helser, T.E., 2005. Abundance, dynamics and mortality rates of the Delaware Bay
564   stock of blue crabs, *Callinectes sapidus*. *Journal of Shellfish Research*, *24*(1), 269-284.
565
566   Kara, M. H., Chaoui, L., 2021. Strong invasion of Mellah lagoon (South-Western Mediterranean)
567   by the American blue crab *Callinectes sapidus* Rathbun, 1896. *Marine Pollution Bulletin*, 164,
568   112089.
569
570   Kevrekidis, K., Kevrekidis, T., Mogias, A., Boubonari, T., Kantaridou, F., *et al.,* 2023. Fisheries
571   Biology and Basic Life-Cycle Characteristics of the Invasive Blue Crab *Callinectes sapidus*
572   Rathbun in the Estuarine Area of the Evros River (Northeast Aegean Sea, Eastern
573   Mediterranean). *Journal of Marine Science and Engineering*, 11(3), 462.
574
575   Laird, P.W., Zijderveld, A., Linders, K., Rudnicki, M.A., Jaenisch, R., Berns, A. 1991.
576   Simplified mammalian DNA isolation procedure. *Nucleic Acids Research*, 19(15), 4293.
577
578   Lamkhalkhal, A., Rahmouni, I., Selfati, M., Hamid, A., Kmentová, N., *et al.,* 2024.
579   *Hematodinium perezi* (Dinophyceae: Syndiniales) in Morocco: The First Record on the African
580   Atlantic Coast and the First Country Record of a Parasite of the Invasive Non-Native Blue Crab
581   *Callinectes sapidus*. *Journal of Marine Science and Engineering*, *12*(7), 1045.
582
583   Lattos, A., Papadopoulos, D. K., Giantsis, I. A., Stamelos, A.,Karagiannis, D., 2024.
584   Histopathology and phylogeny of the dinoflagellate *Hematodinium perezi* and the epibiotic
585   peritrich ciliate *Epistylis* sp. infecting the blue crab *Callinectes sapidus* in the eastern
586   Mediterranean, *Microorganisms*, 12(3), 456.
587
588   Lee, R.F., Frischer, M.E., 2004. Where have the blue crabs gone: Evidence for a drought induced
589   epidemic of a protozoan disease. *American Scientist*, 92, 547-553.
590
591   Mancinelli, G., Carrozzo, L., Marini, G., Pagliara, P., Pinna, M., 2013. The co-occurrence of
592   *Callinectes sapidus* Rathbun, 1896 (Brachyura: Portunidae) and the parasitic dinoflagellate
593   *Hematodinium* sp. (Dinoflagellata: Syndinidae) in two transitional water ecosystems of the
594   Apulia coastline (South-Italy), *Transitional Waters Bulletin*, 7, 32-42.
595
596   Mancinelli, G., Chainho, P., Cilenti, L., Falco, S., Kapiris, K., *et al.,* 2017. The Atlantic blue crab
597   *Callinectes sapidus* in southern European coastal waters: Distribution, impact and prospective
598   invasion management strategies. *Marine Pollution Bulletin*, 119(1), 5-11.
599
600   Mancinelli, G., Bardelli, R., Zenetos, A., 2021. A global occurrence database of the Atlantic
601   blue crab *Callinectes sapidus*, *Scientific Data,* 8(1), 111.
602

Mancinelli, G., Lago, N., Scirocco, T., Lillo, O.A., De Giorgi, R., Doria, L., Mancini, E.,
Mancini, F., Potenza, L. and Cilenti, L., 2024. Abundance, Size Structure, and Growth of the
Invasive Blue Crab *Callinectes sapidus* in the Lesina Lagoon, Southern Adriatic Sea, *Biology*,
13(12), 1051.

Manfrin, C., Comisso, G., Dall'Asta, A., Bettoso, N. and Chung, J.S., 2016. The return of the
Blue Crab, *Callinectes sapidus* Rathbun, 1896, after 70 years from its first appearance in the
Gulf of Trieste, northern Adriatic Sea, Italy (Decapoda: Portunidae), *Check List*, *12*(6).

Messick, G.A., Shields, J.D., 2000. Epizootiology of the parasitic dinoflagellate *Hematodinium*
sp. in the American blue crab *Callinectes sapidus*, *Diseases of Aquatic Organisms*, 43:139-152.

Molto-Martin I, Neil DM, Coates CJ, MacKenzie SA, Bass D, Stentiford GD, Albalat A. 2024.
Infection of Norway lobster (*Nephrops norvegicus*) by the parasite *Hematodinium* sp.: insights
from 30 years of field observations. *Royal Soc Open Sci.,* 17;11(1):231147.

Montanie, H., Bossy, J.P., Bonami, J.R., 1993. Morphological and genomic characterization of
two reoviruses (P and W2) pathogenic for marine crustaceans; do they constitute a novel genus
of the Reoviridae family? *Journal of General Virology*, 74, 1555-1561.

Morais, P., Gaspar, M., Garel, E., Baptista, V., Cruz, J., *et al.,* 2019. The Atlantic blue crab
Callinectes sapidus Rathbun, 1896 expands its non-native distribution into the Ria Formosa
lagoon and the Guadiana estuary (SW-Iberian Peninsula, Europe). *BioInvasions Records,* 8(1),
123-133.

Nagle, L., Place, A.R., Schott, E.J., Jagus, R., Messick, G., Pitula, J.S., 2009. Real-time PCR-
based assay for quantitative detection of *Hematodinium* sp. in the blue crab *Callinectes sapidus,*
Diseases of Aquatic Organisms, 84: 79-87.

Nardelli, L., Fucilli, V., Pinto, H., Elston, J.N., Carignani, A., *et al.,* 2024. Socio-economic
impacts of the recent bio-invasion of *Callinectes sapidus* on small-scale artisanal fishing in
southern Italy and Portugal. *Frontiers in Marine Science*, *11*, 466132.

Nehring, S. (2011) Invasion history and success of the American blue crab *Callinectes sapidus* in
European and adjacent waters. In: Galil BS, Clark PF, Carlton JT (eds), In the Wrong Place -
Alien Marine Crustaceans: Distribution, Biology and Impacts. Springer, The Netherlands, pp
607–624.

Pagenkopp Lohan, K. M., Reece, K. S., Miller, T. L., Wheeler, K. N., Small, H. J., Shields, J. D.,
2012. The role of alternate hosts in the ecology and life history of *Hematodinium* sp., a parasitic
dinoflagellate of the blue crab (*Callinectes sapidus*). *Journal of Parasitology*, *98*(1), 73-84.

Pagliara, P., Mancinelli, G., 2018. Parasites affect hemocyte functionality in the hemolymph of
the invasive Atlantic blue crab *Callinectes sapidus* from a coastal habitat of the Salento
Peninsula (SE Italy), *Mediterranean Marine Science*, 19, 193-200.

Ribeiro, F., Veríssimo, A., 2014. A new record of *Callinectes sapidus* in a western European
estuary (Portuguese coast), *Marine Biodiversity Records*, 7. e36.

Pezy, J. P., Raoux, A., Baffreau, A., Dauvin, J. C., 2019. A well established population of the
Atlantic blue crab *Callinectes sapidus* (Rathbun, 1896) in the English Channel? *Cahiers de*
Biologie Marine, 60, 205-209.

Shields, J.D., Overstreet, R.M., 2003. The Blue Crab: Diseases, Parasites and Other Symbionts
and Other Symbionts, 8, 223-339. In The Blue Crab: *Callinectes sapidus*. Kennedy. College
Park: Maryland Sea Grant College University of Maryland- UM-SG-TS-2007-1.

Schubart, C.D., Deli, T., Mancinelli, G., Cilenti, L., Gil Fernández, A., *et al.,* 2022.
Phylogeography of the Atlantic blue crab *Callinectes sapidus* (Brachyura: Portunidae) in the
Americas versus the Mediterranean Sea: Determining origins and genetic connectivity of a large-
scale invasion, *Biology*, 12(1), 35.

Schulz AN, Lucardi RD, Marsico TD. 2019. Successful invasions and failed biocontrol: the role
of antagonistic species interactions, *Bioscience,* 69: 711–724.

Simberloff, D., Gibbons, L., 2004. Now you see them, now you don't!–population crashes of
established introduced species, *Biological Invasions*, 6, 161-172.

Small HJ, Shields JD, Reece KS, Bateman K, Stentiford GD (2012) Morphological and
molecular characterization of *Hematodinium perezi* (Dinophyceae: Syndiniales), a dinoflagellate
parasite of the harbour crab, *Liocarcinus depurator*, *Journal of Eukaryotic Microbiology,* 59, 54-
66

Spitznagel, M., Small, H.,J, Lively, J.A., Shields, J.D., Schott, E.J., 2019. Investigating risk
factors for mortality and reovirus infection in aquaculture production of soft-shell blue crabs
(*Callinectes sapidus*), *Aquaculture*, 502, 289-295.

Stentiford, G.D., Neil, D.M. 2011. Diseases of Nephrops and Metanephrops: a review. *Journal of*
Invertebrate Pathology, 106(1), 92-109.

Stentiford GD, Shields JD (2005) A review of the parasitic dinoflagellates *Hematodinium*
species and *Hematodinium*-like infections in marine crustaceans, *Diseases of Aquatic Organisms*
66, 47–70.

Taybi, A. F., Mabrouki, Y., 2020. The American blue crab *Callinectes sapidus* Rathbun, 1896
(Crustacea: Decapoda: Portunidae) is rapidly expanding through the Mediterranean coast of
Morocco, *Thalassas: An International Journal of Marine Sciences*, *36*(2), 267-271.

Torchin, M., Lafferty, K.D., Dobson, A.P., McKenzie, V.J., Kuris, A.M. 2003. Introduced
species and their missing parasites. *Nature* 421, 628-630.

Vella, A., Giarrusso, E., Monaco, C., Mifsud, C.M., Darmanin, S.A., *et al.,* 2023. New Records
of *Callinectes sapidus* (Crustacea, Portunidae) from Malta and the San Leonardo River Estuary
in Sicily (Central Mediterranean), *Diversity*, 15(5), 679.

Walton, A., Montanie, H., Arcier, J.M., Smith, V.J., Bonami, J.R., 1999. Construction of a gene
probe for detection of P virus (Reoviridae) in a marine decapod, *Journal of Virological Methods*,
81(1-2),183-192.

Zhao, M., Behringer, D.C., Bojko, J., Kough, A., Plough, L., M., *et al.,* 2020. Climate and season
is associated with prevalence and distribution of trans-hemispheric blue crab reovirus
(*Callinectes sapidus* reovirus 1), *Marine Ecology Progress Series,* 647, 123-133.

Zhao M., Xu L., Bowers H., Schott E. J. 2022. Characterization of Two Novel Toti-Like Viruses
Co-infecting the Atlantic Blue Crab, *Callinectes sapidus*, in Its Northern Range of the United
States, *Frontiers in Microbiology*, 13, 855750.

Zhao, M., Plough, L.V., Behringer, D.C., Bojko, J., Kough, A.S., *et al.,* 2023. Cross-
Hemispheric Genetic Diversity and Spatial Genetic Structure of *Callinectes sapidus* Reovirus 1
(CsRV1), *Viruses,* 15, 563.