Infections in the Atlantic Blue Crab *Callinectes sapidus* (Rathbun, 1896) in coastal Adriatic, Aegean sea, and Atlantic Iberian coast

Authors:
Eric Schott[1]* Marina Brailo Šćepanović[2], Jasna Maršić Lučić[3], Olivia Pares[1], Mingli Zhao[4], Jerko Hrabar[3], Luka Glamuzina[2], Marijana Pećarević[2], Sanja Grđan[2], João Pedro da Silva Encarnação[5], Pedro Morais[6] Ana Pešić[7], Ilija Ćetković[7], Branko Glamuzina[2]

Affiliations
[1]University of Maryland Center for Environmental Science, Institute of Marine and Environmental Technology, 701 E Pratt St. Baltimore MD 21202, USA

[2] University of Dubrovnik, Department of Applied Ecology, Dubrovnik, Croatia

[3] Institute of Oceanography and Fisheries, Laboratory for Aquaculture, Split, Croatia

[4]Royal Veterinary College, University of London, United Kingdom

[5]Centre of Marine Sciences (CCMAR) University of Algarve, Faro, Portugal

[6]Department of Environmental Science, Policy, and Management, University of California, Berkeley, California, USA

[7] University of Montenegro, Institute of Marine Biology, Kotor, Montenegro

* Corresponding author: Eric Schott schott@umces.edu

Abstract

In the Mediterranean Sea, the abundance of the invasive portunid crab, *Callinectes sapidus*, has dramatically increased in recent years. This raises concerns about damage to ecosystems, but also offers opportunities for exploitation of a new fishery. Newly invasive species may escape from pathogens in their endemic range, may introduce new pathogens, or can become host to endemic pathogens. Understanding these factors is important for predicting or managing natural resources in the invaded range. This study investigated the prevalence of two pathogens common in *C. sapidus* in its home range of North America: the reovirus CsRV1 and the protozoan parasite *Hematodinium* sp. In crabs collected from Aegean, Adriatic, and Atlantic waters, the CsRV1 virus was not detected. In contrast, the parasite *Hematodinium* sp. was found in all areas except the Aegean Sea. Sequence analysis of the *Hematodinium* sp. ITS1 gene indicated that the strains observed are most related to *Hematodinium* sp. strains already described in Europe and the Mediterranean, and not to strains from the Americas. The arrival of new species and new potential pathogens is ongoing through transfer of ballast water to the Atlantic and Mediterranean. Although systems are in place to exchange or inactivate ballast water, it is

39 advisable to continue and expand surveillance for pathogens in introduced species, to inform
40 management of movement of these species between regions.
41
42 **Introduction**
43
The Atlantic blue crab *Callinectes sapidus* Rathbun,1896 (Brachyura: Portunidae) is a swimming
crab native to the western Atlantic in coastal and estuarine waters of New England (USA) to
Uruguay, including the Gulf of Mexico and the Caribbean Sea (Mancinelli *et al.,* 2021, 2017). In
recent years it has also emerged as one of the most concerning invasive species in the
Mediterranean Sea. It was first recorded in Europe in 1900 (Bouvier, 1901), and appears to be
established on the Atlantic coast of France and England (Pezy *et al.,* 2019). Other records show
the species has also moved south to the upper Iberian coast (Ribeiro and Veríssimo 2014). In the
Mediterranean Sea it was first recorded in 1949 in the Gulf of Venice, Italy, in the northern
Adriatic Sea (Mancinelli *et al.,* 2017), and possibly as early as 1935 (Enzenroß *et al.,* 1997). It
has become invasive in the eastern-most and western basins of the Mediterranean and the Black
Sea (Vella *et al.,* 2023). Recently, the species was reported in Malta and Sicily (Vella *et al.,*
2023), Sardinia (Culurgioni *et al.,* 2020), Spain (Clavero *et al.,* 2022), the Adriatic and Aegean
seas (Glamuzina *et al.,* 2023, 2021; Cilenti, 2015; Kevrekidis *et al.,* 2023), and the
Mediterranean coast of Africa (Kara and Chaoui, 2021; Taybi and Mabrouki, 2020). Also
recently, *C. sapidus* has appeared in abundance on the southern Atlantic Coast of Portugal, and
has been reported on the Atlantic coast of Morocco, possibly reflecting an expansion through the
straits of Gibraltar (Morais *et al.,* 2019; González-Ortegón *et al.,* 2022; Lamkhalkhal *et al.,*
2024).
62
The life history of *C. sapidus* includes a 28-35 day oceanic planktonic phase before
metamorphosis to benthic postlarvae (megalopa stage) and enter estuaries. Blue crabs are
relatively short-lived, with an estimated longevity of 2-3 years (Kahn and Hestler 2005).
Therefore, in locations with consistently high crab abundance, there must be recurring sources of
larvae, either through local self recruitment or from broodstock sources upstream in oceanic
currents. For example, in the south-eastern Adriatic Sea, a strong anti-clockwise gyre connects
the coast of Italy and Croatia, while in the Aegean Sea, where blue crabs have been abundant for
decades (Holthuis 1961) currents are weak and the many islands ensure promote connectivity
and self recruitment (Estournel *et al.,* 2021). The local-recruitment hypothesis is supported by
Schubart *et al.,* (2022) who found low genetic connectivity between different populations of *C.*
sapidus in the invaded range, consistent with multiple founder introductions.
74
The success of the blue crab in new regions is likely enhanced by its omnivory and ability to
adapt to a wide range of salinity, temperature and dissolved oxygen (Glandon *et al.,* 2019).
Success of non-native species in a new range is often also attributed to an escape from predators
and infectious diseases that suppressed their numbers in the home range (sometimes referred to

as the enemy release hypothesis) (Blakeslee *et al.,* 2009; Torchin *et al.,* 2007; Schultz *et al.,*
2019). While a lot has been reported on occurrence and the invasiveness of *C. sapidus* in the
Mediterranean, there is still little knowledge of the infections that it may acquire in this new
range. The enemy release (pathogen escape) concept assumes that newly arrived species have not
carried their home-range pathogens with them.
84
In its home range, *C. sapidus* is infected by a number of parasites and viruses (Shields and
Overstreet 2007). The most well characterized is the parasitic dinoflagellate, *Hematodinium*
perezi, which is a host generalist that also infect other crustacean species. Different
Hematodinium sp. genotypes and sub-species are characteristic of different hosts and global
region, and identifiable by genomic changes in the ribosomal RNA genes (Small 2012, Messick
and Shields 2000, Pagenkopp-Lohan *et al.,* 2012). In the United States, regional population
declines of *C. sapidus* have coincided with elevated prevalence *Hematodinium* sp. and in
northern Europe (Irish Clyde Sea) the parasite is associated with declines in Norway lobster (Lee
and Frischer 2004, Stentifod and Neil, 2012). In the Mediterranean, there are few reports of this
parasite in any crustacean, however it is reported in *C. sapidus* in the south of Italy (Pagliara and
Mancinelli 2018; Mancinelli *et al.,* 2013). Intriguingly, Lamkhalkhal *et al.,* (2024) recently
described *Hematodinium perezi* in *C. sapidus* from the Atlantic coast of Morocco.
97
The blue crab is also infected across its native range by a virus known as CsRV1, which was first
observed in electron microscopy as "reo-like virus" (RLV) in the 1970s (Johnson 1977). This
virus is associated with the majority of mortality in captive blue crabs in the United States
(Bowers *et al.,* 2010; Spitznagel *et al.,* 2019). Surprisingly, the CsRV1 genome has a region of
high genetic similarity to a reovirus, named P virus, described in the portunid crab *Liocarcinus*
depurator, on the Mediterranean coast of France (Flowers *et al.,* 2016; Montanie *et al.,* 1993;
Walton *et al.,* 1999). There has been no publication on evidence of CsRV1 or P virus in the
Mediterranean in the intervening 25 years. CsRV1 has been studied thoroughly within the crab's
native range (Zhao *et al.,* 2020). CsRV1 is present from Uruguay to New England, but has not
been reported in wild capture of any other species than *C. sapidus* (Zhao *et al.,* 2020). The
genotype of CsRV1 varies in a manner consistent with geographic origins (Zhao *et al.,* 2023).
109
As *C. sapidus* continues to expand and increase in abundance across the Mediterranean and
Atlantic coast of Europe, several authors point out that there are still considerable gaps in our
knowledge of the ecology of this crustacean in the new range, even as the species is being
exploited as a valuable fishery resource in some locations (e.g., Mancinelli *et al.,* 2017; Vella *et*
al., 2023; Azzurro *et al.,* 2024). One interesting question is: What pathogens can be found in *C.*
sapidus in the new range? Given the cosmopolitan nature of *Hematodinium* sp., and the
molecular evidence that CsRV1 may have a close relative in Europe, we investigated the
prevalence of these two known *C. sapidus* pathogens in the Adriatic and Aegean Seas, and on the
newly-invaded coast of Portugal in the Guadiana estuary.

Materials and methods

Crab sampling and environmental data

Thirty adult *Callinectes sapidus* were collected from each of 6 estuaries in the Mediterranean and Portuguese Atlantic coast between 2021 and 2023 (Figure 1), for a total of 180 crabs. Wire crab traps were used in Eastern Adriatic Sea (Parila lagoon in Neretva river delta, Ston saltworks, and Mlini in Župa Bay), and traps in the Aegean sea in Greece and Ada Bojana in Montenegro, while gill nets were used in the Guadiana River estuary in Portugal. Temperature and salinity were measured by YSI Pro30 probe in Parila lagoon and Ston saltworks and by HANNA HI 98194 multiparameter probe in Ada Bojana at the time of the crab sampling, while environmental data for Mlini were obtained from a year round study of the Župa bay (M. Pećarević, personal communication). Temperature for Greece was obtained by daily satellite readings provided by a NOAA, some 70 km southeast from the sampling site. Crabs from Portugal were captured by fishermen in the tidal Guadiana River (bordering Spain), Atlantic Iberian coast using gill nets and fyke nets. Salinity and temperature were collected as a contribution to the International Group for Marine Ecology Time Series (IGMETS) monitoring program, in the Guadiana lower estuary station, southwest Iberian Peninsula. Crabs were transported to shore within 2 hours, and frozen at -20 ˚C until processing in the laboratory.

Table 1. *Callinectes sapidus* sampling sites, dates and environmental conditions

Location	Map ID	Collect date (month-year)	Latitude (°N)	Longitude (°E)	Temp (°C)	Sal (psu)
Parila lagoon, South Adriatic Sea	1	February and March 2021	43.0307	17.4512	15	18
Mlini, South Adriatic Sea	2	February and May 2021	42.8301	17.6975	21.7	38
Ston saltworks, South Adriatic Sea	3	May, June and October 2021	42.6178	18.2031	18.8	38
Ada Bojana, South Adriatic Sea	4	September 2022	41.8538	19.3572	22.5	32
North Euboean Gulf, Aegean Sea	5	April 2023	38.7107	23.2851	15.0	n.a.*
Foz de Odeleite, Guadiana estuary	6	April 2023	37.3534	-7.4409	18.3	13

*n.a.   not available

142
143 **Figure 1.** Sampling locations on the Adriatic, Aegean and Iberian Atlantic coasts. Numbers at
144 each location correspond to numbers in Table 1.
145

Dissection, nucleic acid extraction and polymerase chain reactions

Freshly caught or crabs were cleaned with RNaseZap (Thermo Fisher Scientific) and dissected
with razor blades. Two samples of 50-100 mg muscle tissue was dissected from a walking leg of
each crab and stored at 4°C in 96% ethanol (Gram-Mol, Croatia) for DNA analysis, as well as in
TriReagent solution (Thermo Fisher Scientific) for RNA analysis. All samples were transported
to the Laboratory for Aquaculture of the Institute of Oceanography and Fisheries in Split, Croatia
for nucleic acid extraction and histological analysis see Methods 2.3.
153

DNA extraction

Genomic DNA was extracted following a simplified phenol-chloroform-isoamyl DNA isolation
procedure (Laird *et al.,* 1991). Resulting DNA pellets were dissolved in 30 µl of TE buffer,
checked for purity and concentration by NanoPhotometer (Implen) and stored at -20°C until
shipment. DNA was then re-precipitated with ethanol and shipped as precipitates to the Institute
Marine and Environmental Technology (IMET) in Baltimore, USA for PCR analyses.
160

RNA extraction

For the RNA extraction the tissue was homogenized with 1 mL TriReagent solution (Thermo
Fisher Scientific) with ceramic beads using MagNa Lyser (Roche) and then the procedure
followed the protocol used by Spitznagel *et al.,* (2019). Obtained RNA pellets were suspended in
50 µl of nuclease-free water and checked for purity and concentration by NanoPhotometer
(Implen). RNA was re-precipitated in ethanol and pelleted nucleic acid samples were shipped to

the Institute of Marine and Environmental Technology (IMET) in Baltimore, USA where they
were stored at -20°C until resuspension in nuclease-free water for Rt-qPCR analysis.
169
Rt-qPCR detection of CsRV1

The presence of CsRV1 RNA was investigated using quantitative reverse transcription PCR (Rt-
qPCR) methods adapted from Flowers *et al.,* (2016), using a primer pair designed to amplify a
158-bp region of the ninth genome segment of CsRV1: 5'-TGC GTT GGA TGC GAA GTG
ACA AAG-3' (RLVset1F) and 5'-GCG CCA TAC CGA GCA AGT TCA AAT-3' (RLVset1R).
Positive controls and standard curves consisted of a 10-fold dilution series of purified double-
stranded RNA (dsRNA) purified from crabs infected with CsRV1, and containing 10 to 10^6
CsRV1 genome copies per µl, serially diluted in 25 ng µl^{-1} yeast tRNA carrier. The qPCR
cycling conditions and reagents were as described by Spitznagel *et al.,* (2019), using qScript
XLT One-Step RT-PCR Kit (Quantabio, Beverly MA) in 10 µl reactions containing 0.5 µM of
each primer, on a QuantStudio 3 Real Time Thermocycler. Gene target copies were calculated as
copies per mg of crab muscle, and samples with >500 copies mg^{-1} were recorded as CsRV1-
positive according to Flowers *et al.* (2016), which is based on empirical measurement of
background contamination from field and laboratory procedures. RNA amplification quality was
assessed by using Rt-PCR to amplify a fragment the blue crab small subunit ribosomal RNA
transcript using primers CrabSSUfor 5'-CATGTCTAAGTACAAGCCGAA-3' and
CrabSSURev 5'-GGGTAATTTGCGTGCCT-3', which produce a 405 nucleotide amplicon
(GenBank JAHFWG010003967; Bachvaroff *et al.,* 2021). RNA samples that failed to amplify
with this rRNA primer set were deemed low quality and were not included in the CsRV1
analyses.
190
PCR detection of Hematodinium *sp.*

The presence and amount of *Hematodinium* sp. DNA was assessed by the method of Nagle *et al.,*
(2009), which amplifies a 121-nucleotide region of the small subunit ribosomal RNA gene.
Analyses were performed in duplicate on a QuantStudio 3 Real Time Thermocycler using
PerfeCTa SYBR Green FastMix (Quantabio) with the following thermocycling conditions; 50°C
for 2 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min. PCR
reactions included negative (no template) and positive (cloned target gene) controls. The number
of *Hematodinium* sp. target copies was quantified by comparison to a standard curve (10 to 10^6
copies) of cloned *Hematodinium* sp. DNA (pES103, see Hanif *et al.,* 2013) that was run on each
plate. Based on the fraction of total DNA used in the qPCR reaction and the weight of the tissue
from which it was extracted, the *Hematodinium* sp. infection intensity was expressed as the
number of gene copies per mg of muscle.
203
Sequencing and analysis of Hematodinium *ITS1 locus*

A 750 bp region of the 18S rRNA gene, from the SSU to the first internal transcribed spacer
(ITS1) region, was amplified and sequenced for phylogenetic analysis. Amplification of this
region was performed using the forward primer PlaceF and reverse ITS1 primers (5'-
GACCCAGCCTTCACGATAAA-3'). The PCR was conducted on a BioRad C1000
thermocycler with AccuStart II PCR ToughMix under the following conditions: initial
denaturation at 94°C for 5 minutes, followed by 31 cycles of 94°C for 25 seconds, 58°C for 25
seconds, and 72°C for 1 minute. Amplified products were visualized by 2% (w/v) agarose gel
electrophoresis, stained with SYBR Safe DNA Gel Stain, and examined under ultraviolet light.
Aliquots of each reaction were sequenced using an ABI 3130 genetic analyzer. Quality trimming
and removal of barcode sequences from the raw reads were carried out using CLC Genomics
Workbench 9.5.2 (Qiagen, Hilden, Germany). The assemblies were manually inspected, and
consensus sequences were exported for further analysis. Sequences were aligned using CLC
Workbench, using default parameters, and used to create a neighbor Joining tree with boostrap
values determined using 100 iterations.
219
Histological analysis

At the time of dissection for DNA extractions, pieces of Croatian crab hepatopancreas were
excised from freshly caught crabs and fixed for 24 hours in modified Davidson's fixative,
prepared with filtered seawater to maintain isotonicity. Crabs from Greece and Portugal were not
assessed by histology. After fixation, the samples were transferred to 70% ethanol and stored at
4°C. Selected samples of fixed tissue were dehydrated in graded ethanol series, cleared in xylene
and paraffin-embedded. Tissue sections were cut at 3-5 µm thickness, stained with Mayer's
hematoxylin and eosin (MHE) and coverslipped in NeoMount. Stained sections were inspected
under a Leica DM3000 LED microscope. Images were captured with a Leica DMC 4500 camera
and assembled and annotated in PhotoShop CS5 (Adobe Systems).
230
231 **Results**
232
There was no evidence found for CsRV1 virus RNA in the samples of *C. sapidus* collected in
this study, from the Aegean Sea, Adriatic Sea or Guadiana estuary. This represented a total of
166 individual samples from 6 locations. CsRV1 PCR assays were conducted on all 30 RNA
samples, but of these, several RNA samples, especially from Portugal, did not produce RT-PCR
amplicons with the positive control SSU primers, and so were excluded from analysis.

DNA of the parasite *Hematodinium* sp. was detected in crabs from the Adriatic Sea and
Guadiana estuary, but not from the Aegean Sea. The proportion of infected crabs and the mean
intensity (copies of *Hematodinium* sp. target DNA per mg muscle tissue) are listed in Table 2.
Individual crabs had *Hematodinium* sp. DNA intensity ranging from 30 copies to over 1 million
copies of target DNA per mg of tissue (supplemental material available).

Table 2. Prevalence and mean copy number of *Hematodinium perezi* and CsRV1 as determined by qPCR. Copy number of *Hematodinium* target genes per mg tissue.

Location	*Hematodinium perezi*			CsRV1	
	Sample size	Prevalence (%)	Mean intensity (copies mg^{-1})	Sample size*	Prevalence (%)
Parila lagoon, South Adriatic Sea	30	23	1.28E+03	30	0
Ston saltworks, South Adriatic Sea	30	40	1.54E+05	30	0
Mlini, South Adriatic Sea	30	67	3.03E+05	30	0
Ada Bojana, South Adriatic Sea	30	10	3.72E+03	30	0
North Euboean Gulf, Aegean Sea	30	0	n.a.	26	0
Foz de Odeleite, Guadiana estuary	30	87	3.25E+04	20	0

Histological analysis of the hepatopancreas of a crab from Montenegro that had 1.8 x10e3 gene copies per mg of fresh weight is shown in **Figure 2**. The interstitial spaces between tubules is distended and heavily invaded by parasite cells. Occasionally, cellular aggregates, possibly fixed phagocytes, are seen. Some of the tubules (T) are depleted of lipid inclusions (arrows).

Figure 2. Micrograph (100X magnification), with H&E staining, of *Hematodinium* sp. PCR-
positive *C. sapidus* hepatopancreas. The intertubular space contains numerous parasite cells
(asterisk). Cellular aggregates, possibly fixed phagocytes, are seen (arrowheads). Some tubules
(T) lack lipid inclusions (arrows). **Inset**: 400X magnification showing trophonts and fixed
phagocytes (arrowhead) adjacent to tubular epithelium (curved arrows). Scale bar: 200 μm, inset:
50 μm.
260
Hematodinium sp. infections (PCR signal) in three locations were intense enough to enable
amplification of a 650 nt region of the rRNA gene. Each of these sequences is deposited in
GenBank, under the accession numbers PP854984 (Portugal), PP854699 (Montenegro),
PP854700 (Zupa). A Maximum Likelihood phylogenetic analysis of aligned ITS1 regions of
these sequences with corresponding regions of 256 nucleotides of the *Hematodinium* ITS1 from
other regions of the eastern Atlantic, North America, and Asia shows that the two *Hematodinium*
sp. sequences from the Adriatic are identical, and slightly less similar to the sequence from
Portugal (98% identity). All of them are more similar to *Hematodinium* sp. sequences from
European waters than from Asia or North America. Bootstrap values are below 75 for all nodes
except those at the branchpoints between continental regions.
271
272

273
274   **Figure 3.** Neighbor joining tree from an alignment (256 nt) of *Hematodinium* sp. ITS1 sequences
275   from this study (within boxes) as compared to *Hematodinium* sp. sequences in hosts from the
276   Atlantic coast of Europe and North Africa, Mediterranean, Asia and North America. Numbers at
277   branches are bootstrap values for 100 replicates.
278

279   **Discussion**
280
The spread of *C. sapidus* throughout the Mediterranean and eastern Atlantic is highly
consequential for the ecology and economy of the region. Measuring, predicting and adapting to
the specific impacts of this invasive species is important for practical reasons, and presents a rare
opportunity to evaluate ecological and evolutionary theories about invasive species escaping
predators and disease. Often, invasive species exhibit a surge in abundance, followed by a

decline and leveling of abundance in a new region after some years (Simberloff and Gibbons,
2004). In some cases this could be a consequence of the introduced species disrupting its own
prey or habitat resources. Additionally, within the "enemy escape" framework/concept, this
could be a result of both external predators and internal pathogens adapting to and exploiting the
presence of the new species. There has been some indication that blue crabs can dramatically
decrease the population of aquatic prey (Céspedes *et al.,* 2024; Nardelli *et al.,* 2024).
292
The current study gives insight to the presence of two known infectious agents of the Atlantic
blue crab *C. sapidus* in selected locations of western Mediterranean and Atlantic Iberian
estuaries. The opportunity to analyze crabs from the Aegean Sea, Adriatic, and Guadiana estuary
represents an east-to-west range that corresponds to the possible expansion trajectory of the
species over time (Mancinelli *et al.,* 2021). The two infectious agents studied represent very
different life history strategies for pathogens. The CsRV1 virus is described only in the *C.*
sapidus native range, and has not been yet been reported in any other species of host (Zhao *et al.,*
2022). In contrast, the protozoan parasite (parasitic dinoflagellate) *Hematodinium* sp., is a
globally-distributed crustacean host generalist (Small 2012).
302
There are relatively few studies related to the health status of *C. sapidus* in the Mediterranean
and adjacent waters. To our knowledge, ours is the first study to investigate the virus CsRV1 in
the Mediterranean or eastern Atlantic. Although we did not detect CsRV1 in any of the *C.*
sapidus tested, our limited study cannot be taken as evidence that the virus is entirely absent
from the Mediterranean or eastern Atlantic. Perhaps the most intriguing suggestion of CsRV1
being present in the Mediterranean is the report by Walton *et al.,* (1999) on the sequence of a
segment of the P virus genome in the portunid crab *Liocarcinus depurator*. The segment of P
virus has over 95% identity to segment 4 of CsRV1 (Flowers *et al.,* 2016). This level of identity
is in the range of genome difference between CsRV1 ecotypes in north America versus south
America (Zhao *et al.,* 2020). Future investigations of CsRV1 in the Mediterranean should
include assessments of portunid species such as *Liocarcinus* sp. and *Carcinus* sp.
314
On the North American Atlantic coast, studies have shown that CsRV1 prevalence is highly
variable across time and space, often as low as 0% as well as 50 to 75% prevalence (Flowers *et*
al., 2018). The only consistent trend seen for CsRV1 in the native range is the near absence in
the tropics (Zhao *et al.,* 2020). Flowers *et al.,* (2018) noted that high prevalence of CsRV1was
coincident with higher *C. sapidus* populations in Chesapeake Bay, which is consistent with the
principles of host-dependent prevalence (Cortez and Duffy 2021). All of the locations surveyed
in the current study are temperate, but the overall blue crab population numbers are also
unknown in all of the areas. It will be important to continue to survey for the CsRV1 virus in
areas with high *C. sapidus* populations and where the species has been recorded for a long time.
324

The presence of *Hematodinium* sp. in Mediterranean *C. sapidus* was first reported by Mancinelli *et al.,* (2013) in ecosystems of the Apulia coastline. In that study, over 70% of the crabs examined (*C. sapidus, Eriphia verrucosa, Pachygrapsus marmoratus* and *Carcinus mediterraneus*) in the Torre Colimena (West Aegean sea) basin and Acquatina lagoon (West Adriatic Sea) were infected by *Hematodinium* sp. (Mancinelli *et al.,* 2013). Our study showed that just across the southern Adriatic, prevalence of *Hematodinium* sp. was moderate to high at all locations. These two coasts are highly connected by an anticlockwise gyre in the southern (Estournel *et al.,* 2021), which would have the potential to rapidly convey both crab larvae and *Hematodinium* sp. propagules between them.

In the Aegean Sea, Lattos *et al.,* (2024) recently reported 85% prevalence of *Hematodinium* sp. in *C. sapidus* that were collected during 2020 mortality events. They suggest this high prevalence is responsible for recent declines in *C. sapidus* abundance in that region of Greece. In our study of Aegean Sea crabs harvested in 2021, we detected no *Hematodinium* sp. Our collection was from a local fish market, so lacked specificity of geographic origin. The association of crab declines with *Hematodinium* sp. is still relevant to our current study; there are anecdotal reports from fishermen in Mlini, which had the 2nd highest prevalence (76%), that they didn't encounter blue crabs after the summer of 2021 for almost a year (M. Pećarević, personal communication). While there are many possible reasons for this, one is that infections by *Hematodinium* sp. reduced the abundance of blue crabs, similar to what has been reported in the *C. sapidus* native range (Lee & Frischer 2004).

On the Atlantic coast of north Morroco, Lamkhalkhal *et al.,* (2024) also recently reported *Hematodinium* sp. in *C. sapidus* at 25-60% prevalence. Moreover, they show that the sequence of the ITS1 gene of *Hematodinium* sp. from Morocco is nearly identical to that in strains from Europe and the Mediterranean. Our study is the first in our knowledge to document *Hematodinium* sp. on the Atlantic coast of Portugal. We are aware that a 2021 metagenomic investigation of *C. sapidus* stomach contents from the Rio Formosa estuary near Faro, Portugal also revealed evidence of *Hematodinium* sp. DNA. In this case, the sequence was a match to the mitochondrial COX1 gene (J. Encarnacao and P. Morais, personal communication), so no direct comparison can be made to the ITS1 sequences. Nevertheless, this suggests that the parasite has been in Atlantic coastal estuaries of Portugal since at least 2021.

In the *C. sapidus* native range, the prevalence of *Hematodinium* sp. is typically associated with high salinity (Messick and Shields 2000; Gandy *et al.,* 2015). In our study, there is no strict correspondence between salinity at the sites sampled and the prevalence of *Hematodinium* sp. In fact, the highest prevalence was found in the Guadiana River, which had locally low salinity but is less than 20 km away from the ocean. In the Eastern Adriatic Sea, salinity was above 30 psu except in Parila lagoon (18 psu), which also had the lowest *Hematodinium* sp. prevalence in that region. The only sample without detection of *Hematodinium* sp. was in the Aegean Sea, where

crabs were obtained from a fish market. Therefore precise location and salinity information was
not available. This is in contrast to the recent report by Lattos *et al.,* (2024) of *Hematodinium* sp.
prevalence over 80% across the Aegean Sea, which they note was coincident with a decline in
the abundance of *C. sapidus* as reported by fishermen. The contrasting prevalence found in our
study and that of Lattos *et al.,* is not unusual for marine disease studies, where diverse
environmental factors interact with shifting population size and connectivity to produce highly
variable prevalence.
372
The relationship of *Hematodinium* sp. ITS1 sequences in our study with corresponding regions
of *Hematodinium* sp. strains across the globe is consistent with prior observations that
Hematodinium sp. within different continents are similar to one another, and distinct from other
regions, regardless of host. The high identity of ITS1 sequences between Greece and Portugal
Hematodinium sp. strains stands out, however, and may suggest a rapid movement from one
location to another. One possible mechanism for such transport of a parasite strain is ballast
water. To comply with the International Convention for the Control and Management of Ships'
Ballast Water and Sediments (IMO, 2004), prior to entering the Mediterranean, cargo ships
exchange ballast water in the Atlantic. Therefore, it is possible that a strain of *Hematodinium* sp.
or a host infected with it was transferred in ballast from the Atlantic to the Aegean Sea.
Additional comparisons of *Hematodinium* sp. genetics will be needed to lend support or refute
the idea that that the parasite has recently moved.
385
It is important to understand how endemic infectious agents find or adapt to newly arrived
invasive species for at least two reasons. One is the effect that these agents may have on the
health of native species; will they proliferate in a new reservoir host, or perhaps the new host will
constitute a dead end that lowers transmission of the infectious agent. Evidence is that the host
generalist *Hematodinium* sp. has already adapted to *C. sapidus*. A second reason to understand
infectious agents is the impact on the invasive species over time. In many regions of Europe,
there is still a policy to reduce the number of *C. sapidus* in the invaded range. As income from *C.*
sapidus fisheries grow, however, there may be concern about disease-related mortality if it
affects harvests. In North America, the CsRV1 virus is associated with stress and high
populations of *C. sapidus* (Spitznagel *et al.,* 2019; Flowers *et al.,* 2018). Elsewhere in the
Americas, other viruses have been recently identified in *C. sapidus* (Tavares *et al.*, 2022; Zhao
et al., 2022), illustrating that there is a potential for various pathogens to be transferred to or
within European waters by ballast water transfer or movement of live crustaceans. Surveillance
of *C. sapidus* health is therefore advisable in all regions of Europe.
400
401
402 **Funding and Acknowledgements**
403

404 OP received support from the Smithsonian Environmental Research Center. ES and OP were
405 partially funded by the National Oceanic and Atmospheric Administration, Office of Education

Educational Partnership Program award number (NA21SEC4810005). Its contents are solely the responsibility of the award recipient and do not necessarily represent the official views of the U.S. Department of Commerce, National Oceanic and Atmospheric Administration.

Statements and Declarations

The authors have no financial or non-financial interests that are directly or indirectly related to the work submitted for publication.

References

Azzurro, E., Bonanomi, S., Chiappi, M., De Marco, R., Luna, G.M., *et al.,* 2024. Uncovering unmet demand and key insights for the invasive blue crab (*Callinectes sapidus*) market before and after the Italian outbreak: Implications for policymakers and industry stakeholders. *Marine Policy*, 167, 106295.

Bachvaroff, T.R., McDonald, R.C., Plough, L.V., Chung, J.S., 2021. Chromosome-level genome assembly of the blue crab, *Callinectes sapidus. G3 (Bethesda)* 11 (9), jkab212.

Blakeslee, A.M.H., Keogh, C.L., Byers, J.E., Kuris, A.M., Lafferty, K.D., *et al.,* 2009. Differential escape from parasites by two competing introduced crabs. *Marine Ecology Progress Series*, 393, 83-96.

Bouvier, E. L., 1901. Sur quelques Crustacés du Japon, offerts au Muséum par M. le Dr. Harmand. *Bulletin du Muséum national d'histoire naturelle,* 7, 332-334.

Bowers, H.A., Messick, G.A., Hanif, A., Jagus, R., Carrion, L., *et al.,* 2010. Physicochemical properties of double-stranded RNA used to discover a reo-like virus from blue crab *Callinectes sapidus*. *Diseases of Aquatic Organisms*, 93, 17−29.

Céspedes, V., Bernardo-Madrid, R., Picazo, F., Vilà, M., Rubio, C., *et al.,* 2024. Massive decline of invasive apple snail populations after blue crab invasion in the Ebro River, Spain. *Biological Invasions*, 26, 2387-2395.

Cilenti, L., Pazienza, G., Scirocco, T., Fabbrocini, A., D'Adamo, R., 2015. First record of ovigerous *Callinectes sapidus* (Rathbun, 1896) in the Gargano Lagoons (south-west Adriatic Sea). *BioInvasions Record*, 4(4), 281-287.

Clavero, M., Franch, N., Bernardo-Madrid, R., López, V., Abelló, P., *et al.,* 2022. Severe, rapid and widespread impacts of an Atlantic blue crab invasion. *Marine Pollution Bulletin*, 176, 113479.

447 Cortez, M.H., Duffy, M.A., 2021. The context-dependent effects of host competence, competition,
448 and pathogen transmission mode on disease prevalence. *The American Naturalist*, *198*(2), pp.179-
449 194.
450
451 Culurgioni, J., Diciotti, R., Satta, C. T., Camedda, A., de Lucia, G. A., Pulina, S. *et al.,* 2020.
452 Distribution of the alien species *Callinectes sapidus* (Rathbun, 1896) in Sardinian waters (western
453 Mediterranean). *BioInvasions Record*, 9(1), 65-73.
454
455 Enzenroß, R., Enzenroß, L., Bingel, F., 1997. Occurrence of blue crab, *Callinectes sapidus*
456 (Rathbun, 1896) (Crustacea, Brachyura) on the Turkish Mediterranean and the adjacent Aegean
457 coast and its size distribution in the bay of Iskenderun. *Turkish Journal of Zoology*, 21, 113-122.
458
459 Flowers, E.M., Simmonds, K., Messick, G.A., Sullivan, L., Schott, E.J., 2016. PCR-based
460 prevalence of a fatal reovirus of the blue crab, *Callinectes sapidus* (Rathbun) along the northern
461 Atlantic coast of the USA. *Journal of Fish Diseases*, 39: 705-714.
462
463 Flowers, E.M., Johnson, A.F., Aguilar, R., Schott, E.J. 2018. Prevalence of a pathogenic crab
464 virus *Callinectes sapidus* reovirus 1 in near flow-through aquaculture in Chesapeake Bay, USA.
465 *Diseases of Aquatic Organisms*, 129(2):135-144.
466
467 Gandy, R., Schott, E.J., Crowley, C., Leone, E.H. 2015. Temperature correlates with annual
468 changes in *Hematodinium perezi* prevalence in blue crab *Callinectes sapidus* in Florida, USA.
469 *Diseases of Aquatic Organisms* 113, 235-243.
470
471 Glamuzina, L., Conides, A., Mancinelli, G., Glamuzina, B., 2021. A comparison of traditional
472 and locally novel fishing gear for the exploitation of the invasive Atlantic blue crab in the Eastern
473 Adriatic Sea. *Journal of Marine Science and Engineering*, *9*(9), 1019.
474
475 Glamuzina, L., Pešić, A., Marković, O., Tomanić, J., Pećarević, M., *et al.,* 2023. Population
476 structure of the invasive Atlantic blue crab, *Callinectes sapidus* on the Eastern Adriatic coast
477 (Croatia, Montenegro). *Naše more*. 70. 153-159.
478
479 Glandon, H.L., Paynter, K.T., Rowe, C.L., Miller, T.J., 2019. Resilience of oxygen consumption
480 rates in the juvenile blue crab *Callinectes sapidus* to future predicted increases in environmental
481 temperature and pCO2 in the mesohaline Chesapeake Bay. *Journal of Shellfish Research*, 38(3),
482 711-723.
483
González-Ortegón, E., Berger, S., Encarnação, J., Chairi, H., Morais, P., *et al.,* 2022. Free pass
through the pillars of Hercules? Genetic and historical insights into the recent expansion of the
Atlantic Blue Crab *Callinectes sapidus* to the West and the East of the Strait of Gibraltar. *Frontiers*
in Marine Science, 9, 918026.

Holthuis, L.B. 1961. Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey
and the Balkans. *Zoologische Verhandelingen*, 47, 1-67.

Johnson, P.T., 1977. A viral disease of the blue crab, *Callinectes sapidus*: histopathology and
differential diagnosis. *Journal of Invertebrate Pathology*, 29, 201-209.

Kahn, D.M., Helser, T.E., 2005. Abundance, dynamics and mortality rates of the Delaware Bay
stock of blue crabs, *Callinectes sapidus*. *Journal of Shellfish Research*, *24*(1), 269-284.

Kara, M. H., Chaoui, L., 2021. Strong invasion of Mellah lagoon (South-Western Mediterranean)
by the American blue crab *Callinectes sapidus* Rathbun, 1896. *Marine Pollution Bulletin*, 164,
112089.

Kevrekidis, K., Kevrekidis, T., Mogias, A., Boubonari, T., Kantaridou, F., *et al.,* 2023. Fisheries
Biology and Basic Life-Cycle Characteristics of the Invasive Blue Crab *Callinectes sapidus*
Rathbun in the Estuarine Area of the Evros River (Northeast Aegean Sea, Eastern Mediterranean).
Journal of Marine Science and Engineering, 11(3), 462.

Laird, P.W., Zijderveld, A., Linders, K., Rudnicki, M.A., Jaenisch, R., Berns, A. 1991. Simplified
mammalian DNA isolation procedure. *Nucleic Acids Research*, 19(15), 4293.

Lamkhalkhal, A., Rahmouni, I., Selfati, M., Hamid, A., Kmentová, N., *et al.,* 2024. *Hematodinium*
perezi (Dinophyceae: Syndiniales) in Morocco: The First Record on the African Atlantic Coast
and the First Country Record of a Parasite of the Invasive Non-Native Blue Crab *Callinectes*
sapidus. *Journal of Marine Science and Engineering*, *12*(7), 1045.

Lee, R.F., Frischer, M.E., 2004. Where have the blue crabs gone: Evidence for a drought induced
epidemic of a protozoan disease. *American Scientist*, 92, 547-553.

Mancinelli, G., Carrozzo, L., Marini, G., Pagliara, P., Pinna, M., 2013. The co-occurrence of
Callinectes sapidus Rathbun, 1896 (Brachyura: Portunidae) and the parasitic dinoflagellate
Hematodinium sp. (Dinoflagellata: Syndinidae) in two transitional water ecosystems of the Apulia
coastline (South-Italy). *Transitional Waters Bulletin*, 7, 32-42.

Mancinelli, G., Chainho, P., Cilenti, L., Falco, S., Kapiris, K., *et al.,* 2017. The Atlantic blue crab
Callinectes sapidus in southern European coastal waters: Distribution, impact and prospective
invasion management strategies. *Marine Pollution Bulletin*, 119(1), 5-11.

Mancinelli, G., Bardelli, R., Zenetos, A., 2021. A global occurrence database of the Atlantic blue
crab *Callinectes sapidus*. *Scientific Data,* 8(1), 111. doi.org/10.1038/s41597-021-00888-w
529
530   Messick, G.A., Shields, J.D., 2000. Epizootiology of the parasitic dinoflagellate *Hematodinium*
531   *sp*. in the American blue crab *Callinectes sapidus*. *Diseases of Aquatic Organisms*, 43:139-152.
532
533   Montanie, H., Bossy, J.P., Bonami, J.R., 1993. Morphological and genomic characterization of
534   two reoviruses (P and W2) pathogenic for marine crustaceans; do they constitute a novel genus of
535   the Reoviridae family? *Journal of General Virology*, 74, 1555-1561.
536
537   Morais, P., Gaspar, M., Garel, E., Baptista, V., Cruz, J., *et al.,* 2019. The Atlantic blue crab
538   *Callinectes sapidus* Rathbun, 1896 expands its non-native distribution into the Ria Formosa lagoon
539   and the Guadiana estuary (SW-Iberian Peninsula, Europe). *BioInvasions Records* 8(1), 123-133.
540
541   Nagle, L., Place, A.R., Schott, E.J., Jagus, R., Messick, G., Pitula, J.S., 2009. Real-time PCR-
542   based assay for quantitative detection of *Hematodinium* sp. in the blue crab *Callinectes sapidus.*
543   *Diseases of Aquatic Organisms,* 84: 79-87.
544
545   Nardelli, L., Fucilli, V., Pinto, H., Elston, J.N., Carignani, A., *et al.,* 2024. Socio-economic impacts
546   of the recent bio-invasion of *Callinectes sapidus* on small-scale artisanal fishing in southern Italy
547   and Portugal. *Frontiers in Marine Science*, *11*, 466132.
548
549   Pagliara, P., Mancinelli, G., 2018. Parasites affect hemocyte functionality in the hemolymph of
550   the invasive Atlantic blue crab *Callinectes sapidus* from a coastal habitat of the Salento Peninsula
551   (SE Italy). *Mediterranean Marine Science*. 19, 193-200.
552
553   Ribeiro, F., Veríssimo, A., 2014. A new record of *Callinectes sapidus* in a western European
554   estuary (Portuguese coast). *Marine Biodiversity Records*, 7. e36.
555
556   Pezy, J. P., Raoux, A., Baffreau, A., Dauvin, J. C., 2019. A well established population of the
557   Atlantic blue crab *Callinectes sapidus* (Rathbun, 1896) in the English Channel? *Cahiers de*
558   *Biologie Marine*, 60, 205-209.
559
560   Shields, J.D., Overstreet, R.M., 2003. The Blue Crab: Diseases, Parasites and Other Symbionts
561   and Other Symbionts, 8, 223-339. In The Blue Crab: *Callinectes sapidus*. Kennedy. College Park:
562   Maryland Sea Grant College University of Maryland- UM-SG-TS-2007-1.
563
Schubart, C.D., Deli, T., Mancinelli, G., Cilenti, L., Gil Fernández, A., *et al.,* 2022.
Phylogeography of the Atlantic blue crab *Callinectes sapidus* (Brachyura: Portunidae) in the
Americas versus the Mediterranean Sea: Determining origins and genetic connectivity of a large-
scale invasion. *Biology*, 12(1), 35.

Simberloff, D., Gibbons, L., 2004. Now you see them, now you don't!–population crashes of
established introduced species. *Biological Invasions*, 6, 161-172.

Small, H. J., 2012. Advances in Our Understanding of the Global Diversity and Distribution of
Hematodinium spp. – Significant Pathogens of Commercially Exploited Crustaceans. *Journal of*
Invertebrate Pathology 110 (2), 234-46.

Spitznagel, M., Small, H.,J, Lively, J.A., Shields, J.D., Schott, E.J., 2019. Investigating risk factors
for mortality and reovirus infection in aquaculture production of soft-shell blue crabs (*Callinectes*
sapidus). *Aquaculture*, 502, 289-295.

Stentiford, G.D., Neil, D.M. 2011. Diseases of Nephrops and Metanephrops: a review. *Journal of*
Invertebrate Pathology, 106(1), 92-109.

Taybi, A. F., Mabrouki, Y., 2020. The American blue crab *Callinectes sapidus* Rathbun, 1896
(Crustacea: Decapoda: Portunidae) is rapidly expanding through the Mediterranean coast of
Morocco. *Thalassas: An International Journal of Marine Sciences*, *36*(2), 267-271.

Torchin, M., Lafferty, K.D., Dobson, A.P., McKenzie, V.J., Kuris, A.M. 2003. Introduced species
and their missing parasites. *Nature* 421, 628-630.

Vella, A., Giarrusso, E., Monaco, C., Mifsud, C.M., Darmanin, S.A., *et al.,* 2023. New Records of
Callinectes sapidus (Crustacea, Portunidae) from Malta and the San Leonardo River Estuary in
Sicily (Central Mediterranean). *Diversity*, 15(5), 679.

Walton, A., Montanie, H., Arcier, J.M., Smith, V.J., Bonami, J.R., 1999. Construction of a gene
probe for detection of P virus (Reoviridae) in a marine decapod. *Journal of virological methods*,
81(1-2),183-192.

Zhao, M., Behringer, D.C., Bojko, J., Kough, A., Plough, L., M., *et al.,* 2020. Climate and season
is associated with prevalence and distribution of trans-hemispheric blue crab reovirus (*Callinectes*
sapidus reovirus 1). *Marine Ecology Progress Series,* 647, 123-133.

Zhao M., Xu L., Bowers H., Schott E. J. 2022. Characterization of Two Novel Toti-Like Viruses
Co-infecting the Atlantic Blue Crab, *Callinectes sapidus*, in Its Northern Range of the United
States. *Frontiers in Microbiology*, 13, 855750.

Zhao, M., Plough, L.V., Behringer, D.C., Bojko, J., Kough, A.S., *et al.,* 2023. Cross-Hemispheric
Genetic Diversity and Spatial Genetic Structure of *Callinectes sapidus* Reovirus 1 (CsRV1).
Viruses, 15, 563.