Neurons in auditory cortex integrate information within constrained temporal windows that are invariant to the stimulus context and information rate

Magdalena Sabat[1,2,†], Hortense Gouyette[2], Quentin Gaucher[3], Mateo López Espejo[4], Stephen V. David[4], Sam Norman-Haignere[5,6,*,†], Yves Boubenec[2,*,†]

[1]Laboratoire de Neurosciences Cognitives et Computationnelles, Département d'études cognitives, INSERM, Ecole Normale Supérieure, PSL University, Paris, France

[2]Laboratoire des systèmes perceptifs, Département d'études cognitives, École normale supérieure, PSL University, Paris, France

[3]Sorbonne Université, CNRS, Inserm, Institut de la Vision, F-75012 Paris, France

[4]Oregon Hearing Research Center, Department of Otolaryngology-Head & Neck Surgery, Oregon Health & Science University. Portland, USA.

[5]Department of Biostatistics & Computational Biology, Department of Neuroscience, University or Rochester Medical Center, Rochester, USA

[6]Department of Brain & Cognitive Sciences, Department of Biomedical Engineering, University or Rochester, Rochester, USA

*: these authors equally contributed to the work

†: corresponding authors

magdalena.sabat@ens.psl.eu

Samuel_Norman-haignere@urmc.rochester.edu

yves.boubenec@ens.fr

Data and Code availability

Code and data necessary for replication of the results in this article will be made publicly available upon publication in a peer reviewed journal.

Acknowledgments

We would like to thank Balkis Cadi and Lynda Bourguignon for technical and administrative support throughout the project. This work was supported by the National Science Foundation & Institut Universitaire de France (CRCNS grant to Y.B. and S.V.N.-H; NSF-2308725, ANR-17-EURE-0017, ANR-10-IDEX-0001-02 and ANR-23-NEUC-0001), as well as the National Institutes of Health (R00-DC018051 to S.V.N.-H).

Author contributions

Conceptualization, M.S., S.N-H, and Y.B.; Data collection, M.S, H.G, Q.G., and M.L.E; Formal Analysis, data curation and visualization, M.S, S.N-H and Y.B; Software, M.S, Q.G., S.N-H and Y.B; Writing, first draft M.S, S.N-H and Y.B, all authors reviewed the article before submission; Supervision and funding, S.N-H, S.V.D, and Y.B.

Competing interests

The authors declare no competing interests.

Abstract

Much remains unknown about the computations that allow animals to flexibly integrate across multiple timescales in natural sounds. One key question is whether multiscale integration is accomplished by diverse populations of neurons, each of which integrates information within a constrained temporal window, or whether individual units effectively integrate across many different temporal scales depending on the information rate. Here, we show that responses from neurons throughout the ferret auditory cortex are nearly completely unaffected by sounds falling beyond a time-limited "integration window". This window varies substantially across cells within the auditory cortex (~15 to ~150 ms), increasing substantially from primary to non-primary auditory cortex across all cortical layers, but is unaffected by the information rate of sound. These results indicate that multiscale computation is predominantly accomplished by diverse and hierarchically organized neural populations, each of which integrates information within a highly constrained temporal window.

Introduction

Temporal integration is central to nearly all aspects of auditory perception. A key challenge is that information in natural sounds is distributed across a vast range of timescales. Speech and animal vocalizations, for example, are characterized by time-varying changes in pitch and spectrotemporal structure that unfold over tens of milliseconds, which are then assembled into longer sequences across hundreds of milliseconds or seconds (Kello et al., 2017). Importantly, the rate at which this information varies is itself highly dynamic (Pellegrino et al., 2011). The rate of spectrotemporal changes, for example, is constantly varying, even within a vocalization, and the duration and density of vocalizations that compose a sequence is also highly variable. Thus, to derive information from complex sounds such as speech and animal vocalizations, the auditory system must be able to flexibly integrate across multiple timescales.

One hypothesis is that neurons in the auditory cortex can flexibly vary the time window over which they integrate information, depending on the rate at which information is varying. This hypothesis is motivated, in part, by prior work showing neurons in the auditory cortex perform flexible, context-dependent computations (J. Fritz et al., 2005; J. B. Fritz et al., 2007; Spitmaan et al., 2020). Artificial neural networks increasingly rely on flexible and nonlinear computational units whose temporal integration window can dynamically vary (Keshishian et al., 2020; Keshishian & Norman-Haignere, 2021), and such systems have shown promise as models of neural computation in the brain (Cichy & Kaiser, 2019; Esposito et al., 2024; Kell et al., 2018; Kietzmann et al., 2019; Li et al., 2023; Yamins & DiCarlo, 2016). A second hypothesis is that individual neurons integrate across largely fixed timescales, but instantiate a diverse array of these integration scales across cells so as to support flexible multiscale integration across the neural population (Perez-Nieves et al., 2021). This hypothesis is motivated in part by prior work suggesting that cortical neurons show a diverse range of temporal scales even within a single cortical region (Cavanagh et al., 2020; Elgueda et al., 2019; T. Lu et al., 2001; Ulanovsky, 2004).

Distinguishing between these hypotheses has been challenging, in part due to the difficulty of measuring integration windows from highly nonlinear systems such as the brain. Integration windows are often defined as the time window analyzed by a sensory system, and thus the window within which stimuli can alter the neural response (Chen et al., 2012; Lerner et al., 2011; Poeppel, 2003; Theunissen & Miller, 1995). Although this definition is simple and general, there have been no simple and general methods for estimating integration windows. Neurons in the auditory system are often modeled using

spectrotemporal receptive fields (STRFs), which assume a fixed, linear mapping between a spectrogram
and the neural response. STRFs, however, cannot model dynamic nonlinear processing, which is
prominent in the cortex (Harper et al., 2016; Meyer et al., 2016; Mizrahi et al., 2014; S. V. Norman-
Haignere & McDermott, 2018), and likely plays an important role in flexible, multiscale integration
(Deneux et al., 2016; Dubreuil et al., 2022; Kell et al., 2018; Keshishian et al., 2020).

In this study, we leveraged a recently developed paradigm for measuring integration windows from
nonlinear (or linear) systems (the temporal context invariance or TCI paradigm) (**Figure 1A**). The
paradigm directly tests the definition of an integration window by determining whether there is a stimulus
duration for which the response is invariant to surrounding context stimuli. As a consequence, the method
does not make any assumptions about the features that underlie the response (e.g., a spectrogram) or
the relationship between those features and the neural response (e.g., linear or nonlinear). The TCI
paradigm was recently validated using artificial deep neural networks (Keshishian & Norman-Haignere,
2021) and macroelectrode recordings from the human auditory cortex (S. V. Norman-Haignere et al.,
2022). Macroelectrodes, however, average activity across thousands of neurons within a local area
(Baratham et al., 2022; Leonard et al., 2024; Manning et al., 2009; Ray & Maunsell, 2011), which makes
it impossible to determine if multiscale integration is accomplished by diverse populations of neurons
integrating across largely fixed timescales, or flexible integration within individual neurons.

In this study, we applied the TCI paradigm to microelectrode recordings from the ferret auditory cortex
using ferret vocalizations, speech, and music across three experiments. We first tested whether cells in
the auditory cortex showed evidence of a well-defined integration window, beyond which stimuli have
little effect on the response, and if so, how this window varies across the neural population. We then
examined whether integration windows reflect absolute time or vary with the information rate of sound by
systematically manipulating the information rate using stretched and compressed sounds.

Results

Neurons in the auditory cortex integrate information within constrained temporal windows

We characterized integration timescales of neurons in the auditory cortex of awake ferrets. We presented
sequences of sound segments in different orders, such that each segment was surrounded by distinct
"context" segments (**Figure 1A**). We note that while "context" has many meanings (Angeloni & Geffen,
2018), we operationally define context in our paradigm as the stimuli that surround a segment. We test
the hypothesis that there is a minimum time window beyond which surrounding stimuli no longer alter the
response. If a neuron has a well-defined integration window that is less than the segment duration, there
will be a moment when the window is fully contained within the shared segment and thus by definition
unaffected by the surrounding segments.

We measured context invariance using the "cross-context correlation" (CCC). For a given segment
duration, we aligned the response timecourses surrounding each segment in a segment-by-time matrix,
separately for each of two contexts (**Figure 1B**). We then correlated corresponding columns across
matrices from differing contexts (CCC, connected boxes in **Figure 1B**). For a given time lag relative to
segment onset, the CCC measures the similarity of the response pattern across segments between the
two contexts. At segment onset, the integration window will overlap the preceding segments, which are
random across contexts, and the CCC should be near 0. If the integration window is less than the
segment duration, there will be a time lag when the window is fully contained within the shared segment,

leading to a correlation of 1 for a noise-free response. To account for noise, we measured a ceiling for the CCC by computing the same correlation when the context was identical using repeated presentations of the same sequence. Longer integration windows will cause the peak of the CCC to drop relative to the noise ceiling and will also cause a more gradual build-up and fall-off since it takes longer for the integration window to enter and exit the shared segment.

Figure 1 | Neurons in auditory cortex show context-invariant responses. A, Segments of natural sounds were presented in different orders, leading to the same segment (blue) being surrounded by different "context segments". **Top row,** If the neural response has an integration window that is less than the segment duration, there will be a moment when the window is fully contained within the shared segment yielding the same response across both contexts. Waveforms for different segments are schematically indicated by colored boxes and the integration window is plotted at the moment of maximum overlap between the two contexts. **Bottom row,** If the shared segments are shorter than the integration window, then the surrounding context segments can alter the response. **B,** Schematic of the "cross-context correlation" (CCC) analysis used to estimate context invariance. Response timecourses surrounding each segment are organized as a segment-by-time matrix (aligned to segment onset), separately for each context. We then correlate responses across segments between the two contexts for a given lag, which is equivalent to correlating corresponding columns (connected boxes) across matrices from different contexts (see text for details). A noise ceiling for the CCC is computed by performing the same analysis when the context is identical (using repeated presentations of each sequence). **C,** Example of the CCC (blue) and noise ceiling (black) from example multi-units in primary (MEG, top) and non-primary (PEG, bottom) ferret auditory cortex. For all units, there is a lag and segment duration for which the CCC equals the noise ceiling, indicating a context-invariant response. The segment duration needed to achieve a context-invariant response varies substantially across units.

We plot the CCC and noise ceiling for multi-units from different regions of the ferret auditory cortex (**Figure 1C;** see **Supplementary Figure 1** for additional units). Strikingly, we found that for virtually all cells with a reliable response to sound, there was a segment duration and lag for which the CCC reached the noise ceiling, indicating a nearly fully context-invariant response. Once the CCC reached the noise

ceiling, it tightly tracked the noise ceiling until segment offset, consistent with a temporally compact
window composed of a single mode (i.e., not multimodal). These results indicate that nearly all sound-
responsive cells in the auditory cortex show a temporally compact integration window beyond which
stimuli have little effect on the neural response. Notably, the segment duration and lag needed to achieve
a context-invariant response differed substantially across units and cortical fields (**Figure 1C**). For some
units, segment durations as short as 15 ms were sufficient to yield a context-invariant response, while
other units required longer segment durations of 32, 64, or 128 ms, particularly in non-primary regions of
the auditory cortex (PEG).

Neural integration windows vary widely across the neural population, increasing in non-primary
regions across all cortical layers.

To quantify these effects, we used a previously developed and validated computational model to
estimate the integration window of each identified unit (345 units with reliable response to sounds across
3 animals, test-retest correlation r> 0.1; $p<10^{-5}$ via permutation). The model finds a parametric window
that best predicts the CCC across all segment durations and lags. The main text figures plot the estimated
width of the model window (the smallest interval that contained 75% of the mass). The model also
computes an estimated integration "center" (median of the parametric window), which can be thought of
as the overall delay between the stimulus and the integration window. The center of the window was
highly correlated with the window's width and close to the minimal possible center given the window
width, suggesting that neurons integrate information about as quickly as possible given the time window
being analyzed (**Supplementary Figure 2**).

This analysis revealed a wide range of integration windows across cells spanning approximately 3
octaves (**Figure 2A-D; Supplementary Figure 3** shows analogous results for integration centers). These
integration times showed clear anatomical organization (**Figure 2A**): cells that were anatomically closer
showed more similar windows (**Figure 2B**), and cells further from primary auditory cortex (PAC) showed
longer windows (**Figure 2C**) ($F_{1,343}$ = 84.79, *p* < 0.001, $\beta_{distance\ to\ PAC}$ = 0.35 octaves/mm, CI = [0.275,
0.424]; N=345), leading to a substantial and significant difference between primary (MEG) and
nonprimary (PEG) auditory cortex (**Figure 2D**) ($F_{1,343}$ = 109.06, *p* < 0.001, β_{region} = −0.799 octaves, CI =
[-0.949, -0.648]; N=345).

To complement our model-based analyses, we also measured the average CCC in primary and non-
primary auditory cortex and normalized this metric by the average noise ceiling, also computed separately
for each region (**Figure 2E**). This analysis replicated our model-based results, with non-primary regions
requiring substantially longer segment durations for the CCC to reach the noise ceiling. This analysis
further demonstrated that even in non-primary regions, there is a time-limited window (~150 ms), outside
of which stimuli have very little effect on the neural response, suggesting that there is a compact
integration window that strongly constrains the computations of virtually all neurons in the auditory cortex.
These results held across all sound categories tested (**Supplementary Figure 4**).

A Integration window maps
MEG
PEG
1 mm
Integration width (ms)
7 15 32 64 128
△ Ferret A
□ Ferret B
○ Ferret C

B Spatial clustering
Width difference (octaves)
noise floor
Distance between units (mm)
>1 2 2<

C Hierarchical organization
Width (ms)
128
64
32
16
0 2 4 6
Distance from Primary Auditory Cortex (mm)

D Estimated width by region
Neurons/bin
30
20
10
0
8 32 128
Integration width (ms)

E Normalized cross-context correlation by region
Normalized correlation
Segment duration
15 ms 31 ms 62 ms 88 ms 125 ms 250 ms
1
0
0 200
Time lag from onset of shared segment (ms)

■ Primary Auditory Cortex - MEG
■ Non-primary Auditory Cortex - PEG

Cross-context correlation
▨ Time span of shared segment — Primary Auditory Cortex - MEG
— Noise ceiling — Non-primary Auditory Cortex - PEG

Figure 2 | Diverse, hierarchically organized temporal integration windows across the neural population. A, Anatomical map of model-estimated integration windows in three animals (window width: smallest interval containing 75% of the window's mass). **B**, The difference in integration windows between pairs of units as a function of their spatial distance, demonstrating that nearby units have more similar windows. **C**, Integration windows as a function of distance to the primary auditory cortex (PAC) (see color map in inset). **D,** Histograms of integration windows for primary (MEG) and non-primary auditory cortex (PEG) showing substantial diversity across units and hierarchical organization. **E,** Median noise corrected cross-context correlation for all neurons in primary (MEG) and non-primary (PEG) auditory cortex.

Our first experiment shows that temporal integration is organized hierarchically across regions, but did not enable us to investigate hierarchical organization between different cortical layers. To address this question, we repeated our experiments using laminar probes (231 cells with reliable response to sounds across 5 animals, test-retest correlation r> 0.1; $p<10^{-5}$ via permutation) and grouped cells based on both their region and cortical layer (supragranular, granular, infragranular) using source density analysis (**Figure 3A**). We replicated our prior findings, showing a substantial increase in integration windows between primary (MEG) and non-primary regions (PEG) ($F_{1,21.47}$ = 19.76, p < 0.001, β_{region} = −0.88 octaves, CI = [-1.270, -0.490]; N=231, **Figure 3B**). There was also a significant effect of layer (width: $F_{2,218.86}$ = 6.87 , p = 0.001) driven by shorter integration windows in the infragranular layers ($\beta_{infragranular}$ = −0.486 octaves, CI = [-0.796, -0.175]). However, the magnitude of the effect was small both in absolute terms (ΔM_{width} = −4.73 ms) and compared with the effect of region ($F_{2,37.48}$ = 15.44, p < 0.001). Consistent with these results, the average cross-context correlation was similar between layers but differed across regions with non-primary areas requiring a longer segment duration and lags to reach the noise ceiling (**Figure 3C**). Thus, the dominant form of hierarchical anatomical organization appears to be region-based and not layer-driven.

Figure 3 | Integration windows show hierarchical organization across regions but not layers. A, *Top:*
Schematic illustration of recordings from a laminar probe sampling supragranular (blue), granular (red), or
infragranular (yellow). *Bottom:* Current Source Density (CSD) plot, evoked with a burst of noise for an example
session. Individual markers correspond to recorded units and indicate approximate locations along the laminar
probe. The color corresponds to the laminar classification of each unit. The colorbar reflects the spatial distribution
of current flow in extracellular space used to estimate the sources and sinks of change in measured potential along
the laminar probe. Details of the laminar assignment are described in the method section. **B,** Cumulative distribution
of estimated integration windows for primary (top) and non-primary (bottom) regions broken down by layer class
(blue: supragranular, red: granular, yellow: infragranular). Dashed lines show corresponding points between plots
from different regions so they can be visually compared. Integration windows are longer in non-primary regions but
do not differ substantially between layers. **C,** Median noise corrected cross-context correlation separately for each
layer (supragranular, granular, infragranular) and region (primary - MEG, nonprimary - PEG).

Integration windows reflect absolute time and do not vary with the information rate of sound.

Our analyses so far suggest that neurons in the auditory cortex integrate information within a highly
constrained temporal window beyond which stimuli have little effect on the cortical response and that this
window varies substantially between neurons across the population. We next tested whether these
integration windows reflect absolute time or instead vary with the information rate of the sound. To answer
this question, we systematically altered the duration and rate at which sound information varies by
stretching and compressing sounds by a factor of $\sqrt{2}$ (preserving frequency/pitch) (**Figure 4A**). If the
neural integration window varies with information rate (rate-yoked integration), the window should appear
to scale with the magnitude of stretching and compression. In contrast, if the integration window reflects
absolute time (time-yoked integration), the window will be unchanged by this manipulation. We tested
both speech and ferret vocalization because they contain many different forms of time-varying structure
that are ecologically relevant to the animal (particularly for ferret vocalizations). We focused our
experiments on a non-primary region (PEG) because we expected rate-yoked integration, if present, to
be strongest in higher-order regions (Atiani et al., 2014; Elgueda et al., 2019).

As a simple summary metric, we first measured the average cross-context correlation for all units from
non-primary auditory cortex for each of the different rates, pooling data from both sound categories
(**Figure 4B**). We found that the average CCC was virtually identical across the three rates for all segment
durations tested, suggesting integration windows reflect absolute time and do not vary with the
information rate. We next quantitatively compared the integration windows for different stimulus rates,

separately estimated for each cell using our computational model (**Figure 4C**). If the integration windows
are fully rate-yoked, integration windows for stretched sounds should be twice as long on average as the
integration windows for compressed sounds, yielding an octave shift on a logarithmic scale (**Figure 4C**,
purple line). If the integration windows are time-yoked, the windows should cluster around the line of unity
(**Figure 4C,** green line). We found that the estimated windows clearly clustered around the line of unity,
consistent with our previous analysis suggesting time-yoked integration dominates. These effects were
reliable across both of the two sound categories tested (**Supplementary Figure 5**) and across all pairs
of sound rates (e.g., original vs. compressed, original vs. stretched).

Figure 4 | Temporal integration is invariant to information rate. A, Schematic illustration of the predictions for
time- vs. rate-yoked integration. Stretching and compression will expand and shrink the integration window if it is
yoked to the information rate of sound (purple, left) but not if it reflects absolute time (green, right). **B**, Median
normalized CCC across all recorded units for stretched, original, and compressed stimuli. **C**, Integration windows
for all units for compressed (x-axis) and stretched (y-axis) stimuli. Green and purple lines show the prediction from
a time-yoked and rate-yoked response. **D,** Reliability of rate-yoking index across two independent data splits. **E,**
Reliability of integration windows averaged across stimuli rates for comparison.

There was substantial variation from cell to cell in the estimated integration window for stretched and
compressed speech (**Figure 4A**). To test if this variation reflected genuine variation in rate yoking or just
variation due to unreliable sources of noise, we measured the integration window for each cell and rate
using two independent splits of data. We then computed a measure of rate yoking separately for each
data split, and correlated this measure across units between the two splits (**Figure 4D**). Rate yoking was
measured by subtracting the integration window for stretched and compressed speech on a logarithmic
scale and dividing by the 2-octave difference in rates. The metric thus provides a simple, graded measure
of the extent of time-yoked (0) vs. rate-yoked integration (1) (note that due to noise the metric can exceed
the 0 and 1 bounds). The effect of rate of the stimuli was marginally significant (width: $F_{2,664.58} = 2.53$, $p = 0.08$, $\beta_{stretched} = 0.16$ octaves, CI = [0.019, 0.308], $\beta_{compressed} = 0.105$ octaves, CI = [-0.039, 0.249]; N =
702 (observations for 117 neurons)), however, we found that the variation from cell to cell in the strength

of rate yoking was not reliable across data splits (Spearman rank correlation of 0.09, p = 0.52 via
bootstrapping), indicating that it mostly reflected unreliable source of noise in the data (**Figure 4D**). In
contrast, the overall integration window, averaged across rates, was highly reliable (Spearman rank
correlation of 0.72, p < 0.0001 via bootstrapping) (**Figure 4E**). These results demonstrate that neural
integration windows do not vary with the information rate of sound, and thus that multiscale integration is
accomplished by diverse populations of neurons each of which integrate information within a highly
constrained temporal window.

____

Discussion

Our study demonstrates that neurons in the auditory cortex integrate information across a time-limited
window beyond which sounds have very little influence on the cortical response. This window varies by
approximately 3 octaves (~15 to 150 ms) across cells in the auditory cortex and exhibits clear anatomical
organization, with two to three times longer integration windows in non-primary regions and little
difference within a cortical column. In contrast, these windows do not vary with the information rate and
duration of sound structures (e.g., vocalizations), suggesting that they are a fundamental property of the
auditory system and not a property of the sounds analyzed by the auditory system. These results indicate
that multiscale computation is accomplished by diverse, hierarchically organized populations of neurons,
each of which integrates information across a specific, time-limited, and context-invariant temporal
window.

Implications for models of multiscale computation.

Understanding the computations that enable the auditory system to integrate across the multiscale
temporal structure of natural sounds is an important goal of auditory neuroscience. Multiscale integration
is computationally challenging in part because information in sound is organized across many different
timescales, and these timescales are highly variable and context-dependent.

Our results show that virtually all neurons in the auditory cortex integrate information with a compact,
temporal window beyond which stimuli have very little effect on the neural response. This finding is highly
nontrivial, since highly nonlinear, recurrent systems such as the brain are capable of integrating across
potentially very long timescales (Keshishian & Norman-Haignere, 2021; Mante et al., 2013). Our results
suggest that this window is temporally compact, since multi-peaked windows would produce oscillations
in the cross-context correlation, which were not observed empirically. This window is also about as fast
as it can be with window centers close to the minimum possible for a causal system. These results
suggest that temporal integration windows strongly constrain and shape the computations performed by
the auditory cortex.

Our second key result is that integration windows show substantial heterogeneity across the neural
population, with nearly 3 octaves of variation. We show that this variation is hierarchically organized
across regions, but not across cortical layers. We note that increasing integration windows are not an
inevitable consequence of hierarchical organization (as evidenced by the weak effect of layer). For
example, hierarchical recurrent neural networks can show similar integration windows across their layers
(Keshishian & Norman-Haignere, 2021), and training RNNs on challenging auditory tasks does not
necessarily induce increasing integration windows across layers (S. Norman-Haignere et al., 2024). We
note that while we observe a substantial increase in integration windows between regions there is still
substantial heterogeneity in integration windows within a region that exhibits anatomical clustering,

suggesting that within a region the auditory cortex performs multiscale temporal analysis using spatially
organized neural populations.

Critically, we provide evidence that the temporal windows of neurons in the auditory cortex do not vary
with the rate at which information is varying in natural sounds. In particular, we find that temporal windows
on average are completely unaffected by stretching and compression, even though stretching and
compression changes the rate at which virtually all acoustic structure varies, as well as the duration of all
sound events (e.g., vocalizations). This result is also highly nontrivial. Nonlinear systems are highly
capable of flexibly varying their integration window and our TCI method is sensitive to these types of
changes and can robustly detect rate-yoked integration when present (Keshishian & Norman-Haignere,
2021). Collectively, these results demonstrate that multiscale computation is performed by diverse
populations of neurons, each of which integrate information within a highly constrained temporal window
beyond which stimuli have very little effect on the neural response and which is invariant to the
information. These results suggest that flexible temporal integration is accomplished by flexibility at the
neural population level rather than flexibility within single neurons.

Our finding of hierarchical organization is consistent with a prior study that also observed hierarchical
organization in human auditory cortex (S. V. Norman-Haignere et al., 2022). This prior study used
macroelectrodes which pool activity from thousands of neurons and thus was not able to test whether
individual neurons show a time-limited integration window, characterize the heterogeneity across the
neural population, or examine layer-wise effects. Prior work has found that the functional organization of
the human auditory cortex differs substantially from that of ferrets (Landemard et al., 2020), likely in part
due to the unique importance of speech and music to human hearing, and thus a priori, it was entirely
possible that we might have observed a very different organization in ferrets. Our findings therefore
provide novel evidence that region-based hierarchical integration, in which longer timescale computations
are performed using the output of shorter timescale computations, is a fundamental property of biological
auditory systems that is not limited to the human auditory cortex.

The integration windows observed in this study are shorter than those observed in the human auditory
cortex, which range from about 50 to 400 ms. This difference could reflect differences in the neural
measures employed (e.g., macro- vs. micro-electrodes) or could reflect species differences. For example,
prior work shows that neural populations with longer integration windows (< 200 ms) in the human
auditory cortex show strong selectivity for speech and music ((S. V. Norman-Haignere et al., 2022)),
which is absent from ferret auditory cortex (Landemard et al., 2020), suggesting that these longer
timescale may have evolved or developed to process speech- or music-specific structure.

Relationships to other types of neural timescales.

One approach to characterizing the sensory responses is to attempt to learn an explicit functional
mapping (or "encoding model") between the stimulus and the neural response. Neural responses in the
auditory system have been classically modeled using spectrotemporal receptive fields which fit a linear
mapping between a time-frequency spectrogram-like representation and the neural response (Calabrese
et al., 2011). STRFs, however, are unable to predict much of the neural response variance in the auditory
cortex, particularly for natural sounds, due to prominent nonlinearities between the spectrogram and the
neural response (Sadagopan & Wang, 2009) that are not captured by the model (David et al., 2009;
Keshishian et al., 2020). Some prior studies have fit more complex encoding models, such as deep neural
networks (Harper et al., 2016; Keshishian et al., 2020), but these models still fail to explain much of the

neural response, and it is not clear how to measure integration windows from these types of complex,
nonlinear models. In contrast, our paradigm does not make any assumptions about the features that
underlie the response (e.g., spectrogram) or how these features are mapped to the neural response, and
thus is able to characterize the integration windows from highly nonlinear systems such as deep artificial
neural networks (Keshishian & Norman-Haignere, 2021).

An alternative approach that does not require fitting a complete encoding model is to simply measure
temporal characteristics of the neural response to a sound. For example, prior studies have measured
the rate of temporal modulations in the neural response (Bartlett & Wang, 2007), which has revealed that
the upper frequency limit at which the neural response will "phase-lock" to a stimulus frequency slows
from the periphery to the cortex. Phase locking, however, does not provide a measure of the neural
integration window since for example a fast neural modulation could be produced integrating information
across both a long or short window (e.g., integrating across many cycles of an oscillation or just a single
cycle). Other studies have attempted to measure the time period in the neural response that contains
information about a stimulus or stimulus property (sometimes referred to as the "encoding window")
(Theunissen & Miller, 1995). While useful, this measure is highly stimulus dependent and will vary with
the duration and information rate of sounds. Other studies have measured the extent to which the neural
response timecourse expands or contracts as the stimulus is stretched or compressed (Lerner et al.,
2014). However, the neural response will tend to expand and contract with the stimulus even if the
integration window remains fixed and this metric thus does not provide a good estimate of the integration
window or whether this window varies with the information rate. By contrast, the TCI method can directly
estimate the neural integration window, independent of many stimulus properties such as the information
rate and duration of sounds, as we show here.

Recent research has characterized the "intrinsic timescales" of neural systems, often by measuring the
autocorrelation of the neural response after removing stimulus-driven changes (Chaudhuri et al., 2014;
Fontanier et al., 2022; Manea et al., 2022; Murray et al., 2014; Zeraati et al., 2023). Intrinsic timescales
are distinct from integration windows, since they do not directly specify anything about the relationship
between the stimulus and the neural response. In contrast, integration windows are fundamentally
defined with respect to the stimulus. Empirically, however, intrinsic timescales have been shown to
increase across the cortical hierarchy, potentially analogous to the hierarchical organization that we
observe here. Thus, one possibility is that there are shared underlying neurobiological mechanisms (e.g.,
changes in synaptic integration time constants) that couple intrinsic timescales and stimulus integration
windows, and cause both to increase across the cortical hierarchy. Future research could explore this
link by applying the methods used here to computational models of intrinsic timescales (i.e. (Zeraati et
al., 2022)), which is feasible because our method is effective in nonlinear models.

Methodological choices, limitations, and future directions.

Many prior studies have demonstrated stimulus-driven effects that extend beyond the integration
timescales observed in this study. For example, stimulus-specific adaptation effects have been observed
over multi-second timescales (Herrmann et al., 2015; Ulanovsky, 2004; Ulanovsky et al., 2003; Valerio
et al., 2024), and effects of learning and memory can be observed across minutes, hours, and days
(Elgueda et al., 2019; K. Lu et al., 2018). Thus, there are likely multiple timescales over which stimuli can
influence the neural response, in part reflecting different neurobiological mechanisms (e.g., adaptation,
learning, memory, etc.). Our results demonstrate that the cumulative effects of these longer timescale
processes are small during the processing of natural sounds compared with the effects of stimulus

variation within the window. Indeed, we find that for virtually every unit in the auditory cortex there is a
time period from approximately 15 to 150 milliseconds, where the cross-context correlation closely tracks
the noise ceiling, indicating that surrounding stimuli have virtually no effect on the neural response. One
caveat is that the noise ceiling measured by the TCI paradigm will reflect all responses that are reliable
across a long sequence of sounds. Thus, any neural responses that change between two presentations
of the same sequence will be treated as noise. As a consequence, if there were learning and memory
processes that took place across repetitions of an entire sequence (Bartlett & Wang, 2005) our paradigm
would not be sensitive to them.

We used uniform stretching and compression (preserving pitch) because this alters the information rate
and duration of virtually all sound properties by a known magnitude, which is useful because we do not
know which features are important to the neural response. Recently, we showed that this approach is
able to reveal a near complete transition from time-yoked to rate-yoked information across the layers of
highly nonlinear deep artificial neural network (DANN) trained on challenging speech recognition tasks,
even though the models were only ever exposed to natural speech without any compression or stretching.
This finding demonstrates that our method can cleanly detect rate-yoked integration, even from a highly
complex nonlinear system, where the features and structures that underlie the system's response are
unknown. Nonetheless, a potential limitation of our approach is that uniform stretching and compression
are not entirely natural, and it is possible that biological systems might show greater evidence of rate-
yoked computation for more natural rate manipulations.

Our study focused on characterizing integration windows in the auditory cortex, across multiple regions
and layers. An important question for future research is how integration windows are structured beyond
the auditory cortex in higher-order cognitive and motor regions, as well as in associative multisensory
regions. It is possible that stimulus-driven integration windows continue to grow beyond the auditory
cortex (Chaudhuri et al., 2015; Manea et al., 2022; Murray et al., 2014), but remain yoked to absolute
time. Another possibility is that integration windows become increasingly flexible outside of the auditory
cortex in order to support adaptive computations that underlie perception and behavior (Piet et al., 2018).
These types of questions could be answered by applying our paradigm to neural responses beyond the
auditory cortex, testing longer segment durations and using multimodal stimuli.

───

427 Methods

The data from the paper come from three experiments. Experiment I compared integration windows
between different regions using natural speech, music, and ferret vocalizations. Experiment II examined
layer effects. Experiment III examined the importance of the information rate by comparing original,
stretched, and compressed sound stimuli. Experiments I and III were conducted at the École Normale
Supérieure (ENS) using chronic multi-electrode arrays, and Experiment II was conducted at Oregon
Health & Sciences University (OHSU) using laminar probes. The neurophysiological recording methods
for chronic arrays and laminar probes are described separately.

435 <u>Neurophysiological recordings with chronic arrays at ENS</u>

Animal preparation. Experiments were performed in adult female ferrets (Mustela putorius furo, ferret
A, B, and C) across one or both hemispheres (ferret A: left and right, ferret B, right, and ferret C: left).
The animals were 1–3 years of age, weighing 500–1000 g, and were housed in pairs or trios with a

normal day-night light cycle and unrestricted access to food and water. All ferrets were virgins.
Experiments were approved by the French Ministry of Agriculture (protocol authorization: 21022) and all
manipulations strictly comply with the European directives on the protection of animals used for scientific
purposes (2010/63/EU).

Surgical procedure. Chronic neurophysiological recordings were performed head-fixed after
implantation of a metal headpost attached to the skull (detailed procedure in (Chillale et al., 2023)). After
recovery from the initial headpost surgery, chronic floating multielectrode arrays were implanted in a
subsequent surgery. A 10 mm x 10 mm craniotomy was performed under anesthesia (1% isoflurane)
over auditory cortex. The large craniotomy allowed us to identify the different regions of the auditory
cortex (anterior/middle/posterior ectosylvian gyri) by visual inspection of the pseudosylvian and
suprasylvian sulci and additional landmarks. The animals were then implanted with floating multielectrode
arrays (Platinum/Iridium, MicroProbes, electrodes of impedance of 500 or 750 kΩ, 4 rows of 8 electrodes
with 0.4 µm distance between the electrodes) in primary or non-primary regions using a micromanipulator
and sealed with Vetbond (3M). When possible the dura was pulled over the implanted array. Finally the
removed skull flap was put back and sealed with Kwik-Sil (WPI) and bone cement (Palacos, Heraeus).
Animals could then recover for 1 week, with unrestricted access to food, water, and environmental
enrichment.

Recording procedure. Recordings were performed head-fixed in a soundproof chamber. Continuous
electrophysiological recordings were digitized, amplified, and recorded at 30 kHz using a digital
acquisition system (OpenEphys). We presented sounds to the animals via custom scripts implemented
using a MATLAB toolbox designed for low-latency sound delivery (PsychToolbox). We collected 19
recording sessions in Ferret A, 45 in ferret B, and 40 in ferret C.

Preprocessing. Electrode responses were common-averaged to the grand mean across all functional
electrodes of each array. We applied a 300-6000 Hz band-pass filter (2nd order Butterworth filter with 3dB
cutoff applied forward and backward). We then detected multi-unit activity by thresholding the signal (3
SD) and removing artifacts using PCA--based customized spike sorting routines written in MATLAB (J.
Fritz et al., 2003).

Neurophysiological recordings with acute laminar probes at OHSU

Animal preparation and surgical procedure. Adult male ferrets (aged 6–9 months) were surgically
implanted with a head post to stabilize the head and enable multiple small craniotomies for acute
electrophysiological recordings. Anesthesia was induced with ketamine (35 mg/kg) and xylazine (5
mg/kg) and maintained with isoflurane (0.5–2%) during the surgical procedure. The skin and muscle on
top of the cranium were removed and the surface of the skull was cleaned. Ten to twelve small surgical
screws were placed on the edge of the exposed skull as anchor points. The surface of the skull was
chemically etched (Optibond Universal, Kerr) and a thin layer of UV-cured dental cement (Charisma
Classic, Kultzer) was applied over the exposed surface. Two stainless steel head posts were aligned
along the midline and embedded with additional cement. Finally, cement was used to build a rim
extending out from the edges of the implant. The rim served the dual purpose of holding bandages over
the implant margin wounds and creating wells to hold saline over the recording sites. Once the implant
was finished, excess skin around it was removed, the wound around the implant was closed with sutures
and the animal was bandaged. Antibiotics and analgesics were administered as part of the post-op
recovery.

After at least 2 weeks following surgery, the animals were acclimated to a head-fixed posture, during
intervals starting at 5 min and increased 5–10 min every day. Food and liquid rewards were given during
these acclimation sessions to help the animals relax under restraint. Animals were considered ready for
recording when they could be restrained for more than 3 h without signs of distress (e.g., the animals
being relaxed enough to fall asleep).

Preprocessing. The putative location of A1 and dPEG was determined during the headpost implantation
surgery based on external landmarks: the posterior and medial edges of A1 falling, respectively, 13 mm
anterior to the occipital crest and 8 mm lateral to the center line, and dPEG immediately antero-lateral to
A1 (J. Bizley et al., 2005). To functionally confirm recording locations, we opened small craniotomies <1
mm diameter and performed preliminary mapping with tungsten electrodes (FH–Co. Electrodes, AM
Systems Amp, MANTA software (Englitz et al., 2013)). We measured the tuning of the recording regions
using rapid sequences of 100ms pure tones and used tonotopy to identify cortical fields. We specifically
looked for the frequency tuning inversion: high-low-high moving in an antero-lateral direction, which
marks the boundary between primary (A1) and secondary (dPEG) fields. At tonotopically mapped sites,
we performed acute recordings with 64-channel integrated UCLA probes (Du et al., 2011), digital head-
stages (RHD 128-Channel, Intan technologies) and OpenEphys data acquisition boxes and software
(Siegle et al., 2017). The probes were inserted approximately normal to the cortical surface, up to a depth
of ~1 mm from the dura surface. Due to spatial constraints of the recording site and apparatus,
penetrations deviated from normality up to 20°. The depth of recording sites and their location in
superficial areas A1 or PEG were confirmed by current source density analysis.

Raw voltage traces were processed with Kilosort 2 (Stringer et al., 2019), clustering and assigning spikes
to putative single neurons. The clusters were manually curated with Phy (Rossant et al., 2016). Units
were only kept for analysis if they maintained isolation and a stable firing rate over the course of the
experiment. Unit isolation was quantified as the percent overlap of the spike waveform distribution with
neighboring units and baseline activity. Isolation >95% was considered a single unit and kept for analysis.
We further filtered neurons based on the reliability of their responses, requiring a Pearson's correlation
>0.1 between PSTH responses to the same stimuli (3-10 repetitions, 20 Hz sampling) drawn from random
halves of repeated trials.

Experimental protocols

Experiment I. The goal of this experiment was to measure integration windows using a diverse set of
natural sounds. The stimuli tested included sequences composed of segments of varying durations
(ranging from 15.625 to 500 ms). Segments were excerpted from 24 natural sounds (each 500 ms)
spanning three different categories (8 sounds/category): speech, ferret vocalizations, and music. Each
sound was subdivided to create segments of the desired duration. Each natural sound was RMS-
normalized prior to segmentation. We presented the segments in two pseudorandom sequences, with all
segment durations and categories interleaved. Sequences varied in duration from 2 to 6 minutes
depending on the session and animal. Segments were concatenated using cross-fading to avoid click
artifacts (15.625 ms raised cosine window). Each stimulus was repeated several times to measure a
reliable noise ceiling (8 repetitions for most recording sessions).

Experiment II. The goal of this experiment was to replicate key findings from Experiment I as well as
characterize the organization of integration windows across cortical layers using acute laminar probes.
The stimuli tested were similar to those in Experiment I. The segments were excerpted from 20 natural

sound recordings that included ferret vocalizations, speech, music, and other animal calls/vocalizations
and environmental sounds. The segment durations spanned 16 to 500 milliseconds. The segments for
each duration were presented in a separate sequence, using two different random orders. The frequency
of the sounds was adjusted so that the mean excitation pattern across sounds was approximately white
to ensure sufficient power at all frequencies, including higher frequencies.

Experiment III. The goal of the experiment was to examine whether integration windows reflected
absolute time or the rate at which information varies in sound. Stimuli were similar to those for Experiment
I except that we only tested speech and ferret vocalizations (not music), and we altered the information
rate of the sounds by stretching and compressing the sounds (without altering the pitch). Sounds were
was stretched and compressed by a factor of $\sqrt{2}$ and $\sqrt{\frac{1}{2}}$ prior to subdivision into segments using a phase
vocoder with identity phase locking (Driedger, Müller, et al., 2014; Driedger, Muller, et al., 2014).
Segments were excerpted from 16 natural sounds (500 ms, half speech, half ferret vocalizations) using
the same range of segment durations as Experiment I. We also tested natural rate sounds without any
stretching or compression.

Data analysis

Selecting electrodes with a reliable response to sound. We selected units (single and multi-units)
with a reliable response to sound. Specifically, we measured the split-half Pearson correlation of the
response timecourse of each unit across all stimuli from two non-overlapping sets of stimulus repetitions
(odd vs. even). For Experiments I & II, we selected electrodes with a split-half correlation of at least 0.1,
where this correlation was highly significant ($p < 10^{-5}$). We used a lower absolute correlation threshold
(0.05) in Experiment III because we were focusing on PEG which has sparser and more variable
responses. We identified 693 units across 8 animals and 3 experiments that showed a reliable response
to natural sounds based on these criteria (Exp I: 345, Exp II: 231, and Exp III: 117). Significance was
measured using a permutation test where we permuted 100 ms segments of the neural timecourse (1000
permutations), and re-measured the correlation coefficient to build-up a null distribution. This null
distribution was then fitted with a Gaussian from which we then computed the tail probability of the
observed correlation value. We used a Gaussian fit rather than the empirical distribution so that we could
measure small p values that would be impossible to measure via counting.

Cross-context correlation. We measured integration windows using an analysis developed in prior work
for estimating context invariance. The neural response timecourse to each sequence is organized as a
segment x time matrix, where each row contains the response timecourse surrounding a single segment,
aligned to segment onset (**Figure 1B**). A separate matrix is calculated for each of the two contexts, using
the two different segment orderings/sequences. Corresponding rows contain the response timecourse to
the same segment across the two different contexts. Different columns contain the responses from
multiple segments for a fixed time lag relative to segment onset.

To estimate context invariance, we correlate corresponding columns (i.e. columns with the same lag)
across the matrices from each of the two contexts (**Figure 1B**). If the integration window is less than the
segment duration, there will be a moment when the response is the same across contexts yielding a
correlation of 1. Because the correlation will never be 1 due to noise, we computed a noise ceiling for our
measure by performing the same calculation when the context was identical using repeated presentations
of each sequence.

Model-estimated windows. In our prior work, we developed a computational model that is capable of using the cross-context correlation from multiple segment durations and lags to estimate a single underlying integration window (S. V. Norman-Haignere et al., 2022). We have extensively tested and validated the method in this prior work, showing that we can recover accurate integration window estimates from multiple ground truth models using noisy data.

The complete details of the method are described in our prior paper (S. V. Norman-Haignere et al., 2022). In brief, the integration window is parametrized with a Gamma distribution. We then make a prediction for the noise-free cross-context correlation by measuring the relative overlap of the window with shared central segments vs. surrounding context segments. This noise free prediction is then multiplied by the empirically measured noise ceiling to arrive at a prediction for the measured cross-context correlation. We determine the best fitting window by varying the window's width and center, and selecting the window that yields the best prediction of the cross-context correlation. The width is defined as the central 75% of the window with the smallest mass and the center of the window is defined as the window's median. We also varied the shape of the window, from more exponential to more Gaussian, although we have found that the shape does not substantially influence the predictions and results. The complete model includes several additional details that we have found improve the accuracy of the model in simulations. A detailed description and derivation of the added features is described in our prior paper.

Anatomical analysis - cortical surface maps. There is variability in the functional location of cortical fields across animals. To create a cortical surface representation of all recorded units and to account for functional variability, we localized each electrode array on a standardized template tonotopic map, computed across many ferrets (J. K. Bizley & King, 2009; Nelken et al., 2004). The location of each array was selected based on the frequency selectivity of the electrode responses within that array, the tonotopic gradient found across the array and its position relative to anatomical landmarks. Electrode recordings from different sessions were assumed to sample different cells, and each recording was therefore plotted at the same location with a small amount of jitter so they could be distinguished. The jitter was sampled using polar coordinates (radius was sampled from a uniform distribution from 0 to 360° and distance was sampled from a uniform distribution from 20 to 240 micrometers). Recordings from the left hemisphere were mirrored onto the right hemisphere for visualization purposes. We measured the distance on the cortical surface of each unit to a reference point in the middle of primary auditory cortex.

Anatomical analysis - laminar assignment. Laminar analysis was conducted following a method presented in our previous paper (López Espejo & David, 2024). In summary, cortical layers were categorized into three groups (layers 1–3, layer 4, and layers 5–6) and assigned to each electrode of the silicon probes using a custom graphical user interface. Boundaries between these groups were determined based on characteristic features of the local field potential (LFP) signal. The LFP signal was extracted by applying a zero-phase shift, 4th-order Butterworth low-pass filter (250 Hz, filter-filter method) to the raw electrophysiological data, followed by downsampling to 500 Hz. Key features used for layer identification included the current source density (CSD) sink and source patterns elicited by broadband noise bursts centered on each site's best frequency (BF), which align with established auditory-evoked CSD patterns in the auditory cortex of other species (Schaefer et al., 2015). Additionally, the relative power of high-frequency (>40 Hz) versus low-frequency (<30 Hz) components of the LFP was analyzed to delineate layers 1–3, leveraging prior findings of increased high-frequency power in superficial cortical layers (Mendoza-Halliday et al., 2024). Finally, layer 4 was identified by detecting a drop in LFP coherence between adjacent channels at its upper boundary, consistent with earlier reports (Davis et al.,

2023; Maier et al., 2010). Each unit was assigned to a cortical layer based on the electrode exhibiting the
highest spike amplitude for that unit.

Rate-yoking index. Compressing and stretching change the rate at which information varies by a known
magnitude. We used a stretching and compression factor of $\sqrt{2}$ such that there is an octave difference in
the rate at which information varies between stretched and compressed sounds. Thus, if the integration
window reflects the information rate, we should observe a one-octave change in the integration window.
We therefore computed a rate-yoking index by computing the difference in integration windows on an
octave scale between stretched and compressed speech and dividing this difference by that which would
be expected for a rate-yoked window (i.e., a one-octave change):

616
$$Rate-yoking\ index\ = \frac{log_2\ (i_{stretched}) - log_2\ (i_{compressed})}{log_2(2)}$$

where i is the measured integration window for stretched/compressed sounds. An index of 0 indicates a
fixed, time-yoked window, while a value of 1 indicates a fully rate-yoked window. Note that the
denominator in the above equation happens to be 1, and thus has no effect on the result, since there
was an octave difference in the information rate for stretched and compressed sounds. If we had used a
different stretching and compression factor, then it would be different and would need to be included. The
rate-yoking index is not strictly bound between 0 and 1 and noise will tend to cause the estimate to exceed
these bounds. A value less than 0 indicates that the integration window was longer for compressed
sounds and a value greater than 1 indicates that the difference between stretched and compressed
integration windows exceeded 1 octave.

Statistics. Unless otherwise noted, we used a linear mixed effects (LME) model to evaluate the
significance of our effects (fitlme.m and coefTest.m in MATLAB; Satterthwaite approximation was used
to estimate degrees of freedom when measuring significance). Whenever possible, we included random
effects to account for variation across sessions from day to day, using diagonal covariance matrices to
prevent overfitting/singular results. As is typical, we do not include random effects for animals because
of the small number of animals in our experiment. For categorical variables, the category with the most
samples was always used as the baseline and not modeled explicitly in the linear model. Integration
window estimates were always logarithmically transformed before being fit by the LME model.

For Experiment I, to test whether integration windows increased across the cortical hierarchy, we
modeled the integration window as a linear function of distance to primary auditory cortex, including
random intercepts and slopes for different recording sessions:

637
$$y \sim 1 + distance + (1 + distance\,|\,session)$$

638 where y reflects either the integration width or center (logarithmically transformed). The sample
639 dimension was unit (single or multi-unit).

640 To test whether integration windows varied across regions, we used the following model:

641
$$y \sim 1 + region + (1 + region\,|\,session)$$

642 Region was a binary categorical variable (MEG vs. PEG).

643 For Experiment II, which used laminar probes, we investigated effects of layer and region using the
644 following model:

$$y \sim 1 + layer + region + (1|session)$$

Layer was a categorical variable with three levels (granular, infragranular, supragranular). Region was
again a binary variable (MEG, PEG). A different insertion was used for each recording session, and as a
consequence, session was partially confounded with region and layer and we found that a model with
random slopes did not converge (and these terms were thus removed). We tested whether the effect of
region was significantly different (larger) than the effect of layer by directly comparing their coefficients
against a null model that assumed no difference. To make this a fair comparison, we collapsed our layers
from three levels to two levels to make them comparable to that for each region. To be conservative, we
chose the two levels so that the effect of layers would be maximal thus minimizing the likelihood that the
effect of the region would be even larger than the effect of layer. To this end, we separated out the
infragranular layers from the granular and supragranular layers (infragranular vs.
granular/supragranular), because the infragranular layers showed the most different integration windows
(**Figure 3**).

For Experiment III, we used the following model to investigate the effect of rate, controlling for any
possible difference due to sound category (ferrets vs. speech):

$$y \sim 1 + rate + category + (1|unit) + (1|session)$$

Rate and category were categorical variables with three (original, compressed, speech) and two (speech,
ferrets) levels, respectively. We did not include random slopes for session in this model because the
model was not able to estimate them (variance terms for these random effects were either undefined or
close to 0).

References

Angeloni, C., & Geffen, M. N. (2018). Contextual modulation of sound processing in the auditory cortex.
Current Opinion in Neurobiology, *49*, 8–15. https://doi.org/10.1016/j.conb.2017.10.012
Atiani, S., David, S. V., Elgueda, D., Locastro, M., Radtke-Schuller, S., Shamma, S. A., & Fritz, J. B.
(2014). Emergent selectivity for task-relevant stimuli in higher-order auditory cortex. *Neuron*,
82(2), 486–499. https://doi.org/10.1016/j.neuron.2014.02.029
Baratham, V. L., Dougherty, M. E., Hermiz, J., Ledochowitsch, P., Maharbiz, M. M., & Bouchard, K. E.
(2022). Columnar Localization and Laminar Origin of Cortical Surface Electrical Potentials. *The
Journal of Neuroscience*, *42*(18), 3733–3748. https://doi.org/10.1523/JNEUROSCI.1787-
21.2022
Bartlett, E. L., & Wang, X. (2005). Long-lasting modulation by stimulus context in primate auditory
cortex. *Journal of Neurophysiology*, *94*(1), 83–104. https://doi.org/10.1152/jn.01124.2004
Bartlett, E. L., & Wang, X. (2007). Neural representations of temporally modulated signals in the
auditory thalamus of awake primates. *Journal of Neurophysiology*, *97*(2), 1005–1017.
https://doi.org/10.1152/jn.00593.2006
Bizley, J. K., & King, A. J. (2009). Visual influences on ferret auditory cortex. *Hearing Research*, *258*(1),
55–63. https://doi.org/10.1016/j.heares.2009.06.017
Bizley, J., Nodal, F., Nelken, I., & King, A. (2005). Functional Organization of Ferret Auditory Cortex.
Cerebral Cortex (New York, N.Y. : 1991), *15*, 1637–1653. https://doi.org/10.1093/cercor/bhi042
Calabrese, A., Schumacher, J. W., Schneider, D. M., Paninski, L., & Woolley, S. M. N. (2011). A
Generalized Linear Model for Estimating Spectrotemporal Receptive Fields from Responses to

Natural Sounds. *PLOS ONE*, *6*(1), e16104. https://doi.org/10.1371/journal.pone.0016104

Cavanagh, S. E., Hunt, L. T., & Kennerley, S. W. (2020). A Diversity of Intrinsic Timescales Underlie Neural Computations. *Frontiers in Neural Circuits*, *14*, 615626. https://doi.org/10.3389/fncir.2020.615626

Chaudhuri, R., Bernacchia, A., & Wang, X.-J. (2014). A diversity of localized timescales in network activity. *eLife*, *3*, e01239. https://doi.org/10.7554/eLife.01239

Chaudhuri, R., Knoblauch, K., Gariel, M.-A., Kennedy, H., & Wang, X.-J. (2015). A Large-Scale Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex. *Neuron*, *88*(2), 419–431. https://doi.org/10.1016/j.neuron.2015.09.008

Chen, C., Read, H. L., & Escabí, M. A. (2012). Precise feature based time scales and frequency decorrelation lead to a sparse auditory code. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(25), 8454–8468. https://doi.org/10.1523/JNEUROSCI.6506-11.2012

Chillale, R. K., Shamma, S., Ostojic, S., & Boubenec, Y. (2023). Dynamics and maintenance of categorical responses in primary auditory cortex during task engagement. *eLife*, *12*, e85706. https://doi.org/10.7554/eLife.85706

Cichy, R. M., & Kaiser, D. (2019). Deep Neural Networks as Scientific Models. *Trends in Cognitive Sciences*, *23*(4), 305–317. https://doi.org/10.1016/j.tics.2019.01.009

David, S. V., Mesgarani, N., Fritz, J. B., & Shamma, S. A. (2009). Rapid Synaptic Depression Explains Nonlinear Modulation of Spectro-Temporal Tuning in Primary Auditory Cortex by Natural Stimuli. *Journal of Neuroscience*, *29*(11), 3374–3386. https://doi.org/10.1523/JNEUROSCI.5249-08.2009

Davis, Z. W., Dotson, N. M., Franken, T. P., Muller, L., & Reynolds, J. H. (2023). Spike-phase coupling patterns reveal laminar identity in primate cortex. *eLife*, *12*, e84512. https://doi.org/10.7554/eLife.84512

Deneux, T., Kempf, A., Daret, A., Ponsot, E., & Bathellier, B. (2016). Temporal asymmetries in auditory coding and perception reflect multi-layered nonlinearities. *Nature Communications*, *7*(1), 12682. https://doi.org/10.1038/ncomms12682

Driedger, J., Muller, M., & Ewert, S. (2014). Improving Time-Scale Modification of Music Signals Using Harmonic-Percussive Separation. *IEEE Signal Processing Letters*, *21*(1), 105–109. https://doi.org/10.1109/LSP.2013.2294023

Driedger, J., Müller, M., Fourier, J., & Shannon, C. (2014). *TSM Toolbox: MATLAB Implementations of Time-Scale Modification Algorithms*.

Du, J., Blanche, T. J., Harrison, R. R., Lester, H. A., & Masmanidis, S. C. (2011). Multiplexed, High Density Electrophysiology with Nanofabricated Neural Probes. *PLOS ONE*, *6*(10), e26204. https://doi.org/10.1371/journal.pone.0026204

Dubreuil, A., Valente, A., Beiran, M., Mastrogiuseppe, F., & Ostojic, S. (2022). The role of population structure in computations through neural dynamics. *Nature Neuroscience*, *25*(6), 783–794. https://doi.org/10.1038/s41593-022-01088-4

Elgueda, D., Duque, D., Radtke-Schuller, S., Yin, P., David, S. V., Shamma, S. A., & Fritz, J. B. (2019). State-dependent encoding of sound and behavioral meaning in a tertiary region of the ferret auditory cortex. *Nature Neuroscience*, *22*(3), 447–459. https://doi.org/10.1038/s41593-018-0317-8

Englitz, B., David, S. V., Sorenson, M. D., & Shamma, S. A. (2013). MANTA—an open-source, high density electrophysiology recording suite for MATLAB. *Frontiers in Neural Circuits*, *7*. https://doi.org/10.3389/fncir.2013.00069

Esposito, M., Valente, G., Plasencia-Calaña, Y., Dumontier, M., Giordano, B. L., & Formisano, E. (2024). Bridging auditory perception and natural language processing with semantically informed deep neural networks. *Scientific Reports*, *14*(1), 20994. https://doi.org/10.1038/s41598-024-71693-9

Fontanier, V., Sarazin, M., Stoll, F. M., Delord, B., & Procyk, E. (2022). Inhibitory control of frontal metastability sets the temporal signature of cognition. *eLife*, *11*, e63795. https://doi.org/10.7554/eLife.63795

Fritz, J. B., Elhilali, M., & Shamma, S. A. (2007). Adaptive Changes in Cortical Receptive Fields Induced by Attention to Complex Sounds. *J Neurophysiol*, *98*.

Fritz, J., Elhilali, M., & Shamma, S. (2005). Active listening: Task-dependent plasticity of spectrotemporal receptive fields in primary auditory cortex. *Hearing Research*, *206*(1–2), 159–176. https://doi.org/10.1016/j.heares.2005.01.015

Fritz, J., Shamma, S., Elhilali, M., & Klein, D. (2003). Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. *Nature Neuroscience*, *6*(11), 1216–1223. https://doi.org/10.1038/nn1141

Harper, N. S., Schoppe, O., Willmore, B. D. B., Cui, Z., Schnupp, J. W. H., & King, A. J. (2016). Network Receptive Field Modeling Reveals Extensive Integration and Multi-feature Selectivity in Auditory Cortical Neurons. *PLOS Computational Biology*, *12*(11), e1005113. https://doi.org/10.1371/journal.pcbi.1005113

Herrmann, B., Henry, M. J., Fromboluti, E. K., McAuley, J. D., & Obleser, J. (2015). Statistical context shapes stimulus-specific adaptation in human auditory cortex. *Journal of Neurophysiology*, *113*(7), 2582–2591. https://doi.org/10.1152/jn.00634.2014

Kell, A. J. E., Yamins, D. L. K., Shook, E. N., Norman-Haignere, S. V., & McDermott, J. H. (2018). A Task-Optimized Neural Network Replicates Human Auditory Behavior, Predicts Brain Responses, and Reveals a Cortical Processing Hierarchy. *Neuron*, *98*(3), 630-644.e16. https://doi.org/10.1016/j.neuron.2018.03.044

Kello, C. T., Bella, S. D., Médé, B., & Balasubramaniam, R. (2017). Hierarchical temporal structure in music, speech and animal vocalizations: Jazz is like a conversation, humpbacks sing like hermit thrushes. *Journal of The Royal Society Interface*, *14*(135), 20170231. https://doi.org/10.1098/rsif.2017.0231

Keshishian, M., Akbari, H., Khalighinejad, B., Herrero, J. L., Mehta, A. D., & Mesgarani, N. (2020). Estimating and interpreting nonlinear receptive field of sensory neural responses with deep neural network models. *eLife*, *9*, e53445. https://doi.org/10.7554/eLife.53445

Keshishian, M., & Norman-Haignere, S. V. (2021). Understanding Adaptive, Multiscale Temporal Integration In Deep Speech Recognition Systems. *Advances in Neural Information Processing Systems*.

Kietzmann, T. C., McClure, P., & Kriegeskorte, N. (2019). Deep Neural Networks in Computational Neuroscience. In *Oxford Research Encyclopedia of Neuroscience*. https://doi.org/10.1093/acrefore/9780190264086.013.46

Landemard, A., Bimbard, C., Demené, C., Shamma, S., Norman-Haignere, S., & Boubenec, Y. (2020). *Distinct higher-order representations of natural sounds in human and ferret auditory cortex*. https://doi.org/10.1101/2020.09.30.321695

Leonard, M. K., Gwilliams, L., Sellers, K. K., Chung, J. E., Xu, D., Mischler, G., Mesgarani, N., Welkenhuysen, M., Dutta, B., & Chang, E. F. (2024). Large-scale single-neuron speech sound encoding across the depth of human cortex. *Nature*, *626*(7999), 593–602. https://doi.org/10.1038/s41586-023-06839-2

Lerner, Y., Honey, C. J., Katkov, M., & Hasson, U. (2014). Temporal scaling of neural responses to compressed and dilated natural speech. *Journal of Neurophysiology*, *111*(12), 2433–2444. https://doi.org/10.1152/jn.00497.2013

Lerner, Y., Honey, C. J., Silbert, L. J., & Hasson, U. (2011). Topographic mapping of a hierarchy of temporal receptive windows using a narrated story. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *31*(8), 2906–2915. https://doi.org/10.1523/JNEUROSCI.3684-10.2011

Li, Y., Anumanchipalli, G. K., Mohamed, A., Chen, P., Carney, L. H., Lu, J., Wu, J., & Chang, E. F. (2023). Dissecting neural computations in the human auditory pathway using deep neural networks for speech. *Nature Neuroscience*, *26*(12), 2213–2225. https://doi.org/10.1038/s41593-023-01468-4

López Espejo, M., & David, S. V. (2024). A sparse code for natural sound context in auditory cortex. *Current Research in Neurobiology*, *6*, 100118. https://doi.org/10.1016/j.crneur.2023.100118

Lu, K., Liu, W., Zan, P., David, S. V., Fritz, J. B., & Shamma, S. A. (2018). Implicit Memory for Complex

Sounds in Higher Auditory Cortex of the Ferret. *Journal of Neuroscience*, *38*(46), 9955–9966.
https://doi.org/10.1523/JNEUROSCI.2118-18.2018

Lu, T., Liang, L., & Wang, X. (2001). Temporal and rate representations of time-varying signals in the
auditory cortex of awake primates. *Nature Neuroscience*, *4*(11), 1131–1138.
https://doi.org/10.1038/nn737

Maier, A., Adams, G. K., Aura, C., & Leopold, D. A. (2010). Distinct Superficial and Deep Laminar
Domains of Activity in the Visual Cortex during Rest and Stimulation. *Frontiers in Systems*
Neuroscience, *4*. https://doi.org/10.3389/fnsys.2010.00031

Manea, A. M., Zilverstand, A., Ugurbil, K., Heilbronner, S. R., & Zimmermann, J. (2022). Intrinsic
timescales as an organizational principle of neural processing across the whole rhesus
macaque brain. *eLife*. https://doi.org/10.7554/eLife.75540

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband shifts in local field potential
power spectra are correlated with single-neuron spiking in humans. *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience, *29*(43), 13613–13620.
https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Mante, V., Sussillo, D., Shenoy, K. V., & Newsome, W. T. (2013). Context-dependent computation by
recurrent dynamics in prefrontal cortex. *Nature*, *503*(7474), 78–84.
https://doi.org/10.1038/nature12742

Mendoza-Halliday, D., Major, A. J., Lee, N., Lichtenfeld, M. J., Carlson, B., Mitchell, B., Meng, P. D.,
Xiong, Y. (Sophy), Westerberg, J. A., Jia, X., Johnston, K. D., Selvanayagam, J., Everling, S.,
Maier, A., Desimone, R., Miller, E. K., & Bastos, A. M. (2024). A ubiquitous spectrolaminar motif
of local field potential power across the primate cortex. *Nature Neuroscience*, *27*(3), 547–560.
https://doi.org/10.1038/s41593-023-01554-7

Meyer, A. F., Williamson, R. S., Linden, J. F., & Sahani, M. (2016). Models of Neuronal Stimulus-
Response Functions: Elaboration, Estimation, and Evaluation. *Frontiers in Systems*
Neuroscience, *10*, 109. https://doi.org/10.3389/fnsys.2016.00109

Mizrahi, A., Shalev, A., & Nelken, I. (2014). Single neuron and population coding of natural sounds in
auditory cortex. *Current Opinion in Neurobiology*, *24*(1), 103–110.
https://doi.org/10.1016/j.conb.2013.09.007

Murray, J. D., Bernacchia, A., Freedman, D. J., Romo, R., Wallis, J. D., Cai, X., Padoa-Schioppa, C.,
Pasternak, T., Seo, H., Lee, D., & Wang, X.-J. (2014). A hierarchy of intrinsic timescales across
primate cortex. *Nature Neuroscience*, *17*(12), 1661–1663. https://doi.org/10.1038/nn.3862

Nelken, I., Bizley, J. K., Nodal, F. R., Ahmed, B., Schnupp, J. W. H., & King, A. J. (2004). Large-Scale
Organization of Ferret Auditory Cortex Revealed Using Continuous Acquisition of Intrinsic
Optical Signals. *Journal of Neurophysiology*, *92*(4), 2574–2588.
https://doi.org/10.1152/jn.00276.2004

Norman-Haignere, S., Orhan, P., King, J.-R., & Boubenec, ves. (2024, March). Hierarchical, structure-
yoked integration spontaneously emerges with self-supervised training on speech. *Poster*
Session. Computational and Systems Neuroscience (COSYNE) Conference, Lisbon, Portugal.

Norman-Haignere, S. V., Long, L. K., Devinsky, O., Doyle, W., Irobunda, I., Merricks, E. M., Feldstein,
N. A., McKhann, G. M., Schevon, C. A., Flinker, A., & Mesgarani, N. (2022). Multiscale temporal
integration organizes hierarchical computation in human auditory cortex. *Nature Human*
Behaviour, *6*(3), 455–469. https://doi.org/10.1038/s41562-021-01261-y

Norman-Haignere, S. V., & McDermott, J. H. (2018). Neural responses to natural and model-matched
stimuli reveal distinct computations in primary and nonprimary auditory cortex. *PLOS Biology*,
16(12), e2005127. https://doi.org/10.1371/journal.pbio.2005127

Pellegrino, F., Coupé, C., & Marsico, E. (2011). Across-Language Perspective on Speech Information
Rate. *Language*, *87*(3), 539–558.

Perez-Nieves, N., Leung, V. C. H., Dragotti, P. L., & Goodman, D. F. M. (2021). Neural heterogeneity
promotes robust learning. *Nature Communications*, *12*(1), 5791. https://doi.org/10.1038/s41467-
021-26022-3

Piet, A. T., El Hady, A., & Brody, C. D. (2018). Rats adopt the optimal timescale for evidence integration
in a dynamic environment. *Nature Communications*, *9*(1), 4265. https://doi.org/10.1038/s41467-

018-06561-y

Poeppel, D. (2003). The analysis of speech in different temporal integration windows: Cerebral
lateralization as 'asymmetric sampling in time.' *Speech Communication*, *41*(1), 245–255.
https://doi.org/10.1016/S0167-6393(02)00107-3

Ray, S., & Maunsell, J. H. R. (2011). Different origins of gamma rhythm and high-gamma activity in
macaque visual cortex. *PLoS Biology*, *9*(4), e1000610.
https://doi.org/10.1371/journal.pbio.1000610

Rossant, C., Kadir, S. N., Goodman, D. F. M., Schulman, J., Hunter, M. L. D., Saleem, A. B., Grosmark,
A., Belluscio, M., Denfield, G. H., Ecker, A. S., Tolias, A. S., Solomon, S., Buzsáki, G.,
Carandini, M., & Harris, K. D. (2016). Spike sorting for large, dense electrode arrays. *Nature*
Neuroscience, *19*(4), 634–641. https://doi.org/10.1038/nn.4268

Sadagopan, S., & Wang, X. (2009). Nonlinear Spectrotemporal Interactions Underlying Selectivity for
Complex Sounds in Auditory Cortex. *Journal of Neuroscience*, *29*(36), 11192–11202.
https://doi.org/10.1523/JNEUROSCI.1286-09.2009

Schaefer, M. K., Hechavarría, J. C., & Kössl, M. (2015). Quantification of mid and late evoked sinks in
laminar current source density profiles of columns in the primary auditory cortex. *Frontiers in*
Neural Circuits, *9*. https://doi.org/10.3389/fncir.2015.00052

Siegle, J. H., López, A. C., Patel, Y. A., Abramov, K., Ohayon, S., & Voigts, J. (2017). Open Ephys: An
open-source, plugin-based platform for multichannel electrophysiology. *Journal of Neural*
Engineering, *14*(4), 045003. https://doi.org/10.1088/1741-2552/aa5eea

Spitmaan, M., Seo, H., Lee, D., & Soltani, A. (2020). Multiple timescales of neural dynamics and
integration of task-relevant signals across cortex. *Proceedings of the National Academy of*
Sciences, *117*(36), 22522–22531. https://doi.org/10.1073/pnas.2005993117

Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C. B., Carandini, M., & Harris, K. D. (2019).
Spontaneous behaviors drive multidimensional, brainwide activity. *Science*, *364*(6437),
eaav7893. https://doi.org/10.1126/science.aav7893

Theunissen, F., & Miller, J. P. (1995). Temporal encoding in nervous systems: A rigorous definition.
Journal of Computational Neuroscience, *2*(2), 149–162. https://doi.org/10.1007/BF00961885

Ulanovsky, N. (2004). Multiple Time Scales of Adaptation in Auditory Cortex Neurons. *Journal of*
Neuroscience, *24*(46), 10440–10453. https://doi.org/10.1523/JNEUROSCI.1905-04.2004

Ulanovsky, N., Las, L., & Nelken, I. (2003). Processing of low-probability sounds by cortical neurons.
Nature Neuroscience, *6*(4), 391–398. https://doi.org/10.1038/nn1032

Valerio, P., Rechenmann, J., Joshi, S., De Franceschi, G., & Barkat, T. R. (2024). Sequential
maturation of stimulus-specific adaptation in the mouse lemniscal auditory system. *Science*
Advances, *10*(1), eadi7624. https://doi.org/10.1126/sciadv.adi7624

Yamins, D. L. K., & DiCarlo, J. J. (2016). Using goal-driven deep learning models to understand
sensory cortex. *Nature Neuroscience*, *19*(3), 356–365. https://doi.org/10.1038/nn.4244

Zeraati, R., Engel, T. A., & Levina, A. (2022). A flexible Bayesian framework for unbiased estimation of
timescales. *Nature Computational Science*, *2*(3), 193–204. https://doi.org/10.1038/s43588-022-
00214-3

Zeraati, R., Shi, Y.-L., Steinmetz, N. A., Gieselmann, M. A., Thiele, A., Moore, T., Levina, A., & Engel,
T. A. (2023). Intrinsic timescales in the visual cortex change with selective attention and reflect
spatial connectivity. *Nature Communications*, *14*(1), 1858. https://doi.org/10.1038/s41467-023-
37613-7

Supplementary Figures

Supplementary Figure 1 | Examples of cross-context correlation for multi-units in different regions of the auditory cortex. Examples of the CCC (blue) and noise ceiling (black) from example

multi-units from primary and non-primary ferret auditory cortex. For all units, there is a lag and segment
duration for which CCC equals the noise ceiling, indicating a context-invariant response. The segment
duration needed to achieve a context-invariant response varies substantially across units. The location
of each recorded multi-unit is indicated to the left of each row.

Supplementary Figure 2 | Relationship between the center and width of the integration window.
A, Integration windows were estimated using a parametric window (Gamma distribution) with a varying
width and center. We estimated the width and center that best predicted the cross-context correlation,
excluding parameters that yielded a non-causal window. This panel plots examples of the parameter
window as a function of the window's width (x-axis) and center (y-axis). Combination of parameters that
led to acausal windows were excluded (gray box with red dashed line) because they are not biologically
possible. **B**, Scatter plot of estimated integration centers vs widths for all units with a best-fit linear function
overlaid (orange line). Blue line shows the minimum possible center for a causal, Gamma-distributed
window.

Supplementary Figure 3 | Organisation of integration centers across neural populations in the
auditory cortex. A, Anatomical map of model-estimated integration centers in three animals (window
center: median of the interval containing 75% of the window's mass). B, The difference between the
centers of the integration windows between pairs of units as a function of their spatial distance,
demonstrating that nearby units have more similar centers. C, Centers of the integration windows as a
function of distance to the primary auditory cortex (PAC) (see color map in inset). D, Histograms of the
centers of the integration windows for primary (MEG) and non-primary auditory cortex (PEG) showing
substantial diversity across units and hierarchical organization.

Supplementary Figure 4 | Temporal integration is similar for different stimulus categories. A,
Histograms of model-fitted widths and centers of neurons in primary and non-primary auditory cortex
across the three sound categories tested (Ferret vocalizations, Speech, Music). B, Normalized median
cross-context correlation split by sound category across the two cortical areas. C, Scatter plots of the
model-fitted integration centers and widths as a function of distance to primary auditory cortex for all three
sound categories tested.

A Median cross-context correlation for original, stretched, and compressed sounds across categories

Supplementary Figure 5 | Time-yoked integration is consistent across sound categories. A, Median normalized CCC across all recorded units for stretched, original, and compressed stimuli for both sound categories (Ferret vocalizations and Speech). **B**, Integration windows of all units for compressed (x-axis) and stretched (y-axis) stimuli for both sound categories, separately (speech in yellow and ferret vocalizations in blue). Green and purple lines show the prediction from a time-yoked and rate-yoked response. **C**, Reliability of rate-yoking index across two independent data splits, separately for each category. **D**, Reliability of integration windows averaged across stimuli rates for comparison, again separately for each category.