Title: Tracking changes in birds' interaction milieu

Authors: Stanislas Rigal[1*], Vincent Devictor[2], Vasilis Dakos[2]

[1]TETIS, Université de Montpellier, AgroParisTech, Cirad, CNRS, INRAE, Montpellier, France

[2]ISEM, Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France

* Corresponding author: stanislas.rigal@inrae.fr

Conflict of Interest: The authors have no conflict of interest to disclose.

Author Contributions: S. Rigal, V. Devictor and V. Dakos conceived the ideas and designed

methodology; S. Rigal analysed the data and led the writing of the manuscript. All authors contributed

critically to the drafts and gave final approval for publication.

Abstract

As biodiversity is declining, the dynamics of species interactions is a growing conservation concern.

However, estimating and monitoring explicit species interactions across large spatial and temporal

scales remain challenging. An alternative and yet under-explored approach is to track whether and how

the interaction milieu, defined as the background of all realised interactions, is changing in space and

time. Here, we assess changes in the interaction milieu of common bird communities in France. We

estimate associated species pairs using spatial and temporal information for 109 species monitored

across 1,969 sites during 17 years. We validate the ecological significance of associated species pairs

by testing the relationship between the propensity to be associated and species functional proximity or

shared habitat preference. We reconstruct association networks for these intra-guild bird communities

and track temporal changes in network layout in terms of size, density of links, modularity and degree

distribution. We show that, beyond changes usually documented based on species numbers and

abundances, the interaction milieu is also changing non-randomly. Communities become smaller with a

similar relative number of associations that becomes unevenly distributed through time. These

structural changes vary among bird communities according to their habitat and may impact community

functioning and how communities can cope with global change.

Keywords: Biotic homogenisation, bird community, Convergent Cross Mapping, co-occurrence,

network, species associations.

1. Introduction

Vertebrate communities have been experiencing rapid and profound changes in recent decades, mainly documented through variations in species numbers, abundances, or distributions (Kampichler et al. 2012, Koerner et al. 2017). Changes affecting vertebrate communities have been also analysed through the lens of species ecological traits, e.g. by using community weighted indices that have highlighted a general pattern of homogenisation (Clavel et al. 2011, Davey et al. 2012, Li et al. 2018). However, addressing changes in the structure of the interaction networks underlining these communities have been more challenging (McGill et al. 2006). Any change in sign, strength or number of interactions has an effect on the network structure of a community and can strongly affect community dynamics (Lyons et al. 2016) and functioning (e.g. productivity, efficiency of resource use) (Fontaine and Thébault 2015, Besson et al. 2019). Modifications affecting networks with pair-wise interactions of a given type such as plant-pollinator or predator-prey networks have been related to specific consequences for the conservation of these communities (Tylianakis et al. 2010). However, isolating specific types of interactions remains challenging in many communities, in particular in large-scale studies (Barner et al. 2016). The *interaction milieu* was therefore proposed as a concept to refer to the resultant biotic interaction background to which an individual is exposed when arriving in a community (McGill et al. 2006). This notion thus enables to focus on community changes beyond species loss or turnover when addressing conservation issues in communities for which explicit interactions are out of reach.

Bird communities, in particular, have been extensively studied in particular in terms of changes in species composition at large scale, highlighting a visible and rapid effect of anthropogenic pressures leading to a biotic and functional homogenisation (Kampichler et al. 2012, Lindström et al. 2013, Gaüzère et al. 2015). More locally changes in bird communities have also been assessed through

dedicated network analyses based on food webs (Lurgi et al. 2012), plant-frugivore networks (Ramos-

Robles et al. 2016), competition for nests (Orchan et al. 2013) and specific types of bird communities

such as mixed-species flocks (Goodale et al. 2015, Zou et al. 2018). The observed shifts in species

composition indicate that the interaction milieu in common bird communities is also likely to be

altered, although the extent and direction of this change may depend on the community's environment

(Rigal et al. 2022). In particular, in the context of the sharp decline in the abundance of birds breeding

in open environments (Reif and Hanzelka 2020, Rigal and Knape 2024, Gómez-Catasús et al. 2025),

we can expect specific changes in the interaction milieu of these communities (e.g. in farmland, open

environment) compared to other environments (e.g. forest, urban environment).

The interaction milieu does not aim at mimicking a definite network of interactions, but rather

corresponds to the background of interaction. The distribution of species ecological traits has been

initially proposed to assess the interaction milieu in plant communities (McGill et al. 2006) and later

extended to microbial and soil communities (Jaillard et al. 2018, Davison et al. 2024). For mobile

species such as birds, large-scale monitoring data have been used to infer key elements of the

interaction milieu through species association based on community assembly processes: species with

similar functional traits or habitats may either tend to repulse (or even exclude) each other because of

resource competition, under the limiting similarity hypothesis (Martin and Bonier 2018) or attract each

other through social information exchange and facilitation processes (Gil et al. 2017, Seppänen et al.

2007). While the nature of interaction net effect that emerge as the result of several interacting

processes between species cannot be defined in communities at large scale (Barner et al. 2016),

estimating the interaction milieu becomes more accessible through increasing availability of

monitoring data with large spatial and temporal resolutions (Ovaskainen et al. 2016), providing that

species' associations are correctly described.

Species associations have been estimated using spatial co-occurrences of species (HilleRisLambers et al. 2012, Letten et al. 2017), disentangling the effect of the biotic filter (*i.e.* interactions) from other filters (*e.g.* habitat filtering) on co-occurrence patterns (Morueta-Holme et al. 2016, Elo et al. 2021) or in combination with interactions described in the literature (Lurgi et al. 2012). Species associations have also been obtained from species time-series using dynamical system analyses (Ushio et al. 2018, Barraquand et al. 2021). Using such associations as meaningful proxies of ecological interactions has been criticised, due to numerous unchecked non-biotic processes (*e.g.* dispersal capacity, habitat, phylogeography, Moran effet) (Sander et al. 2017, Freilich et al. 2018, Dallas et al. 2019, Blanchet et al. 2020, Cordero and Jackson 2021). However, spatial co-occurrences correspond to an apparent equilibrium resulting from large-scale processes (i.e. biotic and abiotic filters) (Rigal et al. 2022), while fluctuations in species time-series reflect an non-equilibrium inside community composition (Ushio et al. 2018). Estimating the interaction milieu in bird communities can therefore benefit from combining spatial and temporal information on bird species abundance.

Pairwise species associations per se are not the primary focus of a community analysis based on the interaction milieu (McGill et al. 2006) but can be used to quantify, via network metrics, the structure of the interaction milieu (Kaiser-Bunbury and Blüthgen 2015, Lau et al. 2017). A network with species as nodes and association as edges between nodes, can be analysed at network, intermediate (i.e. by sub-groups) or node levels (Lau et al. 2017). For instance, at network level, the structure can be defined by the number of nodes, number of edges and connectance (i.e. density of edges). Network structure can be analysed at intermediate level, e.g. by motifs (i.e. recurrent patterns), average path length (i.e. average number of steps between all possible pairs of nodes) and modularity (i.e. extent of network division into sub-groups). At node or species level, the network structure can be studied using, for

instance, the degree (i.e. the number of edges by nodes), the centrality (i.e. how influential is a node) or

the partnership diversity (i.e. preference for nodes in a network to attach to other similar nodes)

(Delmas et al. 2019). The interpretation of each metric on its own is neither relevant nor generalisable,

since a change in the index may be due to changes in the size of the network (Heleno et al. 2012), be

correlated with the other indicators (Delmas et al. 2019) and depend on the type of interaction studied

(Tylianakis et al. 2010).

In this study, we seek to understand the changes in the interaction milieu that accompany the changes in

species composition over the last two decades in France. In addition to the overall changes in the

interaction milieu of common bird communities, we expect that there are also variations in the changes

of interaction milieu across habitats, as bird communities are not exposed to the same pressures and

changes in species composition in the different habitats. To do so, we used the French Breeding Bird

Survey from 2001 to 2017 to estimate potential associated species pairs using spatial co-occurrences to

account for large-scale community composition filters (Fig. 1a), and we refined the selection of

associated species pairs using species time-series (Fig. 1b). We then assessed the relevance of the

associated species pairs to infer the biotic background of interaction using species ecological traits (Fig.

1c). Lastly, we tracked the temporal changes in the interaction milieu using network metrics (Fig. 1d)

and analyse the consequences, for the conservation of bird communities, of changes in the interaction

milieu that accompany changes in species composition.

Figure 1: Framework of the estimation of changes in the interaction milieu of bird communities from
large-scale spatio-temporal data. a) In a first step, the effects of large-scale processes on species co-
occurrences are removed using community apparent equilibrium, to estimate the potentially associated
species pairs. b) In a second step, the estimation of associated species pairs is refined by removing the
effects of non-biotic local processes on the non-equilibrium of community composition. c) The
ecological relevance of the estimated associations is tested using the ecological hypothesis on
community assembly and species traits. d) Community composition in space and time are combined
with associated species pairs to compute network metrics quantifying the interaction milieu and track
its changes.

2. Material and methods

2.1 Spatial data and time-series of birds

We used the French Breeding Bird Survey (FBBS) based on a standardised protocol to monitor the
abundance of common bird species. 2,514 sites of 2x2 km were survey in mainland France between
2001 and 2017 (Jiguet et al. 2012). In each site, the abundances of breeding bird species were counted

on 10 sampling points (during 5 min, 1 to 4 h after sunrise) by volunteer experienced ornithologists,

twice a year between 1^{st} of April and 15^{th} of June from 2001 to 2017. Each site represented a species

assemblage changing in abundance and composition over time. Habitats were described for each

sampling point in 19 types (supplementary material 1). The data provided by the French National

History Museum (MNHN) were abundances corrected for the difference in species detectability.

Among the 242 recorded species, we selected the most abundant species representing 99% of the total

abundance in the dataset to avoid over-representation of rare species for which the FBBS is not well

designed. After removing rare species and sites only monitored once, our dataset is composed of 1,969

sites and 109 species.

2.2 Potentially associated species pairs

We used spatial co-occurrence to estimate the species pairs potentially associated (Fig. 1a) by removing

large-scale non-interaction filters (historical dispersal, neutral dispersal and habitat filtering) following

Rigal et al. (2022). We measured the partial correlations between species pairs using their co-

occurrence profile. We accounted for historical and habitat filtering by estimating potentially associated

pairs for each type of habitat and each biogeographic region (see additional details in supplementary

material 1). We limited the influence of random processes and uncontrolled habitat filtering using a null

model by resampling species in a site. This provided 1000 random co-occurrence datasets obtained by

keeping constant the total number of individuals in a given sampling point, and assuming that the

probability for a species to occur in a given sampling point was proportional to its frequency in the

dataset. We calculated the standardised effect size (SES) of the observed partial correlations compared

to the partial correlations from the random datasets. Significant associated pairs were based on p-values

(<0.05) obtained from the rank of the observed partial correlations in the distribution of random partial

correlations (Morueta-Holme et al. 2016).

2.3 Species associations

For each potentially associated species pair in a given habitat and a given biogeographic region, we

determined which species pairs were significantly associated (Fig. 1b) based on the relationship

between species time-series. We used the multispatial Convergent Cross-Mapping (CCM) method (R

package *multispatialCCM*) (Clark et al. 2015), an extension of the CCM method (Sugihara et al. 2012).

In CCM, the time-series of two species are linked if we can predict the dynamics of species X using the

time-series of species Y, *i.e.* if the attractor of X (defined as the set of states reconstructed from the

original and lagged time-series of X) can be estimated from the attractor of Y. If so, species X has a

signature into the dynamics of species Y, meaning that species X is affecting species Y (see detailed in

supplementary material 2 and Tsonis et al. 2018). CCM resulted in a correlation value ρ, depicting the

correlation between the two attractors, associated to a p-value for significance.

We focused on sites monitored during at least six consecutive years to maximise the number of sites

and with long enough time-series in order to assess significant temporal associations between species.

For each species, we reconstructed its time-series at a given site using its recorded abundance, set to

zero when the species was not recorded during a monitoring session of the site. Obtaining pseudo-time-

series (*i.e.* the times-series reconstructed by placing side by side data from multiple sites) longer than

500 time steps (following the empirical example from Clark et al. 2015) requires to select the 32,491

combinations of species pairs, habitats and biogeographic regions that match this criteria among the

89,270 potentially associated species pairs in given habitat and biogeographic region obtained from

spatial co-occurrences. We use 300 replicates for the bootstrap routine due to computational time. The

multispatial CCM provides species pairs for which the time-series of one species was related to the

time-series of the other. As we aimed at estimating the interaction milieu rather than analysing pairwise

associations, we did not account for the direction of the relationship, i.e. if one or both directions was

significant, the pair of species was considered to be associated.

2.4 Testing ecological relevance of species associations

We investigated whether the associated species pairs were relevant to assess the biotic background of

interaction (Fig. 1c) based on trait-based assembly theory and specific expected interactions. According

to trait-based assembly theory, species that share similar functional traits, are phylogenetically clustered

or with significant habitat overlap, are expected to interact more strongly (Fontaine and Thébault 2015,

Mönkkönen et al. 2017). This stronger interaction may be either positive through social information

exchange and facilitation processes (Gil et al. 2017), or negative because of resource competition under

the limiting similarity hypothesis (Abrams 1983), which should result in a funnel pattern when species

interactions are combined with trait-related distances (Mönkkönen et al. 2017). We tested whether the

associations obtained from previous steps are in line with these expectations using two approaches. In a

first test based on distances between species characteristics, we computed functional, phylogenetic and

niche overlap distances (see relationship and complementarity between distances in supplementary

material 3) and relate them to the probability of a species pair to be associated. In a second test that is

specifically focused on expected interactions, we analysed whether species pairs with a similar diet or

nest requirement are more prone to be associated, while controlling for their preferred habitat.

2.4.1 Functional distance

Functional distance was estimated using a life-history trait dataset for the birds of the Palaearctic

(Storchová and Hořák 2018). We computed the Generalized Functional Diversity (GFD) following

(Mouchet et al. 2008) to generate a pairwise functional distance between all pairs of species (Mouchet

et al. 2010). This method calculated all the combinations of the clustering algorithms from a functional

distance matrix and provides the best consensus tree, *i.e.* the one for which the dissimilarity with the

original distance matrix is the lowest according to the 2-norm goodness-of-fit index (Mérigot et al.

2010). We constructed the initial functional distance matrix using the Gower's distance as the trait

matrix contained both qualitative and quantitative data (Gower 1971). The best consensus dendrogram

was built using the Unweighted Pair Group Method using arithmetic Average algorithm (UPGMA) (2-

norm=0.03, threshold=3.49). From this dendrogram, we computed the cophenetic functional distance

(Sokal and Rohlf 1962) between species, using the 242 initial species to assess the functional distances

in the complete functional space of the common bird species.

2.4.2 Phylogenetic distance

Phylogenetic trees of the 242 bird species were obtained from **birdtree.org** (Jetz et al. 2014). We

estimated the consensus tree from 10,000 trees using the *ape* R package (Paradis et al. 2019). We

calculated the phylogenetic distance as the cophenetic distance between each pair of species in the

consensus tree.

2.4.3 Niche overlap distance

We computed niche overlap for each species pair as the overlap between the frequency distribution of

each species within the 19 classes of habitat. To do so, we used the habitat information for each

sampling point and we calculated the frequency distribution of each species over the 19 classes of

habitats in the whole dataset. We then computed the χ^2-distance, widely used to compare ecological

assemblages (Ben-Shahar and Skinner 1988, Flower et al. 1997), to estimate the distance between each

species habitat profile. The χ^2-distance characterises the proximity in habitat distribution profiles

independently of species abundance. We defined the habitat niche overlap distance (*NOD*) using the

similarity between habitat profiles as follows (Eq. 1 and see details on NOD in supplementary material

3):

$\quad NOD_{a,b} = 1 - \chi^2_{dist\,a,b} = \sqrt{\sum_{j=1}^{H} \frac{y_t}{y_h} \left(\frac{y_{ah}}{y_a} - \frac{y_{hb}}{y_b} \right)}$ (Eq. 1)

with $NOD_{a,b}$ the niche overlap distance between species *a* and *b*, $\chi^2_{dist\,a,b}$ the χ^2-distance between habitat

profiles of species *a* and *b*, *H* the number of habitats, y_t the total abundance of species *a* and *b* in all the

habitats, y_h the abundance of species *a* and *b* in habitat *h*, y_{ah} the abundance of species *a* in *h*, y_a the total

abundance of species *a*, y_{hb} the abundance of species *b* in *h* and y_b the total abundance of species *b*.

2.4.4 Test with functional, phylogenetic and niche overlap distances

To test the ecological relevance of the associated species pair, we performed a binary logistic regression

using Generalised Linear Models (GLM). The response variable (*A*) was the presence or absence of

association for each species pair (a species pair is considered to be associated if this has been the case

in at least one habitat and one biogeographic region), with a binomial distribution with parameter *p* and

a logit link function *g*. The explanatory variables were scaled functional distance (*FD*), phylogenetic

distance (*PD*) and niche overlap distance (*NOD*) (FD and PD being slightly correlated but not with

NOD, see supplementary material 3) for species pair *i* (Eq. 2):

262
$$A_i \sim Binomial(p)$$
$$g(p) = \beta_0 + \beta_1 FD_i + \beta_2 PD_i + \beta_3 NOD_i$$
$$g(x) = \log\left(\frac{x}{1-x}\right)$$
(Eq. 2)

2.4.5 Test with expected interactions for food and nest location

Using the life-history trait dataset for the birds of the Palaearctic (Storchová and Hořák 2018), we

recorded for each species pair whether the two species used the same nest type among the five recorded

types (*closed arboreal; on ground directly; in tussock very close to ground but not directly on ground;*

hole in tree, bank, ground, crevice; open arboreal cup in bush, tree, on cliff ledge) and had the same

diet pattern during breeding season based on nine diet descriptors (*at least 10% of diet composed of*

small plants; fruits; grains and seeds; arthropods; other invertebrates; fish; other vertebrates;

carrions; diet composed of similar amount of plants and animals). To assess the effect of these share

traits while controlling for the habitat requirement of the species pair, we recorded whether the two

species of the pair prefer the same habitat among the 19 classes, according to their occurrence

frequency in each habitat class.

A second GLM was performed in which the binomial response variable (*A*) was the presence or

absence of association for each species pair and the explanatory variables were the presence or absence

of a share diet (*Diet*), the presence or absence of a shared nest type (*Nest*) and the presence or absence

of a similar preferred habitat (*Habitat*) for species pair *i* (Eq. 3).

$g(p) = \beta_0 + \beta_1 Diet_i + \beta_2 Nest_i + \beta_3 Habitat_i$ (Eq. 3)

2.5 Analysis of the interaction milieu

To quantify the changes in the interaction milieu, we focus on the number of species, the number of

associations (Heleno et al. 2012, Kaiser-Bunbury and Blüthgen 2015) and a set of relevant metrics at

network, intermediate and node levels. At the network level, we use connectance, a standard metric

whose mathematical relationship with other network metrics is well known and which is therefore

important for analysing other metrics. The relationship between connectance and perturbations is not

consistent, but independently of the type of interactions, connectance increase has been linked to

greater species generalism (Tylianakis et al. 2010). At the intermediate level, we use modularity which

has been shown as a suitable metric for conservation issues: high modularity restrains perturbations and

local extinction within a module (Stouffer and Bascompte 2011). At the node levels, we use evenness

of degree distribution that has been related to higher robustness as it decreases the consequences of a

local extinction for the remaining species and tend to be lower in the context of degraded habitats

(Kaiser-Bunbury and Blüthgen 2015, Fisogni et al. 2021).

2.5.1 Network metrics

We reconstructed a local network (Fig. 1d) at each site and for each year using the species composition

(species as nodes) and the associated species pairs (associations as edges). We calculated the following

metrics from each network j: network size s (i.e. number of species), number of associations a,

connectance c (i.e. density of associations in the network), modularity m (i.e. intensity of network

division into sub-groups) and evenness of degree distribution e (i.e. evenness of probability distribution

of the number of associations by species). Connectance is the ratio between the observed number of

associations a and the potential number of associations a_{max} (Landi et al. 2018) (directly linked to the

number of species pairs $\binom{s}{2}$, Eq. 4):

$$c_j = \frac{a_j}{a_{max\,j}} = \frac{a_j}{\binom{s_j}{2}}$$ (Eq. 4)

Modularity (Eq. 5) was computed using the R package *rnetcarto* (Doulcier and Stouffer 2015) which

enables fast modularity computation (required for the analysis of the amount of networks in this study).

It is based on the *rgraph* library and the simulated Annealing method (Guimerà and Nunes Amaral

2005).

$$m_j = \frac{1}{2a_j} \sum_{x,y} \left(A_{x,y} - \frac{k_x k_y}{2a} \right) \delta(t_x, t_y)$$ (Eq. 5)

with a_j the number of associations in network j, $A_{x,y}$ the element of the adjacency matrix, k the degree

and t the type to which species x or y belongs and the Kroenecker delta function defined as equal to 1

when $t_x = t_y$ and 0 when $t_x \neq t_y$.

The evenness of the degree distribution was calculated using the Shannon entropy of the degree

distribution (Blüthgen et al. 2008) (Eq. 6):

$$e_j = \frac{-\sum_{k=1}^{s_j} p_{kj} \ln p_{kj}}{\ln s_j}$$ (Eq. 6)

with p_k the frequency of degree k.

To explore potential differences in changes of the interaction milieu across habitats, we also conducted

the computation of network metrics for the four main habitat classes: farmland, forest, urban and

natural open land. We grouped the 19 habitats provided at each site and each year by the volunteers

conducting the monitoring into the four habitat classes (see supplementary material 1) and we

attributed to each local community its main habitat class to separate between farmland, forest, urban

and natural open land communities.

2.5.2 Changes in the interaction milieu through time

We calculated the yearly change in the interaction milieu by estimating the effect of years Y (set as an

explanatory variable) on each network metric, while explicitly modelling spatial auto-correlation

between sites using a Generalised Additive Mixed Model (GAMM) (Eq. 7) with a tensor product of a

thin plate regression spline based on geographic coordinates of sites (longitude *lo* and latitude *la*)

(Wood 2003, 2017). As the number of associated pairs and connectance are directly affected by the

number of species (see correlation between network metrics in supplementary material 4) (Blüthgen et

al. 2006, Besson et al. 2019), we added the number of species as a control variable for these two

metrics, to estimate if the observed change in the number of associated pairs was higher or lower than

expected. Similarly, connectance constrains the possible values of modularity and degree distribution

(Poisot and Gravel 2014, Delmas et al. 2019), and thus connectance was added as a control variable for

estimating changes in modularity and evenness of degree distribution. As the number of monitored sites

changes between years, the site identity z was added as a random variable.

$\quad NM_j \sim \alpha_0 + \alpha_1 Y_j + \alpha_2 CV_j + f(lo_j, la_j) + \epsilon_j$ (Eq. 7)

with NM the network metric (number of species s, number of associations a, connectance c, modularity

m and evenness e) and CV the control variable ($\varnothing$ for s, s for a and c, c for m and e) and $\varepsilon_j \sim N(0,\sigma)$.

3. Results

3.1 Species associations

Among the 292,796 combinations between 5,886 pairs of bird species, from four biogeographic regions

and 19 habitats, we found 89,270 potential species associations (Fig. 1a). Among them, 32,491

combinations enable to reconstruct time-series with more than 500 time steps and we found 585

combinations with significantly associated species pairs (Fig. 1b). Among the 5,886 species pairs, this

represents 396 (7%) associated species pairs (see association matrix in supplementary material 5), some

species pairs being significantly associated in several habitats and biogeographic regions.

We found negative relationships between the propensity of a species pair to be associated and

functional (-0.22 ± 0.10, Fig. 2a) and niche overlap distances (-1.73 ± 0.17, Fig. 2a). In other words,

species that are more functionally similar and with higher niche overlap are more likely associated.

However, we found only weak evidence of a relationships between the propensity of a species pair to

be associated and phylogenetic distance (-0.09 ± 0.10, Fig. 2a and model validation in supplementary

material 4) as part of the variation might be already explained by functional distance (slightly

correlated with phylogenetic distance, see supplementary material 3). Concerning specific expected

interactions, we found a positive relationship between the propensity of a species pair to be associated

and diet during breeding season (0.23 ± 0.20, Fig. 2b), a positive relationship with similar preferred

habitat (1.05 ± 0.19, Fig. 2b) and no evidence of a relationship with similar nest type (0.10 ± 0.18, Fig.

2b).

Figure 2: Odd ratios of GLM between propensity of a species pair to be associated and a) in black, the

ecological distances (phylogenetic, niche overlap and functional distances), b) in white, specific

expected interactions (similar nest, similar preferred habitat and similar diet in breeding season). 95%

CI are displayed with grey bars. McFadden R² corresponds to the pseudo-R² for logistic regression

models (McFadden 1974).

3.2 Temporal change in the interaction milieu

Breeding bird communities tend to decrease in size ($s.year^{-1}$ = -0.15 ± 0.02, Fig. 3a, see annual

variations in supplementary material 4), increase in connectance ($c.year^{-1}$ = $5.6.10^{-4}$ ± $2.3.10^{-4}$, Fig. 3c)

and decrease in evenness of the degree distribution ($e.year^{-1}$ = $-1.1.10^{-4}$ ± $0.5.10^{-4}$, Fig. 3d) over the

studied period (2001-2017). This is verified for farmland and forest communities that represent the

majority of communities (55.3% in farmland, 29.7% in forest, 7.9% in natural open land and 7.1% in

urban areas). For farmland and forest communities, the increase in connectance is related to the decline

in the number of species, whereas the number of associations relatively increases compared to the

number of species in natural open land communities. For the interaction milieu in urban communities,

the only significant change is the decline of the number of species (Fig. 3a), the other metrics staying

constant.

396

Figure 3: Temporal changes in the interaction milieu of bird communities between 2001 and 2017. a)

Temporal change in the number of species s, b) temporal change in the number of associations a (while

controlling for the number of species), c) temporal change in connectance c (while controlling for the

number of species), d) temporal change in modularity m (while controlling for connectance), e)

temporal change in evenness of the degree distribution e (while controlling for connectance). Black dot

is for all communities, colours for communities from each habitat: yellow for farmland, green for

forest, red for urban, blue for natural open land. 95% CIs are displayed and significant changes are

shown by $*$.

Consequently, the interaction milieu from the start of the period (Fig. 4a) is modified by a change in

both the number of species (Fig. 4b) and the topology of relationships between species (Fig. 4c). At the

level of the community, the density of associated pairs (i.e. connectance, Fig. 3c) is higher, which is

mainly due to the reduction in the number of species (Fig. 3a), not to a relative increase of the number

of associated pairs (Fig. 3b). The intensity of division of the community into sub-groups (i.e.

modularity) remains stable when accounting for the density of associations. Thus, the major change is

at the species level, where the probability distribution of the number of association by species (i.e.

evenness) tends to become more skewed (i.e. shifting from species with a similar number of association

to species with few or multiple associations). The trend over the period studied is from large

communities with evenly distributed associations (Fig. 4a) to smaller communities with a similar

relative number of associations but unevenly distributed (Fig. 4d), which an focus on species only

would not have revealed.

Figure 4: Observed examples of bird communities illustrating the topological changes in the
interaction milieu between 2001 and 2017 with the values for each network metric. From a) a
community with a given number of species s and network metrics (number of associations a,
connectance c, modularity m and evenness e), b) network metrics may not be affected by a change in
the number of species, c) conversely, a stable number of species may hide a change in network
structure. d) On average, the observed changes between 2001 and 2017 in bird communities show an
interaction milieu with a lower number of species and a similar relative number of associations that are
becoming less evenly distributed, as illustrated by the observed network displayed.

4. Discussion

4.1 Ecological relevance of associated species pairs

The species pairs we have identified as associated share similar functional characteristics or habitat requirements. The significant relationships with functional and niche overlap distances were in line with our hypothesis based on trait-based assembly theory and empirical results on birds (Mönkkönen et al. 2017) and suggest that our approach produced ecologically meaningful associated species pairs. The absence of a significant relationship between associated species pairs and phylogenetic distance can be linked to the phylogenetic diversity already encompassed in functional distance (see supplementary material 3). Despite this result and the strong filters applied in the estimation of associated pairs, some of the associated pairs can still be due to a common response to an environmental variable in space and time (Pollock et al. 2014) and thus an absence of association does not imply an absence of interaction between two species (Cazelles et al. 2016). Yet, we provide an in-depth validation of the overall set of associated pairs using the ecological distances and confirm this by looking at expected interacted pairs of species based on specific interactions during the breeding season (corresponding to the period of the survey) for food and nest location. The similarity in nest location is not significant, which may be due to the fact that competition would be the main expected interaction from this shared characteristic (Martin et al. 2004). On the contrary, a similar diet, that significantly increases the propensity to be associated, could induce both repulsion and attraction (Seppänen et al. 2007, Gil et al. 2017) and thus is expected to induce interactions in a wider range of species pairs and to be more easily detected. This result supports the findings based on functional and niche overlap distances (and see supplementary material 4 for additional detail on specific associations).

4.2 Specificity of the interaction milieu

The changes in the interaction milieu of bird communities estimated from the associated species pairs

show a decrease in the number of species and the evenness of the number of associations by species as

well as an increase in the density of association, with urban and natural open land communities that

differ slightly from this general pattern. These changes in the biotic background of interaction in bird

communities remain estimated from species links that are coarse and static, whereas ecological

interactions are known to be changing though time (Ramos-Robles et al. 2016, Suzuki et al. 2023), due

in particular to changes in environmental conditions (Lurgi et al. 2012, Poisot et al. 2015). In addition,

the interpretation of changes in the interaction milieu is limited by the absence of formal identification

of the interaction nature, compared to explicit ecological networks (e.g. mutualistic plant-pollinators,

food webs). Our finding can therefore only be compared by analogy to changes observed in ecological

networks with specific interactions such as plant-pollinator networks and food webs. In particular, the

changes observed in just two decades in the interaction milieu of bird communities can be considered

as marked compared to analogous communities in which frugivore-plant, pollination networks and

food webs are reported to be stable (Petanidou et al. 2008, Kaartinen and Roslin 2012, Plein et al.

2013). Plant-pollinator networks, food webs and research on bird flocks (closely related to intra-guild

networks) may however bring clues to interpret the observed changes in the interaction milieu of bird

communities.

4.3 Changes in the interaction milieu and consequences for conservation

Based on an analogy between these ecological networks and bird communities (Lane et al. 2014,

Mokross et al. 2014), we can assume in particular that specialists species are supposed to be involved

in fewer but stronger interactions (Rigal et al. 2022). If changes in connectance alone are not

informative for conservation (Heleno et al. 2012), the increase in connectance is related here to a

reduction in the number of species in bird communities that could be interpreted in line with the

observed decline of habitat specialist species (Devictor et al. 2008b). Higher connectance can improve

network persistence facing perturbation (Besson et al. 2019), which might be the case if all the fragile

species and interactions are already wiped out while generalists involved in more associations

relatively expand (Godet et al. 2015). However, if this connectance increase is at the cost of species

removal, according to the biodiversity-ecosystem functioning theory (which is relevant inside a guild),

this will result in a declining efficiency of resource use (Loreau 2010). Based on our results, such a

decline in community functioning is therefore expected for forest and farmland communities, in which

the change in connectance is explained by the decline in species number. Conversely, the structure and

connectance of communities in natural open land are principally modified by the replacement of less

associated species by more associated ones, expected to be generalists.

An increase in connectance also induces a decline in modularity (Delmas et al. 2019) which has been

the case in bird communities, although modularity change is not stronger or weaker than expected (i.e.

the modularity does not change through time when we control for changes in connectance Fig. 3d). A

modular structure is known to reduce the impact and propagation of a perturbation throughout the

network (Stouffer and Bascompte 2010). Therefore, considering consequences for the interaction

milieu, a decline in modularity might affect community ability to cope with environmental changes and

induce a higher sensitivity to cascade and secondary extinctions.

The observed decline in evenness of association degree distribution can be more directly interpreted.

Low evenness in the distribution of interaction strength can be often observed in ecological networks,

due to the numerous weak interactions (Bascompte et al. 2005) known to stabilise community

dynamics. However, when focusing on the presence, not the strength, of a link, the evenness of the

distribution is higher in preserved or restored communities than in degraded habitats (Tylianakis et al.

2010, Fisogni et al. 2021), not only in food webs but also in avian flocks (Mokross et al. 2014). It is

therefore likely that the observed decline in evenness, which is higher than expected by the observed

change in connectance, is the footprint of habitat degradation experienced by bird communities. The

observed decline in evenness in forest and farmland communities may therefore be compared with the

well-studied decline in species dynamics and changes in community composition over the past decades

in these environments (Brotons and Jiguet 2010, Gaüzère et al. 2020, Rigal et al. 2023).

Although our study focuses on the interaction milieu of bird communities, this should not understate

the importance of the marked decline observed in the size of these communities. Beyond the species

loss that is an issue per se, a decline in species diversity and abundance is expected to affect interaction

frequency as the encounter rates become lower. This could lead to the loss of some interactions while

the two species are still present in the vicinity (Staniczenko et al. 2017, Tylianakis and Morris 2017).

Yet, species remains a central unit for conservation purposes and our results support the fact that

change in ecological network structure should be analysed in regard with change in species number and

identity (Heleno et al. 2012).

Changes in the interaction milieu of bird communities highlighted in this study are crucial to

understand how communities will be able to respond to future perturbations, in particular if community

functioning is affected by changes in the network layout (Besson et al. 2019). Community responses

are affected by the loss of species (Loreau and De Mazancourt 2013) which is, additionally, deepened

by the rewiring (Tylianakis and Morris 2017) or the vanishing of associations (Gilljam et al. 2015).

These rapid modifications of bird biodiversity, that would have been hidden by a species-centred

approach, are expected to affect the stability of communities and their capacity to cope with rapid and

multiple global changes.

Data availability: All the analyses are conducted using the R software v.4.2.2 (R Core Team 2018).

Code and data are available on Zenodo https://zenodo.org/doi/10.5281/zenodo.12784005 (Rigal et al.

2024).

References

Abrams, P. 1983. The theory of limiting similarity. - Annu. Rev. Ecol. Syst. 14: 359–376.

Barner, A. K., Hacker, S. D., Menge, B. A. and Nielsen, K. J. 2016. The complex net effect of reciprocal interactions and recruitment facilitation maintains an intertidal kelp community. - J. Ecol. 104: 33–43.

Barraquand, F., Picoche, C., Detto, M. and Hartig, F. 2021. Inferring species interactions using Granger causality and convergent cross mapping. - Theor. Ecol. 14: 87–105.

Bas, Y., Devictor, V., Moussus, J.-P. and Jiguet, F. 2008. Accounting for weather and time-of-day parameters when analysing count data from monitoring programs. - Biodivers. Conserv. 17: 3403–3416.

Bascompte, J., Melián, C. J. and Sala, E. 2005. Interaction strength combinations and the overfishing of a marine food web. - Proc. Natl. Acad. Sci. 102: 5443–5447.

Ben-Shahar, R. and Skinner, J. D. 1988. Habitat Preferences of African Ungulates Derived by Uni- and Multivariate Analyses. - Ecology 69: 1479–1485.

Besson, M., Delmas, E., Poisot, T. and Gravel, D. 2019. Complex ecological networks. - Encycl. Ecol. 1: 536–545.

Blanchet, F. G., Cazelles, K. and Gravel, D. 2020. Co-occurrence is not evidence of ecological interactions. - Ecol. Lett. 23: 1050–1063.

Blüthgen, N., Menzel, F. and Blüthgen, N. 2006. Measuring specialization in species interaction networks. - BMC Ecol. 6: 9.

Blüthgen, N., Fründ, J., Vázquez, D. P. and Menzel, F. 2008. What do interaction network metrics tell us about specialization and biological traits. - Ecology 89: 3387–3399.

Brotons, L. and Jiguet, F. 2010. Bird communities and climate change. - Eff. Clim. Change Birds: 275–294.

Cazelles, K., Araújo, M. B., Mouquet, N. and Gravel, D. 2016. A theory for species co-occurrence in interaction networks. - Theor. Ecol. 9: 39–48.

Clark, A. T., Ye, H., Isbell, F., Deyle, E. R., Cowles, J., Tilman, G. D. and Sugihara, G. 2015. Spatial convergent cross mapping to detect causal relationships from short time series. - Ecology 96: 1174–1181.

Clavel, J., Julliard, R. and Devictor, V. 2011. Worldwide decline of specialist species: toward a global functional homogenization? - Front. Ecol. Environ. 9: 222–228.

Cordero, R. D. and Jackson, D. A. 2021. Abiotic factors influence species co-occurrence patterns of lake fishes. - J. Anim. Ecol. 90: 2859–2874.

Dallas, T., Melbourne, B. A. and Hastings, A. 2019. When can competition and dispersal lead to checkerboard distributions? - J. Anim. Ecol. 88: 269–276.

Davey, C. M., Chamberlain, D. E., Newson, S. E., Noble, D. G. and Johnston, A. 2012. Rise of the generalists: evidence for climate driven homogenization in avian communities. - Glob. Ecol. Biogeogr. 21: 568–578.

Davison, J., Gerz, M., Hiiesalu, I., Moora, M., Semchenko, M. and Zobel, M. 2024. Niche types and community assembly. - Ecol. Lett. 27: e14327.

Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Dalla Riva, G. V., Fortin, M.-J., Gravel, D., Guimarães Jr, P. R., Hembry, D. H. and Newman, E. A. 2019. Analysing ecological networks of species interactions. - Biol. Rev. 94: 16–36.

Devictor, V., Julliard, R., Clavel, J., Jiguet, F., Lee, A. and Couvet, D. 2008a. Functional biotic homogenization of bird communities in disturbed landscapes. - Glob. Ecol. Biogeogr. 17: 252–261.

Devictor, V., Julliard, R. and Jiguet, F. 2008b. Distribution of specialist and generalist species along spatial gradients of habitat disturbance and fragmentation. - Oikos 117: 507–514.

Doulcier, G. and Stouffer, D. 2015. Rnetcarto: Fast network modularity and roles computation by simulated annealing. - R Package Version 02 in press.

Elo, M., Jyrkänkallio-Mikkola, J., Ovaskainen, O., Soininen, J., Tolonen, K. T. and Heino, J. 2021. Does trait-based joint species distribution modelling reveal the signature of competition in stream macroinvertebrate communities? - J. Anim. Ecol. 90: 1276–1287.

Fisogni, A., Massol, F., De Manincor, N., Quaranta, M., Bogo, G., Bortolotti, L. and Galloni, M. 2021. Network analysis highlights increased generalisation and evenness of plant-pollinator interactions after conservation measures. - Acta Oecologica 110: 103689.

Flower, R. J., Juggins, S. and Battarbee, R. W. 1997. Matching diatom assemblages in lake sediment cores and modern surface sediment samples: the implications for lake conservation and restoration with special reference to acidified systems. - Hydrobiologia 344: 27–40.

Fontaine, C. and Thébault, E. 2015. Comparing the conservatism of ecological interactions in plant–pollinator and plant–herbivore networks. - Popul. Ecol. 57: 29–36.

Freilich, M. A., Wieters, E., Broitman, B. R., Marquet, P. A. and Navarrete, S. A. 2018. Species co-occurrence networks: Can they reveal trophic and non-trophic interactions in ecological communities? - Ecology 99: 690–699.

Gaüzère, P., Jiguet, F. and Devictor, V. 2015. Rapid adjustment of bird community compositions to local climatic variations and its functional consequences. - Glob. Change Biol. 21: 3367–3378.

Gaüzère, P., Barbaro, L., Calatayud, F., Princé, K., Devictor, V., Raison, L., Sirami, C. and Balent, G. 2020. Long-term effects of combined land-use and climate changes on local bird communities in mosaic agricultural landscapes. - Agric. Ecosyst. Environ. 289: 106722.

Gil, M. A., Emberts, Z., Jones, H. and St. Mary, C. M. 2017. Social information on fear and food drives animal grouping and fitness. - Am. Nat. 189: 227–241.

Gilljam, D., Curtsdotter, A. and Ebenman, B. 2015. Adaptive rewiring aggravates the effects of species loss in ecosystems. - Nat. Commun. 6: 1–10.

Godet, L., Gaüzere, P., Jiguet, F. and Devictor, V. 2015. Dissociating several forms of commonness in birds sheds new light on biotic homogenization. - Glob. Ecol. Biogeogr. 24: 416–426.

Gómez-Catasús, J., Benítez-López, A., Díaz, M., González del Portillo, D., Bravo, C., et al. 2025. Alarming conservation status of Western European steppe birds and their habitats: An expert-based review of current threats, traits and knowledge gaps. - Biol. Conserv. 311: 111414.

Goodale, E., Ding, P., Liu, X., Martínez, A., Si, X., Walters, M. and Robinson, S. K. 2015. The structure of mixed-species bird flocks, and their response to anthropogenic disturbance, with special reference to East Asia. - Avian Res. 6: 14.

Gower, J. C. 1971. A general coefficient of similarity and some of its properties. - Biometrics: 857–871.

Guimerà, R. and Nunes Amaral, L. A. 2005. Functional cartography of complex metabolic networks. - Nature 433: 895–900.

Heleno, R., Devoto, M. and Pocock, M. 2012. Connectance of species interaction networks and conservation value: Is it any good to be well connected? - Ecol. Indic. 14: 7–10.

HilleRisLambers, J., Adler, P. B., Harpole, W. S., Levine, J. M. and Mayfield, M. M. 2012. Rethinking community assembly through the lens of coexistence theory. - Annu. Rev. Ecol. Evol. Syst. 43: 227–248.

Jaillard, B., Richon, C., Deleporte, P., Loreau, M. and Violle, C. 2018. An a posteriori species clustering for quantifying the effects of species interactions on ecosystem functioning. - Methods Ecol. Evol. 9: 704–715.

Jetz, W., Thomas, G. H., Joy, J. B., Redding, D. W., Hartmann, K. and Mooers, A. O. 2014. Global Distribution and Conservation of Evolutionary Distinctness in Birds. - Curr. Biol. 24: 919–930.

Jiguet, F., Devictor, V., Julliard, R. and Couvet, D. 2012. French citizens monitoring ordinary birds provide tools for conservation and ecological sciences. - Acta Oecologica 44: 58–66.

Kaartinen, R. and Roslin, T. 2012. High temporal consistency in quantitative food web structure in the face of extreme species turnover. - Oikos 121: 1771–1782.

Kaiser-Bunbury, C. N. and Blüthgen, N. 2015. Integrating network ecology with applied conservation: a synthesis and guide to implementation. - AoB PLANTS 7: plv076.

Kampichler, C., Van Turnhout, C. A., Devictor, V. and Van Der Jeugd, H. P. 2012. Large-scale changes in community composition: determining land use and climate change signals. - PloS One 7: e35272.

Koerner, S. E., Poulsen, J. R., Blanchard, E. J., Okouyi, J. and Clark, C. J. 2017. Vertebrate community composition and diversity declines along a defaunation gradient radiating from rural villages in Gabon. - J. Appl. Ecol. 54: 805–814.

Landi, P., Minoarivelo, H. O., Brännström, Å., Hui, C. and Dieckmann, U. 2018. Complexity and stability of adaptive ecological networks: a survey of the theory in community ecology. - In: Systems analysis approach for complex global challenges. Springer, pp. 209–248.

Lane, P. W., Lindenmayer, D. B., Barton, P. S., Blanchard, W. and Westgate, M. J. 2014. Visualization of species pairwise associations: a case study of surrogacy in bird assemblages. - Ecol. Evol. 4: 3279–3289.

Lau, M. K., Borrett, S. R., Baiser, B., Gotelli, N. J. and Ellison, A. M. 2017. Ecological network metrics: opportunities for synthesis. - Ecosphere 8: e01900.

Letten, A. D., Ke, P.-J. and Fukami, T. 2017. Linking modern coexistence theory and contemporary niche theory. - Ecol. Monogr. 87: 161–177.

Li, D., Poisot, T., Waller, D. M. and Baiser, B. 2018. Homogenization of species composition and species association networks are decoupled. - Glob. Ecol. Biogeogr. 27: 1481–1491.

Lindström, Å., Green, M., Paulson, G., Smith, H. G. and Devictor, V. 2013. Rapid changes in bird community composition at multiple temporal and spatial scales in response to recent climate change. - Ecography 36: 313–322.

Loreau, M. 2010. Linking biodiversity and ecosystems: towards a unifying ecological theory. - Philos. Trans. R. Soc. B Biol. Sci. 365: 49–60.

Loreau, M. and De Mazancourt, C. 2013. Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. - Ecol. Lett. 16: 106–115.

Lurgi, M., López, B. C. and Montoya, J. M. 2012. Climate change impacts on body size and food web structure on mountain ecosystems. - Philos. Trans. R. Soc. B Biol. Sci. 367: 3050–3057.

Lyons, S. K., Amatangelo, K. L., Behrensmeyer, A. K., Bercovici, A., Blois, J. L., Davis, M., DiMichele, W. A., Du, A., Eronen, J. T. and Faith, J. T. 2016. Holocene shifts in the assembly of plant and animal communities implicate human impacts. - Nature 529: 80.

Martin, P. R. and Bonier, F. 2018. Species interactions limit the occurrence of urban-adapted birds in cities. - Proc. Natl. Acad. Sci. 115: E11495–E11504.

Martin, K., Aitken, K. E. and Wiebe, K. L. 2004. Nest sites and nest webs for cavity-nesting communities in interior British Columbia, Canada: nest characteristics and niche partitioning. - The condor 106: 5–19.

McFadden, D. 1974. Conditional logit analysis of qualitative choice behavior. - In: Frontiers in econometrics. P. Zarembka. Academic Press, New York.

McGill, B. J., Enquist, B. J., Weiher, E. and Westoby, M. 2006. Rebuilding community ecology from functional traits. - Trends Ecol. Evol. 21: 178–185.

Mérigot, B., Durbec, J.-P. and Gaertner, J.-C. 2010. On goodness-of-fit measure for dendrogram-based analyses. - Ecology 91: 1850–1859.

Mokross, K., Ryder, T. B., Côrtes, M. C., Wolfe, J. D. and Stouffer, P. C. 2014. Decay of interspecific avian flock networks along a disturbance gradient in Amazonia. - Proc. R. Soc. B Biol. Sci. 281: 20132599.

Mönkkönen, M., Devictor, V., Forsman, J. T., Lehikoinen, A. and Elo, M. 2017. Linking species interactions with phylogenetic and functional distance in European bird assemblages at broad spatial scales. - Glob. Ecol. Biogeogr. 26: 952–962.

Morueta-Holme, N., Blonder, B., Sandel, B., McGill, B. J., Peet, R. K., Ott, J. E., Violle, C., Enquist, B. J., Jørgensen, P. M. and Svenning, J.-C. 2016. A network approach for inferring species associations from co-occurrence data. - Ecography 39: 1139–1150.

Mouchet, M., Guilhaumon, F., Villéger, S., Mason, N. W., Tomasini, J.-A. and Mouillot, D. 2008. Towards a consensus for calculating dendrogram-based functional diversity indices. - Oikos 117: 794–800.

Mouchet, M. A., Villéger, S., Mason, N. W. and Mouillot, D. 2010. Functional diversity measures: an overview of their redundancy and their ability to discriminate community assembly rules. - Funct. Ecol. 24: 867–876.

Orchan, Y., Chiron, F., Shwartz, A. and Kark, S. 2013. The complex interaction network among multiple invasive bird species in a cavity-nesting community. - Biol. Invasions 15: 429–445.

Ovaskainen, O., Abrego, N., Halme, P. and Dunson, D. 2016. Using latent variable models to identify large networks of species-to-species associations at different spatial scales. - Methods Ecol. Evol. 7: 549–555.

Paradis, E., Blomberg, S., Bolker, B., Brown, J., Claude, J., Cuong, H. S. and Desper, R. 2019. Package 'ape.' - Anal. Phylogenetics Evol. Version in press.

Petanidou, T., Kallimanis, A. S., Tzanopoulos, J., Sgardelis, S. P. and Pantis, J. D. 2008. Long-term observation of a pollination network: fluctuation in species and interactions, relative invariance of network structure and implications for estimates of specialization. - Ecol. Lett. 11: 564–575.

Plein, M., Längsfeld, L., Neuschulz, E. L., Schultheiß, C., Ingmann, L., Töpfer, T., Böhning-Gaese, K. and Schleuning, M. 2013. Constant properties of plant–frugivore networks despite fluctuations in fruit and bird communities in space and time. - Ecology 94: 1296–1306.

Poisot, T. and Gravel, D. 2014. When is an ecological network complex? Connectance drives degree distribution and emerging network properties. - PeerJ 2: e251.

Poisot, T., Stouffer, D. B. and Gravel, D. 2015. Beyond species: why ecological interaction networks vary through space and time. - Oikos 124: 243–251.

Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., Vesk, P. A. and McCarthy, M. A. 2014. Understanding co-occurrence by modelling species simultaneously with a Joint Species Distribution Model (JSDM) (J McPherson, Ed.). - Methods Ecol. Evol. 5: 397–406.

R Core Team 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www. R-project. org.

Ramos-Robles, M., Andresen, E. and Díaz-Castelazo, C. 2016. Temporal changes in the structure of a plant-frugivore network are influenced by bird migration and fruit availability. - PeerJ 4: e2048.

Reif, J. and Hanzelka, J. 2020. Continent-wide gradients in open-habitat insectivorous bird declines track spatial patterns in agricultural intensity across Europe. - Glob. Ecol. Biogeogr. 29: 1988–2013.

Rigal, S. and Knape, J. 2024. Investigating the heterogeneity within Wild bird indices in Europe. - Biol. Conserv. 290: 110452.

Rigal, S., Devictor, V., Gaüzère, P., Kéfi, S., Forsman, J. T., Kajanus, M. H., Mönkkönen, M. and Dakos, V. 2022. Biotic homogenisation in bird communities leads to large-scale changes in species associations. - Oikos 2022: e08756.

Rigal, S., Dakos, V., Alonso, H., Auniņš, A., Devictor, V., et al. 2023. Farmland practices are driving bird population decline across Europe. - Proc. Natl. Acad. Sci. 120: e2216573120.

Rigal, S., Devictor, V. and Dakos, V. 2024. Code and data for article Rapid structural network changes in bird communities.

Sander, E. L., Wootton, J. T. and Allesina, S. 2017. Ecological network inference from long-term presence-absence data. - Sci. Rep. 7: 7154.

Seppänen, J.-T., Forsman, J. T., Mönkkönen, M. and Thomson, R. L. 2007. Social information use is a process across time, space, and ecology, reaching heterospecifics. - Ecology 88: 1622–1633.

Sokal, R. R. and Rohlf, F. J. 1962. The comparison of dendrograms by objective methods. - Taxon: 33–40.

Staniczenko, P. P., Sivasubramaniam, P., Suttle, K. B. and Pearson, R. G. 2017. Linking macroecology and community ecology: refining predictions of species distributions using biotic interaction networks. - Ecol. Lett. 20: 693–707.

Storchová, L. and Hořák, D. 2018. Life-history characteristics of European birds. - Glob. Ecol. Biogeogr. 27: 400–406.

Stouffer, D. B. and Bascompte, J. 2010. Understanding food-web persistence from local to global scales. - Ecol. Lett. 13: 154–161.

Stouffer, D. B. and Bascompte, J. 2011. Compartmentalization increases food-web persistence. - Proc. Natl. Acad. Sci. 108: 3648–3652.

Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M. and Munch, S. 2012. Detecting causality in complex ecosystems. - Science 338: 496–500.

Suzuki, S. S., Baba, Y. G. and Toju, H. 2023. Dynamics of species-rich predator–prey networks and seasonal alternations of core species. - Nat. Ecol. Evol. 7: 1432–1443.

Tsonis, A. A., Deyle, E. R., Ye, H. and Sugihara, G. 2018. Convergent cross mapping: theory and an example. - In: Advances in Nonlinear Geosciences. Springer, pp. 587–600.

Tylianakis, J. M. and Morris, R. J. 2017. Ecological networks across environmental gradients. - Annu. Rev. Ecol. Evol. Syst. in press.

Tylianakis, J. M., Laliberté, E., Nielsen, A. and Bascompte, J. 2010. Conservation of species interaction networks. - Biol. Conserv. 143: 2270–2279.

Ushio, M., Hsieh, C., Masuda, R., Deyle, E. R., Ye, H., Chang, C.-W., Sugihara, G. and Kondoh, M. 2018. Fluctuating interaction network and time-varying stability of a natural fish community. - Nature 554: 360–363.

Wood, S. N. 2003. Thin plate regression splines. - J. R. Stat. Soc. Ser. B Stat. Methodol. 65: 95–114.

Wood, S. N. 2017. Generalized additive models: an introduction with R. - Chapman and Hall/CRC.

Zou, F., Jones, H., Colorado Z., G. J., Jiang, D., Lee, T.-M., Martínez, A., Sieving, K., Zhang, M., Zhang, Q. and Goodale, E. 2018. The conservation implications of mixed-species flocking in terrestrial birds, a globally-distributed species interaction network. - Biol. Conserv. 224: 267–276.