Benchmarking long-read genome assemblers for three sequencing protocols and three agricultural species

Clément Birbes[1,2], Andreea Dréau[1,2], Denis Milan[2], Carole Iampietro[2], Christine Gaspin[1], Cécile Donnadieu[2], Matthias Zytnicki[1] & Christophe Klopp[1]

[1] Université Fédérale de Toulouse, INRAE, MIAT, BioinfOmics – 31 326, Castanet-Tolosan, France
[2] INRAE, US 1426, GeT-PlaGe, Genotoul, France Genomique, Université Fédérale de Toulouse – 31 326, Castanet-Tolosan, France

Correspondence: christophe.klopp@inrae.fr

Abstract

Today, long read technologies make it possible to produce telomere-to-telomere genome assemblies. These assemblies permit more accurate whole genome analyses, including on repeated regions.

The genome assembly process can chain several steps such as read correction, contigs assembly, contig polishing, scaffolding and gap filling, among others. Each step usually requires at least one software package, and input sequences. For the end user, it is not necessarily clear which combination of tools, together with read types, will work best for a given genome.

In this work, we will focus on contig production, which is a central and complex task of the assembly process. It aims at producing the longest, errorless, sequences, called *contigs*, from reads. While it is possible to produce contigs from short reads, long reads are now widely preferred, since they produce much longer contigs.

In this work, we evaluate several contig producing software packages (usually named *assemblers*), on long reads generated by two sequencers using three protocols, for three eukaryotic, complex species with different characteristics. Our aim is twofold. First, we would like to give readers insight on the impact of sequencing technology and assembler combinations, in order to help them make their choice for a given genome. Second, we would like to present different assembly metrics and provide a critical view on their interpretation.

Keywords: Long read; Benchmark; Evaluation; de novo Assembly

Introduction ₂₁

Today many studied species benefit from at least one reference genome assembly. A genome assembly is a text file usually in FASTA format containing sequences representing chromosomes or chromosome parts. Genome assemblies enable a large set of genomic analysis, including gene expression quantification, genomic variation calling, ChIP-seq peak discovery, methylation state measurement, etc.

With the advent of long read sequencing technologies provided by companies such as PacBio and Oxford Nanopore Technologies, the assembly of small genomes up to tens of megabases is a solved question and the assembly of large genome has been tremendously simplified. It can now be performed by a single researcher for a few thousand dollars compared to the large consortia needed twenty or thirty years ago. It is thus now possible to study new levels of genomic complexities, at a reasonable price.

Genomes can differ in several ways, some of which can be linked. Size has been mentioned before and repeat content is often linked to size. The longer the repeats and the more resembling the more difficult is their disentanglement, and therefore the assembly. Heterozygocity is another characteristic impacting genome assembly. The lower the heterozygocity the easier the assembly, explaining why some projects use double haploids. Recent whole genome duplication will also render assembly more difficult because only few variations are found between chromosomes sharing the same ancestor. These cases being rare, they are often not taken into account in the assembly software packages.

The aim of genome assembly is to produce a *bona fide* representation of the genome. Many strategies can be used, but most now rely on long reads. These reads, which can contain errors at varying degrees, can first be corrected using other reads or on themselves. They are then assembled together, in order to form longer sequences. These sequences can then be polished in order to remove errors and scaffolded into longer pseudo-molecules. In this article, we will only focus on the contig assembly part.

In principle, an assembler takes a set of input reads produced by a sequencer, and stitches them using common sequence part or sequence overlaps, in order to produce the longest possible contigs. Long read assemblers rely on two different methods: Overlap Layout Consensus (OLC) graphs, or de Bruijn graphs. OLC assemblers are comparing the reads to themselves in order to build an overlap graph. De Bruijn assemblers also use a graph, which is built from k-mers, which are overlapping read chunks. In principle, each path should represent chromosome parts called *contigs*, which are extracted from the graph. Although simple at first glance, genome assembly is a highly complex task. Repetition and sequencing errors tend to add spurious information in often very large graphs, which become highly tangled, and detecting this spurious information is far from straight-forward. Moreover, contigs should be as long as possible, and contain as few errors as possible. Likewise, the tools should be fast, and not consume too much memory. As a consequence, each tool implements different strategies in order to solve this multi-objective problem.

Given the wealth of tools, it is necessary to assess their performances through the quality of the produced assemblies. Here again, several metrics can be used. Some metrics only measure the contig sizes. Other metrics compare the assembly with a reference. This reference can be an estimated genome size, a reference genome, a set of reference genes, or the reads themselves. All these references give a fraction of information, which can be used to assess assembly quality. Taken individually, these metrics can be misleading. Our aim is to describe and show their limitations, and to suggest a set of metrics that can faithfully assess the assembly quality.

This article is based on long read datasets produced during the SeqOccIn project which was focused on

species of agronomic interest. This project aimed at comparing long read sequencing technologies to build state of the art pipelines for several applications, including genome assembly, variation and methylation calling as well as meta-genomic analysis. This article will include read sets from three species : quail (*Coturnix Japonica*), pig (*Sus scrofa*) and cattle (*Bos taurus*) and from three sequencing protocols Oxford Nanopore technologies R9.4, PacBio CLR and PacBio HiFi. We present the result of 39 assemblies performed using eight software packages (flye, hifiasm, ipa, lja, NextDenovo, raven, shasta, wtdbg2).

The species have been chosen because of their agronomic interest and their diversity. They have different genome sizes and repeat content. One genome harbors micro-chromosomes which are known to be difficult to sequence and assemble. They all have a reference genome assembly available at the NCBI and EMBL/EBI.

Material and methods

Estimated genome sizes

The three genomes vary in size. Although reference assemblies are available, we wanted to use an external, more reliable, estimation of genome size. We used estimations based on flow cytometry, ultraviolet microscopy or biochemical analysis. Figures provided by the Animal Genome Database (Gregory, 2023) and MaizeGDB (Bennett, 1995) are:

- *Bos taurus*: 3.08–3.84 Gbp

- *Coturnix japonica*: 1.26–1.38 Gbp

- *Zea mays*: 2.41–2.74 Gbp

We chose the average values, which are, respectively, 3.53 Gbp, 1.32 Gbp, and 2.57 Gbp.

Overview of current reference assemblies

The genomes have specific characteristics presented in Table 1.

All three genomes have already been assembled and released several times. The NCBI provides several statistics on these assemblies, some of which are provided in Table 1. These assemblies have all been generated using long read technologies. A first observation is that the assembly sizes are smaller than the expected genome size, which is commonly observed. Second, none of them are assembled telomere-to-telomere, and the numbers of scaffolds range from 687 to 2,531. Third, the percentage of repetition greatly varies between organisms from 8% to 79%. Although we expect highly repeated genomes to be more difficult to assemble, the variability in repetition is crucial: it is much more difficult to assemble genomes that contain exact repetitions than genomes where repetitions have accumulated errors during evolution.

Assembly quality metrics

Several metrics can be used to assess the quality of an assembly. They include the NG50 and the LG50, which assess assembly contiguity: the NG50 will be high if the contigs are very long. However, a high NG50 does not imply a good assembly. An assembler that would stitch together contigs from different *loci* would also produce a high NG50, at the expense of the assembly quality.

Other approaches compare the assembly with a reference that is deemed reliable. We first computed the relative size, which is $\frac{o-e}{e}$, where o is the observed size of the assembly, and e is the expected size, obtained from cytometry. The relative size is positive if the assembly is longer than the expected one, and negative otherwise.

Table 1. Metrics on the current assemblies, computed by the NCBI.

	Bos taurus	*Coturnix japonica*	*Zea mays*
Assembly name	ARS-UCD1.2	Coturnix japonica 2.1	Zm-B73-REFERENCE-NAM-5.0
Assembly size	2,715,853,792	927,656,957	2,182,786,008
# scaffolds	2,211	2,531	687
Scaffold N50	103,308,737	2,975,000	226,353,449
# chromosomes and plasmids	31	32	12
# protein-coding genes	21,039	16,026	34,337
# non protein-coding genes	9,357	4,736	10,366
# mRNAs	63,675	40,881	57,304
% repetition	49.66%	7.82%	78.51%
reference	(NCBI, 2023a)	(NCBI, 2023b)	(NCBI, 2023c)

We also computed coverage metrics, provided by Inspector (Y Chen, Zhang, et al., 2021). It uses a reference assembly, maps the contigs to this assembly, and provided the proportion of the genome which is covered. This metrics is meaningful if the assembly and the reference genome are close enough.

The Merqury package (Rhie, Walenz, et al., 2020) provides several metrics that compare the assembly with the reads themselves. The QV is an estimation of the probability of error for the assembly at the nucleotide level, computed as a Phred score. The completeness estimates the quantity of the nucleotides that have been used in the assembly, divided by the number of nucleotides that have been sequenced. Inspector (Y Chen, Zhang, et al., 2021) also computes the number of assembly errors, categorized as expansions (gaps in read alignment), collapses (extra sequence in reads), and inversions (inverted read alignments). In our analysis, we grouped them as errors. Of note, these metrics based on reads are biased towards the sequencing quality. If the sequencer produces (non-random) errors, these metrics can be high, even though the assembly is far from the actual genome.

The BUSCO tool (Feron and Waterhouse, 2022) uses a list of genes that have been manually curated, and are expected to be present in exactly one copy in the phylum of the species. The *complete*, or BUSCO C, score is the percentage of genes that are found once in the assembly, without major error. The *duplicate*, or BUSCO D, score is the percentage of genes that are found more than once in the assembly. These BUSCO scores vary greatly with the base level quality of the assembly: a few nucleotide errors in an assembled gene may truncate the open reading frame, and hinder BUSCO to detect this gene.

Table 2 links metrics with the types of misassemblies it can detect.

Table 2. Possible sources of errors in genome assemblies.

Type of error	Metrics associated	Possible reasons
Duplication of sequences	BUSCO D	Di- or polyploidy
Collapsed repetitions	relative size	Genome repetitions
Fragmented assembly	NG50, LG50	Genome repetitions
Short sequencing errors	QV	Errors in reads
Long sequencing errors	# errors	Genome repetitions
Unused read sequences	completeness	All the above

The time spent, as well as the memory used by each tool, has also been added to the list of metrics. They can be used to choose whether a given tool can be used, or not, on the available computers. In order to compare the results on the different genomes, they have been divided by the estimated genome size.

We classified each metrics using a color code, in order to ease the table reading. Each cell was colored in green if a criteria is fully satisfied (high quality), red if it is not satisfied (low quality), and orange if it is in between (medium quality). We tried to use a criteria that was previously published. The quality criteria, the

colour code and the corresponding bibliographic references can be found on Table 3.

Table 3. Metrics and thresholds used to evaluate assemblies and assemblers.

	low	medium	high	references
Contig NG50	< 1 Mb	1 Mb $\leq\ \leq 10$ Mb	> 10 Mb	(Rhie, McCarthy, et al., 2021)
Contig LG50	> 200	$50 \leq\ \leq 200$	> 200	
k-mer completeness	$< 80\%$	$80\% \leq\ \leq 95\%$	$> 95\%$	(Rhie, McCarthy, et al., 2021; Rhie, Walenz, et al., 2020)
genome coverage difference	$> 40\%$	$40\% \leq\ \leq 20\%$	$< 20\%$	
reference assembly coverage difference	$> 20\%$	$20\% \leq\ \leq 10\%$	$< 10\%$	
BUSCO complete	$< 80\%$	$80\% \leq\ \leq 95\%$	$> 95\%$	(Rhie, McCarthy, et al., 2021)
BUSCO duplicated	$> 5\%$	$2\% \leq\ \leq 5\%$	$< 2\%$	
Basepair QV	< 30	$30 \leq\ \leq 50$	> 50	(Rhie, McCarthy, et al., 2021; Rhie, Walenz, et al., 2020)
Assembly errors	> 100	$20 \leq\ \leq 100$	< 100	
memory	> 125GB/Gbp	32GB/Gbp $\leq\ \leq 125$GB/Gbp	< 32GB/Gbp	
time	> 1 week on 8 cores per Gbp	1 day on 8 cores per Gbp $\leq\ \leq 1$ week on 8 cores per Gbp	< 1 day on 8 cores per Gbp	

Read sets

The reads generated by the sequencing protocols have also specifics characteristics presented in Table 4.

The *Bos taurus* datasets have been first published by Eché et al., 2023, whereas the other datasets will be published in other data papers. All datasets are available at EMBL-EBI ENA archive. The corresponding accessions can be found in in Table 5.

Table 4. Metrics on long read datasets used in the benchmark.

Species	technology	# reads	# base pairs	mean length (bp)	coverage
Bos taurus	HiFi	7,430,931	117,560,145,736	15,820.4	33.30
Bos taurus	ONT	10,144,221	156,488,832,730	15,426.4	44.33
Bos taurus	CLR	14,765,644	279,331,710,507	18,917.7	102.85
Zea mays	HiFi	5,976,900	75,601,454,246	12,648.9	29.42
Zea mays	ONT	12,698,704	145,838,781,573	11,484.5	56.75
Zea mays	CLR	6,114,831	130,950,539,094	21,415.2	50.95
Coturnix japonica	HiFi	1,851,480	30,108,307,131	16,261.8	22.81
Coturnix japonica	ONT	4,147,888	65,000,070,596	15,670.6	49.24

Table 5. Public accession number of long reads used in the benchmarks.

Species	technology	accessions
Bos taurus	HiFi	ERR10378054 to ERR10378058
Bos taurus	ONT	ERR10386215 to ERR10386219
Bos taurus	CLR	ERR10378053
Zea mays	HiFi	ERR14085295, ERR14085303, ERR14085304
Zea mays	ONT	ERR14056479 to ERR14056485
Zea mays	CLR	ERR14085305, ERR14085306
Coturnix japonica	HiFi	ERR11528759, ERR11528760
Coturnix japonica	ONT	ERR11591487, ERR11591488

Overview of long read de novo assembly methods

Choosing an assembler is not easy because novel software packages are coming out regularly and existing assemblers change version sometimes frequently. The criteria used to chose the assemblers were mainly their actual usage in the community and their efficacy. Canu (Koren et al., 2017) and necat (Y Chen, Nie, et al., 2021) were left out because of the number of computing hours need to assemble a large genome.

Initial long read genome assembly software packages list is given in Table 6.

Table 6. Assembler versions and technologies used in the benchmark.

Assembler	Version	ONT	CLR	HiFi	Reference	Comments
canu	2.2	✗	✗	✗	(Koren et al., 2017)	
flye	2.9	✓	✓	✓	(Kolmogorov et al., 2019)	contig polishing
hifiasm	0.16.1	✗	✗	✓	(Cheng et al., 2022)	read correction
ipa	1.0.3-0	✗	✗	✓	(Sovic et al., 2022)	
lja	V0.1	✗	✗	✓	(Bankevich et al., 2022)	
necat	20200803	✗	✗	✗	(Y Chen, Nie, et al., 2021)	
NextDenovo	2.5.0	✓	✓	✗	(Hu et al., 2023)	
raven	1.8.1	✓	✗	✗	(Vaser and Šikić, 2021)	
shasta	0.5.1	✓	✓	✗	(Shafin et al., 2020)	
wtdbg2	2.5	✓	✓	✗	(Ruan and Li, 2019)	

Not all data types, species, assembler combinations could be run because CLR data were not available for *Coturnix japonica*. Some assemblers are data specific: for example, hifiasm, ipa, lja cannot assemble ONT nor CLR reads. Finally, some combinations did not terminate correctly, despite several tries: shasta on *Zea mays* ONT reads, and ipa on HiFi reads for the same species.

Experimental setup

The benchmark was performed on 2.1GHz Intel Xeon E5-2683 v4 CPUs, on CentOS 7 Linux distributions.

Results

The names of the assemblies presented in this section will join the sequencing protocole (HiFi, CLR, ONT) and the assembler (flye, hifiasm, ipa, lja, NextDenovo, raven, shasta, or wtdbg2).

Bos taurus results

We can note at first glance that three assemblies —namely ONT raven, CLR raven, and CLR shasta— are not satisfactory, observing several metrics such as completeness, genome coverage, BUSCO C, QV, and number

of errors. These assemblies are significantly too short to be useful, and will not be discussed in the rest of the analysis.

type	assembler	NG50	LG50	comp.	rel. size	cov.	B. C	B. D	QV	err.	time	mem.
Ref.		82.40	16	NA	−26%	NA	96%	2%	NA	NA	NA	NA
ONT	flye	26.38	33	93%	−25%	99%	86%	1%	31	25	33d	88.3
ONT	NextDenovo	58.92	21	92%	−25%	99%	85%	2%	31	37	108d	16.9
ONT	raven	12.92	70	37%	−26%	7%	9%	0%	15	14893	6d	27.5
ONT	shasta	18.13	55	91%	−26%	98%	81%	1%	29	779	<1d	205.3
ONT	wtdbg2	14.64	56	87%	−23%	97%	75%	1%	25	146	62d	97.3
CLR	flye	12.29	66	96%	−25%	99%	95%	2%	36	871	55d	90.3
CLR	NextDenovo	42.98	23	95%	−25%	99%	92%	2%	34	571	358d	16.4
CLR	raven	10.69	76	14%	−22%	0%	1%	0%	12	112914	3d	29.8
CLR	shasta	0.09	6843	54%	−43%	16%	35%	0%	19	213	<1d	127.2
CLR	wtdbg2	9.82	80	93%	−25%	97%	90%	1%	29	393	25d	71.6
Hifi	flye	0.72	998	98%	−2%	99%	94%	6%	49	58	20d	62.8
Hifi	hifiasm	76.68	17	96%	−9%	100%	96%	2%	66	13	12d	47.3
Hifi	ipa	32.46	30	96%	−16%	99%	96%	3%	56	67	15d	9.7
Hifi	lja	1.09	942	100%	62%	100%	91%	26%	50	2	18d	77.2
Hifi	raven	2.80	301	92%	−24%	99%	87%	1%	45	342	1d	23.1
Hifi	wtdbg2	18.30	47	93%	−27%	96%	94%	2%	41	52	11d	43

Table 7. Metrics for *Bos taurus*. Description of abbreviations: NG50: NG50 in MB, comp: completeness, cov.: genome coverage, B. C: BUSCO C, B. D: BUSCO D, err.: number of errors, mem.: memory in GB.

NG50 and LG50, together show that HiFi lja or HiFi raven provide the most fragmented assemblies. On the other end, HiFi hifiasm, HiFi ipa, CLR NextDenovo, ONT NextDenovo and ONT flye output the least fragmented genome assemblies.

A striking result is that the genome coverage is very high for all remaining assemblies, meaning that almost all the reference genome is covered by the remaining assemblies. However, completeness, which is based on k-mers instead of alignments, gives another picture: the completeness of most ONT and CLR assemblies (with the notable exception of CLR flye), as well as HiFi raven and HiFi wtdbg2, is below 95%. This suggests that ONT and CLR assemblies, as expected, contain small errors, although flye corrects these errors in its pipe-line. This result is confirmed by the BUSCO C score, the QV, and the number of errors: best scores are given to HiFi assemblies, while CLR assemblies have slightly better scores than ONT. Of note, the error metrics is sensitive to any kind of mutation, even point mutations, which could give rise to frame-shift or early stop codons in the translated proteins.

This is also confirmed, to some extent, by the number of errors, which is higher for CLR and ONT data, compared to HiFi.

Another interesting result is the relative size, compared to cytology analysis. Most assemblies, including the reference one, are significantly shorter than the predicted size. This could suggest that the repeated fraction of the genome is poorly assembled. HiFi flye, HiFi hifiasm and to a lesser degree, HiFi ipa are the only assemblies which sizes are close to the goal. However, the assembly size of HiFi flye is somewhat misleading because it has a BUSCO D of 6% clearly suggesting that it is partly duplicated. If we link this result with the low NG50, and the high LG50, this suggests that HiFi flye kept haplotypes apart, cutting the assembly each time two alleles where present and keeping both of them.

It is also fruitful to compare the results produced by the same tool, and different data. NextDenovo, for instance, has been used on ONT and CLR data. NextDenovo ONT gives a more contiguous assembly (NG50: 59M vs 43M, LG50: 21 vs 23). However, NextDenovo CLR shows less errors (BUSCO C: 92% vs 85%, QV: 34 vs 31), and reflects the reads better (completeness: 95% vs 92%). Other tools where used with the three

technologies: flye, raven and wtdbg2. They also confirm that ONT assemblies are longer, but more error-prone than CLR assemblies. There are fewer errors in HiFi assemblies, when compared to CLR assemblies (for HiFi wtdgb2 and CLR wtdbg2 , BUSCO C: 94% vs 90%, QV: 41 vs 29). The assembly size however, is not directly linked to the type of data: HiFi wtdbg2 assembly has a longer NG50 (18M vs 10M) and a smaller LG50 (47 vs 80), but we observe the opposite for HiFi raven (NG50: 3M vs 13M, LG50: 301 vs 70) and HiFi flye (NG50: 700K vs 26M, LG50: 998 vs 33). This is a probable consequence of the fact that flye and raven separate haplotypes.

Coturnix japonica results

Assemblies of *Coturnix japonica* are much more fragmented than *Bos taurus*, and the NG50 ranges between 500k (for HiFi raven and flye) to 13M (for CLR NextDenovo). The NG50 is actually sensible to the genome size, and the size of the *Coturnix japonica* genome is much smaller than the size of the *Bos taurus* genome. Quail has also few large chromosomes and a large number of micro-chromosomes which limits the possible NG50 size.

type	assembler	NG50	LG50	comp.	rel. size	cov.	B. C	B. D	QV	err.	time	mem.
Ref.		31.42	8	NA	−32%	NA	96%	0%	NA	NA	NA	NA
ONT	flye	9.58	28	86%	−24%	98%	88%	0%	26	129	43d	157.6
ONT	NextDenovo	13.00	21	86%	−24%	98%	88%	0%	27	179	60d	43.4
ONT	raven	5.10	65	39%	−23%	9%	28%	0%	14	3903	2d	37.8
ONT	shasta	5.42	52	84%	−27%	96%	86%	0%	26	745	<1d	212.7
ONT	wtdbg2	5.52	44	78%	−23%	94%	76%	0%	21	187	1d	15.1
Hifi	flye	0.81	465	99%	23%	98%	96%	28%	46	19	12d	129.3
Hifi	hifiasm	4.05	84	91%	−4%	98%	96%	1%	59	60	3d	35.4
Hifi	ipa	1.84	161	95%	1%	98%	97%	17%	56	27	8d	17
Hifi	lja	1.19	304	99%	30%	98%	96%	30%	45	4	3d	40.7
Hifi	raven	0.52	472	87%	−23%	97%	90%	0%	44	63	<1d	22.3
Hifi	wtdbg2	1.89	116	89%	−26%	97%	95%	0%	34	19	3d	47.9

Table 8. Metrics for *Coturnix japonica*.

Akin to the previous data-set, BUSCO C score, QV, and the number of errors, show that ONT assemblies contain more errors, compared to HiFi. Moreover, assembly metrics are longer with ONT data (longest NG50 is 10M for ONT flye, and shortest is 500k for HiFi raven).

The relative size shows that most tools only assemble part of the genome, with score around -24%, which is still higher than the reference assembly (-32%). HiFi flye, HiFi hifiasm, HiFi ipa, and HiFi lja are the only exceptions. Flye lja and, to a lesser degree, HiFi ipa have positive relative sizes with BUSCO D scores up to 30%. This suggests that both haplotypes have been included in the assemblies. HiFi hifiasm is the only assembly with a good relative size, and a modest BUSCO D score of 1%. Its completeness of 91%, which is lower than other assemblies (e.g. HiFi flye, HiFi ipa and HiFi lja) may confirm that only one allele was kept in the final assembly.

Zea mays results

Several assemblers including ONT raven, ONT wtdbg2, CLR raven, HiFi raven, failed on this dataset. They produce very short assemblies. The genome coverage of raven is really low (down to 0%) for both ONT and CLR read types. wtdbg2 NG50 did not exceed 170K on all read types. ONT NextDenovo, as well as HiFi flye, hifiasm, and lja, provided rather contiguous assemblies, with NG50 greater than 10M.

A first observation is that the genome coverage never exceeds 60%. At a first glance, this could imply that roughly half of the reference genomes is not syntenic with the assembled genome. However, the reason could be elsewhere. In order to compute the genome coverage, the assembled genome is mapped to the reference

type	assembler	NG50	LG50	comp.	rel. size	cov.	B. C	B. D	QV	err.	time	mem.
Ref.		217.87	5	NA	−20%	NA	95%	14%	NA	NA	NA	NA
ONT	flye	1.67	379	78%	−16%	49%	94%	13%	20	72	111d	135.7
ONT	NextDenovo	31.31	23	77%	−19%	39%	94%	12%	20	330	244d	25.7
ONT	raven	0.11	6418	23%	−30%	0%	11%	0%	12	0	8d	39.2
ONT	wtdbg2	0.12	6219	65%	37%	22%	84%	12%	15	0	17d	156.1
CLR	flye	4.12	158	98%	−18%	60%	98%	15%	42	138	145d	133.8
CLR	raven	0.75	1000	24%	−5%	0%	4%	0%	12	44982	6d	31.2
CLR	shasta	0.22	2647	87%	-31%	47%	86%	9%	23	429	<1d	189.6
CLR	wtdbg2	0.17	4617	91%	89%	50%	88%	20%	17	2	13d	126.7
Hifi	flye	10.39	63	98%	-17%	55%	98%	15%	63	13	16d	200.3
Hifi	hifiasm	53.62	13	98%	-15%	54%	98%	15%	67	5	7d	46
Hifi	lja	26.53	28	98%	-17%	60%	98%	15%	58	1	38d	63.3
Hifi	raven	0.00	21551	57%	-71%	25%	75%	6%	42	0	2d	24
Hifi	wtdbg2	0.07	8097	92%	-29%	54%	96%	15%	36	0	59d	103.2

Table 9. Metrics for *Zea mays*.

genome. Two copies of the same repeat in the assembled genome will like map to the same copy in the reference. As a consequence, the unmapped copy of the reference genome will be detected as uncovered, and reported as such.

The BUSCO D is also very low. However, this is most probably due to a whole genome duplication which only happened in this species (Schnable et al., 2009). As a consequence, the private copied genes are flagged as duplicated by BUSCO.

Above all, shasta assemblies seem problematic, since the CLR and ONT assemblies do not resemble the reference assembly (genome coverage of 0%). Confirming this observation, the relative size is almost always negative, except for ONT and CLR wtdbg2. On the other side, CLR flye, as well as HiFi flye, hifiasm, and lja, show a completeness of 98%, indicating that almost all the reads have been used in the assembly. This may suggest that the reference genome is very divergent from the assembled one, or that repeats are poorly assembled.

Another striking result is that, although the assemblies are smaller than expected, the BUSCO C score can reach 98% (for CLR flye, as well as HiFi flye, hifiasm, and lja), which is higher than the BUSCO C score for the reference genome of 95%. This suggests that essential genes are found in these assemblies.

The BUSCO D score, however, is not less than 9%, including the reference genome except for raven assemblies. This strongly suggests that the genome is highly repeated, and that repetitions include genes.

As previously observed, HiFi assemblies include less errors than other assemblies.

Discussion

Assembly accuracy

Choice of the sequencing strategy

The type of sequenced reads is critical to obtain a high quality assembly: HiFi reads deliver better assemblies, with longer NG50, smaller LG50, reflecting reference genomes and reads better, with better BUSCO scores. CLR data, on our benchmarks, seem to be slightly better than ONT reads, especially when used with flye, which corrects read errors.

The assemblers also provide very distinct results, depending on the sequencing strategy. Hifiasm, only applicable to HiFi data, seems to provide the best results. Flye provides the best results for CLR reads, and NextDenovo, the best results for ONT reads.

Impact of the genome

Complexity of sequenceed genomes does also impact assembly quality. Assemblies appear to be more fragmented when genomes are heterozygous. As shown in the *Bos taurus* example, some tools may cut the assembly each time they observe several alleles, and possibly report them. This may obviously over-estimate the size of the assembly, especially if alternative allele are not appropriately handled. New pipe-lines, which include verkko (Rautiainen et al., 2023), are specifically designed to assemble diploid genomes, and provide haplotyped assemblies.

Genomes with many repetitions may also be poorly assembled. The different copies of the repetitions may be "collapsed" into a single sequence. In order to solve this problem, assemblers relies on small differences that appear between copies. Discriminating between real differences and sequencing errors is a key feature in order to correctly assemble reads.

For this, the tool should be able to differentiate real differences, and sequencing errors. This is especially difficult for ONT reads, or uncorrected CLR reads, which accumulate errors. For long, highly repetitive, regions, consortium for telomere-to-telomere assemblies (Nurk et al., 2022) suggests the use of ultra-long reads, with size larger than 100kb, in order to assemble them.

Time and memory usage

Concerning time, NextDenovo is by far the slowest tool, and flye usually is the second slowest. shasta is the fastest tool, and raven the second fastest. ipa and lja usually are between the two previous groups. wtdbg2 shows an irregular profile: it can be almost as fast as raven (*Zea mays* with ONT data), and slower than flye (*Bos taurus* with ONT data). This tool seems to be highly dependant on both the species and the data. flye also seems to be much faster on HiFi data.

Concerning memory, shasta requires more memory than other tools, flye is usually second, and wtdbg2, usually third. NextDenovo, raven, and ipa require less memory than other tools.

NextDenovo and shasta exhibit an opposite profile: the former requires more time and less memory, the latter requires less time and more memory. This trade-off is most likely the consequence of implementation choices.

Raven and ipa are both fast, and require less memory than other tools. On the opposite, flye is rather slow, and requires more memory than most other tools.

Metrics

Defining a high-quality assembly is straight-forward: it accurately provides the series of nucleotides that form each chromosome. However, there are many types of low-quality assemblies: missing regions, duplications, etc. The metrics we used make it possible to assess the different possible types of errors in the assemblies. Obviously, most metrics evaluate several criteria at the same time: the NG50 will be low if the assembly is fragmented, and also if the genome is incomplete. The BUSCO C, for instance, may be used to evaluate unused read sequences, duplication of sequences, and short sequencing errors.

Some metrics should be considered with extra caution, especially when the genome is highly repeated. The *Zea mays* genome, for instance, shows very poor genome coverage and BUSCO D results. A telomere-to-telomere assembly (J Chen et al., 2023), which has been published after this analysis was carried out, informed that the genome contains many tandem (near exact) repeats, highly repeated centromeric, telomeric and subtelomeric regions, and numerous satellite and super-long simple-sequence-repeat arrays. This, together with a recent whole genome duplication, are crucial information that make it possible to correctly interpret poor metrics.

Acknowledgements

We are grateful to the GenoToul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, https://doi.org/10.15454/1.5572369328961167E12) for providing help and/or computing and/or storage resources.

We thank members of SeqOccIn Consortium for their feedback before and during this study.

Funding

We thank "La Région Occitanie" and European Union for financing project in the call for projects "Regional Platforms of Research and Innovation" of the Occitanie region on the Operational Program FEDER-FSE MIDI-PYRENEES ET GARONNE 2014–2020. APIS-GENE and the industrial organisations of the ruminant sector CNIEL, Interbev, Eliance, CNE and Institut de l'Elevage contributed to the financing of the cattle analyses. SYSAAF, ADISSEO, and Alliance R&D contributed to the financing of the quail analyses. KWS, Maïs Adour, Euralis, Caussade semences, Syngenta, Limagrain, and RAGT contributed to the financing of the maize analyses. The INRAE experimental unit of Le Pin (https://doi.org/10.15454/1.5483257052131956E12) and Nouzilly (PEAT, https://doi.org/10.15454/1.5572326250887292E12) produced the cattle and quail samples.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, script, code, and supplementary information availability

Assemblies are available on a dedicated repository https://doi.org/10.57745/63UXPG. Tool versions, and commands used are available on https://forgemia.inra.fr/seqoccin/assembler-benchmarking.

References

Bankevich A, AV Bzikadze, M Kolmogorov, D Antipov, and PA Pevzner (Feb. 2022). Multiplex de Bruijn graphs enable genome assembly from long, high-fidelity reads. *Nature Biotechnology* 40, 1075–1081. https://doi.org/10.1038/s41587-022-01220-6.

Bennett MD (1995). Maize GDB.

Chen J, Z Wang, K Tan, W Huang, J Shi, T Li, J Hu, K Wang, C Wang, B Xin, H Zhao, W Song, MB Hufford, JC Schnable, W Jin, and J Lai (June 2023). A complete telomere-to-telomere assembly of the maize genome. *Nature Genetics* 55, 1221–1231. https://doi.org/10.1038/s41588-023-01419-6.

Chen Y, F Nie, SQ Xie, YF Zheng, Q Dai, T Bray, YX Wang, JF Xing, ZJ Huang, DP Wang, LJ He, F Luo, JX Wang, YZ Liu, and CL Xiao (Jan. 2021). Efficient assembly of nanopore reads via highly accurate and intact error correction. *Nature Communications* 12. https://doi.org/10.1038/s41467-020-20236-7.

Chen Y, Y Zhang, AY Wang, M Gao, and Z Chong (Nov. 2021). Accurate long-read de novo assembly evaluation with Inspector. *Genome Biology* 22. https://doi.org/10.1186/s13059-021-02527-4.

Cheng H, ED Jarvis, O Fedrigo, KP Koepfli, L Urban, NJ Gemmell, and H Li (Mar. 2022). Haplotype-resolved assembly of diploid genomes without parental data. *Nature Biotechnology* 40, 1332–1335. https://doi.org/10.1038/s41587-022-01261-x.

Eché C, C Iampietro, C Birbes, A Dréau, C Kuchly, A Di Franco, C Klopp, T Faraut, S Djebali, A Castinel, M Zytnicki, E Denis, M Boussaha, C Grohs, D Boichard, C Gaspin, D Milan, and C Donnadieu (June 2023). A Bos taurus sequencing methods benchmark for assembly, haplotyping, and variant calling. *Scientific Data* 10. ISSN: 2052-4463. https://doi.org/10.1038/s41597-023-02249-1.

Feron R and RM Waterhouse (Feb. 2022). Assessing species coverage and assembly quality of rapidly accumulating sequenced genomes. en. *Gigascience* 11.

Gregory T (2023). Animal Genome Size Database.

Hu J, Z Wang, Z Sun, B Hu, AO Ayoola, F Liang, J Li, JR Sandoval, DN Cooper, K Ye, J Ruan, CL Xiao, DP Wang, DD Wu, and S Wang (Mar. 2023). An efficient error correction and accurate assembly tool for noisy long reads. https://doi.org/10.1101/2023.03.09.531669.

Kolmogorov M, J Yuan, Y Lin, and PA Pevzner (Apr. 2019). Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology* 37, 540–546. https://doi.org/10.1038/s41587-019-0072-8.

Koren S, BP Walenz, K Berlin, JR Miller, NH Bergman, and AM Phillippy (Mar. 2017). Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat separation. *Genome Research* 27, 722–736. https://doi.org/10.1101/gr.215087.116.

NCBI (2023a). NCBI Bos taurus Annotation Release 106.

— (2023b). NCBI Coturnix japonica Annotation Release 101.

— (2023c). NCBI Zea mays Annotation Release 103.

Nurk S et al. (Apr. 2022). The complete sequence of a human genome. *Science* 376, 44–53. https://doi.org/10.1126/science.abj6987.

Rautiainen M, S Nurk, BP Walenz, GA Logsdon, D Porubsky, A Rhie, EE Eichler, AM Phillippy, and S Koren (Feb. 2023). Telomere-to-telomere assembly of diploid chromosomes with Verkko. *Nature Biotechnology* 41, 1474–1482. https://doi.org/10.1038/s41587-023-01662-6.

Rhie A, SA McCarthy, et al. (Apr. 2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature* 592, 737–746. https://doi.org/10.1038/s41586-021-03451-0.

Rhie A, BP Walenz, S Koren, and AM Phillippy (Sept. 2020). Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. en. *Genome Biol.* 21, 245.

Ruan J and H Li (Dec. 2019). Fast and accurate long-read assembly with wtdbg2. *Nature Methods* 17, 155–158. https://doi.org/10.1038/s41592-019-0669-3.

Schnable PS et al. (Nov. 2009). The B73 Maize Genome: Complexity, Diversity, and Dynamics. *Science* 326, 1112–1115. https://doi.org/10.1126/science.1178534.

Shafin K, T Pesout, R Lorig-Roach, M Haukness, HE Olsen, C Bosworth, J Armstrong, K Tigyi, N Maurer, S Koren, FJ Sedlazeck, T Marschall, S Mayes, V Costa, JM Zook, KJ Liu, D Kilburn, M Sorensen, KM Munson, MR Vollger, J Monlong, E Garrison, EE Eichler, S Salama, D Haussler, RE Green, M Akeson, A Phillippy, KH Miga, P Carnevali, M Jain, and B Paten (May 2020). Nanopore sequencing and the Shasta toolkit enable efficient de novo assembly of eleven human genomes. *Nature Biotechnology* 38, 1044–1053. https://doi.org/10.1038/s41587-020-0503-6.

Sovic I, Z Kronenberg, C Dunn, D Barnett, S Kingan, and J Drake (2022). IPA HiFi Genome Assembler. https://github.com/PacificBiosciences/pbipa.

Vaser R and M Šikić (May 2021). Time- and memory-efficient genome assembly with Raven. *Nature Computational Science* 1, 332–336. https://doi.org/10.1038/s43588-021-00073-4.