Sex-related gut microbiota in three geographically separated Norway lobster (*Nephrops norvegicus*) populations

Polina Rusanova[1,2], Eleni Nikouli[3], Michele Casini[2,4], Gioacchino Bono[1,5], Elena Mente[6], Alexandra Meziti[7], Konstantinos Kormas[3*]

[1] Institute for Biological Resources and Marine Biotechnologies, National Research Council (IRBIM-CNR), Mazara del Vallo, Italy

[2] Department of Biological, Geological and Environmental Sciences (BiGeA), Laboratory of Marine Biology and Fisheries, University of Bologna, Bologna, Italy

[3] Department of Ichthyology and Aquatic Environment, School of Agricultural Sciences, 384 46 Volos, Greece

[4] Department of Aquatic Resources, Institute of Marine Research, Swedish University of Agricultural Sciences, Lysekil, Sweden

[5] Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STEBICEF), Palermo University, Palermo, Italy

[6] Laboratory of Ichthyology-Culture and Pathology of Aquatic Animals, School of Veterinary Medicine, Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece

[7] Department of Marine Sciences, University of the Aegean, University Hill, 81100, Mytilene, Greece

[*] Corresponding author: kkormas@uth.gr

32   **ABSTRACT**

Despite the ecological and economic interest of the Norway lobster (*Nephrops norvegicus*),

34   its gut microbiota remains largely understudied. The current study aimed at investigating the gut

bacterial microbiota in three geographically separated *N. norvegicus* populations from the

36   Mediterranean and the North Sea and detecting any potential sex-related microbiota differences, by

high-throughput sequencing of the V3-V4 16S rRNA gene diversity of the gut tissue. From the

38   Greek population, egg-bearing females were also caught. A total of 2,385 operational taxonomic

units (OTUs) were identified and between 417 and 1290 occurred in each population/sex group.

40   The dominant OTUs belonged to the Fusobacteriia and Bacteroidia (Sweden), Bacilli and

Gammaproteobacteria (Italy) and Spirochaetia and Bacilli (Greece) bacterial classes. In the eggs,

42   the Actinobacteria, Alphaproteobacteria and Gammproteobacteria prevailed. Four OTUs related to

the *Oceanispirochaeta*, *Kordiimonas, Desulfovibrio, Carboxylicivirga* genera and one unafilliated

44   OTU were positively correlated (p values between 0.001 and 0.04) to body size indicating their

potential role in the animal's nutrition and growth. No statistically significant differences were

46   found between males-females in each of the three populations. However, statistically significant

differences between populations for each sex, were found for all females (p values between 0.008

48   and 0.032) and for the males between the most distant populations, i.e. Italy-Sweden (p=0.021) and

Greece-Sweden (p=0.015). The eggs microbiota was statistically different from both the adult

50   females (p=0.027) and males (p=0.046) gut microbiota. Overall, this study showed that *N.

norvegicus* gut microbiota are differentiated between geographically distant populations and that

52   sex-related differences are not significant.

54   Keywords: crustacea, gut, microbiome, fisheries, North Sea, Mediterranean Sea, Aegean Sea

1. INTRODUCTION

58   The hologenome and holobiont concepts (Zilber-Rosenberg and Rosenberg, 2008) have

opened a new way we view animal life (Theis et al., 2016; Webster, 2017) for their ontogeny and

60   function (Stencel and Wloch-Salamon, 2022; Troussellier et al., 2017). Thus, during the last few

years, the scientific interest on holobiont research has boomed for animals, as well. However, there

62   are yet animal species from various habitats, whose microbiomes remain largely unknow, despite

their ecological and/or economic importance and the ongoing rapid technological advancements in

64   DNA/RNA technologies.

One such species is the Norway lobster *Nephrops norvegicus* (Linnaeus, 1758) a decapod

66   crustacean, langoustine or scampi, which is a typical clawed lobster with a slender body, long claws

and large dark eyes. *Nephrops norvegicus* is considered one of the most important species for

68   fisheries, and is considered a particularly valuable commercial crustacean species in Europe. It is a

benthic crustacean dependent on muddy-type sediments suitable for the construction of burrows. It

70   is widely distributed across the continental shelves of the northeast Atlantic Ocean and the

Mediterranean Sea, from Iceland and Norway in the north, to Morocco and the Adriatic seas in the

72   south (Bell et al., 2013). Populations dwelling in colder waters around Iceland and the Faroe Islands

undertake a biennial breeding cycle whereas in the Mediterranean follow an annual breeding cycle

74   (Bell et al., 2013). *N. norvegicus* are opportunistic predators and consume a wide variety of prey

species including crustaceans, polychaetes, molluscs and echinoderms (Bell et al., 2013). *N.*

76   *norvegicus* population genetics within the Mediterranean Sea using different methods revealed low

or moderate genetic differentiation between geographical regions (Atlantic vs. Mediterranean) but

78   no geographical pattern of genetic differentiation, thus genetic variability seems to be randomly

distributed among populations (Maltagliati et al., 1998; Passamonti et al., 1997). Nevertheless,

80   different populations were strongly related to spatial and environmental features of the

Mediterranean (Commission et al., 2022).

82   The animal gut microbiome, i.e. the collective genetic material of all microbes found in the

gut, is considered one of the central biological features for the reproduction, development, nutrition,

84   growth, health/immunity of their host (Diwan et al., 2023; Singh et al., 2025) but even in the

conservation of wildlife (Kanika et al., 2025; West et al., 2019). The microbiome concept is so

86   established that the microbiome has been proposed as a means and a target of manipulation for

improving farmed animals production (Luna et al., 2022). Gut microbiomes are also useful at

88   assessing the impact of environmentally induced stress of animals (Evariste et al., 2019) but also as

responders to such disturbances (See et al., 2025). Although decapods have attracted some scientific

90   interest in regards to their microbiome (Foysal, 2023), even to date *N. norvegicus* as a holobiont has

been sporadically studied. The bacterial microbiota of *N. norvegicus* has been investigated both in

natural and experimentally reared populations (Meziti and Kormas, 2013; Meziti et al., 2012; Meziti et al., 2010) while pathogens such as the dinoflagellate *Hematodinium* (Small et al., 2006) have also been investigated. The above render the species' microbiome largely unknown, especially in the framework of the most recently developed and high-throughput sequencing technologies. For these reasons, the current study aimed at (a) investigating the gut bacterial microbiota in three geographically separated *N. norvegicus* populations from the Mediterranean and the North Sea, (b) detecting any potential sex-related microbiota differences and (c) suggesting potential core bacterial taxa for the species populations. It is hypothesized that the animal's gut bacterial communities are highly similar as the animals live in habitats with rather low seasonal variations and very similar environmental conditions and also to the little, if any, genetic differentiation between the populations.

2. MATERIALS AND METHODS

2.1. Sample collection

In the European Union, decapods are not classified as experimental animals under legislation (Directive 2010/63/EU and SFS 2019:66), therefore no animal ethical permits for their collection are required. In Sweden the samples for this study were collected during the ordinary International Bottom Trawl Survey, performed under the EU Data Collection Framework. In Italy the samples for this study were part of the daily landings of standard commercial fishing vessel operating in the Strait of Sicily therefore these samples are subject to IACUC regulations. In Greece the samples were acquired during onboard fishing operations with commercial bottom trawler in the frame of the National Data Collection Framework Program; by the time of capture all samples were already dead. All sampling efforts was limited to minimize large-scale impacts on the populations and is conducted using methodologies to ensure animal welfare and all speciments didn't go through any experimental procedure.

Samples from the North Aegean Sea were collected southwest off Thasos Island (Greece, 40.46ºN 24.6ºE) between 190 and 435 m depth on 31 October 2023 by a bottom otter trawl boat. The Italian samples were collected in the Strait of Sicily (37° 16' N, 13° 02'W, at 350-400 m depth) on 13 September 2023 by a commercial bottom trawler. Samples from the Kattegat, eastern North Sea (56.22ºN 12.16ºE, at 33 m depth), were collected by a research vessel equipped with a bottom trawling fishing net on 28 August 2023. From each sampling site, a total of seven male and seven females were collected for further analysis (Table S1, Figure S1).

2.2. Sample processing and microbiota analysis

For each individual and while on board the fishing vessel, the animals were placed on ice and the abdomen was severed from the cepaholothorax immediately after collection of the animals. The

entire gut tissue was then carefully extracted by holding the abdomen with one hand and gently

128 pulling the telson with the other. The collected gut tissue samples were stored immediately in
DNA/RNA shield (Zymo Research, USA) and stored at -80⁰C after a few days. Gut bacterial

130 communities composition of the gut tissue was determined after bulk DNA was extracted from
approximately 0.25 mg of whole gut tissue (or eggs in the case for specimens from Greece) with the

132 DNeasy PowerSoil Pro Kit (Qiagen, Germany) with no modifications from the suggested protocol.
The V3–V4 regions of the bacterial 16S rRNA genes were amplified from the extracted DNA with

134 the primer pair S-D-Bact-0341-b-S-17 and S-D-Bact-115 0785-a-A-21 (Klindworth et al., 2012).
Sequencing of the amplicons was performed on a MiSeq Illumina instrument (2×300 bp) at the

136 MRDNA Ltd. (Shallowater, TX, USA) sequencing facilities. The raw DNA sequences from this
study have been submitted to the Sequence Read Archive (https://www.ncbi.nlm.nih.gov/sra/) in the

138 BioProject PRJNA1070646 (BioSample SAMN39675726). The standard operating procedure of
MOTHUR software (v.1.48.0) (Schloss et al., 2011; Schloss et al., 2009) was used for processing

140 all the raw 16S rRNA sequence reads. Sequences assigned to mitochondria and chloroplasts, and
single singletons were excluded from further analyses. The operational taxonomic units (OTUs)

142 were determined at 97% cutoff similarity level and were classified with the SILVA database release
138.1 (Quast et al., 2013; Yilmaz et al., 2014). The final OTUs table was normalized to 33,164

144 sequence reads. Rarefaction curves (Figure S2) reached the plateau phase for the number of
sequence reads we used for our analysis. The Nucleotide BLAST (http://blast.ncbi.nlm.nih.gov) tool

146 was used for identifying the closest relatives of the resulting OTUs.

Testing of the differences across all samples were implemented in the PAleontological

148 STudies (PAST) software (Hammer et al., 2001) by applying non-metric multidimensional scaling
(nMDS) based on the unweighted pair group method with the arithmetic mean Bray–Curtis

150 similarity. In addition, Bary-Curtis similarity permutational multivariate analysis of variance
(PERMANOVA) with 9999 permutations, between the three geographical locations and sex of the

152 *N. norvegicus* gut microbiota and eggs, was applied.

154 **3. RESULTS**

The three investigated populations had specimens of different sizes, based on their carapace

156 length and body weight (Table S1). The smallest animals were caught in Italy and the largest in
Greece. Statistically significant differences between females-males, based on PERMANOVA,

158 occurred only for the animals from Sweden (p=0.027, F=19.810). Among animals of the same sex,
the male specimens from Italy differed significantly from those from Sweden (p=0.011, F=125.100)

160 while the females from Italy were different from those from Greece (p=0.015, F=32.600) and
Sweden (p=0.014, F=277.100). The three populations were clearly separated with their male and

female specimens overlapping in terms of their gut microbiota structure (Figure 1). The relative contributions of bacteria at the class-level were also similar between females and males in each of the three populations, with Fusobacteriia and Bacteroidia prevailing in the Swedish specimens, Bacilli and Gammaproteobacteria prevailing in the Italian specimens, and Spirochaetia along with Bacilli prevailing in the female Greece gut samples while the male samples were very diverse. In the eggs samples from the specimens from Greece, the Actinobacteria, Alphaproteobacteria and Gammproteobacteria prevailed (Figure 2).

A total of 2,385 bacterial OTUs were identified across all samples. Females had between 417 (Greece) and 931 (Sweden) OTUs while the respective range for males was between 489 (Italy) and 1290 (Greece), while 180 OTUs were found in eggs from Greece (Table 1). The structure of the eggs bacterial communities was statistically different from both the gut microbiota of both females and males (Table 2). Moreover, there was no overlap between the dominant (relative abundance ≥80%) OTUs in these samples (Table S2). Females had higher number of shared OTUs with the eggs bacterial microbiota (24.9%) compared to males (10.6%) (Figure S3). For each population, no significant differences were observed between females and males, but statistically significant differences occurred for each sex between all populations except between males from Greece and Italy (Table 2).

The populations from Italy and Sweden were highly dominated by the same OTU for both females and males, a Photobacterium-related (OTU-0002) and a *Psychrilyobacter*-related (OTU-0001), respectively (Table 1). The females of the Greek population were dominated (17.8%) by the Spirochaetia-related OTU-0004 while the dominant Cyanobiaceae-related OTU-0018 for males prevailed with 8.4% relative abundance (Table S2).

In each population, a different set of OTUs were found to occur in abundances ≥1% concomitantly in both the females and males, i.e. denoted as "most important OTUs" in the study, in each population (Figure 3, Table 3). Three of these, 17 in total, most important OTUs, namely OTU-0001, -0003 and -0013, were found in all three populations. The population from Sweden had 5/10 most important OTUs occurring exclusively in these samples, while the populations from Italy and Greece had only 2/8 and 1/9, respectively. Regression analysis of the most important OTUs abundances vs. their total body weight, showed two cases of statistically significant negative regressions (OTU-0002, -0005) and five cases of positive regressions (OTU-0004, -0010, -0011, -0013, -0015) (Table 3). There were no "most important" OTUs between the eggs and females or males in the Greek population (Figure S4). In each population, the females:males ratios of abundant OTUs, i.e. relative abundance of ≥1% in each population, which exhibited high (>10) or low (<0.1) values, revealed that various and no overlapping between females and males OTUs contributed differentially to these bacterial communities (Figure 4). Each population had a different set of

OTUs with 0.1<ratio>10 for all OTUs (Figure S5). The population from Italy had the lowest

number of OTUs (8) with females:males ratio >10 and the highest (74) in the population from

Sweden. The population from Sweden showed the lowest numbers of OTUs (14) with ratios <0.1

while the highest number of this ratio occurred in the population from Greece (Figure S5).

4. DISCUSSION

Our study analyzed the gut bacterial communities in three different *Nephrops norvegicus*

populations including both male and female specimens, in order to detect potential differences

between geographic regions and sex. As the exact age/developmental stage of the collected animals

is difficult to safely be determined, we use the total body weight and carapace length as indicative

parameters for growth which can be related to some of the occurred OTUs.

Minor differences were observed between male and female individuals in Sweden and Italy

populations while differences observed between male and female gut samples from Greece in both

Class and OTU level might be associated with females carrying eggs and thus probably consuming

different food sources. Especially for *N. norvegicus* the influence of available feed on gut

microbiota has been investigated and confirmed in previous studies (Meziti et al., 2012; Meziti et

al., 2010).

Overall dominant OTUs detected in this study imply a more algal-related feeding resource

for all populations apart from Italy. More specifically the genus *Psychrilyobacter*, that dominated in

the Swedish population, is a Fusobacteriota bacterium that is very commonly detected in association

with marine animals such as oysters (Fernandez-Piquer et al., 2012), snails (Aronson et al., 2016),

mussels (Santibáñez et al., 2022) and crabs (Zhang et al., 2017) while it also exhibits free-living

lifestyles being a part of the rare biosphere (Liu et al., 2023; Yadav et al., 2021). A recent study

combining cultivation-dependent and independent methods, revealed some functions from an

abalone *Psychrilyobacter* isolate where the specific genus is thriving. Amongst others it was

revealed that the isolate can utilize monosaccharides and disaccharides but not polysaccharides,

implying that it is possibly involved in later fermentation steps but not in the initial food

degradation (Liu et al., 2023). They showed that *Psychrilyobacter* is a versatile fermenter that in

collaboration with other bacteria is very important in digesting the algae consumed by the host.

However, *N. norvegicus* does not feed on algae although remains of plants have been found in its

stomach (Cristo and Cartes, 1998). Previous studies on seasonal changes of the *N. norvegicus* gut

microbiome had suggested that temporal changes in water column leading to increased algal

material concentration in sediment might enhance the presence of alginolytic communities in *N.*

norvegicus gut microbiome (Meziti et al., 2010).

Populations from Greece were characterized by sexual variability leading to the prevalence of a conspicuous Spirochaetaceae representative in females. The genus *Oceanispirochaeta,* observed in this study, contains three fully described species so far, that are obligately anaerobic sediment chemoorganotrophs mainly relying on mono- and disaccharides fermentation (Dubinina et al., 2020; Subhash and Lee, 2017). Although the importance of *Spirochaeta* in the termite gut is well studied (Breznak and Leadbetter, 2006) there is not much information regarding their contribution in marine animals. Recently it has been reported that a spirochaetes associated with a marine sponge mediated terpene production for possible protection of their host against oxidative stress (Waterworth et al., 2024) and has also been speculated to be involved in nitrite metabolism and degradation of complex sugars in the gut of the mudshrimp *Gilvossius tyrrhenus* (formerly *Pestarella tyrrhena*) (Demiri et al., 2009).

The genus *Photobacterium* is commonly detected in *N. norvegicus* and other crustaceans gut microbiome (Foysal, 2023; Jiang et al., 2023; Meziti et al., 2012; Moi et al., 2017). This genus can produce several beneficial compounds such as polyunsaturated fatty acids, lipases, esterases and antimicrobial compounds being also a good candidate for probiotic use (Jiang et al., 2023; Le Doujet et al., 2019). In our study this genus was dominant in the Italian population and important for both Swedish and Italian populations. Its negative correlation with size might imply its prevalence in younger populations. Most importantly, in an older study examining *N. norvegicus* gut microbiota in reared populations receiving different feed, *Photobacterium* was the most abundant genus in mussel fed individuals appearing also as a potential probiotic (Meziti et al., 2012).

Finally, the conspicuous Mycoplasmataceae (ca. Hepatoplasma) representatives were also prevalent in this study similarly to previous studies on *N. norvegicus* as well as on other crustaceans gut microbiome (Meziti et al., 2012). Initially, these Bacteria were detected in the midgut glands (hepatopancreas) of the terrestrial isopod *Porcellio scaber* (Wang et al., 2004) and it has been shown to benefit their hosts under low-nutrient conditions (Fraune and Zimmer, 2008). Today it seems that this bacterial taxon seems to be frequently abundant in various decapods (Foysal, 2023), such as the hepatopancreas of the velvet crab *Necora puber* (Martin et al., 2024), in the gut of deep-sea amphipods (Cheng et al., 2019) juvenile (Sun et al., 2020) and adult individuals (An et al., 2024) of the mitten crab *Eriocher sinensis*, juvenile Caribbean spiny lobsters *Panulirus argus* (Zamora-Briseño et al., 2020), mud crab *Scylla paramamosain* (Jiang et al., 2023) and the vent shrimp *Rimicaris exoculata* (Aubé et al., 2022). Genome sequencing of ca. Hepatoplasma (Collingro et al., 2015) as well as Metagenome Assembled Genomes (MAGs) sequencing (Aubé et al., 2022), showed typical Mycoplasmataceae reduced genome size with the majority of energy producing pathways missing with the exception of glycolysis. Similarly, the majority of nucleotide

and amino acid biosynthesis pathways are not present in the genomes suggesting that ca. Hepatoplasma mainly relies on its host or syntrophic bacteria for its growth (Aubé et al., 2022).

Overall the majority of the dominant and 'important' OTUs in our dataset belonged to groups and genera that had been previously detected in *N. norvegicus* or other Crustaceans gut

microbiome studies (Meziti et al., 2012; Meziti et al., 2010); (Aubé et al., 2022; Jiang et al., 2023). Most importantly, the closest relatives detected, for the majority of the 'important OTUs', were

phylotypes from previous *N. norvegicus* gut microbiome studies mainly performed in Greece more than one decade ago implying the presence of a core gut *N. norvegicus* microbiome regardless of

geographical boundaries. Similar findings were reported for the mud crab *Scylla paramosain,* where ca. Hepatoplasma, *Vibrio, Photobacterium, Carboxylicivirga* were identified as core gut microbiota

genera from different coastal regions in southern China (Jiang et al., 2023).

To date, no known interaction between the *N. norvegicus* sex and its gut microbiota exists.

However, it is known that some vertebrate steroids (e.g. estradiol, progesterone, testosterone) influence and are dependent on the developmental stage and reproduction of the *H. americanus* and

N. norvegicus lobsters; these steroids are most likely species-specific (Burns et al., 1984; Chang, 1997; Fairs et al., 1989). Bacteria seem to have a role in steroid hormone levels in mammals. For

example, the activity of the β-glucuronidase produced by gut bacteria, like *Bacteroides* and *Clostridium,* regulate the levels of active estrogen by breaking glycosidic bonds between glucuronic

acid and estrogen (Cotton et al., 2023). β-glucuronidases are widespread along several bacterial taxa (Lombard et al., 2013; Wardman et al., 2022), some of which are among the most abundant found

in this study (e.g. Clostridia, Sphingomonadaceae, Spirochaetia). Bacteria seem to be also involved in the degradation or the production/reactivation of testosterone while gut microbiota changes can

be accompanied by testosterone levels (Cotton et al., 2023). Other invertebrates, such as cephalopods, have an accessory nidamental gland which is known to harbour specific microbiota

for assisting the defense of their females against pathogens and fouling organisms (Vijayan et al., 2024). In addition, our current dataset cannot provide any potential metabolic features of the

investigated microbial communities as among the most important OTUs several cannot be securely affiliated to any of the known bacterial taxa (see Table S2). These yet-to-be cultivated Bacteria,

which hinder the full characterization of their ecological demands, perhaps refrain us from having a more complete picture for the life cycle of this animal but also for its commercial cultivation.

AKNOWLEDGEMENTS

298 We thank Barbara Bland and Jan-Erik Johansson at the Swedish University of Agricultural Sciences for the collection of the North Sea samples onboard the R/V "SVEA", Danilo Scannella

300 (IRBIM CNR) for the collection of the Strait of Sicily samples and Vasiliki Kousteni (Fisheries

Research Institute, Hellenic Agricultural Organization - Demeter, Kavala, Greece) for the collection

302   of the North Aegean Sea samples.

304   **REFERENCES**

An, H.-E., Malik, A., Lee, J., Mun, M.-H., Lee, K.H., Yoo, H.Y., Kim, C.-B., 2024. Comparison of

306   Bacterial Community Composition in Gut of Chinese Mitten Crabs from Three Distinct

Rivers in Korea. Fishes 9(4), 144. https://doi.org/10.3390/fishes9040144

308   Aronson, H.S., Zellmer, A.J., Goffredi, S.K., 2016. The specific and exclusive microbiome of the

deep-sea bone-eating snail, *Rubyspira osteovora*. FEMS Microbiology Ecology 93(3).

310   https://doi.org/10.1093/femsec/fiw250

Aubé, J., Cambon-Bonavita, M.-A., Velo-Suárez, L., Cueff-Gauchard, V., Lesongeur, F.,

312   Guéganton, M., Durand, L., Reveillaud, J., 2022. A novel and dual digestive symbiosis scales

up the nutrition and immune system of the holobiont Rimicaris exoculata. Microbiome 10(1),

314   189. https://doi.org/10.1186/s40168-022-01380-2

Bell, M., Tuck, I., Dobby, H., 2013. Nephrops Species, in: Lobsters: Biology, Management,

316   Aquaculture and Fisheries. pp. 357-413.

Breznak, J.A., Leadbetter, J.R., 2006. Termite Gut Spirochetes, in: Dworkin, M., Falkow, S.,

318   Rosenberg, E., Schleifer, K.-H., Stackebrandt, E. (Eds.), The Prokaryotes: Volume 7:

Proteobacteria: Delta, Epsilon Subclass. Springer New York, New York, NY, pp. 318-329.

320   Burns, B.G., Sangalang, G.B., Freeman, H.C., McMenemy, M., 1984. Isolation and identification of

testosterone from the serum and testes of the American lobster (Homarus americanus).

322   General and Comparative Endocrinology 54(3), 429-432.

https://doi.org/https://doi.org/10.1016/0016-6480(84)90158-8

324   Chang, E.S., 1997. Chemistry of Crustacean Hormones that Regulate Growth and Reproduction. In:

Fingerman M (ed) Endocrinology and Reproduction Recent Advances in Marine

326   Biotechnology. Taylor & Frnacis Group, p 163-1178.

Cheng, X.-y., Wang, Y., Li, J.-y., Yan, G.-y., He, L.-s., 2019. Comparative analysis of the gut

328   microbial communities between two dominant amphipods from the Challenger Deep, Mariana

Trench. Deep Sea Research Part I: Oceanographic Research Papers 151, 103081.

330   https://doi.org/https://doi.org/10.1016/j.dsr.2019.103081

Collingro, A., Kostanjšek, R., Toenshoff, E.R., Schulz, F., Schuster, L., Domann, D., Horn, M.,

332   2015. Draft Genome Sequence of "*Candidatus* Hepatoplasma crinochetorum" Ps, a Bacterial

Symbiont in the Hepatopancreas of the Terrestrial Isopod *Porcellio scaber*. Genome

334   Announcements 3(4), 10.1128/genomea.00674-00615.

https://doi.org/doi:10.1128/genomea.00674-15

336   Commission, E., European Climate, I., Agency, E.E., Spedicato, M., Cannas, R., Mahé, K.,
        Morales, B., Tsigenopoulos, C., Zane, L., Kavadas, S., Maina, I., Scarcella, G., Sartor, P.,
338   Bandelj, V., Russo, T., Fiorentino, F., 2022. Study on advancing fisheries assessment and
        management advice in the Mediterranean by aligning biological and management units of
340   priority species MED_UNITs – Final report. Publications Office of the European Union.

    Cotton, S., Clayton, C.A., Tropini, C., 2023. Microbial endocrinology: the mechanisms by which
342   the microbiota influences host sex steroids. Trends in Microbiology 31(11), 1131-1142.
        https://doi.org/https://doi.org/10.1016/j.tim.2023.03.010

344   Cristo, M., Cartes, J.E., 1998. A comparative study of the feeding ecology of *Nephrops norvegicus*
        L. (Decapoda: Nephropidae) in the bathyal Mediterranean and the adjacent Atlantic. Scientia
346   Marina 62(S1), 81-90. https://doi.org/10.3989/scimar.1998.62s181

    Demiri, A., Meziti, A., Papaspyrou, S., Thessalou-Legaki, M., Kormas, K.A., 2009. Abdominal
348   setae and midgut bacteria of the mudshrimp *Pestarella tyrrhena*. Central European Journal of
        Biology 4, 558-566. https://doi.org/10.2478/s11535-009-0053-x

350   Diwan, A.D., Harke, S.N., Panche, A.N., 2023. Host-microbiome interaction in fish and shellfish:
        An overview. Fish and Shellfish Immunology Reports 4, 100091.
352   https://doi.org/https://doi.org/10.1016/j.fsirep.2023.100091

    Dubinina, G., Leshcheva, N., Mikheeva, N., Spring, S., Neumann-Schaal, M., Shcherbakova, V.,
354   Grabovich, M., 2020. Description of *Oceanispirochaeta crateris* sp. nov. and reclassification
        of *Spirochaeta perfilievii* as T*hiospirochaeta perfiliev*ii gen. nov., comb. nov. International
356   Journal of Systematic and Evolutionary Microbiology 70(12), 6373-6380.
        https://doi.org/https://doi.org/10.1099/ijsem.0.004544

358   Evariste, L., Barret, M., Mottier, A., Mouchet, F., Gauthier, L., Pinelli, E., 2019. Gut microbiota of
        aquatic organisms: A key endpoint for ecotoxicological studies. Environmental Pollution 248,
360   989-999. https://doi.org/https://doi.org/10.1016/j.envpol.2019.02.101

    Fairs, N.J., Evershed, R.P., Quinlan, P.T., Goad, L.J., 1989. Detection of unconjugated and
362   conjugated steroids in the ovary, eggs, and haemolymph of the decapod crustacean Nephrops
        norvegicus. General and Comparative Endocrinology 74(2), 199-208.
364   https://doi.org/https://doi.org/10.1016/0016-6480(89)90213-X

    Fernandez-Piquer, J., Bowman, J.P., Ross, T., Tamplin, M.L., 2012. Molecular analysis of the
366   bacterial communities in the live Pacific oyster (Crassostrea gigas) and the influence of
        postharvest temperature on its structure. Journal of Applied Microbiology 112(6), 1134-1143.
368   https://doi.org/10.1111/j.1365-2672.2012.05287.x

    Foysal, M.J., 2023. Host habitat shapes the core gut bacteria of decapod crustaceans: A meta-
370   analysis. Heliyon 9(6), e16511. https://doi.org/https://doi.org/10.1016/j.heliyon.2023.e16511

Fraune, S., Zimmer, M., 2008. Host-specificity of environmentally transmitted Mycoplasma-like isopod symbionts. Environmental Microbiology 10(10), 2497-2504. https://doi.org/https://doi.org/10.1111/j.1462-2920.2008.01672.x

Hammer, Ø., Harper, D., Ryan, P., 2001. PAST: Paleontological statistics software package for education and data analysis. Palaeontol Electr 4:9.

Jiang, X., Niu, M., Qin, K., Hu, Y., Li, Y., Che, C., Wang, C., Mu, C., Wang, H., 2023. The shared microbiome in mud crab (*Scylla paramamosain*) of Sanmen Bay, China: core gut microbiome. Frontiers in Microbiology 14. https://doi.org/10.3389/fmicb.2023.1243334

Kanika, N.H., Liaqat, N., Chen, H., Ke, J., Lu, G., Wang, J., Wang, C., 2025. Fish gut microbiome and its application in aquaculture and biological conservation. Frontiers in Microbiology 15. https://doi.org/10.3389/fmicb.2024.1521048

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J.r., Quast, C., Horn, M., Glöckner, F.O., 2012. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. Nucleic Acids Research 41(1), e1. https://doi.org/10.1093/nar/gks808

Le Doujet, T., De Santi, C., Klemetsen, T., Hjerde, E., Willassen, N.-P., Haugen, P., 2019. Closely-related Photobacterium strains comprise the majority of bacteria in the gut of migrating Atlantic cod (Gadus morhua). Microbiome 7(1), 64. https://doi.org/10.1186/s40168-019-0681-y

Liu, M., Wei, G., Lai, Q., Huang, Z., Li, M., Shao, Z., 2023. Genomic and metabolic insights into the first host-associated isolate of *Psychrilyobacter*. Microbiology Spectrum 11(5), e03990-03922. https://doi.org/doi:10.1128/spectrum.03990-22

Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P.M., Henrissat, B., 2013. The carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Research 42(D1), D490-D495. https://doi.org/10.1093/nar/gkt1178

Luna, G.M., Quero, G.M., Kokou, F., Kormas, K., 2022. Time to integrate biotechnological approaches into fish gut microbiome research. Current Opinion in Biotechnology 73, 121-127. https://doi.org/https://doi.org/10.1016/j.copbio.2021.07.018

Maltagliati, F., Camilli, L., Biagi, F., Abbiati, M., 1998. Genetic structure of Norway lobster, Nephrops norvegicus (L.) (Crustacea: Nephropidae), from the Mediterranean Sea. Scientia Marina 62(S1), 91-99. https://doi.org/10.3989/scimar.1998.62s191

Martin, S., Smith, C., Stewart, K., Barr, W., Cheslett, D., O'Connor, I., Swords, F., Ijaz, U.Z., O'Dwyer, K., 2024. The hepatopancreas microbiome of velvet crab, Necora puber. Environmental Microbiology Reports 16(5), e70014. https://doi.org/https://doi.org/10.1111/1758-2229.70014

406   Meziti, A., Kormas, K.A., 2013. Comparison of the Norway lobster (Nephrops norvegicus) gut
          bacterial communities using 16S rDNA clone libraries and pyrosequencing. Anaerobe 23(0),
408   9-11. https://doi.org/http://dx.doi.org/10.1016/j.anaerobe.2013.07.010

       Meziti, A., Mente, E., Kormas, K.A., 2012. Gut bacteria associated with different diets in reared
410   *Nephrops norvegicus*. Systematic and Applied Microbiology 35(7), 473-482.
          https://doi.org/10.1016/j.syapm.2012.07.004

412   Meziti, A., Ramette, A., Mente, E., Kormas, K.A., 2010. Temporal shifts of the Norway lobster
          (Nephrops norvegicus) gut bacterial communities. FEMS Microbiology Ecology 74(2), 472-
414   484. https://doi.org/10.1111/j.1574-6941.2010.00964.x

       Moi, I.M., Roslan, N.N., Leow, A.T.C., Ali, M.S.M., Rahman, R.N.Z.R.A., Rahimpour, A., Sabri,
416   S., 2017. The biology and the importance of Photobacterium species. Applied Microbiology
          and Biotechnology 101(11), 4371-4385. https://doi.org/10.1007/s00253-017-8300-y

418   Passamonti, M., Mantovani, B., Scali, V., Froglia, C., 1997. Allozymic Characterization of Scottish
          and Aegean Populations of Nephrops Norvegicus. Journal of the Marine Biological
420   Association of the United Kingdom 77(3), 727-735.
          https://doi.org/10.1017/S0025315400036158

422   Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
          2013. The SILVA ribosomal RNA gene database project: improved data processing and web-
424   based tools. Nucleic Acids Research 41(D1), D590-D596.
          https://doi.org/10.1093/nar/gks1219

426   Santibáñez, P., Romalde, J., Maldonado, J., Fuentes, D., Figueroa, J., 2022. First characterization of
          the gut microbiome associated with Mytilus chilensis collected at a mussel farm and from a
428   natural environment in Chile. Aquaculture 548, 737644.
          https://doi.org/https://doi.org/10.1016/j.aquaculture.2021.737644

430   Schloss, P.D., Gevers, D., Westcott, S.L., 2011. Reducing the effects of PCR amplification and
          sequencing artifacts on 16S rRNA-based studies. PLoS One 6(12), e27310.

432   Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski,
          R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van
434   Horn, D.J., Weber, C.F., 2009. Introducing mothur: Open-source, platform-independent,
          community-supported software for describing and comparing microbial communities.
436   Applied and Environmental Microbiology 75(23), 7537-7541.
          https://doi.org/10.1128/aem.01541-09

438   See, M.S., Ching, X.L., Khoo, S.C., Abidin, S.Z., Sonne, C., Ma, N.L., 2025. Aquatic microbiomes
          under stress: The role of gut microbiota in detoxification and adaptation to environmental

440   exposures. Journal of Hazardous Materials Advances 17, 100612. https://doi.org/https://doi.org/10.1016/j.hazadv.2025.100612

442 Singh, B.K., Thakur, K., Kumari, H., Mahajan, D., Sharma, D., Sharma, A.K., Kumar, S., Singh, B., Pankaj, P.P., Kumar, R., 2025. A review on comparative analysis of marine and

444   freshwater fish gut microbiomes: insights into environmental impact on gut microbiota. FEMS Microbiology Ecology 101(1), fiae169. https://doi.org/10.1093/femsec/fiae169

446 Small, H.J., Neil, D.M., Taylor, A.C., Atkinson, R.J.A., Coombs, G.H., 2006. Molecular detection of Hematodinium spp. in Norway lobster Nephrops norvegicus and other crustaceans.

448   Diseases of Aquatic Organisms 69(2-3), 185-195.

   Stencel, A., Wloch-Salamon, D., 2022. A pluralistic view of holobionts in the context of process

450   ontology. Frontiers in Microbiology 13. https://doi.org/10.3389/fmicb.2022.911577

   Subhash, Y., Lee, S.-S., 2017. Description of *Oceanispirochaeta sediminicola* gen. nov., sp. nov.,

452   an obligately anaerobic bacterium isolated from coastal marine sediments, and reclassification of *Spirochaeta litoralis* as *Oceanispirochaeta litoralis* comb. nov. International Journal of

454   Systematic and Evolutionary Microbiology 67(9), 3403-3409. https://doi.org/https://doi.org/10.1099/ijsem.0.002130

456 Sun, Y., Han, W., Liu, J., Liu, F., Cheng, Y., 2020. Microbiota comparison in the intestine of juvenile Chinese mitten crab Eriocheir sinensis fed different diets. Aquaculture 515, 734518.

458   https://doi.org/https://doi.org/10.1016/j.aquaculture.2019.734518

   Theis, K.R., Dheilly, N.M., Klassen, J.L., Brucker, R.M., Baines, J.F., Bosch, T.C.G., Cryan, J.F.,

460   Gilbert, S.F., Goodnight, C.J., Lloyd, E.A., Sapp, J., Vandenkoornhuyse, P., Zilber-Rosenberg, I., Rosenberg, E., Bordenstein, S.R., 2016. Getting the Hologenome Concept

462   Right: an Eco-Evolutionary Framework for Hosts and Their Microbiomes. mSystems 1(2), e00028-00016. https://doi.org/10.1128/mSystems.00028-16

464 Troussellier, M., Escalas, A., Bouvier, T., Mouillot, D., 2017. Sustaining rare marine microorganisms: Macroorganisms as repositories and dispersal agents of microbial diversity.

466   Frontiers in Microbiology 8, 947. https://doi.org/10.3389/fmicb.2017.00947

   Vijayan, N., McAnulty, S.J., Sanchez, G., Jolly, J., Ikeda, Y., Nishiguchi, M.K., Réveillac, E.,

468   Gestal, C., Spady, B.L., Li, D.H., Burford, B.P., Kerwin, A.H., Nyholm, S.V., 2024. Evolutionary history influences the microbiomes of a female symbiotic reproductive organ in

470   cephalopods. Applied and Environmental Microbiology 90(3), e00990-00923. https://doi.org/doi:10.1128/aem.00990-23

472 Wang, Y., Stingl, U., Anton-Erxleben, F., Zimmer, M., Brune, A., 2004. 'Candidatus Hepatincola porcellionum' gen. nov., sp. nov., a new, stalk-forming lineage of Rickettsiales colonizing the

474   midgut glands of a terrestrial isopod. Archives of Microbiology 181(4), 299-304.

https://doi.org/10.1007/s00203-004-0655-7

476   Wardman, J.F., Bains, R.K., Rahfeld, P., Withers, S.G., 2022. Carbohydrate-active enzymes

(CAZymes) in the gut microbiome. Nature Reviews Microbiology 20(9), 542-556.

478   https://doi.org/10.1038/s41579-022-00712-1

Waterworth, S.C., Solomons, G.M., Kalinski, J.-C.J., Madonsela, L.S., Parker-Nance, S.,

480   Dorrington, R.A., 2024. The unique and enigmatic spirochete symbiont of latrunculid

sponges. mSphere 9(12), e00845-00824. https://doi.org/doi:10.1128/msphere.00845-24

482   Webster, N.S., 2017. Conceptual and methodological advances for holobiont research.

Environmental Microbiology Reports 9(1), 30-32. https://doi.org/10.1111/1758-2229.12500

484   West, A.G., Waite, D.W., Deines, P., Bourne, D.G., Digby, A., McKenzie, V.J., Taylor, M.W.,

2019. The microbiome in threatened species conservation. Biological Conservation 229, 85-

486   98. https://doi.org/https://doi.org/10.1016/j.biocon.2018.11.016

Yadav, S., Koenen, M., Bale, N., Sinninghe Damsté, J.S., Villanueva, L., 2021. The physiology and

488   metabolic properties of a novel, low-abundance *Psychrilyobacter* species isolated from the

anoxic Black Sea shed light on its ecological role. Environmental Microbiology Reports

490   13(6), 899-910. https://doi.org/https://doi.org/10.1111/1758-2229.13012

Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J.,

492   Ludwig, W., Glöckner, F.O., 2014. The SILVA and "All-species Living Tree Project (LTP)"

taxonomic frameworks. Nucleic Acids Research 42(D1), D643-D648.

494   https://doi.org/10.1093/nar/gkt1209

Zamora-Briseño, J.A., Cerqueda-García, D., Hernández-Velázquez, I.M., Rivera-Bustamante, R.,

496   Huchín-Mian, J.P., Briones-Fourzán, P., Lozano-Álvarez, E., Rodríguez-Canul, R., 2020.

Alterations in the gut-associated microbiota of juvenile Caribbean spiny lobsters Panulirus

498   argus (Latreille, 1804) infected with PaV1. Journal of Invertebrate Pathology 176, 107457.

https://doi.org/https://doi.org/10.1016/j.jip.2020.107457

500   Zhang, N., Song, C., Wang, M., Liu, Y., Hui, M., Cui, Z., 2017. Diversity and characterization of

bacteria associated with the deep-sea hydrothermal vent crab *Austinograea* sp. comparing

502   with those of two shallow-water crabs by 16S ribosomal DNA analysis. PLOS ONE 12(11),

e0187842. https://doi.org/10.1371/journal.pone.0187842

504   Zilber-Rosenberg, I., Rosenberg, E., 2008. Role of microorganisms in the evolution of animals and

plants: the hologenome theory of evolution. FEMS Microbiology Reviews 32(5), 723-735.

506   https://doi.org/10.1111/j.1574-6976.2008.00123.x

508 **Table 1**. Midgut bacterial operational taxonomic units (OTUs) from three *Nephrops norvegicus* populations. Sequence reads=33,164; N=7 for male and

509 female individuals, N=5 for eggs.

		Average carapace length (mm) (coefficient of variation)	Average total body weight (g) (coefficient of variation)	No. of OTUs (average, coefficient of variation)	No. of OTUs with ≥80% relative abundance	Most dominant OTU (% relative abundance)
Greece	Eggs	-	-	180 (86, 4.2%)	17	OTU-0020 (18.5%) *Cutibacterium*
	Females	46.0 (11.6%)	74.0 (30.9%)	417 (128, 24.1%)	11	OTU-0004 (17.8%) *Oceanispirochaeta*
	Males	46.7 (14.6%)	77.0 (42.0%)	1290 (335, 64.2%)	20	OTU-0018 (8.4%) *Cyanobium*
Italy	Females	34.9 (4.5%)	32.4 (10.3%)	434 (136, 24.0%)	7	OTU-0002 (20.2%) *Photobacterium*
	Males	36.9 (4.0%)	36.3 (13.9%)	489 (153, 30,4%)	10	OTU-0002 (20.3%) *Photobacterium*
Sweden	Females	45.0 (0.0%)	64.6 (2.5%)	931 (252, 63.0%)	14	OTU-0001 (24.5%) *Psychrilyobacter*
	Male	45.0 (0.0%)	67.9 (1.6%)	527 (161, 36,7%)	11	OTU-0001 (25.5%) *Psychrilyobacter*

512 **Table 2**. Permutational analysis of variance (PERMANOVA) of eggs, male and female individuals
513 of *Nephrops norvegicus* gut bacterial communities from Greece (GRE), Italy (ITA) and Sweden
514 (SWE). * : $p<0.05$, ** : $p<0.002$, *** : $p<0.01$.

	Eggs-Females	Eggs-Males	Female-Males		Females	Males
GRE	p=0.027*	p=0.046*	p=0.897	GRE-ITA	p=0.032*	p=0.580
	F=3.162	F=2.190	F=1.565		F=2.916	F=1.579
ITA	X	X	p=1.000	ITA-SWE	p=0.027*	p=0.021*
			F=0.616		F=2.981	F=2.668
SWE	X	X	p=0.960	GRE-SWE	p=0.008***	p=0.015*
			F=1.668		F=3.611	F=3.438

518   **Table 3**. Regression of the total body weight vs. the abundance of each of the most important operational taxonomic units (OTU) from three *Nephrops*

519   *norvegicus* populations. N=42; negative and positive regressions are in red and blue letters, respectively; *: $p < 0.05$; G: Greece, I: Italy, S: Sweden.

OTU	Bacterial taxon	Closest (≥98%) relative in NCBI (GenBank accession no.)	p	r
0001 (G/I/S)	*Psychrilyobacter*	Uncultured bacterium clone SS1_B_01_74 (EU050918)	0.165	0.218
0002 (I/S)	*Photobacterium*	*Photobacterium phosphoreum* (AB179540)	0.031*	-0.333
0003 (G/I/S)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092163)	0.219	-0.194
0004 (G/I)	*Oceanispirochaeta*		0.001*	0.487
0005 (G/I)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092262)	0.016*	-0.368
0006 (S)	*Ancyomarina*	Uncultured Bacteroidetes bacterium clone FII-OX025 (JQ579674)	0.131	0.237
0008 (I)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092255)	0.089	-0.266
0009 (I)	*Aliivibrio*	*Aliivibrio wodanis* (LR721750)	0.121	-0.243

0010 (G/S)	*Kordiimonas*	Uncult. Alphaproteobacteria from *N. norvegicus* gut (JN092256)	0.040*	0.319
0011 (G)	*Desulfovibrio*		0.001*	0.503
0012 (S)	Desulfobacteraceae unclass.	Uncult. Desulfobacterales from *N. norvegicus* gut (GQ866071)	0.273	0.173
0013 (G/I/S)	*Carboxylicivirga*	Uncult. Bacteroidia from mud crab *(Scylla paramamosain)* gut (HE610318)	0.033*	0.330
0015 (G)	Bacteria		0.015*	0.373
0018 (G)	Cyanobium	Uncultured bacterium clone 2CE2-5m-91 (GU062169)	0.927	0.015
0021 (S)	*Vibrio*	Uncult. bacterium from *N. norvegicus* gut (JN092213)	0.125	0.240
0024 (S)	*Candidatus* Absconditabacteriales		0.470	0.115
0036 (S)	Flavobacteriaceae unclass.	Uncultured bacterium clone EzyYyy31 (KX172210)	0.464	0.116

523
524

Figure 1. Non-metric multidimensional scaling (nMDS) of male (♂) and female (♀) individuals of
Nephrops norvegicus gut from Greece (GRE), Italy (ITA) and Sweden (SWE).

Figure 2. Relative abundance of class-level operational taxonomic units in the female (F) and male (M) *Nephrops norvegicus* gut from Greece (GR), Italy (IT) and Sweden (SW).

533
534

535
536

537
538

Figure 3. Most important (relative abundance ≥1% in both male and female individuals) bacterial operational taxonomic units (OTUs) of the *Nephrops norvegicus* gut from Greece (GRE), Italy (ITA) and Sweden (SWE). Red dots: bacterial operational taxonomic units (OTUs); blue and pink dots: important OTUs, i.e. ≥1% relative abundance in males and females only, respectively.

Figure 4. Ratio of females:males bacterial operational taxonomic units (OTUs) with ≥1% relative abundance in the *Nephrops norvegicus* gut from three populations. Only closest relatives of OTUs with 0.1≤ratio≥10 is shown. Blue bars indicate OTUs occurrence only in males.