Sex-related gut microbiota in three geographically separated Norway lobster (*Nephrops norvegicus*) populations

Polina Rusanova[*,†], Eleni Nikouli[$], Michele Casini[†,‡], Gioacchino Bono[*,#], Elena Mente[¶], Alexandra Meziti[∥], Konstantinos Kormas[$*]

[*] Institute for Biological Resources and Marine Biotechnologies, National Research Council (IRBIM-CNR), Mazara del Vallo, Italy

[†] Department of Biological, Geological and Environmental Sciences (BiGeA), Laboratory of Marine Biology and Fisheries, University of Bologna, Bologna, Italy

[$] Department of Ichthyology and Aquatic Environment, School of Agricultural Sciences, 384 46 Volos, Greece

[‡] Department of Aquatic Resources, Institute of Marine Research, Swedish University of Agricultural Sciences, Lysekil, Sweden

[#] Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STEBICEF), Palermo University, Palermo, Italy

[¶] Laboratory of Ichthyology-Culture and Pathology of Aquatic Animals, School of Veterinary Medicine, Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece

[∥] Department of Marine Sciences, University of the Aegean, University Hill, 81100, Mytilene, Greece

[1] Corresponding author: kkormas@uth.gr

ABSTRACT

Despite the ecological and economic interest of the Norway lobster (*Nephrops norvegicus*), its gut microbiota remains largely understudied. The current study aimed at investigating the gut

bacterial microbiota in three geographically separated *N. norvegicus* populations from the Mediterranean and the North Sea and detect any potential sex-related microbiota differences, by

high-throughput sequencing of the V3-V4 16S rRNA gene diversity of the gut tissue. From the Greek population, egg-bearing females were also caught. A total of 2,385 operational taxonomic

units (OTUs) were identified and between 417 and 1290 occurred in each population/sex group. The dominant OTUs belonged to the Fusobacteriia and Bacteroidia (Sweden), Bacilli and

Gammaproteobacteria (Italy) and Spirochaetia and Bacilli (Greece) classes. In the eggs, the Actinobacteria, Alphaproteobacteria and Gammproteobacteria prevailed. Four OTUs related to the

Oceanispirochaeta, *Kordiimonas, Desulfovibrio, Carboxylicivirga* genera and one unafilliated OTU were positively correlated (p values between 0.001 and 0.04) to body size indicating their potential

role in the animal's nutrition and growth. No statistically significant differences were found between males-females in any of the populations. However, statistically significant differences

between populations for each sex, were found for all females (p values between 0.008 and 0.032) and for the males between the most distant populations, i.e. Italy and Sweden (p=0.021) and,

Greece and Sweden (p=0.015). The eggs microbiota was statistically different from both the adult females (p=0.027) and males (p=0.046) gut microbiota. Overall, this study showed that *N.*

norvegicus gut microbiota vary for each sex between geographically distant populations.

Keywords: crustacea, microbiome, fisheries, North Sea, Mediterranean Sea, Aegean Sea

INTRODUCTION

The concepts of hologenome and the holobiont (Zilber-Rosenberg and Rosenberg, 2008) has opened a new way we view animal life (Theis et al., 2016; Webster, 2017) for their ontogeny and function (Troussellier et al., 2017; Stencel and Wloch-Salamon, 2022). Thus, during the last few years, the scientific interest on holobiont research has boomed for animals, as well. However, there are yet animal species from various habitats whose microbiomes remain largely unknow, despite their ecological and/or economic importance and the ongoing rapid technological advancements in DNA/RNA technologies.

One such species is the Norway lobster *Nephrops norvegicus* (Linnaeus, 1758) a decapod crustacean, langoustine or scampi, which is a typical clawed lobster with a slender body, long claws and large dark eyes. *Nephrops norvegicus* is considered one of the most important species for fisheries, valuable commercial crustacean species in Europe. It is a benthic crustacean dependent on muddy-type sediments crustacean suitable for the construction of burrows. It is widely distributed across the continental shelves of the northeast Atlantic Ocean and the Mediterranean Sea, from Iceland and Norway in the north, to Morocco and the Adriatic seas in the south (Bell et al., 2013). Populations dwelling in colder waters around Iceland and the Faroe Islands undertake a biennial breeding cycle whereas in the Mediterranean follow an annual breeding cycle (Bell et al., 2013). *N. norvegicus* are opportunistic predators and consume a wide variety of prey species including crustaceans, polychaetes, molluscs and echinoderms (Bell et al., 2013). *N. norvegicus* population genetics within the Mediterranean Sea using different methods revealed low or moderate genetic differentiation between geographical regions (Atlantic vs. Mediterranean) but no geographical pattern of genetic differentiation, thus genetic variability seems to be randomly distributed among populations (Passamonti et al., 1997; Maltagliati et al., 1998). Nevertheless, different populations were strongly related to spatial and environmental features of the Mediterranean (Commission et al., 2022).

The animal gut microbiome, i.e. the collective genetic material of all microbes, is considered one of the central biological features for the reproduction, development, nutrition, growth, health/immunity of their host (Diwan et al., 2023; Singh et al., 2025) but even in the conservation (West et al., 2019; Kanika et al., 2025). The microbiome concept is so established that the microbiome has been proposed as a means and target of manipulation for improving farmed animals production (Luna et al., 2022). Gut microbiomes are also useful at assessing the impact of environmentally induced stress of animals (Evariste et al., 2019) but also as responders to such disturbances (See et al., 2025). Although decapods have attracted some scientific interest in regards to their microbiome (Foysal, 2023), to date *N. norvegicus* as a holobiont has been sporadically

studied. The bacterial microbiota of *N. norvegicus* has been investigated both in natural and

experimentally reared populations (Meziti et al., 2010; Meziti et al., 2012; Meziti and Kormas, 2013) while pathogens such as the dinoflagellate *Hematodinium* (Small et al., 2006) have also been investigated. The above render the species' microbiome largely unknown, especially in the framework of the most recently developed and high-throughput sequencing technologies. For these reasons, the current study aimed at (a) investigating the gut bacterial microbiota in three geographically separated *N. norvegicus* populations from the Mediterranean and the North Sea, (b) detecting any potential sex-related microbiota differences and (c) suggesting potential core bacterial taxa for the species populations. It is hypothesized that the animal's gut bacterial communities are highly similar as the animals live in habitats with rather low seasonal variations and very similar environmental conditions and also to the little, if any, genetic differentiation between the populations.

MATERIALS AND METHODS

Sample collection. Samples from the North Aegean Sea were collected southwest off Thasos Island (Greece, 40.46ºN 24.6ºE) between 190 and 435 m depth on 31 October 2023 by a bottom otter trawl boat. The Italian samples were collected in the Strait of Sicily (37° 16' N, 13° 02'W, at 350-400 m depth) on 13 September 2023 by a commercial bottom trawler. Samples from the Kattegat, eastern North Sea (56.22ºN 12.16ºE, at 33 m depth), were collected by a research vessel equipped with a bottom trawling fishing net on 28 August 2023. From each sampling site, a total of seven male and seven females were collected for further analysis (Table S1).

Sample processing and microbiota analysis. For each individual and while on board the fishing vessel, the animals were placed on ice and the abdomen was severed from the cepaholothorax immediately after collection of the animals. The entire gut tissue was then carefully extracted by holding the abdomen with one hand and gently pulling the telson with the other. The collected gut tissue samples were stored immediately in DNA/RNA shield (Zymo Research, USA) and stored at -80^0C after a few days. Gut bacterial communities composition of the gut tissue was determined after bulk DNA was extracted from approximately 0.25 mg of whole gut tissue (or eggs in the case for specimens from Greece) with the DNeasy PowerSoil Pro Kit (Qiagen, Germany) with no modifications from the suggested protocol. The V3–V4 regions of the bacterial 16S rRNA genes were amplified from the extracted DNA with the primer pair S-D-Bact-0341-b-S-17 and S-D-Bact-115 0785-a-A-21 (Klindworth et al., 2012). Sequencing of the amplicons was performed on a MiSeq Illumina instrument (2×300 bp) at the MRDNA Ltd. (Shallowater, TX, USA) sequencing

128 facilities. The raw DNA sequences from this study have been submitted to the Sequence Read

Archive (https://www.ncbi.nlm.nih.gov/sra/) in the BioProject PRJNA1070646 (BioSample

130 SAMN39675726). The standard operating procedure of MOTHUR software (v.1.48.0) (Schloss et

al., 2009; Schloss et al., 2011) was used for processing all the raw 16S rRNA sequence reads.

132 Sequences assigned to mitochondria and chloroplasts, and single singletons were excluded from

further analyses. The operational taxonomic units (OTUs) were determined at 97% cutoff similarity

134 level and were classified with the SILVA database release 138.1 (Quast et al., 2013; Yilmaz et al.,

2014). The final OTUs table was normalized to 33,164 sequence reads. Rarefaction curves (Fig. S2)

136 reached the plateau phase for the number of sequence reads we used for our analysis. The

Nucleotide BLAST (http://blast.ncbi.nlm.nih.gov) tool was used for identifying the closest relatives

138 of the resulting OTUs.

Testing of the differences across all samples were implemented in the PAleontological

140 STudies (PAST) software (Hammer et al., 2001) by applying non-metric multidimensional scaling

(nMDS) based on the unweighted pair group method with the arithmetic mean Bray–Curtis

142 similarity. In addition, Bary-Curtis similarity permutational multivariate analysis of variance

(PERMANOVA) with 9999 permutations, between the three geographical locations and sex of the

144 *N. norvegicus* gut microbiota and eggs, was applied.

RESULTS

The three investigated populations had specimens of different sizes, based on their carapace

150 length and body weight (Table S1). The smallest animals were caught in Italy and the largest in

Greece. Statistically significant differences between females-males, based on PERMANOVA,

152 occurred only for the animals from Sweden (p=0.027, F=19.810). Among animals of the same sex,

the male specimens from Italy differed significantly from those from Sweden (p=0.011, F=125.100)

154 while the females from Italy were different from those from Greece (p=0.015, F=32.600) and

Sweden (p=0.014, F=277.100). The relative contributions of bacteria at the class-level were also

156 similar between females and males in each of the three populations, with Fusobacteriia and

Bacteroidia prevailing in the Swedish specimens, Bacilli and Gammaproteobacteria prevailing in

158 the Italian specimens, and Spirochaetia along with Bacilli prevailing in the female Greece gut

samples while the male samples were very diverse. In the eggs samples from the specimens from

160 Greece, the Actinobacteria, Alphaproteobacteria and Gammproteobacteria prevailed (Fig. 2).

A total of 2,385 bacterial OTUs were identified across all samples. Females had between

162 417 (Greece) and 931 (Sweden) OTUs while the respective range for males was between 489 (Italy)

and 1290 (Greece), while 180 OTUs were found in eggs from Greece (Tab. 1). The structure of the eggs bacterial communities was statistically different from both the gut microbiota of both females and males (Table 2). Moreover, there was no overlap between the dominant (relative abundance ≥80%) OTUs in these samples (Tab. S2). Females had higher number of shared OTUs with the eggs bacterial microbiota (24.9%) compared to males (10.6%) (Fig. S3). For each population, no significant differences were observed between females and males, but statistically significant differences occurred for each sex between all populations except between males from Greece and Italy (Table 2).

The populations from Italy and Sweden were highly dominated by the same OTU for both females and males, a Photobacterium-related (OTU-0002) and a *Psychrilyobacter*-related (OTU-0001), respectively (Tab. 1). The females of the Greek population were dominated (17.8%) by the Spirochaetia-related OTU-0004 while the dominant Cyanobiaceae-related OTU-0018 for males prevailed with 8.4% relative abundance (Table S2).

In each population, a different set of OTUs were found to occur in abundances ≥1% concomitantly in both the females and males, i.e. denoted as "most important OTUs" in the study, in each population (Fig. 3, Tab. 3). Three of these, 17 in total, most important OTUs, namely OTU-0001, -0003 and -0013, were found in all three populations. The population from Sweden had 5/10 most important OTUs occurring exclusively in these samples, while the populations from Italy and Greece had only 2/8 and 1/9, respectively. Regression analysis of the most important OTUs abundances vs. their total body weight, showed two cases of statistically significant negative regressions (OTU-0002, -0005) and five cases of positive regressions (OTU-0004, -0010, -0011, -0013, -0015) (Table 2). There were no "most important" OTUs between the eggs and females or males in the Greek population (Fig. S4). In each population, the females:males ratios of abundant OTUs, i.e. relative abundance of ≥1% in each population, which exhibited high (>10) or low (<0.1) values, revealed that various and no overlapping between females and males OTUs contributed differentially to these bacterial communities (Fig. 4). Each population had a different set of OTUs with 0.1<ratio>10 for all OTUs (Fig. S5). The population from Italy had the lowest number of OTUs (8) with females:males ratio >10 and the highest (74) in the population from Sweden. The population from Sweden showed the lowest numbers of OTUs (14) with ratios <0.1 while the highest number of this ratio occurred in the population from Greece (Fig. S5).

DISCUSSION

Our study analyzed the gut bacterial communities in three different *Nephrops norvegicus*
populations including both male and female specimens, in order to detect potential differences
between geographic regions and sex. As the exact age/developmental stage of the collected animals
is difficult to safely be determined, we use the total body weight and carapace length as indicative
parameters for growth which can be related to some of the occurred OTUs.

Minor differences were observed between male and female individuals in Sweden and Italy
populations while differences observed between male and female gut samples from Greece in both
Class and OTU level might be associated with females carrying eggs and thus probably consuming
different food sources. Especially for *N. norvegicus* the influence of available feed on gut
microbiota has been investigated and confirmed in previous studies (Meziti et al., 2010; Meziti et
al., 2012).

Overall dominant OTUs detected in this study imply a more algal-related feeding for all
populations apart from Italy. More specifically the genus *Psychrilyobacter*, that dominated in the
Swedish population, is a Fusobacteriota bacterium that is very commonly detected in association
with marine animals such as oysters (Fernandez-Piquer et al., 2012), snails (Aronson et al., 2016),
mussels (Santibáñez et al., 2022) and crabs (Zhang et al., 2017) while it also exhibits free-living
lifestyles being a part of the rare biosphere (Yadav et al., 2021; Liu et al., 2023). A recent study
combining cultivation-dependent and independent methods, revealed some functions from an
abalone *Psychrilyobacter* isolate where the specific genus is thriving. Amongst others it was
revealed that the isolate can utilize monosaccharides and disaccharides but not polysaccharides,
implying that it is possibly involved in later fermentation steps but not in the initial food
degradation (Liu et al., 2023). They showed that *Psychrilyobacter* is a versatile fermenter that in
collaboration with other bacteria is very important in digesting the algae consumed by the host.
However, *N. norvegicus* does not feed on algae although remains of plants have been found in its
stomach (Cristo and Cartes, 1998). Previous studies on seasonal changes of the *N. norvegicus* gut
microbiome had suggested that temporal changes in water column leading to increased algal
material concentration in sediment might enhance the presence of alginolytic communities in *N.
norvegicus* gut microbiome (Meziti et al., 2010).

Populations from Greece were characterized by sexual variability leading to the prevalence
of a conspicuous Spirochaetaceae representative in females. The genus *Oceanispirochaeta,*
observed in this study, contains three fully described species so far, that are obligately anaerobic
sediment chemoorganotrophs mainly relying on mono- and disaccharides fermentation (Subhash
and Lee, 2017; Dubinina et al., 2020). Although the importance of *Spirochaeta* in the termite gut is
well studied (Breznak and Leadbetter, 2006) there is not much information regarding their
contribution in marine animals. Recently it has been reported that a spirochaetes associated with a

marine sponge mediated terpene production for possible protection of their host against oxidative stress (Waterworth et al., 2024) and has also been speculated to be involved in nitrite metabolism

and degradation of complex sugars in the gut of the mudshrimp *Gilvossius tyrrhenus* (formerly *Pestarella tyrrhena*) (Demiri et al., 2009).

The genus *Photobacterium* is commonly detected in *N. norvegicus* and other crustaceans gut microbiome (Meziti et al., 2012; Moi et al., 2017; Foysal, 2023; Jiang et al., 2023). This genus can

produce several beneficial compounds such as polyunsaturated fatty acids, lipases, esterases and antimicrobial compounds being also a good candidate for probiotic use (Le Doujet et al., 2019;

Jiang et al., 2023). In our study this genus was dominant in the Italian population and important for both Swedish and Italian populations. Its negative correlation with size might imply its prevalence

in younger populations. Most importantly, in an older study examining *N. norvegicus* gut microbiota in reared populations receiving different feed, *Photobacterium* was the most abundant

genus in mussel fed individuals appearing also as a potential probiotic (Meziti et al., 2012).

       Finally, the conspicuous Mycoplasmataceae (ca. Hepatoplasma) representatives were also

prevalent in this study similarly to previous studies on *N. norvegicus* as well as on other crustaceans gut microbiome (Meziti et al., 2012). Initially, these Bacteria were detected in the midgut glands

(hepatopancreas) of the terrestrial isopod *Porcellio scaber* (Wang et al., 2004) and it has been shown to benefit their hosts under low-nutrient conditions (Fraune and Zimmer, 2008). Today it

seems that this bacterial taxon seems to be frequently abundant in various decapods (Foysal, 2023), such as the hepatopancreas of the velvet crab *Necora puber* (Martin et al., 2024), in the gut of deep-

sea amphipods (Cheng et al., 2019) juvenile (Sun et al., 2020) and adult individuals (An et al., 2024) of the mitten crab *Eriocher sinensis*, juvenile Caribbean spiny lobsters *Panulirus argus*

(Zamora-Briseño et al., 2020), mud crab *Scylla paramamosain* (Jiang et al., 2023) and the vent shrimp *Rimicaris exoculata* (Aubé et al., 2022). Genome sequencing of ca. Hepatoplasma

(Collingro et al., 2015) as well as Metagenome Assembled Genomes (MAGs) sequencing (Aubé et al., 2022), showed typical Mycoplasmataceae reduced genome size with the majority of energy

producing pathways missing with the exception of glycolysis. Similarly, the majority of nucleotide and amino acid biosynthesis pathways are not present in the genomes suggesting that ca.

Hepatoplasma mainly relies on its host or syntrophic bacteria for its growth (Aubé et al., 2022).

       Overall the majority of the dominant and 'important' OTUs in our dataset belonged to

262   groups and genera that had been previously detected in *N. norvegicus* or other Crustaceans gut microbiome studies (Meziti et al., 2010; Meziti et al., 2012); (Aubé et al., 2022; Jiang et al., 2023).

264   Most importantly, the closest relatives detected, for the majority of the 'important OTUs', were phylotypes from previous *N. norvegicus* gut microbiome studies mainly performed in Greece more

266   than one decade ago implying the presence of a core gut *N. norvegicus* microbiome regardless of

geographical boundaries. Similar findings were reported for the mud crab *Scylla paramosain,* where ca. Hepatoplasma, *Vibrio, Photobacterium, Carboxylicivirga* were identified as core gut microbiota genera from different coastal regions in southern China (Jiang et al., 2023).

To date, no known interaction between the *N. norvegicus* sex and its gut microbiota exists. However, it is known that some vertebrate steroids (e.g. estradiol, progesterone, testosterone) influence and are dependent on the developmental stage and reproduction of the *H. americanus* and *N. norvegicus* lobsters; these steroids are most likely species-specific (Burns et al., 1984; Fairs et al., 1989; Chang, 1997). Bacteria seem to have a role in steroid hormone levels in mammals. For example, the activity of the β-glucuronidase produced by gut bacteria, like *Bacteroides* and *Clostridium,* regulate the levels of active estrogen by breaking glycosidic bonds between glucuronic acid and estrogen (Cotton et al., 2023). β-glucuronidases are widespread along several bacterial taxa (Lombard et al., 2013; Wardman et al., 2022), some of which are among the most abundant found in this study (e.g. Clostridia, Sphingomonadaceae, Spirochaetia). Bacteria seem to be also involved in the degradation or the production/reactivation of testosterone while gut microbiota changes can be accompanied by testosterone levels (Cotton et al., 2023). Other invertebrates, such as cephalopods, have an accessory nidamental gland which is known to harbour specific microbiota for assisting the defense of their females against pathogens and fouling organisms (Vijayan et al., 2024). In addition, our current dataset cannot provide any potential metabolic features of the investigated microbial communities as among the most important OTUs several cannot be securely affiliated to any of the known bacterial taxa (see Table S2). These yet-to-be cultivated Bacteria, which hinder the full characterization of their ecological demands, perhaps refrain us from having a more complete picture for the life cycle of this animal but also for its commercial cultivation.

DISCLOSURES

The authors declare no conflict of interest.

AKNOWLEDGEMENTS

We thank Barbara Bland and Jan-Erik Johansson at the Swedish University of Agricultural Sciences for the collection of the North Sea samples onboard the R/V "SVEA", Danilo Scannella (IRBIM CNR) for the collection of the Strait of Sicily samples and Vasiliki Kousteni (Fisheries

302 Research Institute, Hellenic Agricultural Organization - Demeter, Kavala, Greece) for the collection

of the North Aegean Sea samples.

306 **REFERENCES**

308 An, H.-E., A. Malik, J. Lee, M.-H. Mun, K. H. Lee, H. Y. Yoo, and C.-B. Kim. 2024. Comparison
of Bacterial Community Composition in Gut of Chinese Mitten Crabs from Three Distinct
310 Rivers in Korea. Fishes 9(4):144.

Aronson, H. S., A. J. Zellmer, and S. K. Goffredi. 2016. The specific and exclusive microbiome of
312 the deep-sea bone-eating snail, *Rubyspira osteovora*. FEMS Microbiology Ecology
93(3)doi: 10.1093/femsec/fiw250

314 Aubé, J., M.-A. Cambon-Bonavita, L. Velo-Suárez, V. Cueff-Gauchard, F. Lesongeur, M.
Guéganton, L. Durand, and J. Reveillaud. 2022. A novel and dual digestive symbiosis scales
316 up the nutrition and immune system of the holobiont Rimicaris exoculata. Microbiome
10(1):189. doi: 10.1186/s40168-022-01380-2

318 Bell, M., I. Tuck, and H. Dobby. 2013. Nephrops Species, Lobsters: Biology, Management,
Aquaculture and Fisheries. p. 357-413.

320 Breznak, J. A., and J. R. Leadbetter. 2006. Termite Gut Spirochetes. In: M. Dworkin, S. Falkow, E.
Rosenberg, K.-H. Schleifer and E. Stackebrandt, editors, The Prokaryotes: Volume 7:
322 Proteobacteria: Delta, Epsilon Subclass. Springer New York, New York, NY. p. 318-329.

Burns, B. G., G. B. Sangalang, H. C. Freeman, and M. McMenemy. 1984. Isolation and
324 identification of testosterone from the serum and testes of the American lobster (Homarus
americanus). General and Comparative Endocrinology 54(3):429-432. doi:
326 https://doi.org/10.1016/0016-6480(84)90158-8

Chang, E. S. 1997. Chemistry of Crustacean Hormones that Regulate Growth and Reproduction. In:
328 Fingerman M (ed) Endocrinology and Reproduction Recent Advances in Marine
Biotechnology. Taylor & Frnacis Group:p 163-1178.

330 Cheng, X.-y., Y. Wang, J.-y. Li, G.-y. Yan, and L.-s. He. 2019. Comparative analysis of the gut
microbial communities between two dominant amphipods from the Challenger Deep,
332 Mariana Trench. Deep Sea Research Part I: Oceanographic Research Papers 151:103081.
doi: https://doi.org/10.1016/j.dsr.2019.103081

334 Collingro, A., R. Kostanjšek, E. R. Toenshoff, F. Schulz, L. Schuster, D. Domann, and M. Horn.
2015. Draft Genome Sequence of "<i>Candidatus</i> Hepatoplasma crinochetorum" Ps, a
336 Bacterial Symbiont in the Hepatopancreas of the Terrestrial Isopod <i>Porcellio scaber</i>.
Genome Announcements 3(4):10.1128/genomea.00674-00615. doi:
338 doi:10.1128/genomea.00674-15

Commission, E., I. European Climate, E. E. Agency, M. Spedicato, R. Cannas, K. Mahé, B.
340 Morales, C. Tsigenopoulos, L. Zane, S. Kavadas, I. Maina, G. Scarcella, P. Sartor, V.
Bandelj, T. Russo, and F. Fiorentino. 2022. Study on advancing fisheries assessment and
342 management advice in the Mediterranean by aligning biological and management units of
priority species MED_UNITs – Final report. Publications Office of the European Union.

344 Cotton, S., C. A. Clayton, and C. Tropini. 2023. Microbial endocrinology: the mechanisms by
which the microbiota influences host sex steroids. Trends in Microbiology 31(11):1131-
346 1142. doi: https://doi.org/10.1016/j.tim.2023.03.010

Cristo, M., and J. E. Cartes. 1998. A comparative study of the feeding ecology of *Nephrops*
348 *norvegicus* L. (Decapoda: Nephropidae) in the bathyal Mediterranean and the adjacent
Atlantic. Scientia Marina 62(S1):81-90. doi: 10.3989/scimar.1998.62s181

350 Demiri, A., A. Meziti, S. Papaspyrou, M. Thessalou-Legaki, and K. A. Kormas. 2009. Abdominal
          setae and midgut bacteria of the mudshrimp *Pestarella tyrrhena*. Central European Journal
352   of Biology 4:558-566.

Diwan, A. D., S. N. Harke, and A. N. Panche. 2023. Host-microbiome interaction in fish and
354   shellfish: An overview. Fish and Shellfish Immunology Reports 4:100091. doi:
          https://doi.org/10.1016/j.fsirep.2023.100091

356 Dubinina, G., N. Leshcheva, N. Mikheeva, S. Spring, M. Neumann-Schaal, V. Shcherbakova, and
          M. Grabovich. 2020. Description of *Oceanispirochaeta crateris* sp. nov. and reclassification
358   of *Spirochaeta perfilievii* as T*hiospirochaeta perfiliev*ii gen. nov., comb. nov. International
          Journal of Systematic and Evolutionary Microbiology 70(12):6373-6380. doi:
360   https://doi.org/10.1099/ijsem.0.004544

Evariste, L., M. Barret, A. Mottier, F. Mouchet, L. Gauthier, and E. Pinelli. 2019. Gut microbiota of
362   aquatic organisms: A key endpoint for ecotoxicological studies. Environmental Pollution
          248:989-999. doi: https://doi.org/10.1016/j.envpol.2019.02.101

364 Fairs, N. J., R. P. Evershed, P. T. Quinlan, and L. J. Goad. 1989. Detection of unconjugated and
          conjugated steroids in the ovary, eggs, and haemolymph of the decapod crustacean
366   Nephrops norvegicus. General and Comparative Endocrinology 74(2):199-208. doi:
          https://doi.org/10.1016/0016-6480(89)90213-X

368 Fernandez-Piquer, J., J. P. Bowman, T. Ross, and M. L. Tamplin. 2012. Molecular analysis of the
          bacterial communities in the live Pacific oyster (Crassostrea gigas) and the influence of
370   postharvest temperature on its structure. Journal of Applied Microbiology 112(6):1134-
          1143. doi: 10.1111/j.1365-2672.2012.05287.x

372 Foysal, M. J. 2023. Host habitat shapes the core gut bacteria of decapod crustaceans: A meta-
          analysis. Heliyon 9(6):e16511. doi: https://doi.org/10.1016/j.heliyon.2023.e16511

374 Fraune, S., and M. Zimmer. 2008. Host-specificity of environmentally transmitted Mycoplasma-like
          isopod symbionts. Environmental Microbiology 10(10):2497-2504. doi:
376   https://doi.org/10.1111/j.1462-2920.2008.01672.x

Hammer, Ø., D. Harper, and P. Ryan. 2001. PAST: Paleontological statistics software package for
378   education and data analysis. Palaeontol Electr 4:9

Jiang, X., M. Niu, K. Qin, Y. Hu, Y. Li, C. Che, C. Wang, C. Mu, and H. Wang. 2023. The shared
380   microbiome in mud crab (*Scylla paramamosain*) of Sanmen Bay, China: core gut
          microbiome. Frontiers in Microbiology 14(Original Research) doi:
382   10.3389/fmicb.2023.1243334

Kanika, N. H., N. Liaqat, H. Chen, J. Ke, G. Lu, J. Wang, and C. Wang. 2025. Fish gut microbiome
384   and its application in aquaculture and biological conservation. Frontiers in Microbiology
          15(Mini Review) doi: 10.3389/fmicb.2024.1521048

386 Klindworth, A., E. Pruesse, T. Schweer, J. r. Peplies, C. Quast, M. Horn, and F. O. Glöckner. 2012.
          Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
388   generation sequencing-based diversity studies. Nucleic Acids Research 41(1):e1. doi:
          10.1093/nar/gks808

390 Le Doujet, T., C. De Santi, T. Klemetsen, E. Hjerde, N.-P. Willassen, and P. Haugen. 2019.
          Closely-related Photobacterium strains comprise the majority of bacteria in the gut of
392   migrating Atlantic cod (Gadus morhua). Microbiome 7(1):64. doi: 10.1186/s40168-019-
          0681-y

394 Liu, M., G. Wei, Q. Lai, Z. Huang, M. Li, and Z. Shao. 2023. Genomic and metabolic insights into
          the first host-associated isolate of *Psychrilyobacter*. Microbiology Spectrum 11(5):e03990-
396   03922. doi: doi:10.1128/spectrum.03990-22

Lombard, V., H. Golaconda Ramulu, E. Drula, P. M. Coutinho, and B. Henrissat. 2013. The
398   carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Research
          42(D1):D490-D495. doi: 10.1093/nar/gkt1178

400   Luna, G. M., G. M. Quero, F. Kokou, and K. Kormas. 2022. Time to integrate biotechnological approaches into fish gut microbiome research. Current Opinion in Biotechnology 73:121-
402   127. doi: https://doi.org/10.1016/j.copbio.2021.07.018

    Maltagliati, F., L. Camilli, F. Biagi, and M. Abbiati. 1998. Genetic structure of Norway lobster,
404   Nephrops norvegicus (L.) (Crustacea: Nephropidae), from the Mediterranean Sea. Scientia Marina 62(S1):91-99. doi: 10.3989/scimar.1998.62s191

406   Martin, S., C. Smith, K. Stewart, W. Barr, D. Cheslett, I. O'Connor, F. Swords, U. Z. Ijaz, and K. O'Dwyer. 2024. The hepatopancreas microbiome of velvet crab, Necora puber.
408   Environmental Microbiology Reports 16(5):e70014. doi: https://doi.org/10.1111/1758-2229.70014

410   Meziti, A., and K. A. Kormas. 2013. Comparison of the Norway lobster (Nephrops norvegicus) gut bacterial communities using 16S rDNA clone libraries and pyrosequencing. Anaerobe
412   23(0):9-11. doi: http://dx.doi.org/10.1016/j.anaerobe.2013.07.010

    Meziti, A., E. Mente, and K. A. Kormas. 2012. Gut bacteria associated with different diets in reared
414   *Nephrops norvegicus*. Systematic and Applied Microbiology 35(7):473-482. doi: 10.1016/j.syapm.2012.07.004

416   Meziti, A., A. Ramette, E. Mente, and K. A. Kormas. 2010. Temporal shifts of the Norway lobster (Nephrops norvegicus) gut bacterial communities. FEMS Microbiology Ecology 74(2):472-
418   484. doi: 10.1111/j.1574-6941.2010.00964.x

    Moi, I. M., N. N. Roslan, A. T. C. Leow, M. S. M. Ali, R. N. Z. R. A. Rahman, A. Rahimpour, and
420   S. Sabri. 2017. The biology and the importance of Photobacterium species. Applied Microbiology and Biotechnology 101(11):4371-4385. doi: 10.1007/s00253-017-8300-y

422   Passamonti, M., B. Mantovani, V. Scali, and C. Froglia. 1997. Allozymic Characterization of Scottish and Aegean Populations of Nephrops Norvegicus. Journal of the Marine Biological
424   Association of the United Kingdom 77(3):727-735. doi: 10.1017/S0025315400036158

    Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O. Glöckner.
2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Research 41(D1):D590-D596. doi: 10.1093/nar/gks1219

Santibáñez, P., J. Romalde, J. Maldonado, D. Fuentes, and J. Figueroa. 2022. First characterization of the gut microbiome associated with Mytilus chilensis collected at a mussel farm and from
a natural environment in Chile. Aquaculture 548:737644. doi: https://doi.org/10.1016/j.aquaculture.2021.737644

Schloss, P. D., D. Gevers, and S. L. Westcott. 2011. Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. PLoS One 6(12):e27310.

Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A. Lesniewski, B. B. Oakley, D. H. Parks, C. J. Robinson, J. W. Sahl, B. Stres, G. G.
Thallinger, D. J. Van Horn, and C. F. Weber. 2009. Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing
microbial communities. Applied and Environmental Microbiology 75(23):7537-7541. doi: 10.1128/aem.01541-09

See, M. S., X. L. Ching, S. C. Khoo, S. Z. Abidin, C. Sonne, and N. L. Ma. 2025. Aquatic microbiomes under stress: The role of gut microbiota in detoxification and adaptation to
environmental exposures. Journal of Hazardous Materials Advances 17:100612. doi: https://doi.org/10.1016/j.hazadv.2025.100612

Singh, B. K., K. Thakur, H. Kumari, D. Mahajan, D. Sharma, A. K. Sharma, S. Kumar, B. Singh, P. P. Pankaj, and R. Kumar. 2025. A review on comparative analysis of marine and freshwater
fish gut microbiomes: insights into environmental impact on gut microbiota. FEMS Microbiology Ecology 101(1):fiae169. doi: 10.1093/femsec/fiae169

Small, H. J., D. M. Neil, A. C. Taylor, R. J. A. Atkinson, and G. H. Coombs. 2006. Molecular detection of Hematodinium spp. in Norway lobster Nephrops norvegicus and other
crustaceans. Diseases of Aquatic Organisms 69(2-3):185-195.

Stencel, A., and D. Wloch-Salamon. 2022. A pluralistic view of holobionts in the context of process ontology. Frontiers in Microbiology 13(Hypothesis and Theory) doi: 10.3389/fmicb.2022.911577

Subhash, Y., and S.-S. Lee. 2017. Description of *Oceanispirochaeta sediminicola* gen. nov., sp. nov., an obligately anaerobic bacterium isolated from coastal marine sediments, and reclassification of *Spirochaeta litoralis* as *Oceanispirochaeta litoralis* comb. nov. International Journal of Systematic and Evolutionary Microbiology 67(9):3403-3409. doi: https://doi.org/10.1099/ijsem.0.002130

Sun, Y., W. Han, J. Liu, F. Liu, and Y. Cheng. 2020. Microbiota comparison in the intestine of juvenile Chinese mitten crab Eriocheir sinensis fed different diets. Aquaculture 515:734518. doi: https://doi.org/10.1016/j.aquaculture.2019.734518

Theis, K. R., N. M. Dheilly, J. L. Klassen, R. M. Brucker, J. F. Baines, T. C. G. Bosch, J. F. Cryan, S. F. Gilbert, C. J. Goodnight, E. A. Lloyd, J. Sapp, P. Vandenkoornhuyse, I. Zilber-Rosenberg, E. Rosenberg, and S. R. Bordenstein. 2016. Getting the Hologenome Concept Right: an Eco-Evolutionary Framework for Hosts and Their Microbiomes. mSystems 1(2):e00028-00016. doi: 10.1128/mSystems.00028-16

Troussellier, M., A. Escalas, T. Bouvier, and D. Mouillot. 2017. Sustaining rare marine microorganisms: Macroorganisms as repositories and dispersal agents of microbial diversity. Frontiers in Microbiology 8:947. (Hypothesis and Theory) doi: 10.3389/fmicb.2017.00947

Vijayan, N., S. J. McAnulty, G. Sanchez, J. Jolly, Y. Ikeda, M. K. Nishiguchi, E. Réveillac, C. Gestal, B. L. Spady, D. H. Li, B. P. Burford, A. H. Kerwin, and S. V. Nyholm. 2024. Evolutionary history influences the microbiomes of a female symbiotic reproductive organ in cephalopods. Applied and Environmental Microbiology 90(3):e00990-00923. doi: doi:10.1128/aem.00990-23

Wang, Y., U. Stingl, F. Anton-Erxleben, M. Zimmer, and A. Brune. 2004. 'Candidatus Hepatincola porcellionum' gen. nov., sp. nov., a new, stalk-forming lineage of Rickettsiales colonizing the midgut glands of a terrestrial isopod. Archives of Microbiology 181(4):299-304. doi: 10.1007/s00203-004-0655-7

Wardman, J. F., R. K. Bains, P. Rahfeld, and S. G. Withers. 2022. Carbohydrate-active enzymes (CAZymes) in the gut microbiome. Nature Reviews Microbiology 20(9):542-556. doi: 10.1038/s41579-022-00712-1

Waterworth, S. C., G. M. Solomons, J.-C. J. Kalinski, L. S. Madonsela, S. Parker-Nance, and R. A. Dorrington. 2024. The unique and enigmatic spirochete symbiont of latrunculid sponges. mSphere 9(12):e00845-00824. doi: doi:10.1128/msphere.00845-24

Webster, N. S. 2017. Conceptual and methodological advances for holobiont research. Environmental Microbiology Reports 9(1):30-32. doi: 10.1111/1758-2229.12500

West, A. G., D. W. Waite, P. Deines, D. G. Bourne, A. Digby, V. J. McKenzie, and M. W. Taylor. 2019. The microbiome in threatened species conservation. Biological Conservation 229:85-98. doi: https://doi.org/10.1016/j.biocon.2018.11.016

Yadav, S., M. Koenen, N. Bale, J. S. Sinninghe Damsté, and L. Villanueva. 2021. The physiology and metabolic properties of a novel, low-abundance *Psychrilyobacter* species isolated from the anoxic Black Sea shed light on its ecological role. Environmental Microbiology Reports 13(6):899-910. doi: https://doi.org/10.1111/1758-2229.13012

Yilmaz, P., L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J. Peplies, W. Ludwig, and F. O. Glöckner. 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Research 42(D1):D643-D648. doi: 10.1093/nar/gkt1209

Zamora-Briseño, J. A., D. Cerqueda-García, I. M. Hernández-Velázquez, R. Rivera-Bustamante, J. P. Huchín-Mian, P. Briones-Fourzán, E. Lozano-Álvarez, and R. Rodríguez-Canul. 2020. Alterations in the gut-associated microbiota of juvenile Caribbean spiny lobsters Panulirus argus (Latreille, 1804) infected with PaV1. Journal of Invertebrate Pathology 176:107457. doi: https://doi.org/10.1016/j.jip.2020.107457

Zhang, N., C. Song, M. Wang, Y. Liu, M. Hui, and Z. Cui. 2017. Diversity and characterization of bacteria associated with the deep-sea hydrothermal vent crab *Austinograea* sp. comparing with those of two shallow-water crabs by 16S ribosomal DNA analysis. PLOS ONE 12(11):e0187842. doi: 10.1371/journal.pone.0187842

Zilber-Rosenberg, I., and E. Rosenberg. 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. FEMS Microbiology Reviews 32(5):723-735. doi: 10.1111/j.1574-6976.2008.00123.x

Table 1. Midgut bacterial operational taxonomic units (OTUs) from three *Nephrops norvegicus* populations. Sequence reads=33,164; N=7 for male and female individuals, N=5 for eggs.

		Average carapace length (mm) (coefficient of variation)	Average total body weight (g) (coefficient of variation)	No. of OTUs (average, coefficient of variation)	No. of OTUs with ≥80% relative abundance	Most dominant OTU (% relative abundance)
Greece	Eggs	-	-	180 (86, 4.2%)	17	OTU-0020 (18.5%) *Cutibacterium*
	Females	46.0 (11.6%)	74.0 (30.9%)	417 (128, 24.1%)	11	OTU-0004 (17.8%) *Oceanispirochaeta*
	Males	46.7 (14.6%)	77.0 (42.0%)	1290 (335, 64.2%)	20	OTU-0018 (8.4%) *Cyanobium*
Italy	Females	34.9 (4.5%)	32.4 (10.3%)	434 (136, 24.0%)	7	OTU-0002 (20.2%) *Photobacterium*
	Males	36.9 (4.0%)	36.3 (13.9%)	489 (153, 30,4%)	10	OTU-0002 (20.3%) *Photobacterium*
Sweden	Females	45.0 (0.0%)	64.6 (2.5%)	931 (252, 63.0%)	14	OTU-0001 (24.5%) *Psychrilyobacter*
	Male	45.0 (0.0%)	67.9 (1.6%)	527 (161, 36,7%)	11	OTU-0001 (25.5%) *Psychrilyobacter*

Table 2. Permutational analysis of variance (PERMANOVA) of eggs, male and female individuals of *Nephrops norvegicus* gut bacterial communities from Greece (GRE), Italy (ITA) and Sweden (SWE). * : p<0.05, ** : p<0.002, *** : p<0.01.

	Eggs-Females	Eggs-Males	Female-Males		Females	Males
GRE	p=0.027* F=3.162	p=0.046* F=2.190	p=0.897 F=1.565	GRE-ITA	p=0.032* F=2.916	p=0.580 F=1.579
ITA	X	X	p=1.000 F=0.616	ITA-SWE	p=0.027* F=2.981	p=0.021* F=2.668
SWE	X	X	p=0.960 F=1.668	GRE-SWE	p=0.008*** F=3.611	p=0.015* F=3.438

Table 3. Regression of the total body weight vs. the abundance of each of the most important operational taxonomic units (OTU) from three *Nephrops norvegicus* populations. N=42; negative and positive regressions are in red and blue letters, respectively; *: p<0.05; G: Greece, I: Italy, S: Sweden.

OTU	Bacterial taxon	Closest (≥98%) relative in NCBI (GenBank accession no.)	p	r
0001 (G/I/S)	*Psychrilyobacter*	Uncultured bacterium clone SS1_B_01_74 (EU050918)	0.165	0.218
0002 (I/S)	*Photobacterium*	*Photobacterium phosphoreum* (AB179540)	0.031*	-0.333
0003 (G/I/S)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092163)	0.219	-0.194
0004 (G/I)	*Oceanispirochaeta*		0.001*	0.487
0005 (G/I)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092262)	0.016*	-0.368
0006 (S)	*Ancyomarina*	Uncultured Bacteroidetes bacterium clone FII-OX025 (JQ579674)	0.131	0.237
0008 (I)	Mycoplasmataceae (Ca. Hepatoplasma)	Uncult. bacterium from *N. norvegicus* gut (JN092255)	0.089	-0.266
0009 (I)	*Aliivibrio*	*Aliivibrio wodanis* (LR721750)	0.121	-0.243
0010 (G/S)	*Kordiimonas*	Uncult. Alphaproteobacteria from *N. norvegicus* gut (JN092256)	0.040*	0.319

0011 (G)	*Desulfovibrio*		0.001*	0.503
0012 (S)	Desulfobacteraceae unclass.	Uncult. Desulfobacterales from *N. norvegicus* gut (GQ866071)	0.273	0.173
0013 (G/I/S)	*Carboxylicivirga*	Uncult. Bacteroidia from mud crab *(Scylla paramamosain)* gut (HE610318)	0.033*	0.330
0015 (G)	Bacteria		0.015*	0.373
0018 (G)	Cyanobium	Uncultured bacterium clone 2CE2-5m-91 (GU062169)	0.927	0.015
0021 (S)	*Vibrio*	Uncult. bacterium from *N. norvegicus* gut (JN092213)	0.125	0.240
0024 (S)	*Candidatus* Absconditabacteriales		0.470	0.115
0036 (S)	Flavobacteriaceae unclass.	Uncultured bacterium clone EzyYyy31 (KX172210)	0.464	0.116

Figure 1. Non-metric multidimensional scaling (nMDS) of male (♂) and female (♀) individuals of *Nephrops norvegicus* gut from Greece (GRE), Italy (ITA) and Sweden (SWE).

Figure 2. Relative abundance of class-level operational taxonomic units in the female (F) and male (M) *Nephrops norvegicus* gut from Greece (GR), Italy (IT) and Sweden (SW).

Figure 3. Most important (relative abundance ≥1% in both male and female individuals) bacterial operational taxonomic units (OTUs) of the *Nephrops norvegicus* gut from Greece (GRE), Italy (ITA) and Sweden (SWE). Red dots: bacterial operational taxonomic units (OTUs); blue and pink dots: important OTUs, i.e. ≥1% relative abundance in males and females only, respectively.

Figure 4. Ratio of females:males bacterial operational taxonomic units (OTUs) with ≥1% relative abundance in the *Nephrops norvegicus* gut from three populations. Only closest relatives of OTUs with 0.1≤ratio≥10 is shown. Blue bars indicate OTUs occurrence only in males.