Title: Full inter-hemispheric integration sustained by a fraction of posterior callosal fibers

Authors: Tyler Santander[1,2,*], Selin Bekir[2], Theresa Paul[3], Jessica M. Simonson[1,2], Valerie M. Wiemer[3], Henri Etel Skinner[2], Johanna L. Hopf[4], Anna Rada[4], Friedrich G. Woermann[4,5], Thilo Kalbhenn[4,6], Barry Giesbrecht[1,2], Christian G. Bien[4], Olaf Sporns[7], Michael S. Gazzaniga[2], Lukas J. Volz[3,†], & Michael B. Miller[1,2,†,*]

Affiliations:
[1]Institute for Collaborative Biotechnologies, University of California, Santa Barbara, Santa Barbara, CA, USA
[2]Department of Psychological & Brain Sciences, University of California, Santa Barbara, Santa Barbara, CA, USA
[3]Medical Faculty, University of Cologne, and Department of Neurology, University Hospital Cologne, Cologne, Germany
[4]Department of Epileptology, Krankenhaus Mara, Bethel Epilepsy Center, Medical School OWL, Bielefeld University, Bielefeld, Germany
[5]MRI Department, Society for Epilepsy Research, Krankenhaus Mara, Bethel Epilepsy Center, Bielefeld, Germany
[6]Department of Neurosurgery, Evangelisches Klinikum Bethel, Medical School OWL, Bielefeld University, Bielefeld, Germany
[7]Department of Psychological & Brain Sciences, Indiana University, Bloomington, IN, USA
[†]Authors contributed equally
*Corresponding authors: tsantander@ucsb.edu, mbmiller@ucsb.edu

Abstract: The dynamic integration of the lateralized and specialized capacities of the two cerebral hemispheres constitutes a hallmark feature of human brain function. This inter-hemispheric exchange of information critically depends upon the corpus callosum. Classical descriptions of callosal organization outline a topographic gradient, such that specific fibers integrate distinct aspects of brain function. Here we present a challenge to this conventional model. Using neuroimaging data obtained from a new cohort of adult split-brain patients, we leverage modern network neuroscience techniques to show—for the first time—that full inter-hemispheric integration can be achieved via a small proportion of posterior callosal fibers. These findings motivate a novel mechanistic understanding of synchronized inter-hemispheric neural activity for large-scale human brain function and behavior.

Main Text: In recent decades, network neuroscience has emerged as a powerful framework for modeling functional interactions between remote neural systems. A foundational assumption of this approach is that anatomical links between brain regions constrain neuronal dynamics and support the emergence of large-scale functional networks, which in turn enable the efficient integration of perceptual, motor, and higher-order cognitive information necessary for human behavior (*1-6*). The corpus callosum (CC) is thought to be the central conduit through which specialized processing nodes in each cerebral hemisphere coordinate and synchronize function, with 200-250 million axons enabling direct, long-range communication across the brain. Classical topographic models of CC projections have revealed a well-defined anterior-to-posterior anatomical gradient, corresponding to the spatial organization of canonical brain networks servicing various sensorimotor and cognitive functions (*7, 8*). Accordingly, this model predicts that damage to the CC should result in drastic functional disconnection between hemispheres—with spatially-specific disruptions following the localization of lesioned fibers along the aforementioned anatomical gradient.

Split-brain patients, in whom the CC has been severed in an effort to treat medically-intractable epilepsy, offer a powerful means of testing this core conceit of structure-function correspondence. However, neural data on these patients remain scarce—callosotomy was largely discontinued around the turn of the 21st century, coinciding with the rise of network neuroscience. Extant data largely stem from anesthetized pediatric patients (*9-12*). Taken together, these studies support the hypothesis that full versus partial callosotomy confer differential effects with respect to the location and extent of functional disconnections. At the same time, however, they have provided contradictory findings regarding recovery of function post-operatively (*9, 11, 12*), and various other methodological factors obfuscate more straightforward interpretations. Thus, to date, it remains unknown how severing the largest fiber bundle of the human brain affects the topological organization and intrinsic dynamics of large-scale functional networks.

To address this fundamental question, we present the first investigation to characterize the impacts of full and partial callosotomy on the adult human brain using multiple network neuroscience methods. We hypothesized that full callosotomy would wholly disrupt functional synchrony across the cerebral hemispheres, while the spatial distribution of functional disconnections following partial callosotomy would unveil the anatomical gradient of the CC. As predicted, we found that complete sectioning of the CC broadly obliterated both typical inter-hemispheric network topologies and dynamic patterns of inter-hemispheric temporal synchrony. More strikingly, however, we also found that a small proportion of intact posterior CC fibers (a case of partial callosotomy) was sufficient to support widespread inter-hemispheric coupling. These findings challenge our conventional understanding of the CC's functional topography and sheds new light on the critical mechanisms for inter-hemispheric information integration.

Patient descriptions

Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were tested post-operatively at the Bethel Epilepsy Center (Bielefeld, Germany). The time between surgery and testing varied between one and six years (**Table 1**). In two patients (EM and BT), surgery did not result in a complete callosal transection; they are hereafter referred to as *partial splits*. In patient EM, the surgery was stopped after sectioning the anterior third of the CC; patient BT was spared

91 approximately 1 cm of splenium. The remaining four patients (HY, DT, TJ, and LJ) were
92 completely sectioned and are hereafter referred to as *full splits*.
93
Callosotomy is not commonly associated with neuropsychiatric or neurological disorders. Rather,
split-brain patients often demonstrate *disconnection syndromes*, such that they appear to have two
hemispheres operating independently, unable to communicate information from one half of the
brain to the other (*13-15*). Here, behavioral disconnection syndromes were clinically-observed
using simple, lateralized bedside testing protocols. Comprehensive descriptions of these tests are
described elsewhere (*16*). In brief, all four complete callosotomy patients clearly indicated
behavioral disconnections. Partial splits, in contrast, did not show *any* signs of disconnection
syndromes—including, remarkably, patient BT (with a mere 1 cm of splenium). These findings
contradict our classical understanding of callosal topography, which posits functionally-specific
deficits depending upon which portions of the CC were severed.
104
Inter-hemispheric functional connectivity after callosotomy
All patients underwent resting-state functional magnetic resonance imaging without sedation,
affording whole-brain analyses of resting-state functional connectivity (FC) and quantification of
inter-hemispheric FC (IFC). To benchmark patients against 'neurotypical' patterns of IFC, we used
data from the Human Connectome Project 100 Unrelated Subjects (HCP100) cohort. All HCP100
subjects and callosotomy patients underwent an identical preprocessing pipeline, including
conservative corrections for mitigating head motion artifacts (*17, 18*) (see **Fig. S1** for a quantitative
description of patient head motion). Whole-brain FC in the range of 0.01-0.08 Hz was assessed by
parcellating the brain into 400 cortical regions and 32 subcortical regions (according to the
Schaefer (*19, 20*) and Tian (*21*) atlases, respectively; **Fig. 2a**) and correlating all pairs of regional
time series (**Figs. 2b-c**).
116
All full splits demonstrated marked reductions in IFC relative to both neurotypical controls and
partial splits (**Figs. 2b-c**). Spatial patterns of intra-hemispheric FC, by contrast, were generally
better-matched to controls (**Fig. S2**), suggesting that callosotomy primarily impaired inter-
hemispheric functional coupling with full splits suffering the greatest disruptions. Curiously,
however, while gross IFC was dissolved in full splits, inspection of individual FC matrices
suggested that visual network connectivity was partially preserved (**Fig. 2c**). This may be due to
synchronized perceptual inputs (scans were acquired with eyes open), driving *apparent* IFC
despite the lack of transcallosal connectivity between visual regions and evidence for behavioral
disconnections. We therefore sought to assess differences in the magnitude of IFC between
functional homologues to determine whether any preserved IFC was unique to sensory systems.
Here we computed a series of *network laterality indices*, where a value of zero indicates a perfect
balance between inter- and intra-hemispheric FC, and values closer to -1 or 1 indicate relatively
more inter- or intra-hemispheric FC, respectively. Consistent with observed behavioral
disconnections, all full splits demonstrated more lateralized patterns of FC relative to healthy
controls and partial splits, who typically fell within the control distributions (**Fig. 3**). This was
evident in both higher-order, transmodal systems (e.g., the default mode and dorsal attention
networks) and in somatosensory networks (with the exception of patient TJ who fell within the
upper tail of the control distribution in the somatomotor network). Patient EM was more lateralized
across frontoparietal control nodes, which include a number of regions likely to be structurally-
disconnected given the extent of their callosotomy; however, BT generally showed degrees of

network bilaterality equivalent to healthy adults, despite having a mere fraction of splenium intact. Finally, full splits appeared to show conserved bilateral FC across limbic regions like the orbitofrontal cortex and temporal poles, which are known to be structurally coupled through the preserved anterior commissure as well as the CC (*22-24*). Taken together, these results highlight the dependence of inter-hemispheric networks on CC connectivity—yet, neurotypical levels of functional integration were possible even when only the splenium was left intact.

Lateralized expressions of functional modules

To further elucidate whole-brain differences in functional network topology—agnostic to *a priori* assignment of individual brain regions to canonical functional systems—we employed data-driven community detection techniques to cluster whole-brain patterns of FC into functional *modules* (**Fig. 4**). We hypothesized that the breakdown of IFC in full splits would produce lateralized network topologies, predominantly self-organized within one hemisphere or the other. Here we used a modular laterality index computed according to prior work (*25*): where a value of zero, for example, indicates that all functional modules are perfectly bilateral (i.e., the constituent brain regions belonging to each community are equally-balanced between the left and right hemispheres), and values closer to one indicate progressively more lateralized topologies.

As predicted, we observed much greater degrees of modular laterality in all full splits (**Fig. 5**). Communities with bihemispheric representations primarily spanned visual and somatomotor regions—reflecting a mechanism through which synchronized, external sensory inputs can drive apparent functional integration. With respect to partial splits, patient EM with the most CC showed bilateral patterns of network organization mirroring neurotypical controls. Patient BT also fell within the control distribution—however, they showed a strikingly different pattern of modularity (**Fig. 4**). Here, functional communities appeared to dissociate along the Sylvian fissure, with dorsal modules and ventral modules spanning the anteroposterior axis of the brain. This individual had the longest gap between surgery and testing (~6 years), which may have allowed for widespread network reorganization affording optimal utilization of the minimal callosal fibers left intact. Together, these findings highlight the importance of transcallosal connections for the emergence of coherent inter-hemispheric systems, while further highlighting a role for sensory inputs enabling apparent preservation of bilateral sensorimotor modules.

Inter-hemispheric synchrony in network dynamics

The modular organization of the brain plays a critical role in governing functional network dynamics, with structural architectures orchestrating widespread synchronization of whole-brain activity over time (*26*). Thus, we used *edge time series* (ETS) (*27, 28*) to test the hypothesis that structural connectivity via the CC is necessary for large-scale coordination of function—capturing moment-to-moment changes in the extent to which *many* pairs of brain regions within each hemisphere co-fluctuated in activity (**Fig. 6a**).

Among neurotypical controls, we observed tight temporal coupling between left and right hemisphere dynamics (Median Spearman $r = 0.82$, *MAD* $= 0.08$). In contrast, all four full splits demonstrated significant reductions in inter-hemispheric synchrony, suggesting that network dynamics were more functionally-independent (**Fig. 6b**; see **Fig. S4** for individual networks). This is consistent with the overall attenuation of inter-hemispheric FC and further indicates that brief moments of high-amplitude co-fluctuations (which normally encompass regions in both

hemispheres and are thought to be structurally-mediated (*26*)) occur separately in each. Notably, both partial splits fell within the standard range of the HCP subjects. These results therefore strongly suggest a mechanistic role for CC structural connections in supporting widespread integration of inter-hemispheric network dynamics. This may be achieved with a mere fraction of intact fibers through the splenium and thus might be largely-independent of the presence of direct cortico-cortical connections.

Discussion

We characterized variable disruptions of inter-hemispheric functional connectivity following complete versus partial transection of the corpus callosum. Full splits demonstrated more lateralized network topologies and desynchronized network dynamics, aligning with classical behavioral disconnection syndromes. All patients had intact anterior and posterior commissures along with other subcortical routes of inter-hemispheric communication. These findings therefore suggest that transcallosal connections are the essential pathways through which large-scale inter-hemispheric functional networks emerge. Contrary to our expectations, however, inter-hemispheric functional coupling among partial splits remained intact—these patients additionally showed *no* signs of behavioral disconnection, even when the vast majority of the callosum was severed (*16*). Thus, a fraction of callosal fibers in the splenium may be sufficient to orchestrate inter-hemispheric functional synchrony and integrate behavior. This poses a remarkable challenge to our classical topographic models of callosal connectivity and function, motivating a novel view on inter-hemispheric structure-function relationships: namely, that robust inter-hemispheric synchrony and information transfer can be achieved via polysynaptic connectivity through a relatively small proportion of posterior transcallosal connections, rather than direct structural connectivity between functional homologues.

Complete callosotomy and structure-function relationships

Previous investigations of pediatric callosotomy patients have provided an initial perspective on the extent to which spatially-distributed functional networks depend upon transcallosal connectivity (*9-12*). However, these patients often suffer from severe cognitive and developmental deficits, requiring scans to be completed under sedation—the latter of which is known to reduce dynamical complexity in brain activity and produce functional connectomes more rigidly tied to direct anatomical pathways (*29*). It is further known that functional and structural brain networks undergo profound developmental changes early in life (*30-33*). In other words, neural plasticity in these young individuals may still afford the long-term establishment of alternate inter-hemispheric pathways (perhaps akin to callosal agenesis patients (*34*)), and indeed, it is well-documented that disconnection syndromes in pediatric patients resolve over time (*35, 36*). This likely accounts for differential reports of recovery in inter-hemispheric FC among these individuals (*9, 11, 12*).

The adult human brain also retains neuroplastic potential (*37*); however, given the enormity of the structural disconnection, adult callosotomy patients may lack the capacity to adapt their functional architectures in such a way that allows complete recovery of communication between disconnected brain regions. Our results clearly underscore this hypothesis. Whole-brain patterns of FC appeared highly lateralized, both with respect to canonical network definitions (**Figs. 2-3**) and the topological organization of functional modules (**Figs. 4-5**). Furthermore, we observed considerable reductions in the extent to which multivariate co-fluctuations in activity were synchronized between hemispheres (**Fig. 6**), underscoring a role for the CC in orchestrating whole-brain network

dynamics. Importantly, however, there were some exceptions to these effects, which follow anatomical predictions. Limbic and subcortical regions—the former of which are known to be connected through the anterior commissure (AC) as well as the CC (*22-24*)—showed conserved bilateral FC (**Fig. 3**). Nevertheless, it seems clear that non-callosal pathways were insufficient to support widespread inter-hemispheric synchrony, contrary to previous proposals both in humans (*38-40*) and nonhuman primates (where AC fibers are sufficient to sustain inter-hemispheric FC) (*41*). Finally, visual and sensorimotor areas also appeared to remain somewhat integrated. We cannot fully discount the possibility that subcortical information transfer contributed to this preserved synchrony; however, an alternative hypothesis may lie in the fact that visual and somatosensory inputs to each hemisphere were synchronized during the resting-state scan— producing *apparent* functional coherence despite observed behavioral disconnections.

Preserved inter-hemispheric integration in partial splits
While our observations in full splits clearly emphasize the importance of CC connectivity in orchestrating inter-hemispheric synchrony, partial splits revealed something much more surprising. We predicted that partial splits would show specific deficits in inter-hemispheric FC along an anterior-to-posterior gradient corresponding to the extent of the callosal transection (following previous characterizations of CC topography (*7, 8*)). However, both partial splits consistently demonstrated levels of inter-hemispheric integration in the range of neurotypical controls (**Figs. 2-6**). This was even observed for patient BT, in whom only ~1 cm of posterior callosal fibers were spared. It has become increasingly-evident that structure-function relationships are highly complex and generally do not reflect simple, one-to-one interactions between directly-coupled areas (*42-44*). Still, these remarkable results imply that the functional topography of the CC may be much more flexible than previously-imagined. In particular, there may be a critical threshold at which full functional integration can be sustained given a sufficient proportion of intact fibers—even if the vast majority of the callosum has been severed.

An intriguing extension of this hypothesis is that the *location* of the spared callosal fibers may be critical. Previous cytological work has shown that anterior portions of the CC are primarily populated by thin fibers, while fibers in the tail locally-vary in diameter (*45-47*). Such differences in fiber size and myelination along the CC have been associated with differences in conduction velocity (*47*) and, when considered alongside the topographic organization of the CC, may account for the 'fast' versus 'slow' action of primary sensorimotor and association cortices, respectively. Spared posterior fibers as in BT could therefore be more beneficial than spared anterior fibers. However, from a sheer bandwidth perspective, it is difficult to imagine how such a small proportion of fibers could account for widespread integration of function and cognitive processing. It is also worth noting that previous behavioral studies of partial callosotomy patients *have* shown expected disconnection effects given the specific portions of the CC that were severed, although no functional imaging was performed and these tests were largely conducted within the first year post-operatively (*48-51*).

Accordingly, it appears that functional systems in BT dramatically reorganized in the six years following surgery. In brains that develop normally with the support of the CC, multisensory integration zones in parietal cortex may serve as inter-hemispheric hubs (*52, 53*). By taking advantage of spared posterior callosal pathways connecting these regions, the processing of various intra-hemispheric brain regions can still be integrated and exchanged along polysynaptic

routes running between hemispheres. The striking differences in functional network topology observed in BT, with large modules spanning the anterior-to-posterior axis of the brain (**Fig. 4**), are one potential consequence of this reorganization, reflecting greater functional dependencies between frontal and posterior areas affording more optimal usage of the spared splenial fibers. Of note, the fundamental importance of these posterior inter-hemispheric connections aligns with and may further inform neuroscientific theories of consciousness (*54, 55*). The temporal-parietal-occipital 'hot zone' of regions recently proposed to maximize the singular 'here-and-now' *qualia* of consciousness (*54, 56*) are likely dependent (in part) upon CC connections through the splenium. This may explain why BT maintains an integrated conscious experience while full splits demonstrate inter-hemispheric disintegration (*16*).

Limitations

We acknowledge the small number of patients included in this study, each of whom have limited post-operative data. While previous pediatric studies have recruited larger samples, the availability of patients who underwent callosotomy in adulthood is extremely scarce. It is often difficult to draw robust and reliable statistical inferences from neuroimaging data acquired on such small samples. However, the marked differences observed between full splits and healthy controls suggest that we are not parsing trivial effects on the scale of traditional neuroimaging—rather, they reflect outcomes drawn from fundamentally unique sampling distributions. Furthermore, relative to other lesion studies (often hampered by variability in spatial localization/extent), callosotomy is a highly-selective intervention. By severing a single white matter structure in the brain without any damage to gray matter, callosotomy may afford more homogeneous outcomes. It will be critical for future studies to include larger samples with more imaging data to better characterize the extent of neural variability in callosotomy patients, especially as network architectures reorganize over time.

Conclusion

Following more than a half-century of behavioral neuroscience, split-brain patients continue to provide startling insights into the fundamental nature of the human brain. The rise of network neuroscience has further transformed our understanding of large-scale brain systems, tracing the pathways relating brain structure to function and behavior. Our findings reveal a unique type of criticality in brain function—such that a vanishingly-small proportion of posterior callosal fibers may be sufficient to support the integration of inter-hemispheric functional brain networks and unified cognitive processing. It has long been celebrated that the human brain is a remarkably adaptable complex system. These findings challenge us to rethink our conventional assumptions about structure-function relationships and the means through which the brain enables the mind.

References

1. D. S. Bassett, O. Sporns, Network neuroscience. *Nat Neurosci* **20**, 353-364 (2017).
2. C. J. Honey *et al.*, Predicting human resting-state functional connectivity from structural connectivity. *Proc Natl Acad Sci U S A* **106**, 2035-2040 (2009).
3. M. P. van den Heuvel, R. C. Mandl, R. S. Kahn, H. E. Hulshoff Pol, Functionally linked resting-state networks reflect the underlying structural connectivity architecture of the human brain. *Hum Brain Mapp* **30**, 3127-3141 (2009).

4. M. Fukushima *et al.*, Structure-function relationships during segregated and integrated network states of human brain functional connectivity. *Brain Struct Funct* **223**, 1091-1106 (2018).

5. A. M. Hermundstad *et al.*, Structurally-constrained relationships between cognitive states in the human brain. *PLoS Comput Biol* **10**, e1003591 (2014).

6. A. M. Hermundstad *et al.*, Structural foundations of resting-state and task-based functional connectivity in the human brain. *Proc Natl Acad Sci U S A* **110**, 6169-6174 (2013).

7. S. Hofer, J. Frahm, Topography of the human corpus callosum revisited--comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. *Neuroimage* **32**, 989-994 (2006).

8. Y. Liu *et al.*, Connectivity-Based Topographical Changes of the Corpus Callosum During Aging. *Front Aging Neurosci* **13**, 753236 (2021).

9. S. C. Hung *et al.*, Early recovery of interhemispheric functional connectivity after corpus callosotomy. *Epilepsia* **60**, 1126-1136 (2019).

10. J. M. Johnston *et al.*, Loss of resting interhemispheric functional connectivity after complete section of the corpus callosum. *J Neurosci* **28**, 6453-6458 (2008).

11. J. L. Roland *et al.*, On the role of the corpus callosum in interhemispheric functional connectivity in humans. *Proc Natl Acad Sci U S A* **114**, 13278-13283 (2017).

12. C. E. Pizoli *et al.*, Resting-state activity in development and maintenance of normal brain function. *Proc Natl Acad Sci U S A* **108**, 11638-11643 (2011).

13. M. S. Gazzaniga, Forty-five years of split-brain research and still going strong. *Nat Rev Neurosci* **6**, 653-659 (2005).

14. L. J. Volz, M. S. Gazzaniga, Interaction in isolation: 50 years of insights from split-brain research. *Brain* **140**, 2051-2060 (2017).

15. M. B. Miller, L. J. Volz, J. M. Simonson, M. S. Gazzaniga, in *Cerebral Asymmetries,* C. Papagno, P. Corballis, Eds. (Elsevier BV, San Diego, in press).

16. S. Bekir *et al.*, No disconnection syndrome after near-complete callosotomy. *bioRxiv*. 2025.

17. R. Ciric *et al.*, Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *Neuroimage* **154**, 174-187 (2017).

18. L. Parkes, B. Fulcher, M. Yucel, A. Fornito, An evaluation of the efficacy, reliability, and sensitivity of motion correction strategies for resting-state functional MRI. *Neuroimage* **171**, 415-436 (2018).

19. A. Schaefer *et al.*, Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114 (2018).

20. B. T. Yeo *et al.*, The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).

21. Y. Tian, D. S. Margulies, M. Breakspear, A. Zalesky, Topographic organization of the human subcortex unveiled with functional connectivity gradients. *Nat Neurosci* **23**, 1421-1432 (2020).

22. M. Catani, M. Schotten, *Atlas of Human Brain Connections*. (Oxford University Press, Oxford, UK, 2012).

363   23.   M. D. Patel, N. Toussaint, G. D. Charles-Edwards, J. P. Lin, P. G. Batchelor, Distribution
364   and fibre field similarity mapping of the human anterior commissure fibres by diffusion
365   tensor imaging. *MAGMA* **23**, 399-408 (2010).

366   24.   G. Di Virgilio, S. Clarke, G. Pizzolato, T. Schaffner, Cortical regions contributing to the
367   anterior commissure in man. *Exp Brain Res* **124**, 1-7 (1999).

368   25.   K. W. Doron, D. S. Bassett, M. S. Gazzaniga, Dynamic network structure of
369   interhemispheric coordination. *Proc Natl Acad Sci U S A* **109**, 18661-18668 (2012).

370   26.   M. Pope, M. Fukushima, R. F. Betzel, O. Sporns, Modular origins of high-amplitude
371   cofluctuations in fine-scale functional connectivity dynamics. *Proc Natl Acad Sci U S A*
372   **118**, (2021).

373   27.   J. Faskowitz, F. Z. Esfahlani, Y. Jo, O. Sporns, R. F. Betzel, Edge-centric functional
374   network representations of human cerebral cortex reveal overlapping system-level
375   architecture. *Nat Neurosci* **23**, 1644-1654 (2020).

376   28.   F. Zamani Esfahlani *et al.*, High-amplitude cofluctuations in cortical activity drive
377   functional connectivity. *Proc Natl Acad Sci U S A* **117**, 28393-28401 (2020).

378   29.   P. Barttfeld *et al.*, Signature of consciousness in the dynamics of resting-state brain
379   activity. *Proc Natl Acad Sci U S A* **112**, 887-892 (2015).

380   30.   U. A. Tooley *et al.*, The Age of Reason: Functional Brain Network Development during
381   Childhood. *J Neurosci* **42**, 8237-8251 (2022).

382   31.   V. J. Sydnor *et al.*, Neurodevelopment of the association cortices: Patterns, mechanisms,
383   and implications for psychopathology. *Neuron* **109**, 2820-2846 (2021).

384   32.   J. E. Reynolds, M. N. Grohs, D. Dewey, C. Lebel, Global and regional white matter
385   development in early childhood. *Neuroimage* **196**, 49-58 (2019).

386   33.   K. J. Whitaker *et al.*, Adolescence is associated with genomically patterned consolidation
387   of the hubs of the human brain connectome. *Proc Natl Acad Sci U S A* **113**, 9105-9110
388   (2016).

389   34.   J. M. Tyszka, D. P. Kennedy, R. Adolphs, L. K. Paul, Intact bilateral resting-state
390   networks in the absence of the corpus callosum. *J Neurosci* **31**, 15154-15162 (2011).

391   35.   M. Lassonde, H. Sauerwein, A. J. Chicoine, G. Geoffroy, Absence of disconnexion
392   syndrome in callosal agenesis and early callosotomy: brain reorganization or lack of
393   structural specificity during ontogeny? *Neuropsychologia* **29**, 481-495 (1991).

394   36.   D. Graham, M. M. Tisdall, D. Gill, Corpus callosotomy outcomes in pediatric patients: A
395   systematic review. *Epilepsia* **57**, 1053-1068 (2016).

396   37.   E. Fuchs, G. Flugge, Adult neuroplasticity: more than 40 years of research. *Neural Plast*
397   **2014**, 541870 (2014).

398   38.   J. S. Nomi *et al.*, Diffusion weighted imaging evidence of extra-callosal pathways for
399   interhemispheric communication after complete commissurotomy. *Brain Struct Funct*
400   **224**, 1897-1909 (2019).

401   39.   L. Q. Uddin *et al.*, Residual functional connectivity in the split-brain revealed with
402   resting-state functional MRI. *Neuroreport* **19**, 703-709 (2008).

403   40.   I. Marcantoni *et al.*, Interhemispheric functional connectivity: an fMRI study in
404   callosotomized patients. *Front Hum Neurosci* **18**, 1363098 (2024).

405   41.   J. X. O'Reilly *et al.*, Causal effect of disconnection lesions on interhemispheric functional
406   connectivity in rhesus monkeys. *Proc Natl Acad Sci U S A* **110**, 13982-13987 (2013).

42. C. Seguin, L. S. Mansour, O. Sporns, A. Zalesky, F. Calamante, Network communication
models narrow the gap between the modular organization of structural and functional
brain networks. *Neuroimage* **257**, 119323 (2022).
43. L. E. Suarez, R. D. Markello, R. F. Betzel, B. Misic, Linking Structure and Function in
Macroscale Brain Networks. *Trends Cogn Sci* **24**, 302-315 (2020).
44. B. Misic *et al.*, Network-Level Structure-Function Relationships in Human Neocortex.
Cereb Cortex **26**, 3285-3296 (2016).
45. F. Aboitiz, J. Lopez, J. Montiel, Long distance communication in the human brain: timing
constraints for inter-hemispheric synchrony and the origin of brain lateralization. *Biol Res*
36, 89-99 (2003).
46. F. Aboitiz, A. B. Scheibel, R. S. Fisher, E. Zaidel, Fiber composition of the human corpus
callosum. *Brain Res* **598**, 143-153 (1992).
47. A. Horowitz *et al.*, In vivo correlation between axon diameter and conduction velocity in
the human brain. *Brain Struct Funct* **220**, 1777-1788 (2015).
48. M. G. Funnell, P. M. Corballis, M. S. Gazzaniga, Cortical and subcortical
interhemispheric interactions following partial and complete callosotomy. *Arch Neurol*
57, 185-189 (2000).
49. M. S. Gazzaniga, H. Freedman, Observations on visual processes after posterior callosal
section. *Neurology* **23**, 1126-1130 (1973).
50. G. L. Risse, J. Gates, G. Lund, R. Maxwell, A. Rubens, Interhemispheric transfer in
patients with incomplete section of the corpus callosum. Anatomic verification with
magnetic resonance imaging. *Arch Neurol* **46**, 437-443 (1989).
51. M. S. Gazzaniga, in *Neurobiology of Neocortex,* P. Rakic, W. Singer, Eds. (John Wiley &
Sons, New York, 1988), pp. 385-405.
52. M. P. van den Heuvel, O. Sporns, Rich-club organization of the human connectome. *J*
Neurosci **31**, 15775-15786 (2011).
53. P. Hagmann *et al.*, Mapping the structural core of human cerebral cortex. *PLoS Biol* **6**,
e159 (2008).
54. C. Koch, M. Massimini, M. Boly, G. Tononi, Neural correlates of consciousness:
progress and problems. *Nat Rev Neurosci* **17**, 307-321 (2016).
55. L. Melloni *et al.*, An adversarial collaboration protocol for testing contrasting predictions
of global neuronal workspace and integrated information theory. *PLoS One* **18**, e0268577
(2023).
56. C. Koch, M. Massimini, M. Boly, G. Tononi, Posterior and anterior cortex - where is the
difference that makes the difference? *Nat Rev Neurosci* **17**, 666 (2016).
57. D. C. Van Essen *et al.*, The WU-Minn Human Connectome Project: an overview.
Neuroimage **80**, 62-79 (2013).
58. D. C. Van Essen *et al.*, The Human Connectome Project: a data acquisition perspective.
Neuroimage **62**, 2222-2231 (2012).
59. N. J. Tustison *et al.*, The ANTsX ecosystem for quantitative biological and medical
imaging. *Sci Rep* **11**, 9068 (2021).
60. N. J. Tustison *et al.*, Large-scale evaluation of ANTs and FreeSurfer cortical thickness
measurements. *Neuroimage* **99**, 166-179 (2014).
61. A. C. Evans, A. L. Janke, D. L. Collins, S. Baillet, Brain templates and atlases.
Neuroimage **62**, 911-922 (2012).

62.   M. Jenkinson, C. F. Beckmann, T. E. Behrens, M. W. Woolrich, S. M. Smith, Fsl.
Neuroimage **62**, 782-790 (2012).
63.   K. Hua *et al.*, Tract probability maps in stereotaxic spaces: analyses of white matter
anatomy and tract-specific quantification. *Neuroimage* **39**, 336-347 (2008).
64.   S. Wakana *et al.*, Reproducibility of quantitative tractography methods applied to cerebral
white matter. *Neuroimage* **36**, 630-644 (2007).
65.   K. J. Friston, P. Rotshtein, J. J. Geng, P. Sterzer, R. N. Henson, A critique of functional
localisers. *Neuroimage* **30**, 1077-1087 (2006).
66.   O. Sporns, R. F. Betzel, Modular Brain Networks. *Annu Rev Psychol* **67**, 613-640 (2016).
67.   F. Zamani Esfahlani *et al.*, Modularity maximization as a flexible and generic framework
for brain network exploratory analysis. *Neuroimage* **244**, 118607 (2021).
68.   M. Rubinov, O. Sporns, Weight-conserving characterization of complex functional brain
networks. *Neuroimage* **56**, 2068-2079 (2011).
69.   M. Rubinov, O. Sporns, Complex network measures of brain connectivity: uses and
interpretations. *Neuroimage* **52**, 1059-1069 (2010).
70.   S. Fortunato, M. Barthelemy, Resolution limit in community detection. *Proc Natl Acad*
Sci U S A **104**, 36-41 (2007).
469

470 **Acknowledgements:**
471 **Funding:**
472 Army Research Office under contract W911NF-19-D-0001 for the Institute for Collaborative
473 Biotechnologies (TS, SB, JMS, HES, BG, MBM)
474 German Research Foundation, CRC-1451, Project 431549029 (TP, LJV)
475
Author contributions:
Conceptualization: TS, SB, TP, HES, JLH, AR, BG, CGB, OS, MSG, LJV, MBM
Investigation: TP, VMW, JLH, AR, FGW, TK, CGB, LJV
Data curation: TP, VMW, JLH, FGW, LJV
Methodology: TS, JMS, OS, LJV
Formal analysis: TS, JMS, OS
Visualization: TS
Writing—original draft: TS, TK, OS, MSG, LJV, MBM
Writing—review & editing: TS, SB, TP, JMS, VMW, HES, LJH, AR, FGW, TK, BG, CGB, OS,
MSG, LJV, MBM
Project administration: AR, CGB, LJV, MBM
Funding acquisition: BG, LJV, MBM
488
489 **Competing interests:** Authors declare that they have no competing interests.
490
491 **Data and materials availability:** Patient data will be made available upon reasonable request to
492 the corresponding authors; all relevant processing and analysis code will be uploaded to a Github
493 repository.

494
495 **List of Supplementary Materials:**
496 Materials and Methods
497 Figs. S1 to S5

Fig. 1. Patient anatomical scans. Post-operative T_1 and diffusion MRI scans collected for all six patients at time of testing. White arrows/ellipses for patients EM and BT highlight intact CC (i.e., partial splits). The remaining four patients were fully-sectioned. Note that a series of clinical observations suggested behavioral disconnection syndromes in all full splits, but not in partial splits.

Fig. 2. Inter-hemispheric functional connectivity is broadly-diminished in full (but not partial) splits. (a) Whole-brain functional parcellation comprised of seven canonical resting-state networks (*19, 20*) and subcortical nodes (*21*). **(b)** A standard reference for neurotypical whole-brain functional connectivity (FC), given as the median over HCP100 subjects. FC was estimated by correlating low-frequency brain activity (0.01-0.08 Hz) between all pairs of regions in **(a)**. The large boxes outlining the upper left and lower right quadrants denote *intra-hemispheric* FC; off-diagonal quadrants thus capture *inter-hemispheric* FC (IFC). **(c)** Whole-brain FC estimates for

each callosotomy patient, thresholded at $p < .001$ (uncorrected for multiple comparisons). Typical patterns of IFC were largely abolished in full splits (HY, DT, TJ, LJ), while partial splits (EM, BT) were more similar to healthy controls. Abbreviations: SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network; LH, Left Hemisphere; RH, Right Hemisphere.

Fig. 3. Full callosotomy is associated with more lateralized functional connectivity. Network laterality indices across seven canonical resting-state systems (*19, 20*) and subcortical regions (*21*). Values closer to zero indicate an equal balance between intra- and inter-hemispheric FC, while values closer to one indicate more intra-hemispheric FC relative to inter-hemispheric FC. Note that each subplot captures a unique range of values, given differences in variability observed across neurotypical adults (colored distributions), patients, and networks. Full splits generally demonstrated more strongly-lateralized FC compared to partials, who fell within the control distributions. Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Fig. 4. Modular decomposition of whole-brain functional connectivity. Spatial maps of optimal, consensus community assignments derived via data-driven modularity maximization at a default level of resolution ($\gamma = 1$). Regional coloring indicates assignment to a given functional module. Full splits generally exhibited more lateralized clustering of regions into modules relative to partial splits and healthy adult controls (see **Fig. 5** for quantification of module laterality).

Fig. 5. Functional modules are more strongly-lateralized in full splits. The modular laterality index quantifies the extent to which functional communities exhibit bihemispheric representations (values closer to zero) or are restricted to one hemisphere or the other (values closer to one). At a default level of resolution ($\gamma = 1$), partial splits demonstrated bilateral network topologies similar to healthy controls, whereas full splits were consistently more lateralized. Point intervals under the control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively).

Fig. 6. Large-scale functional network dynamics are desynchronized in full splits. **(a)** An illustration of edge time series in the left and right hemispheres for patient BT (a partial split with 1 cm of intact splenium) and patient TJ (a full split). By taking the root-sum-of-squares (RSS) across all edges at each timepoint, we derived a summary estimate of the total amplitude of co-fluctuating brain activity. Note the bursty, high-amplitude co-fluctuations in each hemisphere that often coincide in time for BT, but not in TJ. **(b)** All four full splits exhibited reduced temporal synchrony (i.e., temporal correlations) between left/right hemisphere RSS time courses. Partial splits, by contrast, fell within the range of healthy adult controls. Point intervals under the control

557 distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median
558 (thick black lines and thin black lines, respectively).
559
560 **Table 1. General overview of callosotomy patients.**

Patient	Age at surgery	Age at testing	Sex	Handedness	Extent of callosotomy	Disconnection syndrome
EM	17	19	M	Right	Partial	Absent
BT	22	28	M	Right	Partial	Absent
HY	21	22	M	Right	Complete	Present
DT	15	20	M	Right	Complete	Present
TJ	49	52	F	Right	Complete	Present
LJ	57	60	M	Right	Complete	Present

561
562

Supplementary Materials for

Full inter-hemispheric integration sustained by a fraction of posterior callosal fibers

Tyler Santander, Selin Bekir, Theresa Paul, Jessica M. Simonson, Valerie M. Wiemer, Henri Etel Skinner, Johanna L. Hopf, Anna Rada, Friedrich G. Woermann, Thilo Kalbhenn, Barry Giesbrecht, Christian G. Bien, Olaf Sporns, Michael S. Gazzaniga, Lukas J. Volz, & Michael B. Miller

Corresponding authors: tsantander@ucsb.edu, mbmiller@ucsb.edu

The PDF file includes:

Materials and Methods
Figs. S1 to S5

608
Materials and Methods
Materials and Methods

<u>Patient recruitment</u>
Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were recruited
during routine post-operative follow-up visits to the Bethel Epilepsy Center (Bielefeld, Germany).
The study was approved by the Ethics Committee of the University of Münster,
Germany (2021–523-f-S). The patients or their legal representatives provided written informed
consent.
616

<u>Callosotomy procedure</u>
Callosotomies were performed with the assistance of intraoperative neuronavigation, with patients
in a supine position and the head inclined by roughly 30° (fixed in the Mayfield clamp). A straight
skin incision was made running slightly next to the midline on the non-language dominant
hemisphere. Craniotomy was then performed by making one drill hole in the midline at
approximately the level of the precentral sulcus. A bone flap was sawn out with an outline of 3 cm
x 5 cm. The dura was incised longitudinally and then opened with two wings under
microneurosurgical conditions—access to the inter-hemispheric fissure was successively created
while preserving the cortical bridging veins. Next, using a self-retaining retractor to carefully shift
the frontal lobe, the level of the corpus callosum was gradually exposed under sharp dissection
with micro scissors.
628

Dissection is often made considerably more difficult by the course of the pericallosal arteries,
which run parallel to the corpus callosum but are often tightly intertwined and twisted in the inter-
hemispheric fissure. In cases where the risk of vascular damage is too great, the surgery may be
halted short of a full disconnection (resulting in partial splits). However, under most conditions,
the inter-hemispheric fissure may be opened along the entire length of the knee of the corpus
callosum. The corpus callosum was then perforated into the lateral ventricle of the ipsilateral
hemisphere. Next, following the pericallosal artery and the anterior cerebral artery down to the
frontal base, the knee of the callosum was split in the midline, continuing backwards to about the
level of the foramen of Monroi. From this point, it is critical to protect the fornix of the language-
dominant hemisphere: to mitigate the risk of postoperative functional impairment, we carefully
ensure that the self-retaining retractor is not placed on the precentral gyrus within the inter-
hemispheric fissure. The disconnection can now be carried out completely into the splenium—the
severing of the splenium was performed subpially to protect all the cisternal vessels. Extensive
hemostasis is not usually necessary, and thus a ventricular drainage is not used. To conclude, the
dura mater was closed and the bone flap reinserted and fixed with titanium plates and screws.
Finally, a subgaleal drainage was inserted and the skin closed in two layers.
645

<u>Healthy adult controls</u>
We used open-access data from the Human Connectome Project 100 Unrelated Subjects (HCP100)
cohort (46 men, 54 women; $M_{Age} = 29.11$, $SD = 3.68$) as a robust benchmark for various network
features estimated in callosotomy patients. Participant recruitment and imaging procedures have
previously been detailed elsewhere (*57, 58*). In brief, data were collected over two sessions on a
custom 3T Siemens Skyra with a 32-channel head coil. T_1-weighted structural scans were acquired
using a 3D magnetization prepared rapid gradient echo (MPRAGE) sequence (*TR* = 2400 ms, *TE*
= 2.14 ms, *TI* = 1000 ms, flip angle = 8º, voxel size = 0.7 mm^3). Resting-state fMRI was collected

with eyes open over four ~14-minute scans, using a multiband echo-planar imaging (EPI) sequence
sensitive to the blood-oxygen-level-dependent (BOLD) contrast (TR = 720 ms, TE = 33.1 ms, flip
angle = 52º, voxel size = 2 mm^3, multiband factor = 8). All participants gave written informed
consent for a protocol approved by the Washington University Institutional Review Board. All
imaging data were downloaded in their raw (unprocessed) state.

<u>Patient neuroimaging procedures</u>
Patient data were collected on a 3T Siemens Vida at the Bethel Epilepsy Center using a 32-channel
head coil. T_1-weighted structural scans were acquired using a 3D MPRAGE sequence (TR = 1900
ms, TE = 2.47 ms, TI = 900 ms, flip angle = 8º, voxel size = 0.8 mm^3). Resting-state fMRI was
collected with eyes open using a single-shot EPI sequence sensitive to the BOLD contrast (TR =
2000 ms, TE = 30 ms, flip angle = 75º, voxel size = 3 x 3 x 3.30 mm, iPAT/GRAPPA factor = 2).

We note that each patient varied somewhat in the amount of resting-state data obtained (from ~8-
20 minutes total) due to differences in clinical scheduling and other imaging protocols not relevant
to the current analysis. This information is summarized in **Fig. S1**.

<u>fMRI preprocessing</u>
All data (from both patients and healthy controls) were run through a unified preprocessing
pipeline. Initial preprocessing relied on functions from the Statistical Parametric Mapping 12
software (SPM12, Wellcome Trust Centre for Neuroimaging, London) in Matlab and Advanced
Normalization Tools (ANTs) (*59*). Resting-state fMRI data were realigned and unwarped to correct
for head motion and motion x susceptibility artifact as well as slice-time corrected. Anatomical
images were denoised (ANTs' `DenoiseImage` with a spatially-adaptive Gaussian filter) and
submitted to ANTs' cortical thickness pipeline, which includes N4 bias field correction and high-
quality image segmentation and brain extraction (*60*). For some patients, the brain extraction mask
included extracerebral components: this was corrected by manually eroding the extraction mask
and re-applying it to the anatomical scan when necessary. Image coregistration parameters were
subsequently estimated between an N3 bias-corrected mean BOLD reference and the skull-
stripped T_1 image via `antsRegistration` (including both linear and nonlinear transformations
to minimize residual signal deformations, cf. 'SyN distortion correction' as implemented by the
widely-used fMRIPrep and QSIPrep pipelines).

Subsequent spatial normalization to a standard stereotaxic space presented a nontrivial problem
for patient data—in the absence of a corpus callosum, generic registration routines (with both
linear and nonlinear stages) will frequently pull structures such as the cingulate gyrus into the
space occupied by the CC in the MNI template image, which misaligns brain regions relative to
their location in our functional parcellation. We mitigated this problem in two ways. First, we
implemented cost-function masking during the nonlinear step to minimize the influence of callosal
regions that were transected by the surgery: these masks were specified in the fixed image space
(i.e., the MNI152NLin6Asym template, as provided by the FMRIB Software Library, FSL (*61,*
62)) because we lacked pre-operative data for each patient, and CC segments were defined by the
JHU-DTI white matter atlas (*63, 64*) (also provided in FSL). For patient EM, the genu was masked
out; for patient BT, all but the splenium was masked out; and for all full splits, the entire cross-
section of the callosum was masked out. Second, in addition to these masks, we specified a series
of registration metric objectives at the nonlinear step. As is typical for `antsRegistration`, we

estimated similarity between the MNI template and T_1 images using both ANTs' neighborhood cross-correlation and Mattes mutual information similarity metrics. We also provided smoothed segmentation posteriors derived from the ANTs cortical thickness pipeline (six in total: white matter, cortical gray matter, CSF, deep/subcortical gray matter, brain stem, and cerebellum) and estimated similarity between the MNI template segments and patient segments using a mean-squares objective, with cortical white matter and CSF segments downweighted in importance relative to the other four segments (by a factor of one third). Thus, the nonlinear step was comprised of eight similarity metrics. The transformation was estimated using BSplineSyN to allow for smoother deformations without sacrificing accuracy.

Ultimately, we concatenated and applied the estimated coregistration and spatial normalization parameters to all fMRI data in a single interpolation via `antsApplyTransforms`, resampled into 2 mm^3 isotropic voxels. Registration quality was visually assessed to ensure appropriate mapping to MNI space. Functional images were then smoothed using a 3D Gaussian kernel with FWHM = 5 mm^3. Prior to functional connectivity estimation, several additional preprocessing steps were performed using in-house Matlab code: this included global signal scaling (median = 1000), linear detrending, and nuisance regression. The nuisance regression model followed a '36-parameter' (36P) approach including the six motion regressors from realignment, the first eigenvariate of white matter signal, the first eigenvariate of CSF signal, and mean global signal—along with their squares, their first derivatives (via backwards-differentiation), and the squares of the derivatives. All nuisance regressors were detrended to match the BOLD data. We further augmented the design matrix of the nuisance model with a series of binary spike regressors: one for each image volume where the relative framewise displacement in head motion (accounting for both translation and rotation) was ≥ 0.50 mm. This '36P + spike regression' model has importantly been shown to outperform alternative confound regression strategies (e.g., motion scrubbing and ICA-based methods) and provide the best control for motion artifact in functional connectivity analyses (*17, 18*). Patients generally showed no more head motion than HCP subjects; for reference, we provide a summary of head motion parameters and the frequency of motion spikes for each patient in **Fig. S1**.

Functional connectivity estimation and network laterality
Functional network nodes were defined by parcellating the brain into 400 cortical regions and 32 subcortical regions, according to the Schaefer (*19, 20*) and Tian (*21*) atlases, respectively. We extracted a summary time course from each parcel by taking the first eigenvariate over time (*65*). These regional time series were bandpass-filtered to the range of 0.01-0.08 Hz. We then constructed 432 x 432 FC matrices by estimating the zero-order Pearson correlation between all pairs of regional time series. For both HCP100 subjects and callosotomy patients with more than one resting-state scan, runs were concatenated prior to FC estimation.

We sought to quantify differences in inter-hemispheric functional connectivity (IFC) between callosotomy patients and healthy controls from a number of network neuroscience perspectives. Pursuant to this, we began with an initial, breadth-first assessment of IFC within each of the seven canonical resting-state networks defined by our cortical parcellation (*19, 20*), as well as regions spanning the subcortex (*21*). Similarly to studies of functional lateralization in task-based fMRI, we derived a *network laterality index* as a simple means of summarizing and quantifying the *relative magnitude* of IFC as compared to the magnitude of intra-hemispheric FC. Each FC matrix

was first thresholded at an uncorrected level of $p < .001$ to avoid overly-conservative and highly-
nonlinear effects of multiple comparisons correction on the distribution of edge weights, while
simultaneously minimizing the influence of weak (potentially spurious) connections. We then
Fisher-Z transformed the thresholded FC matrices and, for each individual and for each network,
estimated a laterality index, LI, as:

$$LI = \frac{(\sum FC_{Intra-LH} + \sum FC_{Intra-RH}) - \sum FC_{Inter}}{\sum FC_{Intra-LH} + \sum FC_{Intra-RH} + \sum FC_{Inter}}$$

Intra-hemispheric connections within the left and right hemispheres are necessarily summed in the
numerator given that there are, by definition, twice as many inter-hemispheric connections as each
hemisphere alone. The result is a ratio on the interval of [-1, 1] that captures the extent to which
each canonical resting-state network is predominantly connected intra- or inter-hemispherically:
in the limit, a value of -1 would suggest that there is *no* intra-hemispheric FC and the network is
solely comprised of inter-hemispheric connections, while a value of 1 would indicate the reverse.
A value of 0 therefore represents a network that is perfectly balanced in the magnitude of intra- vs.
inter-hemispheric connections. Note that the results presented in **Fig. 3** are with respect to
positively-correlated edges only; we provide a comparison of anti-correlated edge lateralization in
Fig. S3.

<u>Community detection and modularity</u>
While the above approach is useful for considering differences among canonical functional
networks, these systems were defined in healthy adult brains—it is therefore perhaps dubious to
assume that these networks follow a similar spatial organization in callosotomy patients. Data-
driven community detection techniques afford assessments of functional network topology
unbeholden to prior assumptions about specific brain regions and their canonical network
memberships (*66, 67*). This is achieved via decomposition of FC matrices using *modularity*
maximization algorithms, which group regions into *functional modules*: these represent regions
that are generally more functionally-coupled to each other than they are connected to other regions.
The optimization objective is subject to a *quality* metric, *Q*—unlike our previous laterality analysis
(which required separating out positive and negative edges), this can be estimated over FC matrices
comprised of both positive and negative correlations (*68*). We performed this decomposition using
the `community_louvain` function from the Brain Connectivity Toolbox
(https://sites.google.com/site/bctnet/) with asymmetric treatment of negative edge weights (*68,*
69). Importantly, modularity maximization is sensitive to spatial scale, subject to a *resolution*
parameter, γ (*70*). Small values of γ typically yield coarse partitions with a small number of large
communities; larger values of γ, by contrast, afford more fine-grained partitions. Because the
solution of this decomposition is dependent upon one's choice for the resolution parameter, we
tested a range of values from $\gamma = [0.10, 2]$ (in steps of 0.10). To further account for variability in
community assignments (e.g., due to algorithmic stochasticity arising from random initializations),
we estimated modularity 1000 times for each value of γ—optimal consensus partitions were
obtained using the `consensus_similarity` function from the Network Community Toolbox
(https://commdetect.weebly.com/).

In similar fashion to the network laterality indices described above, we derived a *module laterality*
index for the resulting partitions according to prior work (*25*). This metric is limited to the [0, 1]

interval, where a value of 0, for example, indicates that all functional communities are perfectly
bilateral (i.e., all of the constituent brain regions belonging to each community are equally-
represented between the left and right hemispheres), and values closer to 1 indicate progressively
more lateralized topologies. We note that in this case, more fine-grained partitions may be
beneficial for assessing modular laterality—nevertheless, the results shown in **Figs. 4-5** reflect a
'default' $\gamma = 1$ and we further illustrate in **Fig. S5** that full splits rapidly diverge from controls even
at smaller values of γ (while partial splits and healthy controls remain relatively consistent,
regardless of γ).
799
<u>Inter-hemispheric network dynamics</u>
Finally, we used edge time series (ETS) (*27*) to assess the extent to which *many* brain regions in
the left and right hemispheres dynamically co-fluctuate in synchrony. The ETS approach has been
used to obtain moment-to-moment, time-resolved estimates of FC and characterize the nature of
high-amplitude events (*28*) that occur when multivariate BOLD signals simultaneously fluctuate
together over brief intervals of time—these bursts of synchronized network dynamics are sufficient
to recapitulate 'static' FC estimates over the course of a scan (*28*), and importantly, they are
independent of head motion (*27*). Current evidence instead suggests these high-amplitude co-
fluctuations are governed, in part, by the modular organization of the brain and its underlying
structural architecture (*26*). An ETS can be computed by taking a pair of BOLD time series from
two regions, *Z*-scoring them, and taking the elementwise product of the signals at each timepoint—
the mean of an ETS is equivalent to the zero-order Pearson correlation between the two time series,
and thus ETS constitute a temporal-unwrapping of FC over the scan (*27*).
813
Here, we first separated out all regions in the left hemisphere and all regions in the right
hemisphere. We constructed a *T* x 23220 ETS matrix for each hemisphere, where *T* gives the
number of timepoints and 23220 gives the total number of edges within each hemisphere. A
summary time course quantifying the total amplitude of intra-hemispheric co-fluctuations at any
given moment (i.e., across *all* edges) was obtained by taking the root-sum-of-squares (RSS) at
each timepoint (*28*). We then computed the Spearman correlation between the left hemisphere RSS
and right hemisphere RSS time series to capture the extent to which network dynamics in each
hemisphere were synchronized.

Patient	Scan num.	Scan duration		Head motion estimates					
		Minutes	Frames	Min FWD	Max FWD	Med FWD	MAD FWD	Num. spikes	Spike %
EM	1	8	240	0.03	2.44	0.13	0.05	9	3.75
BT	1	7	210	0.04	1.09	0.19	0.07	11	5.24
BT	2	7	210	0.05	1.34	0.25*	0.08	9	4.29
HY	1	12	360	0.02	0.44	0.14	0.05	0	0.00
DT	1	12	360	0.02	10.74*	0.15	0.05	31	8.61*
TJ	1	7	210	0.02	0.59	0.13	0.04	2	0.95
TJ	2	7	210	0.03	0.38	0.11	0.04	0	0.00
LJ	1	7	210	0.06	0.55	0.20	0.04	2	0.95
LJ	2	7	210	0.04	0.38	0.18	0.04	0	0.00
LJ	3	7	210	0.06	0.56	0.20	0.05	3	1.43

*Denotes head motion estimate exceeding normal range, see corresponding figure(s) below.

Fig. S1. Scan times and head motion estimates across callosotomy patients. Due to differences
in clinical scheduling and other imaging protocols not relevant to the current analysis, patients
varied in resting-state fMRI scan time. Here we summarize these differences and provide various
estimates of head motion for each scan. Motion parameters are given with respect to *framewise*
displacement (FWD)—the amount of total head motion (in millimeters, considering both
translational displacements and rotations) observed from one image volume to the next. Because
FWD is a time series, we provide the minimum, maximum, and median FWD over each scan (with
the median absolute deviation, *MAD*, over time). We also denote the number of motion spikes,
defined as any frame with FWD $\geq$ 0.50 mm, and the percentage of frames contaminated with
motion spikes—these were used to define individual spike regressors for nuisance signal
regression (*17, 18*). Patients generally did not show any more head motion than healthy adult
controls (see visualized distributions). Note that in rare cases where patients exceeded the range
of controls in one metric, they were typically within the standard range of other metrics. Point
intervals under the control distributions give the median (black dot) and the 80-95% quantiles
around the median (thick black lines and thin black lines, respectively).

Spatial pattern similarity to controls (Dice coefficient)

Fig. S2. Multivariate similarity in spatial patterns of functional connectivity. As an initial estimate of the extent to which full vs. partial callosotomy disrupted intra- vs. inter-hemispheric functional connectivity (FC), we binarized FC matrices at a threshold of $z(r) > 0$ (where $z(r)$ refers to the Fisher-Z transformed correlation coefficients for each network edge) and computed the Dice coefficient between each patient and each individual healthy control—this statistic ranges from [0,1] and quantifies the degree of similarity in two patterns of FC (liberally, whether the same edges show *any* FC > 0, regardless of the exact magnitude of those connections). The distributions for each patient thus show the range of similarity between a given patient and each control. Patient EM (with the most corpus callosum intact) is generally most similar to controls; full splits, on the other hand, show markedly dissimilar patterns of inter-hemispheric FC relative to intra-hemispheric FC. Point intervals under each distribution give the median Dice similarity (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: LH, Left Hemisphere; RH, Right Hemisphere.

Fig. S3. Anti-correlated functional connectivity is generally not more lateralized in callosotomy patients. Network laterality indices across seven canonical resting-state systems (*19, 20*) and subcortical regions (*21*). To contrast against **Fig. 3**, here we estimate lateralization of negative (anti-correlated) connections. Values closer to 0 indicate an equal balance between intra- and inter-hemispheric (negative) FC, while values closer to -1 or 1 indicate more inter-hemispheric FC intra-hemispheric (negative) FC, respectively. Note that, unlike in **Fig. 3**, most distributions skew to the left (indicating that negative FC is predominantly inter-hemispheric, i.e., not lateralized), and both full and partial splits frequently fall within the range of HCP controls. Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: FC; Functional Connectivity; SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Fig. S4. Estimates of large-scale dynamic functional synchrony by network. Decomposing whole-brain comparisons of RSS synchrony in **Fig. 6b** to individual networks reveals nuanced differences between patients and neurotypical adults (colored distributions). As before, partial splits consistently fall within the range of HCP controls. Full splits generally show more desynchronized network dynamics in higher-order cognitive systems, but comparatively-neurotypical levels of inter-hemispheric synchrony in limbic and subcortical networks (which are not strictly dependent upon structural connections through the corpus callosum, also reflecting **Fig. 3**). Point intervals under the control distributions give the HCP100 median (black dot) and 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: RSS, Root-Sum-Of-Squares (see **Materials and Methods**); SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Fig. S5. Lateralization of functional modules is readily-apparent in full splits regardless of resolution. Modularity maximization algorithms are sensitive to a spatial resolution parameter, γ (*70*). A value of γ =1 is often treated as a 'default' level of resolution, whereas values of γ < 1 or γ > 1 produce progressively more coarse or fine-grained solutions, respectively. Here we show that differences in modular laterality between patients and healthy adult controls are not strictly-dependent upon the chosen value of γ. We performed 1000 iterations of modularity maximization at each of 20 levels of γ (ranging from [0.10, 2] in steps of 0.10), derived an optimal consensus partition at each level (see **Materials and Methods**), and estimated a laterality index for the resulting partitions (*25*) (such that values closer to 1 indicate more lateralized network topologies). Full splits (solid-colored lines) rapidly diverged from the distribution of healthy controls (quantile intervals shown in gradations of purple, with the solid black line representing the median) demonstrating more lateralized organization of functional modules, even at coarse levels of γ < 1. Partial splits (dotted colored lines), by contrast, consistently remained within the control distribution.