Full inter-hemispheric integration sustained by a fraction of posterior callosal fibers

Tyler Santander[1,2,*], Selin Bekir[2], Theresa Paul[3], Jessica M. Simonson[1,2], Valerie M. Wiemer[3], Henri Etel Skinner[2], Johanna L. Hopf[4], Anna Rada[4], Friedrich G. Woermann[4,5], Thilo Kalbhenn[4,6], Barry Giesbrecht[1,2], Christian G. Bien[4], Olaf Sporns[7], Michael S. Gazzaniga[2], Lukas J. Volz[3,†], & Michael B. Miller[1,2,†,*]

[1]Institute for Collaborative Biotechnologies, University of California, Santa Barbara, Santa Barbara, CA, USA

[2]Department of Psychological & Brain Sciences, University of California, Santa Barbara, Santa Barbara, CA, USA

[3]Medical Faculty, University of Cologne, and Department of Neurology, University Hospital Cologne, Cologne, Germany

[4]Department of Epileptology, Krankenhaus Mara, Bethel Epilepsy Center, Medical School OWL, Bielefeld University, Bielefeld, Germany

[5]MRI Department, Society for Epilepsy Research, Krankenhaus Mara, Bethel Epilepsy Center, Bielefeld, Germany

[6]Department of Neurosurgery, Evangelisches Klinikum Bethel, Medical School OWL, Bielefeld University, Bielefeld, Germany

[7]Department of Psychological & Brain Sciences, Indiana University, Bloomington, IN, USA

[†]Authors contributed equally

*To whom correspondence should be addressed:

Tyler Santander (tsantander@ucsb.edu)

Michael B. Miller (mbmiller@ucsb.edu)

Acknowledgements

T.S., S.B., J.M.S., H.E.S., B.G., and M.B.M. were sponsored by the Army Research Office under

contract W911NF-19-D-0001 for the Institute for Collaborative Biotechnologies. T.P. and L.J.V

were funded by the German Research Foundation (CRC-1451, Project 431549029).

Author contributions

T.S., S.B., T.P., H.E.S., J.L.H., A.R., B.G., C.G.B., O.S., M.S.G., L.J.V., and M.B.M. conceived

and planned the experiments. T.P., V.M.W., J.L.H., A.R., F.G.W., C.G.B., and L.J.V. assisted with

patient recruitment and were responsible for data collection. T.K. performed the callosotomy

surgeries. T.S., J.M.S., O.S., and L.J.V. conceived the data analysis strategy; T.S., J.M.S., and O.S.

contributed to analysis code and executed the data processing/analysis pipeline. T.S. wrote the

manuscript with contributions from T.K., O.S., M.S.G., L.J.V., and M.B.M. The project was

supervised by A.R., C.G.B., L.J.V., and M.B.M. Funding was secured by B.G., L.J.V., and M.B.M.

All authors assisted with manuscript revision and approved the final submitted version. L.J.V. and

M.B.M. contributed equally.

Conflicts of Interest

The authors declare no competing interests.

Abstract

The dynamic integration of the lateralized and specialized capacities of the cerebral hemispheres constitutes a hallmark feature of human brain function. This inter-hemispheric exchange of information critically depends upon the corpus callosum. Classical descriptions of callosal organization outline a topographic gradient, such that specific fibers integrate distinct aspects of brain function. Here we present a challenge to this conventional model. Using neuroimaging data obtained from a new cohort of adult corpus callosotomy patients, we leverage modern network neuroscience techniques to show—for the first time—that full inter-hemispheric integration can be achieved via a small proportion (~1 cm) of intact posterior callosal fibers. Only complete callosotomy patients demonstrated the expected dissolution of typical inter-hemispheric network architectures, aligning with disconnection syndromes long-thought to reflect diminished information propagation and communication across the brain. These findings motivate a novel mechanistic understanding of synchronized inter-hemispheric neural activity for large-scale human brain function and behavior.

Main

In recent decades, network neuroscience has emerged as a powerful analysis framework for modeling functional interactions between remote neural systems. A foundational assumption of this approach is that anatomical links between brain regions constrain neuronal dynamics and support the emergence of large-scale functional networks, which in turn enable the efficient integration of perceptual, motor, and higher-order cognitive information necessary for human behavior[1,2,3,4,5,6]. The corpus callosum (CC) is thought to be the central conduit through which specialized processing nodes in each cerebral hemisphere coordinate and synchronize function, with 200-250 million axons enabling direct, long-range communication across the brain. Classical topographic models of CC projections have revealed a well-defined anterior-to-posterior anatomical gradient, corresponding to the spatial organization of canonical brain networks supporting various sensorimotor and cognitive functions[7,8]. One consequence of this view is the prediction that damage to the CC should result in drastic functional disconnections between hemispheres—with disruptions following the localization of lesioned fibers along the aforementioned anatomical gradient.

Split-brain patients, in whom the CC has been severed in an effort to treat medically-intractable epilepsy, offer a powerful means of testing this core conceit of structure-function correspondence. However, neural data on these patients remain scarce—callosotomy was largely discontinued around the turn of the 21st century, coinciding with the rise of network neuroscience. Extant data largely stem from anesthetized pediatric patients[9,10,11,12]. Taken together, these studies support the hypothesis that full versus partial callosotomy confer differential effects with respect to the location and extent of functional disconnections. At the same time, however, they have provided contradictory findings regarding recovery of function post-operatively[9,11,12], and various other methodological factors obfuscate more straightforward interpretations. Thus, to date, it remains unknown how severing the largest fiber bundle of the human brain affects the topological organization and intrinsic dynamics of large-scale functional networks.

To address this fundamental question, we present the first investigation characterizing the impacts of full and partial callosotomy on the adult human brain using multiple network neuroscience

methods. We hypothesized that full callosotomy would wholly disrupt functional synchrony across

the cerebral hemispheres, while the spatial distribution of functional disconnections following

partial callosotomy would unveil the anatomical gradient of the CC. As predicted, we found that

complete sectioning of the CC broadly obliterated both typical inter-hemispheric network

topologies and dynamic patterns of inter-hemispheric temporal synchrony. More strikingly,

however, we also found that a small proportion of intact posterior CC fibers (a case of partial

callosotomy) was sufficient to support widespread inter-hemispheric coupling. These findings

challenge our conventional understanding of the CC's functional topography and sheds new light

on the critical mechanisms for inter-hemispheric information integration.

Patient descriptions

Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were tested post-

operatively at the Bethel Epilepsy Center (Bielefeld, Germany). The time between surgery and

testing varied between one and six years (**Table 1**). In two patients (EM and BT), surgery did not

result in a complete callosal transection; they are hereafter referred to as *partial splits*. In patient

EM, the surgery was stopped after sectioning the anterior third of the CC; patient BT was spared

approximately 1 cm of splenium. The remaining four patients (HY, DT, TJ, and LJ) were

completely sectioned and are hereafter referred to as *full splits*. All patients have intact anterior

and posterior commissures regardless of the extent of the callosal section.

Figure 1. Patient anatomical scans. Post-operative T_1 and diffusion MRI scans collected for all

six patients at time of testing. White arrows/ellipses for patients EM and BT highlight intact CC

(i.e., partial splits). The remaining four patients were fully-sectioned. Note that a series of clinical

observations suggested behavioral disconnection syndromes in all full splits, but not in partial
splits.

Table 1. General overview of callosotomy patients.

Patient	Age at surgery	Age at testing	Sex	Handedness	Extent of callosotomy	Disconnection syndrome
EM	17	19	M	Right	Partial	Absent
BT	22	28	M	Right	Partial	Absent
HY	21	22	M	Right	Complete	Present
DT	15	20	M	Right	Complete	Present
TJ	49	52	F	Right	Complete	Present
LJ	57	60	M	Right	Complete	Present

Callosotomy is not commonly associated with post-operative neuropsychiatric or neurological
disorders. Rather, split-brain patients often demonstrate *disconnection syndromes*, such that they
appear to have two hemispheres operating independently, unable to communicate information
from one half of the brain to the other[13,14,15]. Here, behavioral disconnection syndromes were
clinically-observed using simple, lateralized bedside testing protocols. Comprehensive
descriptions of these tests are described elsewhere[16]. In brief, all four complete callosotomy
patients clearly indicated behavioral disconnections. Partial splits, in contrast, did not show *any*
signs of disconnection syndromes—including, remarkably, patient BT (with a mere 1 cm of
splenium). These findings contradict our classical understanding of callosal topography, which
posits functionally-specific deficits depending upon which portions of the CC were severed.
Consequently, a small proportion of posterior callosal fibers may be sufficient to support
information integration across large-scale inter-hemispheric brain networks.

Figure 2. Inter-hemispheric functional connectivity is broadly-diminished in full (but not partial) splits. a, Whole-brain functional parcellation comprised of seven canonical resting-state networks[17,18] and subcortical nodes[19]. **b,** A standard reference for neurotypical whole-brain functional connectivity (FC), given as the median over HCP100 subjects. FC was estimated by correlating low-frequency brain activity (0.01-0.08 Hz) between all pairs of regions in **(a)**. The

large boxes outlining the upper left and lower right quadrants denote *intra-hemispheric* FC; off-
diagonal quadrants thus capture *inter-hemispheric* FC (IFC). **c,** Whole-brain FC estimates for each
callosotomy patient, thresholded at $p < .001$ (uncorrected for multiple comparisons). Typical
patterns of IFC were largely abolished in full splits (HY, DT, TJ, LJ), while partial splits (EM, BT)
were more similar to healthy controls. Abbreviations: SomMot, SomatoMotor Network; DorsAttn,
Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network; LH, Left
Hemisphere; RH, Right Hemisphere.

Inter-hemispheric functional connectivity after callosotomy
All patients underwent resting-state functional magnetic resonance imaging without sedation,
affording whole-brain analyses of resting-state functional connectivity (FC) and quantification of
inter-hemispheric FC (IFC). To benchmark patients against 'neurotypical' patterns of IFC, we used
data from the Human Connectome Project 100 Unrelated Subjects (HCP100) cohort. All HCP100
subjects and callosotomy patients underwent an identical preprocessing pipeline, including
conservative corrections for mitigating head motion artifacts[20,21] (see **Extended Data Fig. 1** for a
quantitative description of patient head motion). Whole-brain FC in the range of 0.01-0.08 Hz was
assessed by parcellating the brain into 400 cortical regions and 32 subcortical regions (according
to the Schaefer[17,18] and Tian[19] atlases, respectively; **Fig. 2a**) and correlating all pairs of regional
time series (**Figs. 2b-c**).

All full splits demonstrated marked reductions in IFC relative to both neurotypical controls and
partial splits (**Figs. 2b-c**). Spatial patterns of intra-hemispheric FC, by contrast, were generally
better-matched to controls (**Extended Data Fig. 2**), suggesting that callosotomy primarily
impaired inter-hemispheric functional coupling with full splits suffering the greatest disruptions.
Curiously, however, while gross IFC was dissolved in full splits, inspection of individual FC
matrices suggested that visual network connectivity was partially preserved (**Fig. 2c**). This may be
due to synchronized perceptual inputs (scans were acquired with eyes open), driving *apparent* IFC
despite the lack of transcallosal connectivity between visual regions and evidence for behavioral
disconnections. We therefore sought to assess differences in the magnitude of IFC between
functional homologues to determine whether any preserved IFC was unique to sensory systems.
Here we computed a series of *network laterality indices*, where a value of zero indicates a perfect

balance between inter- and intra-hemispheric FC, and values closer to -1 or 1 indicate relatively more inter- or intra-hemispheric FC, respectively. Note that we focus here on the distribution of positively-coupled edges (anti-correlated edges are shown in **Extended Data Fig. 3**).

Figure 3. Full callosotomy is associated with more lateralized functional connectivity. Network laterality indices across seven canonical resting-state systems[17,18] and subcortical regions[19]. Values closer to zero indicate an equal balance between intra- and inter-hemispheric FC, while values closer to one indicate more intra-hemispheric FC relative to inter-hemispheric FC. Note that each subplot captures a unique range of values, given differences in variability observed across neurotypical adults (colored distributions), patients, and networks. Full splits generally demonstrated more strongly-lateralized FC compared to partials, who fell within the control distributions. Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Consistent with observed behavioral disconnections, all full splits demonstrated more lateralized

patterns of FC relative to healthy controls and partial splits, who typically fell within the control

distributions (**Fig. 3**). This was evident in both higher-order, transmodal systems (e.g., the default

mode and dorsal attention networks) and in somatosensory networks (with the exception of patient

TJ who fell within the upper tail of the control distribution in the somatomotor network). Patient

EM was more lateralized across frontoparietal control nodes, which include a number of regions

likely to be structurally-disconnected given the extent of their callosotomy; however, BT generally

showed degrees of network bilaterality equivalent to healthy adults, despite having a mere fraction

of splenium intact. Finally, full splits appeared to show conserved bilateral FC across limbic

regions like the orbitofrontal cortex and temporal poles, which are known to be structurally

coupled through the preserved anterior commissure as well as the CC[22,23,24]. In sum, these results

highlight the dependence of inter-hemispheric networks on CC connectivity—yet, neurotypical

levels of functional integration were possible even when only the splenium was left intact.

Lateralized expressions of functional modules

To further elucidate whole-brain differences in functional network topology—agnostic to any *a*

priori assignment of individual brain regions to canonical functional systems—we employed data-

driven community detection techniques to cluster whole-brain patterns of FC into functional

modules (**Fig. 4**). The modular organization of both structural and functional brain networks is

thought to enable specialized information processing while retaining the ability to efficiently

integrate information between disparate neural systems[25,26]. Thus, we hypothesized that the

breakdown of IFC in full splits would produce lateralized network topologies, predominantly self-

organized within one hemisphere or the other.

Here we computed laterality indices for the estimated partitions according to prior work[27]: where

a value of zero, for example, indicates that all functional modules are perfectly bilateral (i.e., the

constituent brain regions belonging to each community are equally-balanced between the left and

right hemispheres), and values closer to one indicate progressively more lateralized topologies.

Importantly, we note that modularity maximization is sensitive to spatial scale, subject to a

resolution parameter, γ[28]. We describe results using a 'default' $\gamma = 1$ below but provide further

consideration of these issues in the **Methods** and **Extended Data Fig. 4**.

Figure 4. Modular decomposition of whole-brain functional connectivity. Spatial maps of optimal, consensus community assignments derived via data-driven modularity maximization at a default level of resolution ($\gamma = 1$). Regional coloring indicates assignment to a given functional module. Full splits generally exhibited more lateralized clustering of regions into modules relative to partial splits and healthy adult controls (see **Fig. 5** for quantification of module laterality).

As predicted, we observed much greater degrees of modular laterality in all full splits (**Fig. 5**). Communities with bihemispheric representations primarily spanned visual and somatomotor regions—reflecting a mechanism through which synchronized, external sensory inputs can drive

apparent functional integration. With respect to partial splits, patient EM with the most CC showed bilateral patterns of network organization mirroring neurotypical controls. Patient BT also fell within the control distribution—however, they showed a strikingly different pattern of modularity (**Fig. 4**). Here, functional communities appeared to dissociate along the Sylvian fissure, with dorsal modules and ventral modules spanning the anteroposterior axis of the brain. This individual had the longest gap between surgery and testing (~6 years), which may have allowed for widespread network reorganization affording optimal utilization of the minimal callosal fibers left intact. These findings underscore the importance of transcallosal connections for the emergence of coherent inter-hemispheric systems, while further highlighting a role for sensory inputs enabling apparent preservation of bilateral sensorimotor modules.

Figure 5. Functional modules are more strongly-lateralized in full splits. The modular laterality index quantifies the extent to which functional communities exhibit bihemispheric representations (values closer to zero) or are restricted to one hemisphere or the other (values closer to one). At a default level of resolution ($\gamma = 1$), partial splits demonstrated bilateral network topologies similar to healthy controls, whereas full splits were consistently more lateralized. Point intervals under the control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively).

Inter-hemispheric synchrony in network dynamics

The modular organization of the brain plays a critical role in governing large-scale functional
dynamics, such that structural architectures orchestrate widespread co-fluctuations in whole-brain
activity over time[29]. Such patterns of coordinated network dynamics have recently been elucidated
using *edge time series* (ETS): in brief, the ETS approach constitutes a temporal unwrapping of the
zero-order Pearson correlation between any given pair of brain regions, capturing moment-to-
moment changes in the extent to which regional pairs are co-fluctuating in activity[30]. Taking the
root-sum-of-squares (RSS) across all ETS at each timepoint yields summary estimates of
multivariate co-fluctuations in functional network dynamics over the course of a scan[31] (**Fig. 6a**).
291
Here we sought to test the hypothesis that structural connectivity via the CC is necessary for large-
scale coordination of function. To that end, we separated edges within the left and right
hemispheres and derived an RSS time course for each. We then computed correlations between
left and right RSS to assess the degree to which dynamic co-fluctuations in brain function were
synchronized across hemispheres. Among neurotypical controls, we observed tight temporal
coupling between left and right hemisphere dynamics (Median Spearman $r = 0.82$, $MAD = 0.08$).
In contrast, all four full splits demonstrated significant reductions in inter-hemispheric synchrony,
suggesting that network dynamics were more functionally-independent (**Fig. 6b**; see **Fig. S4** for
individual networks). This is consistent with the overall attenuation of inter-hemispheric FC and
further indicates that brief moments of high-amplitude co-fluctuations (which normally encompass
regions in both hemispheres and are thought to be structurally-mediated[29]) occur separately in
each. Notably, both partial splits fell within the standard range of the HCP subjects. These results
therefore strongly suggest a mechanistic role for CC structural connections in supporting
widespread integration of inter-hemispheric network dynamics. This may be achieved with a mere
fraction of intact fibers through the splenium and thus might be largely-independent of the presence
of direct cortico-cortical connections.
308

Figure 6. Large-scale functional network dynamics are desynchronized in full splits. **a,** An illustration of edge time series in the left and right hemispheres for patient BT (a partial split with 1 cm of intact splenium) and patient TJ (a full split). By taking the root-sum-of-squares (RSS) across all edges at each timepoint, we derived a summary estimate of the total amplitude of co-fluctuating brain activity. Note the bursty, high-amplitude co-fluctuations in each hemisphere that often coincide in time for BT, but not in TJ. **b,** All four full splits exhibited reduced temporal synchrony (i.e., temporal correlations) between left/right hemisphere RSS time courses. Partial splits, by contrast, fell within the range of healthy adult controls. Point intervals under the control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively).

Discussion

Distinct disruptions of inter-hemispheric functional connectivity were observed following complete versus partial transection of the corpus callosum. Full splits demonstrated more lateralized network topologies and desynchronized network dynamics, aligning with classical behavioral disconnection syndromes. Given that all patients had intact anterior and posterior commissures (along with other subcortical routes of inter-hemispheric communication), these findings underscore the notion that transcallosal connections are the essential pathways through which large-scale inter-hemispheric functional networks emerge. Contrary to our expectations,

however, inter-hemispheric functional coupling among partial splits remained intact. These
patients also showed *no* signs of behavioral disconnection, even when the vast majority of the
callosum was severed[16]. Thus, a fraction of callosal fibers in the splenium may be sufficient to
orchestrate inter-hemispheric functional synchrony and integrate behavior. This poses a
remarkable challenge to our classical topographic models of callosal connectivity and function,
motivating a novel view on inter-hemispheric structure-function relationships: namely, that robust
inter-hemispheric synchrony and information transfer can be achieved via polysynaptic
connectivity through a relatively small proportion of posterior transcallosal connections, rather
than direct structural connectivity between functional homologues.

Complete callosotomy and structure-function relationships
Split-brain patients present a unique opportunity to draw causal inferences regarding the role of
transcallosal structural connections and the extent to which inter-hemispheric functional
synchrony depends upon such connections. Previous investigations of pediatric callosotomy
patients have provided an initial perspective on this question[9,10,11,12]. However, these patients often
suffer from severe cognitive and developmental deficits, requiring scans to be completed under
sedation—which reduces dynamical complexity in brain activity and produces functional
connectomes more rigidly tied to direct anatomical pathways[32]. It is further known that functional
and structural brain networks undergo profound developmental changes early in life[33,34,35,36]. In
other words, neural plasticity in these young individuals may still afford the long-term
establishment of alternate inter-hemispheric pathways (perhaps akin to callosal agenesis
patients[37]), and indeed, it is well-documented that disconnection syndromes in pediatric patients
resolve over time[38,39]. This likely accounts for differential reports of recovery in inter-hemispheric
FC among these individuals[9,11,12].

The adult human brain also retains neuroplastic potential[40]. However, given the enormity of the
structural disconnection, adult callosotomy patients may lack the capacity to adapt their functional
architectures in such a way that allows complete recovery of communication between disconnected
brain regions. Our results clearly underscore this hypothesis. Whole-brain patterns of FC appeared
highly lateralized, both with respect to canonical network definitions (**Figs. 2-3**) and the
topological organization of functional modules (**Figs. 4-5**). Furthermore, we observed considerable

reductions in the extent to which multivariate co-fluctuations in activity were synchronized

between hemispheres (**Fig. 6**), underscoring a role for the CC in orchestrating whole-brain network

dynamics. Importantly, however, there were some exceptions to these effects, which follow

anatomical predictions. Limbic and subcortical regions—the former of which are known to be

connected through the anterior commissure (AC) as well as the CC[22,23,24]—showed conserved

bilateral FC (**Fig. 3**). Nevertheless, non-callosal pathways were insufficient to support widespread

inter-hemispheric synchrony, contrary to previous proposals both in humans[41,42,43] and nonhuman

primates (where AC fibers are sufficient to sustain inter-hemispheric FC)[44]. Finally, visual and

sensorimotor areas also appeared to remain somewhat integrated. We cannot fully discount the

possibility that subcortical information transfer contributed to this preserved synchrony; however,

an alternative hypothesis may lie in the fact that visual and somatosensory inputs to each

hemisphere were synchronized during the resting-state scan—producing *apparent* functional

coherence despite observed behavioral disconnections.

Preserved inter-hemispheric integration in partial splits

While our observations in full splits clearly emphasize the importance of CC connectivity in

orchestrating inter-hemispheric synchrony, partial splits revealed something much more

surprising. We predicted that partial splits would show specific deficits in inter-hemispheric FC

along an anterior-to-posterior gradient localized to the extent of the callosal transection, following

both previous characterizations of CC topography[7,8] and prior behavioral studies[45,46,47,48].

However, both partial splits consistently demonstrated levels of inter-hemispheric integration in

the range of neurotypical controls (**Figs. 2-6**). This was even observed for patient BT, in whom

only ~1 cm of posterior callosal fibers were spared. It has become increasingly-evident that

structure-function relationships are highly complex and generally do not reflect simple, one-to-

one interactions between directly-coupled areas[49,50,51]. Still, these remarkable results imply that the

functional topography of the CC may be much more flexible than previously-imagined. In

particular, there may be a critical threshold at which full functional integration can be sustained

given a sufficient proportion of intact fibers—even if the vast majority of the callosum has been

severed.

An intriguing extension of this hypothesis is that the *location* of the spared callosal fibers may be
critical. Previous cytological work has shown that anterior portions of the CC are primarily
populated by thin fibers, while fibers in the tail locally-vary in diameter[52,53,54]. Such differences in
fiber size and myelination along the CC have been associated with differences in conduction
velocity[54] and, when considered alongside the topographic organization of the CC, may account
for the 'fast' versus 'slow' action of primary sensorimotor and association cortices, respectively.
Spared posterior fibers as in BT could therefore be more beneficial than spared anterior fibers.
However, from a sheer bandwidth perspective, it is difficult to imagine how such a small
proportion of fibers could account for widespread integration of function and cognitive processing.
400
It appears likely that functional systems in BT dramatically reorganized in the six years following
surgery. In brains that develop normally with the support of the CC, multisensory integration zones
in parietal cortex may serve as inter-hemispheric hubs[55,56]. By taking advantage of spared posterior
callosal pathways connecting these regions, the processing of various intra-hemispheric brain
regions can still be integrated and exchanged along polysynaptic routes running between
hemispheres. The striking differences in functional network topology observed in BT, with large
modules spanning the anterior-to-posterior axis of the brain (**Fig. 4**), are one potential consequence
of this reorganization, reflecting greater functional dependencies between frontal and posterior
areas affording more optimal usage of the spared splenial fibers. The fundamental importance of
these posterior inter-hemispheric connections also aligns with and may further inform
neuroscientific theories of consciousness[57,58]. While our results are not suited to evaluate or
endorse any particular theory, it is nevertheless curious that the temporal-parietal-occipital 'hot
zone' of regions recently proposed to maximize the singular 'here-and-now' *qualia* of
consciousness[57,59] are likely dependent (in part) upon CC connections through the splenium. This
may explain why BT maintains an integrated conscious experience while full splits demonstrate
inter-hemispheric disintegration[16].
417

Limitations
We acknowledge the small number of patients included in this study, each of whom have limited
post-operative data. While previous pediatric studies have recruited larger samples, the availability
of patients who underwent callosotomy in adulthood is extremely scarce. It is often difficult to

draw robust and reliable statistical inferences from neuroimaging data acquired on such small

samples. However, the marked differences observed between full splits and healthy controls

suggest that we are not parsing trivial effects on the scale of traditional neuroimaging—rather, they

reflect outcomes drawn from fundamentally unique sampling distributions. Furthermore, relative

to other lesion studies (often hampered by variability in spatial localization/extent), callosotomy

is a highly-selective intervention. By severing a single white matter structure in the brain without

any damage to gray matter, callosotomy may afford more homogeneous outcomes. It will be

critical for future studies to include larger samples with more imaging data to better characterize

the extent of neural variability in callosotomy patients, especially as network architectures

reorganize over time.

Conclusion

Following more than a half-century of behavioral neuroscience, split-brain patients continue to

provide startling insights into the fundamental nature of the human brain. The rise of network

neuroscience has further transformed our understanding of large-scale brain systems, tracing the

pathways relating brain structure to function and behavior. Our findings reveal a unique type of

criticality in brain function—such that a vanishingly-small proportion of posterior callosal fibers

may be sufficient to support the integration of inter-hemispheric functional brain networks and

unified cognitive processing. It has long been celebrated that the human brain is a remarkably

adaptable complex system. These findings challenge us to rethink our conventional assumptions

about structure-function relationships and the means through which the brain enables the mind.

Online Methods

Patient recruitment

Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were recruited

during routine post-operative follow-up visits to the Bethel Epilepsy Center (Bielefeld, Germany).

The study was approved by the Ethics Committee of the University of Münster,

Germany (2021–523-f-S). The patients or their legal representatives provided written informed

consent.

Callosotomy procedure

Callosotomies were performed with the assistance of intraoperative neuronavigation, with patients in a supine position and the head inclined by roughly 30° (fixed in the Mayfield clamp). A straight skin incision was made running slightly next to the midline on the non-language dominant hemisphere. Craniotomy was then performed by making one drill hole in the midline at approximately the level of the precentral sulcus. A bone flap was sawn out with an outline of 3 cm x 5 cm. The dura was incised longitudinally and then opened with two wings under microneurosurgical conditions—access to the inter-hemispheric fissure was successively created while preserving the cortical bridging veins. Next, using a self-retaining retractor to carefully shift the frontal lobe, the level of the corpus callosum was gradually exposed under sharp dissection with micro scissors.

Dissection is often made considerably more difficult by the course of the pericallosal arteries, which run parallel to the corpus callosum but are often tightly intertwined and twisted in the inter-hemispheric fissure. In cases where the risk of vascular damage is too great, the surgery may be halted short of a full disconnection (resulting in partial splits). However, under most conditions, the inter-hemispheric fissure may be opened along the entire length of the knee of the corpus callosum. The corpus callosum was then perforated into the lateral ventricle of the ipsilateral hemisphere. Next, following the pericallosal artery and the anterior cerebral artery down to the frontal base, the knee of the callosum was split in the midline, continuing backwards to about the level of the foramen of Monroi. From this point, it is critical to protect the fornix of the language-dominant hemisphere: to mitigate the risk of postoperative functional impairment, we carefully ensure that the self-retaining retractor is not placed on the precentral gyrus within the inter-hemispheric fissure. The disconnection can now be carried out completely into the splenium—the severing of the splenium was performed subpially to protect all the cisternal vessels. Extensive hemostasis is not usually necessary, and thus a ventricular drainage is not used. To conclude, the dura mater was closed and the bone flap reinserted and fixed with titanium plates and screws. Finally, a subgaleal drainage was inserted and the skin closed in two layers.

Healthy adult controls

We used open-access data from the Human Connectome Project 100 Unrelated Subjects (HCP100)

cohort (46 men, 54 women; M_{Age} = 29.11, SD = 3.68) as a robust benchmark for various network

features estimated in callosotomy patients. Participant recruitment and imaging procedures have

previously been detailed elsewhere[60,61]. In brief, data were collected over two sessions on a custom

3T Siemens Skyra with a 32-channel head coil. T_1-weighted structural scans were acquired using

a 3D magnetization prepared rapid gradient echo (MPRAGE) sequence (TR = 2400 ms, TE = 2.14

ms, TI = 1000 ms, flip angle = 8º, voxel size = 0.7 mm^3). Resting-state fMRI was collected with

eyes open over four ~14-minute scans, using a multiband echo-planar imaging (EPI) sequence

sensitive to the blood-oxygen-level-dependent (BOLD) contrast (TR = 720 ms, TE = 33.1 ms, flip

angle = 52º, voxel size = 2 mm^3, multiband factor = 8). All participants gave written informed

consent for a protocol approved by the Washington University Institutional Review Board. All

imaging data were downloaded in their raw (unprocessed) state.

Patient neuroimaging procedures

Patient data were collected on a 3T Siemens Vida at the Bethel Epilepsy Center using a 32-channel

head coil. T_1-weighted structural scans were acquired using a 3D MPRAGE sequence (TR = 1900

ms, TE = 2.47 ms, TI = 900 ms, flip angle = 8º, voxel size = 0.8 mm^3). Resting-state fMRI was

collected with eyes open using a single-shot EPI sequence sensitive to the BOLD contrast (TR =

2000 ms, TE = 30 ms, flip angle = 75º, voxel size = 3 x 3 x 3.30 mm, iPAT/GRAPPA factor = 2).

We note that each patient varied somewhat in the amount of resting-state data obtained (from ~8-

20 minutes total) due to differences in clinical scheduling and other imaging protocols not relevant

to the current analysis. This information is summarized in **Extended Data Fig. 1**.

fMRI preprocessing

All data (from both patients and healthy controls) were run through a unified preprocessing

pipeline. Initial preprocessing relied on functions from the Statistical Parametric Mapping 12

software (SPM12, Wellcome Trust Centre for Neuroimaging, London) in Matlab and Advanced

Normalization Tools (ANTs)[62]. Resting-state fMRI data were realigned and unwarped to correct

for head motion and motion x susceptibility artifact as well as slice-time corrected. Anatomical

images were denoised (ANTs' `DenoiseImage` with a spatially-adaptive Gaussian filter) and

submitted to ANTs' cortical thickness pipeline, which includes N4 bias field correction and high-quality image segmentation and brain extraction[63]. For some patients, the brain extraction mask included extracerebral components: this was corrected by manually eroding the extraction mask and re-applying it to the anatomical scan when necessary. Image coregistration parameters were subsequently estimated between an N3 bias-corrected mean BOLD reference and the skull-stripped T_1 image via `antsRegistration` (including both linear and nonlinear transformations to minimize residual signal deformations, cf. 'SyN distortion correction' as implemented by the widely-used fMRIPrep and QSIPrep pipelines).

Subsequent spatial normalization to a standard stereotaxic space presented a nontrivial problem for patient data—in the absence of a corpus callosum, generic registration routines (with both linear and nonlinear stages) will frequently pull structures such as the cingulate gyrus into the space occupied by the CC in the MNI template image, which misaligns brain regions relative to their location in our functional parcellation. We mitigated this problem in two ways. First, we implemented cost-function masking during the nonlinear step to minimize the influence of callosal regions that were transected by the surgery: these masks were specified in the fixed image space (i.e., the MNI152NLin6Asym template, as provided by the FMRIB Software Library, FSL[64,65]) because we lacked pre-operative data for each patient, and CC segments were defined by the JHU-DTI white matter atlas[66,67] (also provided in FSL). For patient EM, the genu was masked out; for patient BT, all but the splenium was masked out; and for all full splits, the entire cross-section of the callosum was masked out. Second, in addition to these masks, we specified a series of registration metric objectives at the nonlinear step. As is typical for `antsRegistration`, we estimated similarity between the MNI template and T_1 images using both ANTs' neighborhood cross-correlation and Mattes mutual information similarity metrics. We also provided smoothed segmentation posteriors derived from the ANTs cortical thickness pipeline (six in total: white matter, cortical gray matter, CSF, deep/subcortical gray matter, brain stem, and cerebellum) and estimated similarity between the MNI template segments and patient segments using a mean-squares objective, with cortical white matter and CSF segments downweighted in importance relative to the other four segments (by a factor of one third). Thus, the nonlinear step was comprised of eight similarity metrics. The transformation was estimated using BSplineSyN to allow for smoother deformations without sacrificing accuracy.

Ultimately, we concatenated and applied the estimated coregistration and spatial normalization

parameters to all fMRI data in a single interpolation via `antsApplyTransforms`, resampled into

2 mm^3 isotropic voxels. Registration quality was visually assessed to ensure appropriate mapping

to MNI space. Functional images were then smoothed using a 3D Gaussian kernel with FWHM =

5 mm^3. Prior to functional connectivity estimation, several additional preprocessing steps were

performed using in-house Matlab code: this included global signal scaling (median = 1000), linear

detrending, and nuisance regression. The nuisance regression model followed a '36-parameter'

(36P) approach including the six motion regressors from realignment, the first eigenvariate of

white matter signal, the first eigenvariate of CSF signal, and mean global signal—along with their

squares, their first derivatives (via backwards-differentiation), and the squares of the derivatives.

All nuisance regressors were detrended to match the BOLD data. We further augmented the design

matrix of the nuisance model with a series of binary spike regressors: one for each image volume

where the relative framewise displacement in head motion (accounting for both translation and

rotation) was ≥ 0.50 mm. This '36P + spike regression' model has importantly been shown to

outperform alternative confound regression strategies (e.g., motion scrubbing and ICA-based

methods) and provide the best control for motion artifact in functional connectivity analyses[20,21].

Patients generally showed no more head motion than HCP subjects; for reference, we provide a

summary of head motion parameters and the frequency of motion spikes for each patient in

Extended Data Fig. 1.

Functional connectivity estimation and network laterality

Functional network nodes were defined by parcellating the brain into 400 cortical regions and 32

subcortical regions, according to the Schaefer[17] and Tian[19] atlases, respectively. We extracted a

summary time course from each parcel by taking the first eigenvariate over time[68]. These regional

time series were bandpass-filtered to the range of 0.01-0.08 Hz. We then constructed 432 x 432

FC matrices by estimating the zero-order Pearson correlation between all pairs of regional time

series. For both HCP100 subjects and callosotomy patients with more than one resting-state scan,

runs were concatenated prior to FC estimation.

We sought to quantify differences in inter-hemispheric functional connectivity (IFC) between

callosotomy patients and healthy controls from a number of network neuroscience perspectives.

Pursuant to this, we began with an initial, breadth-first assessment of IFC within each of the seven

canonical resting-state networks defined by our cortical parcellation[17,18], as well as regions

spanning the subcortex[19]. Similarly to studies of functional lateralization in task-based fMRI, we

derived a *network laterality index* as a simple means of summarizing and quantifying the *relative*

magnitude of IFC as compared to the magnitude of intra-hemispheric FC. Each FC matrix was

first thresholded at an uncorrected level of $p < .001$ to avoid overly-conservative and highly-

nonlinear effects of multiple comparisons correction on the distribution of edge weights, while

simultaneously minimizing the influence of weak (potentially spurious) connections. We then

Fisher-*Z* transformed the thresholded FC matrices and, for each individual and for each network,

estimated a laterality index, *LI*, as:

588
$$LI = \frac{(\sum FC_{Intra-LH} + \sum FC_{Intra-RH}) - \sum FC_{Inter}}{\sum FC_{Intra-LH} + \sum FC_{Intra-RH} + \sum FC_{Inter}}$$

Intra-hemispheric connections within the left and right hemispheres are necessarily summed in the

numerator given that there are, by definition, twice as many inter-hemispheric connections as each

hemisphere alone. The result is a ratio on the interval of [-1, 1] that captures the extent to which

each canonical resting-state network is predominantly connected intra- or inter-hemispherically:

in the limit, a value of -1 would suggest that there is *no* intra-hemispheric FC and the network is

solely comprised of inter-hemispheric connections, while a value of 1 would indicate the reverse.

A value of 0 therefore represents a network that is perfectly balanced in the magnitude of intra- vs.

inter-hemispheric connections. Note that the results presented in **Fig. 3** are with respect to

positively-correlated edges only; we provide a comparison of anti-correlated edge lateralization in

Extended Data Fig. 3.

Community detection and modularity

While the above approach is useful for considering differences among canonical functional

networks, these systems were defined in healthy adult brains—it is therefore perhaps dubious to

assume that these networks follow a similar spatial organization in callosotomy patients. Data-

driven community detection techniques afford assessments of functional network topology
unbeholden to prior assumptions about specific brain regions and their canonical network
memberships[26,69]. This is achieved via decomposition of FC matrices using *modularity*
maximization algorithms, which group regions into *functional modules*: these represent regions
that are generally more functionally-coupled to each other than they are connected to other regions.
The optimization objective is subject to a *quality* metric, Q—unlike our previous laterality analysis
(which required separating out positive and negative edges), this can be estimated over FC matrices
comprised of both positive and negative correlations[70]. We performed this decomposition using
the `community_louvain` function from the Brain Connectivity Toolbox
(https://sites.google.com/site/bctnet/) with asymmetric treatment of negative edge weights[70,71].
Importantly, modularity maximization is sensitive to spatial scale, subject to a *resolution*
parameter, γ[28]. Small values of γ typically yield coarse partitions with a small number of large
communities; larger values of γ, by contrast, afford more fine-grained partitions. Because the
solution of this decomposition is dependent upon one's choice for the resolution parameter, we
tested a range of values from $\gamma = [0.10, 2]$ (in steps of 0.10). To further account for variability in
community assignments (e.g., due to algorithmic stochasticity arising from random initializations),
we estimated modularity 1000 times for each value of γ—optimal consensus partitions were
obtained using the `consensus_similarity` function from the Network Community Toolbox
(https://commdetect.weebly.com/).
624
In similar fashion to the network laterality indices described above, we derived a *module laterality*
index for the resulting partitions according to prior work[27]. This metric is limited to the [0, 1]
interval, where a value of 0, for example, indicates that all functional communities are perfectly
bilateral (i.e., all of the constituent brain regions belonging to each community are equally-
represented between the left and right hemispheres), and values closer to 1 indicate progressively
more lateralized topologies. We note that in this case, more fine-grained partitions may be
beneficial for assessing modular laterality—nevertheless, the results shown in **Figs. 4-5** reflect a
'default' $\gamma = 1$ and we further illustrate in **Extended Data Fig. 4** that full splits rapidly diverge
from controls even at smaller values of γ (while partial splits and healthy controls remain relatively
consistent, regardless of γ).
635

Inter-hemispheric network dynamics

Finally, we used edge time series (ETS)[30] to assess the extent to which *many* brain regions in the left and right hemispheres dynamically co-fluctuate in synchrony. The ETS approach has been used to obtain moment-to-moment, time-resolved estimates of FC and characterize the nature of high-amplitude events[31] that occur when multivariate BOLD signals simultaneously fluctuate together over brief intervals of time—these bursts of synchronized network dynamics are sufficient to recapitulate 'static' FC estimates over the course of a scan[31], and importantly, they are independent of head motion[30]. Current evidence instead suggests these high-amplitude co-fluctuations are governed, in part, by the modular organization of the brain and its underlying structural architecture[29]. An ETS can be computed by taking a pair of BOLD time series from two regions, *Z*-scoring them, and taking the elementwise product of the signals at each timepoint—the mean of an ETS is equivalent to the zero-order Pearson correlation between the two time series, and thus ETS constitute a temporal-unwrapping of FC over the scan[30].

Here, we first separated out all regions in the left hemisphere and all regions in the right hemisphere. We constructed a *T* x 23220 ETS matrix for each hemisphere, where *T* gives the number of timepoints and 23220 gives the total number of edges within each hemisphere. A summary time course quantifying the total amplitude of intra-hemispheric co-fluctuations at any given moment (i.e., across *all* edges) was obtained by taking the root-sum-of-squares (RSS) at each timepoint[31]. We then computed the Spearman correlation between the left hemisphere RSS and right hemisphere RSS time series to capture the extent to which network dynamics in each hemisphere were synchronized.

Data and code availability

Patient data will be made available upon reasonable request to the corresponding authors; all relevant processing and analysis code will be uploaded to a Github repository.

References

1.   Bassett DS, Sporns O. Network neuroscience. *Nat Neurosci* **20**, 353-364 (2017).

667 2.   Honey CJ, *et al.* Predicting human resting-state functional connectivity from structural
668   connectivity. *Proc Natl Acad Sci U S A* **106**, 2035-2040 (2009).
669

670 3.   van den Heuvel MP, Mandl RC, Kahn RS, Hulshoff Pol HE. Functionally linked resting-
671   state networks reflect the underlying structural connectivity architecture of the human
672   brain. *Hum Brain Mapp* **30**, 3127-3141 (2009).
673

674 4.   Fukushima M, Betzel RF, He Y, van den Heuvel MP, Zuo XN, Sporns O. Structure-
675   function relationships during segregated and integrated network states of human brain
676   functional connectivity. *Brain Struct Funct* **223**, 1091-1106 (2018).
677

678 5.   Hermundstad AM, *et al.* Structurally-constrained relationships between cognitive states
679   in the human brain. *PLoS Comput Biol* **10**, e1003591 (2014).
680

681 6.   Hermundstad AM, *et al.* Structural foundations of resting-state and task-based functional
682   connectivity in the human brain. *Proc Natl Acad Sci U S A* **110**, 6169-6174 (2013).
683

684 7.   Hofer S, Frahm J. Topography of the human corpus callosum revisited--comprehensive
685   fiber tractography using diffusion tensor magnetic resonance imaging. *Neuroimage* **32**,
686   989-994 (2006).
687

688 8.   Liu Y, *et al.* Connectivity-Based Topographical Changes of the Corpus Callosum During
689   Aging. *Front Aging Neurosci* **13**, 753236 (2021).
690

691 9.   Hung SC, *et al.* Early recovery of interhemispheric functional connectivity after corpus
692   callosotomy. *Epilepsia* **60**, 1126-1136 (2019).
693

694 10.   Johnston JM, *et al.* Loss of resting interhemispheric functional connectivity after
695   complete section of the corpus callosum. *J Neurosci* **28**, 6453-6458 (2008).
696

697 11.   Roland JL, *et al.* On the role of the corpus callosum in interhemispheric functional
698   connectivity in humans. *Proc Natl Acad Sci U S A* **114**, 13278-13283 (2017).
699

700 12.   Pizoli CE, *et al.* Resting-state activity in development and maintenance of normal brain
701   function. *Proc Natl Acad Sci U S A* **108**, 11638-11643 (2011).
702

703 13.   Gazzaniga MS. Forty-five years of split-brain research and still going strong. *Nat Rev
704   Neurosci* **6**, 653-659 (2005).
705

706 14.   Volz LJ, Gazzaniga MS. Interaction in isolation: 50 years of insights from split-brain
707   research. *Brain* **140**, 2051-2060 (2017).
708

709 15.   Miller MB, Volz LJ, Simonson JM, Gazzaniga MS. Split-brain patients: A clinical versus
710   experimental perspective. In: *Cerebral Asymmetries* (eds Papagno C, Corballis P).
711   Elsevier BV (in press).
712

16. Bekir S*, et al.* No disconnection syndrome after near-complete callosotomy. Preprint at https://www.biorxiv.org/content/10.1101/2025.02.16.638524v1 (2025).

17. Schaefer A*, et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114 (2018).

18. Yeo BT*, et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).

19. Tian Y, Margulies DS, Breakspear M, Zalesky A. Topographic organization of the human subcortex unveiled with functional connectivity gradients. *Nat Neurosci* **23**, 1421-1432 (2020).

20. Ciric R*, et al.* Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *Neuroimage* **154**, 174-187 (2017).

21. Parkes L, Fulcher B, Yucel M, Fornito A. An evaluation of the efficacy, reliability, and sensitivity of motion correction strategies for resting-state functional MRI. *Neuroimage* **171**, 415-436 (2018).

22. Catani M, Schotten M. *Atlas of Human Brain Connections*. Oxford University Press (2012).

23. Patel MD, Toussaint N, Charles-Edwards GD, Lin JP, Batchelor PG. Distribution and fibre field similarity mapping of the human anterior commissure fibres by diffusion tensor imaging. *MAGMA* **23**, 399-408 (2010).

24. Di Virgilio G, Clarke S, Pizzolato G, Schaffner T. Cortical regions contributing to the anterior commissure in man. *Exp Brain Res* **124**, 1-7 (1999).

25. Bassett DS, Gazzaniga MS. Understanding complexity in the human brain. *Trends Cogn Sci* **15**, 200-209 (2011).

26. Sporns O, Betzel RF. Modular Brain Networks. *Annu Rev Psychol* **67**, 613-640 (2016).

27. Doron KW, Bassett DS, Gazzaniga MS. Dynamic network structure of interhemispheric coordination. *Proc Natl Acad Sci U S A* **109**, 18661-18668 (2012).

28. Fortunato S, Barthelemy M. Resolution limit in community detection. *Proc Natl Acad Sci U S A* **104**, 36-41 (2007).

29. Pope M, Fukushima M, Betzel RF, Sporns O. Modular origins of high-amplitude cofluctuations in fine-scale functional connectivity dynamics. *Proc Natl Acad Sci U S A* **118**, (2021).

30. Faskowitz J, Esfahlani FZ, Jo Y, Sporns O, Betzel RF. Edge-centric functional network representations of human cerebral cortex reveal overlapping system-level architecture. *Nat Neurosci* **23**, 1644-1654 (2020).

31. Zamani Esfahlani F, *et al.* High-amplitude cofluctuations in cortical activity drive functional connectivity. *Proc Natl Acad Sci U S A* **117**, 28393-28401 (2020).

32. Barttfeld P, Uhrig L, Sitt JD, Sigman M, Jarraya B, Dehaene S. Signature of consciousness in the dynamics of resting-state brain activity. *Proc Natl Acad Sci U S A* **112**, 887-892 (2015).

33. Tooley UA, *et al.* The Age of Reason: Functional Brain Network Development during Childhood. *J Neurosci* **42**, 8237-8251 (2022).

34. Sydnor VJ, *et al.* Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. *Neuron* **109**, 2820-2846 (2021).

35. Reynolds JE, Grohs MN, Dewey D, Lebel C. Global and regional white matter development in early childhood. *Neuroimage* **196**, 49-58 (2019).

36. Whitaker KJ, *et al.* Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. *Proc Natl Acad Sci U S A* **113**, 9105-9110 (2016).

37. Tyszka JM, Kennedy DP, Adolphs R, Paul LK. Intact bilateral resting-state networks in the absence of the corpus callosum. *J Neurosci* **31**, 15154-15162 (2011).

38. Lassonde M, Sauerwein H, Chicoine AJ, Geoffroy G. Absence of disconnexion syndrome in callosal agenesis and early callosotomy: brain reorganization or lack of structural specificity during ontogeny? *Neuropsychologia* **29**, 481-495 (1991).

39. Graham D, Tisdall MM, Gill D. Corpus callosotomy outcomes in pediatric patients: A systematic review. *Epilepsia* **57**, 1053-1068 (2016).

40. Fuchs E, Flugge G. Adult neuroplasticity: more than 40 years of research. *Neural Plast* **2014**, 541870 (2014).

41. Nomi JS, *et al.* Diffusion weighted imaging evidence of extra-callosal pathways for interhemispheric communication after complete commissurotomy. *Brain Struct Funct* **224**, 1897-1909 (2019).

42. Uddin LQ, *et al.* Residual functional connectivity in the split-brain revealed with resting-state functional MRI. *Neuroreport* **19**, 703-709 (2008).

43. Marcantoni I, *et al.* Interhemispheric functional connectivity: an fMRI study in callosotomized patients. *Front Hum Neurosci* **18**, 1363098 (2024).

44. O'Reilly JX, *et al.* Causal effect of disconnection lesions on interhemispheric functional connectivity in rhesus monkeys. *Proc Natl Acad Sci U S A* **110**, 13982-13987 (2013).

45. Funnell MG, Corballis PM, Gazzaniga MS. Cortical and subcortical interhemispheric interactions following partial and complete callosotomy. *Arch Neurol* **57**, 185-189 (2000).

46. Gazzaniga MS, Freedman H. Observations on visual processes after posterior callosal section. *Neurology* **23**, 1126-1130 (1973).

47. Risse GL, Gates J, Lund G, Maxwell R, Rubens A. Interhemispheric transfer in patients with incomplete section of the corpus callosum. Anatomic verification with magnetic resonance imaging. *Arch Neurol* **46**, 437-443 (1989).

48. Gazzaniga MS. Interhemispheric integration. In: *Neurobiology of Neocortex* (eds Rakic P, Singer W). John Wiley & Sons (1988).

49. Seguin C, Mansour LS, Sporns O, Zalesky A, Calamante F. Network communication models narrow the gap between the modular organization of structural and functional brain networks. *Neuroimage* **257**, 119323 (2022).

50. Suarez LE, Markello RD, Betzel RF, Misic B. Linking Structure and Function in Macroscale Brain Networks. *Trends Cogn Sci* **24**, 302-315 (2020).

51. Misic B, *et al.* Network-Level Structure-Function Relationships in Human Neocortex. *Cereb Cortex* **26**, 3285-3296 (2016).

52. Aboitiz F, Lopez J, Montiel J. Long distance communication in the human brain: timing constraints for inter-hemispheric synchrony and the origin of brain lateralization. *Biol Res* **36**, 89-99 (2003).

53. Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus callosum. *Brain Res* **598**, 143-153 (1992).

54. Horowitz A, Barazany D, Tavor I, Bernstein M, Yovel G, Assaf Y. In vivo correlation between axon diameter and conduction velocity in the human brain. *Brain Struct Funct* **220**, 1777-1788 (2015).

55. van den Heuvel MP, Sporns O. Rich-club organization of the human connectome. *J Neurosci* **31**, 15775-15786 (2011).

56. Hagmann P, *et al.* Mapping the structural core of human cerebral cortex. *PLoS Biol* **6**, e159 (2008).

57. Koch C, Massimini M, Boly M, Tononi G. Neural correlates of consciousness: progress and problems. *Nat Rev Neurosci* **17**, 307-321 (2016).

852   58.   Melloni L*, et al.* An adversarial collaboration protocol for testing contrasting predictions
853   of global neuronal workspace and integrated information theory. *PLoS One* **18**, e0268577
854   (2023).
855

856   59.   Koch C, Massimini M, Boly M, Tononi G. Posterior and anterior cortex - where is the
857   difference that makes the difference? *Nat Rev Neurosci* **17**, 666 (2016).
858

859   60.   Van Essen DC*, et al.* The WU-Minn Human Connectome Project: an overview.
860   *Neuroimage* **80**, 62-79 (2013).
861

862   61.   Van Essen DC*, et al.* The Human Connectome Project: a data acquisition perspective.
863   *Neuroimage* **62**, 2222-2231 (2012).
864

865   62.   Tustison NJ*, et al.* The ANTsX ecosystem for quantitative biological and medical
866   imaging. *Sci Rep* **11**, 9068 (2021).
867

868   63.   Tustison NJ*, et al.* Large-scale evaluation of ANTs and FreeSurfer cortical thickness
869   measurements. *Neuroimage* **99**, 166-179 (2014).
870

871   64.   Evans AC, Janke AL, Collins DL, Baillet S. Brain templates and atlases. *Neuroimage* **62**,
872   911-922 (2012).
873

874   65.   Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. Fsl. *Neuroimage* **62**,
875   782-790 (2012).
876

877   66.   Hua K*, et al.* Tract probability maps in stereotaxic spaces: analyses of white matter
878   anatomy and tract-specific quantification. *Neuroimage* **39**, 336-347 (2008).
879

880   67.   Wakana S*, et al.* Reproducibility of quantitative tractography methods applied to cerebral
881   white matter. *Neuroimage* **36**, 630-644 (2007).
882

883   68.   Friston KJ, Rotshtein P, Geng JJ, Sterzer P, Henson RN. A critique of functional
884   localisers. *Neuroimage* **30**, 1077-1087 (2006).
885

886   69.   Zamani Esfahlani F*, et al.* Modularity maximization as a flexible and generic framework
887   for brain network exploratory analysis. *Neuroimage* **244**, 118607 (2021).
888

889   70.   Rubinov M, Sporns O. Weight-conserving characterization of complex functional brain
890   networks. *Neuroimage* **56**, 2068-2079 (2011).
891

892   71.   Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and
893   interpretations. *Neuroimage* **52**, 1059-1069 (2010).
894

Extended Data

Patient	Scan num.	Scan duration		Head motion estimates					
		Minutes	Frames	Min FWD	Max FWD	Med FWD	MAD FWD	Num. spikes	Spike %
EM	1	8	240	0.03	2.44	0.13	0.05	9	3.75
BT	1	7	210	0.04	1.09	0.19	0.07	11	5.24
BT	2	7	210	0.05	1.34	0.25*	0.08	9	4.29
HY	1	12	360	0.02	0.44	0.14	0.05	0	0.00
DT	1	12	360	0.02	10.74*	0.15	0.05	31	8.61*
TJ	1	7	210	0.02	0.59	0.13	0.04	2	0.95
TJ	2	7	210	0.03	0.38	0.11	0.04	0	0.00
LJ	1	7	210	0.06	0.55	0.20	0.04	2	0.95
LJ	2	7	210	0.04	0.38	0.18	0.04	0	0.00
LJ	3	7	210	0.06	0.56	0.20	0.05	3	1.43

*Denotes head motion estimate exceeding normal range, see corresponding figure(s) below.

Extended Data Fig. 1. Scan times and head motion estimates across callosotomy patients. Due to differences in clinical scheduling and other imaging protocols not relevant to the current analysis, patients varied in resting-state fMRI scan time. Here we summarize these differences and provide various estimates of head motion for each scan. Motion parameters are given with respect to *framewise displacement* (FWD)—the amount of total head motion (in millimeters, considering both translational displacements and rotations) observed from one image volume to the next. Because FWD is a time series, we provide the minimum, maximum, and median FWD over each scan (with the median absolute deviation, *MAD*, over time). We also denote the number of motion spikes, defined as any frame with FWD ≥ 0.50 mm, and the percentage of frames contaminated with motion spikes—these were used to define individual spike regressors for nuisance signal regression[20,21]. Patients generally did not show any more head motion than healthy adult controls (see visualized distributions). Note that in rare cases where patients exceeded the range of controls in one metric, they were typically within the standard range of other metrics. Point intervals under

the control distributions give the median (black dot) and the 80-95% quantiles around the median

(thick black lines and thin black lines, respectively).

Extended Data Fig. 2. Multivariate similarity in spatial patterns of functional connectivity.

As an initial estimate of the extent to which full vs. partial callosotomy disrupted intra- vs. inter-

hemispheric functional connectivity (FC), we binarized FC matrices at a threshold of $z(r) > 0$

(where $z(r)$ refers to the Fisher-Z transformed correlation coefficients for each network edge) and

computed the Dice coefficient between each patient and each individual healthy control—this

statistic ranges from [0,1] and quantifies the degree of similarity in two patterns of FC (liberally,

whether the same edges show *any* FC > 0, regardless of the exact magnitude of those connections).

The distributions for each patient thus show the range of similarity between a given patient and

each control. Patient EM (with the most corpus callosum intact) is generally most similar to

controls; full splits, on the other hand, show markedly dissimilar patterns of inter-hemispheric FC

relative to intra-hemispheric FC. Point intervals under each distribution give the median Dice

similarity (black dot) and the 80-95% quantiles around the median (thick black lines and thin black

lines, respectively). Abbreviations: LH, Left Hemisphere; RH, Right Hemisphere.

Extended Data Fig. 3. Anti-correlated functional connectivity is generally not more lateralized in callosotomy patients. Network laterality indices across seven canonical resting-state systems[17,18] and subcortical regions[19]. To contrast against **Fig. 3**, here we estimate lateralization of negative (anti-correlated) connections. Values closer to 0 indicate an equal balance between intra- and inter-hemispheric (negative) FC, while values closer to -1 or 1 indicate more inter-hemispheric FC intra-hemispheric (negative) FC, respectively. Note that, unlike in **Fig. 3**, most distributions skew to the left (indicating that negative FC is predominantly inter-hemispheric, i.e., not lateralized), and both full and partial splits frequently fall within the range of HCP controls. Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: FC; Functional Connectivity; SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Extended Data Fig. 4. Lateralization of functional modules is readily-apparent in full splits regardless of resolution. Modularity maximization algorithms are sensitive to a spatial resolution parameter, γ[28]. A value of $\gamma = 1$ is often treated as a 'default' level of resolution, whereas values of $\gamma < 1$ or $\gamma > 1$ produce progressively more coarse or fine-grained solutions, respectively. Here we show that differences in modular laterality between patients and healthy adult controls are not strictly-dependent upon the chosen value of γ. We performed 1000 iterations of modularity maximization at each of 20 levels of γ (ranging from [0.10, 2] in steps of 0.10), derived an optimal consensus partition at each level (see **Methods**), and estimated a laterality index for the resulting partitions[27] (such that values closer to 1 indicate more lateralized network topologies). Full splits (solid-colored lines) rapidly diverged from the distribution of healthy controls (quantile intervals shown in gradations of purple, with the solid black line representing the median) demonstrating more lateralized organization of functional modules, even at coarse levels of $\gamma < 1$. Partial splits (dotted colored lines), by contrast, consistently remained within the control distribution.

Extended Data Fig. 5. Estimates of large-scale dynamic functional synchrony by network.
Decomposing whole-brain comparisons of RSS synchrony in **Fig. 6b** to individual networks
reveals nuanced differences between patients and neurotypical adults (colored distributions). As
before, partial splits consistently fall within the range of HCP controls. Full splits generally show
more desynchronized network dynamics in higher-order cognitive systems, but comparatively-
neurotypical levels of inter-hemispheric synchrony in limbic and subcortical networks (which are
not strictly dependent upon structural connections through the corpus callosum, also reflecting **Fig.
3**). Point intervals under the control distributions give the HCP100 median (black dot) and 80-95%
quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations:
RSS, Root-Sum-Of-Squares (see **Methods**); SomMot, SomatoMotor Network; DorsAttn, Dorsal
Attention Network; SalVentAttn, Salience/Ventral Attention Network.