Full inter-hemispheric integration sustained by a fraction of posterior callosal fibers

Tyler Santander[1,2,*], Selin Bekir[2], Theresa Paul[3], Jessica M. Simonson[1,2], Valerie M. Wiemer[3], Henri Etel Skinner[2], Johanna L. Hopf[4], Anna Rada[4], Friedrich G. Woermann[4,5], Thilo Kalbhenn[4,6], Barry Giesbrecht[1,2], Christian G. Bien[4], Olaf Sporns[7], Michael S. Gazzaniga[2], Lukas J. Volz[3,†], & Michael B. Miller[1,2,†,*]

[1]Institute for Collaborative Biotechnologies, University of California, Santa Barbara, Santa Barbara, CA, USA

[2]Department of Psychological & Brain Sciences, University of California, Santa Barbara, Santa Barbara, CA, USA

[3]Medical Faculty, University of Cologne, and Department of Neurology, University Hospital Cologne, Cologne, Germany

[4]Department of Epileptology, Krankenhaus Mara, Bethel Epilepsy Center, Medical School OWL, Bielefeld University, Bielefeld, Germany

[5]MRI Department, Society for Epilepsy Research, Krankenhaus Mara, Bethel Epilepsy Center, Bielefeld, Germany

[6]Department of Neurosurgery, Evangelisches Klinikum Bethel, Medical School OWL, Bielefeld University, Bielefeld, Germany

[7]Department of Psychological & Brain Sciences, Indiana University, Bloomington, IN, USA

[†]Authors contributed equally

*To whom correspondence should be addressed:

Tyler Santander (tsantander@ucsb.edu)

Michael B. Miller (mbmiller@ucsb.edu)

Acknowledgements

T.S., S.B., J.M.S., H.E.S., B.G., and M.B.M. were sponsored by the Army Research Office under contract W911NF-19-D-0001 for the Institute for Collaborative Biotechnologies. T.P. and L.J.V were funded by the German Research Foundation (CRC-1451, Project 431549029).

Author contributions

T.S., S.B., T.P., H.E.S., J.L.H., A.R., B.G., C.G.B., O.S., M.S.G., L.J.V., and M.B.M. conceived and planned the experiments. T.P., V.M.W., J.L.H., A.R., F.G.W., C.G.B., and L.J.V. assisted with patient recruitment and were responsible for data collection. T.K. performed the callosotomy surgeries. T.S., J.M.S., O.S., and L.J.V. conceived the data analysis strategy; T.S., J.M.S., and O.S. contributed to analysis code and executed the data processing/analysis pipeline. T.S. wrote the manuscript with contributions from T.K., O.S., M.S.G., L.J.V., and M.B.M. The project was supervised by A.R., C.G.B., L.J.V., and M.B.M. Funding was secured by B.G., L.J.V., and M.B.M. All authors assisted with manuscript revision and approved the final submitted version. L.J.V. and M.B.M. contributed equally.

Conflicts of Interest

The authors declare no competing interests.

Abstract

The dynamic integration of the lateralized and specialized capacities of the two cerebral hemispheres constitutes a hallmark feature of human brain function. This inter-hemispheric exchange of information is thought to critically depend upon the corpus callosum. Classical anatomical descriptions of callosal organization outline a topographic gradient from front to back, such that specific transcallosal fibers support distinct aspects of integrated brain function. Here we present a challenge to this conventional model. Using neuroimaging data obtained from a new cohort of adult callosotomy patients, we leverage modern network neuroscience techniques to show—for the first time—that full inter-hemispheric integration can be achieved via a small proportion of posterior callosal fibers. Partial callosotomy patients with spared callosal fibers retained widespread patterns of inter-hemispheric functional connectivity and showed no signs of behavioral disconnection syndromes, even with only 1 cm of the splenium intact. Conversely, only complete corpus callosotomy patients demonstrated sweeping disruptions of inter-hemispheric network architectures, aligning with disconnection syndromes long-thought to reflect diminished information propagation and communication across the brain. These findings motivate a novel mechanistic understanding of synchronized inter-hemispheric neural activity for large-scale human brain function and behavior.

Main

In recent decades, network neuroscience has emerged as a powerful analysis framework for modeling functional interactions between remote neural systems. A foundational assumption of this approach is that anatomical links between brain regions constrain neuronal dynamics and support the emergence of large-scale functional networks, which in turn enable the efficient integration of perceptual, motor, and higher-order cognitive information necessary for human behavior[1,2,3,4,5,6]. Yet, across the whole-brain, structural connectivity (i.e., the direct white matter pathways tethering brain regions together) often only accounts for a fraction of the variance in functional connectivity (i.e., statistical dependencies in regional brain activity over time)[2,7,8]. For some interregional connections, however, structure seems to predetermine function more closely: when considering pairs of homotopic brain regions—mirrored areas of the cortex in the left and right hemispheres—individual differences in the microstructure of the corpus callosum (CC) predict the strength of functional coupling between such regions[9]. On a broader level, the topographic organization of CC fibers has been characterized along a well-defined anterior-to-posterior anatomical gradient, corresponding to the spatial organization of canonical brain networks servicing various sensorimotor and cognitive functions[10,11]. Differential patterns of myelination along callosal fibers (among other cytological properties) have further been attributed to differences in the 'fast' vs. 'slow' action of primary sensory vs. association cortices, respectively, and distinguish between homotopic and heterotopic connections between regions[12,13,14,15]. A confluence of evidence has thus suggested that CC connectivity is the central conduit through which inter-hemispheric brain networks coordinate and synchronize function. One consequence of this view is the prediction that damage to the CC should result in drastic functional disconnection between hemispheres—with spatially-specific disruptions in accordance with the localization of lesioned fibers along the aforementioned anatomical gradient. A corollary to this prediction is that disruptions may resolve in the long-term, such that neuroplastic and reorganizational processes enable recovery of function given a sufficient length of time.

Split-brain patients offer a powerful means of testing this core conceit of structure-function correspondence. These are individuals in whom the CC has been severed in an effort to treat medically-intractable epilepsy. However, neural data on these patients remains scarce—

callosotomy was largely discontinued around the turn of the 21st century, coinciding with the rise

of network neuroscience. Extant data largely stem from anesthetized pediatric patients[16,17,18,19].

These studies have lent support to the hypothesis that full versus partial callosotomy confer

differential effects with respect to the location and extent of functional disconnections; at the same

time, however, they have also provided contradictory interpretations regarding recovery of

function post-operatively[16,18,19]. While the exact mechanisms affording preserved inter-

hemispheric synchrony in split-brain patients are yet to be elucidated, these outcomes have

previously been proposed to arise from functional reorganization along alternative inter-

hemispheric pathways, including the anterior commissure and/or other subcortical

structures[16,20,21,22]. Thus, to date, it remains unknown how severing the largest fiber bundle of the

human brain affects the topological organization and intrinsic dynamics of large-scale functional

networks.

To address this fundamental question, we present the first investigation using multiple network

neuroscience methods to characterize the impacts of full and partial callosotomy on the adult

human brain. We hypothesized that full callosotomy would wholly disrupt functional synchrony

across the cerebral hemispheres, while the spatial distribution of functional disconnections

following partial callosotomy would unveil the anatomical gradient of the CC. As predicted, we

found that complete sectioning of the CC broadly obliterates both typical inter-hemispheric

network topologies and dynamic patterns of inter-hemispheric temporal synchrony. More

strikingly, however, we also found that a small proportion of intact posterior CC fibers (a case of

partial callosotomy) was sufficient to support widespread inter-hemispheric coupling. This finding

challenges our conventional understanding of the CC's functional topography and sheds new light

on the critical mechanisms for inter-hemispheric information integration.

Patient descriptions

Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were tested post-

operatively at the Bethel Epilepsy Center (Bielefeld, Germany). The time between surgery and

testing varied between one and six years (**Table 1**). In two patients (EM and BT), surgery did not

result in a complete callosal transection; they are hereafter referred to as *partial splits*. In patient

EM, the surgery was stopped after sectioning the anterior third of the CC; patient BT was spared

approximately 1 cm of splenium. The remaining four patients (HY, DT, TJ, and LJ) were completely sectioned and are hereafter referred to as *full splits*. All patients have intact anterior and posterior commissures regardless of the extent of the callosal section.

Figure 1. Patient anatomical scans. Post-operative T_1 and diffusion MRI scans collected for all six patients at time of testing. White arrows/ellipses for patients EM and BT highlight intact CC (i.e., partial splits). The remaining four patients were fully-sectioned. Note that a series of clinical observations suggested behavioral disconnection syndromes in all full splits, but not in partial splits.

Table 1. General overview of callosotomy patients.

Patient	Age at surgery	Age at testing	Sex	Handedness	Extent of callosotomy	Disconnection syndrome
EM	17	19	M	Right	Partial	Absent
BT	22	28	M	Right	Partial	Absent
HY	21	22	M	Right	Complete	Present
DT	15	20	M	Right	Complete	Present
TJ	49	52	F	Right	Complete	Present
LJ	57	60	M	Right	Complete	Present

Callosotomy is not commonly associated with post-operative neuropsychiatric or neurological disorders—and indeed, all of these individuals previously tested normal on standard clinical neuropsychological and intelligence assessments. Rather, split-brain patients often demonstrate *disconnection syndromes*, such that they appear to have two hemispheres operating independently,

unable to communicate information from one half of the brain to the other[23,24,25]. In the current set
of patients, behavioral disconnection syndromes were clinically-observed using simple, lateralized
bedside testing protocols. This included visual tasks (finger perimetry), somesthetic tasks
(topognosis, stereognosis), and motor tasks (bimanual coordination). Comprehensive descriptions
of these tests are described elsewhere[26]. In brief, however: as predicted, all four complete
callosotomy patients clearly indicated behavioral disconnections. Partial splits, in contrast, did not
show *any* signs of disconnection syndromes—including, remarkably, patient BT (with a mere 1
cm of splenium). We confirmed this by testing BT more extensively, using tasks intended to cover
various cognitive domains and modalities along the anteroposterior axis of the CC[26]. Although
behavioral performance is not the focus of this report, these findings contradict our classical
understanding of callosal topography, which posits functionally-specific deficits depending on
which portions of the CC were severed. Consequently, a small proportion of posterior callosal
fibers may be sufficient to support information integration across large-scale inter-hemispheric
brain networks.

Inter-hemispheric functional connectivity after callosotomy
All patients underwent structural and functional magnetic resonance imaging at the time of their
visit. Here we describe results from whole-brain analyses of resting-state functional connectivity,
specifically with regard to inter-hemispheric functional connectivity (IFC). Resting-state scans
slightly differed in length between patients (see **Extended Data Fig. 1** for details); all scans were
completed without sedation and patients were asked to simply lie still with their eyes open.

To benchmark patient data against 'typical' patterns of IFC observed in healthy adults, we used
resting-state scans from the Human Connectome Project 100 Unrelated Subjects (HCP100) cohort.
All HCP100 subjects and callosotomy patients underwent an identical preprocessing pipeline,
including conservative corrections for mitigating head motion artifacts[27,28] (see **Methods** for
detailed processing procedures and **Extended Data Fig. 1** for a quantitative description of patient
head motion). Whole-brain resting-state FC in the range of 0.01-0.08 Hz was assessed by
parcellating the brain into 400 cortical regions and 32 subcortical regions, according to the
Schaefer[29] and Tian[30] atlases, respectively (**Fig. 2a**). We then correlated all pairs of regional time
series to construct a 432 x 432 FC matrix for each patient and healthy control (**Figs. 2b-c**).

Figure 2. Inter-hemispheric functional connectivity is broadly-diminished in full (but not partial) splits. (a) Whole-brain functional parcellation comprised of seven canonical resting-state networks[29,31] and subcortical nodes[30]. **(b)** A standard reference for neurotypical whole-brain functional connectivity (FC), given as the median over HCP100 subjects. FC was estimated by

correlating low-frequency brain activity (0.01-0.08 Hz) between all pairs of regions in **(a)**. The large boxes outlining the upper left and lower right quadrants denote *intra-hemispheric* FC; off-diagonal quadrants thus capture *inter-hemispheric* FC (IFC). **(c)** Whole-brain FC estimates for each callosotomy patient, thresholded at $p < .001$ (uncorrected for multiple comparisons). Typical patterns of IFC were largely abolished in full splits (HY, DT, TJ, LJ), while partial splits (EM, BT) were more similar to healthy controls. Abbreviations: SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network; LH, Left Hemisphere; RH, Right Hemisphere.

All full splits demonstrated marked reductions in IFC relative to both healthy adult controls and partial splits (**Figs. 2b-c**). To quantify this, we first compared multivariate spatial patterns of IFC between each patient and each healthy control: the FC matrices in **Fig. 2c** were binarized to distinguish between positive and negatively-correlated edges, and we estimated similarity in the spatial extent of IFC (relative to controls) using the Dice coefficient (*DSC*). As a point of reference, the *minimum* Dice similarity in IFC from one healthy control to another was ~0.46—all full splits were considerably less similar to controls, suggesting a sweeping disruption of IFC following complete callosotomy (**Extended Data Fig. 2**; HY: Median *DSC* = 0.28, 95% QI = [0.26, 0.30]; DT: 0.27 [0.25, 0.29]; TJ: 0.27 [0.25, 0.29]; LJ: 0.27 [0.26, 0.28]). Patient EM—a partial split with approximately two-thirds of their CC intact—showed a greater degree of similarity to controls, reflecting more typical, intact patterns of IFC (Median *DSC* = 0.46 [0.42, 0.49]). Finally, patient BT (with a small sparing of fibers through the splenium) demonstrated fragmented, albeit still widespread, patterns of IFC: thus, this individual was less similar to controls than EM but still greater than full splits (Median *DSC* = 0.36 [0.34, 0.38]). With regard to spatial patterns of intra-hemispheric FC, all patients were better-matched to controls (**Extended Data Fig. 2**), suggesting that callosotomy most impairs inter-hemispheric functional coupling with full splits suffering the greatest disruptions.

Curiously, while IFC was largely dissolved in full splits, inspection of individual FC matrices suggested that visual network connectivity might be preserved to some extent (**Fig. 2c**). This may be due to synchronized perceptual inputs (scans were completed with eyes open), driving *apparent* IFC despite the lack of transcallosal connectivity between visual regions and evidence for

behavioral disconnections. We therefore sought to assess differences in the magnitude of IFC between functional homologues to determine whether any preserved IFC was unique to sensory networks. Here we computed a series of *network laterality indices*—one for each network—where a value of zero indicates a perfect balance between inter- and intra-hemispheric FC, and values closer to -1 or 1 indicate relatively more inter- or intra-hemispheric FC, respectively (see **Methods** for additional details). As before, we focused on positively-coupled regions, but comparisons of anti-correlated edges within each network are shown in **Extended Data Fig. 3**. Consistent with the observed behavioral disconnection effects, all full splits demonstrated more lateralized patterns of functional connectivity relative to healthy controls and partial splits, who typically fell within the control distributions (**Fig. 3**). This was evident in both higher-order, transmodal systems (e.g., the default mode and dorsal attention networks) and in somatosensory networks (with the exception of patient TJ who fell within the upper tail of the control distribution in the somatomotor network). Patient EM was more lateralized across frontoparietal control nodes, which include a number of regions that are likely to be structurally disconnected given the extent of their callosotomy; however, BT generally showed similar degrees of network bilaterality as healthy controls, despite having a mere fraction of splenium intact. Finally, full splits also appeared to show more bilateral FC across limbic regions like the orbitofrontal cortex and temporal poles, which are known to be structurally coupled through the anterior commissure as well as the CC[32,33,34]. Taken together, these results highlight the dependence of inter-hemispheric networks on CC connectivity—yet, neurotypical levels of inter-hemispheric integration are possible even without a fully-intact CC.

Figure 3. Full callosotomy is associated with more lateralized functional connectivity. Network laterality indices across seven canonical resting-state systems[29,31] and subcortical regions[30]. Values closer to zero indicate an equal balance between intra- and inter-hemispheric FC, while values closer to one indicate more intra-hemispheric FC relative to inter-hemispheric FC. Note that each subplot captures a unique range of values, given differences in variability observed across neurotypical adults (colored distributions), patients, and networks. Full splits generally demonstrated more strongly-lateralized FC compared to partials, who fell within the control distributions. Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.

Lateralized expressions of functional modules

To further elucidate whole-brain differences in functional network topology—agnostic to any *a priori* assignment of individual brain regions to any given functional system—we employed data-

driven community detection techniques to cluster whole-brain patterns of FC into functional

modules (**Fig. 4**). The modular organization of both structural and functional brain networks is

thought to enable specialized information processing while retaining the ability to efficiently

integrate information between disparate neural systems[35,36]. In practice, modular decompositions

of FC reveal groups of brain regions that are strongly connected to each other while being weakly

connected (or not at all connected) to other groups, constituting coherent functional

communities[36,37]. Thus, we hypothesized that the breakdown of IFC in full splits would produce

lateralized network topologies, predominantly self-organized within one hemisphere or the other.

Here we use a variant of the *Q*-metric for modularity optimization well-suited to correlation

matrices (with negative weights)[38] and compute laterality indices for the resulting partitions

according to prior work[39], where a value of zero, for example, indicates that all functional modules

are perfectly bilateral (i.e., all of the constituent brain regions belonging to each community are

equally-balanced between the left and right hemispheres), and values closer to 1 indicate

progressively more lateralized topologies. Importantly, we note that modularity maximization is

sensitive to spatial scale, subject to a *resolution parameter*, γ[40]. We describe results using a

'default' $\gamma = 1$ below but provide further consideration of these methodological issues in the

Methods and **Extended Data Fig. 4**.

Figure 4. Modular decomposition of whole-brain functional connectivity. Spatial maps of optimal, consensus community assignments derived via data-driven modularity maximization at a default level of resolution ($\gamma = 1$). Regional coloring indicates assignment to a given functional module. Full splits generally exhibited more lateralized clustering of regions into modules relative to partial splits and healthy adult controls (see **Fig. 5** for quantification of module laterality).

As predicted, we observed much greater degrees of modular laterality in all full splits (**Fig. 5**). Communities with bihemispheric representations most often spanned visual and somatomotor regions—reflecting a mechanism through which synchronized, external sensory inputs can drive apparent functional coherence without direct callosal coupling. Interestingly, regions of the default

mode network also often showed some bilateral symmetry, both along the medial wall and lateral
temporal-parietal areas, although this may be due in part to signal being smoothed across the
midline. With respect to partial splits, patient EM with the most CC showed neurotypical, bilateral
patterns of network organization largely mirroring healthy controls. Patient BT also fell within the
control distribution—however, they showed a strikingly different pattern of bilateral modular
organization (**Fig. 4**). Here, functional communities appeared to dissociate along the Sylvian
fissure, with dorsal modules and ventral modules spanning the anteroposterior axis of the brain.
This individual had the longest gap between surgery and testing (~6 years), which may have
allowed for widespread network reorganization affording optimal utilization of the minimal
callosal fibers left intact. Overall, callosotomy's impact on the intrinsic modular organization of
the brain highlights the importance of transcallosal connections for the emergence of coherent
functional systems, while also suggesting a contributory role for sensory inputs enabling apparent
preservation of bilateral sensorimotor modules.

Figure 5. Functional modules are more strongly-lateralized in full splits. The modular
laterality index quantifies the extent to which functional communities exhibit bihemispheric
representations (values closer to zero) or are restricted to one hemisphere or the other (values closer
to one). At a default level of resolution ($\gamma = 1$), partial splits demonstrated bilateral network
topologies similar to healthy controls, whereas full splits were consistently more lateralized. Point
intervals under the control distribution give the HCP100 median (black dot) and the 80-95%
quantiles around the median (thick black lines and thin black lines, respectively).

Inter-hemispheric synchrony in network dynamics

The modular organization of the brain plays a critical role in governing large-scale functional dynamics, such that structural architectures orchestrate widespread co-fluctuations in whole-brain activity over time[41]. Such patterns of coordinated network dynamics have recently been elucidated using *edge time series*: in brief, the edge time series approach constitutes a temporal unwrapping of the zero-order Pearson correlation between any given pair of brain regions, capturing moment-to-moment changes in the extent to which regional pairs are co-fluctuating in activity[42]. Taking the root-sum-of-squares (RSS) across all edge time series at each timepoint yields summary estimates of whole-brain co-fluctuations in functional network dynamics over the course of a scan[43] (**Fig. 6a**).

Here we sought to test the hypothesis that structural connectivity via the CC is necessary for large-scale coordination of function. To that end, we separated edges within the left and right hemispheres and derived an RSS time course for each. We then computed correlations between left and right RSS to assess the degree to which dynamic co-fluctuations in brain function were synchronized across hemispheres. Among healthy controls with a fully-intact CC, we observed tight temporal coupling between left and right hemisphere RSS time courses (Median Spearman *r* = 0.82, *MAD* = 0.08). In contrast, all four full splits demonstrated significant reductions in RSS synchrony—each hemisphere appeared to be operating much more independently on the whole without callosal orchestration of function (**Fig. 6b**; see **Extended Data Fig. 5** for individual networks). This finding is consistent with the overall attenuation of inter-hemispheric FC and further indicates that brief moments of high-amplitude co-fluctuations (which normally encompass regions in both hemispheres and are thought to be structurally-mediated[41]) occur separately in each. Notably, both partial splits fell within the standard range of the HCP subjects—including patient BT. These results therefore strongly suggest a mechanistic role for CC structural connections in supporting widespread integration of inter-hemispheric network dynamics. This may be achieved with a mere fraction of intact fibers through the splenium and thus might be largely-independent of the presence of direct cortico-cortical connections.

Figure 6. Large-scale functional network dynamics are desynchronized in full splits. **(a)** An illustration of edge time series in the left and right hemispheres for patient BT (a partial split with 1 cm of intact splenium) and patient TJ (a full split). By taking the root-sum-of-squares (RSS) across all edges at each timepoint, we derived a summary estimate of the total amplitude of co-fluctuating brain activity. Note the bursty, high-amplitude co-fluctuations in each hemisphere that often coincide in time for BT, but not in TJ. **(b)** All four full splits exhibited reduced temporal synchrony (i.e., temporal correlations) between left/right hemisphere RSS time courses. Partial splits, by contrast, fell within the range of healthy adult controls. Point intervals under the control distribution give the HCP100 median (black dot) and the 80-95% quantiles around the median (thick black lines and thin black lines, respectively).

Discussion

Here we characterized variable disruptions of inter-hemispheric functional connectivity following complete versus partial transection of the corpus callosum. Full splits demonstrated more lateralized network topologies and desynchronized network dynamics which align with classical behavioral disconnection syndromes. All patients had intact anterior and posterior commissures along with other subcortical routes of inter-hemispheric communication: these findings therefore suggest that transcallosal connections are the essential pathways through which large-scale inter-hemispheric functional networks emerge. Contrary to our expectations, however, inter-

hemispheric functional coupling among partial splits remained intact—these patients additionally
showed *no* signs of behavioral disconnection, even when the vast majority of the callosum was
severed[26]. Thus, a fraction of callosal fibers in the splenium may be sufficient to orchestrate inter-
hemispheric functional synchrony and integrate behavior, largely independent of presumed
anatomical constraints. This poses a remarkable challenge to our classical topographic models of
callosal connectivity and function, motivating a novel view on inter-hemispheric structure-
function relationships: namely, that robust inter-hemispheric synchrony and information transfer
can be achieved via polysynaptic connectivity through a relatively small proportion of posterior
transcallosal connections, rather than direct structural connectivity between functional
homologues.

Complete callosotomy and structure-function relationships
Conventional wisdom in network neuroscience suggests that structural connectivity between brain
regions both constrains and enables function. Healthy adult brains frequently demonstrate strong
patterns of functional connectivity between mirrored, homotopic regions across the left and right
hemispheres—a phenomenon which is traditionally attributed to the spatial topography of
structural connections traversing the CC[10,11,15,44,45,46]. One study has suggested that, for as many as
90% of homotopic regional pairs, individual differences in callosal microstructure can predict
differences in the strength of functional connectivity between these regions[9]. It has become
increasingly-evident, however, that the relationships between structure and function are highly
complex and generally do not reflect simple, one-to-one interactions between directly-coupled
areas[47,48,49]. Split-brain patients present a unique opportunity to draw causal inferences regarding
the role of transcallosal structural connections and the extent to which inter-hemispheric functional
synchrony depends upon such connections. While callosal agenesis cases have highlighted that
inter-hemispheric functional connectivity and information transfer can emerge from a variety of
subcortical and non-callosal structural architectures[50,51], split-brain patients are individuals in
whom the brain developed with support of the CC and thus provide direct insights into the
functional consequences of losing this major structural scaffolding.

Previous investigations of pediatric callosotomy patients have provided an initial perspective on
the extent to which inter-hemispheric FC depends upon CC structural connectivity[16,17,18,19].

However, a number of methodological factors in these studies obfuscate straightforward
conclusions. For example, these patients often also have severe cognitive and developmental
deficits, and they require scans to be completed under sedation—the latter of which is known to
reduce dynamical complexity in brain activity and produce functional connectomes more rigidly
tied to existing anatomical connectivity[52]. It is further known that functional and structural brain
networks undergo profound developmental changes early in life[53,54,55,56]. In other words, neural
plasticity in these young individuals may still afford the long-term establishment of alternate inter-
hemispheric pathways similar to callosal agenesis patients[51], and in fact, it is well-documented that
disconnection syndromes in pediatric patients resolve over time[57,58]. This likely accounts for
differential reports of recovery in inter-hemispheric FC among these individuals[16,18,19].

The adult human brain also retains neuroplastic potential[59]; however, given the enormity of the
structural disconnection, adult callosotomy patients may lack the capacity to adapt their functional
architectures in such a way that allows complete recovery of communication between disconnected
brain regions. Our results clearly underscore this hypothesis: we observed dramatically diminished
functional integration between the cerebral hemispheres across multiple metrics. Whole-brain
patterns of FC became increasingly-more lateralized, both with respect to canonical resting-state
network definitions (**Figs. 2-3**) and the topological organization of functional modules revealed
using data-driven community detection (**Figs. 4-5**). Furthermore, we observed considerable
reductions in the extent to which multivariate co-fluctuations in function were synchronized
between hemispheres (**Fig. 6**), underscoring a role for the CC in orchestrating whole-brain network
dynamics. Importantly, however, there were some exceptions to these effects, which also follow
anatomical predictions. Limbic regions showed conserved bilateral FC (**Fig. 3**), for example—
these regions are known to be connected through the anterior commissure (AC) as well as the
CC[32,33,34], which was left intact in these patients. Functional connectivity between subcortical
regions also may be preserved. Nevertheless, it seems clear that such non-callosal and subcortical
pathways were insufficient to support more widespread patterns of inter-hemispheric synchrony,
perhaps contrary to previous proposals both in humans[20,21,22] and nonhuman primates (where AC
fibers are sufficient to sustain inter-hemispheric FC)[60]. Finally, an additional exception to inter-
hemispheric network disintegration was observed between some visual and sensorimotor areas.
We cannot fully discount the possibility that subcortical information transfer contributes to this

preserved synchrony; however, an alternative hypothesis may simply lie in the fact that visual and

somatosensory inputs to each hemisphere were synchronized during the resting-state scan—

producing *apparent* synchrony in function despite the behavioral disconnections observed in our

bedside testing protocol.

Preserved inter-hemispheric integration in partial splits

While our observations in full splits clearly emphasize the importance of CC connectivity in

orchestrating inter-hemispheric synchrony, partial splits revealed something much more

surprising. We initially predicted that partial splits would show specific deficits in inter-

hemispheric FC along an anteroposterior gradient corresponding to the extent of the callosal

transection (following previous characterizations of CC topography[10,11]). Contrary to our

expectation, however, both partial splits demonstrated levels of inter-hemispheric integration

largely in the range of healthy controls, regardless of the network metric considered (**Figs. 2-6**).

This was even observed for patient BT, in whom only ~1 cm of posterior callosal fibers were

spared. These remarkable outcomes imply that the functional topography of the CC may be much

more flexible than previously imagined. Moreover, there may be some critical threshold at which

full functional integration can be sustained given a sufficient proportion of intact fibers—even if

the vast majority of the callosum has been severed.

An intriguing extension of this hypothesis is that the *location* of the spared callosal fibers is also

critical. Previous cytological work has shown that anterior portions of the CC are primarily

populated by thin fibers, while fibers in the tail locally-vary in diameter—rostral portions of the

splenium have a relatively-higher density of thin fibers, while the posterior pole shows a higher

concentration of large-diameter fibers[12,13,14]. Such differences in fiber size and myelination along

the CC have been associated with differences in conduction velocity[12] and, when considered

alongside the topographic organization of the CC, may account for the 'fast' versus 'slow' action

of primary sensorimotor and association cortices, respectively. Spared posterior fibers as in BT

could therefore be more beneficial than spared anterior fibers, but from a sheer bandwidth

perspective, it is difficult to imagine how such a small proportion of fibers could account for

widespread integration of function and cognitive processing, regardless of the relative 'speed' of

these fibers. It is also worth noting that previous behavioral studies of partial callosotomy patients

have shown expected behavioral disconnection effects given the specific portions of the CC that
were severed, although no functional imaging was performed and these tests were largely
conducted within the first year post-operatively[61,62,63,64].

Accordingly, it appears that functional systems in BT have dramatically reorganized in the six
years following surgery. In brains that develop normally with the support of the CC, multisensory
integration zones in parietal cortex may serve as inter-hemispheric hubs[65,66]—we suggest that by
taking advantage of spared posterior callosal pathways connecting these regions, the processing of
various intra-hemispheric brain regions can still be integrated and exchanged along polysynaptic
routes running between hemispheres. The striking differences in functional network topology
observed in BT, with large modules spanning the anteroposterior axis of the brain (**Fig. 4**), are one
potential consequence of this reorganization, reflecting greater functional dependencies between
frontal and posterior areas affording more optimal usage of the spared splenial fibers. The
fundamental importance of these posterior inter-hemispheric connections also aligns with and may
further inform theories of consciousness[67,68]. While the purpose of this study is not to evaluate or
endorse any particular theory, it is nevertheless curious that the temporal-parietal-occipital 'hot
zone' of regions recently proposed to maximize the singular 'here-and-now' *qualia* of
consciousness[67,69] are likely dependent (in part) upon CC connections through the splenium. This
may explain why BT maintains an integrated conscious experience while full splits demonstrate
inter-hemispheric disintegration[26].

Limitations

An important limitation of this work is the lack of pre-operative or otherwise longitudinal
assessments of these patients. While our results generally align with prior pre-/post-operative
findings in children—namely, in suggesting a causal role for the CC in maintaining inter-
hemispheric functional connectivity—we can only conjecture about brain states both at baseline
and in the intervening months/years between surgery and testing. For example, as discussed above,
the atypical organization of bilateral functional modules in patient BT may be a result of various
neuroplastic processes occurring over the six years from surgery, including repurposing of residual
callosal fibers, subcortical/non-callosal commissures, and perhaps even intra-hemispheric
association tracts such as the superior and inferior longitudinal fasciculi. The question of network

reorganization, both intra- and inter-hemispherically, is of central interest, and future longitudinal
work is essential to characterize these processes as they evolve over time.

We also acknowledge the small number of callosotomy patients included in this study, each of
whom have limited neuroimaging data. While previous pediatric studies have recruited larger
samples, the availability of patients who underwent callosotomy in adulthood is extremely scarce.
It is often difficult to draw robust and reliable statistical inferences from neuroimaging data
acquired on such small samples, particularly in the presence of individual differences. However,
the marked differences observed between full splits and healthy controls suggest that we are not
parsing trivial effects on the scale of traditional neuroimaging analyses—rather, they reflect
outcomes drawn from fundamentally unique sampling distributions. Furthermore, relative to other
patient studies (that are often hampered by variability in the location of lesions, e.g., following
stroke), callosotomy is a highly-selective intervention. By severing a single white matter structure
in the brain without any damage to gray matter, callosotomy may afford more homogeneous
outcomes. It will be critical for future studies to include larger samples with more imaging data to
better capture and quantify the extent of neural variability in callosotomy patients. Nevertheless,
we believe that these results (wherein full splits consistently lie well beyond the expected range of
variability in healthy controls) are a sensible representation of the robust differences between splits
and healthy adult brains.

Conclusion
Following more than a half-century of behavioral neuroscience, split-brain patients continue to
provide startling insights into the fundamental nature of the human brain. The rise of network
neuroscience has similarly transformed our understanding of large-scale brain systems, tracing the
pathways relating brain structure to function and behavior. And yet, the human brain frequently
dismantles these conceits. Our results reveal a unique type of criticality in brain function—such
that a vanishingly-small proportion of posterior callosal fibers may be sufficient to support the
integration of inter-hemispheric functional brain networks and unified cognitive processing. It has
long been celebrated that the human brain is a remarkably adaptable complex system. These
findings challenge us to rethink our conventional assumptions about structure-function
relationships and the means through which the brain enables the mind.

                                          Methods

Patient recruitment

Six adult callosotomy patients (5 men, 1 woman; aged 19-60; **Fig. 1**, **Table 1**) were recruited

during routine post-operative follow-up visits to the Bethel Epilepsy Center (Bielefeld, Germany).

The study was approved by the Ethics Committee of the University of Münster,

Germany (2021–523-f-S). The patients or their legal representatives provided written informed

consent.

Callosotomy procedure

Callosotomies were performed with the assistance of intraoperative neuronavigation, with patients

in a supine position and the head inclined by roughly 30° (fixed in the Mayfield clamp). A straight

skin incision was made running slightly next to the midline on the non-language dominant

hemisphere. Craniotomy was then performed by making one drill hole in the midline at

approximately the level of the precentral sulcus. A bone flap was sawn out with an outline of 3 cm

x 5 cm. The dura was incised longitudinally and then opened with two wings under

microneurosurgical conditions—access to the inter-hemispheric fissure was successively created

while preserving the cortical bridging veins. Next, using a self-retaining retractor to carefully shift

the frontal lobe, the level of the corpus callosum was gradually exposed under sharp dissection

with micro scissors.

Dissection is often made considerably more difficult by the course of the pericallosal arteries,

which run parallel to the corpus callosum but are often tightly intertwined and twisted in the inter-

hemispheric fissure. In cases where the risk of vascular damage is too great, the surgery may be

halted short of a full disconnection (resulting in partial splits). However, under most conditions,

the inter-hemispheric fissure may be opened along the entire length of the knee of the corpus

callosum. The corpus callosum was then perforated into the lateral ventricle of the ipsilateral

hemisphere. Next, following the pericallosal artery and the anterior cerebral artery down to the

frontal base, the knee of the callosum was split in the midline, continuing backwards to about the

level of the foramen of Monroi. From this point, it is critical to protect the fornix of the language-

dominant hemisphere: to mitigate the risk of postoperative functional impairment, we carefully

ensure that the self-retaining retractor is not placed on the precentral gyrus within the inter-
hemispheric fissure. The disconnection can now be carried out completely into the splenium—the
severing of the splenium was performed subpially to protect all the cisternal vessels. Extensive
hemostasis is not usually necessary, and thus a ventricular drainage is not used. To conclude, the
dura mater was closed and the bone flap reinserted and fixed with titanium plates and screws.
Finally, a subgaleal drainage was inserted and the skin closed in two layers.
573
Healthy adult controls
We used open-access data from the Human Connectome Project 100 Unrelated Subjects (HCP100)
cohort (46 men, 54 women; $M_{Age} = 29.11$, $SD = 3.68$) as a robust benchmark for various network
features estimated in callosotomy patients. Participant recruitment and imaging procedures have
previously been detailed elsewhere[70,71]. In brief, data were collected over two sessions on a custom
3T Siemens Skyra with a 32-channel head coil. T_1-weighted structural scans were acquired using
a 3D magnetization prepared rapid gradient echo (MPRAGE) sequence ($TR = 2400$ ms, $TE = 2.14$
ms, $TI = 1000$ ms, flip angle = 8º, voxel size = 0.7 mm^3). Resting-state fMRI was collected with
eyes open over four ~14-minute scans, using a multiband echo-planar imaging (EPI) sequence
sensitive to the blood-oxygen-level-dependent (BOLD) contrast ($TR = 720$ ms, $TE = 33.1$ ms, flip
angle = 52º, voxel size = 2 mm^3, multiband factor = 8). All participants gave written informed
consent for a protocol approved by the Washington University Institutional Review Board. All
imaging data were downloaded in their raw (unprocessed) state.
587
Patient neuroimaging procedures
Patient data were collected on a 3T Siemens Vida at the Bethel Epilepsy Center using a 32-channel
head coil. T_1-weighted structural scans were acquired using a 3D MPRAGE sequence ($TR = 1900$
ms, $TE = 2.47$ ms, $TI = 900$ ms, flip angle = 8º, voxel size = 0.8 mm^3). Resting-state fMRI was
collected with eyes open using a single-shot EPI sequence sensitive to the BOLD contrast ($TR = $
2000 ms, $TE = 30$ ms, flip angle = 75º, voxel size = 3 x 3 x 3.30 mm, iPAT/GRAPPA factor = 2).
594
595 We note that each patient varied somewhat in the amount of resting-state data obtained (from ~8-
596 20 minutes total) due to differences in clinical scheduling and other imaging protocols not relevant
597 to the current analysis. This information is summarized in **Extended Data Fig. 1**.

fMRI preprocessing

All data (from both patients and healthy controls) were run through a unified preprocessing pipeline. Initial preprocessing relied on functions from the Statistical Parametric Mapping 12 software (SPM12, Wellcome Trust Centre for Neuroimaging, London) in Matlab and Advanced Normalization Tools (ANTs)[72]. Resting-state fMRI data were realigned and unwarped to correct for head motion and motion x susceptibility artifact as well as slice-time corrected. Anatomical images were denoised (ANTs' `DenoiseImage` with a spatially-adaptive Gaussian filter) and submitted to ANTs' cortical thickness pipeline, which includes N4 bias field correction and high-quality image segmentation and brain extraction[73]. For some patients, the brain extraction mask included extracerebral components: this was corrected by manually eroding the extraction mask and re-applying it to the anatomical scan when necessary. Image coregistration parameters were subsequently estimated between an N3 bias-corrected mean BOLD reference and the skull-stripped T_1 image via `antsRegistration` (including both linear and nonlinear transformations to minimize residual signal deformations, cf. 'SyN distortion correction' as implemented by the widely-used fMRIPrep and QSIPrep pipelines).

Subsequent spatial normalization to a standard stereotaxic space presented a nontrivial problem for patient data—in the absence of a corpus callosum, generic registration routines (with both linear and nonlinear stages) will frequently pull structures such as the cingulate gyrus into the space occupied by the CC in the MNI template image, which misaligns brain regions relative to their location in our functional parcellation. We mitigated this problem in two ways. First, we implemented cost-function masking during the nonlinear step to minimize the influence of callosal regions that were transected by the surgery: these masks were specified in the fixed image space (i.e., the MNI152NLin6Asym template, as provided by the FMRIB Software Library, FSL[74,75]) because we lacked pre-operative data for each patient, and CC segments were defined by the JHU-DTI white matter atlas[76,77] (also provided in FSL). For patient EM, the genu was masked out; for patient BT, all but the splenium was masked out; and for all full splits, the entire cross-section of the callosum was masked out. Second, in addition to these masks, we specified a series of registration metric objectives at the nonlinear step. As is typical for `antsRegistration`, we estimated similarity between the MNI template and T_1 images using both ANTs' neighborhood cross-correlation and Mattes mutual information similarity metrics. We also provided smoothed

segmentation posteriors derived from the ANTs cortical thickness pipeline (six in total: white
matter, cortical gray matter, CSF, deep/subcortical gray matter, brain stem, and cerebellum) and
estimated similarity between the MNI template segments and patient segments using a mean-
squares objective, with cortical white matter and CSF segments downweighted in importance
relative to the other four segments (by a factor of one third). Thus, the nonlinear step was
comprised of eight similarity metrics. The transformation was estimated using BSplineSyN to
allow for smoother deformations without sacrificing accuracy.

Ultimately, we concatenated and applied the estimated coregistration and spatial normalization
parameters to all fMRI data in a single interpolation via `antsApplyTransforms`, resampled into
$2\ mm^3$ isotropic voxels. Registration quality was visually assessed to ensure appropriate mapping
to MNI space. Functional images were then smoothed using a 3D Gaussian kernel with FWHM =
$5\ mm^3$. Prior to functional connectivity estimation, several additional preprocessing steps were
performed using in-house Matlab code: this included global signal scaling (median = 1000), linear
detrending, and nuisance regression. The nuisance regression model followed a '36-parameter'
(36P) approach including the six motion regressors from realignment, the first eigenvariate of
white matter signal, the first eigenvariate of CSF signal, and mean global signal—along with their
squares, their first derivatives (via backwards-differentiation), and the squares of the derivatives.
All nuisance regressors were detrended to match the BOLD data. We further augmented the design
matrix of the nuisance model with a series of binary spike regressors: one for each image volume
where the relative framewise displacement in head motion (accounting for both translation and
rotation) was ≥ 0.50 mm. This '36P + spike regression' model has importantly been shown to
outperform alternative confound regression strategies (e.g., motion scrubbing and ICA-based
methods) and provide the best control for motion artifact in functional connectivity analyses[27,28].
Patients generally showed no more head motion than HCP subjects; for reference, we provide a
summary of head motion parameters and the frequency of motion spikes for each patient in
Extended Data Fig. 1.

Functional connectivity estimation and network laterality
Functional network nodes were defined by parcellating the brain into 400 cortical regions and 32
subcortical regions, according to the Schaefer[29] and Tian[30] atlases, respectively. We extracted a

summary time course from each parcel by taking the first eigenvariate over time[78]. These regional

time series were bandpass-filtered to the range of 0.01-0.08 Hz. We then constructed 432 x 432

FC matrices by estimating the zero-order Pearson correlation between all pairs of regional time

series. For both HCP100 subjects and callosotomy patients with more than one resting-state scan,

runs were concatenated prior to FC estimation.

We sought to quantify differences in inter-hemispheric functional connectivity (IFC) between

callosotomy patients and healthy controls from a number of network neuroscience perspectives.

Pursuant to this, we began with an initial, breadth-first assessment of IFC within each of the seven

canonical resting-state networks defined by our cortical parcellation[29,31], as well as regions

spanning the subcortex[30]. Similarly to studies of functional lateralization in task-based fMRI, we

derived a *network laterality index* as a simple means of summarizing and quantifying the *relative*

magnitude of IFC as compared to the magnitude of intra-hemispheric FC. Each FC matrix was

first thresholded at an uncorrected level of $p < .001$ to avoid overly-conservative and highly-

nonlinear effects of multiple comparisons correction on the distribution of edge weights, while

simultaneously minimizing the influence of weak (potentially spurious) connections. We then

Fisher-Z transformed the thresholded FC matrices and, for each individual and for each network,

estimated a laterality index, *LI*, as:

$$LI = \frac{(\sum FC_{Intra-LH} + \sum FC_{Intra-RH}) - \sum FC_{Inter}}{\sum FC_{Intra-LH} + \sum FC_{Intra-RH} + \sum FC_{Inter}}$$

Intra-hemispheric connections within the left and right hemispheres are necessarily summed in the

numerator given that there are, by definition, twice as many inter-hemispheric connections as each

hemisphere alone. The result is a ratio on the interval of [-1, 1] that captures the extent to which

each canonical resting-state network is predominantly connected intra- or inter-hemispherically:

in the limit, a value of -1 would suggest that there is *no* intra-hemispheric FC and the network is

solely comprised of inter-hemispheric connections, while a value of 1 would indicate the reverse.

A value of 0 therefore represents a network that is perfectly balanced in the magnitude of intra- vs.

inter-hemispheric connections. Note that the results presented in **Fig. 3** are with respect to

positively-correlated edges only; we provide a comparison of anti-correlated edge lateralization in

Extended Data Fig. 3.

Community detection and modularity

While the above approach is useful for considering differences among canonical functional

networks, these systems were defined in healthy adult brains—it is therefore perhaps dubious to

assume that these networks follow a similar spatial organization in callosotomy patients. Data-

driven community detection techniques afford assessments of functional network topology

unbeholden to prior assumptions about specific brain regions and their canonical network

membership[36,37]. This is achieved via decomposition of FC matrices using *modularity*

maximization algorithms, which group regions into *functional modules*: these represent regions

that are generally more functionally-coupled to each other than they are connected to other regions.

The optimization objective is subject to a *quality* metric, Q—unlike our previous laterality analysis

(which required separating out positive and negative edges), this can be estimated over FC matrices

comprised of both positive and negative correlations[38]. We performed this decomposition using

the `community_louvain` function from the Brain Connectivity Toolbox

(https://sites.google.com/site/bctnet/) with asymmetric treatment of negative edge weights[38,79].

Importantly, modularity maximization is sensitive to spatial scale, subject to a *resolution*

parameter, γ[40]. Smaller values of γ typically yield coarse partitions with a small number of large

communities; larger values of γ, by contrast, afford more fine-grained partitions. Because the

solution of this decomposition is dependent upon one's choice for the resolution parameter, we

tested a range of values from $\gamma = [0.10, 2]$ (in steps of 0.10). To further account for variability in

community assignments (e.g., due to algorithmic stochasticity arising from random initializations),

we estimated modularity 1000 times for each value of γ —optimal consensus partitions were

obtained using the `consensus_similarity` function from the Network Community Toolbox

(https://commdetect.weebly.com/).

In similar fashion to the network laterality indices described above, we derived a *module laterality*

index for the resulting partitions according to prior work[39]. This metric is limited to the [0, 1]

interval, where a value of 0, for example, indicates that all functional communities are perfectly

bilateral (i.e., all of the constituent brain regions belonging to each community are equally-

represented between the left and right hemispheres), and values closer to 1 indicate progressively
more lateralized topologies. We note that in this case, more fine-grained partitions may be
beneficial for assessing modular laterality—nevertheless, the results shown in **Figs. 4-5** reflect a
'default' $\gamma = 1$ and we further illustrate in **Extended Data Fig. 4** that full splits rapidly diverge
from controls even at smaller values of γ (while partial splits and healthy controls remain relatively
consistent, regardless of γ).

Inter-hemispheric network dynamics

Finally, we used edge time series (ETS)[42] to assess the extent to which *many* brain regions in the
left and right hemispheres dynamically co-fluctuate in synchrony. The ETS approach has been
used to obtain moment-to-moment, time-resolved estimates of FC and characterize the nature of
high-amplitude events[43] that occur when multivariate BOLD signals simultaneously fluctuate
together over brief intervals of time—these bursts of synchronized network dynamics are sufficient
to recapitulate 'static' FC estimates over the course of a scan[43], and importantly, they are
independent of head motion[42]. Current evidence instead suggests these high-amplitude co-
fluctuations are governed, in part, by the modular organization of the brain and its underlying
structural architecture[41]. An ETS can be computed by taking a pair of BOLD time series from two
regions, *Z*-scoring them, and taking the elementwise product of the signals at each timepoint—the
mean of an ETS is equivalent to the zero-order Pearson correlation between the two time series,
and thus ETS constitute a temporal-unwrapping of FC over the scan[42].

Here, we first separated out all regions in the left hemisphere and all regions in the right
hemisphere. We constructed a T x 23220 ETS matrix for each hemisphere, where T gives the
number of timepoints and 23220 gives the total number of edges within each hemisphere. A
summary time course quantifying the total amplitude of intra-hemispheric co-fluctuations at any
given moment (i.e., across *all* edges) was obtained by taking the root-sum-of-squares (RSS) at
each timepoint[43]. We then computed the Spearman correlation between the left hemisphere RSS
and right hemisphere RSS time series to capture the extent to which network dynamics in each
hemisphere were synchronized.

Data and code availability

Patient data will be made available upon reasonable request; all relevant processing and analysis code will be uploaded to a Github repository.

References

1. Bassett DS, Sporns O. Network neuroscience. *Nat Neurosci* **20**, 353-364 (2017).

2. Honey CJ*, et al.* Predicting human resting-state functional connectivity from structural connectivity. *Proc Natl Acad Sci U S A* **106**, 2035-2040 (2009).

3. van den Heuvel MP, Mandl RC, Kahn RS, Hulshoff Pol HE. Functionally linked resting-state networks reflect the underlying structural connectivity architecture of the human brain. *Hum Brain Mapp* **30**, 3127-3141 (2009).

4. Fukushima M, Betzel RF, He Y, van den Heuvel MP, Zuo XN, Sporns O. Structure-function relationships during segregated and integrated network states of human brain functional connectivity. *Brain Struct Funct* **223**, 1091-1106 (2018).

5. Hermundstad AM*, et al.* Structurally-constrained relationships between cognitive states in the human brain. *PLoS Comput Biol* **10**, e1003591 (2014).

6. Hermundstad AM*, et al.* Structural foundations of resting-state and task-based functional connectivity in the human brain. *Proc Natl Acad Sci U S A* **110**, 6169-6174 (2013).

7. Damoiseaux JS, Greicius MD. Greater than the sum of its parts: a review of studies combining structural connectivity and resting-state functional connectivity. *Brain Struct Funct* **213**, 525-533 (2009).

8. Honey CJ, Thivierge JP, Sporns O. Can structure predict function in the human brain? *Neuroimage* **52**, 766-776 (2010).

9. Mollink J*, et al.* The spatial correspondence and genetic influence of interhemispheric connectivity with white matter microstructure. *Nat Neurosci* **22**, 809-819 (2019).

10. Hofer S, Frahm J. Topography of the human corpus callosum revisited--comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. *Neuroimage* **32**, 989-994 (2006).

11. Liu Y*, et al.* Connectivity-Based Topographical Changes of the Corpus Callosum During Aging. *Front Aging Neurosci* **13**, 753236 (2021).

12. Horowitz A, Barazany D, Tavor I, Bernstein M, Yovel G, Assaf Y. In vivo correlation between axon diameter and conduction velocity in the human brain. *Brain Struct Funct* **220**, 1777-1788 (2015).

796 13. Aboitiz F, Lopez J, Montiel J. Long distance communication in the human brain: timing
797   constraints for inter-hemispheric synchrony and the origin of brain lateralization. *Biol Res*
798   **36**, 89-99 (2003).
799

800 14. Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus
801   callosum. *Brain Res* **598**, 143-153 (1992).
802

803 15. Stark DE*, et al.* Regional variation in interhemispheric coordination of intrinsic
804   hemodynamic fluctuations. *J Neurosci* **28**, 13754-13764 (2008).
805

806 16. Hung SC*, et al.* Early recovery of interhemispheric functional connectivity after corpus
807   callosotomy. *Epilepsia* **60**, 1126-1136 (2019).
808

809 17. Johnston JM*, et al.* Loss of resting interhemispheric functional connectivity after
810   complete section of the corpus callosum. *J Neurosci* **28**, 6453-6458 (2008).
811

812 18. Roland JL*, et al.* On the role of the corpus callosum in interhemispheric functional
813   connectivity in humans. *Proc Natl Acad Sci U S A* **114**, 13278-13283 (2017).
814

815 19. Pizoli CE*, et al.* Resting-state activity in development and maintenance of normal brain
816   function. *Proc Natl Acad Sci U S A* **108**, 11638-11643 (2011).
817

818 20. Nomi JS*, et al.* Diffusion weighted imaging evidence of extra-callosal pathways for
819   interhemispheric communication after complete commissurotomy. *Brain Struct Funct*
820   **224**, 1897-1909 (2019).
821

822 21. Uddin LQ*, et al.* Residual functional connectivity in the split-brain revealed with resting-
823   state functional MRI. *Neuroreport* **19**, 703-709 (2008).
824

825 22. Marcantoni I*, et al.* Interhemispheric functional connectivity: an fMRI study in
826   callosotomized patients. *Front Hum Neurosci* **18**, 1363098 (2024).
827

828 23. Gazzaniga MS. Forty-five years of split-brain research and still going strong. *Nat Rev*
829   *Neurosci* **6**, 653-659 (2005).
830

831 24. Volz LJ, Gazzaniga MS. Interaction in isolation: 50 years of insights from split-brain
832   research. *Brain* **140**, 2051-2060 (2017).
833

834 25. Miller MB, Volz LJ, Simonson JM, Gazzaniga MS. Split-brain patients: A clinical versus
835   experimental perspective. In: *Cerebral Asymmetries* (eds Papagno C, Corballis P).
836   Elsevier BV (in press).
837

838 26. Bekir S*, et al.* [Preprint] No disconnection syndrome after near-complete callosotomy.
839   *bioRxiv*, (2024).
840

841   27.   Ciric R*, et al.* Benchmarking of participant-level confound regression strategies for the
842   control of motion artifact in studies of functional connectivity. *Neuroimage* **154**, 174-187
843   (2017).
844

845   28.   Parkes L, Fulcher B, Yucel M, Fornito A. An evaluation of the efficacy, reliability, and
846   sensitivity of motion correction strategies for resting-state functional MRI. *Neuroimage*
847   **171**, 415-436 (2018).
848

849   29.   Schaefer A*, et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
850   Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114 (2018).
851

852   30.   Tian Y, Margulies DS, Breakspear M, Zalesky A. Topographic organization of the human
853   subcortex unveiled with functional connectivity gradients. *Nat Neurosci* **23**, 1421-1432
854   (2020).
855

856   31.   Yeo BT*, et al.* The organization of the human cerebral cortex estimated by intrinsic
857   functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).
858

859   32.   Catani M, Schotten M. *Atlas of Human Brain Connections*. Oxford University Press
860   (2012).
861

862   33.   Patel MD, Toussaint N, Charles-Edwards GD, Lin JP, Batchelor PG. Distribution and
863   fibre field similarity mapping of the human anterior commissure fibres by diffusion
864   tensor imaging. *MAGMA* **23**, 399-408 (2010).
865

866   34.   Di Virgilio G, Clarke S, Pizzolato G, Schaffner T. Cortical regions contributing to the
867   anterior commissure in man. *Exp Brain Res* **124**, 1-7 (1999).
868

869   35.   Bassett DS, Gazzaniga MS. Understanding complexity in the human brain. *Trends Cogn*
870   *Sci* **15**, 200-209 (2011).
871

872   36.   Sporns O, Betzel RF. Modular Brain Networks. *Annu Rev Psychol* **67**, 613-640 (2016).
873

874   37.   Zamani Esfahlani F*, et al.* Modularity maximization as a flexible and generic framework
875   for brain network exploratory analysis. *Neuroimage* **244**, 118607 (2021).
876

877   38.   Rubinov M, Sporns O. Weight-conserving characterization of complex functional brain
878   networks. *Neuroimage* **56**, 2068-2079 (2011).
879

880   39.   Doron KW, Bassett DS, Gazzaniga MS. Dynamic network structure of interhemispheric
881   coordination. *Proc Natl Acad Sci U S A* **109**, 18661-18668 (2012).
882

883   40.   Fortunato S, Barthelemy M. Resolution limit in community detection. *Proc Natl Acad Sci*
884   *U S A* **104**, 36-41 (2007).
885

886 41. Pope M, Fukushima M, Betzel RF, Sporns O. Modular origins of high-amplitude
887 cofluctuations in fine-scale functional connectivity dynamics. *Proc Natl Acad Sci U S A*
888 **118**, (2021).
889

890 42. Faskowitz J, Esfahlani FZ, Jo Y, Sporns O, Betzel RF. Edge-centric functional network
891 representations of human cerebral cortex reveal overlapping system-level architecture.
892 *Nat Neurosci* **23**, 1644-1654 (2020).
893

894 43. Zamani Esfahlani F*, et al.* High-amplitude cofluctuations in cortical activity drive
895 functional connectivity. *Proc Natl Acad Sci U S A* **117**, 28393-28401 (2020).
896

897 44. Mancuso L*, et al.* The homotopic connectivity of the functional brain: a meta-analytic
898 approach. *Sci Rep* **9**, 3346 (2019).
899

900 45. Jarbo K, Verstynen T, Schneider W. In vivo quantification of global connectivity in the
901 human corpus callosum. *Neuroimage* **59**, 1988-1996 (2012).
902

903 46. De Benedictis A*, et al.* New insights in the homotopic and heterotopic connectivity of the
904 frontal portion of the human corpus callosum revealed by microdissection and diffusion
905 tractography. *Hum Brain Mapp* **37**, 4718-4735 (2016).
906

907 47. Seguin C, Mansour LS, Sporns O, Zalesky A, Calamante F. Network communication
908 models narrow the gap between the modular organization of structural and functional
909 brain networks. *Neuroimage* **257**, 119323 (2022).
910

911 48. Suarez LE, Markello RD, Betzel RF, Misic B. Linking Structure and Function in
912 Macroscale Brain Networks. *Trends Cogn Sci* **24**, 302-315 (2020).
913

914 49. Misic B*, et al.* Network-Level Structure-Function Relationships in Human Neocortex.
915 *Cereb Cortex* **26**, 3285-3296 (2016).
916

917 50. Paul LK*, et al.* Agenesis of the corpus callosum: genetic, developmental and functional
918 aspects of connectivity. *Nature Reviews Neuroscience* **8**, 287-299 (2007).
919

920 51. Tyszka JM, Kennedy DP, Adolphs R, Paul LK. Intact bilateral resting-state networks in
921 the absence of the corpus callosum. *J Neurosci* **31**, 15154-15162 (2011).
922

923 52. Barttfeld P, Uhrig L, Sitt JD, Sigman M, Jarraya B, Dehaene S. Signature of
924 consciousness in the dynamics of resting-state brain activity. *Proc Natl Acad Sci U S A*
925 **112**, 887-892 (2015).
926

927 53. Tooley UA*, et al.* The Age of Reason: Functional Brain Network Development during
928 Childhood. *J Neurosci* **42**, 8237-8251 (2022).
929

930 54. Sydnor VJ*, et al.* Neurodevelopment of the association cortices: Patterns, mechanisms,
931 and implications for psychopathology. *Neuron* **109**, 2820-2846 (2021).

55. Reynolds JE, Grohs MN, Dewey D, Lebel C. Global and regional white matter development in early childhood. *Neuroimage* **196**, 49-58 (2019).

56. Whitaker KJ*, et al.* Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. *Proc Natl Acad Sci U S A* **113**, 9105-9110 (2016).

57. Lassonde M, Sauerwein H, Chicoine AJ, Geoffroy G. Absence of disconnexion syndrome in callosal agenesis and early callosotomy: brain reorganization or lack of structural specificity during ontogeny? *Neuropsychologia* **29**, 481-495 (1991).

58. Graham D, Tisdall MM, Gill D. Corpus callosotomy outcomes in pediatric patients: A systematic review. *Epilepsia* **57**, 1053-1068 (2016).

59. Fuchs E, Flugge G. Adult neuroplasticity: more than 40 years of research. *Neural Plast* **2014**, 541870 (2014).

60. O'Reilly JX*, et al.* Causal effect of disconnection lesions on interhemispheric functional connectivity in rhesus monkeys. *Proc Natl Acad Sci U S A* **110**, 13982-13987 (2013).

61. Funnell MG, Corballis PM, Gazzaniga MS. Cortical and subcortical interhemispheric interactions following partial and complete callosotomy. *Arch Neurol* **57**, 185-189 (2000).

62. Gazzaniga MS, Freedman H. Observations on visual processes after posterior callosal section. *Neurology* **23**, 1126-1130 (1973).

63. Risse GL, Gates J, Lund G, Maxwell R, Rubens A. Interhemispheric transfer in patients with incomplete section of the corpus callosum. Anatomic verification with magnetic resonance imaging. *Arch Neurol* **46**, 437-443 (1989).

64. Gazzaniga MS. Interhemispheric integration. In: *Neurobiology of Neocortex* (eds Rakic P, Singer W). John Wiley & Sons (1988).

65. van den Heuvel MP, Sporns O. Rich-club organization of the human connectome. *J Neurosci* **31**, 15775-15786 (2011).

66. Hagmann P*, et al.* Mapping the structural core of human cerebral cortex. *PLoS Biol* **6**, e159 (2008).

67. Koch C, Massimini M, Boly M, Tononi G. Neural correlates of consciousness: progress and problems. *Nat Rev Neurosci* **17**, 307-321 (2016).

68. Melloni L*, et al.* An adversarial collaboration protocol for testing contrasting predictions of global neuronal workspace and integrated information theory. *PLoS One* **18**, e0268577 (2023).

979   69.   Koch C, Massimini M, Boly M, Tononi G. Posterior and anterior cortex - where is the
980   difference that makes the difference? *Nat Rev Neurosci* **17**, 666 (2016).
981

982   70.   Van Essen DC*, et al.* The WU-Minn Human Connectome Project: an overview.
983   *Neuroimage* **80**, 62-79 (2013).
984

985   71.   Van Essen DC*, et al.* The Human Connectome Project: a data acquisition perspective.
986   *Neuroimage* **62**, 2222-2231 (2012).
987

988   72.   Tustison NJ*, et al.* The ANTsX ecosystem for quantitative biological and medical
989   imaging. *Sci Rep* **11**, 9068 (2021).
990

991   73.   Tustison NJ*, et al.* Large-scale evaluation of ANTs and FreeSurfer cortical thickness
992   measurements. *Neuroimage* **99**, 166-179 (2014).
993

994   74.   Evans AC, Janke AL, Collins DL, Baillet S. Brain templates and atlases. *Neuroimage* **62**,
995   911-922 (2012).
996

997   75.   Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. Fsl. *Neuroimage* **62**,
998   782-790 (2012).
999

1000   76.   Hua K*, et al.* Tract probability maps in stereotaxic spaces: analyses of white matter
1001   anatomy and tract-specific quantification. *Neuroimage* **39**, 336-347 (2008).
1002

1003   77.   Wakana S*, et al.* Reproducibility of quantitative tractography methods applied to cerebral
1004   white matter. *Neuroimage* **36**, 630-644 (2007).
1005

1006   78.   Friston KJ, Rotshtein P, Geng JJ, Sterzer P, Henson RN. A critique of functional
1007   localisers. *Neuroimage* **30**, 1077-1087 (2006).
1008

1009   79.   Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and
1010   interpretations. *Neuroimage* **52**, 1059-1069 (2010).
1011

1012

1013

1014

1015

1016

1017

1018

1019

Patient	Scan num.	Scan duration		Head motion estimates					
		Minutes	Frames	Min FWD	Max FWD	Med FWD	MAD FWD	Num. spikes	Spike %
EM	1	8	240	0.03	2.44	0.13	0.05	9	3.75
BT	1	7	210	0.04	1.09	0.19	0.07	11	5.24
BT	2	7	210	0.05	1.34	0.25*	0.08	9	4.29
HY	1	12	360	0.02	0.44	0.14	0.05	0	0.00
DT	1	12	360	0.02	10.74*	0.15	0.05	31	8.61*
TJ	1	7	210	0.02	0.59	0.13	0.04	2	0.95
TJ	2	7	210	0.03	0.38	0.11	0.04	0	0.00
LJ	1	7	210	0.06	0.55	0.20	0.04	2	0.95
LJ	2	7	210	0.04	0.38	0.18	0.04	0	0.00
LJ	3	7	210	0.06	0.56	0.20	0.05	3	1.43

*Denotes head motion estimate exceeding normal range, see corresponding figure(s) below.

Extended Data Fig. 1. Scan times and head motion estimates across callosotomy patients. Due to differences in clinical scheduling and other imaging protocols not relevant to the current analysis, patients varied in resting-state fMRI scan time. Here we summarize these differences and provide various estimates of head motion for each scan. Motion parameters are given with respect to *framewise displacement* (FWD)—the amount of total head motion (in millimeters, considering both translational displacements and rotations) observed from one image volume to the next. Because FWD is a time series, we provide the minimum, maximum, and median FWD over each scan (with the median absolute deviation, *MAD*, over time). We also denote the number of motion spikes, defined as any frame with FWD $\geq$ 0.50 mm, and the percentage of frames contaminated with motion spikes—these were used to define individual spike regressors for nuisance signal regression[27,28]. Patients generally did not show any more head motion than healthy adult controls (see visualized distributions). Note that in rare cases where patients exceeded the range of controls in one metric, they were typically within the standard range of other metrics. Point intervals under

1036 the control distributions give the median (black dot) and the 80-95% quantiles around the median

1037 (thick black lines and thin black lines, respectively).

1038

1039

Extended Data Fig. 2. Multivariate similarity in spatial patterns of functional connectivity.

As an initial estimate of the extent to which full vs. partial callosotomy disrupted intra- vs. inter-

hemispheric functional connectivity (FC), we binarized FC matrices at a threshold of $z(r) > 0$

(where $z(r)$ refers to the Fisher-Z transformed correlation coefficients for each network edge) and

computed the Dice coefficient between each patient and each individual healthy control—this

statistic ranges from [0,1] and quantifies the degree of similarity in two patterns of FC (liberally,

whether the same edges show *any* FC > 0, regardless of the exact magnitude of those connections).

The distributions for each patient thus show the range of similarity between a given patient and

each control. Patient EM (with the most corpus callosum intact) is generally most similar to

controls; full splits, on the other hand, show markedly dissimilar patterns of inter-hemispheric FC

relative to intra-hemispheric FC. Point intervals under each distribution give the median Dice

similarity (black dot) and the 80-95% quantiles around the median (thick black lines and thin black

lines, respectively). Abbreviations: LH, Left Hemisphere; RH, Right Hemisphere.

1053

 Extended Data Fig. 3. Anti-correlated functional connectivity is generally not more
 lateralized in callosotomy patients. Network laterality indices across seven canonical resting-
 state systems[29,31] and subcortical regions[30]. To contrast against **Fig. 3**, here we estimate
 lateralization of negative (anti-correlated) connections. Values closer to 0 indicate an equal balance
 between intra- and inter-hemispheric (negative) FC, while values closer to -1 or 1 indicate more
 inter-hemispheric FC intra-hemispheric (negative) FC, respectively. Note that, unlike in **Fig. 3**,
 most distributions skew to the left (indicating that negative FC is predominantly inter-hemispheric,
 i.e., not lateralized), and both full and partial splits frequently fall within the range of HCP controls.
 Point intervals for each control distribution give the HCP100 median (black dot) and the 80-95%
 quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations:
 FC; Functional Connectivity; SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention
 Network; SalVentAttn, Salience/Ventral Attention Network.

Extended Data Fig. 4. Lateralization of functional modules is readily-apparent in full splits regardless of resolution. Modularity maximization algorithms are sensitive to a spatial resolution parameter, γ[40]. A value of $\gamma = 1$ is often treated as a 'default' level of resolution, whereas values of $\gamma < 1$ or $\gamma > 1$ produce progressively more coarse or fine-grained solutions, respectively. Here we show that differences in modular laterality between patients and healthy adult controls are not strictly-dependent upon the chosen value of γ. We performed 1000 iterations of modularity maximization at each of 20 levels of γ (ranging from [0.10, 2] in steps of 0.10), derived an optimal consensus partition at each level (see **Methods**), and estimated a laterality index for the resulting partitions[39] (such that values closer to 1 indicate more lateralized network topologies). Full splits (solid-colored lines) rapidly diverged from the distribution of healthy controls (quantile intervals shown in gradations of purple, with the solid black line representing the median) demonstrating more lateralized organization of functional modules, even at coarse levels of $\gamma < 1$. Partial splits (dotted colored lines), by contrast, consistently remained within the control distribution.

Extended Data Fig. 5. Estimates of large-scale dynamic functional synchrony by network. Decomposing whole-brain comparisons of RSS synchrony in **Fig. 6b** to individual networks reveals nuanced differences between patients and neurotypical adults (colored distributions). As before, partial splits consistently fall within the range of HCP controls. Full splits generally show more desynchronized network dynamics in higher-order cognitive systems, but comparatively-neurotypical levels of inter-hemispheric synchrony in limbic and subcortical networks (which are not strictly dependent upon structural connections through the corpus callosum, also reflecting **Fig. 3**). Point intervals under the control distributions give the HCP100 median (black dot) and 80-95% quantiles around the median (thick black lines and thin black lines, respectively). Abbreviations: RSS, Root-Sum-Of-Squares (see **Methods**); SomMot, SomatoMotor Network; DorsAttn, Dorsal Attention Network; SalVentAttn, Salience/Ventral Attention Network.