cxcl18b-defined transitional state-specific nitric oxide drives injury-induced

Müller glia cell-cycle re-entry in the zebrafish retina

Aojun Ye[1,3], Shuguang Yu[2], Meng Du[4], Dongming Zhou[4], Jie He[2,*] and Chang

Chen[1,3,5,*]

[1]National Laboratory of Biomacromolecules, CAS Center for Excellence in

Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing

100101, China

[2]Institute of Neuroscience, State Key Laboratory of Neuroscience, CAS Center for

Excellence in Brain Science and Intelligence Technology, Chinese Academy of

Sciences, Shanghai 200031, China

[3]University of Chinese Academy of Sciences, Beijing 100049, China

[4]Tianjin Medical University, Tianjin 300070, China

[5]Lead contact

*Correspondence: jiehe@ion.ac.cn, changchen@moon.ibp.ac.cn

Abstract:

In lower vertebrates, retinal Müller glia (MG) exhibit a life-long capacity of cell-

cycle re-entry to regenerate neurons following the retinal injury. However, the

mechanism driving such injury-induced MG cell-cycle re-entry remains incompletely

understood. Combining single-cell transcriptomic analysis and *in-vivo* clonal analysis,

we identified previously undescribed *cxcl18b*-defined MG transitional states as

essential routes towards MG proliferation following green/red cone (G/R cone)

ablation. Inflammation blockage abolished the triggering of these transitional states,

which expressed the gene modules shared by cells of the ciliary marginal zone (CMZ),

where life-long adult neurogenesis takes place. Functional studies of the redox

properties of these transitional states further demonstrated the regulatory role of nitric

oxide (NO) produced by *Nos2b* in injury-induced MG proliferation. Finally, we

developed a viral-based strategy to specifically disrupt *nos2b* in *cxcl18b*-defined MG

transitional states and revealed the effect of transitional state-specific NO signaling.

Our findings elucidate the precision redox mechanism underlying injury-induced MG

cell-cycle re-entry, providing insights into species-specific mechanisms for vertebrate

retina regeneration.

Key Words: Müller glia, zebrafish retina, cell cycle re-entry, nitric oxide, precision

redox, *cxcl18b*.

Introduction:

Unlike mammalian counterparts, retinal Müller glia (MG) in aquatic and

amphibian species can enter injury-induced regeneration program(**Goldman, 2014;**

Lahne et al., 2020b; Wan and Goldman, 2016), which unfolds a set of temporal

events from glial reactivation, cell-cycle re-entry, new neuron generation, and circuit

integration(**Abraham et al., 2024; Powell et al., 2016**). In zebrafish, MG respond to

injury by glial reactivation, which refers to the initial response of quiescent MG,

characterized by morphological changes and upregulation of reactive markers (*gfap,*

ascl1a, lin28a) and activation of signaling pathways such as Notch, Jak/Stat, and Wnt

(Lahne et al., 2020b; Pollak et al., 2013; Sifuentes et al., 2016; Yao et al., 2016). This

is followed by the proliferation phase, in which MG initiate a transcriptional shift

toward a progenitor-like state and cell cycle re-entry. Many genes induced by glial

reactivation, such as *lin28a*, *sox2*, and *mycb/h*, are required for cell-cycle re-entry and

subsequent neurogenesis **(Gorsuch et al., 2017; Jorstad et al., 2017; Lee et al., 2024;**

Ramachandran et al., 2010). Diverse signaling pathways have been identified to

participate in this initial regenerative response**(Campbell et al., 2021; Wan and**

Goldman, 2017).

In contrast, although forced expression of factors like *Ascl1* or deletion of

repressive chromatin regulators can partially reprogram MG into interneuron-like

cells in the injured mouse retina**(Hoang et al., 2020; Jorstad et al., 2017)**, the

knowledge of the upstream signals that trigger injury-induced cell-cycle re-entry, a

critical step for producing neurogenic progenitors, remains limited**(Lenkowski and**

Raymond, 2014; Vihtelic and Hyde, 2000; Wu et al., 2001). The mechanistic

understanding of MG cell-cycle re-entry in the injured zebrafish retina will provide

critical knowledge for inspiring new strategies for *in-vivo* re-programming MG to

repair the human retina.

The intrinsic and environmental factors contribute to injury-induced MG

proliferation**(Gao et al., 2021; Lahne et al., 2020b; Xiao et al., 2023)**. Recent efforts

in single-cell RNA-sequencing (scRNA-Seq) have identified *ascl1α*,

*clcf1/crlf1a***(Boyd et al., 2023)**, and *mycb/mych***(Lee et al., 2024)** as key intrinsic

drivers of injury-induced MG proliferation in the zebrafish retina. Also, the elegant

study of MG injury responses across multiple species, including zebrafish, chicken,

and mice, revealed the *hmga1/yap1* signaling network as a key regulator of MG

reactivation and neurogenesis in the regenerating retina**(Hoang et al., 2020).**

Interestingly, previous studies further reported that post-injury MG in the zebrafish is

capable of re-acquiring the regeneration program that largely recapitulates the

embryonic retinal developmental program(**Celotto et al., 2023; Hoang et al., 2020;**

Lahne et al., 2020a; Lyu et al., 2023).

In terms of extrinsic cues, two sources, those from damaged neurons and injury-

recruited microglia, are considered indispensable for injury-induced MG

proliferation(**Leach et al., 2021; White et al., 2017**). Previous studies have shown

that signals derived from various retinal neuron injury models trigger MG

regeneration(**Lyu et al., 2023**). Meanwhile, accumulating evidence increasingly

appreciates the role of inflammatory signals derived from injury-recruited microglia,

such as *TNFα*(**Conner et al., 2014; Nelson et al., 2013**), cytokines IL-1β, and IL-

10(**Lu and Hyde, 2024**). Notably, inflammatory responses are intrinsically associated

with redox signaling, which is involved in the regeneration processes in various non-

neuronal tissues(**Breus and Dickmeis, 2021; Jaeschke, 2000; Zhang et al., 2024**).

However, the involvement of the redox signaling resulting from microglia-derived

inflammatory signals in MG cell-cycle re-entry following retina damage remains to be

clarified. Nitric oxide (NO), a redox signal, has participated in regenerating various

tissues in zebrafish, including the heart(**Rochon et al., 2020; Yu et al., 2024**), spinal

cord(**Bradley et al., 2010**), and fin(**Matrone et al., 2021**). Endogenous NO is derived

from three forms of NO synthase (NOS) in mammals: neuronal NOS (nNOS),

endothelial NOS (eNOS), and inducible NOS (iNOS). The zebrafish comprises *Nos1*,

a form of neuronal *Nos*, and *Nos2a/2b*, two inducible *Nos*.

In this study, we performed single-cell RNA sequencing to characterize the post-

injury MG states in the zebrafish retina following the specific ablation of green/red

cone. This comprehensive analysis led us to define a set of previously unknown

cxcl18b-positive transitional states as an essential route of MG cell-cycle re-entry in

response to injury, and the inflammatory was indispensable for the induction of these

transitional states. Intriguingly, we found that these *cxcl18b*-defined transitional states

exhibited gene patterns shared by cell states in the CMZ, favoring the idea that

cxcl18b-defined MG transitional states might represent the developmental program

conserved by neuronal regeneration beyond embryonic development. A remarkable

phenotype of enriched redox features in these gene patterns suggested the importance

of redox signaling. The subsequent screening of redox-related genes revealed the

essential role of nitric oxide, produced by *Nos2b*, in triggering MG cell-cycle re-entry

after the retinal injury. Notably, we developed a sophisticated viral-based approach to

achieve the *cxcl18b*-defined MG transitional state-specific knockout of *nos2b*

successfully and verified the requirement of transitional state-specific NO signaling in

injury-induced MG proliferation. These findings provide novel cellular and molecular

insights into this species-specific post-injury MG cell-cycle re-entry process, with

potential implications for the development of regenerative medicine strategies.

Results

Single-cell transcriptome analysis reveals the landscape of injury-induced Müller

glia states

Zebrafish Müller glia can respond to retina injury by cell cycle re-entry, a critical

step evolutionarily absent from their mammalian counterparts but essential for neuron

regeneration**(Goldman, 2014; Powell et al., 2016)**. We created a zebrafish retina

injury model by crossing *Tg(opn1lws2: nfsb-mCherry)^{uom3}* (referred to as *Tg(lws2:*

nfsb-mCherry)) with *Tg(mpeg1: GFP)* fish, in which the bacterial nitroreductase

(NTR) enzyme was specifically expressed in G/R cone. We selectively ablated the

G/R cone starting at 5 days post-fertilization (dpf) by a subsequent 120 hours of

metronidazole (MTZ) exposure**(Curado et al., 2007; Curado et al., 2008) (Figure**

1A and Figure 1-figure supplement 1A). G/R cone became significantly reduced in

number since 48 hours post-injury (hpi) and was mostly depleted at 120 hpi **(Figure**

1B). Meanwhile, a number of microglia (marked by *Tg(mpeg1: GFP)*) migrated to the

outer nuclear layer (ONL) as early as 48 hpi, peaked at 72 hpi, and began to reduce in

number at 96 hpi and onward **(Figure 1C)**. To confirm the identity of these

proliferating cells, we performed BLBP immunostaining and observed that the

PCNA□ cells were also BLBP□ (**Figure 1–figure supplement 1B**), indicating their

MG origin. Notably, in response to G/R cone ablation, the proliferative MG

population increased starting at 48 hpi, peaked at 72 hpi, and began to decline since

96 hpi and forward **(Figure 1D)**. Considering the result that the number of

proliferative MG peaked at 72 hpi, we focused on the 72-hpi time point for further

exploration of MG proliferative behaviors following G/R cone ablation **(Figure 1E)**.

By revisiting previously obtained scRNA-seq data of MG enriched from *Tg(lws2:*

nfsb-mCherry) crossed with *Tg(gfap: EGFP)* and *Tg(her4.1: dRFP)* retina before and

after G/R cone ablation at 72 hpi**(Krylov et al., 2023)**, we selected 5,932 and 3,999

MG cells and their derived progenies from the uninjured and 72-hpi retina,

respectively **(Details in Methods; Figure 1-figure supplement 1C)**. By clustering

these cells, we identified 13 clusters, 8 out of which (Clusters 2, 3, 5, 6, 9, 10, 11, and

12) with an increased proportion in response to the G/R cone ablation **(Figure 1-**

figure supplement 1D). Subsequently, we re-clustered cells of these 8 clusters (695

cells of uninjured retinae and 3,477 cells of 72-hpi retinae), aggregating into 10 new

clusters. After the quality control procedure, we did not consider Clusters 7/8/9 due to

their small populations with ribosomal, dendritic, and doublet features, resulting in 7

clusters for further analysis. (Details in Methods; **Figure 1F)**

We performed the pseudo-time trajectory analysis to reveal the progression of

these MG clusters following the cone ablation **(Figure 1G)**. Cluster 4 cells were

highly expressing genes related to mature MG (*glula*, *slc1a2b*, *apoeb,* and

rlbp1a)**(Bernardos and Raymond, 2006; Raymond et al., 2006; Thummel et al.,**

2008; Yurco and Cameron, 2005), the quiescent state (*cx43*)**(Dermietzel et al., 2000;**

Janssen-Bienhold et al., 1998), and major MG population marker (*fgf24*)**(Krylov et**

al., 2023) (Figure 1-figure supplement 1E). Furthermore, Cluster 4 began to express

s100α10b and *gfap,* reactive state makers**(Celotto et al., 2023; Hoang et al., 2020)**, in

the injured retinae, but not in the uninjured retina **(Figure 1-figure supplement 1E)**.

Thus, we set Cluster 4 as the early transitional MG state.

To examine the transition of these 7 MG clusters, the pseudo-time trajectory

showed that the main developmental branch consisted of Clusters 4/1/2 and then

became divergent into two sub-branches, including Cluster 0 and 5/3/6 **(Figure 1G)**.

In contrast to the sub-branch of Cluster 0, the sub-branch of Clusters 5/3/6 was highly

expressing proliferative cell markers (*pcna, mki67,* and *mcm2*). Within the latter sub-

branch, while Cluster 3 and 6 with the highest levels of proliferative cell markers,

Cluster 6 began to express neuronal differentiation factors (*otx5, crx,* and

pde6gb)**(Abalo et al., 2020; Asaoka et al., 2014; Shen and Raymond, 2004)**

(Figure 1-figure supplement 1E). Thus, we identified 6 major post-injury MG states,

from the early transitional state (Cluster 4) to three transitional states (Cluster 1/2/5),

to finally two proliferative states (Cluster 3/6).

Remarkably, chemokine (C-X-C motif) ligand 18b *(cxcl18b*), an inflammatory

chemokine, was uniquely expressed in three transitional states but largely absent from

the early transitional Cluster 4 and two proliferative Cluster 3/6. Specifically, while

the first $cxcl18b^+$ transitional state (Cluster 1) was expressing *cx43*, a maker for MG

quiescence**(Dermietzel et al., 2000; Janssen-Bienhold et al., 1998)**, the last

transitional state (Cluster 5) began to show a weak induction of *ascl1α*

(Ramachandran et al., 2010). Immediately following this last transitional state,

Cluster 3 started with high *ascl1α* expression and entered the proliferative state with

the expression of *pcna* and *mik67* **(Figure 1H)**. Our analysis highlighted a new set of

cxcl18b-defined MG transitional states preceding *ascl1α* induction, bridging MG from

the most original quiescence state to injury-induced proliferation.

Clonal analysis reveals injury-induced MG proliferation via *cxcl18b*-defined

transitional states associated with inflammation

To directly verify the presence of *cxcl18b*-defined MG transitional states, we first

examined the temporal relationship of *cxcl18b* expression and MG proliferation after

the cone ablation using *in situ* hybridization combined with immunostaining of either

BLBP (an MG maker) or PCNA (a proliferative cell maker) **(Figure 2A and Figure**

2-figure supplement 1A). The result showed that as early as 24 hpi, the number of

cxcl18b$^+$ MG was rapidly peaked with no emergence of proliferative MG **(11 ± 4,**

n=10 in *cxcl18b*$^+$ MG; mean ± SEM), and then *cxcl18b*$^+$ MG continued declining in

number over time and reached the lowest level since 96 hpi **(1 ± 1, n=7; mean ±**

SEM; Figure 2B). In contrast, the number of proliferative MG (PCNA$^+$) peaked at 72

hpi and decreased to the lowest level at 120 hpi **(9 ± 2, n=11 in 72-hpi retina; 1 ± 1,**

n=5 in 120-hpi retina; mean ± SEM; Figure 2B). Note that *cxcl18b*$^+$ MG was

mostly proliferative at 72 hpi **(Figure 2-figure supplement 1B)**.

To further verify the temporal expression of *cxcl18b* in MG following the cone

ablation, we created a new transgenic reporter *Tg(cxcl18b: GFP)* by cloning a 3k-bp-

long cis-element of 5'UTR with GFP, allowing real-time monitoring of injury-induced

cxcl18b expression *in vivo* **(Figure 2C and Figure 2-figure supplement 1C-D)**.

Combining this line with PCNA immunostaining, we confirmed the remarkable

increase of *cxcl18b* expression (GFP$^+$) at 48 hpi **(0 ± 1, n=4 in uninjured retinas vs**

11 ± 4, n=7 in 48-hpi; mean ± SEM; Figure 2D and E). Due to the prolonged stay

of GFP protein, we could also observe that some GFP$^+$ MG were also PCNA$^+$ **(Figure**

2D). The result of proliferative MG as a subpopulation of *cxcl18b*$^+$ MG led to an

outstanding question as to whether *cxcl18b*-defined MG transitional states represented

an essential route to injury-induced proliferation.

To address it, we created a new transgene *Tg(cxcl18b: Cre-vmhc: mCherry::*

ef1α: loxP-DsRed-loxP-EGFP:: lws2: nfsb-mCherry) to perform the clonal analysis

of historical *cxcl18b*-expressing in MG after the G/R cone ablation **(Details in**

Method; Figure 2-figure supplement 1E). After ablating the G/R cone by MTZ

treatment at 6 dpf for 3 consecutive days, we found that the number of *cxcl18b*

lineage-traced MG (marked by GFP$^+$) and PCNA$^+$ MG was significantly increased at

72 hpi **(Figure 2F)**. Further analysis showed that all PCNA$^+$ MG were *cxcl18b*

lineage-traced MG, indicating the *cxcl18b* lineage-traced MG were the ones who

could eventually enter into the cell cycle **(13 ± 2 cells for PCNA$^+$ vs 13 ± 2 cells**

PCNA$^+$ and GFP$^+$MG; percentage of PCNA$^+$ and *cxcl18b$^+$* vs PCNA$^+$ = 93 ± 2%;

n=14; p=0.19; Mean ± SEM; Figure 2G). Further analysis showed that at 120 hpi,

the time point that the MG proliferation has largely creased, GFP$^+$ MG with the

lineage history of injury-induced *cxcl18b* expression constituted about 44% of GS$^+$

MG at the central retina, indicating that only about half of MG could enter *cxcl18b$^+$*

transitional states following the cone ablation **(14 ± 3 in GS$^+$ and GFP$^+$ MG; 28 ± 4**

in GS$^+$ and GFP$^-$ MG, n=14; p<0.05; mean ± SEM; Figure 2I). Together, our

clonal analysis demonstrated that proliferative MG mostly originated from *cxcl18b*-

defined MG transitional states, and 44% central MG could become *cxcl18b* positive.

To investigate whether *cxcl18b* was required for MG proliferation following G/R cone

ablation, we employed CRISPR-Cas9-mediated gene disruption, using two sgRNAs

targeting *cxcl18b* **(Figure 2-figure supplement 1F and G)**. We found that *cxcl18b*

knockout did not reduce MG proliferation after G/R cone ablation at 72 hpi (**13 ± 3,**

n=11 in WT; 11 ± 3, n=7 in *scramble* sgRNA-injected; 13 ± 3, n=7 in *cxcl18b*

sgRNA-injected; mean ± SEM), suggesting that *cxcl18b* per se does not regulate

MG proliferation directly (**Figure 2-figure supplement 1H and I**). This led us to

wonder about the induction of *cxcl18b*-defined MG transitional states.

As an inflammatory chemokine, *cxcl18b* serves as a reliable marker of

inflammation and regulates neutrophil recruitment to injury sites(**Goumenaki et al.,**

2024; Torraca et al., 2017). Inflammation has been previously shown to be critical

for inducing regenerative responses in adult zebrafish, where it promotes reactive

microglia/macrophages and Müller glial proliferation in the retina(**Iribarne and**

Hyde, 2022; Kyritsis et al., 2012). Notably, suppressing the immune response using

dexamethasone (Dex) in zebrafish retina reduced microglial reactivation and

significantly decreased the number of proliferative MG(**Silva et al., 2020; Zhang et**

al., 2020). In our study, we identified the *cxcl18b*-defined transitional states as the

essential routing for MG proliferation after G/R cone ablation. These results prompted

us to investigate whether the inflammatory responses mediated by recruited microglia

are indispensable for the formation of these *cxcl18b*-defined transitional states.

To address this, we examined *cxcl18b* expression using *Tg(cxcl18b: GFP)* after

inhibiting inflammation using Dex(**Iribarne and Hyde, 2022**), and observed a

significant reduction in the number of $cxcl18b^+$ MG (GFP^+ cells) at 72 hpi (**11 ± 4,**

n=10 in Dex treated retina vs 19 ± 5, n=6 in DMSO treatment; and 18 ± 3, n=14

in the 72-hpi retina; mean ± SEM) (Figure 2K). Consistent with earlier reports, we

observed that Dex treatment inhibited the migration of microglia **(indicated by**

*Tg(mpeg1: GFP)***; 16 ± 4, n=14 in Dex treated retina vs 27 ± 5, n=7 in DMSO**

treatment; and 28 ± 4, n=9 in the 72-hpi retina; mean ± SEM) to the ONL and

significantly reduced the number of proliferative MG **(PCNA$^+$; 8 ± 2, n=14 in Dex**

treated retina vs 17 ± 2, n=7 in DMSO treatment; and 14 ± 3, n=9 in the 72-hpi

retina; mean ± SEM) at 72 hpi after G/R cone ablation **(Figure 2L and M and**

Figure 2-figure supplement 1J). These findings suggest that microglia-mediated

inflammation may contribute to the activation of *cxcl18b*-defined transitional states

that precede MG proliferation, although a causal relationship remains to be

established. While dexamethasone suppressed both microglial recruitment and

cxcl18b$^+$ MG generation, its broad anti-inflammatory action precludes definitive

conclusions about microglial causality. Dissecting this relationship would require

concurrent ablation of microglia and cone photoreceptors using a triple-transgenic

strategy, which is beyond the scope of the current study. Targeted approaches will be

necessary to resolve the specific role of microglia in initiating *cxcl18b* expression.

cxcl18b-**defined MG transitional states recapitulate molecular features of retinal**

stem cells in the ciliary marginal zone

Interestingly, we observed the *cxcl18b* expression in the CMZ after the cone

ablation besides its high expression in the MG **(Figure 2A)**. We were then curious

about the *cxcl18b* expression in the developmental retina as well as in the CMZ

without the injury. Notably, *in situ* results showed that *cxcl18b* was largely absent

from the central region but presented in the peripheral region of 30-hpf retinae,

whereas it was highly expressed in the most peripheral region of the CMZ, where

fabp11a and *col15α1b*, two putative markers for post-embryonic retinal stem cells

(RSCs) are located**(Gonzalez-Nunez et al., 2010; Raymond et al., 2006) (Figure**

3A-B and E-F, and Figure 3-figure supplement 1A). The transgenic line of

Tg(cxcl18b: GFP) also showed a robust *cxcl18b* expression in the CMZ **(Figure 3-**

figure supplement 1D). Consistently, our scRNAseq data of CMZ cells also

confirmed the co-expression of *cxcl18b*, *fabp11a*, and *col15α1b* **(Figure 3C-D and**

G-H, and Figure 3-figure supplement 1B). Furthermore, cluster 1 MG, at the earlier

stage of transitional states, has the highest *cxcl18b* expression with *col15α1b*

expression **(Figure 1F and Figure 3-figure supplement 1C)**. All these results

suggested that *cxcl18b*-defined transitional states, at least to some extent, represent

the developmental state of retinal stem cells in the CMZ, but not that of embryonic

retinal progenitors.

***Nos2b* is required for MG entry into the proliferation via NO signaling**

Previous studies have demonstrated the involvement of redox signaling in cell

regeneration processes in various tissues across species**(Han et al., 2014; Hunter et**

al., 2018; Matrone et al., 2021; Yoo et al., 2012). All these evidences led us to

directly test the roles of redox genes in serving as the molecular mechanism

underlying injury-induced MG proliferation. Thus, we first examined the expression

levels of a comprehensive list of redox genes in *cxcl18b*-defined MG transitional

states (Clusters 1/2/5) in our scRNA-seq data **(Figure 1H)** and screened the influence

of 18 genes from each major category of redox signaling on injury-induced MG

proliferation using CRISPR-Cas9-mediated gene disruption **(Figure 4-figure**

supplement 1A). We focused on the nitric oxide (NO) signaling pathway, targeting

three genes encoding nitric oxide synthases (*Nos*): neuronal *Nos* (*nos1*) and two

inducible forms (*nos2a* and *nos2b*), as well as the gene encoding S-nitrosoglutathione

reductase (*gsnor* or *adh5*), which modulates reactive NO signaling **(Figure 4A and**

Figure 4-figure supplement 1B). The consequence and efficiency of gene disruption

were verified by DNA sequencing **(Figure 4-figure supplement 1C)**. Notably, the

disruption of *nos2b* resulted in a significant reduction of PCNA$^+$ MG at 72 hpi **(6±2,**

n=22 in *nos2b*-disrupted vs 11±3, n=7 in *scramble* sgRNA-injected; Mean ± SEM)

(Figure 4B). Noted that *nos* gene disruption did not significantly alter microglia

recruitment or G/R cone ablation at 72 hpi, suggesting that the influence of NO on

injury-induced MG proliferation was not via inflammatory reactions of recruited

microglia or injury degree **(Figure 4-figure supplement 1D-F)**.

To further examine the function of *Nos2b* via NO, we employed various *Nos*

inhibitors (L-NG-nitro arginine methyl ester, L_NAME; L-NG-Monomethyl Arginine,

L_NMMA; 1400W) and NO scavengers (carboxy-PTIO, C-PTIO)**(Goldstein et al.,**

2003; Hong et al., 2012; Moore et al., 1990; Rees et al., 1990). We performed the

intraocular injection of the drugs (PBS as control) into the zebrafish eye from 2 days

before cell ablation until 72 hpi **(Figure 4C)**. Notably, the NO scavenger C-PTIO

mostly suppressed MG proliferation, indicating the involvement of NO **(cell number**

313 **of PCNA$^+$ MG: 3±2, n=27 after blocking NO by C-PTIO vs 14±4, n=16 in PBS-**

314 **injected retina; Mean ± SEM) (Figure 4D and E)**. Moreover, 1400W (an inhibitor

315 specific to inducible *Nos*, **7±2, n=10; Mean ± SEM**) and L_NAME (a broad inhibitor

316 to all three *Nos* forms, **8±2, n=12; Mean ± SEM**) could also significantly reduce the

317 number of proliferative MG after the ablation, whereas L_NMMA (the inhibitor

318 specific to neuronal *Nos*) did not influence MG proliferation (**11±3, n=12; Mean ±**

319 **SEM**) **(Figure 4E)**. Taken together, these results highlight the critical role of NO

320 signaling in regulating injury-induced MG proliferation.

321 We further generated nitric oxide pathway mutant zebrafish (*nos1*, *nos2a*, *nos2b*,

322 and *gsnor*) to investigate the role of NO in MG proliferation following G/R cone

323 ablation. Utilizing CRISPR/Cas9-mediated gene disruption, we successfully screened

324 out *nos1*, *nos2a*, *nos2b*, and *gsnor* mutants, characterized by deletions of 133-bp, 13-

325 bp, 220-bp, and 11-bp coding sequences, respectively (see Methods for details,

326 **Figure 4-figure supplement 2A and B**). Consistent with the gene disruption

327 experiments described above, we observed a significant reduction in the number of

328 proliferative (PCNA□) MG at 72 hpi following G/R cone ablation in both

329 heterozygous and homozygous *nos2b* mutants (*nos2b*□/□: 3□±□1, *n*□=□20;

330 *nos2b*□/□: 6□±□2, *n*□=□20) compared to WT controls (11□±□2, *n*□=□14;

331 mean□±□SEM). In contrast, no significant changes in MG proliferation were

332 observed in *nos1*, *nos2a*, or *gsnor* mutants compared to wild type **(Figures 4F-4I)**.

333 Interestingly, the reduction in proliferative MG was more pronounced in *nos2b*

334 heterozygous mutants (*nos2b$^{+/-}$*) than in homozygous mutants (*nos2b$^{-/-}$*) **(*nos2b$^{+/-}$* vs**

nos2b^-/-^; p<0.001; Mean ± SEM) (Figure 4J). We observed no significant difference

in the loss of cone photoreceptor at 72 hpi between *nos2b* mutants and WT, indicating

that the reduced MG proliferation observed in *nos2b* mutants is independent of the

injury (WT: 45□±□8 remaining cones, *n*□=□24; *nos2b*□/□: 49□±□12, *n*□=□20;

nos2b□/□: 46□±□9, *n*□=□20; mean□±□SEM) (Figure 4K). This unexpected result

suggests a concentration-dependent effect of NO on proliferative MG. Specifically,

compared to homozygous mutants, heterozygous mutants with intermediate NO levels

more effectively suppressed MG proliferation, whereas wild-type animals with higher

NO levels promoted MG proliferation. This concentration-response pattern highlights

the role of NO as a regulator, rather than a mediator, of injury-induced MG

proliferation.

Specific *nos2b* expression in *cxcl18b*-defined transitional states MG after G/R

cone ablation

To further examine whether the expression of *nos2b* in *cxcl18b*^+^ MG, we used

the newly generated transgene fish line in our study *Tg(cxcl18b: GFP)*, enabling us to

monitor *cxcl18b*^+^ MG after the G/R cone ablation in a real-time manner. By crossing

different fish lines, we fluorescently sorted out three post-injury MG populations (72-

hpi MG, 72-hpi PCNA^+^ MG, and 72-hpi *cxcl18b*^+^ MG) and three control groups

(uninjured MG, 72-hpi retinal cells other than MG and 72-hpi G/R cone) **(Figure 5A-**

C). The real-time quantitative PCR (RT-qPCR) analysis showed that compared to the

three control groups, the expression of *nos2b* was significantly higher in *cxcl18b*^+^ MG

than in 72-hpi MG and 72-hpi PCNA$^+$ MG (**relative expression of *nos2b*: 89±32,**

repeats n=7 in *cxcl18b*$^+$ MG; 14±17, repeats n=5 in 72-hpi MG; and 1±1, repeats

n=6 in PCNA$^+$ MG; Mean ± SEM) (Figure 5D and Figure 5-figure supplement

1D-F). Together with the fact that *cxcl18b*$^+$ MG contained PCNA$^+$ and PCNA$^-$

populations in *Tg(cxcl18b: GFP)* (**Figure 2D**), our result indicated that *nos2b* was

specifically expressed in *cxcl18b*$^+$ non-proliferative MG, which agreed with the

scRNA-seq result of the specific expression of *cxcl18b* in three transitional states

(Clusters 1, 2, and 5; **Figure 1H**). *In situ* hybridization using an HCR molecular probe

in the *Tg(lws2: nfsb-mCherry:: cxcl18b: GFP)* fish line also suggests that injury-

induced *nos2b* expression was specific in the *cxcl18b*-defined transitional state MG

(**Figure 5-figure supplement 1A-C$_3$**). Thus, our analysis reveals that *nos2b* was

specifically expressed in *cxcl18b*$^+$ transitional MG states, particularly in non-

proliferative cells.

NO produced by *nos2b* determines *cxcl18b*$^+$ MG entry for proliferation

To further explore the role of *cxcl18b*$^+$ MG-specific *nos2b* in regulating the entry

of MG proliferation after the G/R cone ablation, we developed a sophisticated clonal

analysis of *cxcl18b*$^+$ MG with genetic disruption of *nos2b* using CRISPR-Cas9

method. To achieve glial type-specific gene manipulation, we employed an

adenovirus strain that specifically infects radial glia in zebrafish(**Jia et al., 2019; Liu**

et al., 2021), and confirmed that it could faithfully mark MG in the zebrafish retina

(**Figure 6A and B**). For the clonal analysis, we performed the intraocular injection of

two viruses packaged with elements of *cxcl18b: gal4* and *UAS: Cas9-T2A-Cre-u6:*

sgRNA (nos2b) into the eye of *Tg(lws2: nfsb-mCherry:: ef1α: loxP-DsRed-loxP-*

EGFP) at 5 dpf, virus packaged with the element of *UAS: Cas9-T2A-Cre-u6: empty*

as the control **(Figure 6C-D and Figure 6-figure supplement 1A)**. One day after the

virus infection, the fish was treated with MTZ for 3 consecutive days to ablate the

G/R cone **(Figure 6E)**.

At 72 hpi, we collected the clones (GFP[+]) derived from *cxcl18b* lineage-traced

MG and analyzed their proliferative property **(Figure 6F and G)**. The analysis of

virus-infected clones (GFP[+]) cell cycle re-entry (PCNA[+]) after G/R cone ablation

revealed that only ~23% of GFP[+] MG clones remained PCNA[+] in the *nos2b* knockout

group, compared to ~75% in the control group **(Figure 6H)**. Note that the efficiency

of *nos2b* sgRNAs was confirmed in terms of mutation types and knockout efficiency

by sequencing **(Figure 6-figure supplement 1B-D)**.

Taken together, these findings suggested that it is the NO produced by *Nos2b*

within *cxcl18b*-defined transitional state MG that specifically drives MG from the

transitional state into proliferation following injury, highlighting a pivotal mechanism

underlying injury-induced regenerative processes.

NO decreased Notch activity that is responsible for injury-induced MG

proliferation

Previous studies have shown that *ascl1α* and Notch signaling are essential for

MG proliferation in the injured zebrafish retina**(Conner et al., 2014; Wan et al.,**

2012). Regarding Notch signaling, a high Notch3 expression is reported to maintain

MG quiescence. In response to the injury, *notch3* expression is downregulated, but

notch1a is necessary for the continued proliferation of the progenitors**(Campbell et**

al., 2022). In consistent, we observed that *notch3* and *hey* (the Notch downstream

target) were highly expressed in uninjured MG clusters and became reduced from the

early transitional state (Cluster4) to the proliferative states (Cluster 3 and 6), whereas

notch1a/1b and *ascl1α* were prominently expressed in the late stage of $cxcl18b^+$

transitional states (clusters 5) and the proliferative states (Cluster 3; **Figure 1H and**

7A). Interestingly, upstream regulators of Notch signaling activation, such as *fgf8a*,

*fgf8b***(Wan and Goldman, 2017)**, and *tgfb3***(Lee et al., 2020)**, were predominantly

expressed in clusters 4 and 1, preceding the expression of *cxcl18b* **(Figure 7A)**. These

results led us to wonder whether NO regulated *cxcl18b*-defined transitional state MG

cell cycle re-entry via the Notch signaling pathway.

To examine the influence of NO signaling blockage on Notch activity dynamics

following the cone ablation, we employed a reporter line *Tg(Tp1bglob: EGFP)*

(referred to as *Tg(Tp1:EGFP))*, in which EGFP is driven by the TP1 element, the

direct target of the intracellular domain of Notch receptors (NICD) that is generated

upon Notch activation**(Parsons et al., 2009; Quillien et al., 2014)**. We treated fish

with the NO scavenger C-PTIO and MTZ, starting at 5 dpf for 5 continuous days and

at 7 dpf for 3 continuous days, respectively, followed by immunostaining for EGFP

and PCNA **(Figure 7B)**. Interestingly, *Tp1*: EGFP$^+$ MG were significantly reduced at

all three injury time points **(cell number of EGFP$^+$ clones: 34±3, n=9 in 24-hpi**

retina; 29±3, n=4 in 48-hpi retina; 33±4, n=6 in 72-hpi retina vs 42±1, n=4 in the

uninjured retina; Mean ± SEM), demonstrating a decrease in Notch activity

following G/R cone ablation **(Figure 7C)**. Notably, this reduction in Notch activation

was further rescued by NO blocking using C-PTIO (**cell number of EGFP$^+$ clones:**

42±1, n=6 in C-PTIO-treated retina at 24 hpi; 42±2, n=5 at 48 hpi; 44±2, n=7 at

72hpi; Mean ± SEM), suggesting that NO modulates Notch signaling **(Figure 7C)**.

Meanwhile, C-PTIO treatment significantly reduced the number of proliferative MG

(marked by PCNA) **(Figure 7D)**. These findings indicated that injury-induced NO

suppresses Notch signaling activation, which potentially drives MG to exit quiescence

and enter proliferation. Together, our results highlight that NO signaling drove the

injury-induced *cxcl18b*-defined transitional state MG to enter proliferation, which

may be mediated by Notch pathway regulation.

Discussion:

Our study provides the single-cell transcriptomic landscape of Müller glial state progression following the cone ablation at the larval stage. Combined with clonal analysis, we identified a previously unreported *cxcl18b*-defined transitional MG state as the essential routing for MG cell cycle re-entry. It led to our further genetic analysis, revealing the cell type-specific and concentration-dependent regulatory mechanism of nitric oxide derived from *nos2b* underlying injury-induced MG proliferation. Furthermore, NO signaling accounted for a decreased Notch activity, which has been previously reported to be essential for MG proliferation after the injury. Finally, cell state-specific gene disruption revealed that *cxcl18b*-defined MG transitional state-specific *nos2b* is required for injury-induced MG proliferation, which is a prime example of previously reported "5R" redox regulation(Meng et al., 2021). Thus, our study provides the novel precision redox mechanism underlying MG proliferation in response to cone ablation in the zebrafish retina, opening exciting possibilities for future research and potential therapeutic interventions.

Identification of *cxcl18b*-defined MG transitional states following the cone ablation

Combined scRNA-seq analysis and clonal analysis, we identified a previously unreported transitional MG state, marked by the expression of *cxcl18b,* as the essential routing for MG to re-enter the cell cycle following the retina injury **(Figure 1H, 2F, and 2G)**. To our knowledge, it is the first transitional state verified by *in vivo* clonal analysis to show a faithful prediction of injury-induced MG proliferation. Notably, this *cxcl18b* induction in MG depends on microglial recruitment in response to cone ablation, suggesting this transitional state is an MG response to the signals derived from the inflammatory reaction **(Figure 2K)**. The underlying mechanism of this crosstalk is crucial to be addressed. Interestingly, the *cxcl18b*-containing gene module was also expressed in the CMZ, a region crucial for adult retina neurogenesis for a lifetime **(Figure 3A-D)**. However, its expression is mainly absent from the central regions of the developing retina **(Figure 3F)**. It suggests that the *cxcl18b-*

defined transitional state might represent a developmental state used by constitutive

neuron generation programs and injury-induced neuron regeneration programs beyond

embryonic development. Furthermore, the *cxcl18b*-defined transitional state exhibits

robust redox-related characteristics, such as the expression of *sod1*, *sod2*, and *catalase*

(Figure 4-figure supplement 1A). It led to the critical discovery of this study, which

showed the essential role of *nos2b* in regulating injury-induced MG proliferation.

Unfortunately, our preliminary effects on gene disruption using CRISPR/Cas9 in F0

founders did not observe a significant reduction in the number of proliferative MG

after *cxcl18b* disruption **(Figure 2-figure supplement 1I)**. However, a recent study

reported the essential role of *cxcl18b* in heart regeneration using mutant fish,

providing a mechanism of *cxcl18b* as innate immune signaling in injury-induced

tissue regeneration**(Goumenaki et al., 2024)**. Their results raised concern about the

efficiency of *cxcl18b* disruption in our system. It is essential to use mutant fish to re-

examine the role of *cxcl18b* in injury-induced MG proliferation in the future. Also, we

cannot rule out the possibility that other co-factors are involved in the action of

cxcl18b in MG regeneration, which is another critical issue that needs to be solved in

the future.

The possible mechanism of NO signaling underlying injury-induced MG

proliferation

Our study, for the first time, demonstrated an essential role of NO signaling in

regulating MG proliferation after the cone ablation. However, we still need to

understand more about the underlying mechanism. There are two well-characterized

molecular events responsible for MG proliferation following the retina injury: a

decreased Notch activity and an increased *ascl1α* expression**(Fausett et al., 2008;**

Jorstad et al., 2017; Ramachandran et al., 2010). Previous studies have shown that

Notch3 is responsible for this decreased Notch activity, leading to increased Ascl1a

through the de-depression mechanism**(Campbell et al., 2021; Sahu et al., 2021).**

Interestingly, unlike *notch3*, *notch1a* has been reported to stimulate MG

proliferation**(Campbell et al., 2021; Campbell et al., 2022)**. Our scRNA-seq analysis

also showed that as MG progressed into the proliferative states after the cone ablation, *notch3* expression gradually declined **(Figure 7A)**. In contrast, *ascl1α, notch1a*, *and notch1b* expressions were up-regulated and peaked at the proliferative states **(Figure 7A)**. Thus, both previous studies and our current analysis support the idea that the transcriptional regulation of Notch expression accounts for the decreased Notch activity after the injury. Intriguingly, NO has been reported to activate the Notch1 signaling cascade by promoting the release and accumulation of the Notch1 intracellular domain (NICD) through nitration reactions, subsequently enhancing tumorigenesis and stem-like features in various cellular systems**(Charles et al., 2010; López-Juárez et al., 2017; Villegas et al., 2018)**. It raises the possibility that NO signaling regulates injury-induced MG proliferation through the post-translational modification of Notch3. Previous studies have demonstrated two significant forms of NO-mediated post-modification: S-nitrosylation on cysteines and nitration on tyrosine(Wu et al., 2014). Our preliminary analysis showed that all 11 cysteine residues within the putative γ-secretase-dependent cleavage sites are conserved between Notch1a and Notch3, while notable differences were observed at four tyrosine residues. It leads to an outstanding question of whether NO regulated injury-induced MG proliferation by decreasing Notch activity via tyrosine nitration of Notch3, which is worthwhile to elucidate.

The production of NO signaling in MG following retina damage

Previous studies reported that *cxcl18b* is a reliable inflammatory marker**(Goumenaki et al., 2024; Torraca et al., 2017)** and different inflammation responses modulate MG proliferation in the damaged zebrafish retina**(Iribarne and Hyde, 2022)**. The induction of *cxcl18b* may represent the inflammatory responses of MG after the cone ablation, pointing out the potential link between the inflammatory response and the emergence of NO signaling in MG. Previous studies have demonstrated that iNOS is induced in various tissues by proinflammatory cytokines**(Förstermann and Sessa, 2012; Pacher et al., 2007)**. One of the

approaches to test the role of inflammatory responses is to manipulate the levels of inflammatory responses in MG to see the NO production and MG proliferative behaviors. Also, previous studies appreciate the essential role of electrical activity in tissue regeneration(**Levin, 2009; Qin et al., 2023**); in particular, calcium signaling has been shown to regulate various molecular pathways for liver regeneration, including the hepatocyte growth factor-Met-tyrosine kinase (HGF-Met) transduction pathway(Bedi et al., 2024) and the epidermal growth factor receptor (EGFR) signaling(Kimura et al., 2023). It is interesting to speculate that the abnormal electrical activity of MG in the injured retina may result in an elevated level of intracellular calcium, which activates calmodulin and induces the conformation change of NOS to NO production(**Hanson et al., 2018; Jones et al., 2007**). A similar mechanism has been proposed in the long-term potentiation of excitatory post-synaptic structure(**Grover and Teyler, 1990; Kawamoto et al., 2012; Park, 2018**) as well as in the glutamate neurotoxicity model(**Ashpole et al., 2013; Ashpole et al., 2012**). Thus, the production of NO derived from *Nos* may be the product of the interplay between the inflammatory responses and the electrical activity in MG after the retina damage.

Limitations of this study

A few limitations of this study should be acknowledged: 1. Our study focuses on larval zebrafish, in which the core features of MG and immune responses are conserved compared to the adult. However, we acknowledge that the adult retina—

with its fully matured differentiated retina and immune response — provides irreplaceable biological insight. Nevertheless, larval models offer a powerful platform to uncover conserved regenerative mechanisms and serve as a valuable complement for identifying age-dependent differences in MG-mediated regeneration. 2. While our data suggest that injury-induced NO suppresses Notch signaling activation and promotes MG proliferation, the use of a general NO scavenger (C-PTIO) does not allow us to determine whether this regulation occurs in an autocrine or paracrine manner. The specific role of NO signaling within *cxcl18b*☐ MG requires further

validation using MG-specific NO depletion. 3. The current study's description of the
landscape of MG transitional states is based on single-cell transcriptomic data
obtained at a single time point (72 hpi) after the cone ablation, which may not provide
a complete picture of the state transition of post-injury MG. Future scRNA-seq
analysis of MG at multiple post-injury time points is necessary to clarify this issue. 4.
While we demonstrated a critical role of NO in injury-induced MG proliferation, the
potential contribution of microglia-derived NO was not directly examined. 5. We did
not perform direct measurements of NO levels specifically within *cxcl18b*-defined
MG cells, leaving open the question of localized NO signaling.
564

Acknowledgments

This research was supported by grants from the National Key R&D Program of China (2022YFA1303000 to C.C.), the Chinese Academy of Sciences Strategic Priority Research Program B grants (XDB39000000 to C.C.), National Natural Science Foundation of China-Key program project (Grant No. 91849203 to C.C.), the Creative Research Groups of the National Natural Science Foundation of China (32321003 to J.H.), STI2030-Major Projects (2021ZD0204500 to J.H.), the National Key Research and Development Program of China (2020YFA0112700 to J.H.), National Natural Science Foundation of China (32471029 to J.H.).

Author Contributions

A.Y., design and experimental investigation, visualization, data analysis, methodology, writing-original draft, writing-review and editing. S.Y., M.D., and D.Z., investigation, methodology. J.H. and C.C., conceptualization, experimental design, data analysis, supervision, funding acquisition, writing-original draft, project administration, writing-review and editing.

Declaration of Interests

The authors declare no competing interests.

STAR METHODS

Detailed methods are provided in this version of this paper and include the following:

KEY RESOURCES TABLE

RESOURCE AVAILABILITY

Lead contact

Materials availability

Data and code availability

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Single-cell RNA sequencing data analysis

Pseudo-time trajectory analysis

Zebrafish husbandry and transgenic fish lines

Generation of *Tg(cxcl18b: GFP)*

Generation of *Tg(cxcl18b: Cre-vmhc: mCherry:: ef1α: loxP-DsRed-loxP-EGFP::*

lws2: nfsb-mCherry)

Generation of mutant fish lines

Plasmid construction

Clone analysis of *cxcl18b* lineage-traced MG

Metronidazole treatment

Gene disruption via CRISPR/Cas9 system

Pharmacological Treatment with NO scavenger and dexamethasone or *Nos*

Inhibitors intraocular injection

The real-time quantitative PCR after FACS-sorted MG

Construction and injection of adenoviral-based MG-specific knockout system in

the zebrafish retina

Tissue preparation and immunostaining

In situ hybridization

Imaging

Quantifications and statistical analysis

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (Danio rerio)	Wild type	Dr William		AB
strain, strain background (Danio rerio)	*lws2: nfsb-mCherry*	(Krylov et al., 2023)		*Tg(opn1lws2: nfsb-mCherry)^{uom3}*
strain, strain background (Danio rerio)	*her4.1: dRFP*	(Yeo et al., 2007)	ZDB-TGCONSTRCT-070612–2	*Tg(her4.1: dRFP)*
strain, strain background (Danio rerio)	*gfap: EGFP*	(Bernardos and Raymond, 2006)	ZDB-FISH-150901–29307	*Tg(gfap: EGFP)*
strain, strain background (Danio rerio)	*mpeg1: GFP*	(Ellett et al., 2011)	ZDB-TGCONSTRCT-170801–5	*Tg(mpeg1: GFP)*
strain, strain background (Danio rerio)	*Tp1bglob: EGFP*	(Yu and He, 2019)		*Tg(Tp1bglob: EGFP)*
strain, strain background (Danio rerio)	*pcna: GFP*	(Xu et al., 2020)	ZDB-LAB-070129-2	*Tg(pcna: GFP)*
strain, strain background (Danio rerio)	*ef1α: loxP-DsRed-loxP-EGFP*	(Hans et al., 2009)		*Tg(ef1α: loxP-DsRed-loxP-EGFP)^{zf284}*
strain, strain background (Danio rerio)	*cxcl18b: Cre-vmhc: ECFP*	This manuscript		*Tg(cxcl18b: Cre-vmhc: ECFP; ef1α: loxP-DsRed-loxP-EGFP; lws2: nfsb-mCherry)*
strain, strain background (Danio rerio)	*cxcl18b: GFP*	This manuscript		*Tg(cxcl18b: GFP)*

antibody	Mouse monoclonal anti-PCNA	Abcam	Cat#Ab29; RRID: AB_303394	IF(1:500)
antibody	Rabbit polyclonal anti-BLBP	Abcam	Cat#ab32423; RRID: AB_880078	IF(1:1000)
antibody	Mouse monoclonal anti-Glutamine Synthetase	BD Biosciences	Cat# 610518; RRID: AB_397880	IF(1:1000)
antibody	Chicken monoclonal anti-GFP	Abcam	Cat# ab13970; RRID: AB_300798	IF(1:2000)
antibody	Rabbit polyclonal anti-GFPtag	Rabbit polyclonal anti-GFPtag	Cat#50430–2-AP; RRID: AB_11042881	IF(1:500)
antibody	Rabbit polyclonal anti-DsRed2	Takara Bio	Cat#632496; RRID: AB_10013483	IF(1:1000)
antibody	Alexa Fluor 488 AffiniPure Goat Anti-Mouse IgG (H+L)	Yeasen Biotech	Cat# 33206ES; RRID: AB_3662603	IF(1:1000)
antibody	Alexa Fluor 488 AffiniPure Donkey Anti-Rabbit IgG (H+L)	Yeasen Biotech	Cat# 34206ES60; RRID: AB_2909605	IF(1:1000)
antibody	Alexa Fluor 488 AffiniPure Donkey Anti-Chicken IgY (IgG) (H+L)	Jackson ImmunoResearch Labs	Cat# 703-545-155; RRID: AB_2340375	IF(1:1000)
antibody	Alexa Fluor 594 AffiniPure Donkey Anti-Mouse IgG (H+L)	Yeasen Biotech	Cat# 34112ES; RRID: AB_3661960	IF(1:1000)

antibody	Alexa Fluor 594 AffiniPure™ Goat Anti-Rabbit IgG (H+L)	Yeasen Biotech	Cat# 33112ES; RRID: AB_3661961	IF(1:1000)
antibody	Alexa Fluor 647-AffiniPure Goat Anti-Mouse IgG + IgM (H+L)	Jackson ImmunoResearch Labs	Cat# 115-605-044; RRID: AB_2338906	IF(1:1000)
recombinant DNA reagent	pTol2-cxcl18b: GFP	This manuscript		We made this plasmid by ligated the cxcl18b promoter and GFP element
recombinant DNA reagent	pTol2-cxcl18b: Cre-vmhc: mcherry	This manuscript		We made this plasmid by ligated the cxcl18b promoter and Cre element
recombinant DNA reagent	pTol2-cxcl18b: gal4FF	This manuscript		We made this plasmid by ligated the cxcl18b promoter and gal4FF element
recombinant DNA reagent	pUAS: Cas9T2ACre; U6: sgRNA1; U6: sgRNA2	(Di Donato et al., 2016)	Addgene plasmid #74010; RRID: Addgene_74010	
recombinant DNA reagent	pTol2-UAS: Cas9-T2A-Cre-U6: nos2b sgRNA1; U6: nos2b sgRNA2	This manuscript		We made this plasmid by insert two sgRNAs of *nos2b* in 10xUAS backbone
commercial assay or kit	MEGAscriptTM T7 High Yield Transcription Kit	Invitrogen	Cat# AM1334	
commercial assay or kit	ClonExpress MultiS One Step Cloning Kit	Vazyme	Cat# C113-02	

commercial assay or kit	DIG RNA labeling kit	Roche	Cat# 11277073910	
chemical compound, drug	N(ω)-nitro-L-arginine methyl ester	Sigma-Aldrich	Cat# N5751-1G	10 mM
chemical compound, drug	N(ω)-Methyl-L-arginine acetate salt	Sigma-Aldrich	Cat# M7033-5MG	10 mM
chemical compound, drug	1400W dihydrochloride	MedChemExpress	Cat# HY-18730	200 nM
chemical compound, drug	Phenyl-4,4,5,5-tetramethyl imidazoline-1-oxyl 3-oxide	Sigma-Aldrich	Cat# P5084-25MG	10 mM
chemical compound, drug	Dexamethasone	Sigma-Aldrich	Cat# D1756	10 mM
chemical compound, drug	Metronidazole	Sigma-Aldrich	Cat# M3761-100G	10 mM
software, algorithm	FV10-ASW 4.0 Viewer	Olympus	www.olympus-global.com	Analysis image
software, algorithm	GraphPad Prism V 9.0.0	GraphPad Software	www.graphpad.com	Data analysis
software, algorithm	Cell Ranger Single Cell Software Suite (v2.1.0)	10x genomics	https://support.10xgenomics.com	scRNA-seq data analysis
software, algorithm	RStudio	RStudio IDE	https://posit.co/	scRNA-seq data analysis
software, algorithm	R 4.4.1	R-project	https://www.r-project.org/	scRNA-seq data analysis

software, algorithm	Seurat	Satijalab	https://satijalab.org/seurat/	scRNA-seq data analysis

Resource availability

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Chang Chen (changchen@moon.ibp.ac.cn).

Materials availability
All unique reagents generated in this study will be made available from the contact. We may require a completed MTA if there is potential for commercial application.

Data and code availability
This paper does not report new raw scRNA-seq data and original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Experimental Model and Subject Details

Single-cell RNA sequencing data analysis

In this study, single-cell RNA sequencing (scRNA-seq) raw data of Müller glia enriched after G/R cone ablation at 72 hpi are from(**Krylov et al., 2023**), embryonic RPCs at 24 hpf, and postembryonic retinal stem cells (RSCs) at 14 dpf are from(**Xu et al., 2020**). We re-processed the single-cell FASTQ sequencing reads (Novogene) and converted to digital gene expression matrices using the Cell Ranger software (version 3.1.0) provided by 10X Genomics after mapping to the zebrafish GRCz11 (Ensembl release-96) genome assembly. An average of 47,545 mean reads per cell with 1,343 median genes per cell in no ablation control, 63,177 mean reads per cell with 1,556 median genes per cell in G/R cone ablation (lws2_72 hpi), 61,772 mean reads per cell with 2,474 median genes per cell in the embryonic RPCs at 24 hpf, and 49,243 mean reads per cell with 1,181 median genes per cell in the postembryonic retinal stem cells (RSCs) at 14 dpf were obtained.

We then used the *cellranger aggr* (version 6.0, https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/using/aggregate) to aggregate the *cellranger* counts from the two datasets and normalize them to the same sequencing depth. The feature barcode matrix was recalculated and analyzed on the combined data. This resulted in a dataset of 19,706 aggregated cells, with 100% of the no ablation control reads retained and 71.4% of the *lws2_3 dpi* reads retained. The aggregated matrix for the no ablation control and lws2_72 hpi samples was loaded into the Seurat R package (version 4.3.0, https://satijalab.org/seurat/). Cells with gene expression of more than 200 and less than 4000, lower than 5% mitochondrial content were filtered for further analysis. The filtered data were normalized, scaled, and clustered using principal component analysis (PCA) with a significance threshold of $p < 0.001$ (FindClusters, resolution = 0.5), and a UMAP was computed using scanpy.tl. umap, resulting in 20 distinct clusters. MG clusters were identified by high expression of marker genes, including *rlbp1a*, *fabp7a*, *slc1a2b*, *glula*, *glulb*, *gfap*, and *her4.1*. Proliferative MG clusters were identified from proliferative cell clusters by high expression of

proliferative cell markers, including *pcna*, *mki67*, *gfap*, *her4.1*, and low expression of

the CMZ markers, including *fabp11a*, *col15α1b*, *fabp7b*, *rx2*. We further analyzed the

MG and proliferative MG clusters to obtain 13 clusters, including no ablation control

5,932 cells and G/R cone ablation 72 hpi 3,999 cells in the UMAP plot (**Figure 1-**

figure supplement 1C). We compared the proportion of cell numbers in each cluster

before and after injury and chose clusters with a high proportion of cell numbers in

G/R cone ablation 72 hpi (clusters 2, 3, 5, 6, 9, 10, 11, 12, and 13) to further cluster

into 10 clusters (**Figure 1-figure supplement 1D**).

Single-cell RNA sequencing data of embryonic RPCs at 24 hpf and

postembryonic RCSs at 14 dpf were generated into 10 and 11 clusters, respectively.

Each cluster is identified by the cluster-specific marker gene and different

development stage markers, including *fabp11a*, *col15α1b*, *cxcl18b*, *her4.2*, *npm1a*,

her9, *fabp7a*, *dla*, *atoh7*, *vsx1*, *mafba*, *otx5*, and *rem1* (**Figure 3A, 3C and Figure 3-**

figure supplement 1A and B).

Pseudo-time trajectory analysis

676   After the UMAP cluster analysis of the single-cell data, trajectory analysis was

677   performed to investigate the pseudo-time transcriptomic change of these ten clusters

678   using the 'monocle 2' R package, which revealed three distinct states (**Figure 1G**).

Zebrafish husbandry and transgenic fish lines

All experimental zebrafish embryos, larvae, and adults were produced, grown,

and maintained at 28°C according to standard protocols. Embryos and larvae were

kept in embryo medium (E3; 5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl$_2$·2H$_2$O, 0.33

mM MgSO$_4$•7 H$_2$O, 1.3×10^{-5} % w/v methylene blue in RO water) at 28.5 °C, under

a 14:10 light: dark cycle. Animal procedures performed in this study were approved

by the Animal Use Committee of the Institute of Neuroscience, Chinese Academy of

Sciences (NA-069-2023).

Published fish lines used in this study include the following: AB (Wild type, WT)

and transgenic lines (Tg): *Tg(opn1lw2: NTR-mCherry)^{uom3}* (ZDB-ALT-201012-

2)**(Wang et al., 2020)** in this study named *Tg(lws2: nfsb-mCherry)*, *Tg(gfap:*

EGFP)^{mi2001} (ZDB-FISH-150901–29307)**(Bernardos and Raymond, 2006)**,

Tg(her4.1: dRFP) (ZDB-TGCONSTRCT-070612–2)**(Yeo et al., 2007)**, *Tg(mpeg1:*

GFP) (ZDB-TGCONSTRCT-170801–5)**(Ellett et al., 2011)**, *Tg(pcna: GFP)* (ZDB-

LAB-070129-2)**(Xu et al., 2020)**, *Tg(ef1α: loxP-DsRed-loxP-EGFP)***(Hans et al.,**

2009), *Tg(Tp1bglob: EGFP)***(Yu and He, 2019)**.

Newly generated transgenic fishline contained: *Tg(cxcl18b: GFP)*, *Tg(cxcl18b:*

Cre-vmhc: mCherry:: ef1α: loxP-DsRed-loxP-EGFP:: lws2: nfsb-mCherry), and

mutants *nos1$^{+/-}$*, *nos2a$^{+/-}$*, *nos2b$^{+/-}$*, *gsnor$^{+/-}$*, *nos1$^{-/-}$*, *nos2a$^{-/-}$*, *nos2b$^{-/-}$*, *gsnor$^{-/-}$*.

Generation of *Tg(cxcl18b: GFP)*

The plasmid of *pTol2-cxcl18b: GFP* (10 ng/ μl) and *tol2* mRNA (50 ng/ μl) were

co-injected into a wild-type embryo at the one-cell stage. Zebrafish embryos with

green fluorescent were grown at the E3 medium according to standard protocol and

selected as the founder (F0). The adult F0 fish crossed with the wild-type selected the

fish with GFP^{+} offspring as F1. In this stable transgenic line, *cxcl18b^{+}* cells expressed

the fluorescent protein GFP and the full name of this line is *Tg(cxcl18b: GFP)*.

Generation of *Tg(cxcl18b: Cre-vmhc: mCherry:: ef1α: loxP-DsRed-loxP-*
EGFP:: lws2: nfsb-mCherry)

The plasmid of *pTol2-cxcl18b: Cre-vmhc: mCherry* (10 ng/ μl) and *tol2* mRNA

(50 ng/ μl) were co-injected into the embryo obtained from *Tg(lws2: nfsb-mCherry)*

cross *Tg(ef1α: loxP-DsRed-loxP-EGFP)* at the one-cell stage. Zebrafish embryos

were maintained at E3 medium according to standard protocol. Larvae with *mCherry*

expression in the heart and eyes were selected and grown as the founder (F0) for

further construction. The adult founder was crossed with wild-type (AB), considering

the expression of *cxcl18b* during zebrafish development, larvae with mCherry and

GFP double positive were selected and grown as the F1. In this zebrafish line, EGFP

permanently marks all cells that have expressed *cxcl18b*, as well as their entire lineage

of progeny. The full name of this line is *Tg(cxcl18b: Cre-vmhc: mCherry:: ef1α:*

loxP-DsRed-loxP-EGFP:: lws2: nfsb-mCherry).

Generation of mutant fish lines

The CRISPR/Cas9 system was employed to efficiently and precisely generate mutant zebrafish lines**(Wu et al., 2018)**. Two sgRNAs targeting gene coding sequences involved in nitric oxide metabolism signaling (nitric oxide synthase, *Nos*; and S-nitrosoglutathione reductase, GSNOR) were designed based on the zebrafish GRCz11 genome assembly.

Primers were designed approximately 200 bp from the sgRNA target sites for genotyping (sgRNAs and genotyping primers are listed in **Supplementary Table S1**). We mixed the two sgRNAs (100 ng/µl) for each target gene with Cas9 protein (400 ng/µl) and injected them into transgenic fishline *Tg(lws2: nfsb-mCherry)* embryos at the one-cell stage. Adult fish were crossed with wild-type, and offspring displaying *mCherry* expression in the eyes at 5 dpf were selected for further genotyping. Mutants were confirmed by sequencing, and the fish with open reading frame (ORF) shift were collected as heterozygous fish for further studies.

Plasmid construction

We cloned the *cxcl18b* regulatory element (3048 bp upstream of the start codon including the *5'UTR*) as the *cxcl18b* promoter from the zebrafish genomic DNA **(Figure 2C)**. The primers used for amplification were 5'-GCATTTGTCTCCTCATGCATTGACTAC-3' (forward primer) and 5'-TTGCTGCAAACTATATGTAGGAAATGCTG-3' (reversed primer). For constructing the plasmids *pTol2-cxcl18b: GFP*, *pTol2-cxcl18b: Cre-vmhc: mCherry*, and *pTol2-cxcl18b: gal4FF*, each DNA elements cassette was inserted into the *pDestTol2pA2* vector**(Kwan et al., 2007)**. The plasmid of *pUAS: Cas9T2AGFP; U6: sgRNA1; U6: sgRNA2* was kindly provided by Professor Filippo Del Bene.**(Di Donato et al., 2016)**. To prepare for plasmid *pTol2-UAS: Cas9-T2A-Cre-U6-empty* and *pTol2-UAS: Cas9-T2A-Cre-U6: nos2b sgRNA1; U6: nos2b sgRNA2*, the same two *nos2b* sgRNAs used to construct *nos2b* mutant fish were inserted into *pUAS: Cas9T2AGFP; U6: sgRNA1; U6: sgRNA2* plasmid, with the *GFP* element replaced by *Cr*e using

ClonExpressMultiS One Step Cloning Kit (Vazyme, C113□02). For the plasmid

construction of the MG-specific knockout system based on recombinant adenoviral

vectors, regulatory elements were inserted into adenoviral vectors *pAdc68-S*

according to the standard protocol outlined by Jia et al.(**Jia et al., 2019**). The final

plasmids generated included, *pAdC68XY2-E1-CMV-GFP pAdC68XY2-E1-10xUAS:*

Cas9-2a-Cre-U6: sgRNA1(nos2b); U6: sgRNA2(nos2b), pAdC68XY2-E1-cxcl18b:

gal4FF and *pAdC68XY2-E1-10xUAS: Cas9-2a-Cre-U6: empty* as the control. Plasmid

construction primer sequences are listed in **Supplementary Table S2**.

Metronidazole treatment

Zebrafish larvae were exposed to a 10 mM metronidazole (MTZ) solution

(Sigma-Aldrich, M3761-100G) in standard fish water. For *Tg(lws2: nfsb-mCherry)*

larvae processed for immunohistochemical analysis, MTZ treatment was initiated at 6

dpf to ablate green and red cones. Larvae were maintained in the MTZ solution at

densities of fewer than 50 larvae per 50 mL petri dish and kept at 28.5°C. The MTZ

solution was refreshed every 24 hours to ensure continuous cone ablation until

fixation in 4% PFA, while control larvae were kept in standard fish water.

Clone analysis of *cxcl18b* lineage-traced MG

Clone analysis of $cxcl18b^+$ lineage-traced MG was restricted to cells located in

the central and dorsal region of the zebrafish retina after G/R cone ablation in Figure

2, Figure 6, and their figure supplement. This spatial restriction strongly suggests that

the proliferative MG originate from local mature MG, although we cannot completely

rule out the possibility that CMZ-derived progenitors contribute to the generation of

proliferative MG in the peripheral retina.

Gene disruption via CRISPR/Cas9 system

We used the CRISPR/Cas9 system for efficient gene disruption. Two sgRNAs

were designed to target the coding sequences of genes involved in the nitric oxide

signaling pathway (including *nos1*, *nos2a*, *nos2b*, *gsnor*) and *cxcl18b*. The sgRNAs

were in vitro transcribed and purified using the LiCl precipitation approach

(MEGAscript T7 Transcription Kit, Invitrogen, AM1334). A mixture of the two

sgRNAs (in total 200 ng/μl) with Cas9 protein (400 ng/μl, Novoprotein,

E365 01A) were co-injected into the embryo of *Tg(lws2: nfsb-mCherry)* at the one-

cell stage. The efficiency of gene knockout for each sgRNA was validated, as

presented in **Figure 4-figure supplement 1C**.

Pharmacological treatment with NO scavenger and dexamethasone or
***Nos* Inhibitors intraocular injection**

Zebrafish larvae at 6 dpf were anesthetized in 0.04% MS222 (Sigma, A5040) for

30 to 45 seconds. Inhibitor solutions of 2 μL were prepared as follows: 10 mM *N(ω)-*

nitro-L-arginine methyl ester (L-NAME, Sigma-Aldrich, N5751-1G), a broad *Nos*

inhibitor, diluted in sterile PBS from a 100 mM stock; 10 mM N(ω)-Methyl-L-

arginine acetate salt (L-NMMA, Sigma-Aldrich, M7033-5MG), a specific inhibitor of

nNos, diluted in sterile PBS from a 20 mM stock; and 200 nM 1400W

dihydrochloride (SW1400, MedChemExpress, HY-18730), a specific inhibitor of

iNos, diluted in sterile PBS from a 39.97 mM stock. PBS was used as the control.

Each solution was loaded into glass capillaries prepared using a micropipette puller

(Narishige, PC-10) and connected to a microinjector (Applied Scientific

Instrumentation, MPPI-3). Intraocular injection of the inhibitor solutions were

administered to the zebrafish eyes starting 2 days before G/R cone ablation and

continued until 72 hpi (**Figure 4C**).

Additionally, zebrafish were pre-treated with 10 mM of the NO scavenger,

Phenyl-4,4,5,5-tetramethyl imidazoline-1-oxyl 3-oxide (C-PTIO, Sigma-Aldrich,

P5084-25MG), diluted in standard water (from a 100 mM stock), or dexamethasone

(Dex, Sigma-Aldrich, D1756) which diluted in DMSO, starting 2 days before G/R

cone ablation and continuing until 72 hpi (**Figure 2J and 4C**).

The real-time quantitative PCR after FACS-sorted MG

We crossed *Tg(lws2: nfsb-mCherry)* with *Tg(pcna: GFP)*, *Tg(her4.1: dRFP)*,

Tg(gfap: EGFP), and *Tg(cxcl18b: GFP)* to collect retina cell. the retinas were

dissected and the cells dissociated for sorting and enrichment of signal-positive cells

using fluorescence-activated cell sorting (FACS, Beckman, MoFlo XDP). After 72

hours post G/R cone ablation, GFP$^+$/RFP$^+$ cells were collected from the retinas of

Tg(lws2: nfsb-mCherry x *pcna: GFP)* as the proliferative MG group. For the

uninjured MG group, GFP$^+$/RFP$^+$ cells were collected from *Tg(lws2: nfsb-mCherry* x

gfap: EGFP x *her4.1: dRFP)* uninjured retinas, while GFP$^+$/RFP$^+$ cells from injured

retinas represented the 72-hpi MG group, while GFP$^-$/RFP$^-$ cells were enriched as the

other retinal cell types group in the 72 hpi retinas. Additionally, from *Tg(lws2: nfsb-*

mCherry x *cxcl18b: GFP)* retinas, GFP$^+$/RFP$^+$ cells were collected as the *cxcl18b$^+$*

MG group and single RFP$^+$ cells as the G/R cone group at 72 hpi **(Figure 5A)**.

Interesting cells were collected into a 1.5 ml tube containing lysis buffer

(20 mg/ml PK in TE buffer) and total RNA was isolated using this lysis buffer.

cDNA synthesis was conducted by adding RT mix contains: 200 U Superscript II

reverse transcriptase (Invitrogen, 18064 014), 1 × First strand buffer (Invitrogen,

18064 014), 5 mM DTT (Invitrogen, 18064 014), 20 U Recombinant RNase

inhibitor (Clontech, 2313A), 6 mM MgCl2 (Sigma, M8266), 1 μM TSO**(Picelli et**

al., 2013); 8% PEG8000 (Sigma, P1458). PCR amplification was performed as

previously described**(Picelli et al., 2013)**.

Real-time quantitative PCR (RT-qPCR) reactions were carried out using TB

Green Premix Ex Taq (Takara, RR420A) on a LightCycler 480 II real-time PCR

detection system (Roche). The qPCR primers are listed in **Supplementary Table S3**.

The ΔΔCt method was used to determine the relative expression of mRNAs in

different group retinae cells and normalized to *actin* mRNA levels. Each group

comparison was performed using a two-way ANOVA followed by Tukey's HSD test.

Error bars represented SEM. ****$p<0.0001$, ***$p<0.001$; **$p<0.01$; *$p<0.05$; ns,

$p>0.05$.

839 Construction and injection of adenoviral-based MG-specific knockout
840 system in the zebrafish retina

The type of AdC68 used in our study is a replication-deficient chimpanzee

adenovirus. The plasmids used for viral packaging were described above, and the

amplification and purification of recombinant chimpanzee adenovirus followed the

standard protocol(**Jia et al., 2019; Liu et al., 2021**). We generated four adenoviruses

for this study, including *AdC68XY2-E1-CMV-GFP pAdC68XY2-E1-10xUAS: Cas9-*

2a-Cre-U6: sgRNA1(nos2b); *U6: sgRNA2(nos2b)* (referred to as *UAS: Cas9-2a-Cre-*

sgnos2b, infectious titer: 2.00 x 10^{13} ifu/ml)*, pAdC68XY2-E1-cxcl18b: gal4FF*

(referred to as *cxcl18b: gal4*, infectious titer: 1.90 x 10^{13} ifu/ml) and *pAdC68XY2-E1-*

10xUAS: Cas9-2a-Cre-U6: empty (referred to as *UAS: Cas9-2a-Cre*, infectious titer:

1.75 x 10^{13} ifu/ml) as a control.

All adenoviruses were diluted in sterile PBS to a final infectious titer: 1.00 x 10^{13}

ifu/ml. The *UAS: Cas9-2a-Cre-sgnos2b* and *cxcl18b: gal4* adenoviruses were mixed

at a 1:1 ratio for the MG-specific knockout system, while the *UAS: Cas9-2a-Cre* with

cxcl18b: gal4 adenoviruses were similarly mixed as a control. To achieve the *cxcl18b$^+$*

MG cell-specific knockout *nos2b*, we performed the intraocular injection of the

adenovirus mixtures into the eyes of *Tg(ef1α: loxP-DsRed-loxP-EGFP* x *lws2: nfsb-*

mCherry) at 5 dpf. One day post-injection, the fish were treated with MTZ for 3

consecutive days to ablate the G/R cone (**Figure 6E**).

Tissue preparation and immunostaining

Zebrafish were fixed in 4% paraformaldehyde (PFA, Electron Microscopy

Services, 157–8) overnight, then cryoprotected in 30% sucrose for 6 hours, flash-

frozen, and cryosectioned at a thickness of 14 μm. Immunostaining was performed

following the protocol described by Tang et al.(**Tang et al., 2017**). The primary

antibody including mouse anti-PCNA (Abcam, ab29) at a 1:500 dilution, as well as

mouse anti-GS (glutamine synthase, BD Transduction Laboratories, 610518), rabbit

anti-BLBP (Abcam, ab32423), rabbit anti-DsRed (Clontech, 632496), rabbit anti-

TaqGFP (Proteintech, 50430-2-AP), and chicken anti-GFP (Abcam, ab13970) each at

a 1:1000 dilution. Secondary antibodies conjugated to Alexa Fluor 488, 594, or 647

(Jackson ImmunoResearch Laboratories Inc.) were used at a 1:1000 dilution. Primary

antibodies were incubated overnight at 4°C, while Alexa Fluor secondary antibodies

were incubated at room temperature for 2 hours. DAPI staining was performed
according to the standard protocol.

In situ hybridization

In this study, three digoxigenin (DIG)-labeled RNA probes targeting endogenous
cxcl18b, fabp11a, and *col15α1b* were synthesized using the MEGAscript™ T7 High
Yield Transcription Kit (Invitrogen, AM1334) and the DIG RNA Labeling Kit (Roche,
11277073910), following the manufacturer's instructions. The cDNA of each gene
was amplified by PCR using the following primers: *cxcl18b* -F: 5'-
ATGGCATTCACACCCAAAGCG-3'; *cxcl18b* -R: 5'-
TAATACGACTCACTATAGGGATTGGCCCTGCTGTTTTTGTG-3'; *fabp11a*-F:
5'-GTTGGAAACCGGACCAAACC-3'; *fabp11a*-R: 5'-
TAATACGACTCACTATAGGGACGGCTCGTTGAGCTTGAAT-3'; *col15a1b* -F:
5'-CCTCAATGGAGGTCCTAAAGGT-3'; *col15α1b* -R: 5'-
TAATACGACTCACTATAGGGACCAGCTTCTGAGACCAAGC-3'.

Following the *in situ* hybridization protocol described by Tang et al.**(Tang et al.,**
2017), fresh zebrafish retinal sections were incubated overnight in the HybEZ system
(Advanced Cell Diagnostics, 310013) at 65°C with 200 ng of probe for each slide.
The next day, slides were sequentially washed in 5x SSC buffer and incubated
overnight at 4°C with an anti-DIG-POD antibody (Roche, 11093274910) diluted
1:500 in TNB buffer (TN buffer with 0.5% blocking reagent; Roche). On the third day,
the signal was detected using the TSA™ Plus Cyanine 3 (PerkinElmer,
NEL744001KT) or Cyanine 5 (PerkinElmer, NEL745001KT)/Fluorescein System.

Imaging

Images were taken using an inverted confocal microscope system (FV1200,
Olympus) confocal microscope using $40 \times$ (silicon oil, 1.05 NA), or $60 \times$ (silicon oil,
1.3 NA) objectives.

Quantifications and statistical analysis

All quantification and visualization were performed with FV31S-SW 2.3.1.163

Viewer (Olympus), and ImageJ. ***Z intensity projection*** was used to process the Z-

stack images acquired from 14-μm-thick sections of zebrafish retina for statistical

analysis. For cell counting, PCNA$^+$ MG and their lineage in this study were defined as

one proliferative MG, *cxcl18b*$^+$ MG and their lineage were counted as one *cxcl18b*$^+$

MG.

To perform the statistical analysis, p values were calculated with GraphPad

Prism 8 (or Microsoft Excel). The unpaired, non-parametric Wilcoxon test was

applied for comparison of two groups. The one-way ANOVA, followed by Tukey's

HSD test was applied for comparison of different groups with one treatment. The two-

way ANOVA followed by Tukey's HSD test was applied for comparison of four

groups with two treatments. Error bars represent SEM. ****$p<0.0001$, ***$p<0.001$;

**$p<0.01$; *$p<0.05$; ns, $p>0.05$.

Main Figure Titles and Legends

Figure 1. Single-cell RNA seq reveals injury-induced the *cxcl18b*-defined MG

transitional states

(**A**) Schematic showing the experimental procedure: five consecutive days of MTZ treatment in *Tg(lws2: nfsb-mCherry* x *mpeg1: GFP)* fish to ablate green and red (G/R) cone, starting at 6 dpf and continuing until 11 dpf. The MTZ solution was refreshed every 24 hours, followed by fish fixation for further immunostaining.

(**B-D**) Quantitative plots showing the dynamic changes in the number of G/R cone (**B**), recruited microglia (**C**), and proliferative MG (PCNA$^+$) (**D**) at different time points after MTZ treatment (uninjured: collected retina number n=14; 24-hpi: n=10; 48-hpi: n=10; 72-hpi: n=11; 96-hpi: n=12; 120-hpi: n=16; mean ± SEM; ****p<0.0001, ***p<0.001, **p<0.01, ns, p>0.05; one-way ANOVA followed by Tukey's HSD test).

(**E**) Representative images showing microglia recruitment (*mpeg1: GFP*, green), G/R cone ablation (*lws2: nfsb-mCherry*, red), and injury-induced MG proliferation (PCNA, white) in uninjured (**E₁**) and 72-hpi (**E₂**) retinas. The high-magnification images of the boxed area (**E₃-E₄**). Scale Bars: 20 μm (**E₁, E₂**), 10 μm (**E₃**), 2 μm (**E₄**).

(**F**) The UMAP plot of 4,172 MG cells was sorted with an increased proportion in response to the G/R cone ablation. Cells were further aggregated into 10 clusters based on previously published scRNA-seq data(**Krylov et al., 2023**).

(**G**) Pseudo-time developmental trajectory of MG states identified by Monocle2 analysis shows a main developmental branch originating from Cluster 4 (*cx43$^+$*), which diverges into two sub-branches: Cluster 0 and Clusters 5/3/6 (*pcna$^+$*).

(**H**) Violin plots showing the expression levels of key genes (*cx43*, *glula*, *gfap*, *cxcl18b*, *ascl1α*, *mki67*, *pcna*, and *notch3*) in the main developmental branch clusters, progressing from the most original MG states (Cluster 4) to transitional MG states (Clusters 1/2/5), and proliferative MG states (Clusters 3/6).

Figure 2. Clonal analysis reveals the proliferative MG mostly originated from

***cxcl18b$^+$* MG transitional states**

(**A**) Representative images show dynamic expression of *cxcl18b* (red, *in situ* hybridization) and PCNA (white, immunostaining) in *Tg(lws2: nfsb-mCherry)* retina at different time points following the G/R cone ablation. Scale bars: 20 μm.

(**B**) Quantitative plots showing the number of *cxcl18b$^+$* (red curve, significance shown above the curve) and PCNA$^+$ MG (blue curve, significance shown below the curve) in uninjured (n=11) and injured retina at 24-hpi (n=10), 48-hpi (n=12), 72-hpi (n=11), 96-hpi (n=7), and 120-hpi (n=5). Each injured time point was compared to the uninjured retina (Mean ± SEM; ****p<0.0001, **p<0.01, ns, p>0.05; one-way ANOVA followed by Tukey's HSD test).

(**C**) Schematic diagram of the *cxcl18b* promoter was used to construct the reporter

fish line *Tg(cxcl18b: GFP)* and clonal analysis fish line *Tg(cxcl18b: Cre-vmhc-*
mCherry:: ef1α: loxP-DsRed-loxP-EGFP; lws2: nfsb-mCherry).

(D) Immunostaining of injury-induced $cxcl18b^+$ (green, indicated by *Tg(cxcl18b:*
GFP)) and proliferative ($PCNA^+$, white) MG showing overlapping in the central
retina area (yellow arrows, $GFP^+/PCNA^+$ MG) at 48 hpi. The high-magnification
images of the boxed area (**D$_3$'-D$_3$'''**). The area of the retina is labeled with a dashed
line, and each layer structure is labeled with dashed lines and marked with the outer
nuclear layer (ONL), inner nuclear layer (INL), inner plexiform layer (IPL), ganglion
cell layer (GCL) and lens. Scale Bars: 20 µm (**D$_1$-D$_2$**) and 5 µm (**D$_3$'-D$_3$'''**).

(E) Quantitative plots showing the number of $cxcl18b^+$ MG (red) and proliferative
MG ($PCNA^+$, blue) in (**D$_1$**) uninjured (n=4) and (**D$_2$**) 48-hpi retina (n=7) (Mean ±
SEM; ****$p<0.0001$, ns, $p>0.05$; two-way ANOVA followed by Tukey's HSD test).

(F) Clonal analysis of injury-induced $cxcl18b^+$ MG in transgenic fish line *Tg(cxcl18b:*
Cre-vmhc: mCherry:: ef1α: loxP-DsRed-loxP-EGFP:: lws2: nfsb-mCherry) at 72 hpi
retina showing overlapping between proliferative ($PCNA^+$, white) MG with $cxcl18b^+$
(green, yellow arrows). The high-magnification images of the boxed area (**F$_3$'-F$_3$'''**).
Scale Bars: 20 µm (**F$_1$-F$_2$**) and 5 µm (**F$_3$'-F$_3$'''**).

(G) Quantitative analysis at 72 hpi shows no significance in the number of
proliferative MG (PCNA+, blue) and double-positive ($PCNA^+$ / $cxcl18b^+$, red) MG
(n=14; Mean ± SEM; ns, $p>0.05$; Unpaired t-test) in (**F**), with 93±2% of $PCNA^+$ MG
also being $cxcl18b^+$.

(H) Representative images show that not all mature MG stained with glutamate
synthase (GS^+, magenta) are $cxcl18b^+$ (green, labeled by *cxcl18b*: Cre) in the central
retinal area (white dashed lines identified a 45° angular region originating from the
optic nerve). The high-magnification images of the boxed area (**H$_2$'-H$_2$'''**). Scale Bars:
20 µm (**H$_1$**) and 5 µm (**H$_2$'-H$_2$'''**).

(I) Quantification of GS^+/*cxcl18b*: Cre^+ double-positive (blue, yellow arrows in **H**)
and GS^+/*cxcl18b*: Cre^- single-positive (red, open yellow arrowheads in **H**) MG (n=14,
Mean ± SEM; *$p<0.05$; one-way ANOVA followed by Tukey's HSD test), and the
proportion of *cxcl18b*: $Cre^{+ or -}$ MG within the total population of mature (GS^+) MG in
the central retina.

(J$_1$-J$_3$) Representative images showing injury-induced $cxcl18b^+$ MG (green) in
Tg(lws2: nfsb-mCherry x cxcl18b: GFP) fish retina treated with dexamethasone (Dex)
or DMSO at 72 hpi. Scale bars: 20 µm.

(K-M) Quantitative plots showing the number of $cxcl18b^+$ MG (72 hpi: n=14; DMSO:
n=6, Dex: n=10) in **J$_1$-J$_3$**; recruited microglia in **Figure 2-figure supplement 1J$_1$-J$_3$**
and proliferative MG in **Figure 2-figure supplement 1J$_4$-J$_6$** (72 hpi: n=9; DMSO:
n=7, Dex: n=14) in *Tg(mpeg1: GFP; lws2: nfsb-mCherry)* retinas after DMSO or Dex
treatment at 72 hpi (Mean ± SEM; ****$p<0.0001$, ***$p<0.001$, ns, $p>0.05$; one-way
ANOVA followed by Tukey's HSD test).

Figure 3. The *cxcl18b*-defined MG transitional states recapitulate the

developmental states of RSCs in the CMZ

(A, C) UMAP plots display 5,368 retinal progenitor cells (RPCs) at 24 hpf and 4,625
CMZ progenitor cells at 14 dpf. Clusters are indicated by their cluster-specific marker
genes based on previously published scRNA-seq data**(Xu et al., 2020)**.
(B, D) UMAP plots showing expression of *fabp11a*, *cxcl18b*, and *col15α1b* at r24-hpf
RPCs (embryonic states) **(B)** and r14-dpf CMZ-progenitors (post-embryonic states)
(D).
(E-H$_3$) *In situ* hybridization images showing the expression of *fabp11a* (green),
col15α1b (white), two putative makers for post-embryonic retinal stem cells (RSCs),
and *cxcl18b* at 30-hpf **(E-F$_3$)** and 14-dpf **(G-H$_3$)** retina. The high-magnification
images of the boxed area **(F-F$_3$, H-H$_3$)**. The area of these three in situ signal trouble
positive is labeled with a dashed yellow line. Scale Bars: 20 μm **(E, G)** and 3 μm **(F-
F$_3$, H-H$_3$)**.
1010

Figure 4. The nitric oxide metabolic pathway regulates injury-induced MG

proliferation

(A) Representative images of microglia recruitment (green, marked by *Tg(mpeg1:*
GFP)), G/R cone ablation (red, marked by *Tg(lws2: nfsb-mCherry)*) and proliferative
MG (white, marked by PCNA$^+$) at 72 hpi with nitric oxide metabolism pathway genes
disruption (*nos1/nos2a/nos2b/gsnor*). Scale Bars: 20 μm.
(B) Quantitative analysis of the number of proliferative MG (PCNA$^+$) at 72 hpi in **(A)**.
In total, we collected 11 retinas for WT (n=11), *scramble* sgRNA-injection (n=7),
nos1 sgRNA-injection (n=23), *nos2a* sgRNA-injection (n=15), *nos2b* sgRNA-
injection (n=22), and *gsnor* sgRNA-injection (n=13) (Mean ± SEM; ****p<0.0001,
**p<0.01, *p<0.05, ns, p>0.05; one-way ANOVA followed by Tukey's HSD test).
(C) Schematic showing the experimental procedure of nitric oxide synthase (*Nos*)
inhibitors injection or NO scavengers treatment in *Tg(lws2: nfsb-mCherry* x *mpeg1:*
GFP) retina starting from 5 dpf for five consecutive days to 10 dpf with three
consecutive days of MTZ treatment for G/R cone ablation from 7 dpf to 10 dpf. *Nos*
inhibitors, NO scavengers, and MTZ solution were refreshed every 24 hours, and fish
fixation was at 10 dpf for further immunostaining.
(D) Representative images of microglial recruitment (green, marked by *Tg(mpeg1:*
GFP)), G/R cone ablation (red, marked by *Tg(lws2: nfsb-mCherry)*) and proliferative
MG (white, marked by PCNA$^+$) at 72 hpi following L-NMMA (10 mM), L-NAME
(10 mM), 1400W (200 nM) intraocular injection, and PBS as control, or C-PTIO (10
mM) treatment. Scale Bars: 20 μm.
(E) Quantitative plots showing the number of proliferative (PCNA$^+$) MG at 72 hpi in
(D). Retinas analyzed WT (n=11), PBS-injected (n=16), L_NMMA-injected (n=12),
L_NAME-injected (n=12), 1400W-injected (n=10), and C-PTIO treatment (n=27)
(Mean ± SEM; ****p<0.0001, ns, p>0.05; one-way ANOVA followed by Tukey's
HSD test).
(F-G) Representative images of proliferative MG (PCNA$^+$, white) and G/R cone
ablation (marked by *Tg(lws2: nfsb-mCherry)*, red) at 72 hpi in heterozygous (*nos1$^{+/-}$*,

nos2a^{+/-}, *nos2b^{+/-}*, *gsnor^{+/-}*) **(F)** and homozygous mutants **(G)** of nitric oxide
metabolism pathway genes (*nos1^{-/-}*, *nos2a^{-/-}*, *nos2b^{-/-}*, *gsnor^{-/-}*). Scale Bars: 20 μm.
(H-I) Quantitative plots showing the number of proliferative MG (white, PCNA$^+$) at
72 hpi in *nos* and *gsnor* mutant fish. In heterozygous **(H)**, analyzed retinas include
WT (n=14), *nos1^{+/-}* (n=19), *nos2a^{+/-}* (n=13), *nos2b^{+/-}* (n=20), *gsnor^{+/-}* (n=12). For
homozygous **(I)**, analyzed retinas include *nos1^{-/-}* (n=18), *nos2a^{-/-}* (n=20), *nos2b^{-/-}*
(n=20), *gsnor^{-/-}* (n=27) (Mean ± SEM; ****$p<0.0001$, ns, $p>0.05$; one-way
ANOVA followed by Tukey's HSD test).
(J-K) Quantitative plots showing the number of proliferative MG (white, PCNA$^+$) **(J)**
and photoreceptor cells remain **(K)** at 72 hpi in *nos2b* hetero- or homozygous mutants.
(Mean ± SEM; ****$p<0.0001$, ***$p<0.001$, ns, $p>0.05$; one-way ANOVA followed
by Tukey's HSD test).
1052

Figure 5. RT-qPCR analysis reveals *nos2b* cell-specific expression in the injury-

induced *cxcl18b*$^+$ MG

(A) Schematic showing the workflow for isolating and enriching for three post-injury
MG populations (72-hpi MG, 72-hpi PCNA$^+$ MG, and 72-hpi *cxcl18b*$^+$ MG) and three
control groups (uninjured MG, 72-hpi retinal cells other than MG, and 72-hpi G/R
cones) using fluorescence-activated cell sorting (FACS).
(B-C) RT-qPCR analysis of *nos1*, *nos2a*, *nos2b*, and *gsnor* expression in different cell
populations. Expression levels are shown relative to 72-hpi G/R cones **(B)** and other
72-hpi retinal cell types **(C)**. A total of six independent replicates were performed for
cell population enrichment and cDNA template preparation (n=6, Mean ± SEM;
****$p<0.0001$, ns, $p>0.05$; two-way ANOVA followed by Tukey's HSD test).
(D) RT-qPCR analysis comparing *nos1*, *nos2a*, *nos2b*, and *gsnor* expression in
distinct MG states relative to uninjured MG (repeats n=7 in *cxcl18b*$^+$ MG; n=5 in 72-
hpi MG; and repeats n=6 in PCNA$^+$ MG; n=6 in the uninjured retina; Mean ± SEM;
****$p<0.0001$, ns, $p>0.05$; two-way ANOVA followed by Tukey's HSD test).
1068

Figure 6. NO produced by *nos2b* in *cxcl18b*$^+$ MG regulates injury-induced

proliferation

(A-C) Representative images showing the adenovirus-mediated infection (green,
indicated by Y$_2$-GFP) specifically target MG (red, GS staining) in the zebrafish retina.
The virus was intraocularly injected into the right eye of *Tg(lws2: nfsb-mCherry* x
ef1α: loxP-DsRed-loxP-EGFP) fish **(A$_2$, C)**, with the left eye as a wild-type (WT)
control **(A$_1$)**. The high-magnification images of the boxed area (B-B$_3$). Scale Bars:
20 μm (A$_1$, A$_2$) and 5 μm (B-B$_3$).
(D) Schematic showing the design of the *cxcl18b*$^+$ MG-specific *nos2b* knockout
system. The viral construct consists of three plasmids: (1) *gal4* expression driven by
the *cxcl18b* promoter; (2) *UAS*-derived *Cas9* and *Cre* elements, and (3) *U6* promoters

driving two sgRNAs targeting *nos2b*, with a non-targeting sgRNA as the control.

(E) Schematic showing the procedure of injury process and intraocular viral injection

in *Tg(lws2: nfsb-mCherry* x *ef1α: loxP-DsRed-loxP-EGFP)* fish.

(F-G) Representative images showing proliferative MG (PCNA$^+$, white) with

cxcl18b$^+$ MG specific knockout *nos2b* in **(G)** and control in **(F)**, (GFP$^+$, green, yellow

arrows) are defined as virus-infected clones. Upper panels show WT retina (no virus

injected),

Bottom panels show retinas injected with the virus (two sgRNA targets as *nos2b*

knockout and without sgRNA targets as control). Scale Bars: 20 μm.

(H) Quantification of proliferative (PCNA$^+$/GFP$^+$, red bars) and non-proliferative

(PCNA$^-$/GFP$^+$, gray bars) MG clones in **(F)**. For control, ~75% of virus-infected

clones entered the cell cycle (PCNA$^+$; red bars), with 90/120 clones analyzed across 8

independent experiments (n=8). For *nos2b* knockout clones, ~23% entered the cell

cycle (PCNA$^+$; blue bars), with 22/103 clones analyzed across 6 independent

experiments (n=6) (Mean ± SEM; ****p<0.0001; two-way ANOVA followed by

Tukey's HSD test).

1096

Figure 7. NO regulates MG proliferation by suppressing Notch signaling

(A) Dot plot showing the Notch signaling-related gene expression in different MG

states. The average expression levels of these genes for all cells in each cluster are

coded by the gray level. The percentage of cells expressing each gene within each

cluster is coded by dot size.

(B) Representative images showing the dynamic changes of Notch signaling activity

(green, indicated by *Tg(Tp1: EGFP)*) and proliferative MG (white, PCNA$^+$) following

injury with nitric oxide (NO) blockade using C-PTIO. Scale Bars: 20 μm.

(C-D) Quantitative plots showing the number of Notch activation (*tp1*: GFP$^+$ MG) in

(C) and proliferative MG (PCNA$^+$ MG) in **(D)** at different time points post-G/R cone

ablation. We collected uninjured retina (24-hpi, n=4; 48-hpi, n=4; 72-hpi, n=3),

injured retina (24-hpi, n=9; 48-hpi, n=4; 72-hpi, n=6), and retina treated with C-PTIO

(24-hpi, n=6; 48-hpi, n=5; 72-hpi, n=7) (Mean ± SEM; ****p<0.0001, ***p<0.001,

**p<0.01, *p<0.05, ns, p>0.05; two-way ANOVA followed by Tukey's HSD test).

1111

1112

Figure 1-figure supplement 1. Clusters with increased proportion are identified

from the scRNA-seq data

(A) Representative images showing microglia recruitment (green, indicated by *Tg(mepg1: GFP)*), G/R cone ablation (red, *Tg(lws2: nfsb-mCherry)*), and injury-induced MG proliferation (white, marked by PCNA) following the injury process from uninjured retina to 120 hpi. Scale Bars: 20 μm.

(B) Representative images showing PCNA+ (white, immunostaining) cells co-expressed with BLBP (red, immunostaining) at 72 hpi. Scale Bars: 20 μm (B_1-B_3)

and 5 μm (**B₄**).

(C) UMAP plot showing 5,932 cells from uninjured retinas (blue) and 3,999 cells

from 72-hpi (red) retinas were obtained, which were further aggregated into 13

clusters.

(D) Percentage of MGs from uninjured and 72-hpi retinas in each cluster. Eight

clusters (6, 5, 9, 11, 12, 3, 2, 10) were identified and showed an increased proportion

for further analysis.

(E) Dot plot showing the expression levels of marker genes for quiescent MG (qMG),

reactive MG (rMG), proliferative MG (pMG), and differentiated MG (dMG). The

average expression levels of these genes for all cells in each cluster are coded by the

gray level. The percentage of cells expressing each gene within each cluster is coded

by dot size.

1133

1134

Figure 2-figure supplement 1. Gene description of *cxcl18b* does not reduce MG

proliferation

(A-A$_3$) *In situ* hybridization images show injury-induced *cxcl18b* (red, marked by *in situ* hybridization) with MG-like morphology, cell-specific expression in the BLBP$^+$ cells (green, labeled by immunostaining) at 24 hpi in the transgenic fish line *Tg(lws2: nfsb-mCherry)* retina. In these clones, *cxcl18b$^+$* MG are red$^+$/BLBP$^+$ cells (yellow arrowheads). The high-magnification images of the boxed area **(A$_1$-A$_3$)**. The area of the retina structure is labeled with a dashed line, and each layer structure is labeled with dashed lines and marked with the outer nuclear layer (ONL), inner nuclear layer (INL), inner plexiform layer (IPL), ganglion cell layer (GCL), and lens. Scale Bars: 20 μm **(A)** and 5 μm **(A$_1$-A$_3$)**.

(B-B$_3$) Representative images showing *cxcl18b* (red, *in situ* hybridization) co-expressed with PCNA (white, immunostaining) at 72 hpi. In these clones, *cxcl18b$^+$*

MG was identified as red$^+$/PCNA$^+$ cells (yellow arrowheads). The high-magnification
images of the boxed area **(B$_1$-B$_3$)**. Scale Bars: 20 μm **(B)** and 5 μm **(B$_1$-B$_3$)**.
(C) Schematic showing the design of the *cxcl18b* reporter fish line and the Cre-loxP
transgenic fish line used for clone analysis. The *cxcl18b* promoter drives GFP or Cre
expression, with the heart labeled by mCherry under the *vmhc* promoter for fish
screening.
(D-E$_3$) Representative images showing *cxcl18b* expression labeled by *Tg(cxcl18b:*
GFP) (green) in **(D-D$_3$)** at 48 hpi, or three transgenic fish *Tg(cxcl18b: Cre-vmhc:*
mCherry:: ef1α: loxP-DsRed-loxP-EGFP:: lws2: nfsb-mCherry) (green) in **(E-E$_3$)** at
24 hpi fish retina, merged with *cxcl18b* in situ hybridization signal (red) **(D-E$_3$)**. The
high-magnification images of the boxed area **(D$_1$-D$_3$, E$_1$-E$_3$)**. Scale Bars: 20 μm **(D,**
E) and 5 μm **(D$_1$-D$_3$, E$_1$-E$_3$)**.
(F) Schematic showing the two sgRNA target sites of the *cxcl18b* genome.
(G) Summary table showing the indel and knockout efficiency of the two sgRNAs
targeting *cxcl18b*.
(H) Representative images of proliferative MG (PCNA$^+$, white) at 72 hpi in *Tg(lws2:*
nfsb-mCherry) fish retina with *cxcl18b* disruption. Scale Bars: 20 μm.
(I) Quantitative plots showing the number of proliferative MG (PCNA$^+$) at 72 hpi in
H. In total, retinas were collected from wild-type (WT, n=11), *scramble* sgRNA-
injected (*scrambled*, n=7), and *cxcl18b* sgRNA-injected (*cxcl18b* KO, n=7) (Mean ±
SEM; ns, $p > 0.05$; one-way ANOVA followed by Tukey's HSD test).
(J$_1$-J$_6$) Representative images showing injury-induced microglia recruitment (green,
marked by *Tg(mpeg1: GFP)*) **(J$_1$-J$_3$)**, and proliferative MG (white, marked by PCNA)
(J$_4$-J$_6$) after dexamethasone (Dex) or DMSO treatment in the in *Tg(mpeg1: GFP x*
lws2: nfsb-mCherry) fish retina at 72 hpi. Scale bars: 20 μm.
1173

1174

Figure 3-figure supplement 1. UMAP plots show the expression of cluster-specific

marker genes utilized to identify developmental states

(A, B) UMAP plots showing the expression pattern of embryonic developmental stage cluster-specific marker genes (*npm1a*, *her9*, *her4.2*, *fabp7a*, *dla*, and *atoh7*) in **(A)** and post-embryonic developmental stage marker genes (*her4.2*, *atoh7*, *vsx1*, *mafba*, *otx5*, and *rem1*) in **(B)**.

(C) UMAP plots showing the expression pattern of *fabp11a*, *cxcl18b*, and *col15α1b* at MG clusters with increased population after G/R cone ablation at 72 hpi in Figure 1F.

(D) Representative images showing the *cxcl18b* expression (green, marked by *Tg(cxcl18b: GFP)*) are located in the most peripheral region of the CMZ (white, marked by PCNA) at 5 dpf. The high-magnification images of the boxed area **(D$_1$-D$_3$)**. Scale Bars: 20 µm **(D)** and 5 µm **(D$_1$-D$_3$)**.

1189

Figure 4-figure supplement 1. Genetic disruption of the nitric oxide pathway

modulates G/R cone ablation and microglial recruitment

(**A**) Dot plot showing the redox gene expression profiles of regulating glutathione (GSH), hydrogen peroxide (H_2O_2), nicotinamide adenine dinucleotide phosphate (NADPH), nitric oxide (NO), metal ion, and sulfhydryl group (-SH) in the uninjured and $cxcl18b^+$ MG in 72-hpi scRNA-seq data (**Krylov et al., 2023**). The average expression levels of these genes for all cells in each cluster were coded by the gray level. The percentage of cells expressing each gene within each cluster was coded by dot size.

(**B**) Schematic showing the two sgRNA target sites of the nitric oxide metabolism pathway genome (*nos1/nos2a/nos2b/gsnor*).

(**C**) Summary table showing the indel and knockout efficiency of the two sgRNAs targeting *nos1/nos2a/nos2b/gsnor*.

(**D**) Representative images of microglial recruitment (marked by *Tg(mpeg1: GFP)*, green) and G/R cone ablation (marked by *Tg(lws2: nfsb-mCherry)*, red) at 72 hpi with *nos1/nos2a/nos2b/gsnor* disruption, with *scrambled* sgRNA-injected as the control.

Scale Bars: 20 μm.

(E-F). Quantitative plots showing the number of recruited microglia (GFP[+]) **(D)** and

G/R cone ablation (mCherry[+]) **(E)** at 72 hpi with *nos1/nos2a/nos2b/gsnor* disruption.

In total, we collected 11 retinas for WT (n=11), *scrambled* sgRNA-injected (n=7),

nos1 sgRNA-injected (n=10), *nos2a* sgRNA-injected (n=6), *nos2b* sgRNA-injected

(n=12), and *gsnor* sgRNA-injected (n=13) (Mean ± SEM; $*p<0.05$; ns, $p>0.05$;

one-way ANOVA followed by Tukey's HSD test).

1213

1214

Figure 4-figure supplement 2. The genotype of nitric oxide metabolism pathway

mutants

(A) Sequence alignment of WT nitric oxide metabolism pathway genes
(*nos1*/*nos2a*/*nos2b*/*gsnor*) and their CRISPR-cas9-induced mutant alleles.
(B) Summary table showing the premature stop codon and unknown polypeptides
sides of each mutant.
1221

Figure 5-figure supplement 1. *In situ* **hybridization reveals** *nos2b* **cell-specific**

expression in the *cxcl18b***-defined transitional MG states**

(A-A₃) Representative *in situ* hybridization images showing *nos2b* expression (red, *in situ* hybridization) cell-specific in the injury-induced *cxcl18b⁺* MG (GFP⁺, green) in *Tg(lws2: nfsb-mCherry* x *cxcl18b: GFP)* fish retina at 72 hpi. Clones with green/red/white trouble-positive are *cxcl18b⁺/nos2b⁺*/PCNA⁺ MG cells (yellow arrowheads). The high-magnification images of the boxed area **(C-C₃)**. Scale Bars: 20 μm (A, B) and 5 μm **(C₁-C₃)**.

(D-E) RT-qPCR analysis of *glula* (pink bar) and *glulb* (green bar) expression in different cell populations relative to 72-hpi G/R cone **(D)**, or other retinal cell types **(E)**. The significance of 72-hpi G/R cone as control is noted by * for comparisons with 72-hpi G/R cones and # for comparisons with uninjured MG. In total, we do 6 independent replicates to enrich the different cell populations and get the cDNA templates (Mean ± SEM; **** and ####p<0.0001, **p<0.001, **p<0.01, ns, p>0.05; two-way ANOVA followed by Tukey's HSD test).

(F) RT-qPCR analysis of *cxcl18b* (blue bar) and *pcna* (orange bar) expression in different MG states. (Mean ± SEM; ****p<0.0001, **p<0.01, *p< 0.05, ns, p>0.05; two-way ANOVA followed by Tukey's HSD test).

1242

Figure 6-figure supplement 1. Analysis of cell-specific knockout efficiency for

nos2b **in virus-infected clones**

(A) Schematic showing the cell-specific gene knockout system design. Injury-induced
cxcl18b promoter directs the expression of *gal4*, which in combination with the *UAS*
element, drives the cell-specific expression of Cas9 and Cre. The *U6* promoter
enables the broad expression of sgRNAs targeting *nos2b*, and Cas9 mediates gene
knockout within the targeted cells through these sgRNAs. The *ef1α* promoter controls
the *loxP-DsRed-loxP-EGFP* reporter system, allowing for the switch from DsRed to
EGFP upon Cas9 and Cre activity, and GFP[+] MG clones are identified with successful
nos2b knockout. This system operates with high specificity via double adenovirus
infection.
(B) Schematic showing the procedure of virus-infected clones are sorted and enriched
by FACS for further knockout efficiency analysis.
(C) Representative gel electrophoresis image showing *nos2b* deletions in knockout
cells compared to controls. Single-virus injections were used to assess Cre system
leakage. While minimal Cre system leakage was observed, it did not result in
detectable *nos2b* knockout.
(D) Summary table showing the knockout efficiency of each virus-injected clone.

1261

1262 **Supplementary Table S1. sgRNA sequences and genotyping primers, related to STAR**
1263 **Methods.**

sgRNA targets sequences	
Gene symbol	**Primer forward**
nos1 sgRNA target-1	TAATACGACTCACTATAGGCTCCCAGGAAAGAGTCGCGTTTTAGAGCTAGAAATAGC
nos1 sgRNA target-2	TAATACGACTCACTATAGGTCACCAGGCCTGTCAACCGTTTTAGAGCTAGAAATAGC
nos2a sgRNA target-1	TAATACGACTCACTATAGGAGGAACTCGCTTTTGGGGGTTTTAGAGCTAGAAATAGC
nos2a sgRNA target-2	TAATACGACTCACTATAGGTGGCTGGTGCCACCGATGGTTTTAGAGCTAGAAATAGC
nos2b sgRNA target-1	TAATACGACTCACTATAGGAGTATTTCAGAAACACATGTTTTAGAGCTAGAAATAGC
nos2b sgRNA target-2	TAATACGACTCACTATAGGATGTGTTACATCACAGAGGTTTTAGAGCTAGAAATAGC
gsnor sgRNA target-1	TAATACGACTCACTATAGGCCCATGAAGTTCGAGTGAGTTTTAGAGCTAGAAATAGC
gsnor sgRNA target-2	TAATACGACTCACTATAGGTCATCCTGGGTCATGAGGGTTTTAGAGCTAGAAATAGC

1264

Genotyping primers for gene disruption		
Gene symbol	**Primer forward**	**Primer reverse**
nos1-sgTarget-1	ATCGTCTCGGACCTCATCCG	ATAGGTTTTCTATTGCCCTGAGCTC
nos1-sgTarget-2	ATGCGAACGATGCTCGC	ATCCTTCCTATCGATGACCCTCTGG
nos2a-sgTarget-1	ATTACTGTTCTGTCGACTTGATAAACACG	ATACTTAAGATGAGTGCACAAGGCATG
nos2a-sgTarget-2	ATTTAGCCACTCTGAAATTCAAGCC	ATCAGTCCTTTGAAGCTGACCTTGC
nos2b-sgTarget-1	ATTGCAGCACGTTATACTCTCTTAGC	AATGGGCACTGCTTGACTTTAT
nos2b-sgTarget-2	ACTCACAAGAGCGAACTACTGC	ATTGACCCTTCACAAATCTTAGACG
gsnor-sgTarget-1	ATTAATTTTCAGGTGATCAAATGTAAGGC	ATGGGGTGAAGTTGAGTCAGC
gsnor-sgTarget-2	ATGTATTCAAAGTGTGTAATGTAGTCGG	TACCTGATTTTCTGGCACAGG

1265

1266

1267 **Supplementary Table S2. Plasmid construction primer sequences, related to STAR**
1268 **Methods.**

Plasmid construction primer sequences			
Plasmid	**Elements**	**Primer forward**	**Primer reverse**
pTol2-cxcl18b: GFP	cxcl18b-promoter	TAGGGCGAATTGGGCATTTGT CTCCTCATGCATTGACTAC	AATTGCTGCAAACTATATGT AGGAAATGCTG
	GFP	ATGGTGAGCAAGGGC	TTACTTGTACAGCTCGTCC ATGC
pTol2-cxcl18b: gal4FF	gal4FF	ATGAAGCTACTGTCTTCTATC GAACAAGC	CTACCCACCGTACTCGTCA ATTCC
pTol2-cxcl18b: Cre-vmhc: mCherry	Cre	ATGGCCAATTTACTGACCGTA CACC	CTAATCGCCATCTTCCAGC AGGC
	vmhc: mCherry	CATGTGTCCTAAATTCTGAAA ACACAGTAAGCC	TTACTTGTACAGCTCGTCC ATG
pTol2-UAS: Cas9-T2A-Cre-U6: nos2b sgRNA1; U6: nos2b sgRNA2	nos2b sgRNA1	CTCGGGTCGCTCTTGTGAGT GACC	AAACGGTCACTCACAAGA GCGACC
	nos2b sgRNA2	AGCTCGGGCTGGTGCCGCCA ATGTCG	TAAACGACATTGGCGGCAC CAGCCCG

1269
1270

1271 **Supplementary Table S3. qPCR primer sequences, related to STAR Methods.**

Gene symbol	Primer forward	Primer reverse
cxcl18b	TGCTGCTCGCGGTAGTTTAC	TCTGCAGTAATTGGCCCTGC
glula	CGCATTACAGAGCCTGCCTA	ATTCCAGTTGCCTGGGATCG
glulb	ATGGTCCAGAACCAGCATCC	TCACAAGGGCCAACTTGGAA
pcna	CAGAAACCTAGCCATGGGGG	GTCTTGGACAGAGGAGTGGC
nos1	TGAGCACGTTTGCATAGGGT	CACTGGATTCGTCCCACACA
nos2a	CCAGATAACCACTGCTCTGCT	CTTGTCTTCCCATCTTGCTAAAA
nos2b	GAGGCAAAATTCCACCACCAG	TGCAAAGTGGTCATGGGACA
gsnor	ACAAGAAATCTCCACCCGGC	GACGGCTCGGATACTCTTCC
β-actin	ATTGCTGACAGGATGCAGAAG	GATGGTCCAGACTCATCGTACTC

1272

1273

Reference: Uncategorized References

Abalo, X.M., Lagman, D., Heras, G., Del Pozo, A., Eggert, J., and Larhammar, D. (2020).

Circadian regulation of phosphodiesterase 6 genes in zebrafish differs between cones and

rods: Implications for photopic and scotopic vision. Vision research *166*, 43-51.

Abraham, E., Hartmann, H., Yoshimatsu, T., Baden, T., and Brand, M. (2024). Restoration of

cone-circuit functionality in the regenerating adult zebrafish retina. Developmental cell *59*,

2158-2170.e2156.

Asaoka, Y., Hata, S., Namae, M., Furutani-Seiki, M., and Nishina, H. (2014). The Hippo

pathway controls a switch between retinal progenitor cell proliferation and photoreceptor cell

differentiation in zebrafish. PloS one *9*, e97365.

Ashpole, N.M., Chawla, A.R., Martin, M.P., Brustovetsky, T., Brustovetsky, N., and Hudmon,

A. (2013). Loss of calcium/calmodulin-dependent protein kinase II activity in cortical

astrocytes decreases glutamate uptake and induces neurotoxic release of ATP. The Journal

of biological chemistry *288*, 14599-14611.

Ashpole, N.M., Song, W., Brustovetsky, T., Engleman, E.A., Brustovetsky, N., Cummins, T.R.,

and Hudmon, A. (2012). Calcium/calmodulin-dependent protein kinase II (CaMKII) inhibition

induces neurotoxicity via dysregulation of glutamate/calcium signaling and hyperexcitability.

The Journal of biological chemistry *287*, 8495-8506.

Bedi, O., Sapra, V., Kumar, M., and Krishan, P. (2024). Newer mitochondrial dynamics and

their role of calcium signalling in liver regeneration. Mitochondrion *79*, 101969.

Bernardos, R.L., and Raymond, P.A. (2006). GFAP transgenic zebrafish. Gene expression

patterns : GEP *6*, 1007-1013.

Boyd, P., Campbell, L.J., and Hyde, D.R. (2023). Clcf1/Crlf1a-mediated signaling is

neuroprotective and required for Müller glia proliferation in the light-damaged zebrafish retina.

Frontiers in cell and developmental biology *11*, 1142586.

Bradley, S., Tossell, K., Lockley, R., and McDearmid, J.R. (2010). Nitric oxide synthase

regulates morphogenesis of zebrafish spinal cord motoneurons. The Journal of neuroscience :

the official journal of the Society for Neuroscience *30*, 16818-16831.

Breus, O., and Dickmeis, T. (2021). Genetically encoded thiol redox-sensors in the zebrafish

model: lessons for embryonic development and regeneration. Biological chemistry *402*, 363-

378.

Campbell, L.J., Hobgood, J.S., Jia, M., Boyd, P., Hipp, R.I., and Hyde, D.R. (2021). Notch3

and DeltaB maintain Müller glia quiescence and act as negative regulators of regeneration in

the light-damaged zebrafish retina. Glia *69*, 546-566.

Campbell, L.J., Levendusky, J.L., Steines, S.A., and Hyde, D.R. (2022). Retinal regeneration

requires dynamic Notch signaling. Neural regeneration research *17*, 1199-1209.

Celotto, L., Rost, F., Machate, A., Bläsche, J., Dahl, A., Weber, A., Hans, S., and Brand, M.

(2023). Single-cell RNA sequencing unravels the transcriptional network underlying zebrafish

retina regeneration. eLife *12*.

Charles, N., Ozawa, T., Squatrito, M., Bleau, A.M., Brennan, C.W., Hambardzumyan, D., and

Holland, E.C. (2010). Perivascular nitric oxide activates notch signaling and promotes stem-

like character in PDGF-induced glioma cells. Cell stem cell *6*, 141-152.

Conner, C., Ackerman, K.M., Lahne, M., Hobgood, J.S., and Hyde, D.R. (2014). Repressing

notch signaling and expressing TNFα are sufficient to mimic retinal regeneration by inducing

Müller glial proliferation to generate committed progenitor cells. The Journal of neuroscience :

the official journal of the Society for Neuroscience *34*, 14403-14419.

Curado, S., Anderson, R.M., Jungblut, B., Mumm, J., Schroeter, E., and Stainier, D.Y. (2007).

Conditional targeted cell ablation in zebrafish: a new tool for regeneration studies.

Developmental dynamics : an official publication of the American Association of Anatomists

236, 1025-1035.

Curado, S., Stainier, D.Y., and Anderson, R.M. (2008). Nitroreductase-mediated cell/tissue

ablation in zebrafish: a spatially and temporally controlled ablation method with applications in

developmental and regeneration studies. Nature protocols *3*, 948-954.

Dermietzel, R., Kremer, M., Paputsoglu, G., Stang, A., Skerrett, I.M., Gomes, D., Srinivas, M.,

Janssen-Bienhold, U., Weiler, R., Nicholson, B.J.*, et al.* (2000). Molecular and functional

diversity of neural connexins in the retina. The Journal of neuroscience : the official journal of

the Society for Neuroscience *20*, 8331-8343.

Di Donato, V., De Santis, F., Auer, T.O., Testa, N., Sánchez-Iranzo, H., Mercader, N.,

Concordet, J.P., and Del Bene, F. (2016). 2C-Cas9: a versatile tool for clonal analysis of gene

function. Genome research *26*, 681-692.

Ellett, F., Pase, L., Hayman, J.W., Andrianopoulos, A., and Lieschke, G.J. (2011). mpeg1

promoter transgenes direct macrophage-lineage expression in zebrafish. Blood *117*, e49-56.

Fausett, B.V., Gumerson, J.D., and Goldman, D. (2008). The proneural basic helix-loop-helix

gene ascl1a is required for retina regeneration. The Journal of neuroscience : the official

journal of the Society for Neuroscience *28*, 1109-1117.

Förstermann, U., and Sessa, W.C. (2012). Nitric oxide synthases: regulation and function.

European heart journal *33*, 829-837, 837a-837d.

Gao, H., A, L., Huang, X., Chen, X., and Xu, H. (2021). Müller Glia-Mediated Retinal

Regeneration. Molecular neurobiology *58*, 2342-2361.

Goldman, D. (2014). Müller glial cell reprogramming and retina regeneration. Nature reviews

Neuroscience *15*, 431-442.

Goldstein, S., Russo, A., and Samuni, A. (2003). Reactions of PTIO and carboxy-PTIO with

*NO, *NO2, and O2-*. The Journal of biological chemistry *278*, 50949-50955.

Gonzalez-Nunez, V., Nocco, V., and Budd, A. (2010). Characterization of drCol 15a1b: a

novel component of the stem cell niche in the zebrafish retina. Stem cells (Dayton, Ohio) *28*,

1399-1411.

Gorsuch, R.A., Lahne, M., Yarka, C.E., Petravick, M.E., Li, J., and Hyde, D.R. (2017). Sox2

regulates Müller glia reprogramming and proliferation in the regenerating zebrafish retina via

Lin28 and Ascl1a. Experimental eye research *161*, 174-192.

Goumenaki, P., Günther, S., Kikhi, K., Looso, M., Marín-Juez, R., and Stainier, D.Y.R. (2024).

The innate immune regulator MyD88 dampens fibrosis during zebrafish heart regeneration.

Nature cardiovascular research *3*, 1158-1176.

Grover, L.M., and Teyler, T.J. (1990). Two components of long-term potentiation induced by

different patterns of afferent activation. Nature *347*, 477-479.

Han, P., Zhou, X.H., Chang, N., Xiao, C.L., Yan, S., Ren, H., Yang, X.Z., Zhang, M.L., Wu, Q.,

Tang, B., *et al.* (2014). Hydrogen peroxide primes heart regeneration with a derepression

mechanism. Cell research *24*, 1091-1107.

Hans, S., Kaslin, J., Freudenreich, D., and Brand, M. (2009). Temporally-controlled site-

specific recombination in zebrafish. PloS one *4*, e4640.

Hanson, Q.M., Carley, J.R., Gilbreath, T.J., Smith, B.C., and Underbakke, E.S. (2018).

Calmodulin-induced Conformational Control and Allostery Underlying Neuronal Nitric Oxide

Synthase Activation. Journal of molecular biology *430*, 935-947.

Hoang, T., Wang, J., Boyd, P., Wang, F., Santiago, C., Jiang, L., Yoo, S., Lahne, M., Todd,

L.J., Jia, M., *et al.* (2020). Gene regulatory networks controlling vertebrate retinal regeneration.

Science (New York, NY) *370*.

Hong, J., Kim, B.K., Lim, H., Lee, S., and Lee, S.J. (2012). Identification and characterization

of triamcinolone acetonide, a microglial-activation inhibitor. Immunopharmacology and

immunotoxicology *34*, 912-918.

Hunter, M.V., Willoughby, P.M., Bruce, A.E.E., and Fernandez-Gonzalez, R. (2018). Oxidative

Stress Orchestrates Cell Polarity to Promote Embryonic Wound Healing. Developmental cell

47, 377-387.e374.

Iribarne, M., and Hyde, D.R. (2022). Different inflammation responses modulate Müller glia

proliferation in the acute or chronically damaged zebrafish retina. Frontiers in cell and

developmental biology *10*, 892271.

Jaeschke, H. (2000). Reactive oxygen and mechanisms of inflammatory liver injury. Journal

of gastroenterology and hepatology *15*, 718-724.

Janssen-Bienhold, U., Dermietzel, R., and Weiler, R. (1998). Distribution of connexin43

immunoreactivity in the retinas of different vertebrates. The Journal of comparative neurology

396, 310-321.

Jia, W., Channappanavar, R., Zhang, C., Li, M., Zhou, H., Zhang, S., Zhou, P., Xu, J., Shan,

S., Shi, X., *et al.* (2019). Single intranasal immunization with chimpanzee adenovirus-based

vaccine induces sustained and protective immunity against MERS-CoV infection. Emerging

microbes & infections *8*, 760-772.

Jones, R.J., Jourd'heuil, D., Salerno, J.C., Smith, S.M., and Singer, H.A. (2007). iNOS

regulation by calcium/calmodulin-dependent protein kinase II in vascular smooth muscle.

American journal of physiology Heart and circulatory physiology *292*, H2634-2642.

Jorstad, N.L., Wilken, M.S., Grimes, W.N., Wohl, S.G., VandenBosch, L.S., Yoshimatsu, T.,

Wong, R.O., Rieke, F., and Reh, T.A. (2017). Stimulation of functional neuronal regeneration

from Müller glia in adult mice. Nature *548*, 103-107.

Kawamoto, E.M., Vivar, C., and Camandola, S. (2012). Physiology and pathology of calcium

signaling in the brain. Frontiers in pharmacology *3*, 61.

Kimura, M., Moteki, H., and Ogihara, M. (2023). Role of Hepatocyte Growth Regulators in

Liver Regeneration. Cells *12*.

Krylov, A., Yu, S., Veen, K., Newton, A., Ye, A., Qin, H., He, J., and Jusuf, P.R. (2023).

Heterogeneity in quiescent Müller glia in the uninjured zebrafish retina drive differential

responses following photoreceptor ablation. Frontiers in molecular neuroscience *16*, 1087136.

Kwan, K.M., Fujimoto, E., Grabher, C., Mangum, B.D., Hardy, M.E., Campbell, D.S., Parant,

J.M., Yost, H.J., Kanki, J.P., and Chien, C.B. (2007). The Tol2kit: a multisite gateway-based

construction kit for Tol2 transposon transgenesis constructs. Developmental dynamics : an

official publication of the American Association of Anatomists *236*, 3088-3099.

Kyritsis, N., Kizil, C., Zocher, S., Kroehne, V., Kaslin, J., Freudenreich, D., Iltzsche, A., and

Brand, M. (2012). Acute inflammation initiates the regenerative response in the adult

zebrafish brain. Science (New York, NY) *338*, 1353-1356.

Lahne, M., Brecker, M., Jones, S.E., and Hyde, D.R. (2020a). The Regenerating Adult

Zebrafish Retina Recapitulates Developmental Fate Specification Programs. Frontiers in cell

and developmental biology *8*, 617923.

Lahne, M., Nagashima, M., Hyde, D.R., and Hitchcock, P.F. (2020b). Reprogramming Müller

Glia to Regenerate Retinal Neurons. Annual review of vision science *6*, 171-193.

Leach, L.L., Hanovice, N.J., George, S.M., Gabriel, A.E., and Gross, J.M. (2021). The

immune response is a critical regulator of zebrafish retinal pigment epithelium regeneration.

Proceedings of the National Academy of Sciences of the United States of America *118*.

Lee, M.S., Jui, J., Sahu, A., and Goldman, D. (2024). Mycb and Mych stimulate Müller glial

cell reprogramming and proliferation in the uninjured and injured zebrafish retina.

Development (Cambridge, England) *151*.

Lee, M.S., Wan, J., and Goldman, D. (2020). Tgfb3 collaborates with PP2A and notch

signaling pathways to inhibit retina regeneration. eLife *9*.

Lenkowski, J.R., and Raymond, P.A. (2014). Müller glia: Stem cells for generation and

regeneration of retinal neurons in teleost fish. Progress in retinal and eye research *40*, 94-123.

Levin, M. (2009). Bioelectric mechanisms in regeneration: Unique aspects and future

perspectives. Seminars in cell & developmental biology *20*, 543-556.

Liu, J., Xu, K., Xing, M., Zhuo, Y., Guo, J., Du, M., Wang, Q., An, Y., Li, J., Gao, P.*, et al.*

(2021). Heterologous prime-boost immunizations with chimpanzee adenoviral vectors elicit

potent and protective immunity against SARS-CoV-2 infection. Cell discovery *7*, 123.

López-Juárez, A., Titus, H.E., Silbak, S.H., Pressler, J.W., Rizvi, T.A., Bogard, M., Bennett,

M.R., Ciraolo, G., Williams, M.T., Vorhees, C.V., *et al.* (2017). Oligodendrocyte Nf1 Controls

Aberrant Notch Activation and Regulates Myelin Structure and Behavior. Cell reports *19*, 545-

557.

Lu, C., and Hyde, D.R. (2024). Cytokines IL-1β and IL-10 are required for Müller glia

proliferation following light damage in the adult zebrafish retina. Frontiers in cell and

developmental biology *12*, 1406330.

Lyu, P., Iribarne, M., Serjanov, D., Zhai, Y., Hoang, T., Campbell, L.J., Boyd, P., Palazzo, I.,

Nagashima, M., Silva, N.J., *et al.* (2023). Common and divergent gene regulatory networks

control injury-induced and developmental neurogenesis in zebrafish retina. Nat Commun *14*,

8477.

Matrone, G., Jung, S.Y., Choi, J.M., Jain, A., Leung, H.E., Rajapakshe, K., Coarfa, C., Rodor,

J., Denvir, M.A., Baker, A.H., *et al.* (2021). Nuclear S-nitrosylation impacts tissue regeneration

in zebrafish. Nat Commun *12*, 6282.

Meng, J., Lv, Z., Zhang, Y., Wang, Y., Qiao, X., Sun, C., Chen, Y., Guo, M., Han, W., Ye, A.,

et al. (2021). Precision Redox: The Key for Antioxidant Pharmacology. Antioxidants & redox

signaling *34*, 1069-1082.

Moore, P.K., al-Swayeh, O.A., Chong, N.W., Evans, R.A., and Gibson, A. (1990). L-NG-nitro

arginine (L-NOARG), a novel, L-arginine-reversible inhibitor of endothelium-dependent

vasodilatation in vitro. British journal of pharmacology *99*, 408-412.

Nelson, C.M., Ackerman, K.M., O'Hayer, P., Bailey, T.J., Gorsuch, R.A., and Hyde, D.R.

(2013). Tumor necrosis factor-alpha is produced by dying retinal neurons and is required for

Muller glia proliferation during zebrafish retinal regeneration. The Journal of neuroscience :

the official journal of the Society for Neuroscience *33*, 6524-6539.

Pacher, P., Beckman, J.S., and Liaudet, L. (2007). Nitric oxide and peroxynitrite in health and

disease. Physiological reviews *87*, 315-424.

Park, M. (2018). AMPA Receptor Trafficking for Postsynaptic Potentiation. Frontiers in cellular

neuroscience *12*, 361.

Parsons, M.J., Pisharath, H., Yusuff, S., Moore, J.C., Siekmann, A.F., Lawson, N., and Leach,

S.D. (2009). Notch-responsive cells initiate the secondary transition in larval zebrafish

pancreas. Mechanisms of development *126*, 898-912.

Picelli, S., Björklund Å, K., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg, R.

(2013). Smart-seq2 for sensitive full-length transcriptome profiling in single cells. Nature

methods *10*, 1096-1098.

Pollak, J., Wilken, M.S., Ueki, Y., Cox, K.E., Sullivan, J.M., Taylor, R.J., Levine, E.M., and

Reh, T.A. (2013). ASCL1 reprograms mouse Muller glia into neurogenic retinal progenitors.

Development (Cambridge, England) *140*, 2619-2631.

Powell, C., Cornblath, E., Elsaeidi, F., Wan, J., and Goldman, D. (2016). Zebrafish Müller glia-

derived progenitors are multipotent, exhibit proliferative biases and regenerate excess

neurons. Scientific reports *6*, 24851.

Qin, C., Qi, Z., Pan, S., Xia, P., Kong, W., Sun, B., Du, H., Zhang, R., Zhu, L., Zhou, D., *et al.*

(2023). Advances in Conductive Hydrogel for Spinal Cord Injury Repair and Regeneration.

International journal of nanomedicine *18*, 7305-7333.

Quillien, A., Moore, J.C., Shin, M., Siekmann, A.F., Smith, T., Pan, L., Moens, C.B., Parsons,

M.J., and Lawson, N.D. (2014). Distinct Notch signaling outputs pattern the developing

1472 arterial system. Development (Cambridge, England) *141*, 1544-1552.

1473 Ramachandran, R., Fausett, B.V., and Goldman, D. (2010). Ascl1a regulates Müller glia

1474 dedifferentiation and retinal regeneration through a Lin-28-dependent, let-7 microRNA

1475 signalling pathway. Nature cell biology *12*, 1101-1107.

1476 Raymond, P.A., Barthel, L.K., Bernardos, R.L., and Perkowski, J.J. (2006). Molecular

1477 characterization of retinal stem cells and their niches in adult zebrafish. BMC developmental

1478 biology *6*, 36.

1479 Rees, D.D., Palmer, R.M., Schulz, R., Hodson, H.F., and Moncada, S. (1990).

1480 Characterization of three inhibitors of endothelial nitric oxide synthase in vitro and in vivo.

1481 British journal of pharmacology *101*, 746-752.

1482 Rochon, E.R., Missinato, M.A., Xue, J., Tejero, J., Tsang, M., Gladwin, M.T., and Corti, P.

1483 (2020). Nitrite Improves Heart Regeneration in Zebrafish. Antioxidants & redox signaling *32*,

1484 363-377.

1485 Sahu, A., Devi, S., Jui, J., and Goldman, D. (2021). Notch signaling via Hey1 and Id2b

1486 regulates Müller glia's regenerative response to retinal injury. Glia *69*, 2882-2898.

1487 Shen, Y.C., and Raymond, P.A. (2004). Zebrafish cone-rod (crx) homeobox gene promotes

1488 retinogenesis. Developmental biology *269*, 237-251.

1489 Sifuentes, C.J., Kim, J.W., Swaroop, A., and Raymond, P.A. (2016). Rapid, Dynamic

1490 Activation of Müller Glial Stem Cell Responses in Zebrafish. Investigative ophthalmology &

1491 visual science *57*, 5148-5160.

1492 Silva, N.J., Nagashima, M., Li, J., Kakuk-Atkins, L., Ashrafzadeh, M., Hyde, D.R., and

1493 Hitchcock, P.F. (2020). Inflammation and matrix metalloproteinase 9 (Mmp-9) regulate

photoreceptor regeneration in adult zebrafish. Glia *68*, 1445-1465.

Tang, X., Gao, J., Jia, X., Zhao, W., Zhang, Y., Pan, W., and He, J. (2017). Bipotent

progenitors as embryonic origin of retinal stem cells. The Journal of cell biology *216*, 1833-

1847.

Thummel, R., Kassen, S.C., Enright, J.M., Nelson, C.M., Montgomery, J.E., and Hyde, D.R.

(2008). Characterization of Müller glia and neuronal progenitors during adult zebrafish retinal

regeneration. Experimental eye research *87*, 433-444.

Torraca, V., Otto, N.A., Tavakoli-Tameh, A., and Meijer, A.H. (2017). The inflammatory

chemokine Cxcl18b exerts neutrophil-specific chemotaxis via the promiscuous chemokine

receptor Cxcr2 in zebrafish. Developmental and comparative immunology *67*, 57-65.

Vihtelic, T.S., and Hyde, D.R. (2000). Light-induced rod and cone cell death and regeneration

in the adult albino zebrafish (Danio rerio) retina. Journal of neurobiology *44*, 289-307.

Villegas, S.N., Gombos, R., García-López, L., Gutiérrez-Pérez, I., García-Castillo, J., Vallejo,

D.M., Da Ros, V.G., Ballesta-Illán, E., Mihály, J., and Dominguez, M. (2018). PI3K/Akt

Cooperates with Oncogenic Notch by Inducing Nitric Oxide-Dependent Inflammation. Cell

reports *22*, 2541-2549.

Wan, J., and Goldman, D. (2016). Retina regeneration in zebrafish. Current opinion in

genetics & development *40*, 41-47.

Wan, J., and Goldman, D. (2017). Opposing Actions of Fgf8a on Notch Signaling Distinguish

Two Muller Glial Cell Populations that Contribute to Retina Growth and Regeneration. Cell

reports *19*, 849-862.

Wan, J., Ramachandran, R., and Goldman, D. (2012). HB-EGF is necessary and sufficient for

Müller glia dedifferentiation and retina regeneration. Developmental cell *22*, 334-347.

Wang, H., Yang, Z., Li, X., Huang, D., Yu, S., He, J., Li, Y., and Yan, J. (2020). Single-cell in

vivo imaging of cellular circadian oscillators in zebrafish. PLoS biology *18*, e3000435.

White, D.T., Sengupta, S., Saxena, M.T., Xu, Q., Hanes, J., Ding, D., Ji, H., and Mumm, J.S.

(2017). Immunomodulation-accelerated neuronal regeneration following selective rod

photoreceptor cell ablation in the zebrafish retina. Proceedings of the National Academy of

Sciences of the United States of America *114*, E3719-e3728.

Wu, D.M., Schneiderman, T., Burgett, J., Gokhale, P., Barthel, L., and Raymond, P.A. (2001).

Cones regenerate from retinal stem cells sequestered in the inner nuclear layer of adult

goldfish retina. Investigative ophthalmology & visual science *42*, 2115-2124.

Wu, K., Ren, R., Su, W., Wen, B., Zhang, Y., Yi, F., Qiao, X., Yuan, T., Wang, J., Liu, L., *et al.*

(2014). A novel suppressive effect of alcohol dehydrogenase 5 in neuronal differentiation. The

Journal of biological chemistry *289*, 20193-20199.

Wu, R.S., Lam, II, Clay, H., Duong, D.N., Deo, R.C., and Coughlin, S.R. (2018). A Rapid

Method for Directed Gene Knockout for Screening in G0 Zebrafish. Developmental cell *46*,

112-125.e114.

Xiao, X., Liao, Z., and Zou, J. (2023). Genetic and epigenetic regulators of retinal Müller glial

cell reprogramming. Advances in ophthalmology practice and research *3*, 126-133.

Xu, B., Tang, X., Jin, M., Zhang, H., Du, L., Yu, S., and He, J. (2020). Unifying developmental

programs for embryonic and postembryonic neurogenesis in the zebrafish retina.

Development (Cambridge, England) *147*.

Yao, K., Qiu, S., Tian, L., Snider, W.D., Flannery, J.G., Schaffer, D.V., and Chen, B. (2016).

Wnt Regulates Proliferation and Neurogenic Potential of Müller Glial Cells via a Lin28/let-7

miRNA-Dependent Pathway in Adult Mammalian Retinas. Cell reports *17*, 165-178.

Yeo, S.Y., Kim, M., Kim, H.S., Huh, T.L., and Chitnis, A.B. (2007). Fluorescent protein

expression driven by her4 regulatory elements reveals the spatiotemporal pattern of Notch

signaling in the nervous system of zebrafish embryos. Developmental biology *301*, 555-567.

Yoo, S.K., Freisinger, C.M., LeBert, D.C., and Huttenlocher, A. (2012). Early redox, Src family

kinase, and calcium signaling integrate wound responses and tissue regeneration in zebrafish.

The Journal of cell biology *199*, 225-234.

Yu, C., Li, X., Ma, J., Liang, S., Zhao, Y., Li, Q., and Zhang, R. (2024). Spatiotemporal

modulation of nitric oxide and Notch signaling by hemodynamic-responsive Trpv4 is essential

for ventricle regeneration. Cellular and molecular life sciences : CMLS *81*, 60.

Yu, S., and He, J. (2019). Stochastic cell-cycle entry and cell-state-dependent fate outputs of

injury-reactivated tectal radial glia in zebrafish. eLife *8*.

Yurco, P., and Cameron, D.A. (2005). Responses of Müller glia to retinal injury in adult

zebrafish. Vision research *45*, 991-1002.

Zhang, H., Hao, J., Hong, H., Gu, W., Li, Z., Sun, J., Zhan, H., Wei, X., and Zhou, L. (2024).

Redox signaling regulates the skeletal tissue development and regeneration. Biotechnology &

genetic engineering reviews *40*, 2308-2331.

Zhang, Z., Hou, H., Yu, S., Zhou, C., Zhang, X., Li, N., Zhang, S., Song, K., Lu, Y., Liu, D., *et*

al. (2020). Inflammation-induced mammalian target of rapamycin signaling is essential for

retina regeneration. Glia *68*, 111-127.

1559

A r24 hpf_RPCs

fabp11a her4.2 2 4 UMAP_2 / UMAP_1 5,368 cells

B *fabp11a* *cxcl18b* *col15a1b*

C r14 dpf_CMZ progenitors

mafba *otx5* *fabp11a* her4.2 *atoh7* *vsx1* *rem1* UMAP_2 / UMAP_1 4,625 cells

D

In situ hybridzation DAPI / *fabp11a* / *cxcl18b* / *col15α1b*

E 30 hpf **F** **F₁** *fabp11a* **F₂** *cxcl18b* **F₃** *col15α1b*

G 14 dpf **H** **H₁** *fabp11a* **H₂** *cxcl18b* **H₃** *col15α1b*

A1 Left retina **A2** Right retina — DAPI / Virus-Y2-EGFP / GS — Non-virus-injected — Virus-injected

B1 merge **B2** Y2-EGFP **B3** GS

C Virus injection (Fish right retina)

Tg(lws2: nfsb-mCherry x ef1α: loxP-DsRed-loxP-EGFP)

D

Y2-cxcl18b: gal4FF — cxcl18b: promoter / gal4 / VP16AD / PolyA

Y2-U6: UAS: cas9-2a-cre-U6 — 5x UAS / nls / cas9 / nls / t2A / cre / U6 / U6 / sgRNA scaffold

Y2-U6: sgnos2b-T1-UAS: cas9-2a-cre-U6: sgnos2b-T2 — 5x UAS / nls / cas9 / nls / t2A / cre / U6 / T2 / U6 / T1

sgRNA scaffold
T1 — *sgRNA nos2b-Target_1*
T2 — *sgRNA nos2b-Target_2*

E *Tg(lws2: nfsb-mCherry x ef1α: loxP-DsRed-loxP-EGFP)*

dpf: 0 | 5 | 6 | 7 | 8 | 9 | 10 | 11

MTZ | MTZ | MTZ

▽ Virus injection
⚡ G/R cone ablation
🔶 MTZ treatment
△ Fixation
△ Immunostaining

F *UAS*: Cas9-2a-Cre + *cxcl18b*: gal4
Uninjected
Virus-injected — merge / Virus infected-MG / PCNA

G *UAS*: Cas9-2a-Cre-*sgnos2b* + *cxcl18b*: gal4
Uninjected
Virus-injected — merge / Virus infected-MG / PCNA

H
75 ± 13%
23 ± 8%
****

■ PCNA+ / GFP+ (Ctrl)
■ PCNA+ / GFP+ (KO *nos2b*)
■ PCNA- / GFP+