Integrative Analysis of Single-cell and Bulk RNA Sequencing

Reveals Macrophage Heterogeneity in Lyme Arthritis

Yan Dong[1,2#], Yantong Chen[1#], Yanshuang Luo[1], Meng Liu[1], Yingchen Zhou[3] ,Yutian

Yang[4], Xuesong Chen[1], Xiaorong Liu[1], Fusong Yang[1], Zhi Liang[1], Liye Zi[1],

Hongbing Gao[1], Zhichao Cao[1], Fucun Gao[1], Chao Song[1*], Guozhong Zhou[1,2*]

[1]Department of Pain Medicine, The Affiliated Anning First People's Hospital of

Kunming University of Science and Technology, Kunming, Yunnan Province, 650302,

China;

[2]School of Basic Medical Sciences, Kunming University of Science and Technology,

Kunming, Yunnan Province, 650500, China;

[3]Clinical Medical College, Kunming University, Kunming, Yunnan Province, 650500,

China;

[4] First School of Clinical Medicine, Kunming Medical University, Kunming, Yunnan

Province, 650500, China;

[#]Yan Dong and Yantong Chen contributed equally to this work.

[*]*Correspondence to:*

Chao Song, Department of Pain Medicine, The Affiliated Anning First People's

Hospital of Kunming University of Science and Technology, Kunming, Yunnan

Province, 650302, China

Guozhong Zhou, Department of Pain Medicine, The Affiliated Anning First People's

Hospital of Kunming University of Science and Technology, Kunming, Yunnan

Province, 650302, China; School of Basic Medical Sciences, Kunming University of

Science and Technology, Kunming, Yunnan Province, 650500, China;

Abstract

Introduction Lyme arthritis (LA) is a common late-stage manifestation of Lyme

disease caused by *Borrelia burgdorferi* (Bb) infection. Macrophages play a crucial

role in LA pathogenesis, yet their heterogeneity and functional states within the

arthritic microenvironment remain poorly understood.

Methods We integrated single-cell RNA sequencing data from ankle joint tissues of

Bb-infected C57BL/6 mice with bulk RNA sequencing data from both infected ankle

joints and bone marrow-derived macrophages. We developed a LA mouse model and

validated key findings through histopathological examination and RT-qPCR analysis.

Results We identified nine distinct cell types in the joint microenvironment with

significant compositional shifts during disease progression. Analysis of 16,617

Macrophages/Monocytes revealed seven functionally distinct subpopulations with

differential responses to infection. Pseudo-time trajectory analysis demonstrated three

principal differentiation pathways from monocyte-derived macrophages toward

anti-inflammatory, TNF+, and IL1b+ inflammatory phenotypes. Cell-cell

communication analysis revealed significantly altered TGF-β signaling networks in

LA, with the Tgfb1-(Tglbr1+TgIbr2) axis playing a critical role. GSVA and pathway

analyses showed metabolic reprogramming from oxidative phosphorylation toward

glycolysis in macrophages during infection. Through integrated analysis and LASSO

regression, we identified four characteristic genes (SIRPB1C, FABP5, MMP14, and

EGR1) as potential LA biomarkers and developed a diagnostic nomogram with high

predictive accuracy.

Conclusion Our integrative analysis provides comprehensive insights into

macrophage heterogeneity and functional plasticity in LA, identifying characteristic

biomarkers and potential therapeutic targets. The metabolic reprogramming and

altered TGF-β signaling networks we identified may contribute to disease

pathogenesis and offer new avenues for intervention strategies.

1. Introduction

Lyme arthritis (LA), the most common and persistent late-stage manifestation of

Lyme disease, develops from infection with the tick-borne spirochete Borrelia

burgdorferi (Bb), typically emerging months after initial infection[1,2]. Characterized

by intermittent or persistent swelling and pain predominantly in large joints, especially the knee, LA significantly impairs quality of life. Without antibiotic treatment, approximately 60% of Lyme disease patients develop arthritis[3,4]. The condition's complexity arises from pathogen-host immune system interactions that trigger inflammation and, in antibiotic-resistant cases, chronic arthritis[5,6]. Despite extensive research into immune responses, bacterial components, and genetic susceptibility, the complete pathophysiological mechanisms of LA remain incompletely understood.

Extensive research has shown that macrophages play a central role in the pathogenesis of LA[7,8]. Upon Bb infection, they secrete CCL4 and CCL2 chemokines that recruit monocytes and T cells to infection sites while amplifying inflammation through phagocytosis and antigen presentation[9,10]. Both animal and human studies confirm that macrophages induce arthritis directly and through pro-inflammatory cytokines (IL-1β, TNF-α, IFN-γ), which maintain and exacerbate joint inflammation[11-13]. However, macrophages also contribute to *Borrelia* clearance and inflammation resolution[14]. Their activity is regulated by polarization states, cytokine environment, and immune cell networks, making a comprehensive understanding of their role in LA pathogenesis essential.

With the rapid development of single-cell RNA sequencing (scRNA-seq) technology, researchers have been able to analyze tissue heterogeneity at the single-cell level[15-17], facilitating precise studies on the roles of individual cells such as macrophages in the progression of LA. However, studies integrating scRNA-seq with bulk RNA sequencing to investigate macrophage roles in LA remain scarce. In this study, we established a Bb-infected C57BL/6 mouse model and performed transcriptome sequencing of ankle joint tissues to analyze gene expression changes during arthritis development. We implemented a multi-omics integrative approach, correlating mouse joint tissue transcriptome data with publicly available bone marrow-derived macrophage transcriptome and scRNA-seq datasets to construct a comprehensive gene expression network elucidating host immune responses to Bb infection. This multidimensional approach enhances our understanding of

macrophages' central role in LA pathogenesis and provides theoretical foundation for

developing targeted therapies.

2. Methods

2.1 Single-cell RNA Sequencing Data Quality Control and Processing

The scRNA-seq data of LA was accessed from the GSE233850 dataset, which is

established in the GEO database corresponding to a published article "Single-cell

RNA sequencing of murine ankle joints over time reveals distinct transcriptional

changes following *Borrelia burgdorferi* infection" [18]. The single-cell RNA

sequencing data processing workflow is as follows: Raw data was obtained from

ankle joint samples of 1 normal and 4 *Borrelia burgdorferi*-infected C57BL/6J mice

from the GSE233850 dataset. Quality control criteria include: more than 500

expressed genes per cell, each gene expressed in at least 3 cells, and mitochondrial

RNA content less than 20%[19]. Data processing using the Seurat package:

Normalization was performed using the NormalizeData function (natural logarithm of

expression values multiplied by total gene expression ×10000), followed by

identification of 2000 highly variable genes using FindVariableFeatures, data

centering using ScaleData, and finally data integration using FindIntegrationAnchors

(RunHarmony) with 2000 anchors to eliminate batch effects. A total of 42,385 cells

were obtained for subsequent analysis[20].

2.2 Dimensionality Reduction and Data Clustering

To visualize high-dimensional gene expression data effectively[21,22], we

implemented a sequential dimensionality reduction and clustering protocol. We first

performed principal component analysis on highly variable genes using RunPCA,

followed by data integration with RunHarmony using 2000 anchor points. We then

applied FindNeighbors and FindClusters functions for clustering analysis. Finally, we

employed RunUMAP (dims=1:14, resolution=0.5) for dimensionality reduction

visualization to display cell clustering results.

2.3 DEG, Marker Gene and Active Subgroups Identification

To characterize each cluster, a comprehensive manual annotation was performed.

Using the Seurat package's FindAllMarkers function, we identified differentially

expressed genes (DEGs) with |log2FoldChange|>1 and adjusted P-value<0.05, and

determined cell type-specific marker genes. We employed the AUCell R package to

calculate pathway activity scores in single-cell data[23]. After extracting the RNA

expression matrix from our Seurat object with GetAssayData, we ran AUCell_run

with predefined gene sets to quantify pathway enrichment per cell. For pathways of

interest, we extracted area under the curve (AUC) values and incorporated them into

the Seurat metadata.

2.4 GO and KEGG Pathway Enrichment Analysis

We conducted GO functional enrichment and KEGG pathway enrichment analyses on

differentially expressed genes using the clusterProfiler package (v3.14.3) and

visualized results with the ggplot2 package[24,25].

2.5 Pseudo-time Analysis

To analyze the continuous dynamic changes in cell types and associated gene

expression, we used Monocle 3 to reconstruct cellular developmental trajectories from

single-cell data[26]. We extracted six macrophage subtypes (Tnf+ inflammatory,

anti-inflammatory, monocyte-derived, Il1b+ inflammatory, tissue-repair, and

monocyte-derived macrophages) to create a cell_data_set object. Following PCA

preprocessing, we removed batch effects with the align_cds function using orig.ident

as the alignment group. We performed dimensionality reduction via

reduce_dimension and applied low-resolution clustering to ensure distinct trajectories

per cluster. We constructed the cellular trajectory network using learn_graph and

employed a custom function to automatically identify monocyte-derived macrophages

as developmental origin points. We then ordered cells along the developmental

trajectory with order_cells and visualized both pseudo-time progression and

expression patterns of the top five genes using plot_cells and

plot_genes_in_pseudotime functions.

2.6 Cell-Cell Communication Analysis

Using the CellChat package (v1.0.0) to analyze intercellular communication networks,

cell communication relationships are constructed based on expression data of known

ligand-receptor pairs to infer interaction patterns and their abundance between

different cell types[27].

2.7 C57BL/6 Mouse LA Modeling and Transcriptome Sequencing

Mice: Female C57BL/6J mice (22-24g, 4-6 weeks old) from Beijing Vital River

Laboratory Animal Technology Co., Ltd. were housed in SPF-grade facilities at

Kunming Medical University's School of Basic Medicine with 12-hour light/dark

cycles at 21.5°C, free food and water access, and daily bedding changes. The study

was approved by Anning First People's Hospital Medical Ethics Committee (Approval

No.2024-046-01). Bacteria propagation and infection: *Borrelia burgdorferi* strain B31

(ATCC-35210) was cultured in complete BSK-II medium at 37 °C with 5 % CO_2 and

100 % humidity. Experimental mice received bilateral footpad injections of spirochete

suspension ($1 \times 10^{\hat{} }6/50\mu l$ per side), while controls received equivalent PBS volumes.

Ankle joint swelling was measured biweekly using calipers, with X-ray examinations

performed on randomly selected mice. Sample Collection and Transcriptome Analysis:

On day 16 post-surgery, we sampled 6 mice per group. Half the samples underwent

histopathological examination with 4 μm paraffin sections for HE staining. The

remaining samples (approximately 80 mg of ankle joint tissue) were sent to LC Bio

Technology CO., Ltd. (Hangzhou, China) for transcriptome sequencing using the

Illumina Novaseq 6000 platform with paired-end sequencing (PE150 mode).

Bioinformatics analysis was conducted on the standardized clean data.

2.8 Transcriptome Data Integration

2.8 Transcriptome Data Integration

Based on our research focus on the role of macrophages in LA synovial tissue, we

obtained transcriptome sequencing dataset (GSE125503, platform GPL17021) of

Bb-infected macrophages from the Gene Expression Omnibus (GEO) database. This

dataset includes 14 samples derived from bone marrow-derived macrophages

(BMDMs) of C57BL/6 mice, comprising 4 control samples and 10 Bb-infected

samples. Subsequently, we integrated our team's Bulk-transcriptome data with the

GSE125503 Bulk-transcriptome data. We merged different datasets by columns using

the cbind function and obtained common genes through Reduce(intersect, geneList).

Batch effects were then corrected using the ComBat function. Following this, we

established criteria for identifying DEGs using the R package DESeq2. Specifically,

we applied a threshold of $|\log_2$ FoldChange$|> 1$ and adjusted P-value <0.05 to select

DEGs for subsequent analysis[28,29].

2.9 Least Absolute Shrinkage and Selection Operator (LASSO) Regression

Analysis

We identified intersection genes between macrophage-related single-cell sequencing

DEGs and our merged transcriptome dataset DEGs. Using these common differential

genes, we constructed and validated a model using the Least Absolute Shrinkage and

Selection Operator (LASSO) method[30].

2.10 Quantitative Reverse Transcription PCR (qRT-PCR)

Cell Culture and Treatment: RAW264.7 cells were purchased from Shanghai

Zhongqiao Xinzhou Biotechnology Co., Ltd. The cells were infected with Bb (MOI=1)

for 6 hours, or treated with vehicle control (PBS). Samples were stored at -80°C until

analysis. RNA Extraction and qRT-PCR Analysis: Total RNA was extracted using

MolPure$^®$ Cell/Tissue Total RNA Kit (#19221ES50, ShangHai Yeasen Biotechnology)

and reverse transcribed into complementary DNA using Hifair$^®$III 1st Strand cDNA

Synthesis SuperMix for qPCR (#11141ES60, ShangHai Yeasen Biotechnology).

qRT-PCR was performed on the Gentier 96R Real-Time PCR System (Tian Long)

using Hieff unicon universal blue Qpcr SYBR Master mix (#11184ES08, ShangHai

Yeasen Biotechnology) under the following conditions: 95°C for 2 minutes, followed

by 40 cycles of 95°C for 1 second and 60°C for 30 seconds. Expression of target

genes was normalized to Gapdh using the $2^{(-\Delta\Delta CT)}$ method. The primers were

synthesized by Tsingke Biotechnology (Beijing). The primer sequences (5'-3') used to

amplify SIRPB1C were forward, TCAGCAGACAGTGGCCTTTA and reverse,

CTGAGAGCGGACATCCCTAG.   The primers used to amplify FABP5 were

forward,   CGGGAAGGAGAGCACGATAA   and   reverse,

TTGTTGCATTTGACCGCTCA. The primers used to amplify MMP14 were forward,

AGGCCAATGTTCGGAGGAAG and reverse, GTGGCACTCTCCCATACTCG. The

primers used to amplify EGR1 were forward, AGTGATGAACGCAAGAGGCA and

reverse TAGCCACTGGGGATGGGTAA. The primers used to amplify GAPDH were

207 forward, TGTTGCCATCAATGACCCCT and reverse

208 TCGCCCCACTTGATTTTGGA.

2.11 Construction and Testing of LA Diagnostic Nomogram Model

210 Using the "pROC" and "rms" R packages[31,32], we developed a nomogram for LA

211 diagnosis. Risk scores were calculated based on the expression levels of each

212 candidate optimal feature gene, with the total risk score defined as the sum of all

213 individual gene risk scores. The diagnostic value of this nomogram for LA was

214 evaluated through decision trees, calibration curves, and ROC curves.

2.12 Statistical Analysis

216 Analyses and visualizations were performed using R (version 4.2.1) and GraphPad

217 Prism (version 9.0.2). Data are presented as mean ± SEM. We compared groups using

218 two-tailed Student's t-test or Mann-Whitney U test, with significance defined as $P <$

219 0.05 (*), $P < 0.01$ (**), and ns for not significant.

3. Results

3.1 Single-cell Transcriptome of LA

We analyzed 5 samples from the GSE233850 dataset, including ankle joint samples

from 1 uninfected normal C57 mouse and 4 paired Bb-infected C57 mice. After

removing low-quality data, 42,385 cells were retained for analysis. Following

normalization and PCA, we retained the first 14 dimensions and visualized clustering

using UMAP, dividing cells into 20 clusters (Figure 2A). SingleR package analysis

identified 9 distinct cell types (Figure 2B). Figures 2C and 2D showed the single-cell

transcriptome atlas of normal and LA samples and across samples, respectively. The

cell population comprised: Macrophages/Monocytes (16,617 cells, 39.2%; high

expression of Csf1r, C1qa, Adgre1, Syngr1), Neutrophils (4,068 cells, 9.6%; S100a9,

S100a8, Csf3r), T/NK Cells (2,472 cells, 5.8%; Ccl5, Nkg7, Cd3d, Cd3e), B Cells

(7,582 cells, 17.9%; Cd79a, Cd79b, Ms4a1, Ighm), DC Cells (2,492 cells, 5.9%;

H2-Eb1, Klrd1, Slamf7), Endothelial Cells (608 cells, 1.43%; Cldn5, Pecam1, Cdh5),

Fibroblasts (5,999 cells, 14.2%; Col1a1, Col3a1, Dcn), Synoviocytes (475 cells, 1.1%;

Prg4, Htra4, Anxa8), and Cycling Cells (2,072 cells, 4.9%; Mki67, Top2a, Pclaf) as

shown in Figures 2E-2G. Figure 2H showed their proportions between groups, with

the LA model exhibiting higher percentages of Fibroblasts, Cycling cells, and

Endothelial cells, while Macrophages/Monocytes, B cells, T/NK cells, Neutrophils,

DC cells, and Synoviocytes showed relatively lower proportions.

3.2 Single-Cell GSVA Pathway Analysis of KEGG Metabolic Signatures

Figure 3 showed GSVA enrichment analysis using the AUCell algorithm, featuring

differential gene expression across cell subpopulations (Figure 3A). We analyzed 20

KEGG signaling pathway activities within various cellular subsets (Figure 3B), with

particular focus on glycolysis/gluconeogenesis and oxidative phosphorylation

pathways visualized through violin plots (Figure 3C-D), UMAP projections (Figure

3E, 3G), and box plots (Figure 3F, 3H). Results showed increased

glycolytic/gluconeogenic activity and decreased oxidative phosphorylation in

Macrophages/Monocytes following Bb infection, indicating a metabolic shift from

aerobic to anaerobic metabolism and highlighting metabolic reprogramming in host

cells during pathogenic challenge.

3.3 Single-cell Transcriptome Landscape

Due to Macrophages/Monocytes' critical immunoregulatory role in LA, we analyzed

16,617 of these cells from all samples. Figure 4 showed scRNA-seq analysis

identified 7 distinct subtypes: Anti-inflammatory macrophages (3,331 cells; high

expression of Olfml3, Sparc, Vsig4), TNF+ Inflammatory macrophages (3,186 cells;

Tnf, Cxcl2, Ccl7), Tissue-Repair macrophages (2,052 cells; Vegfa, Arg1, Adam8),

Mono-derived macrophages (2,030 cells; Mrc1, Gas6), Il1b+ Inflammatory

macrophages (1,924 cells; Il1b, Clec4n, Ptgs2), Ly6c+ inflammatory Mono (1,034

cells; Ly6c2, Hp, Thbs1), and Progenitor cells/activated Mono (1,033 cells; Plac8,

Slfn1). Comparison between normal and LA model groups (Figures 4C-4G) revealed

significant compositional differences: the LA model showed decreased proportions of

Anti-inflammatory macrophages, Mono-derived macrophages, and Ly6c+

inflammatory monocytes, while TNF+ Inflammatory macrophages, Il1b+

Inflammatory macrophages, Tissue-repair macrophages, and Progenitor

cells/activated monocytes increased (Figure 4H).

3.4 Macrophage-Related Enrichment Analysis

To elucidate biological functions of Macrophage/Monocyte marker genes, we

performed GO and KEGG enrichment analyses. GO functional enrichment of

differentially expressed genes (Figure 5A-B) revealed: Biological Processes (BP)

primarily related to endosomal transport, precursor metabolite and energy generation,

hydrolase activity upregulation, cellular respiration, vesicle organization, and

oxidative phosphorylation. Cellular Components (CC) mainly associated with

mitochondrial protein complexes, vacuolar membranes, early endosomes, and

lysosomal/endosomal structures. Molecular Functions (MF) included phospholipid

binding, GTPase regulatory activity, nucleoside-triphosphatase regulation, active

transmembrane transport, phosphatidylinositol binding, and GTPase interactions.

KEGG pathway analysis (Figure 5C) showed signature genes enriched in

inflammatory response pathways linked to energy metabolism regulation, including

bacterial epithelial invasion, osteoclast differentiation, Toll-like receptor signaling,

rheumatoid arthritis, carbon metabolism, HIF-1 signaling, glycolysis/gluconeogenesis,

arginine/proline metabolism, and adherens junctions. Figure 5D shows differential

analysis within Macrophages/Monocytes, while Figure 5E presents a diagonal

volcano plot. GO enrichment bar plots for upregulated and downregulated genes

(Figure 5F) revealed distinct patterns: upregulated genes showed significant

enrichment in pathways governing precursor metabolite generation, nucleotide

metabolism, phospholipid processes, and lipid/small molecule metabolic regulation.

Downregulated genes showed predominant enrichment in oxidative phosphorylation,

aerobic and cellular respiration, energy derivation through organic compound

oxidation, and mitochondrial respiratory chain/NADH dehydrogenase complex

assembly.

3.5 Pseudo-time Analysis

Using Monocle 3 R, we analyzed macrophage differentiation trajectories. Figure 6A

showed dynamic RNA expression changes across five macrophage subgroups, while

Figure 6B illustrateed pseudo-time progression from dark to light-colored cells. Our

analysis identified monocyte-derived macrophages as the primary origin point,

diverging into three distinct developmental paths: toward anti-inflammatory

macrophages, Tnf+ inflammatory macrophages, and Il1b+ inflammatory macrophages.
The trajectory map highlighted macrophage plasticity during disease progression,
with tissue-repair macrophages primarily functioning as an intermediate state
evolving toward Tnf+ inflammatory phenotypes rather than maintaining reparative
functions (Figure 6C-6D). Figure 6E showed pseudo-time dynamics of C1qa, C1qb,
and C1qc genes during LA-related macrophage development. These genes exhibit
similar expression patterns: high expression in early pseudo-time (0-5), particularly in
mono-derived, anti-inflammatory, and TNF+ inflammatory macrophages, followed by
gradual decline in mid-to-late pseudo-time (5-10), with tissue-repair macrophages
showing the most pronounced decrease and Il1b+ inflammatory macrophages
maintaining intermediate expression.

3.6 Cell-to-Cell Communication Between Macrophages and Other Cells

To investigate macrophage influence on other cells during LA pathogenesis, we
analyzed cell-to-cell communication through ligand-receptor interactions using
scRNA-seq data (Figure 7). Interactions between macrophages and other cells were
stronger in LA tissue than in normal tissue, in both number and interaction strength
(Figure 7A-7C). CXCL, VEGF, VISFATIN, MIF, GALECTIN, COMPLEMENT, and
OSM showed enhanced receptor binding in LA, while PTN, CHEMERIN, TNF, IGF,
GAS, SEMA3, BAFF, TGF-β, PDGF, CSF, EGF, and CCL exhibited decreased
binding (Figure 7D). TGF-β signaling pathway-related genes showed decreased
expression between various cell types in LA compared to normal tissues (Figure 7E).
In the TGF-β signaling network, IL1b+ inflammatory macrophages were the primary
signal senders, with tissue-repair and Th1+ inflammatory macrophages also showing
strong signaling capacity. Anti-inflammatory macrophages functioned as the
predominant signal receivers while simultaneously serving as the most significant
network mediators (Figure 7F). TGF-β signaling pathways were widely present across
cell types, with Tgfb1, Tglbr1, and TgIbr2 showing higher expression in normal
groups (Figure 7G). Tissue-repair macrophages exhibited high intercellular
communication volume in LA (Figure 7H-7I), with the Tgfb1-(Tglbr1+TgIbr2) axis
contributing most significantly to TGF-β signaling (Figure 7J). The crosstalk of this

ligand-receptor pair between 12 major cell types confirmed decreased TGF-β

signaling pathway-related gene expression in LA compared to normal tissues (Figures

7K,7L).

3.7 Bulk-transcriptome sequencing of C57BL/6 Mouse Ankle Joints

To examine macrophage specificity in LA, we merged transcriptome data from

C57BL/6 mouse ankle joints with public data from bone marrow-derived

macrophages (GSE125503, platform GPL17021) and conducted differential gene

analysis. Figure 8A-8B show the heatmap and volcano plot of differential genes. We

identified 397 DEGs and intersected them with 222 macrophage-related DEGs from

scRNA-seq, yielding 15 overlapping genes (Figure 8C): SIRPB1C, CD14, FABP5,

IL2RG, MCU, MMP14, MXRA8, EGR1, MSR1, CLEC4D, PIRB, IL1B, H2-AA,

GYPC, and ITGA6. After normalizing the expression matrix, LASSO regression

analysis identified four candidate optimal feature genes: SIRPB1C, FABP5, MMP14,

and EGR1 (Figure 8D-8E), with their expression profiles shown in Figures 8F-G-H-I.

3.8 Construction and testing of a signature gene-based nomogram for predicting

LA

Using the "RMS" R package, we constructed a diagnostic nomogram for LA based on

the four characteristic genes (Figure 8J) and evaluated its predictive capability. The

calibration curve showed minimal difference between actual and predicted LA risk,

indicating high model accuracy (Figure 8K). Decision curve analysis (DCA)

demonstrated potential patient benefit from this nomogram (Figure 8L), while ROC

curve analysis further confirmed the model's accuracy (Figure 8M).

3.9 Validation of Candidate Optimal Feature Genes

In our LA mouse model, C57BL/6 mice developed hind leg ankle joint swelling 2

days after Bb injection, peaking at approximately 2 weeks before gradually subsiding

after 4-5 weeks. H&E staining showed inflammatory cell infiltration and substantial

edema in joint synovial and surrounding soft tissues, while bone and cartilage

remained intact (Figure 9A). By day 16, experimental mice displayed significant joint

swelling and mild stiffness compared to PBS controls, without skin ulceration or

blackening (Figure 9B). X-rays revealed soft tissue edema (Figure 9C). RT-qPCR

analysis confirmed significantly increased expression of SIRPB1C, FABP5, MMP14,

and EGR1 in Bb-infected samples from both RAW264.7 cells and joints of C57BL/6

mice compared to controls.

4. Discussion

LA is one of the most common late-stage manifestations of Lyme disease. Without

timely diagnosis and treatment, it can lead to persistent arthritis and long-term

functional impairment [33]. While antibiotics are generally effective, 10-20% of

patients develop antibiotic-refractory LA (ARLA) [34,35]. Single-cell RNA

sequencing (scRNA-seq) now enables analysis of disease pathogenesis at

unprecedented resolution [36,37], revealing cellular composition and transcriptomic

changes associated with LA. Macrophages, as key effector cells of the innate immune

system, play a central role in the pathogenesis of LA [38], regulating inflammatory

responses and influencing disease progression through cytokine and chemokine

secretion. C57BL/6 mice provide an ideal model for studying transcriptional and

cellular changes during infection due to their uniform genetic background. In this

model, arthritis symptoms resolve spontaneously despite persistent bacteria,

effectively simulating host immune adaptation to chronic infection[39-42]. As the

disease progresses, bacterial numbers gradually decrease while the host immune

response changes, achieving long-term control of infection[18]. This self-healing

process provides valuable insights into the self-limiting nature of LA. This study

analyzed macrophage differential gene expression in LA using GSE233850

scRNA-seq data from C57BL/6 mouse models. By integrating bulk transcriptome

datasets through joint analysis and machine learning, we identified and validated core

macrophage-related genes in LA. The synovial microenvironment (SME), comprising

immune cells, synovial cells, and stromal cells[43], critically influences LA

progression and prognosis[44]. Antibiotic-refractory LA patients often experience

poor outcomes[5], with some showing persistent macrophage activation despite Bb

clearance[45]. Since synovial macrophages can either promote or inhibit

inflammation through M1/M2 polarization, understanding these mechanisms is

essential for improving treatment[46]. Single-cell RNA sequencing has become vital

for characterizing synovial immune cells in inflammatory arthritis[47,48]. Therefore,

systematically studying macrophage functional states and regulatory networks in LA,

and developing macrophage-based prognostic models, has significant clinical

relevance.

Single-cell transcriptome analysis of ankle joints from uninfected and Bb-infected

C57BL/6 mice identified 9 cell types, with significant shifts during LA development:

Fibroblasts increased while Macrophages/Monocytes decreased. We found seven

Macrophage/Monocyte subsets with notable compositional differences between

normal and LA models. LA samples showed decreased anti-inflammatory

macrophages, monocyte-derived macrophages, and Ly6c+ inflammatory monocytes,

with increased TNF+ and Il1b+ inflammatory macrophages, tissue-repair

macrophages, and progenitor cells. These subsets serve distinct functions:

anti-inflammatory macrophages maintain homeostasis by suppressing

inflammation[49-51]; TNF+ inflammatory macrophages mediate inflammatory and

chemokine signaling[52-54]; tissue-repair macrophages orchestrate inflammation

regulation and tissue remodeling[55,56]. Monocyte-derived macrophages play a crucial

regulatory role in various diseases, particularly in liver regeneration, where they

facilitate tissue repair through migration, proliferation, and apoptosis[57].In

neuroinflammation, such as stroke, these cells migrate to the brain via CCR4 and

CCR5 receptors, emerging as potential therapeutic targets[58]; IL-1β+ inflammatory

macrophages initiate inflammatory cascade [59-61]; Ly6c+ inflammatory monocytes

drive inflammation and angiogenesis[62,63]; and monocytes/progenitor cells respond

to systemic inflammation in various diseases[64-66]. Despite the overall reduction in

Macrophages/Monocytes under pathological conditions, IL1b+ inflammatory and

tissue-regenerative macrophages significantly increased while anti-inflammatory

macrophages declined, indicating functional reprogramming within the macrophage

spectrum. Despite overall Macrophage/Monocyte reduction, IL1b+ inflammatory and

tissue-regenerative macrophages increased while anti-inflammatory macrophages

declined, indicating functional reprogramming. Pseudo-temporal trajectory analysis

revealed three main differentiation pathways from monocyte-derived macrophages:

anti-inflammatory, TNF+ inflammatory, and IL1b+ inflammatory lineages.

Cell-cell communication analysis identified significant changes in Transforming

growth factor-β (TGF-β) -receptor interactions during pathological states, suggesting

potential therapeutic targets for LA. Heat map analyses showed decreased expression

of TGF-β signaling pathway-related genes between various cell types in LA compared

to normal tissues. We identified the critical ligand-receptor pair

Tgfb1-(Tglbr1+TgIbr2) within TGF-β signaling networks showing enhanced

intercellular communication across diverse cell types in LA. TGF-β, a cytokine that

regulates inflammation and promotes fibrosis[67], plays a dual role in Lyme disease.

TGF-β1 levels are higher in early/non-chronic patients compared to chronic patients,

who show impaired TGF-β1 synthesis when stimulated by spirochetes[68]. This

difference may explain the mechanism of disease progression to chronicity: elevated

TGF-β1 levels represent an effective anti-inflammatory immune response, promoting

pathogen clearance; whereas the reduction of TGF-β1 in chronic patients leads to

insufficient persistent anti-inflammatory response[69]. Patients with

antibiotic-refractory LA show excessive inflammation and fibrosis in synovial

tissue[70]. TGF-β affects vascular reconstruction and may promote vascular

occlusion[71], suggesting occlusive microvascular lesions in antibiotic-refractory LA

patients may be linked to TGF-β activity.

GSVA and KEGG pathway analyses showed that Bb infection shifted macrophage

metabolism from oxidative phosphorylation to glycolysis. Previous studies confirm

metabolic reconfiguration's critical role in inflammation: lipopolysaccharide-induced

metabolic transition is essential for inflammatory response[72]; while increased

glycolysis enhances macrophage inflammatory functions in ankylosing spondylitis

and myocardial infarction[73] [74]. These findings support our conclusions about

metabolic reprogramming of Monocytes/Macrophages in LA.

We developed a LA model in C57BL/6 mice and analyzed transcriptome data from

infected ankle joints combined with public dataset GSE125503 to investigate

macrophage roles in LA. Through differential gene expression and LASSO regression,

we identified four characteristic genes: SIRPB1C, FABP5, MMP14, and EGR1.

(Signal Regulatory Protein Beta 1C (SIRPB1C) belongs to the signal regulatory protein family involved in inflammation and immunity. SIRPB1C is a key member of the signal regulatory protein beta family. Studies show SIRPB1 knockout reduces inflammatory cytokines (IL1RA, CCL2, IL-8) [75], while another family member, SIRPα, suppresses phagocytosis and immune cell migration, with SIRPα-targeted antibodies reducing inflammation[76,77]. Fatty acid-binding protein 5 (FABP5), a lipid chaperone involved in inflammatory pain modulation, is present in human synovium[78,79]. Its inhibition significantly reduces cytokine and chemokine secretion, suggesting potential impact on osteoarthritis-related inflammatory pain[80]. FABP5 is upregulated in intestinal macrophages during inflammatory bowel diseases, and its inhibition prevents inflammatory M1 macrophage differentiation, reducing colitis in mice[81].

MMP14 (membrane type I matrix metalloproteinase or MT1-MMP) is a collagenase that degrades extracellular matrix, particularly type II collagen in cartilage[82]. In rheumatoid arthritis, MMP14 on synoviocytes and macrophages promotes cartilage degradation[83]. Studies show MMP14 inhibition preserves joints: the inhibitor FR255031 reduces cartilage degradation in arthritis models, while selective MT1-MMP inhibition protects joints and enhances anti-tumor necrosis factor therapy efficacy, suggesting potential combination treatments[84,85].

Early Growth Response 1 (EGR1), a transcription factor, plays a crucial role in host-pathogen interactions[86]. While no direct EGR1-LA link has been previously documented, this factor is rapidly induced by pathogens, cytokines, and stressors, and regulates inflammation, angiogenesis, and matrix remodeling[87,88]. EGR1 promotes cartilage angiogenesis through the NT-1/DCC-VEGF pathway, accelerating cartilage deterioration[89]. As an inflammatory mediator, it enhances pro-inflammatory IL-1 effects while suppressing anti-inflammatory PPARγ[90]. EGR1 also regulates extracellular matrix genes, affecting cartilage integrity[91]. Our research suggests EGR1 likely plays an essential role in LA inflammatory regulation.

Although our research provides valuable insights, it has several limitations that need to be addressed in future studies. First, the sample size of the integrated bulk RNA

sequencing data is relatively limited. Second, regarding the exploration of molecular mechanisms: while we validated the transcriptional changes of core differential genes through in vitro and vivo experiments, we have not fully understood the functional roles of these genes in LA pathogenesis. Future research should aim to validate these findings in larger cohorts and employ functional experiments to confirm the roles of these cells, genes, and pathways in disease progression.

5. Conclusion

Integrative single-cell and bulk RNA sequencing revealed seven macrophage subtypes in LA, a glycolytic shift, and disrupted TGF-β signaling. Pseudo-time analysis traced monocyte-derived macrophages toward distinct anti-inflammatory or TNF+/IL1β+ inflammatory fates, underscoring functional plasticity. Four candidate optimal feature genes-SIRPB1C, FABP5, MMP14 and EGR1-accurately discriminated diseased joints, forming a nomogram with high predictive value. These findings deepen understanding of macrophage heterogeneity in LA and highlight metabolic and signaling nodes that may guide biomarker-driven diagnosis and macrophage-targeted therapy.

Author Affiliations

[1]Department of Pain Medicine, The Affiliated Anning First People's Hospital of Kunming University of Science and Technology, Kunming, Yunnan Province, 650302, China.

[2]School of Basic Medical Sciences, Kunming University of Science and Technology, Kunming, Yunnan Province, 650500, China.

Acknowledgement We thank Professor Yi☐Qun Kuang for providing the Biosafety Laboratory at the Research Center for Clinical Medicine, First Affiliated Hospital of Kunming Medical University, and acknowledge the GEO databases for publicly available datasets.

Contributors Yan Dong and Yantong Chen performed bioinformatics analyses and drafted the manuscript; Yanshuang Luo, Meng Liu, Yingchen Zhou , Yutian Yang,

Xuesong Chen, Xiaorong Liu, Fusong Yang, Zhi Liang, Liye Zi, Hongbing Gao, Zhichao Cao, and Fucun Gao conducted experiments, interpreted data; Guozhong Zhou and Chao Song designed the study and supervised the project. All authors reviewed and approved the final manuscript.

Funding This study was supported by the Yunnan Fundamental Research Projects (grant NO. 202301BE070001-036) and internal projects of the First People's Hospital of Anning, affiliated with Kunming University of Science and Technology (grant NO. 2024AYY001 and 2024AYY005).

Competing interests None declared.

Ethical approval Animal experiments were conducted in accordance with the Declaration of Helsinki principles and approved by the Kunming Medical University Experimental Animal Ethics Committee (Approval No.: KMMU20211590).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Publicly available datasets that support the findings of this study are openly available in the Gene Expression Omnibus (GEO) database (https://www.ncbi.nlm.nih.gov/geo/info/datasets.html) ; GSE233850 and GSE125503.

Code availability Additional information for data re-analysis is available from the lead contact upon request.

Publisher's note Claims expressed in this article are solely those of the authors and do not necessarily represent affiliated organizations or the publisher. Products evaluated or claims made by manufacturers are not guaranteed or endorsed by the publisher.

Figure legends

Figure 1. Comprehensive workflow diagram of this investigation.

Figure 2. Single-cell atlas of normal and Lyme arthritis C57BL/6 mouse samples. (A) UMAP clustering of single cells into 20 clusters; (B) Nine cell types identified by

marker gene expression; (C) UMAP comparison between normal and Lyme arthritis

model groups; (D) UMAP distribution across 5 samples; (E) Bubble plot of marker

gene expression in 9 cell types. Bubble plots display marker gene expression in cell

subpopulations. Bubble color intensity (red) corresponds to expression level, while size reflects

the number of expressing cells; (F) Heatmap of differential gene expression among cell

types; (G) Histogram shows the cumulative distribution of selected marker genes of

cell types; (H) Cell type proportion comparison between normal and Lyme arthritis

groups.

Figure 3. GSVA Enrichment Analysis Between Specific Subgroup Cells and Groups.

(A) Volcano plot of differential gene expression across cellular subpopulations. Red

dots indicate significantly upregulated cluster-specific genes; blue dots show

significantly downregulated genes; (B) GSVA enrichment analysis among 9 cell types;

(C) Violin plot of AUC scores for glycolysis and gluconeogenesis pathway-related

gene sets; (D) Violin plot of AUC scores for oxidative phosphorylation

pathway-related gene sets; (E) UMAP colorogram showing glycolysis and

gluconeogenesis pathway activity, with brighter colors indicating higher activity; (F)

Box plot of AUC scores for glycolysis and gluconeogenesis pathway-related gene sets;

(G) UMAP colorogram showing oxidative phosphorylation pathway activity, with

brighter colors indicating higher activity; (H) Box plot of AUC scores for oxidative

phosphorylation pathway-related gene sets.

Figure 4. Single-cell transcriptome atlas of Macrophages/Monocytes. (A) UMAP plot

showing 7 distinct Macrophage/Monocyte clusters; (B) UMAP plot with 7

Macrophage/Monocyte subtypes identified by signature molecular markers; (C)

UMAP comparison between normal and Lyme arthritis model groups; (D) UMAP

distribution across 5 samples; (E) Bubble plot showing differential expression patterns

of signature genes across 7 cell subgroups; (F) Heatmap of differential gene

expression among the 7 cell subgroups; (G) UMAP visualization of marker gene

expression patterns for each cell subgroup; (H) Comparative proportions of cell

subgroups between normal and Lyme arthritis groups.

Figure 5. Enrichment Analysis Between Macrophage/Monocyte Subgroups. (A-B) GO functional enrichment analysis of differentially expressed genes in Macrophages/Monocytes, categorized by biological processes (BP), molecular functions (MF), and cellular components (CC); (C) KEGG pathway enrichment analysis showing functional distribution of differentially expressed genes across macrophage/monocyte populations; (D) Scatter plot comparing differential expression between Macrophage/Monocyte groups; (E) Diagonal volcano plots illustrating differential gene expression between Macrophage/Monocyte subpopulations; (F) Bar plots showing GO enrichment analysis of upregulated and downregulated genes across Macrophage/Monocyte subpopulations.

Figure 6. Trajectory analysis of macrophage subtypes. (A) UMAP plot showing 5 macrophage subtypes identified by signature molecular markers; (B) Macrophages' differentiation trajectories on pseudo-time, with darker purple cells transitioning to lighter orange cells, indicating progression from earlier to later differentiation stages; (C-D) Trajectory analysis revealing dynamic processes across macrophage subpopulations, with RNA expression profiles changing continuously along anatomical axes; numbered circles represent differentiation sequence, where gray-filled circles indicate different endpoints, white-filled circles mark branch points where cells can follow multiple fates, and black-filled numbered circles show different differentiation nodes; (E) Absolute expression of top 3 genes in pseudo-time, where each point represents a cell, different colors represent different clusters, and the y-axis shows gene expression levels.

Figure 7. Cellular communication analysis. (A-B) Network diagrams showing cell-cell interaction number and strength in normal and Lyme arthritis model groups respectively; (C) Comparative analysis of cellular communication number and strength between groups; (D) Total pathway analysis comparing Lyme arthritis model and normal groups; (E) Heatmap of ligand-receptor interaction patterns in TGF-β

signaling pathway across cell types in both groups; (F) TGF-β signal sending-receiving intensity map in Lyme arthritis group; (G) Expression profiles of representative TGF-β signaling pathway genes across cell types between groups; (H) Interaction network between tissue-repair macrophage and other cell types in Lyme arthritis group; (I) Interaction dynamics among different cell types; (J) Contribution of each TGF-β ligand-receptor pair; (K-L) Cell-cell communication mediated by Tgfb1-(Tglbr1+Tglbr2) ligand-receptor pair in normal and Lyme arthritis model groups.

Figure 8. Machine learning prediction of Lyme arthritis candidate optimal feature genes and nomogram for disease occurrence prediction. (A) Heatmap of differentially expressed genes in merged transcriptome: blue means low expression; red means high expression; (B) Volcano plot of differentially expressed genes in merged transcriptome: green means low expression; red means high expression; (C) Venn diagram identified the 15 overlapping genes between merged transcriptome DEGs and Macrophage/Monocyte-related genes from single-cell data; (D) LASSO regression model construction using merged transcriptome samples as training set; (E) Model parameter optimization cross-validation (left line indicating optimal λ value), determined the four candidate optimal feature genes (SIRPB1C, FABP5, MMP14 and EGR1); (F-J-H-I) Expression levels of the 4 candidate optimal feature genes in merged transcriptome samples; (J) Nomogram for Lyme arthritis occurrence prediction. Line segments represent the predicted contribution of genes to outcome events, with the total score signifying the cumulative sum of all corresponding individual scores across variable values; (K) Calibration curve evaluating nomogram model predictive ability; (L) DCA curve assessing nomogram model value; (M) ROC curve evaluating nomogram model performance in merged transcriptome samples. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 9. Validation of 4 candidate optimal feature genes. (A-B-C) Edema observation, X-ray examination, and H&E staining of joint synovium in Lyme arthritis mouse model versus control, scale bar = 750 μm; (D-K) RT-qPCR validation

625 of these genes in *Borrelia burgdorferi*–infected RAW264.7 cells and C57BL/6 mice
626 versus controls. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.
627

References:
1.   Lochhead, R.B.; Strle, K.; Arvikar, S.L.; Weis, J.J.; Steere, A.C. Lyme arthritis: linking
infection, inflammation and autoimmunity. *Nat. Rev. Rheumatol.* **2021**, 17, 449-461,
doi:10.1038/s41584-021-00648-5.
2.   Kanjana, K.; Strle, K.; Lochhead, R.B.; Pianta, A.; Mateyka, L.M.; Wang, Q.; Arvikar, S.L.;
Kling, D.E.; Deangelo, C.A.; Curham, L. et al. Autoimmunity to synovial extracellular matrix
proteins in patients with   postinfectious Lyme arthritis. *J. Clin. Invest.* **2023**, 133,
doi:10.1172/JCI161170.
3.   Nadelman, R.B.; Hanincová, K.; Mukherjee, P.; Liveris, D.; Nowakowski, J.; McKenna, D.;
Brisson, D.; Cooper, D.; Bittker, S.; Madison, G. et al. Differentiation of reinfection from
relapse in recurrent Lyme disease. *N. Engl. J. Med.* **2012**, 367, 1883-1890,
doi:10.1056/NEJMoa1114362.
4.   Coburn, J.; Garcia, B.; Hu, L.T.; Jewett, M.W.; Kraiczy, P.; Norris, S.J.; Skare, J. Lyme Disease
Pathogenesis. *Curr. Issues Mol. Biol.* **2021**, 42, 473-518, doi:10.21775/cimb.042.473.
5.   Brouwer, M.; van de Schoor, F.R.; Vrijmoeth, H.D.; Netea, M.G.; Joosten, L. A joint effort:
The interplay between the innate and the adaptive immune system   in Lyme arthritis. *Immunol.*
Rev. **2020**, 294, 63-79, doi:10.1111/imr.12837.
6.   Steere, A.C. Posttreatment Lyme disease syndromes: distinct pathogenesis caused by
maladaptive   host responses. *J. Clin. Invest.* **2020**, 130, 2148-2151, doi:10.1172/JCI138062.
7.   Lasky, C.E.; Olson, R.M.; Brown, C.R. Macrophage Polarization during Murine Lyme
Borreliosis. *Infect. Immun.* **2015**, 83, 2627-2635, doi:10.1128/IAI.00369-15.
8.   Shin, J.J.; Strle, K.; Glickstein, L.J.; Luster, A.D.; Steere, A.C. Borrelia burgdorferi stimulation
of chemokine secretion by cells of monocyte   lineage in patients with Lyme arthritis. *Arthritis*
Res. Ther. **2010**, 12, R168, doi:10.1186/ar3128.
9.   Codolo, G.; Bossi, F.; Durigutto, P.; Bella, C.D.; Fischetti, F.; Amedei, A.; Tedesco, F.; D'Elios,
S.; Cimmino, M.; Micheletti, A. et al. Orchestration of inflammation and adaptive immunity in
Borrelia   burgdorferi-induced arthritis by neutrophil-activating protein A. *Arthritis Rheum*
2013, 65, 1232-1242, doi:10.1002/art.37875.
10.   Skovsbo, C.A.; Ørbæk, M.; Renee, P.R.; Oestrup, J.P.; Lebech, A.M.; Kjaer, A.
(64)Cu-DOTATATE Positron Emission Tomography (PET) of Borrelia Burgdorferi   Infection:
In Vivo Imaging of Macrophages in Experimental Model of Lyme   Arthritis. *Diagnostics* **2020**,
10, doi:10.3390/diagnostics10100790.
11.   DuChateau, B.K.; Munson, E.L.; England, D.M.; Lovrich, S.D.; Callister, S.M.; Jensen, J.R.;
Schell, R.F. Macrophages interact with enriched populations of distinct T lymphocyte subsets
for the induction of severe destructive Lyme arthritis. *J. Leukoc. Biol.* **1999**, 65, 162-170,
doi:10.1002/jlb.65.2.162.
12.   Benjamin, S.J.; Hawley, K.L.; Vera-Licona, P.; La Vake, C.J.; Cervantes, J.L.; Ruan, Y.; Radolf,
J.D.; Salazar, J.C. Macrophage mediated recognition and clearance of Borrelia burgdorferi
elicits   MyD88-dependent and -independent phagosomal signals that contribute to
phagocytosis   and   inflammation.   *BMC Immunol.*   **2021**,   22,   32,

doi:10.1186/s12865-021-00418-8.

13.   Harjacek, M.; Diaz-Cano, S.; Alman, B.A.; Coburn, J.; Ruthazer, R.; Wolfe, H.; Steere, A.C.
Prominent expression of mRNA for proinflammatory cytokines in synovium in patients with
juvenile rheumatoid arthritis or chronic Lyme arthritis. *J. Rheumatol.* **2000**, 27, 497-503.

14.   Sonderegger, F.L.; Ma, Y.; Maylor-Hagan, H.; Brewster, J.; Huang, X.; Spangrude, G.J.;
Zachary, J.F.; Weis, J.H.; Weis, J.J. Localized production of IL-10 suppresses early
inflammatory cell infiltration and subsequent development of IFN-γ-mediated Lyme arthritis.
J. Immunol. **2012**, 188, 1381-1393, doi:10.4049/jimmunol.1102359.

15.   Liu, W.; Jia, J.; Dai, Y.; Chen, W.; Pei, G.; Yan, Q.; Zhao, Z. Delineating COVID-19
immunological features using single-cell RNA sequencing. *Innovation (Camb)* **2022**, 3, 100289,
doi:10.1016/j.xinn.2022.100289.

16.   Onuora, S. Single-cell RNA sequencing sheds light on cell-type specific gene expression in
immune cells. *Nat. Rev. Rheumatol.* **2022**, 18, 363, doi:10.1038/s41584-022-00802-7.

17.   Su, C.; Xu, Z.; Shan, X.; Cai, B.; Zhao, H.; Zhang, J. Cell-type-specific co-expression inference
from single cell RNA-sequencing data. *Nat. Commun.* **2023**, 14, 4846,
doi:10.1038/s41467-023-40503-7.

18.   Helble, J.D.; Walsh, M.J.; McCarthy, J.E.; Smith, N.P.; Tirard, A.J.; Arnold, B.Y.; Villani, A.C.;
Hu, L.T. Single-cell RNA sequencing of murine ankle joints over time reveals distinct
transcriptional changes following Borrelia burgdorferi infection. *iScience* **2023**, 26, 108217,
doi:10.1016/j.isci.2023.108217.

19.   Butler, A.; Hoffman, P.; Smibert, P.; Papalexi, E.; Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.*
2018, 36, 411-420, doi:10.1038/nbt.4096.

20.   Stuart, T.; Butler, A.; Hoffman, P.; Hafemeister, C.; Papalexi, E.; Mauck, W.R.; Hao, Y.;
Stoeckius, M.; Smibert, P.; Satija, R. Comprehensive Integration of Single-Cell Data. *Cell* **2019**,
177, 1888-1902, doi:10.1016/j.cell.2019.05.031.

21.   Gong, B.; Zhou, Y.; Purdom, E. Cobolt: integrative analysis of multimodal single-cell
sequencing data. *Genome Biol.* **2021**, 22, 351, doi:10.1186/s13059-021-02556-z.

22.   Stuart, T.; Satija, R. Integrative single-cell analysis. *Nat. Rev. Genet.* **2019**, 20, 257-272,
doi:10.1038/s41576-019-0093-7.

23.   Li, C.; Hao, R.; Li, C.; Liu, L.; Ding, Z. Integration of single-cell and bulk RNA sequencing data
using machine learning identifies oxidative stress-related genes LUM and PCOLCE2 as
potential biomarkers for heart failure. *Int. J. Biol. Macromol.* **2025**, 300, 140793,
doi:10.1016/j.ijbiomac.2025.140793.

24.   Wu, T.; Hu, E.; Xu, S.; Chen, M.; Guo, P.; Dai, Z.; Feng, T.; Zhou, L.; Tang, W.; Zhan, L. et al.
clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)*
2021, 2, 100141, doi:10.1016/j.xinn.2021.100141.

25.   Sherman, B.T.; Hao, M.; Qiu, J.; Jiao, X.; Baseler, M.W.; Lane, H.C.; Imamichi, T.; Chang, W.
DAVID: a web server for functional enrichment analysis and functional annotation of gene
lists (2021 update). *Nucleic. Acids. Res.* **2022**, 50, W216-W221, doi:10.1093/nar/gkac194.

26.   Jean-Baptiste, K.; McFaline-Figueroa, J.L.; Alexandre, C.M.; Dorrity, M.W.; Saunders, L.;
Bubb, K.L.; Trapnell, C.; Fields, S.; Queitsch, C.; Cuperus, J.T. Dynamics of Gene Expression
in Single Root Cells of Arabidopsis thaliana. *Plant. Cell.* **2019**, 31, 993-1011,
doi:10.1105/tpc.18.00785.

27.   Jin, S.; Plikus, M.V.; Nie, Q. CellChat for systematic analysis of cell-cell communication from
single-cell transcriptomics. *Nat. Protoc.* **2024**, doi:10.1038/s41596-024-01045-4.

28.   Love, M.I.; Huber, W.; Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **2014**, 15, 550, doi:10.1186/s13059-014-0550-8.

29.   Liu, S.; Wang, Z.; Zhu, R.; Wang, F.; Cheng, Y.; Liu, Y. Three Differential Expression Analysis
Methods for RNA Sequencing: limma, EdgeR, DESeq2. *J. Vis. Exp.* **2021**, doi:10.3791/62528.

30.   Tibshirani, R. The lasso method for variable selection in the Cox model. *Stat. Med.* **1997**, 16,
385-395, doi:10.1002/(sici)1097-0258(19970228)16:4<385::aid-sim380>3.0.co;2-3.

31.   Jaume, S.P.; Skaltsa, K.; Pallarès, N.; Carrasco, J.L. ThresholdROC: Optimum Threshold
Estimation Tools for Continuous Diagnostic Tests in R. *J. Stat. Softw.* **2017**, 82, 0-0,
doi:10.18637/jss.v082.i04.

32.   Yee, J. rlsm: R package for least squares Monte Carlo. *arXiv (Cornell University)* **2018**, 0, 0-0,
doi:null.

33.   Keyßer, G. Lyme disease and Lyme arthritis. *Aktuelle Rheumatol.* **2024**, 0, 0-0,
doi:10.1055/a-2215-0830.

34.   Steere, A.C.; Strle, F.; Wormser, G.P.; Hu, L.T.; Branda, J.A.; Hovius, J.W.; Li, X.; Mead, P.S.
Lyme borreliosis. *Nat. Rev. Dis. Primers* **2016**, 2, 0-0, doi:10.1038/nrdp.2016.90.

35.   Onuora, S. Lyme arthritis: spirochaete remnants could explain antibiotic-refractory Lyme
arthritis. *Nat. Rev. Rheumatol.* **2012**, 8, 440, doi:10.1038/nrrheum.2012.115.

36.   Van de Sande, B.; Lee, J.S.; Mutasa-Gottgens, E.; Naughton, B.; Bacon, W.; Manning, J.; Wang,
Y.; Pollard, J.; Mendez, M.; Hill, J. et al. Applications of single-cell RNA sequencing in drug
discovery and development. *Nat. Rev. Drug Discov.* **2023**, 22, 496-520,
doi:10.1038/s41573-023-00688-4.

37.   Papalexi, E.; Satija, R. Single-cell RNA sequencing to explore immune cell heterogeneity. *Nat.*
Rev. Immunol. **2018**, 18, 35-45, doi:10.1038/nri.2017.76.

38.   Arvikar, S.L.; Crowley, J.T.; Sulka, K.B.; Steere, A.C. Autoimmune Arthritides, Rheumatoid
Arthritis, Psoriatic Arthritis, or Peripheral Spondyloarthritis Following Lyme Disease.
Arthritis Rheumatol. **2017**, 69, 194-202, doi:10.1002/art.39866.

39.   Koloski, C.W.; Hurry, G.; Foley-Eby, A.; Adam, H.; Goldstein, S.; Zvionow, P.; Detmer, S.E.;
Voordouw, M.J. Male C57BL/6J mice have higher presence and abundance of Borrelia
burgdorferi in their ventral skin compared to female mice. *Ticks Tick.-Borne Dis.* **2024**, 15,
102308, doi:10.1016/j.ttbdis.2024.102308.

40.   Hammond, E.M.; Baumgarth, N. CD4 T cell responses in persistent Borrelia burgdorferi
infection. *Curr. Opin. Immunol.* **2022**, 77, 102187, doi:10.1016/j.coi.2022.102187.

41.   Bernard, Q.; Wang, Z.; Di Nardo, A.; Boulanger, N. Interaction of primary mast cells with
Borrelia burgdorferi (sensu stricto): role in transmission and dissemination in C57BL/6 mice.
Parasites Vectors **2017**, 10, 313, doi:10.1186/s13071-017-2243-0.

42.   Malovrh, T.; Skoberne, M.; Gruntar, I.; Kotnik, V. The cell-mediated immune response to
Borrelia afzelii, garinii and burgdorferi in C57BL/6 mice is dependent on antigen specificity.
FEMS Immunol Med Microbiol **2000**, 28, 233-240, doi:10.1111/j.1574-695X.2000.tb01482.x.

43.   Akin, E.; Aversa, J.; Steere, A.C. Expression of adhesion molecules in synovia of patients with
treatment-resistant lyme arthritis. *Infect. Immun.* **2001**, 69, 1774-1780,
doi:10.1128/IAI.69.3.1774-1780.2001.

44.   Steere, A.C.; Duray, P.H.; Butcher, E.C. Spirochetal antigens and lymphoid cell surface markers

in Lyme synovitis. Comparison with rheumatoid synovium and tonsillar lymphoid tissue. *Arthritis Rheum* **1988**, 31, 487-495, doi:10.1002/art.1780310405.

45.	Shin, J.J.; Glickstein, L.J.; Steere, A.C. High levels of inflammatory chemokines and cytokines in joint fluid and synovial tissue throughout the course of antibiotic-refractory lyme arthritis. *Arthritis Rheum* **2007**, 56, 1325-1335, doi:10.1002/art.22441.

46.	Zerrouk, N.; Alcraft, R.; Hall, B.A.; Augé, F.; Niarakis, A. Large-scale computational modelling of the M1 and M2 synovial macrophages in rheumatoid arthritis. *npj Syst. Biol. Appl.* **2024**, 10, 10, doi:10.1038/s41540-024-00337-5.

47.	Bruno, D.; Somma, D.; Tolusso, B.; Di Mario, C.; Coletto, L.A.; Elmesmari, A.; Petricca, L.; Gigante, M.R.; Perniola, S.; Paglionico, A. et al. POS0425 SINGLE-CELL RNA SEQUENCING OF SYNOVIAL TISSUE-DERIVED MYELOID CELLS IN PSORIATIC ARTHRITIS PATIENTS ACROSS DISEASE PHASES. *null* **2023**, 0, 0-0, doi:10.1136/annrheumdis-2023-eular.3047.

48.	Edalat, S.G.; Gerber, R.; Houtman, M.; Kuret, T.; Izanc, N.; Micheroli, R.; Buerki, K.; Burja, B.; Pauli, C.; Rotar, Ž. et al. A reference single-cell map of freshly dissociated human synovium in inflammatory arthritis with an optimized dissociation protocol for prospective synovial biopsy collection. *bioRxiv (Cold Spring Harbor Laboratory)* **2022**, 0, 0-0, doi:10.1101/2022.06.01.493823.

49.	Siebeler, R.; de Winther, M.; Hoeksema, M.A. The regulatory landscape of macrophage interferon signaling in inflammation. *J. Allergy. Clin. Immunol.* **2023**, 152, 326-337, doi:10.1016/j.jaci.2023.04.022.

50.	Arandjelovic, S.; Ravichandran, K.S. Phagocytosis of apoptotic cells in homeostasis. *Nat. Immunol.* **2015**, 16, 907-917, doi:10.1038/ni.3253.

51.	Guha, R.A.; Odum, O.P.; Wiseman, D.; Weinstock, A. The diverse roles of macrophages in metabolic inflammation and its resolution. *Front. Cell. Dev. Biol.* **2023**, 11, 1147434, doi:10.3389/fcell.2023.1147434.

52.	Lechner, A.; Henkel, F.; Hartung, F.; Bohnacker, S.; Alessandrini, F.; Gubernatorova, E.O.; Drutskaya, M.S.; Angioni, C.; Schreiber, Y.; Haimerl, P. et al. Macrophages acquire a TNF-dependent inflammatory memory in allergic asthma. *J. Allergy. Clin. Immunol.* **2022**, 149, 2078-2090, doi:10.1016/j.jaci.2021.11.026.

53.	Luecke, S.; Adelaja, A.; Guo, X.; Sen, S.; Spreafico, R.; Singh, A.; Liu, Y.; Taylor, B.; Diaz, J.; Cheng, Q. et al. Tonic TNF conditioning of macrophages safeguards stimulus-specific inflammatory responses. *EMBO Rep.* **2023**, 24, e55986, doi:10.15252/embr.202255986.

54.	Lee, J.W.; Chun, W.; Lee, H.J.; Min, J.H.; Kim, S.M.; Seo, J.Y.; Ahn, K.S.; Oh, S.R. The Role of Macrophages in the Development of Acute and Chronic Inflammatory Lung Diseases. *Cells* **2021**, 10, doi:10.3390/cells10040897.

55.	Meizlish, M.L.; Kimura, Y.; Pope, S.D.; Matta, R.; Kim, C.; Philip, N.H.; Meyaard, L.; Gonzalez, A.; Medzhitov, R. Macrophages sense ECM mechanics and growth factor availability through cytoskeletal remodeling to regulate their tissue repair program. *bioRxiv (Cold Spring Harbor Laboratory)* **2023**, 0, 0-0, doi:10.1101/2023.06.28.545586.

56.	Park, M.H.; Lee, E.D.; Chae, W.J. Macrophages and Wnts in Tissue Injury and Repair. *Cells* **2022**, 11, doi:10.3390/cells11223592.

57.	Ma, R.Y.; Zhang, H.; Li, X.F.; Zhang, C.B.; Selli, C.; Tagliavini, G.; Lam, A.D.; Prost, S.; Sims, A.H.; Hu, H.Y. et al. Monocyte-derived macrophages promote breast cancer bone metastasis

outgrowth. *J. Exp. Med.* **2020**, 217, doi:10.1084/jem.20191820.

58. Grinberg, M.V.; Lokhonina, A.; Vishnyakova, P.; Макаров, А.B.; Kananykhina, E.; Eremina, I.Z.; Глинкина, B.B.; Elchaninov, A.; Fatkhudinov, T. Migration, proliferation and cell death of regenerating liver macrophages in an experimental model. *RUDN Journal of Medicine* **2023**, 27, 449-458, doi:10.22363/2313-0245-2023-27-4-449-458.

59. Yang, S.; Zhao, M.; Jia, S. Macrophage: Key player in the pathogenesis of autoimmune diseases. *Front. Immunol.* **2023**, 14, 0-0, doi:10.3389/fimmu.2023.1080310.

60. Yadav, S.; Prasannan, A.; Venkatachalam, K.; Binesh, A. Exploring the mechanism and crosstalk between IL-6 and IL- 1β on M2 macrophages under metabolic stress conditions. *Cytokine* **2025**, 186, 156852, doi:10.1016/j.cyto.2024.156852.

61. Lee, J.W.; Chun, W.; Lee, H.J.; Min, J.H.; Kim, S.M.; Seo, J.Y.; Ahn, K.S.; Oh, S.R. The Role of Macrophages in the Development of Acute and Chronic Inflammatory Lung Diseases. *Cells* **2021**, 10, doi:10.3390/cells10040897.

62. Chousterman, B.G.; Boissonnas, A.; Poupel, L.; Baudesson, D.C.C.; Adam, J.; Tabibzadeh, N.; Licata, F.; Lukaszewicz, A.C.; Lombès, A.; Deterre, P. et al. Ly6Chigh Monocytes Protect against Kidney Damage during Sepsis via a CX3CR1-Dependent Adhesion Mechanism. *J. Am. Soc. Nephrol.* **2016**, 27, 792-803, doi:10.1681/ASN.2015010009.

63. Tran, T.; Lavillegrand, J.R.; Lereverend, C.; Esposito, B.; Cartier, L.; Montabord, M.; Tran-Rajau, J.; Diedisheim, M.; Gruel, N.; Ouguerram, K. et al. Mild dyslipidemia accelerates tumorigenesis through expansion of Ly6C(hi) monocytes and differentiation to pro-angiogenic myeloid cells. *Nat. Commun.* **2022**, 13, 5399, doi:10.1038/s41467-022-33034-0.

64. Singh, R.; Hemati, H.; Bajpai, M.; Yadav, P.; Maheshwari, A.; Kumar, S.; Agrawal, S.; Sevak, J.K.; Islam, M.; Mars, J.S. et al. Sustained expression of inflammatory monocytes and activated T cells in COVID-19 patients and recovered convalescent plasma donors. *IMMUN. INFLAMM. DIS.* **2021**, 9, 1279-1290, doi:10.1002/iid3.476.

65. Booiman, T.; Wit, F.W.; Maurer, I.; De Francesco, D.; Sabin, C.A.; Harskamp, A.M.; Prins, M.; Garagnani, P.; Pirazzini, C.; Franceschi, C. et al. High Cellular Monocyte Activation in People Living With Human Immunodeficiency Virus on Combination Antiretroviral Therapy and Lifestyle-Matched Controls Is Associated With Greater Inflammation in Cerebrospinal Fluid. *Open Forum Infect. Dis.* **2017**, 4, ofx108, doi:10.1093/ofid/ofx108.

66. Xu, Y.; Murphy, A.J.; Fleetwood, A.J. Hematopoietic Progenitors and the Bone Marrow Niche Shape the Inflammatory Response and Contribute to Chronic Disease. *Int. J. Mol. Sci.* **2022**, 23, doi:10.3390/ijms23042234.

67. Bailey, K.N.; Alliston, T. At the Crux of Joint Crosstalk: TGFβ Signaling in the Synovial Joint. *Curr. Rheumatol. Rep.* **2022**, 24, 184-197, doi:10.1007/s11926-022-01074-6.

68. Grygorczuk, S.; Chmielewski, T.; Zajkowska, J.; Swierzbińska, R.; Pancewicz, S.; Kondrusik, M.; Tylewska-Wierzbanowska, S.; Hermanowska-Szpakowicz, T. Concentration of TGF-beta1 in the supernatant of peripheral blood mononuclear cells cultures from patients with early disseminated and chronic lyme borreliosis. *Adv. Med. Sci.* **2007**, 52, 174-178.

69. Widhe, M.; Grusell, M.; Ekerfelt, C.; Vrethem, M.; Forsberg, P.; Ernerudh, J. Cytokines in Lyme borreliosis: lack of early tumour necrosis factor-alpha and transforming growth factor-beta1 responses are associated with chronic neuroborreliosis. *Immunology* **2002**, 107, 46-55, doi:10.1046/j.1365-2567.2002.01500.x.

70. Londoño, D.; Cadavid, D.; Drouin, E.E.; Strle, K.; McHugh, G.; Aversa, J.M.; Steere, A.C.
Antibodies to endothelial cell growth factor and obliterative microvascular lesions in the
synovium of patients with antibiotic-refractory lyme arthritis. *Arthritis Rheumatol.* **2014**, 66,
2124-2133, doi:10.1002/art.38618.

71. Aarts, J.; van Caam, A.; Chen, X.; Marijnissen, R.J.; Helsen, M.M.; Walgreen, B.; Vitters, E.L.;
van de Loo, F.A.; van Lent, P.L.; van der Kraan, P.M. et al. Local inhibition of TGF-β1 signaling
improves Th17/Treg balance but not joint pathology during experimental arthritis. *Sci. Rep.*
2022, 12, 3182, doi:10.1038/s41598-022-07075-w.

72. Erlich, J.R.; To, E.E.; Luong, R.; Liong, F.; Liong, S.; Oseghale, O.; Miles, M.A.; Bozinovski,
S.; Brooks, R.D.; Vlahos, R. et al. Glycolysis and the Pentose Phosphate Pathway Promote
LPS-Induced NOX2 Oxidase- and IFN-β-Dependent Inflammation in Macrophages.
Antioxidants **2022**, 11, doi:10.3390/antiox11081488.

73. Jiang, Y.; Weng, W.; Liu, Y.; Zhong, S.; Pan, Y.; Gu, J. POS0426 PKM2 PROMOTES
PRO-INFLAMMATORY MACROPHAGE ACTIVATION IN ANKYLOSING
SPONDYLITIS. *null* **2023**, 0, 0-0, doi:10.1136/annrheumdis-2023-eular.2926.

74. Mouton, A.J.; Aitken, N.M.; Moak, S.P.; Do, C.J.; Da, S.A.; Omoto, A.; Li, X.; Wang, Z.;
Schrimpe-Rutledge, A.C.; Codreanu, S.G. et al. Temporal changes in glucose metabolism
reflect polarization in resident and monocyte-derived macrophages after myocardial infarction.
Front. Cardiovasc. Med. **2023**, 10, 1136252, doi:10.3389/fcvm.2023.1136252.

75. Tang, J.M.; Wan, X.; ZHANG, J.; Diao, N.; Zhang, C.; Gao, X.; Ren, D. A frameshift variant in
the SIRPB1 gene confers susceptibility to Crohn's disease in a Chinese population. *Front.*
Genet. **2023**, 14, 0-0, doi:10.3389/fgene.2023.1130529.

76. Kanazawa, Y.; Saito, Y.; Supriatna, Y.; Tezuka, H.; Kotani, T.; Murata, Y.; Okazawa, H.;
Ohnishi, H.; Kinouchi, Y.; Nojima, Y. et al. Role of SIRPα in regulation of mucosal immunity in
the intestine. *Genes Cells* **2010**, 15, 1189-1200, doi:10.1111/j.1365-2443.2010.01453.x.

77. Kong, X.N.; Yan, H.X.; Chen, L.; Dong, L.W.; Yang, W.; Liu, Q.; Yu, L.X.; Huang, D.D.; Liu,
S.Q.; Liu, H. et al. LPS-induced down-regulation of signal regulatory protein {alpha}
contributes to innate immune activation in macrophages. *J. Exp. Med.* **2007**, 204, 2719-2731,
doi:10.1084/jem.20062611.

78. Zhou, Y.; Elmes, M.W.; Sweeney, J.M.; Joseph, O.M.; Che, J.; Hsu, H.C.; Li, H.; Deutsch, D.G.;
Ojima, I.; Kaczocha, M. et al. Identification of Fatty Acid Binding Protein 5 Inhibitors Through
Similarity-Based Screening. *Biochemistry.* **2019**, 58, 4304-4316,
doi:10.1021/acs.biochem.9b00625.

79. Bando, Y.; Sakashita, H.; Nagasaka, A.; Sakiyama, K.; Tokuda, N.; Iseki, S.; Owada, Y.; Amano,
O. Septoclasts expressing epidermal fatty acid-binding protein (E-FABP, FABP5) in
endochondral ossification. *J. Oral Biosci.* **2022**, 64, 18-25, doi:10.1016/j.job.2021.12.003.

80. Gordon, C.; Trainor, J.; Shah, R.J.; Studholme, K.M.; Gelman, A.; Doswell, F.; Sadar, F.;
Giovannetti, A.; Gershenson, J.; Khan, A. et al. Fatty acid binding protein 5 inhibition
attenuates pronociceptive cytokine/chemokine expression and suppresses osteoarthritis pain: A
comparative human and rat study. *Osteoarthritis Cartilage* **2023**, 0, 0-0,
doi:10.1016/j.joca.2023.11.010.

81. Xu, J.; Zheng, B.; Xie, C.; Zhao, Y.; Wu, H.; Wang, Y.; Guan, X.; Lei, X.; Liu, D.; Lou, X. et al.
Inhibition of FABP5 attenuates inflammatory bowel disease by modulating macrophage
alternative activation. *Biochem. Pharmacol.* **2024**, 219, 115974,

doi:10.1016/j.bcp.2023.115974.

82. Yeung, C.C.; Garva, R.; Pickard, A.; Lu, Y.; Mallikarjun, V.; Swift, J.; Taylor, S.H.; Rai, J.; Eyre, D.R.; Chaturvedi, M. et al. Mmp14 is required for matrisome homeostasis and circadian rhythm in fibroblasts. *Matrix Biol.* **2023**, 124, 8-22, doi:10.1016/j.matbio.2023.10.002.

83. Li, R.L.; Duan, H.X.; Liang, Q.; Huang, Y.L.; Wang, L.Y.; Zhang, Q.; Wu, C.J.; Liu, S.Q.; Peng, W. Targeting matrix metalloproteases: A promising strategy for herbal medicines to treat rheumatoid arthritis. *Front. Immunol.* **2022**, 13, 1046810, doi:10.3389/fimmu.2022.1046810.

84. Ishikawa, T.; Nishigaki, F.; Miyata, S.; Hirayama, Y.; Minoura, K.; Imanishi, J.; Neya, M.; Mizutani, T.; Imamura, Y.; Naritomi, Y. et al. Prevention of progressive joint destruction in collagen-induced arthritis in rats by a novel matrix metalloproteinase inhibitor, FR255031. *Br. J. Pharmacol.* **2005**, 144, 133-143, doi:10.1038/sj.bjp.0706054.

85. Kaneko, K.; Williams, R.O.; Dransfield, D.T.; Nixon, A.E.; Sandison, A.; Itoh, Y. Selective Inhibition of Membrane Type 1 Matrix Metalloproteinase Abrogates Progression of Experimental Inflammatory Arthritis: Synergy With Tumor Necrosis Factor Blockade. *Arthritis Rheumatol.* **2016**, 68, 521-531, doi:10.1002/art.39414.

86. Banerji, R.; Saroj, S.D. Early growth response 1 (EGR1) activation in initial stages of host-pathogen interactions. *Mol. Biol. Rep.* **2021**, 48, 2935-2943, doi:10.1007/s11033-021-06305-0.

87. Lehman, C.W.; Smith, A.; Kelly, J.; Jacobs, J.L.; Dinman, J.D.; Kehn-Hall, K. EGR1 Upregulation during Encephalitic Viral Infections Contributes to Inflammation and Cell Death. *Viruses* **2022**, 14, doi:10.3390/v14061210.

88. Pang, Z.; Raudonis, R.; McCormick, C.; Cheng, Z. Early Growth Response 1 Deficiency Protects the Host against Pseudomonas aeruginosa Lung Infection. *Infect. Immun.* **2019**, 88, doi:10.1128/IAI.00678-19.

89. Sheng, J.; Liu, D.; Kang, X.; Chen, Y.; Jiang, K.; Zheng, W. Egr-1 increases angiogenesis in cartilage via binding Netrin-1 receptor DCC promoter. *J. Orthop. Surg. Res.* **2018**, 13, 125, doi:10.1186/s13018-018-0826-x.

90. Nebbaki, S.S.; El, M.F.; Afif, H.; Kapoor, M.; Benderdour, M.; Duval, N.; Pelletier, J.P.; Martel-Pelletier, J.; Fahmi, H. Egr-1 contributes to IL-1-mediated down-regulation of peroxisome proliferator-activated receptor γ expression in human osteoarthritic chondrocytes. *Arthritis Res. Ther.* **2012**, 14, R69, doi:10.1186/ar3788.

91. Havis, E.; Duprez, D. EGR1 Transcription Factor is a Multifaceted Regulator of Matrix Production in Tendons and Other Connective Tissues. *Int. J. Mol. Sci.* **2020**, 21, doi:10.3390/ijms21051664.

A

Bb-infected group

Control group

B

Bb-infected group

Control group

C

Bb-infected group

Control group

D

Relative mRNA level of *SIRPB1C*

```
8 ┤
  │              **
6 ┤           ┌──────┐
  │              ┬
4 ┤              │
  │              │
2 ┤              │
  │   ┌──┐       │
0 ┴───┴──┴───────┴──┴──
     PBS      Bb infection
```

E

Relative mRNA level of *FABP5*

```
2.0 ┤
    │            *
1.5 ┤         ┌─────┐
    │            ┬
1.0 ┤  ┌──┐      │
    │  │  │      │
0.5 ┤  │  │      │
    │  │  │      │
0.0 ┴──┴──┴──────┴──┴──
      PBS    Bb infection
```

F

Relative mRNA level of *MMP14*

```
2.0 ┤
    │            *
    │         ┌─────┐
1.5 ┤            ┬
    │            │
1.0 ┤  ┌──┐      │
    │  │  │      │
0.5 ┤  │  │      │
    │  │  │      │
0.0 ┴──┴──┴──────┴──┴──
      PBS    Bb infection
```

J

Relative mRNA level of *EGR1*

```
2.5 ┤
    │            *
2.0 ┤         ┌─────┐
    │            ┬
1.5 ┤            │
    │  ┌──┐      │
1.0 ┤  │  │      │
    │  │  │      │
0.5 ┤  │  │      │
0.0 ┴──┴──┴──────┴──┴──
      PBS    Bb infection
```

H

Relative mRNA level of *SIRPB1C*

```
2.0 ┤
    │            *
1.5 ┤         ┌─────┐
    │  ┌──┐      ┬
1.0 ┤  │  │      │
    │  │  │      │
0.5 ┤  │  │      │
    │  │  │      │
0.0 ┴──┴──┴──────┴──┴──
      PBS    Bb infection
```

I

Relative mRNA level of *FABP5*

```
2.0 ┤
    │            *
    │         ┌─────┐
1.5 ┤            ┬
    │  ┌──┐      │
1.0 ┤  │  │      │
    │  │  │      │
0.5 ┤  │  │      │
0.0 ┴──┴──┴──────┴──┴──
      PBS    Bb infection
```

J

Relative mRNA level of *MMP14*

```
6 ┤
  │              **
  │           ┌──────┐
4 ┤              ┬
  │              │
  │              │
2 ┤              │
  │   ┌──┐       │
0 ┴───┴──┴───────┴──┴──
     PBS      Bb infection
```

K

Relative mRNA level of *EGR1*

```
5 ┤
  │              *
4 ┤           ┌──────┐
  │              ┬
3 ┤              │
  │              │
2 ┤              │
  │   ┌──┐       │
1 ┤   │  │       │
0 ┴───┴──┴───────┴──┴──
     PBS      Bb infection
```