Hierarchical encoding of natural sounds mixtures in ferret auditory cortex

Agnès Landemard[1,2], Célian Bimbard[1,2], and Yves Boubenec[1, *]

[1]Laboratoire des systèmes perceptifs, Département d'études cognitives, École normale supérieure, PSL University, Paris, France
[2]UCL Institute of Ophthalmology, University College London, London, UK
[*]Corresponding author: boubenec@ens.fr

Abstract

Extracting relevant auditory signals from complex natural scenes is a fundamental challenge for the auditory system. Sounds from multiple sources overlap in time and frequency. In particular, dynamic "foreground" sounds are often masked by more stationary "background" sounds. Human auditory cortex exhibits a hierarchical organization where background-invariant representations are progressively enhanced along the processing stream, from primary to non-primary regions. However, we do not know whether this organizational principle is conserved across species and which neural mechanisms drive this invariance. To address these questions, we investigated background invariance in ferret auditory cortex using functional ultrasound imaging (fUSI), which enables large-scale, high-resolution recordings of hemodynamic responses. We measured responses across primary, secondary, and tertiary auditory cortical regions as ferrets passively listened to mixtures of natural sounds and their components in isolation. We found a hierarchical gradient of background invariance, mirroring findings in humans: responses in primary auditory cortex reflected contributions from both foreground and background sounds, while background invariance increased in higher-order auditory regions. Using a spectrotemporal filter-bank model, we found that in ferrets, this hierarchical structure could be largely explained by tuning to low-order acoustic features. However, this model failed to fully account for background invariance in human non-primary auditory cortex, suggesting that additional, higher-order mechanisms are crucial for background segregation in humans.

Introduction

In the intricate auditory landscapes of our surroundings, the brain quickly extracts meaningful information from a cacophony of overlapping sources. Humans are remarkably good at detecting speech in complex noise: two people in a bar can easily have a conversation despite the ambient chatter. This ability is thought to rely on the different properties of background and foreground sounds, which are mixed in a single waveform at the level of the cochlea. Dynamic "foreground" sounds, such as speech or vocalizations, fluctuate over short timescales, and these fluctuations convey new information over time (Singh & Theunissen, 2003). In contrast, stationary "background" sounds, like ambient chatter or the rustle of wind, fluctuate over longer periods, and are thus more predictable and less vital for signaling sudden events. They can be synthesized realistically based solely on their time-averaged acoustic statistics (McWalter & McDermott, 2018; McDermott & Simoncelli, 2011; McWalter & Dau, 2017), which the brain uses to perceptually fill short acoustic gaps (McWalter & McDermott, 2019). Despite these differences in the nature of foreground and background

sounds, we still lack a general understanding of where background invariance occurs in the brain and what neural mechanisms underlie it.

In humans, multiple stages of processing build up background-invariant representations. In fMRI recordings of auditory cortex, non-primary areas are more background-invariant than primary areas (Kell & McDermott, 2019). In addition, intracranial cortical responses show rapid adaptation to new background noise, which improves representations of foreground speech (Khalighinejad et al., 2019). This effect is stronger for neurons in non-primary auditory cortex. This progressive invariance is unlikely to be fully explained by the parallel hierarchical organization of speech and music processing in humans (Norman-Haignere et al., 2015; Norman-Haignere & McDermott, 2018), since it is still present when excluding speech and music from foreground sounds (Kell & McDermott, 2019). Moreover, background invariance appears to be independent of selective attention (Kell & McDermott, 2019; Khalighinejad et al., 2019). Thus, the hierarchical organization of background invariance may not be specific to humans but rather a generic principle shared with other animals.

Background invariance can also be found in non-human animals. In fact, neurons that exhibit background-invariant responses are present throughout the auditory pathway in multiple species, increasingly from subcortical nuclei to the auditory cortex (Souffi et al., 2020; Rabinowitz et al., 2013). Background-invariant representations in auditory cortex are necessary for behavioral discrimination of natural sounds in noise (Town et al., 2023). Neurons with foreground-invariant responses are also found in the non-human auditory cortex (Bar-Yosef, 2007; Ni et al., 2017; Hamersky et al., 2023), but a critical transformation of representations within the cortical hierarchy could specifically enhance foreground representations (Schneider & Woolley, 2013; Saderi et al., 2020; Carruthers et al., 2015). However, these studies differ by species, regions, state (awake vs. anesthetized), type of background noise (synthetic vs. natural) and signal-to-noise ratio. Thus, it remains unclear whether a cortical gradient in background invariance, as seen in humans, is present in other animals.

Mechanistically, background invariance could come from the simple tuning properties of neurons. Auditory cortical neurons are known to be tuned to frequency and spectrotemporal modulations (Miller et al., 2002; Shamma, 2009), which are acoustic features that differ between background and foreground sounds (Elie & Theunissen, 2015). In ferrets, this low-level spectrotemporal tuning can explain large-scale responses to natural sounds in both primary and non-primary regions of auditory cortex (Landemard et al., 2021). Thus, differences in background invariance across regions could result from regional differences in such low-order tuning (Moore et al., 2013; Ivanov et al., 2022). In humans, this tuning can explain responses in primary auditory cortex (Santoro et al., 2014), but not in non-primary fields (Norman-Haignere & McDermott, 2018). Consequently, the sensitivity of human non-primary cortex to acoustic high-order acoustic features may underlie the hierarchical gradient in background invariance. However, it remains untested whether background invariance in the auditory cortex relies on neural mechanisms that are conserved across species.

Here, we asked (1) whether background invariance is hierarchically organized in the auditory cortex of a non-human mammal, ferret, and (2) whether hierarchical background invariance could be explained by tuning to low-order acoustic properties that differentiate foreground-like and background-like natural sounds. We defined foreground and background sounds based on an acoustic criterion of stationarity. We measured large-scale hemodynamic responses in ferret auditory cortex as ferrets passively listened to isolated and combined foregrounds and backgrounds. Leveraging the spatial resolution and coverage of functional ultrasound imaging (fUSI), we showed that background invariance increased from primary to secondary and tertiary areas of auditory cortex. We could predict these effects using a model based on frequency and spectrotemporal modulations, suggesting that the organization of background invariance in the ferret brain can be derived from neural spectrotemporal tuning. Finally, we showed that the same model could not explain as well the patterns of background invariance found in human auditory cortex,

suggesting that additional mechanisms underlie background invariance in humans.

Results

Using functional ultrasound imaging (fUSI), we measured hemodynamic responses in the auditory cortex of ferrets passively listening to continuous streams of natural sounds. We used three types of stimuli: foregrounds, backgrounds, and combinations of those. To define foregrounds and backgrounds, we computed each sound's stationarity, defined as the stability in time of a sound's acoustic statistics (McWalter & McDermott, 2018). Backgrounds were chosen as the most stationary and foregrounds as the least stationary sounds (Figure 1A, table 1). Different sound segments were concatenated to form continuous streams (Figure 1B). In the isolated condition, a sound change occurred every 9.6 seconds. In the mixture condition, the same sequences overlapped with a lag of 4.8 s between the foreground and background streams, leading to a change in one or the other stream every 4.8 s.

Figure 1: **Hemodynamic activity reflects encoding of foregrounds and backgrounds.** (**A**) Stationarity for foregrounds (squares) and backgrounds (diamonds). (**B**) Sound presentation paradigm, with example cochleagrams. We created continuous streams by concatenating 9.6-second foreground (cold colors) and background segments (warm colors) following the illustrated design. Each foreground (resp. background) stream was presented in isolation and with two different background (resp. foreground) streams. (**C**) We measured cerebral blood volume (CBV) in coronal slices (blue plane) of the ferret auditory cortex (black outline) with functional ultrasound imaging. We imaged the whole auditory through successive slices across several days. (**D**) Average changes in CBV (normalized to silent baseline) in auditory cortex aligned to sound changes, for different conditions, averaged across all ferrets. Shaded area represents standard error of the mean across sound segments. (**E**) Test-retest cross-correlation for each condition. Voxel responses for two repeats of sounds are correlated with different lags. Resulting matrices are then averaged across all responsive voxels (ΔCBV > 2.5%).

Hemodynamic responses stably encoded sound identity after transient responses. We continuously measured hemodynamic activity in ferret auditory cortex (Figure 1C) throughout the sound sequences and normalized cerebral blood volume (CBV) relative to blocks of silent baseline. Sound changes elicited transient increases in activity before reaching a sustained level (Figure 1D). Sustained activity was lower for backgrounds in isolation than for foregrounds and mixtures. We asked whether sound encoding was stable over time despite these changes in amplitude. We focused on the sound-specific activity of responsive voxels (ΔCBV > 2.5%) by subtracting their average timecourse across all sounds in each category. For each voxel, we cross-correlated vectors of responses to sounds of each category across two repetitions (Figure

1E). Voxels stably encoded sounds of each category throughout their duration despite the observed variation in response amplitude. Thus, we summarized voxels' activity by their time-averaged activity over an interval spanning 2 to 4.8 s after sound change.

Invariance to background sounds is hierarchically organized in ferret auditory cortex

We investigated the spatial organization of sound responses throughout the whole ferret auditory cortex. We found that reliable sound-evoked responses were not homogeneous but confined to the central part of ventral gyrus of the auditory cortex (Figure 2A, B). Hereafter, We focused on reliable sound-responsive voxels (test-retest correlation > 0.3 for sounds in at least one condition). We assigned voxels to three regions of interest in ferret auditory cortex (Figure 2C): MEG (primary), dPEG (secondary) and VP (tertiary, see Elgueda et al., 2019; Landemard et al., 2021; Radtke-Schuller, 2018).

Invariance to background sounds was stronger in non-primary than in primary auditory cortex. To probe invariance to background sounds for each voxel, we compared responses to mixtures and isolated foregrounds across sound segments (Figure 2D). Responses to mixtures and isolated foregrounds tended to be more similar for voxels in non-primary regions. We quantified this effect by computing a noise-corrected version of the Pearson correlation between responses to mixtures and foregrounds for each voxel, which we call background invariance (Kell & McDermott, 2019). Background invariance emerged shortly after a sound change and was stable throughout the duration of each sound snippet (Figure S1). When going up the hierarchy, correlations between mixtures and foregrounds increased (tertiary (VP) > secondary (dPEG) > primary (MEG), $p = 0.001$ with a permutation test, see table 2 for detailed statistics), indicating a stronger invariance to background sounds in higher-order areas, especially in VP (Figures 2E, F, figure S2).

To test whether foregrounds were processed differently than backgrounds, and at which stage, we also compared responses to mixtures and backgrounds in isolation. We defined a "foreground invariance", symmetric to the background invariance, by correlating responses to mixtures and isolated backgrounds across sound segments (Figure 2G). In primary auditory cortex (MEG), there was no significant difference between values of foreground invariance and background invariance ($p = 0.063$, obtained by randomly permuting the sounds' background and foreground labels, 1000 times). However, foreground invariance tended to decrease from primary to non-primary areas (Figure 2H, I, VP < dPEG: $p = 0.001$, dPEG < MEG: $p = 0.136$ with a permutation test). As a consequence, foreground invariance became lower than background invariance in both dPEG ($p = 0.001$) and VP ($p = 0.001$).

A model of auditory processing predicts hierarchical differences

By definition, foregrounds and backgrounds are separated in the acoustic domain. Low-level acoustic features computed from their waveforms can well discriminate both categories. To show this, we used a standard two-stage filter-bank model of auditory processing (Chi et al., 2005). This model (1) computes a sound's cochleagram, (2) convolves the resulting cochleagram through a bank of spectrotemporal modulation filters, tuned to a range of frequencies and spectrotemporal modulations (Figure 3A). Foregrounds and backgrounds differed in their pattern of energy in the modulation space (Figure 3B, C). In particular, the axis of temporal modulation was highly discriminative: foregrounds tended to have higher energy in the low rates (< 8 Hz), and backgrounds in higher rates (> 8 Hz).

In ferret auditory cortex, large-scale cortical responses to natural sounds are well predicted by the spectrotemporal model, even when controlling for stimuli correlations (Landemard et al., 2021). To test whether differences in neural invariance could be explained by tuning to low-order acoustic features, we predicted the voxels' responses using this model. We derived voxels weights for model features using cross-validated ridge regression of trial-averaged responses to all sounds (Figure 3D, E).

Tuning to frequency and spectrotemporal modulations showed large-scale spatial structure. Through

Figure 2: **Invariance to background sounds is hierarchically organized in ferret auditory cortex.**
(**A**) Map of average response for an example hemisphere (ferret L). Responses are expressed in percent changes in CBV relative to baseline activity, measured in periods of silence. Values are averaged across depth to obtain this surface view of auditory cortex. (**B**) Map of test-retest reliability. In the following maps, only reliably responding voxels are displayed (test-retest > 0.3 for at least one category of sounds) and the transparency of surface bins in the maps is determined by the number of (reliable) voxels included in the average. (**C**) Map of considered regions of interest (ROIs), based on anatomical landmarks. The arrows indicate the example slices shown in **D** (orange: primary; green: non-primary example). (**D**) Responses to isolated and combined foregrounds. Bottom: Responses to mixtures and foregrounds in isolation for example voxels (left:primary; right:non-primary). Each dot represents the voxel's time-averaged response to every foreground (x-axis) and mixture (y-axis), averaged across two repetitions. r indicates the value of the Pearson correlation. Top: Maps show invariance, defined as noise-corrected correlation between mixtures and foregrounds in isolation, for the example voxel's slice with values overlaid on anatomical images representing baseline CBV. Example voxels are shown with white squares. (**E**) Map of background invariance for the same hemisphere (see Figure S2 for other ferrets). (**F**) Quantification of background invariance for each ROI. The crosses (+) indicate median values across all voxels of each ROI, across animals. Grey dots represent median values across primary (MEG) and non-primary (dPEG + VP) voxels for each animal. The size of each dot is proportional to the number of voxels across which the median is taken. The thicker line corresponds to the example ferret L. ***: $p <= 0.001$ for comparing the average background invariance across animals for pairs of ROIs, obtained by a permutation test of voxel ROI labels within each animal. (**G-I**) Same as D-F for foreground invariance (comparing mixtures to backgrounds in isolation). *AEG*, anterior ectosylvian gyrus; *MEG*, medial ectosylvian gyrus; *dPEG*, dorsal posterior ectosylvian gyrus; *VP*, ventral posterior auditory field.

the model weights, we first explored whether tuning to acoustic features was topographically organized. We extracted the voxels' preferred frequency, spectral and temporal modulations by marginalizing over the other features. We retrieved the tonotopic organization of the auditory cortex, confirming our anatomical segmentation into functional areas (Elgueda et al., 2019). The voxels' preferred spectral and temporal modulations were also spatially organized (Figure 3F, figure S3 for all ferrets). To directly examine the differences of tuning between non-primary and primary areas, we computed the differences in weights between dPEG

Figure 3: **Simple spectrotemporal tuning explains spatial organization of background invariance.**
(**A**) Presentation of the two-stage filter-bank, or spectrotemporal model. Cochleagrams (shown for an example foreground and background) are convolved through a bank of spectrotemporal modulation filters.
(**B**) Energy of foregrounds and backgrounds in spectrotemporal modulation space, averaged across all frequency bins. (**C**) Average difference of energy between foregrounds and backgrounds in the full acoustic feature space (frequency * temporal modulation * spectral modulation). (**D**) We predicted time-averaged voxel responses using sound features derived from the spectrotemporal model presented in **A** with ridge regression. For each voxel, we thus obtain a set of weights for frequency and spectrotemporal modulation features, as well as cross-validated predicted responses to all sounds. (**E**) Average model weights for MEG. (**F**) Maps of preferred frequency, temporal and spectral modulation based on the fit model. To calculate the preferred value for each feature, we marginalized the weight matrix over the two other dimensions. (**G**) Average differences of weights between voxels of each non-primary (dPEG and VP) and primary (MEG) region. (**H**) Background invariance (left) and foreground invariance (right) for voxels tuned to low (< 8 Hz) or high (> 8 Hz) temporal modulation rates within each ROI. ***: $p <= 0.001$ for comparing the average background invariance across animals for voxels tuned to low vs. high rates, obtained by a permutation test of tuning within each animal.

vs. MEG and VP vs. MEG, throughout the whole three-dimensional acoustic space (Figure 3G). The weights of voxels in non-primary areas differed from those in primary areas in a complex manner, in contrast to the simple separation of the acoustic space between foregrounds and backgrounds (Figure 3C). Compared to MEG, dPEG was preferentially tuned to high frequencies and higher spectral modulations (Figure 3G top), while VP tuning was biased towards low frequencies and intermediate spectrotemporal modulations (Figure 3G bottom).

We sought to assess whether these differences in tuning underlaid differences in background in-

variance across regions. Since the temporal modulation axis was the most informative to discriminate foregrounds from backgrounds (Figure 3C), we first compared the invariance properties of voxels tuned to low or high temporal modulation rates. Within each region, voxels tuned to lower modulation rates displayed higher background invariance and lower foreground invariance than those tuned to higher rates (Figure 3H, $p < 0.002$ within each ROI by permuting voxels' tuning, 2000 times). However, within each tuning group, differences were still present across regions (primary vs. non-primary, $p = 0.001$ within voxels tuned to high rates and within voxels tuned to low rates, see table 2 for more detailed comparisons between regions), confirming that tuning to temporal modulation was not the only mechanism at play.

Figure 4: **A model of auditory processing predicts hierarchical differences in ferret auditory cortex.** Same as in figure 2 using cross-validated predictions from the spectrotemporal model. (**A**) Predicted responses to mixtures and foregrounds in isolation for example voxels (primary, left and non-primary, right). Each dot represents the voxel's predicted response to foregrounds (x-axis) and mixtures (y-axis). r indicates the value of the Pearson correlation. Maps above show predicted invariance values for the example voxel's slice overlaid on anatomical images representing baseline CBV. Example voxels are shown with white squares. (**B**) Maps of predicted background invariance, defined as the correlation between predicted responses to mixtures and foregrounds in isolation. (**C**) Binned scatter plot representing predicted vs. measured background invariance across voxels. Each line corresponds to one animal, using 0.1 bins of measured invariance. (**D**) Predicted background invariance for each ROI. The crosses (+) indicate median value across all voxels of each ROI, across animals. Grey dots represent median values across primary (MEG) and non-primary (dPEG + VP) voxels for each animal. The size of each dot is proportional to the number of voxels across which the median is done. The thicker line corresponds to example ferret L. *: $p <= 0.05$; ***: $p <= 0.001$ for comparing the average predicted background invariance across animals for pairs of ROIs, obtained by a permutation test of voxel ROI labels within each animal. (**E-H**) Same as **A-D** for predicted foreground invariance, i.e. comparing predicted responses to mixtures and backgrounds in isolation.

To take into account tuning across all dimensions, we then directly looked at the invariance of responses predicted from the spectrotemporal model. The model was able to predict voxel responses accurately (Figure S5) and recapitulated invariance properties across single voxels (Figure 4A-C, E-G). To verify

how well the model could explain the observed invariance, we directly compared measured and predicted invariance across voxels for each animal, regardless of ROI (Figure 4C, G). Predictions explained a large fraction of the reliable variability in background invariance (mean noise-corrected correlation between measured and predicted invariance: $r = 0.65 \pm 0.007$ standard error of the mean across animals) and foreground invariance (mean $r = 0.43 \pm 0.04$). Predicted responses recapitulated the gradient from primary to non-primary areas for background invariance (Figure 4D ; MEG < dPEG, $p = 0.022$; dPEG < VP, $p = 0.001$, permutation test) and foreground invariance (Figure 4H; MEG > dPEG: $p = 0.014$, dPEG > VP: $p = 0.001$, permutation test). Thus, hierarchical differences in invariance were largely explained by differences in tuning to frequency and spectrotemporal modulation between regions.

Species difference in background invariance

Figure 5: **The spectrotemporal model is a poor predictor of human background invariance.** (**A**) We replicated our analyses with a dataset of a similar experiment measuring fMRI responses in human auditory cortex (Kell & McDermott, 2019). We compared responses in primary and non-primary auditory cortex, as delineated in Kell & McDermott (2019). (**B**) Responses to mixtures and foregrounds in isolation for example voxels (left: primary; right: non-primary). Each dot represents the voxel's response to foregrounds (x-axis) and mixtures (y-axis), averaged across repetitions. r indicates the value of the Pearson correlation. (**C-D**) Quantification of background invariance for each ROI, for measured responses (**C**) and responses predicted from the spectrotemporal model (**D**). The crosses (+) indicate median value across all voxels of each ROI, across subjects. Grey dots represent median values for each ROI and subject. The size of each dot is proportional to the number of (reliable) voxels across which the median is done. (**E**) Binned scatter plot representing predicted vs measured background invariance across voxels. Each line corresponds to one subject, using 0.1 bins of measured invariance. (**F-I**) Same as **B-E** for foreground invariance, i.e. comparing predicted responses to mixtures and backgrounds in isolation.

In ferrets, background invariance is hierarchically organized, and this organization can largely be explained by simple spectrotemporal tuning. Is it also the case for the humans' hierarchical organization of invariance? To test this, we compared background invariance previously obtained in humans with what

we found in ferrets. We used a comparable published dataset (Kell & McDermott, 2019), in which combinations of foregrounds and backgrounds were presented to human subjects, in isolation and in mixtures. The BOLD signal from the auditory cortex was acquired with fMRI (Figure 5A). Applying our analyses to this dataset, we reproduced the results of Kell & McDermott (2019) and showed that background invariance was stronger in non-primary than in primary auditory cortex (Figure 5B, C, non-primary > primary, $p = 0.001$, permutation test on the average across 7 subjects). In parallel, we found that foreground invariance decreased from primary to non-primary areas (Figure 5F, G, non-primary < primary, $p = 0.001$, permutation test). As in ferrets, foreground and background invariance values were similar in primary auditory cortex ($p = 0.577$, permutation test), but significantly differed in non-primary auditory cortex ($p = 0.002$, permutation test).

We next tested whether the spectrotemporal model could predict human voxel responses. The model performed similarly in humans and ferrets (Figure S5, $r = 0.42 \pm 0.01$ standard error of the mean across subjects, humans vs ferrets: $p = 0.0667$, Wilcoxon rank-sum test). However, the model explained less of the measured background invariance in humans than in ferrets (Figure 5D, noise-corrected correlation between measured and predicted invariance: $r = 0.23 \pm 0.04$, $p = 0.0167$ compared to ferrets, $p = 0.0156$ compared to 0, Wilcoxon rank-sum test). While predicted background invariance differed significantly between primary and non-primary regions (Figure 5E, primary < non-primary, $p = 0.001$, permutation test), the predicted difference was much smaller than observed. Furthermore, the model failed to predict the levels of foreground invariance in both primary and non-primary human auditory cortex (Figure 5H, I, noise-corrected correlation between measured and predicted invariance: $r = -0.17 \pm 0.06$, $p = 0.0167$ compared to ferrets, $p = 0.0312$ compared to 0, Wilcoxon rank-sum test). Thus, the spectrotemporal model performed worse in explaining the patterns of cortical invariance in humans than in ferrets. This suggests that additional higher-order mechanisms contribute to background segregation in humans.

Discussion

In this work, we investigated large-scale background and foreground invariance in auditory cortex, using mixtures of natural sounds. In ferrets, we found a hierarchical gradient of invariance, where primary cortical areas encode both background and foreground sounds, while non-primary regions show increasing invariance to background sounds. A spectrotemporal filter-bank model could account for a large portion of the hierarchical changes in ferrets, suggesting that low-level tuning to frequency and modulation properties explain how stationary "background" and more dynamic "foreground" signals are segregated in hemodynamic signal. When we applied the same model to human data, we found that although humans also display a hierarchical increase in background invariance, the model failed to capture a substantial fraction of the observed effect in non-primary regions. This suggests that mechanisms beyond simple spectrotemporal tuning are at play in human auditory cortex.

Our results show that foregrounds and backgrounds are similarly represented in primary auditory cortex, both in ferrets and humans; at this stage, we found no bias towards foregrounds. This is in line with previous work in non-human animals, including birds, where neurons in various regions up to primary auditory cortex display a continuum of background-invariant and foreground-invariant responses (Ni et al., 2017; Bar-Yosef, 2007; Schneider & Woolley, 2013; Hamersky et al., 2023). This diversity of responses may play a functional role in background denoising. Studies show complex synergetic interactions between target sounds and background noise in primary auditory cortex, even with simple stimuli. On the one hand, white noise can sharpen neurons' tuning curves, enhancing tone discrimination (Christensen et al., 2019), and improve the representations of frequency sweeps (Malone et al., 2017). On the other hand, background information can be more easily decoded from population responses when presented with a target sound

than when in isolation (Malone et al., 2017). Computationally, background-related information could facilitate further extraction of foregrounds. For instance, in a convolutional neural network trained to recognize digits in noise, when local feedback is implemented, early layers encode background properties, while later layers represent clean signals (Lindsay et al., 2022). Similarly, primary auditory areas could be recruited to maintain background representations, enabling downstream cortical regions to use these representations to specifically suppress background information and enhance foreground representations.

We found that the main difference regarding background invariance between humans and ferrets was in how well invariance could be explained by a model based on low-level acoustic features. There could be multiple reasons for this difference.

First, methodological differences could weigh in the observed species difference. In our ferret experiment, sounds were presented continuously and in longer segments, an experimental design enabled by fUSI but impractical for fMRI due to scanner noise. However, this is unlikely to have a major impact, as our analyses focused on the sustained response component, similar to the fMRI experiment. Furthermore, while fUSI and fMRI both provide hemodynamic signals, they differ in their physiological basis (CBV vs. BOLD). Studies comparing CBV-weighted and BOLD fMRI indicate that the CBV signal provides a more spatially and temporally precise than BOLD (Silva et al., 2007), improving localization of neural activity. In addition, fUSI provides a higher spatial resolution than classical fMRI (100 μm against ~2 mm voxel size). Thus, fUSI recordings can compensate for differences in brain size, allowing for a detailed mapping of the ferret auditory cortex with a greater number of voxels than typically achieved in fMRI studies of the human auditory cortex. Last, to minimize potential biases, we applied the same criteria for voxel inclusion and noise-correction procedures across datasets. Thus, while minor differences in data quality and experimental protocols exist, they are unlikely to fully account for the species differences we observed.

Second, ferret and human paradigms differed by their control of attention. Ferrets were passively listening to the sounds while human participants had to perform a simple loudness-based task to ensure attention to the sounds. Even though specific auditory attention is not necessary for the gradient in invariance (Kell & McDermott, 2019), the task ensured that subjects remained in a state of engagement or general attention. This state might in turn refine auditory representations via top-down processes (Elhilali et al., 2009; Saderi et al., 2021; Schwartz & David, 2018). Thus, we cannot exclude that both species possess a similar feedforward acoustic analysis (well modeled by simple spectrotemporal filters) as well as more context-driven networks in non-primary auditory cortex, which were more strongly engaged in humans than in ferrets due to experimental procedures. In addition, the presented sounds might have been more salient for humans than ferrets. Despite including ferret vocalizations and environmental sounds that are more ecologically relevant for ferrets, it is not clear whether ferrets would behaviorally categorize foregrounds and backgrounds as humans do. Examining how ferrets naturally orient or respond to foreground and background sounds under more ecologically valid conditions, potentially with free exploration or spontaneous listening paradigms, could help address this issue.

Third, human non-primary regions could host higher-order computations that cannot be predicted by simple acoustic features and are not present in ferrets. Human and non-human primates have a complex functional architecture in the auditory belt and parabelt regions (Hackett, 2007), likely supporting elaborate computations. These computations might be related to conspecific calls and required for advanced vocal communication. In particular, the unique importance of speech for humans has shaped auditory circuits throughout the brain. Neural circuits in human non-primary auditory cortex encode higher-order acoustic features, such as phonemic patterns, which might help to segregate meaningful content from background noise (Norman-Haignere et al., 2015; Mesgarani et al., 2014). Last, other processes might critically shape background-invariant representations in humans, such as dynamic adaptation to backgrounds that could be faster in humans than in ferrets (Khalighinejad et al., 2019).

The divergence in background invariance mechanisms between humans and ferrets may reflect

differences in the depth and complexity of their auditory cortical hierarchies. Our previous work (Lande-mard et al., 2021) showed that large-scale representations of natural sounds are not clearly distinct between primary and non-primary auditory regions in the ferret cortex. However, Sabat et al. (2025) found a hier-archical organization of temporal integration in ferrets, reminiscent of humans (Norman-Haignere et al., 2022). In the present study, we found a gradient of background invariance that was similarly structured as in humans, yet relying on different mechanisms. Altogether, this raises a fundamental question: does the ferret auditory cortex correspond to the different early auditory fields in the human core regions, with deeper hierarchical stages simply being absent? Or do these differences reflect evolutionary divergence, with partly shared computational principles and homologous hierarchy, but additional cortical modules in humans, enabling for instance more sophisticated background segregation or dedicated speech and mu-sic processing (Norman-Haignere & McDermott, 2018)? Addressing these questions will require a finer characterization of cross-species homologies, leveraging high-resolution functional imaging and electro-physiological recordings to uncover whether hierarchical transformations in auditory processing follow a conserved or species-specific trajectory.

Materials & Methods

Animal preparation

Experiments were performed in three female ferrets (B, L and R), across one or both hemispheres (left for all ferrets + right for R). Experiments were approved by the French Ministry of Agriculture (protocol au-thorization: 21022) and strictly comply with the European directives on the protection of animals used for scientific purposes (2010/63/EU). Animal preparation and fUS imaging were performed as in Bimbard et al. (2018). Briefly, a metal headpost was surgically implanted on the skull under anesthesia. After recovery from surgery, a craniotomy was performed over auditory cortex and then sealed with an ultrasound-transparent Polymethylpentene (TPX) cover, embedded in an implant of dental cement. Animals could then recover for 1 week, with unrestricted access to food, water, and environmental enrichment.

fUS data were collected using an Iconeus One system (Iconeus, Paris, France) and a linear IcoPrime ultrasonic probe (15 MHz central frequency, 70% bandwidth, 0.110 mm pitch, 128 elements). Complex frames are acquired at 500 Hz. To remove tissue motion, a spatio-temporal clutter filter is applied using sets of 200 contiguous frames (Demene et al., 2015). The Power Doppler signal is then obtained by taking the mean power over these 200 frames, thus getting an effective 2.5 Hz sampling rate. The Power Doppler signal is approximately proportional to cerebral blood volume (Macé et al., 2011).

In each recording day, we imaged activity in response to the entire sound set sequentially in two coronal slices. The probe was angled at ~30-35 degrees relative to the vertical axis in order for it to be roughly parallel to the surface of the brain. Through several recording sessions, we covered primary to ter-tiary regions of auditory cortex. Coronal slices were 14 mm wide, and spaced 0.4 mm apart. The resolution of each voxel was 0.1 × 0.1 × ~0.4 mm (the latter dimension, called elevation, is dependent on the depth of the voxel).

Sound presentation

Ferrets were awake, head-fixed and passively listening to different streams of natural sounds: foregrounds in isolation, backgrounds in isolation, or mixtures of both.

Sound stimuli

To choose foreground vs. background sounds, we used a previously established method based on sounds' stationarity (Kell & McDermott, 2019; McWalter & McDermott, 2019). We reproduce here a description of this algorithm from McWalter & McDermott (2019).

To quantify the stationarity of real-world sound recordings, we computed the standard deviation of the texture statistics measured across successive segments of a signal, for a variety of segment lengths (0.125, 0.25, 0.5, 1, and 2 s). For each segment length, we performed the following steps. First, we divided a signal into adjacent segments of the specified duration and measured the statistics in each segment. Second, we computed the standard deviation across the segments. Third, we averaged the standard deviation across statistics within each statistic class (envelope mean, envelope coefficient of variation, envelope skewness, envelope correlations, modulation power, and modulation correlations). Fourth, we normalized the average standard deviation for a class by its median value across sounds (to put the statistic classes on comparable scales, and to compensate for any differences between statistics in intrinsic variability). Fifth, we averaged the normalized values across statistics classes. Sixth, we averaged the resulting measure for the five segment lengths to yield a single measure of variability for each real-world sound recording. Finally, we took the negative logarithm of this quantity, yielding the stationarity measure.

We then defined arbitrary thresholds to define foregrounds and backgrounds, so that both groups were well separated in the stationarity axis (Figure 1A). Sounds were taken from Kell & McDermott (2019); Norman-Haignere et al. (2015) or downloaded from online resources (see table 1 for the full list of sounds).

Sound design and presentation

Foreground and background streams were composed of sequences of 6 different sound snippets of 9.6 s duration, repeated twice, so that a same snippet would occur at two distinct times of the stream (Figure 1B). For each condition, 6 such streams were created by drawing randomly from a pool of 36 individual snippets. Foreground and background streams were presented in isolation and in mixtures. In mixtures, both streams were overlaid with a delay of 4.8 s so that a change, either of foreground or background, occurred every 4.8 s. Each foreground stream was presented with two different background streams, and vice versa. In this design, each foreground snippet was presented with 4 different backgrounds, and in two history contexts.

Recording sessions were composed of three "runs" in which two foreground streams and two background streams were presented in all possible combinations, each preceded by 20 s of silence to establish baseline activity. These runs were presented twice. The order of sounds within each run was fixed, but the relative order of runs was randomized for each recording session. A list of sounds used is available in table 1.

Sounds were presented at 65 dB SPL. In mixtures, foregrounds and backgrounds were presented with a similar gain (0 dB SNR). Each snippet was normalized by its root mean square prior to integration in the sequence.

Data analysis

Denoising and normalization

To correct for large movements of the brain or deformations throughout the session, we used NoRMcorre, a non-rigid motion correction algorithm developed for two-photon imaging (Pnevmatikakis & Giovannucci, 2017).

Data were then denoised with canonical correlation analysis (CCA) to remove components shared between regions inside and outside of the brain, as described in Landemard et al. (2021) with the difference

that data was not recentered beforehand.

We normalized the signal in each voxel by subtracting and then dividing by average baseline activity. Baseline activity was estimated during 20 s segments of silence occurring every ~20 min, before each run. The obtained value was called ΔCBV and expressed in percent changes in cerebral blood volume.

Cross-correlation matrices

For each voxel, we computed the Pearson correlation between the vectors of responses to sounds of a given category (either foregrounds, backgrounds, or mixtures) for two repeats, and with different lags. We then averaged these matrices across all voxels to obtain the cross-correlation matrices shown in figure 1E. In the rest of the analyses, we use time-averaged responses (2.4 to 4.8 s after sound change). In figure S1, we show similar cross-correlation matrices between responses to mixtures and either foregrounds or backgrounds, to confirm that background and foreground invariance are also stable in time.

We computed the voxelwise test-retest correlation by correlating the vector of time-averaged responses to sounds across two repeats, for each category (Figure 2B). We then focused on voxels with a test-retest correlation above 0.3 in at least one category (foregrounds, backgrounds or mixtures).

Voxel-wise correlations

For each voxel, we defined background invariance (resp. foreground invariance) as the noise-corrected correlation between mixtures and corresponding foregrounds (resp. backgrounds), across sounds. We excluded the first and last sound snippet of each block, in which either foreground or background was missing in half of the snippet (see design in figure 1B).

We used a noise-corrected version of Pearson correlation, to take into account differences in test-retest reliability across voxels. Specifically, the noise-corrected correlation should provide an estimate of the correlation we would measure in the limit of infinite data (Kell & McDermott, 2019). The noise-corrected correlation values were obtained using the following equation:

$$r_{noise-corrected}(\boldsymbol{x}, \boldsymbol{y}) = \frac{0.5 * (corr(\boldsymbol{x}_1, \boldsymbol{y}_1) + corr(\boldsymbol{x}_2, \boldsymbol{y}_2))}{\sqrt{corr(\boldsymbol{x}_1, \boldsymbol{x}_2) * corr(\boldsymbol{y}_1, \boldsymbol{y}_2)}} \tag{1}$$

where $\boldsymbol{x}_1, \boldsymbol{x}_2$ (resp. $\boldsymbol{y}_1, \boldsymbol{y}_2$) are two repetitions of $\boldsymbol{x}$ (resp. $\boldsymbol{y}$).

ROI analyses

We manually defined regions of interest (ROIs) by combining anatomical markers and sound responsiveness. We found CBV responses to be organized in distinct patches in auditory cortex (Figure 2A). Using anatomical landmarks (Radtke-Schuller, 2018), we defined boundaries between the different functional regions from primary (MEG, medial ectosylvian gyrus) to secondary (dPEG, dorsal ectosylvian gyrus) and tertiary (VP, ventral posterior auditory field).

Map display

For surface views, we averaged values over the depth of the slice. The transparency (alpha value) of each surface bin was determined by the number of voxels included in the average. Thus, white sections in maps correspond to regions in which no reliable voxels were recorded. All maps in the main figures show the left auditory cortex of ferret L. Maps for other ferrets are shown in figure S2.

Cochlear and spectrotemporal modulation energy estimation

Spectrotemporal modulation energy was calculated by measuring the strength of modulations in filtered cochleagrams (which highlight modulations at a particular temporal rate and/ or spectral scale) (Chi et al., 2005). Specifically, modulation strength was computed as the standard deviation across time of each frequency channel of the filtered cochleagram, using the same 400 ms windows as for fUSI data. The bank of spectrotemporal filters used was the same as in Landemard et al. (2021) (scales: 0, 0.25, 0.5, 1, 2, 4, 8 cyc/oct; rates: 0.5, 1, 2, 4, 8, 16, 32, 64 Hz). Modulation energy was then averaged in 10 frequency bins (centers: 27, 50, 91, 167, 307, 1032, 3470 Hz).

We then averaged modulation energy using the same temporal window as for fUSI sound-evoked data (2.4-4.8 s after sound change), after shifting it by -0.8 s to account for hemodynamic delays. We thus obtained a feature matrix (frequency * temporal modulation * spectral modulation) for each sound segment.

We fit a linear model using ridge regression to predict voxel responses from sound features. The feature matrix was z-scored as part of the regression. We used 6-fold cross-validation: we split the sound set in 6 folds (keeping blocks of temporally adjacent sounds in the same fold). One fold was left out and the others were used to fit the weights. The lambda hyperparameter was defined using an inner cross-validation loop. The fitted weights were used to predict responses to left-out sounds. We thus built a matrix of cross-validated predicted responses and applied the same analyses as were applied to the true data (Figure 4). Prediction accuracy was defined as the Pearson correlation between measured and cross-validated predicted responses (Figure S5).

To examine the fitted weights, we averaged the weights obtained in all cross-validation folds and multiplied them by the standard deviation of the feature matrix. To extract voxels' preferred value over one feature axis (e.g. frequency), we first averaged the weight matrix along the other feature axes (e.g. spectral and temporal modulation) and used the value of the feature for which the maximum was reached. For plotting the tuning maps and the analysis of voxels tuned to low or high rates, we only considered voxels with average prediction accuracy in left-out folds higher than 0.3.

Human analyses

To directly compare our results with those obtained in humans, we used the dataset from Kell & McDermott (2019). In brief, 7 female human subjects were presented with 30 natural foregrounds, 30 natural backgrounds, and mixtures of both. Foregrounds and backgrounds were determined based on a criterion of stationarity, similar to our study. For each subject, backgrounds and foregrounds were matched at random (so that the mixtures were different for each subject). Each sound was presented 4 times. The SNR (ratio of power of the foreground versus the background noise) was -6 dB (vs 0 dB in our case). To encourage attention to each stimulus, participants performed a sound intensity discrimination task on the stimuli. We used a similar criterion for voxel inclusion and conducted the same analyses as for ferrets. Ridge regression was performed using 10-fold cross-validation. For each subject, (foreground, background, mixture) triplets were kept together in the same fold.

Statistical testing

To test the difference between background and foreground invariance for each ROI, we ran a permutation test. We built the null distribution by randomly permuting foreground and background labels for each sound and computing invariance for each voxel, 1000 times. We then compared the median difference between background and foreground invariance across voxels, to the values obtained in this null distribution and used the proportion of null values higher than the observed difference to obtain our p-value.

To compute statistical significance between ROIs, we also performed a permutation test. We did

the following procedure for each pair of ROI (MEG vs. dPEG, dPEG vs. VP). We built a null distribution by shuffling ROI labels for voxels belonging to the pair of ROIs within each animal, 2000 times. We then compared the observed average across animals of the difference of median correlations between ROIs across animals to this null distribution and used the proportion of null values more extreme than the observed difference to obtain our p-value. We also computed a p-value for each ferret separately (see table 2). We also ran this test by first focusing on voxels tuned to high or low rates within each ROI.

To test for differences in invariance between voxels tuned to low (< 8Hz) or high (> 8Hz) temporal modulation rates within each ROI, we used a similar permutation test in which we compared the difference of invariance between voxels tuned to low vs. high rates to a null distribution in which we shuffled the tuning values across voxels within each animal and ROI.

To compare results between ferrets and humans, we performed Wilcoxon rank-sum tests between ferrets and human subjects for different metrics: median prediction accuracy across voxels, median noise-corrected correlation between measured and predicted invariance. For the latter, we also tested the difference to zero using a sign-rank test.

Resource availability

Data and code will be publicly released upon publication.

Acknowledgements

We thank Shihab Shamma and Sam V. Norman-Haignere for fruitful discussions, Alex J.E. Kell for providing and helping with the human data, Balkis Cadi and Lynda Bourguignon for technical and administrative support throughout the project. This work was supported by the Institut Universitaire de France, ANR-17-EURE-0017, and ANR-10-IDEX-0001-02, as well as an AMX doctoral fellowship to A.L.

Author contributions

Conceptualization, A.L., C.B., Y.B.; Methodology, A.L., C.B., Y.B; Formal Analysis, A.L.; Investigation, A.L.; Writing - Original Draft, A.L.; Writing - Review & Editing, A.L, C.B., Y.B.; Visualization, A.L; Supervision, Y.B.; Funding Acquisition, Y.B.

Declaration of interests

The authors declare no competing interests.

Bibliography

Bar-Yosef, O. (2007). The effects of background noise on the neural responses to natural sounds in cat primary auditory cortex. *Frontiers in Computational Neuroscience*, 1. https://doi.org/10.3389/neuro.10.003.2007

Bimbard, C., Demene, C., Girard, C., Radtke-Schuller, S., Shamma, S., Tanter, M., & Boubenec, Y. (2018). Multi-scale mapping along the auditory hierarchy using high-resolution functional UltraSound in the awake ferret. *eLife*, 7, e35028. https://doi.org/10.7554/eLife.35028

Carruthers, I. M., Laplagne, D. A., Jaegle, A., Briguglio, J. J., Mwilambwe-Tshilobo, L., Natan, R. G., & Geffen, M. N. (2015). Emergence of invariant representation of vocalizations in the auditory cortex. *Journal of neurophysiology*, 114(5), 2726–2740.

Chi, T., Ru, P., & Shamma, S. A. (2005). Multiresolution spectrotemporal analysis of complex sounds. *The Journal of the Acoustical Society of America*, 118(2), 887–906. https://doi.org/10.1121/1.1945807

Christensen, R. K., Lindén, H., Nakamura, M., & Barkat, T. R. (2019). White Noise Background Improves Tone Discrimination by Suppressing Cortical Tuning Curves. *Cell Reports*, 29(7), 2041–2053.e4. https://doi.org/10.1016/j.celrep.2019.10.049

Demene, C., Deffieux, T., Pernot, M., Osmanski, B.-F., Biran, V., Gennisson, J.-L., Sieu, L.-A., Bergel, A., Franqui, S., Correas, J.-M., Cohen, I., Baud, O., & Tanter, M. (2015). Spatiotemporal Clutter Filtering of Ultrafast Ultrasound Data Highly Increases Doppler and fUltrasound Sensitivity. *IEEE Transactions on Medical Imaging*, 34(11), 2271–2285. https://doi.org/10.1109/TMI.2015.2428634

Elgueda, D., Duque, D., Radtke-Schuller, S., Yin, P., David, S. V., Shamma, S. A., & Fritz, J. B. (2019). State-dependent encoding of sound and behavioral meaning in a tertiary region of the ferret auditory cortex. *Nature Neuroscience*, 22(3), 447–459. https://doi.org/10.1038/s41593-018-0317-8

Elhilali, M., Xiang, J., Shamma, S. A., & Simon, J. Z. (2009). Interaction between Attention and Bottom-Up Saliency Mediates the Representation of Foreground and Background in an Auditory Scene. *PLoS Biology*, 7(6), e1000129. https://doi.org/10.1371/journal.pbio.1000129

Elie, J. E. & Theunissen, F. E. (2015). Meaning in the avian auditory cortex: neural representation of communication calls. *European Journal of Neuroscience*, 41(5), 546–567. http://onlinelibrary.wiley.com/doi/10.1111/ejn.12812/full

Hackett, T. A. (2007). Organization and correspondence of the auditory cortex of humans and nonhuman primates.

Hamersky, G. R., Shaheen, L. A., Espejo, M. L., Wingert, J. C., & David, S. V. (2023). *Unexpected suppression of neural responses to natural foreground versus background sounds in auditory cortex*. https://doi.org/10.1101/2023.11.20.567922

Ivanov, A. Z., King, A. J., Willmore, B. D., Walker, K. M., & Harper, N. S. (2022). Cortical adaptation to sound reverberation. *eLife*, 11, e75090. https://doi.org/10.7554/eLife.75090

Kell, A. J. E. & McDermott, J. H. (2019). Invariance to background noise as a signature of non-primary auditory cortex. *Nature Communications*, 10(1), 3958. https://doi.org/10.1038/s41467-019-11710-y

Khalighinejad, B., Herrero, J. L., Mehta, A. D., & Mesgarani, N. (2019). Adaptation of the human auditory cortex to changing background noise. *Nature Communications*, 10(1), 2509. https://doi.org/10.1038/s41467-019-10611-4

Landemard, A., Bimbard, C., Demené, C., Shamma, S., Norman-Haignere, S., & Boubenec, Y. (2021). Distinct higher-order representations of natural sounds in human and ferret auditory cortex. *eLife*, 10, e65566. https://doi.org/10.7554/eLife.65566

Lindsay, G. W., Mrsic-Flogel, T. D., & Sahani, M. (2022). Bio-inspired neural networks implement different recurrent visual processing strategies than task-trained ones do. *bioRxiv*, 2022.03.07.483196. https://doi.org/10.1101/2022.03.07.483196

Macé, E., Montaldo, G., Cohen, I., Baulac, M., Fink, M., & Tanter, M. (2011). Functional ultrasound imaging of the brain. *Nature Methods*, 8(8), 662–664. https://doi.org/10.1038/nmeth.1641

Malone, B. J., Heiser, M. A., Beitel, R. E., & Schreiner, C. E. (2017). Background noise exerts diverse effects on the cortical encoding of foreground sounds. *Journal of Neurophysiology*, 118(2), 1034–1054. https://doi.org/10.1152/jn.00152.2017

McDermott, J. & Simoncelli, E. (2011). Sound Texture Perception via Statistics of the Auditory Periphery: Evidence from Sound Synthesis. *Neuron*, 71(5), 926–940. https://doi.org/10.1016/j.neuron.2011.06.032

McWalter, R. & Dau, T. (2017). Cascaded Amplitude Modulations in Sound Texture Perception. *Frontiers in Neuroscience*, 11, 485. https://doi.org/10.3389/fnins.2017.00485

McWalter, R. & McDermott, J. H. (2018). Adaptive and Selective Time Averaging of Auditory Scenes. *Current Biology*, 28(9), 1405–1418.e10. https://doi.org/10.1016/j.cub.2018.03.049

McWalter, R. & McDermott, J. H. (2019). Illusory sound texture reveals multi-second statistical completion in auditory scene analysis. *Nature Communications*, 10(1), 5096. https://doi.org/10.1038/s41467-019-12893-0

Mesgarani, N., Cheung, C., Johnson, K., & Chang, E. F. (2014). Phonetic Feature Encoding in Human Superior Temporal Gyrus. *Science*, 343(6174), 1006–1010. https://doi.org/10.1126/science.1245994

Miller, L. M., Escabí, M. A., Read, H. L., & Schreiner, C. E. (2002). Spectrotemporal receptive fields in the lemniscal auditory thalamus and cortex. *Journal of Neurophysiology*, 87(1), 516–527. https://doi.org/10.1152/jn.00395.2001

Moore, R. C., Lee, T., & Theunissen, F. E. (2013). Noise-invariant Neurons in the Avian Auditory Cortex: Hearing the Song in Noise. *PLoS Computational Biology*, 9(3), e1002942. https://doi.org/10.1371/journal.pcbi.1002942

Ni, R., Bender, D. A., Shanechi, A. M., Gamble, J. R., & Barbour, D. L. (2017). Contextual effects of noise on vocalization encoding in primary auditory cortex. *Journal of Neurophysiology*, 117(2), 713–727. https://doi.org/10.1152/jn.00476.2016

Norman-Haignere, S., Kanwisher, N., & McDermott, J. (2015). Distinct Cortical Pathways for Music and Speech Revealed by Hypothesis-Free Voxel Decomposition. *Neuron*, 88(6), 1281–1296. https://doi.org/10.1016/j.neuron.2015.11.035

Norman-Haignere, S. V., Long, L. K., Devinsky, O., Doyle, W., Irobunda, I., Merricks, E. M., Feldstein, N. A., McKhann, G. M., Schevon, C. A., Flinker, A., & Mesgarani, N. (2022). Multiscale temporal integration organizes hierarchical computation in human auditory cortex. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-021-01261-y

Norman-Haignere, S. V. & McDermott, J. H. (2018). Neural responses to natural and model-matched stimuli reveal distinct computations in primary and nonprimary auditory cortex. *PLOS Biology*, 16(12), e2005127. https://doi.org/10.1371/journal.pbio.2005127

Pnevmatikakis, E. A. & Giovannucci, A. (2017). NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. *Journal of Neuroscience Methods*, 291, 83–94. https://doi.org/10.1016/j.jneumeth.2017.07.031

Rabinowitz, N. C., Willmore, B. D. B., King, A. J., & Schnupp, J. W. H. (2013). Constructing Noise-Invariant Representations of Sound in the Auditory Pathway. *PLoS Biology*, 11(11), e1001710. https://doi.org/10.1371/journal.pbio.1001710

Radtke-Schuller, S. (2018). *Cyto-and myeloarchitectural brain atlas of the ferret (Mustela putorius) in MRI aided stereotaxic coordinates*. Springer. https://doi.org/10.1007/978-3-319-76626-3

Sabat, M., Gouyette, H., Gaucher, Q., Lopez Espejo, M., David, S. V., Norman-Haignere, S., & Boubenec, Y. (2025). Neurons in auditory cortex integrate information within constrained temporal windows that are invariant to the stimulus context and information rate. *bioRxiv*, 2025.02.14.637944. https://doi.org/10.1101/2025.02.14.637944

Saderi, D., Buran, B. N., & David, S. V. (2020). Streaming of Repeated Noise in Primary and Secondary Fields of Auditory Cortex. *The Journal of Neuroscience*, 40(19), 3783–3798. https://doi.org/10.1523/JNEUROSCI.2105-19.2020

Saderi, D., Schwartz, Z. P., Heller, C. R., Pennington, J. R., & David, S. V. (2021). Dissociation of task engagement and arousal effects in auditory cortex and midbrain. *eLife*, 10, e60153. https://doi.org/10.7554/eLife.60153

Santoro, R., Moerel, M., De Martino, F., Goebel, R., Ugurbil, K., Yacoub, E., & Formisano, E. (2014). Encoding of Natural Sounds at Multiple Spectral and Temporal Resolutions in the Human Auditory Cortex. *PLoS Computational Biology*, 10(1), e1003412. https://doi.org/10.1371/journal.pcbi.1003412

Schneider, D. & Woolley, S. (2013). Sparse and Background-Invariant Coding of Vocalizations in Auditory Scenes. *Neuron*, 79(1), 141–152. https://doi.org/10.1016/j.neuron.2013.04.038

Schwartz, Z. P. & David, S. V. (2018). Focal Suppression of Distractor Sounds by Selective Attention in Auditory Cortex. *Cerebral Cortex*, 28(1), 323–339. https://doi.org/10.1093/cercor/bhx288

Shamma, S. (2009). Auditory perception through the cortical analysis of spectrotemporal modulations. *The Journal of the Acoustical Society of America*, 125(4_Supplement), 2564–2564. https://doi.org/10.1121/1.4783707

Silva, A. C., Koretsky, A. P., & Duyn, J. H. (2007). Functional MRI impulse response for BOLD and CBV contrast in rat somatosensory cortex. *Magnetic Resonance in Medicine*, 57(6), 1110–1118. https://doi.org/10.1002/mrm.21246

Singh, N. C. & Theunissen, F. E. (2003). Modulation spectra of natural sounds and ethological theories of auditory processing. *The Journal of the Acoustical Society of America*, 114(6), 3394–3411. https://doi.org/10.1121/1.1624067

Souffi, S., Lorenzi, C., Varnet, L., Huetz, C., & Edeline, J.-M. (2020). Noise-Sensitive But More Precise Subcortical Representations Coexist with Robust Cortical Encoding of Natural Vocalizations. *The Journal of Neuroscience*, 40(27), 5228–5246. https://doi.org/10.1523/JNEUROSCI.2731-19.2020

Town, S. M., Poole, K. C., Wood, K. C., & Bizley, J. K. (2023). Reversible Inactivation of Ferret Auditory Cortex Impairs Spatial and Nonspatial Hearing. *The Journal of Neuroscience*, 43(5), 749–763. https://doi.org/10.1523/JNEUROSCI.1426-22.2022