CITEgeist: Accurate deconvolution of spatial transcriptomics with same-slide

proteomics reveals midkine as a secreted microenvironment modulator in ESR1

mutant breast cancer.

Alexander Chih-Chieh Chang [1,2,*], Brent T. Schlegel [1,2,*], Neil Carleton [1,2], Priscilla F.

McAuliffe [1,3], Hunter Waltermire [1,2], Steffi Oesterreich [1,4], Russell Schwartz [5,6], and

Adrian V. Lee [1,4,7,#]

[1] Women's Cancer Research Center, UPMC Hillman Cancer Center, Magee-Womens

Research Institute, Pittsburgh, PA, USA

[2] University of Pittsburgh, School of Medicine, Pittsburgh, PA, USA

[3] Department of Surgery, Division of Breast Surgical Oncology, University of Pittsburgh

School of Medicine, Pittsburgh, PA, USA

[4] Department of Pharmacology and Chemical Biology, University of Pittsburgh,

Pittsburgh, PA, USA

[5] Ray and Stephanie Lane Computational Biology Department, Carnegie Mellon

University, Pittsburgh, PA, USA

[6] Department of Biological Sciences, Carnegie Mellon University, Pittsburgh, PA, USA

[7] Institute of Precision Medicine, Pittsburgh, PA, USA

[*] This author contributed equally to this manuscript

[#] Corresponding Author.

Abstract

Background: Dysplastic tissue architecture in estrogen receptor-positive (ER+) breast

cancer across therapy-naïve and therapy-exposed cancer tissues presents unique

challenges in the analysis of spatial transcriptomics. Many tools for deconvolution are

developed on well-structured tissue architectures such as the 10x Genomics mouse

hippocampus dataset. Spatial transcriptomics analysis could offer valuable insights into

treatment response, but faces limitations in cellular resolution.

Methods: To address this problem, we developed CITEgeist, a computational tool for

spatial transcriptomic deconvolution using integrated proteomics data from the same

slide. Visium Antibody Capture technology was applied alongside our novel algorithm to

analyze the tumor microenvironment. We demonstrate the reliability of our method

using pre- and post-treatment samples from six breast cancer cases.

Results: Our approach revealed previously undetectable cellular interactions within the

tumor microenvironment. By taking an interoperable approach to software development

and grounding our algorithm in interpretable variables, we demonstrate how CITEgeist

deconvolution is not only accurate but robust enough to be directly used as input in

external analytical tools developed by other research teams. We then applied this

approach to a set of specimens from a prospective trial our group ran and further

validated the findings in a series of in vitro experiments as a demonstrated use case of

the utility, necessity, and flexibility of CITEgeist; and the potential of our method to

rapidly translate novel clinical samples to new biological insights.

Conclusions: CITEgeist addresses a critical technical gap in spatial multi-omics

analysis through an integrated, multi-disciplinary approach. This work demonstrates the

value of combining clinical, translational, and computational expertise to identify novel

mechanisms of treatment resistance, potentially transforming therapeutic strategies for

resistant disease.

Keywords: Breast cancer, Spatial transcriptomics, Proteomics, Tumor

microenvironment, ESR1 mutation, Elderly patients, Treatment resistance,

Computational deconvolution, Midkine signaling, Multi-omics

Background

Estrogen receptor-positive (ER+) breast cancer accounts for approximately 85% of new breast cancer diagnoses [1], with surgery and anti-estrogen therapy established as the first-line treatment [2]. How breast cancer changes post-treatment is of vital importance to disentangling how treatment resistance develops, and spatial transcriptomics offer a unique opportunity to dissect the mechanisms of cancer evolution in native tissue architectural contexts. Additionally, many techniques allow for the joint collection of proteomic information from the same slide, allowing integration of multi-omic informational landscapes, and hence, deeper insights.

However, no tool is currently capable of integrating this proteomic information for analysis, and there are also limitations in various single-cell reference-dependent deconvolution tools that have been the dominant design choice in the field.

Single-cell reference-based deconvolution methods have several disadvantages. First, it is costly to generate a single cell reference to match each spatial dataset, at approximately double the cost [3]. Second, external references or atlases are not always available, especially for heterogeneous post-treatment cancer samples, which in themselves are rare and difficult to collect routinely [4]. Third, biosamples such as core biopsies often provide too little tissue for single-cell sequencing [8], and, even if available, the biopsy tissue used for the reference may not capture the heterogeneity of the cancer slice used in the spatial transcriptomics. [5,6]. Fourth, there is a significant risk of data hallucination, where gene expression or cell-type abundance, as estimated from single-cell data, is artificially imposed in a spatial context, erroneously granting spatial properties to transcriptomic data when they are actually artifacts [7]. Finally,

many methods suffer from a lack of interpretability, requiring complex assumptions and

architectures to integrate the two informational landscapes of single-cell transcriptomics

and spatial transcriptomics [8], which often is incompatible with other downstream

analysis options and tools popular in the field, i.e., by failing to preserve raw counts

needed in DeSeq [9].

These technical limitations prompted the development of CITEgeist, a novel

computational tool that enables spatial transcriptomic deconvolution using integrated

proteomics data obtained from identical tissue sections via integer linear programming.

We demonstrate the accuracy of CITEgeist using a simulated reference test.

Additionally, we run this tool on post-treatment samples from a clinical trial examining

endocrine therapy in older women who forego surgery to emphasize the rapid cutting-

edge applicability of this method. Furthermore, we develop intrinsic tests on the non-

simulated real tissue data to confirm the confidence of CITEgeist outputs on post-

treatment tumor data. Lastly, we demonstrate how our output is fully compatible with

tools developed by other groups, and use them to identify a novel environmental effect

in ESR1 mutant breast cancer, which we then validate in cell culture in the laboratory.

Methods

CITEgeist Model

CITEgeist deconvolutes spatial transcriptomics data by leveraging antibody capture from the same tissue section, eliminating dependency on external single-cell RNA sequencing references. For the complete derivation, please see the Supplementary Methods. The method consists of three key components: (1) protein-based deconvolution using an Expectation-Maximization (EM) algorithm, (2) regional refinement with neighborhood optimization, and (3) spatial gene deconvolution with variable-sized neighborhoods.

Software

CITEgeist is implemented in Python 3.10 using Gurobi 11.0.2 for Integer Linear Programming (ILP) optimization. Data preprocessing includes gene filtering (> 0 counts in ≥ 1% spots, mean expression > 1.1) and normalization (target sum 10,000). Antibody data undergoes winsorization for the top and bottom 5% of data points, followed by global centered log ratio (CLR) normalization. Complete mathematical derivations and implementation details are provided in Supplementary Methods.

Single-Cell RNA-seq Reference

To establish a baseline for gene expression profiles — both for deconvolution tasks and the simulation of spatially resolved transcriptomic datasets — we utilized a publicly available atlas of single-cell transcriptomic data from human breast cancers, explicitly focusing on estrogen receptor-positive (ER+) tumors. This dataset served as the reference for deconvolution methods like Seurat [10] and cell2location [8], as well as the

foundation for our simulations, enabling us to generate in-silico spatial datasets with a

quantifiable ground truth of cell type proportions and gene expression levels.

We selected 12,000 cells from ER+ tumor samples across 11 ER+/HER2- cases in the

Wu et al. breast cancer reference dataset [11], maintaining cell type proportions through

random down-sampling from the primary atlas according to the annotations of 9 distinct

cell types. Due to the computational constraints of the scCube autoencoder framework

[12], we used this down-sampled version of the atlas as our primary reference for

spatial transcriptomic simulations.

For the benchmarking of reference-based deconvolution methods, we utilized two

distinct permutations of our reference, the first of which retained the complete set of

ER+ tumor cells from the 11 patients from the Wu et al. atlas [11]. This ensured that the

reference expression profiles for cell types were as comprehensive as possible,

avoiding potential biases introduced by down-sampling. We then performed the same

analysis using our second permutation of the reference, down-sampled from ~30,000 to

8,000 cells to reflect both cancer heterogeneity and a realistic simulation of a single-

sample scRNA-seq reference. By using both approaches, we ensured that our

benchmarking fairly evaluated the deconvolution performance without artificial

constraints on the reference profiles, while simultaneously reflecting realistic conditions

in our simulated experiments.

Spatial Transcriptomic Simulation

We utilized the scCube (v2.0.0) framework [12] to simulate Visium-like spatial

transcriptomic data from the downsampled ER+ scRNA-seq atlas. A Variational

Autoencoder (VAE) model with a hidden layer size of 128, a learning rate of 0.0001, and

10,000 epochs generated single-cell gene expression profiles for the nine major

reference cell types. Cells were generated based on the proportions of the annotated

cell types in the Wu atlas, with a batch size of 512, executed on a CUDA-enabled A100

GPU. The trained model was saved for future use.

To capture the spatial distribution of cells, each sample was modeled on a 50 × 50

spatial unit grid, divided into hexagonal Visium-like spots, with an average of five cells

per spot. This structure enabled the modeling of clustered and infiltrative patterns typical

of the breast tumor microenvironment. Clusters of cells representing various types were

assigned specific spatial parameters such as shape, twist, and scale, while infiltration

patterns were intentionally varied within and across clusters. Background cell types

were also incorporated, with proportions reflecting approximate tumor heterogeneity.

Spatial Antibody Capture Simulation

Spatial antibody capture data were simulated for each replicate by first defining two

simulated protein markers for each unique cell type in the dataset, resulting in a set of

cell-type-specific markers. For each simulated spatial spot, the proportion of each cell

type was retrieved, and the expected expression for each marker was calculated by

scaling the cell type proportion by a random factor drawn from a uniform distribution

between 20 and 50. Expression values for each marker were then simulated using a

negative binomial distribution with a mean proportional to the expected expression and

a dispersion parameter of 0.5. A 5% dropout rate was applied to the simulated

expression values, which set the expression to zero to simulate additional technical

noise. Low-level background expression was drawn from a negative binomial

distribution with lower-scale parameters for spots where a given cell type was absent. In

addition to the cell type-specific markers, nonspecific proteins were simulated across all

spots, drawing from a negative binomial distribution with a constant mean expression of

10. The final output was a matrix of simulated protein expression values for each

marker across all spatial spots.

To comprehensively evaluate deconvolution methods, we simulated distinct sets of

mixed and highly segmented populations, acknowledging that actual cancer data spans

a continuum between these two states. By showcasing performance across these

diverse tissue architectures, we offer a thorough assessment encompassing the full

spectrum of tissue heterogeneity. Our analysis emphasizes both population types: the

mixed populations, which are designed to more closely resemble complex tumor

microenvironments, and highly segmented populations, which have been foundational

in prior deconvolution method development.

In order to account for spatial variability, we generated five distinct spatial replicates for

each sample type ("high_seg" and "mixed"), each exhibiting similar yet unique spatial

patterns. This approach yielded spatial gene expression matrices at single-cell and

spot-level resolutions, where each spot's gene expression values reflect the aggregated

profiles of the cells within it. Additionally, we recorded the cell-type proportions per spot

to provide a ground truth for evaluating spatial deconvolution accuracy. These datasets,

featuring diverse spatial configurations and infiltration patterns, aim to offer a more

realistic foundation for assessing different deconvolution methods on tumor-like spatial

data.

Cell Culture

Establishment of MCF-7 and T47D mutant *ESR1* cell lines was previously reported [13].

Individual MCF-7 and T47D wild-type and CRISPR-edited D538G ESR1-mutant cell

clones were evenly pooled and maintained in Dulbecco's modified Eagle medium

(ThermoFisher 11965118) or RPMI (ThermoFisher 11875119), respectively, plus 10%

fetal bovine serum, and 1x penicillin-streptomycin (Sigma-Aldrich P4333) at 37°C in a

humidified incubator with 5% CO_2. For hormone treatment experiments, cells were

deprived of steroid hormones by placement in phenol red-free Improved Minimum

Essential Medium (Richter's Mod.) (Fisher Scientific MT10026CV) plus 10% charcoal

stripped serum (CSS). 17β-Estradiol (E2) (Sigma-Aldrich E2758-1G) was used at a final

concentration of 1nM for +E2 conditions. ELISA was performed using the Invitrogen

Human MDK ELISA Kit (EH319RB) according to manufacturer instructions.

Immunofluorescence

MCF-7 and T47D cells were plated in 8-well chamber slides (FisherScientific 08-774-26)

at 500 cells/well in full medium as described. After adhering to the wells for 24 hours,

the medium was changed to phenol red-free Improved Minimum Essential Medium

(Richter's Mod.) (Fisher Scientific MT10026CV) plus 10% CSS for 5 days, with the

medium changed 2 times per day. For +E2 conditions, 1nM E2 was added to the

medium for 24 hours. The cells were fixed in 4% PFA (FisherScientific 50-980-487),

permeabilized for 10 minutes in 0.2% Triton X-100 (Sigma-Aldrich X100-100ML) in PBS

(Fisher Scientific MT21031CV), then blocked for 30 minutes in 1% BSA (Sigma-Aldrich

A9647-500G), 10% goat serum (Sigma-Aldrich G9023), 0.1% Tween20 (Sigma P7949-

100ML) in PBS. Primary antibodies for anti-MDK (1:1,000 ThermoFisher MA5-32538)

and anti-E-cadherin (1:200 BD Biosciences 610182) were diluted in 1% BSA, 0.1%

Tween20 in PBS and incubated overnight at 4 C in a humidified chamber. The cells

were then incubated in secondary antibodies for anti-rabbit Alexa Fluor 647 (Invitrogen

A32733) and anti-mouse Alexa Fluor 555 (Invitrogen A32727) for 1 hour at room

temperature in the dark. Counterstaining was performed with Hoechst 33342

(ThermoScientific 62249) in PBS for 10 minutes, and cells were imaged on a Nikon AX

confocal microscope with 10 image z-stacks in a minimum of 3 locations per sample at

60x oil magnification. Quantification was performed with CellProfiler and ImageJ.

Code Availability

The CITEgeist package, including the tool itself, benchmarking scripts, and code for

simulating test datasets, is available on GitHub at

https://github.com/leeoesterreich/CITEgeist. The repository provides the complete

implementation for generating the spatially resolved CITE-seq data simulations and

evaluating deconvolution accuracy and gene expression (GEX) profile inference

performance. It includes workflows for benchmarking CITEgeist against state-of-the-art

methods and instructions for replicating the analysis and customizing simulations for

new datasets. Comprehensive documentation and example scripts are included to

ensure usability and reproducibility. A persistent version of CITEgeist will be made

available via FigShare at the time of publication (doi: 10.6084/m9.figshare.28385675).

Clinical Trial Sample Collection

As part of a prospective, pragmatic, hybrid de-centralized non-randomized clinical trial

(NCT05914792) [14] designed to evaluate circulating tumor DNA (ctDNA) levels as a

226 means to augment longitudinal monitoring of older patients with ER+ breast cancer

227 receiving primary endocrine therapy (pET), tumor tissue samples were collected at

228 baseline and at the time of surgical intervention [15] . In brief, patients aged 70 years

229 and older with ER+/HER2- non-metastatic breast cancer signed informed consent

230 (protocol approved by the University of Pittsburgh IRB under STUDY21100091 and

231 conducted through the UPMC Hillman Cancer Center under protocol 22-088) and chose

232 to forego upfront surgery in favor of pET. Because this trial focused on surgical de-

233 escalation, most patients did not undergo surgery during the study follow-up. However,

234 a subset of six patients, four responding and two progressing, based on imaging

235 assessment and ctDNA results, underwent surgery after 3-50 months of pET (12 total

236 specimens). We collected matched core biopsy and surgical specimens from these six

237 patients (12 total specimens) for this analysis, providing a use case for the CITEgeist

238 tool. To preserve patient anonymity, all samples are labelled HCC22-088-P#-S#, for

239 HillmanCancerCenter trial 22-088 -Patient#-Sample#. Sample number is chronological;

240 in all cases, the biopsy samples are one, and the surgical samples are 2.

241 **For specific statistical and sequencing methods, please refer to our attached**

242 **Supplementary Methods.**

Results

CITEgeist is uniquely suited for accurate deconvolution in mixed dysplastic cancer architectures

We developed CITEgeist, a computational method designed specifically for accurate cell-type deconvolution in heterogeneous cancer tissues, inspired by a unique opportunity from a clinical trial at our institution (Figure 1A). As illustrated in Figure 1B, our method integrates multiple data layers to achieve high-precision cell-type identification. The core algorithm implements a neighborhood-based approach that utilizes the same-slide proteomic and transcriptomic data to capture cellular context at local, regional, and global scales, providing a comprehensive view of the tumor microenvironment.

Unlike many existing tools, which are primarily validated on homogeneous tissues such as the mouse brain [8,16] we rigorously evaluated CITEgeist on cancer tissues with their characteristic heterogeneity. To this end, we generated a series of simulated datasets with varying degrees of cellular admixture and spatial segmentation (Figure 1C). These simulated benchmarks represent the complex architecture typical of dysplastic tissues, with intermingled malignant and non-malignant cells exhibiting different levels of differentiation. We then validate novel findings in cancer cell line models via ELISA and immunofluorescent imaging as outlined in Figure 1D.

Benchmarking using simulated data demonstrates that CITEgeist outperforms state-of-the-art algorithms in deconvoluting cellular proportions.

Using these simulated datasets, our comparative analysis revealed increased error in

the mixed datasets compared to the segmented datasets with state-of-the-art methods

such as Cell2Location and Seurat, both seeing increases in error from 0.08 RMSE to

0.167 RMSE and 0.10 RMSE to 0.133 RMSE for deconvoluting cellular proportions,

respectively, when comparing highly segmented test sets vs mixed test sets. CITEgeist

notably does not suffer from this drop, performing with a 0.08 RMSE in both

circumstances.

To test the sensitivity of different methods to the size of the single cell reference, we

also performed tests with reference datasets corresponding to single sample paired

single cell sequencing (8,000 cells) or atlas-level objects (30,000 cells). Based on this

test, RCTD was noted to have a severe reference dependency based on the

comprehensiveness of the reference dataset used. All methods were first run with the

comprehensive 30,000 cells, which in cost terms is about 4x the size of a standard

single-cell sequencing run, and then run with 8,000 cells sampled from that distribution

to capture what would be approximately as if a spatial sample had been submitted

jointly with a singular single-cell sequencing kit. Most methods had minimal alterations

in performance (Supplemental Files 2); however, RCTD showed deterioration under this

test in both text contexts. With RMSEs of 0.21 on average with the 8,000 cell reference,

compared to 0.05 with the 30,000 cell reference. This confirmed our hypothesis

regarding the difficulty in evaluating performance on real cancer data when there are no

clear guidelines on the necessary size and quality of single-cell references.

Tangram also notably and visibly proved a different issue of over-calling T-cells due to

their known over-representation in single-cell datasets [17] (Figure 2B). It reported a

JSD on average of 0.56 in the highly segmented dataset, and 0.53 in the mixed dataset,

compared to an average of 0.15 and 0.34 in all other methods.

CITEgeist deconvolutes gene expression counts and outperforms state-of-the-art

algorithms.

We then extended our benchmarking to evaluate the performance to deconvolute the

spatially-derived gene expression of each cell type, a key function of spatial

transcriptomics. At the time, only two methods had documented methods for

deconvoluting RNA signals per cell type from their model, Cell2Location and Tangram

(Figure 3).

In the mixed cancer-similar test set, CITEgeist had at least a 4-fold improvement over

the other methods with an average NRMSE of 0.04, while Cell2Location had an

average NRMSE of 0.16, and Tangram had an average NRMSE of 0.20 (Figure 3). We

show a 2- to 4-fold increase in performance in the highly segmented test set (CITEgeist:

0.04 average NRMSE, Cell2Location: 0.066 average NRMSE, Tangram: 0.17 average

NRMSE).

The magnitude of this significant performance improvement is visualized in Figure 3,

which quantitatively illustrates CITEgeist's accuracy metrics across different tissue

composition scenarios compared to existing methods.

Interpretability in CITEgeist output

A key issue in spatial deconvolution is also an over-reliance on simulated benchmarks

[18]. Hence, interpretability and verifiable results are key, and thus we developed

several intrinsically-validating tests for CITEgeist deconvoluted samples, as

demonstrated in Figure 4, which allow for confidence in CITEgeist output on a sample-

by-sample basis, independent of simulated testing or ground truth. Figure 4A illustrates

our deconvoluted cell-type distributions, which were subsequently reviewed in

conjunction with the underlying H&E image, demonstrating consistency in the results

and expected proportions; with cancer cell proportion predictions matching prominent

clusters of epithelial cells, fibroblasts matching fibrotic regions, and CD4 and CD8 T

cells in expected proportions consistent with the known biology of these cell types.

Additionally, to demonstrate the accuracy of the gene expression deconvolution, we

took the gene layer for the three most abundant cell types: macrophages, cancer cells,

and fibroblasts, and used the GSEApy package to assess against the CellMarker_2024

gene set, and showed excellent concurrence with 60%, 70%, and 60% gene set overlap

respectively, as well as 3/3 aligned cell type calls for the top most pathway.

These methods demonstrate how even in the absence of ground truth, CITEgeist

interpretability allows for excellent traceable results for downstream analysis.

CITEgeist preservation of spatial context and RNA informational landscape

allows for seamless integration with additional analysis tools

A key strength of CITEgeist is the interpretability of its output, facilitating seamless

integration with various downstream analytical tools due to its use of linear assignment.

Because we utilize strict integer assignments of RNA transcripts, we maintain what are

effectively raw counts for programs such as DeSeq2 [19]. In Figure 4C, we illustrate the

functional effect of CITEgeist deconvolution within the interoperable AnnData format. By

stacking the deconvoluted compartments vertically and maintaining integer counts, we

connect CITEgeist objects to COMMOT [19], a signaling prediction package designed

for spatial data. We validated the biological relevance of CITEgeist results by examining

all detected secretion signals and comparing them with established biological findings

(Figure 4). To validate the accuracy of this output, we conducted a correlation analysis

using the output from our samples for secreted signals and their assigned cell types, as

well as known secretion amounts from the Human Protein Atlas (Figure 4E). This

analysis showed a consistent correlation (Spearman Rho = 0.46, p-value = 2.86e-37).

Signaling analysis using CITEgeist deconvoluted spatial transcriptomics

identifies an upregulation in MDK signaling, confirmed with in-vitro experiments

After validating CITEgeist performance on simulated ground truth datasets and also

real-world samples for interpretability, we applied CITEgeist to analyze clinical samples,

focusing on a breast cancer case with an *ESR1 D538G* mutation initially identified

through pilot bulk-RNA sequencing and subsequently confirmed by droplet digital PCR

(ddPCR) (Supplementary Figure 1). This mutation was of particular interest, given the

clinical relevance of *ESR1* mutations, and our laboratory's previous research in this area

[13].

We have previously shown that *ESR1* mutant breast cancers express basal

cytokeratins, and using CITEgeist on spatial transcriptomics from a case with an *ESR1*

mutation, we confirmed the presence of basal genes in the Cancer Cell layer (Figure

5A). However, we were intrigued to note that these signals were not clustered in one

location among the cancer cells as one would expect from a single subclone (Figure

5B), despite exhibiting all of the same gene mutation signatures from Estrogene [20]

(Figure 5C) and known pathway alterations from our previous work (Figures 5D, 5E)

[13]. As one of the first spatial samples of *ESR1* mutant breast cancer in the native

breast microenvironment, we examined the signaling effect of these mutant breast

cancer cells. We thus employed the validated COMMOT signaling analysis on

CITEgeist-deconvoluted data, which revealed increased MDK, PTN, and MIF signaling

in the mutant regions (Figure 5F). MDK has recently been shown to have a role in

breast cancer , age, and tumorigenesis [21,22], and to validate our in silico findings, we

performed ELISA on media from *ESR1* D538G MCF7 cell lines, and conducted analysis

of the MDK immunofluorescence of MCF7 *ESR1* mutant cell lines. ELISA confirmed that

the mutant cell line secreted double the MDK in environments both with and without

estradiol (p < 0.0001) (Figure 5H). Additionally, the literature review identified that

midkine has a strong pericellular effect, with a significant proportion being retained near

the cell in non-cancerous settings [23]. Hence, we conducted image analysis of

immunofluorescence on the same cell lines, which showed that in the vehicle, estrogen-

deprived environment (akin to a treated patient), the mutant cell lines have

approximately double the midkine at the cell membrane (p < 0.001) as well as in the cell

as a whole (p < 0.01) (Figure 5I).

Discussion

Our study demonstrates that CITEgeist offers a robust computational approach for deconvoluting complex cellular architectures in cancer tissues. Here, we discuss the broader implications of our findings and contextualize them within the current landscape of spatial transcriptomics analysis.

CITEgeist avoids many of the difficulties in using single-cell references in heterogeneous cancer environments.

The integration of same-slide proteomics with our deconvolution method significantly enhances both interpretability and analytical robustness. By simultaneously capturing transcriptomic and proteomic profiles from the same tissue section, CITEgeist mitigates the technical variability inherent in comparing sequential slides, which is particularly valuable in heterogeneous cancer tissues where cellular composition can vary substantially across minimal spatial distances. This multi-modal approach provides internal validation and enables more confident cell-type assignment, especially in ambiguous cases where transcriptional profiles alone may be insufficient for accurate classification.

For example, we generated the reference size dependence test because Cell2Location has previously been demonstrated to be reasonably robust even in cases where the reference may not come from the exact same informational distribution [8], and hence wanted to replicate this test to test for the performance of each method in circumstances mimicking the effects of cancer heterogeneity. RCTD was noticeably challenged by this test, with a 4-fold increase in error – emphasizing the unique

challenge of applying tools benchmarked on common public normal tissue datasets to a

heterogeneous cancer dataset.

CITEgeist is designed with interoperability and clinical considerations for

maximum utility in analysis that allow for interpretable laboratory validation

A key contribution of our work lies in the methodology's capacity to not only confirm

existing biological knowledge but also to uncover novel insights during tool

development. The identification of upregulated MDK signaling in *ESR1*-mutated breast

cancer exemplifies how computational tools can drive biological discovery when

designed with interpretability as a core principle. This finding emerged from our

analytical pipeline rather than being a predetermined hypothesis, highlighting the value

of data-driven approaches in uncovering unexpected biological mechanisms.

The *ESR1* D538G mutation identified in our study has significant implications for

understanding tumor microenvironment dynamics. This mutation, occurring in the

ligand-binding domain of the estrogen receptor, is known to confer constitutive activity

and resistance to endocrine therapies. Our findings suggest that beyond its cell-

autonomous effects, this mutation influences the broader tumor ecosystem through

altered paracrine signaling, particularly via MDK. The increased MDK secretion

observed in mutant cells potentially modulates immune cell recruitment, angiogenesis,

and stromal remodeling, creating a microenvironment more conducive to tumor

progression and therapy resistance. This expanded understanding of mutation effects

beyond cancer cell-intrinsic properties represents an important shift in how we

conceptualize the consequences of genomic alterations.

Limitations

There are several limitations associated with CITEgeist. It is currently dependent on the variety and accuracy of the protein panel sequenced with spatial transcriptomics. It must be ordered together, which limits its utility in already sequenced datasets that lack protein information. However, as a first-in-class tool, it fills a necessary gap in the field where such samples already exist and have been sequenced, but researchers lack the tool to integrate proteomics and transcriptomics.

Secondly, our clinical trial has a small number of samples and is limited to breast cancer; hence, there may be other concerns or issues that emerge with other cancer types or tissue samples processed differently from other centers, and thus, careful usage and incorporation of feedback are critical to the long-term sustainability of the method.

Lastly, we also repeated our ELISA and imaging analysis in T47D, a different breast cancer cell line, which did not recapitulate the results (Supplemental Figure S3). However, previous work has established that the effects of these mutants are often context-specific [24,25]. It is important to note that breast cancer is a notably heterogeneous disease [26], hence, the purpose of our wet lab validation is not to find something true for all breast cancer, but to demonstrate that CITEgeist output can be directly input into external tools developed by others, and still provide analysis that can be confirmed in an isogenic model, proving that CITEgeist deconvolution provides a robust statistical foundation for further abstractions via interoperable tool usage. Further work is needed to establish the whole pathway that supports increased midkine in certain ESR1 mutant breast tumors.

Future Directions

Looking forward, several promising directions emerge from this work. First, the integration of temporal dynamics into spatial analysis would provide valuable insights into how signaling patterns evolve during disease progression and treatment response. Second, expanding our approach to incorporate additional data modalities, such as chromatin accessibility or metabolomic profiles, could further refine cell-type characterization and functional state assessment. Finally, the application of CITEgeist to larger patient cohorts across multiple cancer types would enable the identification of conserved and cancer-specific microenvironmental signatures, potentially revealing new therapeutic vulnerabilities.

Additionally, the computational framework established here could be adapted for other complex tissue architectures beyond cancer, such as in inflammatory conditions or developmental contexts, where cellular heterogeneity and spatial organization profoundly influence tissue function. The principles of neighborhood-based analysis and multi-modal integration are broadly applicable across diverse biological systems.

Conclusions

In this study, we present CITEgeist, a computational method designed explicitly for accurate cell-type deconvolution in heterogeneous cancer tissues. Our approach demonstrates superior performance in mixed cellular architectures compared to existing tools, particularly in the challenging context of dysplastic cancer tissues. The integration of neighborhood context at multiple scales enables robust cell-type identification even in regions of high cellular admixture.

The interpretability of CITEgeist outputs facilitates seamless integration with complementary analytical tools, enhancing flexibility in downstream analyses. This interoperability allowed us to leverage CITEgeist's deconvolution capabilities to identify and validate a novel biological finding: increased MDK signaling associated with the ESR1 D538G mutation, with potential implications for understanding endocrine resistance mechanisms.

By combining computational innovation with biological validation, our work contributes both a valuable methodological advancement and a specific mechanistic insight relevant to cancer biology. CITEgeist represents a step toward more comprehensive characterization of the complex cellular ecosystems in cancer, ultimately supporting the development of more effective therapeutic strategies that account for the full complexity of the tumor microenvironment.

List of abbreviations

ESR1 = Estrogen Receptor 1

CITEgeist = Cellular Indexing of Transcriptomes and Epitopes for Guided Exploration of Intrinsic Spatial Trends

MDK = Midkine

Declarations

Ethics approval and consent to participate

The protocol for the related clinical trial was approved by the University of Pittsburgh

IRB under STUDY21100091 and conducted through the UPMC Hillman Cancer Center

under protocol 22-088

Consent for publication

Not applicable.

Availability of data and materials

The Wu et al. breast cancer atlas can be accessed according to the instructions in their

original publication; the processed scRNA-seq data utilized in this study can be

downloaded through GEO (accession number: GSE176078, accessed 10/25/24). The

downsampled reference and simulated test datasets will be available via FigShare at

publication (doi: 10.6084/m9.figshare.28385675).

Visium datasets used in this paper will be published at the time of publication and are

currently embargoed. Reviewers can access the data via the link and private access

token provided to the journal in the cover letter at the time of submission.

All correspondence should be addressed to Adrian V. Lee at leeav@upmc.edu.

Competing interests

All authors have no competing interests to declare.

Funding

This research was partly supported by the University of Pittsburgh Center for Research

Computing (RRID: SCR 022735) through the resources provided. We utilized the HTC

cluster, supported by NIH award S10OD028483, and the H2P cluster, supported by

NSF award OAC-2117681. Additionally, this work was funded by US NIH Grant

R01HG010589, NIH Grant 5F30CA264963-03, and a Developmental Pilot Award from

the UPMC Hillman Cancer Center, supported through NIH grant P30CA047904.

Histology sectioning was performed in the Pitt Biospecimen Core (RRID: SCR 025229),

and the services and instruments used in this project were supported, in part, by the

University of Pittsburgh and the Office of the Senior Vice Chancellor for Health

Sciences. Work conducted at the UPMC Hillman Cancer Center Tissue and Research

Pathology Services (TARPS) Shared Resource Facility was also supported, in part, by

the University of Pittsburgh and the National Cancer Institute of the NIH under Award

Number P30CA047904.

Lastly, 10x Visium library generation, Takara Smart Stranded Total RNA library

generation, and Illumina sequencing were performed in the Health Sciences

Sequencing Core (RRID: SCR 023116) at UPMC Children's Hospital, Rangos Research

Center. These services and instruments were generously supported, in part, by the

University of Pittsburgh, the Office of the Senior Vice Chancellor for Health Sciences,

the Department of Pediatrics, the Institute for Precision Medicine, and the Richard K

Mellon Foundation for Pediatric Research.

Authors' contributions

A.C.C. conceived the project, designed the computational framework, wrote the CITEgeist code, conducted the bioinformatic analyses, and prepared the initial manuscript draft. B.T.S. performed the CITEgeist benchmarking, simulation studies, and statistical analysis and contributed to discussing benchmarking and simulation strategies. N.C. facilitated the collection of clinical samples and coordinated the clinical trial. P.F.A. served as the clinical trial's principal investigator and performed breast surgery, collecting the samples. H.W completed the wet lab validation of MDK signaling in ESR1 mutant cell lines. S.O. oversaw the overall project management and shaped the project's goals. R.S. contributed to manuscript revisions and provided expertise in mathematical modeling and computational methodologies. A.V.L. guided the project design and objectives. All authors reviewed and edited the manuscript.

Acknowledgements

We want to thank Jingyang Qian for their gracious guidance in utilizing the scCube method to simulate and benchmark spatially resolved Visium-like datasets from a single-cell reference. We also thank Jian Chen from the Lee-Oesterreich Lab for her help with RNA sequencing prep and ddPCR.

We acknowledge the assistance of the physicians and clinical support staff at the University of Pittsburgh Medical Center health system for their contributions to the clinical trial—special thanks to Natera Inc. for their support with the clinical trial NCT05914792.

Supplementary Files

Supplementary File 1: Full mathematical derivation of the CITEgeist method

Supplementary Files 2: Tables reporting the metrics of all compared and benchmarked methods.

Supplementary Files 3: Raw data from all wet lab experiments (ELISA and IF).

Supplementary Files 4: Supplementary Statistical and Sequencing Methods

References

1. Huang H, Wei T, Zhang A, Zhang H, Kong L, Li Y, et al. Trends in the incidence and
survival of women with hormone receptor-positive breast cancer from 1990 to 2019: a
large population-based analysis. Sci Rep. 2024;14:23690.

2. Patel R, Klein P, Tiersten A, Sparano JA. An emerging generation of endocrine therapies
in breast cancer: a clinical perspective. Npj Breast Cancer. 2023;9:1–12.

3. Regan C, Preall J. Practical Considerations for Single-Cell Genomics. Curr Protoc.
2022;2:e498.

4. Chang AC-C, Balic M, Bartholow T, Bhargava R, Brown DD, Brown L, et al. Hope for
OTHERS (Our Tissue Helping Enhance Research & Science): research results from the
University of Pittsburgh rapid autopsy program for breast cancer. Breast Cancer Res
[Internet]. 2025 [cited 2025 July 9];27. Available from:
https://link.springer.com/epdf/10.1186/s13058-025-02014-9

5. Wess M, Andersen MK, Midtbust E, Guillem JCC, Viset T, Størkersen Ø, et al. Spatial
integration of multi-omics data from serial sections using the novel Multi-Omics Imaging
Integration Toolset. GigaScience. 2025;14:giaf035.

6. Zaidi M, Fu F, Cojocari D, McKee TD, Wouters BG. Quantitative Visualization of Hypoxia
and Proliferation Gradients Within Histological Tissue Sections. Front Bioeng Biotechnol
[Internet]. 2019 [cited 2025 Aug 23];7. Available from:
https://www.frontiersin.org/journals/bioengineering-and-
biotechnology/articles/10.3389/fbioe.2019.00397/full

7. Chen J, Liu W, Luo T, Yu Z, Jiang M, Wen J, et al. A comprehensive comparison on cell-
type composition inference for spatial transcriptomics data. Brief Bioinform.
2022;23:bbac245.

8. Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, et al. Cell2location
maps fine-grained cell types in spatial transcriptomics. Nat Biotechnol. 2022;40:661–71.

9. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

10. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol. 2018;36:411–20.

11. Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A, et al. A single-cell
and spatially resolved atlas of human breast cancers. Nat Genet. 2021;53:1334–47.

12. Qian J, Bao H, Shao X, Fang Y, Liao J, Chen Z, et al. Simulating multiple variability in
spatially resolved transcriptomics with scCube. Nat Commun. 2024;15:5021.

13. Li Z, McGinn O, Wu Y, Bahreini A, Priedigkeit NM, Ding K, et al. ESR1 mutant breast
cancers show elevated basal cytokeratins and immune activation. Nat Commun.
2022;13:2011.

14. McAuliffe P. Longitudinal ctDNA Monitoring in Older Women With ER+ Breast Cancer
Who Forego Upfront Surgery in Favor of Primary Endocrine Therapy [Internet].
clinicaltrials.gov; 2025 Apr. Report No.: NCT05914792. Available from:
https://clinicaltrials.gov/study/NCT05914792

15. Carleton N, Chang AC, Chen F, Puhalla SL, Foldi J, Waltermire H, et al. Longitudinal
ctDNA Surveillance in Older Women with ER+ Breast Cancer to Facilitate Surgical De-
Escalation: A Prospective, Hybrid-Decentralized Trial with Correlative Studies [Internet].
medRxiv; 2025 [cited 2025 Sept 11]. p. 2025.08.23.25332468. Available from:
https://www.medrxiv.org/content/10.1101/2025.08.23.25332468v1

16. Biancalani T, Scalia G, Buffoni L, Avasthi R, Lu Z, Sanger A, et al. Deep learning and
alignment of spatially resolved single-cell transcriptomes with Tangram. Nat Methods.
2021;18:1352–62.

17. Li X, Wang C-Y. From bulk, single-cell to spatial RNA sequencing. Int J Oral Sci.
2021;13:36.

18. Li H, Zhou J, Li Z, Chen S, Liao X, Zhang B, et al. A comprehensive benchmarking with
practical guidelines for cellular deconvolution of spatial transcriptomics. Nat Commun.
2023;14:1548.

19. Anders S, Huber W. Differential expression analysis for sequence count data. Genome
Biol. 2010;11:R106.

20. Li Z, Chen F, Chen L, Liu J, Tseng D, Hadi F, et al. EstroGene2.0: A multi-omic database
of response to estrogens, ER-modulators, and resistance to endocrine therapies in breast
cancer. bioRxiv. 2024;2024.06.28.601163.

21. Visvader JE. Midkine links aging with breast cancer—A new predictor of cancer risk.
Cancer Cell. 2024;42:1815–7.

22. Yan P, Jimenez ER, Li Z, Bui T, Seehawer M, Nishida J, et al. Midkine as a driver of age-
related changes and increase in mammary tumorigenesis. Cancer Cell. 2024;42:1936-
1954.e9.

23. Muramatsu T. Midkine, a heparin-binding cytokine with multiple roles in development,
repair and diseases. Proc Jpn Acad Ser B Phys Biol Sci. 2010;86:410–25.

24. Arnesen S, Blanchard Z, Williams MM, Berrett KC, Li Z, Oesterreich S, et al. Estrogen
receptor alpha mutations in breast cancer cells cause gene expression changes through
constant activity and secondary effects. Cancer Res. 2021;81:539–51.

25. Li Z, Wu Y, Yates ME, Tasdemir N, Bahreini A, Chen J, et al. Hotspot ESR1 Mutations Are
Multimodal and Contextual Modulators of Breast Cancer Metastasis. Cancer Res.
2022;82:1321–39.

26. Guo L, Kong D, Liu J, Zhan L, Luo L, Zheng W, et al. Breast cancer heterogeneity and its
implication in personalized precision therapy. Exp Hematol Oncol. 2023;12:3.

Figure 1: Graphical abstract of the CITEgeist development process

Figure 1. (A) Graphical abstract describing the clinical trial tissue collection process. (B)
Graphical abstract describing the CITEgeist approach. (C) Graphical abstract describing
the complete simulation pipeline for benchmarking CITEgeist against other state of the art
single cell reference based spatial deconvolution methods. (D) Graphical abstract
describing the wet lab validation process.

Figure 2: Performance evaluation of CITEgeist and reference-based cell type
proportion deconvolution methods on simulated Visium datasets.

Figure 2. (A) Quantitative comparison of deconvolution accuracy across different spatial
patterns ("Highly Segmented" and "Mixed") and reference atlas sizes (30,000 or 8,000
cells). Metrics include Pearson's Correlation (Corr, higher values indicate better
performance), Jensen-Shannon Divergence (JSD), Mean Absolute Error (MAE), and Root
Mean Square Error (RMSE), with lower values indicating better performance for the latter
three error metrics. Error bars represent standard error (n = 5 independent replicates per
condition). (B) Representative deconvolution results from all benchmarked methods
showing cell-type proportion estimates across spatial spots for both Highly Segmented
(top row) and Mixed (bottom row) tissue patterns. Each spot is colored according to the
estimated proportions of nine cell types (legend at bottom). Ground truth (leftmost
column) shows the actual cell-type distributions, while subsequent columns display
predictions from each benchmarked method. Samples were selected from n = 5
independent simulations per condition to represent typical performance.

Figure 3: Metric comparison of gene expression (GEX) deconvolution per spot,

evaluated across all benchmarked methods and reference cell types.

Figure 3. Metric comparison of gene expression (GEX) deconvolution per spot, evaluated
across all benchmarked methods and reference cell types. Metrics include Pearson's
Correlation (Corr, higher values indicate better performance), Jensen-Shannon Divergence
(JSD), Mean Absolute Error (MAE), and Root Mean Square Error (RMSE), with lower values
indicating better performance for the latter three error metrics. Error bars represent
standard error (n = 5 independent replicates per condition).

Figure 4: Demonstration of CITEgeist validity when applied to real cancer data.

Figure 4. (A) Spatial plots showing the predicted proportions from CITEgeist output of the celltypes in sample HCC22-088-P1-S2 (B) CellMarker 2024 GSEApy calls, showing the top predicted celltypes for their respective 'assigned' gene expression layer. (C) Graphical abstract demonstrating how AnnData objects permit stacking of cell 'compartments' in the same adata spatial spot, and how COMMOT calculates using this information (D) Spatial plot showing an example of the detected sender and receiver signal for pathway MIF-CD74 CD44. (E). Global correlation regression plot showing the correlation between cell type assigned sender signal and said ligands expression in the Human Protein Atlas.

Figure 5: Wet lab validation of predictions based on CITEgeist output using external analytical packages.

Figure 5. (A) Spatial plots showing the heterogenous deconvoluted result of elevated basal cytokeratin expression in the (B) Cancer Cells layer in the surgical sample from the same patient in sample HCC22-088-P4-S2, which has an ESR1 D538G mutation. (C) Combined box and scatter plot showing the average ESR1 Mutant Gene Signature Score between ESR1 D538G mutant cancer cells and WT cells. (T-test value 3.1229, p-value 2.3206e−03 (D) Gene pathways upregulated in the D538G cancer spots from the MSigDB Hallmark 2020 gene set. Red text pathways are previously confirmed upregulated activities in ESR1 mutant cells from Li et al. [20] (E) Gene pathways upregulated in the D538G cancer spots from the KEGG 2021 Human gene set. Red text pathways are previously confirmed upregulated activities from Li et al. (F) Bar plot showing the statistically significant COMMOT sender signal differences between D538G cancer spots and WT cancer spots. A Mann-Whitney U test was performed to compare pathway-specific sender signals between cells with and without the D538G mutation. Error bars are SEM. The two-way test statistic and p-value were computed, and pathways were ranked by significance. The mean difference in signal between mutation-positive and mutation-negative cells was calculated. Benjamini-Hochberg FDR correction was applied to control for multiple comparisons, and pathways with FDR < 0.05 were identified as significantly different. (G) Quantification of midkine ELISA results from MCF7 wild type and with ESR1 D538G mutation. Statistical testing was done with one-way ANOVA with Tukey's post-hoc multiple

comparison (H) Image examples of MCF7 WT and ESR1 D538G with and without 24 hours
of E2 treatment after 5 days of E2 starvation. (I) Quantification of image examples
assessing the midkine signal at the cell membrane and for the total cell signal, with values
compared and tested via unpaired T-test. CellProfiler was used to assess values for each
cell after Z Max Projection across 5 ROIs per slide.

Supplementary Figures

Figure S1: ddPCR confirming ESR1 mutation in HCC22-088-P4-S2

Figure S1. ddPCR confirmation of D538G mutation in Sample HCC22-088-P4-S2 (A) DNA,
(B) RNA

83 **Figure S2: Runtime comparison of CITEgeist against other SOTA methods.**

84
Figure S2. Runtime comparison across all benchmarked methods, in both the "Highly
Segmented" **(left)** and "Mixed" **(right)** simulated conditions. Statistical significance was
calculated via ANOVA followed by Tukey's Honest Significant Difference test.

Figure S3: Quantification of midkine ELISA and immunofluorescence signals in T47D

Figure S3. (A) Quantification of midkine ELISA results from T47D wild type and with ESR1
D538G mutation. Statistical testing was done with one-way ANOVA with Tukey's post-hoc
multiple comparison. (B) Quantification of T47D midkine immunofluorescence assessing
the midkine signal at the cell membrane and for the total cell signal, with values compared
and tested via an unpaired T-test. CellProfiler was used to assess values for each cell after
Z Max Projection.