Spatial and single-cell expression analyses reveal complex expression domains in early

wheat spike development

Xiaosa Xu[1][*][‡], Huiqiong Lin[2, 3][*], Junli Zhang[2][*], German Burguener[2, 3], Francine Paraiso[2, 3], Kun

Li[2, 3], Connor Tumelty[2], Chengxia Li [2, 3], Yuchen Liu[1] and Jorge Dubcovsky[2, 3][‡].

[1] Department of Plant Biology, University of California, Davis, CA 95616, USA

[2] Department of Plant Sciences, University of California, Davis, CA 95616, USA

[3] Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA

[*] The first three authors contributed equally to this paper.

[‡] Co-corresponding authors

Co-corresponding authors

Jorge Dubcovsky, Email: jdubcovsky@ucdavis.edu

Xiaosa Xu, Email: xjkxu@ucdavis.edu

Emails from other authors:

Huiqiong Lin: hqlin@ucdavis.edu

Junli Zhang: psjzhang@ucdavis.edu

German Burguener: gfburguener@ucdavis.edu

Francine Paraiso: fjparaiso@ucdavis.edu

Kun Li: kzli@ucdavis.edu

Connor Tumelty: cwtumelty@ucdavis.edu

Chengxia Li: chxli@ucdavis.edu

Yuchen Liu: yvcliu@ucdavis.edu

ABSTRACT

Wheat is important for global food security and understanding the molecular mechanisms driving spike and spikelet development can inform the development of more productive varieties. In this study, we integrated single-molecule fluorescence *in situ* hybridization (smFISH) and single-cell RNA sequencing (scRNA-seq) to generate an atlas of cell clusters and expression domains during the early stages of wheat spike development. We characterized spatiotemporal expression of 99 genes by smFISH in 48,225 cells at the early transition, late double ridge, and floret primordia stages. These cells were grouped into 21 different expression domains, including four in the basal region of the developing spikelets and three different meristematic regions, which were consistent across spikelets and sections. Using induced mutants, we revealed functional roles associated with the specific expression patterns of *LFY* in intercalary meristems, *SPL14* in inflorescence meristems, and *FZP* in glume axillae. Complementary scRNA-seq profiling of 26,009 cells from W2.5 and W3.5 stages identified 23 distinct cell clusters. We used the scRNA-seq information to impute the expression of 74,464 genes into the spatially anchored smFISH-labelled cells and generated a public website to visualize them. We used experimental and imputed expression profiles, together with co-expression studies and correlation matrices, to annotate the scRNA-seq clusters. From co-expression analyses, we uncovered genes associated with boundary genes *TCP24* and *FZP*, as well as the meristematic genes *AGL6* and *ULT1*. The smFISH and scRNA-seq studies provided complementary tools for dissecting gene networks that regulate spike development and identifying new co-expressed genes for functional characterization.

Keywords

Wheat, spike development, single-molecule fluorescence *in situ* hybridization, single-cell RNA sequencing, spatial transcriptomics, co-expression analysis, gene expression, transcription factors.

BACKGROUND

In 2023, the world produced approximately 799 million tons of wheat grain [1], yet further increases are necessary to feed a growing human population. Wheat grains develop within inflorescences called spikes, thus improving spike productivity is an important breeding objective. Traditional and molecular breeding methods have made small incremental advances in spike productivity, but a deeper understanding of the regulatory mechanisms underlying spike development could accelerate the engineering of higher-yielding wheat varieties.

Wheat spike development starts with the transition of a shoot apical meristem (SAM) that produces leaves to an inflorescence meristem (IM) that produces spikelets. During the vegetative phase, the SAM produces lateral meristems that develop into leaves (Fig.S1A), whereas the subtended axillary meristems generally fail to grow out immediately [2]. By contrast, the wheat IM produces lateral meristems (Fig.S1B), that later differentiate into two distinct ridges (the double ridge stage, Fig. S1C). The lower ridges (also known as leaf ridges) are suppressed, whereas the upper ridges rapidly develop into spikelet meristems (SMs) that generate sessile spikelets attached directly to the inflorescence axis (rachis. Fig. S1D). Therefore, the transition from the vegetative to the reproductive phase involves both a drastic shift in the relative developmental rates of bract and axillary meristems and a change in the identity of the axillary meristems.

Wheat MADS-box genes from the *SQUAMOSA* clade (*VRN1*, *FUL2*, and *FUL3*) play an essential role in the repression of the lower ridge. Wheat plants with simultaneous loss-of-function mutations in these three genes show a normal transition of the SAM to the double ridge stage and stem elongation, but the lower ridges are derepressed and develop into leaves or bracts [3]. Bract outgrowth has also been reported in wheat plants with loss-of-function mutations in the *SQUAMOSA PROMOTER BINDING PROTEIN LIKE17* (*SPL17*) and *SPL14*, two regulators of the *SQUAMOSA* genes [4]. The maize orthologs of these genes (*TSH4*, *UB2*, and *UB3*) play a similar role [5, 6], suggesting a conserved function across grass species.

The wheat *SQUAMOSA* genes also play a critical role in the determination of the meristem identity of the upper ridge. In the absence of these genes, the upper ridges develop into vegetative meristems rather than SMs, resulting in a "spike of tillers" [3]. Taken together, these results indicate that the *SQUAMOSA* genes are essential for conferring SM identity to the

axillary meristem and suppressing bract development, but not for the initial transition of the

SAM to the double ridge stage and the subsequent stem elongation [3].

The MADS-box genes from the *SVP* clade (*VRT2, SVP1, SVP3*) play similar roles as the

SQUAMOSA genes during the initial stages of spike development. Individual loss-of-function

mutations in *VRN1* or *FUL2* [3], or simultaneous mutations in *VRT2* and *SVP1* [7], both result in

delayed heading, shorter stems and increased spikelet number per spike (SNS). However, at later

stages of spike development, the SQUAMOSA proteins transcriptionally repress the *SVP*-clade

genes to facilitate interactions with SEPALLATA1 proteins, which are required for normal

spikelet development [7, 8].

The normal development of the wheat lateral spikelets begins with the formation of two sterile

bracts called glumes, followed by slightly different bracts called lemmas that subtend the floral

organs. Differences between glumes and lemmas are regulated by the *APETALA2-like* genes

AP2L5 (*Q* gene) and *AP2L2*, which are cleaved by the microRNA miR172. Mutations in the

miR172 binding sites of these genes result in increased transcript levels and conversions of

sterile glumes into fertile lemmas [9, 10]. Conversely, loss-of-function mutation in *AP2L5* and

AP2L2 result in conversions of fertile lemmas into sterile glumes [11]. These results indicate an

important role of *AP2L5* and *AP2L2* in the regulation of glume and lemma organ identities and

the development of the axillary floret meristem (FM).

In grasses, the first whorl of floral organs generated by the FM is the palea, which is regulated by

the MADS-box gene *AGL6* [12]. Development of the internal floral whorls (lodicules, stamens,

and pistils) is severely disrupted in wheat plants with loss-of-function mutations in *LEAFY* (*LFY*)

or *WHEAT ORTHOLOGUE of APO1* (*WAPO1*), two genes that are expressed in partially

overlapping domains at the FM and encode interacting proteins [13, 14]. In addition to the

disruptions in floral development, loss-of-function mutations in either or both genes significantly

reduce the rate at which the IM generates lateral meristems, leading to drastic reductions in SNS

[13]. Natural alleles with increased SNS have been identified for both *WAPO1* [14-17] and *LFY*

[18], and allele combinations that maximize SNS have been also reported [18].

Despite progress in the genetic characterization of wheat spike development, the diversity of

expression domains within the wheat spike and their associated gene networks remain poorly

understood. In this study, we characterized the spatial expression of 99 wheat genes during the

initial SAM-IM transition, the late double ridge and floret primordia stages using single-

molecule fluorescence *in situ* hybridization (smFISH). To expand the list of characterized genes,

we integrated the smFISH data with single-cell RNA sequencing (scRNA-seq) from protoplasts

isolated during the late double ridge and floret primordia stages. These complementary datasets

provided complementary resources to study the gene networks regulating spike development and

revealed opportunities to engineer more productive wheat spikes.

RESULTS AND DISCUSSION

1. Spatial expression and cell segmentation

To identify the distinct cell clusters and expression domains involved in early wheat spike

development, we analyzed the expression of 99 genes simultaneously (Data S1) by smFISH at

three early stages of spike development (W1.5, W2.5, and W3.5 as defined by the Waddington

scale [19]). W1.5 marks the initial transition of the shoot apical meristem (SAM) to the

reproductive stage (Fig. 1A). W2.5 is defined as the late double ridge stage before glume

primordia initiation (Fig. 1B and S1A), and W3.5 as the floret primordia stage (Fig. 1C and

S1B). At W3.5, the inflorescence meristem (IM) begins its transition to a terminal spikelet

(IM→TS), and the most advanced spikelet meristems (SMs) at the center of the spike initiate

floret meristems (FMs, Figs. 1C and S1B).

We analyzed two sections from each developmental stage (Figs. 1, S2) using the cell

segmentation program QuPath [20] and identified a total of 48,225 cells. In the same sections,

cell walls were visualized with calcofluor-white (Figs. S3, S4). Based on the expression of 99

genes (Data S1), these cells were grouped into 21 clusters, represented by consistent colors

across all three developmental stages and replications (Figs. 1, S2). The individual cell clusters

are presented separately in Fig. S5 and can be also visualized, together with heatmaps of the 99

genes, at https://dubcovskylab.ucdavis.edu/JD99-wheat-spike-smFISH.

We characterized 7,624 cells at W1.5, 16,400 at W2.5, and 24,210 at W3.5 (Fig. 1 and S2),

showing a clear increase in cell number during spike development. These cells also exhibited

increasing differentiation, with 14 clusters at W1.5, 19 at W2.5, and 21 at W3.5 (Fig. 1 and S2,

Data S2). At the two later stages, cell clusters formed at the SMs and beneath them were

consistent across spikelets within the same spike and between sections in Figs. 1 and S2,

demonstrating good reproducibility. The gene expression profiles obtained with Molecular

Cartography also showed good reproducibility with previous *in situ* hybridization results (Fig.

S6) and recently published wheat spatial transcriptomics results generated using MERFISH [21]

(Fig. S7).

Fig. 1. Cell clusters based on the expression of 99 genes analyzed by smFISH at three stages of wheat spike

development. Cell clusters were generated from the three sections shown in this figure and replicate sections

presented in Fig. S2. Identical colors represent the same clusters across the six sections. **A** W1.5: initial transition

from the vegetative to the reproductive stage (section C1-1). **B** W2.5: late double ridge stage (section B1-3). **C**

W3.5: floret primordia stage (section A1-2). Scale bars = 100 μm. A detailed description of the cell clusters, cell

counts, and preferentially expressed genes in each cluster is available in Data S2. IM= inflorescence meristem, SM=

spikelet meristem, FM= floral meristem, TS= terminal spikelet.

Annotation of the cell clusters and their preferentially expressed genes are summarized in Data
S2, whereas the correlations among cell clusters are presented in Data S3. The five cell clusters
detected in the leaves (c1, c2, c5, c12, and c20) were highly correlated ($R \geq 0.78$, Data S3) and
are not discussed further in this study. To organize the description of these clusters and their
changes during spike development, we separated the developing spike into five regions: the basal
region below the spike, the spike central region, the region below the SMs, the SMs, and IM.

1.1. The basal region below the spike

The basal region below the spike consists of cells that will later differentiate into nodes and
internodes, forming the stem and contributing to important agronomic traits such as plant height
and stem strength. This region is primarily composed of cells grouped in cluster c0, which are
located below the epidermis and outside the vascular tissue and are referred hereafter as ground
tissue. At W2.5 and W3.5 this basal region is interrupted by alternating bands of cells from
clusters c4 (vasculature) and c10 (stem intercalary meristem, Figs. 1, S2, S5).

In longitudinal sections stained with calcofluor-white at W2.5 and W3.5 (Figs. S3, S4), two
alternating zones are visible at the basal region: one with shorter cells likely corresponding to the
nodes, and the other with longer cells likely corresponding to the internodes of the future stem.
Enriched expression of the procambial marker *LAX4* [22] and the phloem marker *APL* [23] was
observed in the smaller cells, supporting their association with future nodes (Fig. 2A). These
cells, together with those expressing the vascular genes in the leaf and spike vascular tissues
were all included in cluster c4 (Fig. 2B). Based on the location of c4 cells in the nodes and the
orientation of the vascular bundles we hypothesize that these cells may interconnect the parallel
vascular bundles of the internodes.

Cells classified as cluster c10 (Fig. 2B) appeared as incomplete bands in the future internode
regions at W2.5 and W3.5 (Figs. 1, S5). In the three-dimensional space, c10 cells may form a
transversal disc across the stem located at the point where the abaxial base of the leaves joins the
stem. These c10 cells were characterized by the differential expression of *LFY*, *PLATZ1* (Fig.
2A), *NAL1*, *ULT1*, and *SVP1* (Fig. S8, Data S2). Previous studies have demonstrated that loss-of-
function mutations in wheat *PLATZ1* [24] or *SVP1* [7] significantly reduced plant height, and
that overexpression of these genes partially restored normal plant height. The functional
validation of the roles of *PLATZ1* and *SVP1* in stem development suggested that cluster c10 may

correspond to the stem intercalary meristem, a zone at the base of grass internodes that

contributes new cells to the elongating internodes [25]. This hypothesis was further supported by

our finding that *LFY* also plays an important role in wheat stem development.

Fig. 2. Single-molecule fluorescence *in-situ* hybridization (smFISH) characterization of the basal region below the
spike at W3.5. Cell walls were stained with calcofluor-white. **A** Vasculature patterning genes *LAX4* (green) and *APL*
(red) are concentrated in the region with shorter cells (likely future nodes). *LFY* (purple) and *PLATZ1* (yellow) mark
a band in the region with longer cells (likely future internodes). **B** Cell segmentation with highlighted clusters c4 in
green (vascular tissue) and c10 in orange (stem intercalary meristem). Scale bar= 100 μm.

Using the *lfy*-null mutant developed in a previous study [13], we showed here that the mutant has

no significant effect on stem length, but resulted in significantly wider nodes (22.3% ± 1.8%)

and internodes (11.5% ± 2.0%). Moreover, the mutants produced more solid stems, showing a

62% ± 17% reduction of the diameter of the hollow central region relative to the WT (Fig. S9,

Data S4). The *Short and Solid Culm* (*ssc*) rice mutant is a *LFY* allele with a D16N amino acid

polymorphism and a 3-fold higher transcript levels than the wildtype, which has been associated

with reduced stem length and diameter [26]. These results revealed a shared role of *LFY* on stem

development between wheat and rice and are a good example of the value of the smFISH data to

generate functional hypothesis.

Since the gibberellic acid (GA) hormone plays a critical role in stem elongation, we examined

the spatial distribution of two genes involved in GA biosynthesis (*GA20ox1* and *GA20ox2*) and

one involved in GA degradation (*GA2ox1*). At W2.5 and W3.5, the two GA biosynthetic genes

were enriched in c0 at the basal region below the spike (Fig. S10B-C, Data S2). Loss-of-function

mutations in either *GA20ox1* or *GA20ox2* reduced the levels of bioactive GA_1 and GA_4, resulting

in shorter plants than the wildtype [27]. These findings support a positive role of these genes in

wheat stem elongation, consistent with their spatial expression adjacent to the stem intercalary

meristem region (c10).

The catabolic gene *GA2ox1* showed higher levels of expression in the central region of the

developing spikes at W1.5 (cluster c15) than at later stages (cluster c3, Fig. S10A, Data S2). We

speculate that the elevated *GA2ox1* expression at early spike development prevents premature

cell elongation and induction of *SOC1* and *LFY*, which are associated with increased GA levels

[28]. At W2.5 and W3.5, *GA2ox1* expression was restricted to the periphery of the region

enriched in GA biosynthetic genes, possibly forming a ring of reduced GA abundance around the

elongating cells. The negative role of *GA2ox1* in stem elongation has been functionally validated

in rice plants by both over-expression and ectopic expression of *GA2ox1* under the *OsGA3ox2*

promoter [29]. We hypothesize that the spatial distribution of GA metabolic and catabolic genes

described in this study may play an important role in normal wheat stem elongation.

1.2. The transition zone between leaf and spike

At W2.5 and W3.5, a group of cells within cluster c19 was detected in the transition region

between leaves and spike (Fig. S11A-B). The genes preferentially expressed in cluster c19

include the *SPL* transcription factors *SPL2*, *SPL14*, and *SPL17* (Fig. S11C-D), the *TCP*

transcription factor (*TEOSINTE BRANCHED1/CYCLOIDEA/PROLIFERATING CELL*

FACTOR) *TB1* and *TB2* (Fig. S11E-F), and the MADS-box genes *FUL2* and *SOC1-1* (Data S2).

Several of these genes have previously been associated with the regulation of spike architecture.

Mutations in the *SPL14* and *SPL17* genes result in the formation of ectopic bracts at the base of

the inflorescence in rice [30], maize [5] and hexaploid wheat [4]. Based on the basal location of

these bracts, we hypothesize that the preferential expression of *SPL14* and *SPL17* in cluster c19

may contribute to bract repression in the transition zone. *TSH4,* the maize ortholog of *SPL17* and

its close paralogues *UB2* and UB3 (*SPL14*) work together to target genes responsible for axillary

meristem suppression such as *TB1* and auxin response regulators [31].

In rice, *SPL14* is a direct regulator of *TB1* [32], which in our study was also preferentially

expressed in cluster c19. *TB1* reduces lateral branching, and mutants of this gene in maize [33]

and rice [34] show increased tillering. *TB1* mutants in barley and wheat also show increased

tillering and produce additional lateral spikelets in barley [35] and paired spikelets in wheat [36,

37]. These findings indicate a conserved role of *TB1* in the regulation of branching.

SQUAMOSA genes are direct targets of the SPL proteins [38, 39] and combined loss-of-function

mutations in the wheat *SQUAMOSA* genes *VRN1* and *FUL2* result in de-repression of the lower

ridge, leading to bract outgrowth below the spikelets at the base of the spike [3]. Individual *vrn1*

or *ful2* mutants show no bract outgrowth, demonstrating that *FUL2* and *VRN1* play redundant

roles in bract repression.

Based on the physical location of the c19 cells and the function of the genes preferentially

expressed in this region (Data S2), we hypothesize that the C19 cells contribute to the transition

between the proximal region (where leaf meristems exhibit rapid development and axillary

meristems undergo a temporary arrest) and the distal region (where the subtending bracts are

suppressed and upper ridges rapidly develop into spikelet meristems).

1.3. The central region of the spike

The ground tissue cells in the central region of developing spikes were primarily assigned to

cluster c15 at W1.5 and cluster c3 at both W2.5 and W3.5 (Figs. 1, S5), which were highly

correlated with each other (*R*= 0.70). These two clusters also showed strong correlations with the

ground tissue at the adjacent basal region beneath the spike (cluster c0, Data S3). Cells from

cluster c15 showed stronger correlations with the leaf mesophyll (*R*= 0.74) than cluster c3 (*R*=

0.46, Data S3), suggesting progressive differentiation of the ground tissue at the center of the

spike. Several genes were preferentially expressed at the spike central region across all three

developmental stages, including *AP3*, *bZIPC4* (Fig. S12A-C), *IDD4* and *OSH1* (synonymous

with *KNOX3*, Fig. S12D-F). The homeobox gene *OSH1* was expressed at the base and center of

the spikelets, but not in developing glumes and lemmas. This expression pattern resembles that of its orthologs in maize and rice [40].

In Fig. 3, we present the expression profiles of four vascular markers: the auxin influx carrier *LAX4* [41], the phloem marker *APL* [23], and the duplicated auxin efflux carriers *PIN1a* and *PIN1b* [22, 42] (Fig. 3). In *Brachypodium*, *PIN1b* plays a role in sink-finding and connecting new primordia to existing vasculature, whereas *PIN1a* forms narrower canals than *PIN1b* and is likely involved in vein patterning [42]. A similar pattern was observed in wheat, where *PIN1b* was expressed in a broader area than *PIN1a* (Fig. 3).

PIN1a and *PIN1b* were preferentially expressed in clusters c4 and c13 (Data S2, Fig. S13A), which were annotated as vasculature clusters. *APL* was among the preferentially expressed genes in both c4 and c13 (Data S2) but most of its expression was outside these clusters (Fig. S13B). The expression of *CEN5* overlapped well with c13 (Fig. S13B) whereas the expression of *LAX4* overlapped better with c4 (Fig. S13C). *PIN1a* showed extensive overlap with both clusters (Fig. S13D). The early accumulation of *PIN1a* and *LAX4* (=*ZmLAX2*) in maize leaves marked the initial procambium specification [22], suggesting that c13 and c4 represent early stages of vascular development.

In maize, *LAX4* and *PIN1a* have antagonistic functions and are frequently co-expressed in pro-vascular cells [22, 41], an overlap that was also apparent in wheat spikes at all three developmental stages (Fig. 3). The phloem marker *APL* was detected at developing vascular tissue of leaves at W1.5 and W2.5 (Fig. 3A-B), and in the central region of the spike at W3.5 (Fig. 3A).

In summary, an important function of the c4 and c13 vascular cells in the central region of the spike is to provide an efficient vascular system that transports water and nutrients to the developing spikelets. The central c3 cells are green from early developmental stages, suggesting that they contribute photosynthetic products to the developing spikes (see section 2.5) and also provide structural support to flowers and grains.

Fig. 3. Single-molecule fluorescence *in-situ* hybridization characterization of the spike central region (cluster c3). Spike central region (cluster c3). **A** W1.5, **B** W2.5, **C** W3.5. In all three panels, *LAX4* hybridization signal is shown in blue, *APL* in green, *PIN1a* in yellow, and *PIN1b* in red. Cell walls are stained with calcofluor-white.

1.4. Region proximal to the spikelet meristem (SM)

At W2.5 (Fig. 4A), cortex cells proximal to the spikelet meristem were primarily classified into

clusters c6 (suppressed bract), c9 (central SM basal region), and c16 (adaxial boundary). These

clusters remained evident at W3.5, with the addition of clusters c7 (glume and lemma) and c14

(glume axilla). A few cells from clusters c7 and c14 were also detected at W2.5 (Fig. 4A).

1.4.1. Cluster c6 "suppressed bract". Cluster c6 includes the suppressed bract region (Fig. S14).

This cluster showed a relatively high correlation with cluster c19 in the transition zone (R= 0.71,

Fig. S14 and Data S3), which was also associated with leaf ridge repression. Both clusters

showed preferential expression of *SPL14* (Fig. S14A-B)*, FUL2*, and *LFY* (Data S2), but cluster

c6 differed from cluster c19 by its preferential expression of *SPL13* (Fig. S14C-D) and lower

expression of *TB1* and *TB2*. In both c6 and c19 clusters, preferential expression of *SPL* genes

was associated with higher *VRN1* and *FUL2* expression (Fig. S14E-F). Combined *vrn1 ful2*

mutations resulted in bract outgrowth at the basal spikelets (Fig. S14G), suggesting a role of c6

and c19 cells in bract repression [3].

1.4.2. Cluster c9 "Spikelet base-center". Cluster c9 is located at the center of the region

proximal to the SM (Fig. 4) and showed high correlations (R= 0.75-0.80) with the surrounding

clusters c3, c7, c8, c14, and c16 (Data S3). This central cluster was enriched in *PIN1a* (Fig. 4A-

D), *PIN1b,* and *LAX4* (Data S2), suggesting a role in vascular development and the transport of

water and nutrients to developing spikelets. Additional genes preferentially expressed in c9

included *SOC1-1*, *SEP1-4, OSH1* and *TB1*.

1.4.3. Cluster c7 "Glume and lemma". The glume and lemma primordia, which became visible

at W3.5, were both included in cluster c7. Both organs showed preferential expression of *AP2L5*

(Fig. S6B), *FUL2* and *VRN1* (Fig. S14E-F) and reduced expression of *OSH1* (Fig. S12E) and

could be differentiated by the preferential expression of *SEP1-2* in lemmas and *SEP1-4* stronger

expression in glumes than lemmas (Figs. 4C-D, S15B). The latter result is consistent with

previous reports in rice [43]. Cluster c7 showed the highest correlation with the early meristem

cluster c8 (R= 0.81) from which glumes and lemmas originate, and also relatively high

correlations with the leaf cell clusters (R= 0.63-0.75, Data S3). This latter result supports the

hypothesis that glumes and lemmas are modified leaves [44]. Similar to leaves, glumes and

lemmas later turn green, contributing photosynthetic products to flowers and developing grains,

in addition to additional protection to the reproductive organs.

Fig. 4. Expression profiles of genes differentially expressed during spikelet development. **A-B, E-F** W2.5 (late double ridge stage). (**C-D, G**) W3.5 (floret primordia stage). **A, C** Cell segmentation and clustering. **B, D** Genes differentially expressed at the base of the SM. **E-G** Genes differentially expressed in the SMs. In **G**, the more basal SMs are at an earlier stage (cluster c8) than the central spikelets, which have already transitioned to the floret development stage (FM, cluster c18), and are already expressing floral homeotic genes *AG1* and *AG2*. *SEP1-4* marks the developing glume (stronger) and lemma, *SEP1-2* the developing lemma, *PIN1a* the developing vasculature, *SPL14* the suppressed bract, *FZP* the glume axilla and *TCP24* the adaxial boundary. Complete spike images showing expression of these genes are presented in Figs. S15-S18. SM= spikelet meristem, FM= floret meristem, le.= lemma, gl.= glume.

1.4.4. Cluster c16 "Adaxial boundary". This cluster was located adaxial to c9 (Figs. 4A-D, S15) and was defined by the differential expression of the transcription factor *TCP24* (Fig. S15, Data S2). *TCP24* and its closest paralogs *TCP22*, *TB1*, and *TB2* belong to the CYC/TB1 subclass of TCP transcription factors (Fig. S11E-F) and can form homo and heterodimers [45]. In the maize male inflorescence [46] and the *Brachypodium distachyon* spike [47], the orthologue of *TCP24, BRANCH ANGLE DEFECTIVE 1* (*BAD1*), is preferentially expressed in the pulvinus and affects branch or spikelet angle, respectively. The pulvinus is located at the axilla of the tassel branches in maize and at the spikelet axilla in *Brachypodium* but is absent in the *Triticeae species*. Loss-of-function mutations of *TCP24* in rice (=*OsTB2* = *RETARTDED PALEA 1*) are associated with mild reductions in the size of the palea with no other obvious effects on inflorescence or flower development [48]. By contrast, loss-of-function mutations in the barley ortholog *HvTB2* (also known as *COMPOSITUM1*) result in spikes with branches in the basal region [45], suggesting functional differentiation among grass species.

1.4.5. Cluster c14 "Glume axilla". Cluster c14 is defined by the expression of *FRIZZY PANICLE* (*FZP,* Fig. S15B, Data S2), which is expressed at the axillae of the developing glumes (Fig. 4C-D). A similar localization has been observed in previous *in situ* hybridization studies in rice and maize [49, 50]. In spike longitudinal sections, c14 cells frequently appear as two small cell clusters flanking c9, with the adaxial one adjacent to c16 (Figs. 4C-D, S15, S16). Cluster c14 also showed preferential expression of the homeobox domain gene *WOX7* (Fig. S16A) and the PINOID-like serine/threonine kinase *BIF2* (Fig. S16B), although these genes are also expressed outside of c14. Previous studies have shown that *FZP* plays an important role in the repression of the glumes' axillary meristem. Natural or induced amino acid changes in FZP result in the formation of branches at the base of the spikes in "miracle wheat" (*bh1* mutant) and barley (*com2* mutant) [51, 52].

To characterize the function of *FZP* in tetraploid wheat, we generated loss-of-function mutations in both *FZP* homeologs by CRISPR (Fig. S17). Eight independent T_0 lines with mutations in both *FZP* homeologs showed spikelets replaced by dense arrays of nested glumes (Fig. S17B and E). Dissection of these structures revealed new axes emerging from the axillae of the initial glumes, each producing multiple secondary glumes. The axillary meristems of these secondary glumes repeat the cycle but produce fewer and smaller tertiary glumes. The patterns repeat until the meristems were exhausted (Fig. S17F). Poorly developed floral organs were occasionally

observed among these recurrent glumes (Fig. S17G). The rachilla continued its development but

only vestigial florets were observed in its nodes (Fig. S17H). A diagram describing the

development of this structure is presented in Fig S17I, and the dissection of a wildtype spikelet

and the corresponding diagram in Fig. S17J-K. In some distal nodes, we observed lemmas with

longer awns subtending more normal floral organs. These results indicate that *FZP* is required to

suppress the axillary meristems in the glumes to ensure normal floret development. Similar

phenotypes have been described for *fzp* loss-of-function mutants in rice and maize [49, 50] ,

suggesting a conserved function in grasses.

In summary, the different expression domains detected in the spikelet region proximal to the SM

is consistent with recent grass inflorescence models proposing that the basal region of the grass

spikelets as an important signaling center for spikelet development [53].

1.5. Spikelet meristem (SM) and floret meristem (FM) regions

At all three developmental stages, cells in the IM and early SMs were assigned to cluster c8. At

W2.5, this cluster exhibited preferential expression of *HAF1*, *BIF2*, (Figs. 4E, S18A), *ULT1* and

WOX7 (Figs. 4F, S18C-D). *HAF1*, a C3HC4 RING domain-containing E3 ubiquitin ligase, has

no reported role in inflorescence development [54, 55], but the other three genes are well-

established regulators of meristem function. *BIF2* regulates auxin transport and is required for

axillary meristem initiation and lateral primordia development in maize and rice [56]. *ULT1*

encodes a SAND domain transcription factor that controls shoot and FM activity in Arabidopsis

[57, 58], whereas *WOX7* is a *WUSCHEL*-related homeobox gene that contributes to

inflorescence architecture in rice [59]. Based on their physical location and preferential gene

expression, we propose that c8 cells contribute to meristem function.

At W3.5, the more developed central spikelets showed the formation of floral meristems

(Fig.S1B) that were assigned to cluster c18 (FM, Figs. 1, S2). This cluster was defined by the

expression of floral homeotic genes *SEP3-1*, *SEP3-2* (Fig. S18E), *PI1*, *AP3* (Fig. S18F), *AG1*

and *AG2* (Figs. 4G, S18G, Data S2), strongly supporting a role in floral development. The SMs

are not visible in the sections of the central spikelets (Fig S1), either because the larger FMs

displaced the adjacent SMs from the section plane or because the section was not perfectly

centered (Fig. S1E).

Cluster c18 also showed preferential expression of *WAPO1*, a gene encoding a protein that interacts with LFY (Fig. S18H). Loss-of-function mutations in wheat *WAPO1* or *LFY* result in downregulation of floral homeotic genes and floral defects in wheat [13, 14] and other grasses [60-62].

The floral meristems in c18 are separated from the lemma primordia (c7) by a group of cells forming a distinct band assigned to cluster c17 (Figs. 1, 4C). This cluster was defined by the preferential and high expression of *LFY* (Fig. S18H, Data S2). A previous study suggested that the overlap between the *WAPO1* and *LFY* expression domains (Fig. S18H) provides critical spatial information for correct flower development in wheat [13]. The dynamic changes in gene expression during spike development are described later in section 1.7.

1.6. Changes within the inflorescence meristem (IM) during the transition to a terminal spikelet (IM→TS) and *SPL14* functional validation

In wheat, the rate of lateral meristem formation and/or the timing of IM transition to a terminal spikelet (IM→TS) determine SNS. Our previous studies showed that loss-of-function mutations in *LFY* or *WAPO1* reduce the rate of SM formation and SNS [13]. The reductions in SNS were comparable in the single and combined *lfy wapo1* mutants, suggesting that the LFY-WAPO1 protein complex is essential for establishing and/or maintaining a normal rate of SM generation. However, *LFY* and *WAPO1* are co-expressed in the IM only during the earliest stages of spike development (W1.5), suggesting that the early presence of the LFY-WAPO1 protein complex may induce a persistent change in IM function [13].

Since changes in the timing of the IM→TS transition can also influence SNS, we focused on genes involved in the early stages of this transition. We were unable to identify marker genes that clearly distinguished IMs from early SMs (both grouped in cluster c8). Because spikelets are "small spikes", SMs are also inflorescence meristem, making the identification of differential markers between IMs and early SMs more challenging. However, as SMs begin to differentiate into spikelets, an increasing number of genes become differentially expressed relative to the IM.

We manually delineated the apical meristem region and compared the hybridization density of the 99 genes in this region before the IM→TS (W2.5) and at the early stages of the transition (W3.5). These are within-gene comparisons and, therefore, were not influenced by probe

hybridization efficiency. For each stage, we analyzed four sections (four biological replications

for W2.5 and two for W3.5, each with two subsamples, see M&M). To verify that the selected

W3.5 sections had already initiated the IM→TS transition, we confirmed the expression of *FZP*

in the youngest lateral meristems (LM, Fig. S19). The selected region encompassed both the

terminal spike meristem (TM) and the two youngest LMs (henceforth, TM+2LM). At W2.5 the

TM corresponds to the IM and LMs to the SMs, whereas at W3.5 the TM corresponds to the SM

of the terminal spikelet and the LMs the future glumes. Using this approach, we previously

observed a significant decrease in *LFY* expression and a significant increase in *FUL2* expression

between W2.5 and W3.5 in the TM+2LM region [13].

In this study, we identified 11 additional genes with significantly altered hybridization densities

(spots/100 μm^2) in the selected region between W2.5 and W3.5 ($P < 0.05$, Fig. S20, Data S5).

Among them, *SPL14* exhibited a highly significant decrease in hybridization density during the

transition, (6.7-fold reduction, $P = 0.0031$, Figs. S19, S20, Data S5). We also identified two

additional MADS-box genes (*SOC1-2* and *SEP1-4*), four bZIP transcription factors (*bZIPC1*,

bZIPC3, *FDL2*, and *FDL6*), one bHLH transcription factor (*BC1*), one auxin efflux carrier

(*PIN1b*), and one receptor-like kinase (*CRN*) involved in the CLV-WUS negative feedback loop

(Fig. S20) that were significantly upregulated during the transition. Several of these genes have

been associated with inflorescence phenotypes. For example, mutations in *bZIPC1* in wheat

drastically reduce SNS [63], whereas mutations in the rice ortholog of *SEP1-4* (*OsMADS5*)

impact inflorescence branching [64].

To explore the phenotypic effect of reduced *SPL14* expression during the IM→TS transition, we

generated loss-of-function mutations in both *SPL-A14* and *SPL-B14* using CRISPR (Fig. S21A-

B). Plants with homozygous mutations for both homeologs showed a premature IM→TS

transition and a 42% reduction in SNS (Fig. S21C and E, Data S6). These results suggest that

SPL14 plays an important role in delaying the timing of the IM→TS transition in durum wheat.

In addition, the *spl14* mutant plants were 12 cm taller (Fig. S21D and F), headed two days later

(Fig. S21G), and produced 2.5 more leaves than the wildtype (Fig. S21H, Data S6), indicating

that *SPL14* plays additional roles. Similar effects on SNS and heading time have been previously

reported in *spl14* CRISPR mutants in hexaploid wheat, and natural alleles have been identified

with favorable effects on wheat grain yield [4]

Since SPL proteins directly regulate the expression of *SQUAMOSA* MADS-box genes, we

hypothesize that the reduced *SPL14* expression may be linked to the correlated increase in *FUL2*

and *VRN1* expression in the IM observed during the IM→TS transition. Supporting this

hypothesis, we previously showed that a mutation in a predicted *SPL*-binding site in the *VRN1*

promoter was associated with a significant increase in *VRN1* expression during early spike

development [65]. In summary, we identified multiple wheat genes involved in the early

IM→TS transition and validated the association between *SPL14* and the final number of

spikelets per spike.

1.7. Time and pseudotime changes in spike development

Spike and spikelet developmental changes over time were explored across three developmental

stages (W1.5, W2.5 and W3.5), along the longitudinal axis of the spike, and through trajectory

analysis (Fig. S22). At the early double ridge stage (W1.5), the IM and the upper ridge cells were

included in cluster c8 (early meristem) and a few cells in cluster c6 (bract repression, Fig. 1A

and S5). These two cell clusters expanded at the late double ridge stage (W2.5, Fig. 1B), and a

gradient of spikelet differentiation started to form along the longitudinal axis, with the most

advanced spikelets located at the center of the spike. This developmental gradient became more

pronounced at W3.5, when well-developed floral meristems were observed at the central

spikelets (Fig. 1C, clusters c17 and c18).

These dynamic changes were captured in a trajectory analysis (Fig. S22), which used cluster c8

(IM and early SM meristems) as the root. From c8, two trajectories diverged in opposite

directions: one leading to the formation of floral meristems (c17 and c18) and the other

branching towards the formation of the transition zone (c19) , the repressed bract (c6) and the

central zone (c3). A third trajectory diverged into three branches: one leading to the central

spikelet (c9), another to the glume and lemma primordia (c7), and a third one to the boundary

clusters c14 and then c16 (Fig. S22). Cluster c9 was also connected to the suppressed bract

trajectory (Fig. S22).

These predicted trajectories reflected many of the developmental changes observed during spike

development but also showed a few discrepancies. Although the c14 cluster appeared earlier than

c16 in the predicted trajectory, c16 cells were observed at the base of the developing spikelets

earlier than those assigned to c14, both across developmental stages and along the longitudinal

gradient of the spike (Fig. 1 and S2). A closer examination of Fig. S22 showed that multiple c16

cells connect directly to the meristematic region (c8), suggesting an alternative trajectory from c8

to c16 independent of c14.

The comparison of the different spikelets within a spike provided a precise view of the dynamic

gene changes leading to the formation of the FM. The meristematic markers *HAF1* and *ULT1*

were expressed in both early and advanced meristems, whereas *BIF2* and *WOX7* expression in

the meristems was restricted to the youngest SMs (Fig. S18B and D). In more advanced SMs or

FMs, *BIF2* and *WOX7* were detected in c14 and other clusters (Fig. S16). In SMs slightly more

developed than those expressing *BIF2* and *WOX7*, we detected the MADS-box gene *AGL6*,

which is known to play a role in paleae and lodicule development [12]. Finally, in the more

developed central spikelets, *AGL6* expression was restricted to a region between the lemma

primordia and the FM that overlapped with c17, and the FM showed expression of floral

homeotic genes and *WAPO1* (Fig. S18E-H).

2. Single-cell RNA sequencing (scRNA-seq) of the developing wheat spike

To identify additional genes regulating wheat spike development, we conducted single-cell

RNA-seq (scRNA-seq) using protoplasts isolated from *Kronos* spikes at the W2.5 and W3.5

stages (Fig. 5A). We optimized the cell wall digestion method to enhance meristem cell

recovery, following a previous single-cell study on maize inflorescence stem cell [66] (see

Methods). Raw scRNA-seq data are available in the NCBI Sequence Read Archive (SRA) under

BioProject accession number PRJNA1199502. Reads were mapped to the Kronos reference

genome v1.1 (annotation v1.0) [67].

In the initial analysis, we profiled 12,304 and 13,705 individual cells from three independent

replicates at W2.5 and W3.5 stages, respectively (Data S7). Good integration was observed

among both replicates and stages (Fig. S23A). We detected expression of 67,629 genes at W2.5

and 70,047 genes at W3.5 (Data S7), comparable to the number of genes detected in bulk RNA-

seq of Kronos spikes at the same developmental stages [68]. The mean and median unique

molecular indices (UMIs), the number of detected genes per cell, and other statistics are reported

in Fig. S23A and Data S7. Cell clustering of 26,009 cells from both stagers using Seurat v5 [69]

identified 25 distinct clusters in the initial analysis (Fig. S23A). Nine clusters were identified

with cells at either the G2/M or S phase, marked by *CYCB2* and *HIS2A1* expression, respectively (Fig. S23B-C).

To improve the resolution of biologically relevant clusters, cells from the nine cell-cycle clusters in the initial analyses were removed, and the remaining 17,283 cells were re-clustered into the final 23 clusters described in this study (sc0 to sc22, Fig. 5B). The final single-cell clusters were visualized using two-dimensional (2D) uniform manifold approximation and projection (UMAP) plots (Fig. 5B) and a 3D-UMAP (https://dubcovskylab.ucdavis.edu/content/kronos-scrna-3d-clusters). We also developed a Shiny App (https://dubcovskylab.ucdavis.edu/content/kronos-integrated-clusters) to visualize gene expression across clusters. To quantify the relationships among clusters, we calculated the average normalized gene expression across all cells for each cluster and used these expression profiles to calculate Pearson correlations for all cluster pairs (Data S8).

Fig. 5. Single-cell RNA-seq (scRNA-seq) analysis of wheat inflorescence spikes at the W2.5 and W3.5 stages.
A Experimental workflow, including wheat protoplast isolation and subsequent scRNA-seq. **B** UMAP plots showing

cell clustering and their annotations, **C** Cartoon with inferred positions of the different scRNA-seq clusters based on information discussed in the next sections. The cartoon is based on the same two central spikelets shown in Fig. 4C, allowing for direct comparisons with the smFISH clusters.

2.1. Characterization of scRNA-seq cell clusters and integration with spatial clusters

To annotate the scRNA-seq clusters, we identified and annotated genes preferentially expressed in each cluster (Data S9). We used this annotation to identify known marker genes (Data S1) and to perform KEGG pathway analyses (Data S10). We also incorporated spatial information into the scRNA-seq clusters using RNA-seq data from the basal, middle, and top sections of Kronos developing spikes at W3.0 (Fig. S24 and Data S11), as well as from correlations with published spatial transcriptomics clusters from hexaploid wheat (Data S13-15) [21].

To integrate the scRNA-seq and smFISH data, we used an imputation approach recently published in barley that incorporates scRNA-seq information into smFISH-labelled cells while retaining their spatial information [70]. Briefly, we analyzed the cells from both studies together using the 91 common genes and for each smFISH-labelled cell we identified the five closest scRNA-seq cells. We then used the average expression of these five cells (weighted by their distances) to impute the scRNA-seq expression into the smFISH-labelled cell (see Materials and Methods). We developed a web tool to visualize the spatial expression profiles of the 74,464 imputed genes in two spike sections at W2.5 and two at W3.5 (https://dubcovskylab.ucdavis.edu/imputed-genes). This tool greatly expanded the spatial information available to annotate the scRNA-seq clusters.

We used a similar approach to project the smFISH cluster ID into the scRNA-seq cells (see Materials and Methods). Since the number of genes in the smFISH clusters affected the number of identified cells, we divided the number of projected cells by the total number of cells in the smFISH clusters and generated a table including the "%-contribution" of each smFISH cluster to each scRNA-seq cluster (Data S12).

The different sources of information used to annotate the scRNA-seq clusters are summarized in Data S16 along with the final cluster annotations. The use in these annotations of similar names as the smFISH clusters indicates a better overlap than with other clusters and not a perfect one-to-one correspondence. The tentative physical locations of the scRNA-seq clusters are presented

561 schematically in Fig. 5C as a reference for the evidence discussed in the sections below, which
562 are grouped by similar functions.

2.2. Cell cycle enriched clusters

Although cells from the initial cell-cycle clusters were removed, four of the 23 final clusters
exhibited enrichment in cell-division marker genes combined with markers for other tissues. The
S-phase markers *HIS2A1-1* and *HIS2A1-2* were enriched in clusters sc0, sc16, sc19, and sc21
(Fig. S25). Cluster sc16 was also enriched in G2/M phase cyclin markers (*CYC7*), cyclin-
dependent kinases (*CDKB2*), and the *GROWTH REGULATING FACTORs GRF2*, *GRF4*, and
their cofactor *GIF1* (Fig. S25). Gene IDs and references for all marker genes are provided in
Data S1.

Among the genes preferentially expressed in cluster sc21 (Data S9) that exhibited significant
differences in the RNA-seq study, 65% were expressed at higher levels in the base of the spike
(Data S11-S16) suggesting a predominantly basal location for sc21. The smFISH cluster with the
highest contribution to sc21 was c3 (central spike) supporting the previous result (Data S12). The
basal and central region of the Kronos spike was characterized by a green coloration (Fig. 5A),
indicative of the presence of chlorophyll. This observation suggested that sc21 was enriched in
genes within the photosynthetic pathway, a hypothesis supported by the KEGG analysis (Data
S10). Based on these findings, cluster sc21 was annotated as "cell cycle / photosynthetic ground
tissue / basal region" (see section 2.5 for additional support).

Most of the genes preferentially expressed in sc0, sc16, and sc19 (Data S9) were expressed at
higher levels in the distal third of the developing spikes (Data S16). This enrichment was
strongest in sc0 (100% of the genes), intermediate in sc16 (78%), and weak but significant in
sc19 (47%). Based on its distal position and enrichment in meristematic markers (section 2.10),
sc0 was annotated as "cell cycle / meristematic / distal region". Among the smFISH clusters, the
one with the highest %-contribution to sc0 was c8 (early meristem) supporting the previous
result (Data S12). Cluster sc19 was also enriched in vascular markers (see section 2.4) , and
among the smFISH clusters projected into sc19 the highest value was for c13 (vasculature, Data
S12). Based on these results, sc19 was annotated as "cell cycle / vasculature / enriched in spike
tip". Finally, sc16 showed multiple epidermal markers (see section 2.3) and KEGG enrichment
in cutin, suberin, and wax biosynthesis (Data S10),which supported its annotation as "cell-cycle /

epidermis / enriched in spike tip". Additional information for sc16 is provided in the next section
on epidermal markers.

2.3. Epidermal clusters

A similar enrichment in cutin, suberin, and wax biosynthesis was detected in the KEGG analysis
for cluster sc15, supporting its annotation as an epidermal cluster. Clusters sc5, sc7, and sc13
were also annotated as epidermal clusters based on the preferential expression of known
epidermal markers, including *ONI* (synonymous with *KCS10*), *PEL3*, *SCR*, *ROC2*, *ROC3*, *ROC7*
(Data S1, Fig. S26). The Kronos imputed expression data confirmed the epidermal expression of
these six genes (Fig. S26H). Since no epidermal markers were included in our smFISH
experiment, these results exemplify the power of the imputation approach.

We compared the single-cell epidermal clusters to a recent hexaploid spatial transcriptomics
study that identified two expression domains, ED1 and ED4, associated with epidermal cells
[21]. These two epidermal expression domains had the highest correlations with sc7, sc13, sc15,
and sc16 (Data S13-15), further supporting the epidermal annotation of these clusters. However,
their correlations with sc5 were intermediate (Data S13-15). To identify markers that
differentiate the five epidermal clusters, we performed a separate comparison including only
these clusters and summarized the results in Data S17.

Epidermal clusters sc16 and sc7 were highly correlated (R= 0.79) and were both enriched in
genes with higher expression levels at the spike tip (sc7= 50% and sc16= 69%). These clusters
also showed high expression levels of ribosomal genes (Data S16), suggesting rapid cell growth
and division. Cluster sc16 showed preferential expression of multiple cell-cycle genes (Fig.
S25A-G, Data S17) and higher correlations with cell-cycle clusters sc0, sc19 and sc21 than all
other epidermal clusters (Data S8). Cluster sc16 was also enriched in meristematic markers
WOX3a and *WOX3b* (Data S17)*.* The smFISH cluster c8 (early meristem) showed the highest %-
contribution to sc16 relative to other smFISH clusters, supporting the previous result (Data S12).
Taken together, these results justify the annotation of sc16 as "cell cycle / epidermis / enriched in
spike tip".

Epidermal cluster sc7 was enriched in both meristematic and lateral organ markers. The enriched
meristematic markers included *AGL6* (Fig. S18B and S27A), and the imputed *WOX3b* (Fig.

S27J) and *CLE33* (Fig. S27K). Cluster sc7 was also enriched in *YAB3* and *YAB4* genes (Fig.

S27C, Data S17), which were detected at the lower ridge and lateral organs in hexaploid wheat

[21]. This expression pattern was validated in our imputed expression data (Fig. 27L).

Correlations between epidermal clusters sc7 and sc16 and the hexaploid wheat expression

domains ED4 (meristem epidermis) and ED1 (glume and lemma epidermis) [21] were the

highest in both directions and in all three developmental stages (Data S13-S15). In addition, the

smFISH clusters with the highest %-contribution to sc7 were c7 (glume and lemma) and c8

(early meristem) relative to other smFISH clusters (Data S12). Taken together, these results

support the annotation of epidermal cluster sc7 as "epidermis / meristem and lateral organs"

(Figs. 5C).

Epidermal cluster sc15 showed the second highest correlations with ED1/ED4 after sc7/sc16 in

hexaploid wheat (Data S13-15). However, sc15 differed from sc7 by its lower expression of

genes encoding ribosomal proteins and in its more uniform distribution along the spike (Data

S16). Among the markers genes identified for sc15, *SPL13* and *SPL14* (Fig. S26G and S27D-E)

suggest that sc15 may correspond to the abaxial base of the spikelets including the suppressed

bract region (Fig. S14A-D). This hypothesis is also supported by the higher correlation observed

between sc15 and the suppressed bract cluster sc1 ($R= 0.63$) than with meristematic clusters sc14

and sc20 (Data S8). Consistent with the previous result, the smFISH cluster with the highest

relative %-contribution to sc15 was c6 (suppressed bract, Data S12). Based on these results, sc15

was annotated as "epidermis / suppressed bract" (Fig. 5C).

Epidermal cluster sc13 showed a relatively high correlation with sc15 ($R= 0.77$) but differed by

its preferential expression of *TCP24*, a marker gene for the adaxial boundary domain (Fig. 4A

and C). Cluster sc13 was enriched in additional boundary markers such as *TCP22, CUC3, LAX1*,

and *DP1*, which have been found to be preferentially expressed in the boundary domain in

hexaploid wheat [21] and in our imputed expression profiles (Figs. S27M-P, see also later

section 2.12). The sc13 cluster sowed higher correlations with the hexaploid wheat expression

domain ED12 (boundary domain) [21] than with epidermal clusters ED1/ED4 and all other

clusters (Data S13-15). Relative to other smFISH clusters, c16 (adaxial boundary) showed the

highest %-contribution to sc13 (Data S12). Based on these results, sc13 was annotated as

"Epidermis / adaxial boundary" Fig. 5C.

Finally, sc5 was the most divergent of the epidermal cluster, showing the lowest correlation with

other epidermal clusters, the lowest expression levels of epidermal markers *SCR*, *PEL* and *ONI*

(Fig. 26G), and the largest number of differentially expressed genes (Data S17) with two

examples in (Fig. S27H-I). The smFISH clusters with the highest contribution to sc5 was c19

(transition zone), suggesting an enrichment in epidermal cells from the transition zone. However,

based on sc5 intermediate/high levels of protoplasting induced genes (Data S16), high

correlations with unknown clusters sc3 and sc22 (Data S8), and uniformly low correlations with

spatial clusters in hexaploid wheat (Data S13-15), we designated sc5 as "Epidermis / unknown".

2.4.Vascular clusters

The vascular genes *APL*, *LAX4*, *CEN5*, *PIN1a*, and *PIN1b* characterized by smFISH (Fig. 3)

were all preferentially expressed in sc18 and sc19, along with additional known vascular marker

SHR1, phloem markers *JUL1* and *DOF19*, and xylem markers *TMAAT, TMO5L1, TMO5L3,*

WAT1, and *XCP1*, *LOG10b*, and *NAC52* (Fig. S28, Data S1). The vascular location of the last ten

markers was validated in the Kronos imputed expression profiles (Fig. S28H), together with *APL*

and *LAX4* controls characterized also by smFISH (Fig. 3). The smFISH cluster with the highest

contribution to sc18 and sc19 was c13 (vasculature, Data S12). These results support the

annotation of sc18 and sc19 as vascular clusters (Fig. 5C).

Both vascular clusters shared relatively high expression levels of *OSH1* and genes encoding

ribosomal proteins (Data S16). However, the genes preferentially expressed in sc19 (Data S9)

were expressed at higher levels at the tip of the spike (Data S16), whereas those preferentially

expressed in sc18 (Data S9) were expressed at higher levels at the base and middle of the

developing spikes (Data S16). Since sc19 was enriched in *HIS2A* expression and sc18 was not

(Fig. S25), sc19 was annotated as "cell cycle / vasculature / enriched in spike tip" and sc18 as

"vasculature / enriched in spike base and middle".

Cluster sc18 showed two distinct regions: one preferentially expressing phloem markers (Figs.

S28B and D) and the other expressing xylem markers (Figs. S28C, E-F). To compare these

regions, we further divided sc18 into three subclusters: sc18_0 with the highest correlation to the

ground tissue cluster c3 (*R*= 0.625), sc18_1 including phloem markers *JUL1* and *DOF19*, and

sc18_2 including xylem markers *WAT1*, *XCP1*, *TMO5L1*, *LOG10b* and *NAC52* (Fig. S28H).

Genes differentially expressed in sc18_1 and sc18_2 are available in Data S18.

Consistent with the key role of hormones in vascular development [71, 72], sc18 showed

preferential expression of both homeologs of the auxin signaling genes *PIN1*, and *ARF11*, and

the JA biosynthetic gene *OPR3*. Cluster sc18 also showed preferential expression of the *CPS1*

gene encoding a GA biosynthesis enzyme, and genes *CKX-B3*, *CKX-B5*, and *CKX-B11*, all

involved in cytokinin degradation (Fig. S28I).

Interestingly, the floral homeotic genes *AP3* and *PI1* (Fig. S18F) were also detected in cluster

sc18 (Figs. S29A-B). A closer examination of the smFISH profiles revealed that these two

MADS-box genes were co-expressed with the vascular marker *APL* at all three developmental

stages (Figs. S29C-E). Expression of class-B MADS-box genes in the developing vasculature

has been reported in a few eudicot plant species [73]. Our findings extend this conserved

function to monocots.

2.5. Central spike clusters

In addition to vascular and pro-vascular cells, the central region of the developing spike contains

ground tissue cells, which correspond to cluster c3 (Fig. 1 central spike) in our smFISH study,

and expression domains ED3 (developing rachis) and ED5 (rachilla) in the hexaploid wheat

study [21]. These ground tissue cells showed a green coloration indicative of chlorophyll

presence, which was stronger at the base of the spike (Fig. 5A). The KEGG analysis indicated

that genes preferentially expressed in single-cell clusters sc6 and sc10 were enriched in

photosynthesis pathways (Data S10). Eighty two percent of the top 50 genes preferentially

expressed in sc6 were annotated with photosynthesis-related functions (Data S9), including

CAB2R and *LHCB1.1*, which were detected in the central spike region in hexaploid wheat [21]

and were validated in the Kronos imputed expression data (Fig. S30A-H). The expression levels

of these photosynthetic genes across all clusters are presented in a heat-map in Fig. S31.

The highest average enrichment in photosynthetic genes was detected for sc6 followed by cell-

cycle cluster sc21 (Fig. S31, Data S16). This finding supported the previous annotation of sc21

in section 2.2 as "cell cycle / photosynthetic ground tissue / basal region". Enrichment in

photosynthetic genes was also observed in sc10 and sc12, but at decreasing levels relative to sc6

(Data S16 and Fig. S31). A large proportion of the genes preferentially expressed in sc10 (44%)

and sc12 (36%) were also preferentially expressed in sc6 (Fig. S32), supporting the connections

among these clusters.

The genes preferentially expressed in sc6, sc10, and sc12 showed higher levels of expression at

the basal region of the developing spikes based on the RNA-seq data (Data S11-S16). This

location was further supported by the enrichment of *FWL3* and *SVP1* in both sc6 and sc10,

which are expressed in the central and basal parts of the spike (Figs. S32A-C and H). In addition,

both sc10 and sc12 were enriched in *OSH1*, a gene highly expressed in the central spike region

(Fig. S12). Cluster sc6 was enriched in additional genes expressed in the central spike region,

including *AP3*, *bZIPC3*, *FDL15*, *FT2*, *OSH1*, *SPL3*, and *VRT2* (Figs. S12 and S30), as well as

CAB2R and *LHCB1.1* both in hexaploid wheat and in our imputed expression data (Fig. S30H).

The smFISH cluster with the highest relative %-contribution to sc6, sc10 and sc12 was c3 (spike

central, Data S12), supporting a central spike location for the cells of these three clusters. Based

on the above information, clusters sc6 and sc10 were annotated as "central spike / photosynthetic

ground tissue / spike base", and sc12 as "tentative ground tissue / enriched spike base" (Fig. 5C).

2.6. Transition zone between leaves and spike

The transition zone between the leaves and the spike is represented by the smFISH cluster c19

(Figs. 1, S5), which shared several preferentially expressed genes with scRNA-seq cluster sc8.

These genes included *TB1*, *TB2*, *SPL17*, *GNI2*, *IDD10*, and *GA20ox1* (Fig. S33A) from our

smFISH study. This region was also enriched in *LEC1* and *TRD1*, two genes identified in the

hexaploid wheat data [21] and validated in our imputed expression data (Fig. S33B-F). Cluster

sc8 also showed high enrichment in jasmonic acid signaling genes *MYC2* and *JAZ8*, although

both genes were also expressed in other clusters (Fig. S33E-F).

Among the genes preferentially expressed in sc8, *SPL17* and its downstream target *TRD1* [30]

(ortholog of *OsNL1* and *ZmTSH1*) were previously shown to play important roles in bract

suppression in the transition zone. Loss-of function mutations in *SPL17/SPL14* in hexaploid

wheat [4], and its orthologs in rice [30] and or maize [5] were associated with bract outgrowth at

the base of the inflorescences. A similar phenotype was observed in *TRD1* mutants in barley

[74], rice *OsNL1* [75], or maize *ZmTSH1* [76]. This phenotype was consistent with the

significantly higher expression at the base of the spike of *SPL17*, *TRD1* (Data S11). Among the

genes in the RNA-seq experiment with significant differences among the three sections of the

spike, 66% of the ones expressed in sc8 showed higher levels at the base of the spike (Data S16).

Among the smFISH clusters, the one with the highest contribution to sc8 was c19 (transition

leaf/spike) followed by c3 (central spike, Data S12). Based on these results, sc8 was annotated as

"transition leaf-spike / spike base".

2.7. Suppressed bract cluster

Cluster c6, which includes the suppressed bract cluster, was consistently detected in the smFISH

study at the base of each developing spikelet (Fig. 1). This cluster showed strong differential

expression of *SPL13* and *SPL14* (Data S2), two markers that were also enriched in the scRNA-

seq cluster sc1 (Fig. S34) and that supported its annotation as "suppressed bract region" (Fig.

5C). This annotation was also supported by smFISH cluster c6 (suppressed bract) showing the

highest contribution to sc1 among the different smFISH clusters (Data S12). The suppressed

bract clusters c6 and sc1 also showed high correlations with the transition zone clusters c19 and

sc8, respectively (Data S3 and S8), probably reflecting their shared function in repressing the

leaf ridge.

The overlap between c6 and sc1 is likely incomplete. Cluster sc1 also showed a relatively high

contribution of smFISH cluster c3 (Fig Data S12), suggesting that it may include cells from both

the suppressed bract and ground tissue. Conversely, the c6 cluster includes epidermal cells (Fig.

4C-D), which were included in the "epidermal / suppressed bract" cluster sc15 in the scRNA-seq

study (section 2.3).

2.8. Boundary clusters

Among the genes analyzed in the smFISH experiment, *FZP* and *TCP24* showed the most

specific expression domains (Figs. 4, S15, Data S2). *FZP* was expressed in the glume axilla

(c14), while *TCP24* was expressed adjacent to *FZP* at the adaxial boundary region below each

spikelet (c16, Fig. 4C-D and S15-16). In addition to *FZP,* the hexaploid wheat spatial expression

data showed enriched expression in the adaxial boundary for the *WUSCHEL-RELATED*

HOMEOBOX9c (*WOX9c*, Fig. S35A), *TCP22*, *LAX1*, and *DP1* (synonymous with *BAF1*) [21].

We validated this expression profiles in the Kronos imputed expression data (Fig. S27M, O-P).

Mutations in wheat *MULTI-FLORET SPIKELET 1* (*MFS1*, synonymous with *DUO* and

ERF130) wheat have been previously associated with mild supernumerary spikelets and

increased grain number per spike, highlighting its role in wheat spike development [77]. This

motivated us to include *MFS1* in the smFISH experiment, which showed high levels of

expression in clusters c14 and c16, and lower expression in other clusters (Fig. S35B-C). The

high expression in c14 and c16 correlated with high *MFS1* expression in the scRNA-seq clusters

sc9 and sc11 (Fig. S35D), prompting us to investigate a potential overlap among these clusters.

Three lines of evidence supported the annotation of single-cell cluster sc11 as "glume axilla"

(Fig. 5C). First, glume axillary markers *FZP* and *WOX9c* were both preferentially expressed in

sc11 (Fig. S36). Second, c14 (glume axilla) showed the highest %-contribution to sc11 among

the smFISH clusters, (Data S12). Finally, RNA-seq results showed that sc11 was significantly

enriched in genes with higher expression in the center of the spike (Data S16), mirroring the

expression patterns of *FZP* and *WOX9c*.

We annotated the scRNA-seq cluster sc9 as "adaxial boundary" based on multiple shared

markers with smFISH cluster c16. *TCP24* was preferentially expressed in sc9 and was also

abundant in sc13 (Fig. S37), which includes the epidermal cell layer of the adaxial boundary

domain (section 2.3). Other markers preferentially expressed in the adaxial domain, such as

TCP22, LAX1, and *DP1* were also preferentially expressed in sc9 and/or sc13 (Fig. S37). The

smFISH cluster c16 (adaxial boundary domain) %-contribution to sc9 was more than two-fold

higher than the second highest (c9 central-base spikelet), which was also more than two-fold

higher than the other smFISH cluster (Data S12).Taken together, these results support the

annotation of sc9 as adaxial boundary domain.

2.9. Cortex cells below developing spikelets

In the wheat spike, the cortex consists of a layer of cells situated between the vascular tissues and

developing spikelets (Fig. 1). The parenchymatic cells in this region are expected to resemble

those in the central ground tissue. As shown in Fig. 4C, the cell layer between the vasculature

and the SM base includes ground tissue cells from the central cluster c3 as well as the innermost

cells from clusters c6 (suppressed bract), c7 (glume – lemma), c9 (SM base-center), c16 (adaxial

boundary) and c14 (glume axilla). Since cortical cells show no greening and are not

meristematic, the scRNA-seq clusters including these cells are expected to show low enrichment

in photosynthetic genes and reduced expression of *OSH1* or ribosomal protein-encoding genes.

Clusters sc2, sc4, and sc17 exhibited high correlations with the spatial clusters located proximal

to the developing spikelets in hexaploid wheat (ED0, ED3, ED5, and ED11, Data S13-S15) [21].

In addition, we found relatively high %-contributions of the smFISH clusters c16 and c6 to sc2,

from clusters c9 and c7 to sc4 and from cluster c16 to sc17 (Data S12). They also exhibited low

expression levels of photosynthetic and meristematic genes (Data S16) and uniform expression

along the spike based on the RNA-seq data. Taken together, these results suggest that sc2, sc4,

and sc17 may correspond to cortical cells below the spikelets (Fig. 5C).

The annotation of sc2 as a cortex cluster was further supported by the preferential expression of

LAX1 and *DEPRESSED PALEA1* (*DP1*) (Fig. S38), two genes previously shown to be

preferentially expressed in the adaxial boundary below the spikelets [78-81] and validated in the

Kronos imputed data (Fig. S27O-P). Two additional genes, *KAN1* and *RL9*, expressed in the

adaxial boundary region below the spikelet in hexaploid wheat [21], were also preferentially

expressed in sc2 (Fig. S38), and their imputed expression is presented in Fig. S38H. Additional

markers enriched in the correlated sc2 and sc13 clusters (Data S8) included *ABCG11* [82] and

SOT17 [83] (Data S9) and were validated in Fig. S38H.

A more central location of sc4, relative to sc2 and sc17, was supported by the higher %-

contributions of the smFISH cluster c9 (central region below spikelet) to sc4 than to the other

two spatial clusters (Data S12). This hypothesis was also supported by the 2.4 to 3.4-fold higher

expression of *OSH1* in sc4 than in sc17 and sc2, respectively (Figs. S12, S39, Data S16). In

addition, the cortex marker gene *NPF3.1* (also known as *NPF6.4-LIKE2* in maize [84]) was

enriched in sc4 (Fig. S39F-G), suggesting that this cluster corresponds to cortical cells.

Overall, the scRNA-seq cluster sc17 showed lower correlations with the spatial clusters in

hexaploid wheat than sc2, but they were still higher for the contiguous cortex clusters than for

the rest of the spatial clusters (Data S13-15). The highest %-contribution from the smFISH

clusters to sc17 was from c16 (adaxial boundary, adjacent to cluster, Data S12). We annotated

sc17 as "tentative cortex region below spikelet lateral regions" due to the lack of validation with

marker genes of known spatial distribution. Additional genes preferentially expressed in sc4 and

sc17 are listed in Fig. S39 and Data S9.

2.10. Meristem clusters

The RNA-seq clusters sc14 and sc20 showed preferential expression of meristematic genes

ULT1, *AGL6* and *SEP3-1* (Fig. S40A-D) suggesting that they included meristematic cells. This

hypothesis received additional support from the differential expression of the two homeologs of

ULT1. The differences between *ULT-A1* and *ULT-B1* were evident in the bubble plot in Fig.

S40G, in the UMAPS for the separate homeologs (Fig. S40H-I), and in the imputed expression

profiles (Fig. S40J-K). In the bubble plots and UMAPS, *ULT-A1* showed high expression in

meristematic clusters sc14 and sc20 (Fig. S40H) whereas *ULT-B1* did not (*ULT-B1* showed

differential expression in the central clusters sc6 and sc10, Data S9). These differences were

reflected in the imputed expression profiles, with *ULT-A1* expression concentrated in the early

(c8) and floral meristems (c17 and c18) and *ULT-B1* expression concentrated in the central part

of the spike (Fig. S40J-K). The correspondence between the spatial location and *ULT1*

homeologs and their scRNA-seq cluster assignment in the UMAPS and bubble plots supported

the conclusion that sc20 and sc14 include meristematic cells. More generally, the *ULT1* results

suggest that the expression imputation method developed by Demesa-Arevalo et al. [70] in

diploid barley, can be also used differentiate expression profiles between wheat close homeologs

in polyploid species.

The presence of meristematic cells in sc20 and sc14 was also supported by high levels of

expression of ribosomal genes (Data S16), intermediate expression levels of *OSH1* (Fig. S40G)

and by the higher %-contributions of the smFISH cluster c8 to both sc20 (1.5% = 33 cells) and

sc14 (5.3% = 114 cells) relative to other smFISH clusters (Data S12). Taken together these

results support the annotation of sc14 and sc20 as meristematic clusters.

AGL6 was detected in early meristems (c8) immediately before their transition to floral

meristems (Fig. S40L), after which it became more abundant at the base of the FM (c17, Fig.

S40L). This result is consistent with the role of *AGL6* role in paleae and lodicule development

[12]. *SEP3-1* showed a similar profile but its expression started slightly later than *AGL6* and was

detected in both the FM (c18) and its base (c17, Fig. S40M). Fig. S40N-P shows three additional

genes differentially expressed in sc14 (Data S9). *SEP1-2* is preferentially expressed in lemmas

and a few meristematic cells (*SEP1-2*, Fig. S40N and Fig. 4). The rice homolog of *SEP1-2*,

LHS1/OsMADS1, controls floret meristem specification [85]. The second gene, *JLO*, encodes a

LOB domain protein gene expressed in Arabidopsis at the base of the floral meristem (c17, Fig.

S40O) [86]. Finally, *FCP1* was expressed in the floral meristem (c17 and c18) and c14 at the

glume axilla (Fig. S40P). In rice, *FCP1* is a negative regulator of *WOX4* and of meristem

maintenance [87]. Based on these results sc14 was annotated as "meristem proximal region".

Cluster sc20 was enriched in genes differentially expressed in the distal one-third of the spike

(Data S16), so it was annotated as "meristem / enriched in spike tip".

Many of the genes enriched in sc14 and sc20, also show some enrichment in epidermal clusters

sc7 and sc16 (Fig. S40G, small arrows), supporting their previous annotation as "epidermis /

meristem and lateral organs". The imputed expression data showed enriched expression of

WOX3b and *CLE33* in c8 including the epidermal cells (Fig. S27J-K). Additional genes

preferentially expressed in sc14 and sc20 are listed in Data S9.

2.11. Protoplasting affected clusters

The last two scRNA-seq clusters, sc3 and sc22 (smallest cluster with only 163 cells), showed

evidence of being affected by the protoplasting. First, these clusters showed the highest

proportion of preferentially expressed genes that were not detected in RNA-seq studies of

Kronos developing spikes (this study and [68]). Approximately 19% of the preferentially

expressed genes in sc3 and 28% in sc22, were not detected at the same stages in Kronos spikes

under normal conditions (Data S16). The 235 genes detected only in the scRNA-seq were

enriched in stress-related GO terms (Data S19), further supporting their induction by

protoplasting. The preferentially expressed genes in sc3 and sc22 also showed a relatively high

overlap with a list of genes previously annotated as potentially affected by protoplasting [84]

(sc3= 11% and sc22= 13%, Data S16). The percentage of genes identified by these two criteria

across the 23 single-cell cluster showed a high correlation (R= 0.81), indicating a similar

underlying process. Cluster sc3 showed a high correlation with sc10 (R= 0.85, Data S8) and sc22

with cluster sc14 (R= 0.88, Data S8), suggesting that these clusters may contain cells related to

those clusters but affected by protoplasting. Based on these results, we annotated sc3 and sc22 as

"unknown".

2.12. Co-expression analyses

The generation of smFISH and scRNA-seq datasets from the same genotype, tissues, and

developmental stages provided the opportunity to identify additional genes co-expressed in

specific expression domains. We focused on four genes with specific expression domains —*FZP*

and *TCP24* in the boundary domains and *ULT1* and *AGL6* in the meristem— and calculated

Pearson correlations between each of them and all other genes across all cells in the scRNA-seq

dataset. The genes with *R* values exceeding four standard deviations above the mean (close to

zero as expected) were selected as co-expressed genes for further analyses (Data S20-S24).

2.12.1. Genes co-expressed with boundary genes FZP (c14) and TCP24 (c16)

The genes with the highest co-expression values for *FZP* and *TCP24* were their respective

homeologs, serving as positive controls (Data S20-S21). We found 608 genes significantly co-

expressed with *FZP* and 606 with *TCP24*, with 226 shared (988 total).

Among the shared co-expressed genes, we identified the well-studied gene *CUP-SHAPED*

COTYLEDON 3 (*CUC3*, Data S20-S21), a NAC-domain transcription factor involved in

meristem / organ boundary specification in rice [88] and Arabidopsis [89]. *In situ* and scRNA-

seq experiments in maize demonstrated that *CUC3* is expressed in both the *FZP* and *TCP24*

domains [81, 84, 90], a result validated by our imputed expression data (Fig. S27N). The *MFS1*

gene also showed a significant correlation with both *FZP* and *TCP24* (Data S20-S21) and was

co-expressed with both genes in the smFISH and imputed expression data (Fig. S35B-C).

Among the genes showing significant co-expression with *FZP* and not with *TCP24*, we

identified *WOX-9c* and *CLE33* (Data S20. Fig. S35). *WOX-9c* is a close paralog of *WOX7* [91,

92] with a similar expression profile as *FZP* (Fig. S35A). *CLE33* showed a similar expression

profile as *WOX7* (Figs S16 and S18C), changing from expression in the early meristems (Fig.

S27K) to an overlap with *FZP* in the more developed spikelets with FMs (Fig. S35E). In

Arabidopsis, RNA interference of *WOX9* in reproductive meristems resulted in the conversion of

flowers into inflorescence-like structures [93], resembling the conversion of spikelets into

branches observed in *fzp* natural mutants in wheat and other grass species [49-52]. The rice loss-

of function mutations in *WOX7* (*dwt1*) resulted in larger panicles, suggesting that these two

related *WOX* genes have functional roles in inflorescence development [59].

TCP24 showed high co-expression with *TCP22* (the close paralog of *TCP24*), *DP1*, and *LAX1*

(Data S21), three genes previously confirmed to be expressed in the adaxial boundary domain in

hexaploid wheat [21] and validated in our imputed expression data (Fig. S27M-P). *DP1* is an

ortholog of the maize gene *BARREN STALK FASTIGIATE1* (*BAF1*, Data S21), which is

expressed in a narrow domain adaxial to developing axillary meristems in maize [90]. Mutations

in *BAF1* resulted in the absence of ears or their fusion to the main stalk, supporting its role in

demarcating the boundary of the developing axillary meristem. *LAX1* is required for the initiation

of aerial lateral meristems and is expressed in the adaxial boundary region of the spikelet

meristems in both rice and maize (*BARREN STALK1*, *ZmBA1*) [78, 79]. A strong correlation was

also observed between *TCP24* and *RAMOSA2* (*RA2*, Data S21), which is preferentially expressed

in cluster sc9 in the imputed expression data (Fig. S35F). In maize, *RA2* is expressed in a group

of cells that predict the position of the axillary meristem formation [94]. Additional genes co-

expressed with *TCP24* are listed in Data S21. Based on the physical location of *TCP24* and its

co-expressed wheat genes, together with the known functions of the maize homologs, we

hypothesize that c16 cells contribute to delimiting the wheat SM and act as an important

signaling center for spikelet development [95].

To investigate the interactions centered on *TCP24* and *FZP*, we generated a gene regulatory

network using GENIE3 [96], focusing on the 988 genes co-expressed with *FZP* (Data S20)

and/or *TCP24* (Data S21) in the scRNA-seq dataset. From the top-500 significant interactions,

we constructed a network visualized in Cytoscape 3.10.3 (Fig. S42). The 95 genes included in

this network were all transcription factors (Data S22) suggesting that *FZP* and *TCP24* are part of

a broad regulatory network including many other transcription factors. The analysis revealed

connections between the *TCP24* homeologs, which share multiple targets, including their closest

paralog *TCP22*. *TCP24* also showed regulatory links with *LAX1* homologs and *RA2*, supporting

their coordinated expression in the adaxial boundary (Fig. S42, Data S22). Similarly, the two

FZP homeologs were connected and shared common links including *WOX9c*. Additionally,

TCP24 and *FZP* shared several downstream targets, such as the boundary regulator *CUC3*,

suggesting a potential functional convergence (Fig. S42, Data S22). This regulatory network

analysis provides a framework to investigate the genetic interactions that define the functions of

these boundary domains.

2.12.2. Genes co-expressed with meristem genes ULT1 and AGL6

The smFISH results for *ULT1* showed expression in the IM, SM and FM meristems as well as in

the central spike region (Fig. 4F and S18C-D). However, the co-expression analysis revealed

limited correlation between *ULT-A1* and *ULT-B1* homeologs (Data S23) supporting their

functional differentiation described in section 2.10 and Fig. S40H-K. Interestingly, among the

100 genes showing the highest correlations with *ULT-A1*, 82 encoded ribosomal proteins (Data

S23), suggesting a higher rate of protein production and rapid cell growth and division. This

enrichment was not observed for *ULT-B1*, supporting the conclusion that *ULT-A1* is the

homeolog predominantly expressed in meristematic cells. A heatmap based on the 45 ribosomal

genes with the highest correlation values with *ULT-A1* revealed that cluster sc20 exhibited the

highest expression level among the 23 single-cell clusters (Fig. S43). Cluster sc14 also showed

relatively high levels of ribosomal genes, but lower than those in sc20.

Co-expression analysis of *AGL6* revealed significant correlations with floral homeotic genes

SEP3-1, SEP3-2, *WAPO1*, *AG1*, and *PI1* (Data S24). Among the 100 genes showing the highest

co-expression values with *AGL6*, 43 encode ribosomal proteins (Data S24). A similar enrichment

of ribosomal protein encoding genes has been reported in a single-cell study of maize stem cells

[66]. This co-expression studies also revealed genes with previously uncharacterized roles in

inflorescence development (Data S23-S24). We explored the imputed expression of these genes

and prioritized three that were expressed in the spike meristematic regions for further functional

characterization: the remodeling genes *DDM1a* [97] (Fig. S41B) and *BTBN16* (Fig. S41C), and

the auxin response gene *SHI1* [98] (Fig. S41D).

2.13. scRNA-seq trajectory analysis

To summarize the relationships among the scRNA-seq clusters, we performed a trajectory

analysis (Fig. S44) starting from the early meristematic cluster sc14. In the lower region of the

graph, one branch of the trajectory connected sc14 with the vascular clusters sc18 and sc19 (and

the cell division clusters), and the other branch with the epidermal clusters (Fig. S44). The early

connection of sc14 with sc5 (epidermis unknown) and sc7 (epidermis meristem and lateral

organ) suggested that these clusters may correspond to early stages of differentiating epidermis

cells. The same trajectory later connected with sc15 (epidermis repressed bract) and sc13

(epidermis adaxial boundary).

In the opposite direction, sc14 was linked to two separate loops, each with multiple internal

connecting trajectories. The first loop included the ground tissue clusters sc6, sc10 and sc12 and

the transition zone sc8. The second interconnected loop included the cortex clusters sc2, sc4 and

sc17, and the related repressed bract cluster sc1 and adaxial boundary sc9. The glume axilla

cluster sc11 remained disconnected from the rest despite its proximity to cortex cluster sc4 and

epidermal cluster sc13. Overall, the predicted trajectories aligned relatively well with the

expected relationships among the different cell types derived from the meristematic cells in the
wheat spike.

3. CONCLUSIONS

The use of the same germplasm, tissues, and developmental stages in our smFISH and scRNA-
seq experiments allowed us to integrate the two datasets, which had positive synergistic effects.
The precise spatial information provided by the smFISH experiment was useful for improving
the annotation of the scRNA-seq clusters. However, the number of genes that can be included in
smFISH experiments is limited. We show here that this problem can be mitigated by using the
scRNA-seq data to impute the expression of a large number of genes into spatially anchored cells
from a spatial transcriptomics experiment. These imputed expression profiles can be used to
characterize the genes differentially expressed in each scRNA-seq cluster, thereby improving
their annotation. This imputation method can be also used to prioritize genes for functional
validation.

The method performed well both in diploid barley [70] and in tetraploid wheat, for which it
correctly predicted the locations of epidermal genes even when no epidermal markers were
included in the selected smFISH genes. The method also enabled us to distinguish expression
profiles between closely related wheat homeologs, confirming that this method can be useful for
studying expression in polyploid species with similar genomes.

The co-expression analyses, combined with smFISH and imputed expression profiles helped us
identify wheat genes with expression domains similar to those of their orthologs from rice and
maize, two crops with more extensive functional information than wheat. Comparative smFISH
studies of inflorescence development across different grass species using common sets of genes
can strengthen information bridges among crops. The analysis of co-expressed genes using gene
networks provided another layer of information on gene relationships in a cell-type dependent
manner. These localized networks can be useful for understanding the function of specific
expression domains such as the glume axillae and the adaxial boundary domain. Finally, these
tools can be used to identify genes with interesting expression profiles or interactions and
prioritize them for functional characterization.

In addition to providing useful datasets and tools, this study offered interesting biological insights into the function of *LFY*. For example, the discovery of *LFY* expression in the stem intercalary meristem motivated us to study its role in stem development and lead to the discovery of a previously unknown role of *LFY* in stem development. The smFISH experiment also revealed consistent and previously unknown expression domains in the region proximal to the developing wheat spikelets, which likely correspond to the "signaling centers" proposed in maize to regulate SM development and determinacy [95].

In summary, the integrated smFISH and scRNA-seq dataset provides a dynamic view of gene expression during the early stages of wheat spike development with improved spatiotemporal resolution. All data and tools generated in this study are publicly available to empower future studies of wheat spike development and support comparative genomic studies across grass species.

4. MATERIALS AND METHODS

4.1. Spatial Expression.

4.1.1. Sample preparation

Developing wheat spikes from the tetraploid wheat Kronos were collected at three developmental stages: early transition from the vegetative to the reproductive stage (W1.5), late double ridge stage before the development of glume primordia (W2.5), and floret primordia stage, when the central spikelets initiated the transition to the floral meristem (W3.5). The different tissue sections used for W1.5 and W2.5 correspond to different plants, but the W3.5 sections A1-1 and A2-1 correspond to the same spike and the sections A1-2 and A2-2 to a single spike from a different plant. Although the two sections from the same spike contain different cells, they were averaged and treated as technical replicates in the statistical analyses. After tissue collection, samples were fixed in 4% paraformaldehyde (PFA), dehydrated, and embedded in paraffin. Longitudinal sections (10 μm-thick) from the central plane of the developing spikes were positioned within the capture areas of the slides. The slides were dried overnight at 37°C, followed by a 10-minute bake at 50°C to enhance section adhesion. The sections were then deparaffinized, permeabilized, and refixed. After complete dehydration, they were mounted

using SlowFade-Gold Antifade reagent, covered with a thin glass coverslip, and shipped on dry ice to Resolve BioSciences (Monheim am Rhein, Germany) for analysis.

Spatial transcriptomics analyses were performed using Molecular Cartography^{TM} and a 100-plex combinatorial single-molecule fluorescence in-situ hybridization. Upon arrival, tissue sections were washed twice in 1x PBS for two minutes, followed by one-minute washing in 50% and 70% ethanol at room temperature. Ethanol was removed by aspiration, and DST1 buffer was added followed by tissue priming and hybridization. Tissues were primed for 30 minutes at 37°C, followed by hybridization for 48 h using probes specific to the target genes. The following day, samples were washed to remove excess probes and fluorescently tagged in a two-step color development process. Regions of interest were imaged as described below, and fluorescent signals were removed during decolorization. Color development, imaging, and decolorization were repeated across multiple cycles to generate a unique combinatorial code for each target gene, derived from raw images as described below.

4.1.2. Probe design

We used our published RNA-seq data [68] to select genes expressed at appropriate levels at the three selected spike developmental stages and prioritized those that showed dynamic changes across these stages. We also prioritized MADS-box genes and other transcription factors previously shown to be involved in inflorescence or floral development in other plant species. Finally, we also included marker genes to facilitate cluster annotations.

For each of the selected tetraploid wheat genes, we selected the homeolog with higher expression based on published RNA-seq data for Kronos spike development [68]. We also provided Resolve Biosciences with the sequences of the lower-expressing homeolog to exclude them during probe specificity evaluations conducted against all the coding sequences of the wheat genome (Ref Seq v1.1, Data S1). Therefore, although probes were designed based on the sequence of the highest-expressing homeolog, they are not genome-specific and can detect both homeologs for each gene. Probes were compared with all the wheat expressed genes, and those that could amplify other genes were discarded. If no specific probes could be identified for a gene, it was excluded from the experiment.

Probes for the 99 genes were designed using Resolve Biosciences' proprietary design algorithm and gene annotations from the Chinese Spring RefSeqv1.1 database. In brief, probe design was conducted at the gene level. For each targeted gene, all full-length protein-coding transcript sequences from the ENSEMBL database were utilized as design targets. To identify potential off-target sites, searches were confined to the coding regions. Each target sequence underwent a single scan for all k-mers, favoring regions with rare k-mers as seeds for full probe design. A probe candidate was generated by extending a seed sequence until reaching a certain target stability. Sequences known to cause problems experimentally were eliminated. Following these initial screens, each retained probe candidate was aligned with the background transcriptome, and probes with stable off-target hits were discarded. Specific probes were then scored based on the number of on-target matches (isoforms), giving preference to principal isoforms over others. A bonus was added if the binding site was within the protein-coding region. From the pool of accepted probes, the final set was selected from the highest-scoring probes.

4.1.3. Imaging

Samples were imaged by Resolve Biosciences using a Zeiss Celldiscoverer 7 (CD7) with a 50× Plan Apochromat water immersion objective having an NA of 1.2 and a 0.5× magnification changer, resulting in a final magnification of 25×. Standard CD7 LED excitation light sources, filters, and dichroic mirrors were used in conjunction with customized emission filters optimized for detecting specific signals. The excitation time per image was fixed at 1,000 ms for each channel, 20 ms for DAPI, and 1 ms for Calcofluor White. A z-stack was captured at each region with a z-slice spacing according to the Nyquist-Shannon sampling theorem. A custom CD7 CMOS camera (Zeiss Axiocam Mono 712, 3.45 μm pixel size) was employed. For each region, a z-stack was acquired for each of two fluorescent colors in each imaging round. A total of 8 imaging rounds were performed for each position, resulting in 16 z-stacks per region. The entirely automated imaging process per round, which included water immersion generation and precise relocation of regions for imaging in all three dimensions, was implemented through a custom Python script utilizing the scripting API of the Zeiss ZEN software (Open application development). Imaging for the cell wall-specific stain, Calcofluor White, was conducted after the primary 8-round imaging.

4.1.4. Spot segmentation and preprocessing

The algorithms for spot segmentation were written in Java and are based on the ImageJ library

functionalities. Only the iterative closest point algorithm is written in C++ based on the

libpointmatcher library (https://github.com/ethz-asl/libpointmatcher).

In the initial image processing steps, all images underwent background fluorescence correction.

The allowed number of maxima, representing the brightest points in the images, was determined

based on the slice's area and an empirically optimized factor. Each image slice was processed

independently. Maxima without neighboring counterparts in adjacent slices (referred to as z-

groups) were discarded. An iterative filtering process adjusted thresholds for factors like absolute

brightness, local background, and background of the periphery, all based on 3D image volume.

Only maxima remaining in z-groups of at least 2 were retained. These maxima, with the highest

absolute brightness within their z-groups, were documented as 3D-point cloud features.

For signal segmentation and decoding, the raw data images were aligned using transformation

matrices derived from the feature point clouds. An iterative closest point cloud algorithm

minimized error between point clouds. Point clouds from each round were aligned to the round

one reference, and the corresponding point clouds were stored for downstream use. Aligned

images were used to create pixel profiles with 16 values, representing two color channels across

eight imaging rounds. Profiles were filtered for variance from zero, normalized by total profile

brightness.

Matched pixel profiles were assigned IDs, and neighboring pixels with the same ID formed

groups. These pixel groups underwent further filtering, considering group size, adjacent pixel

count, and dimensions with a size of two pixels. Local 3D-maxima in groups were identified as

potential transcript locations. Maxima were filtered based on expectations from raw data images.

The remaining maxima were evaluated by fitting to corresponding codes, and they were

documented as transcripts. Specificity, assessed by the ratio of experiment-matching signals to

non-experiment-matching signals, was used to estimate false positives. This comprehensive

process enabled precise image analysis and transcript identification. Final image analysis was

performed in ImageJ using the Polylux tool plugin from Resolve BioSciences to examine

specific Molecular Cartography^{TM} signals. Images, programs, and gene expression results are

deposited in Zenodo [99].

4.1.5. Cell segmentation, clustering, and cell trajectories

We identified cell boundaries from the DAPI-stained images using the cell segmentation program QuPath [20]. To account for different cell sizes and shapes, we defined different regions in each image, applying region-specific parameter sets. Across all regions we used the parameters pixel size = 0, background radius = 8 μm, intensity threshold = 5, and opening by reconstruction with the "include cell nucleus" parameter disabled. For regular, well-defined cells, "split by shape" was activated, and the cell expansion was set to 7 μm, while for irregular, elongated cells, "split by shape" was turned off, and the cell expansion was set to 5 μm. The cell walls, visualized with Calcofluor White in the same sections, served as controls. The six images were merged into a single composite image and converted into a format suitable for clustering using ImageJ.

Hybridization signals per cell and clustering were calculated using the Resolve Biosciences "Recognize" software with the following parameters: Min cell = 5, Min Features = 1, Min transcripts = 1, Dimensions = 20, Resolution = 0.75, and Selected genes = 99. "Recognize" utilizes Seurat v4 [100] as the backend, with the following methods: NormalizeData (normalization.method= "LogNormalize", scale.factor= 1), FindVariableFeatures, ScaleData, RunPCA, FindNeighbors (dims= 1:20), FindClusters (resolution= 0.75), RunUMAP (dims= 1:20, n.components= 3, and FindAllMarkers. The R script to replicate the analysis using Seurat v5 can be found in Zenodo [99]. The average log2 fold changes of marker genes were generated locally using Seurat v5 [69], which implements a pseudocount at the group level instead of the cell level as in Seurat v4.

Cells from the three developmental stages and two replications were clustered together. Clusters are represented with the same color across stages and replicates. To facilitate visualization of the individual cell clusters, we also present them separately in Fig. S5. The Seurat object was deposited in Zenodo [99].

To find subclusters within scRNA-seq vasculature, we used the Seurat function "*FindSubCluster*" with parameter '*resolution* = 0.01*, graph.name* = "RNA_snn", *algorithm* = 2. We then used the Seurat function "*FindMarkers*" to find differentially expressed genes between the subclusters including xylem and phloem markers.

We reconstructed cellular trajectories using the computer program Monocle 3 [101]. The 21 cell clusters from the spatial analysis and the 23 cell clusters from the scRNA-seq were leveraged to

constrain two separate trajectory inferences. The initial cluster object was transferred into a Monocle SingleCellExperiment object using the as.cell_data_set function. Dimensionality reduction was carried out using the UMAP embeddings previously computed with RunUMAP(dims=1:30) into the Monocle object. Then function "cluster_cells" was applied using the default random seed and UMAP as the reduction method to preserve cluster consistency, followed by "learn_graph" to construct the principal graph, and "order_cells" to infer pseudotime trajectories, specifying meristematic cluster c8 as the root cluster in the spatial trajectory and sc14 in the scRNA-seq trajectory.

4.2. RNA-seq of different parts of the developing wheat spike

Kronos spikes at the glume primordia stage (W3.0) were sectioned into three parts (Fig. S24) and placed into 500 µl of DNA/RNA Shield solution in 1.5-ml Eppendorf tubes. Approximately 20 to 25 apices were used per replicate, and three replicates were collected for each section. Samples in DNA/RNA Shield were stored at 4℃ prior to RNA extraction.

We followed the published protocol by Marina Millán Blánquez for RNA extraction (dx.doi.org/10.17504/protocols.io.36wgq7kj3vk5/v1) with minor modifications. Briefly, 400 µl of DNA/RNA Shield solution was removed, and the tissues were homogenized with plastic pestles in the remaining 100 µl of solution. RNAs were then extracted using TRIzol reagent, cleaned, and concentrated with the RNA Clean & Concentrator-5 kit from Zymo Research. RNA samples were shipped to Novogene for quality control, library preparation, and sequencing on the NovaSeq X Plus (PE150) platform.

Data have been deposited in bioproject number PRJNA1217025. The three replicates from each region were clustered together in a principal component analysis, indicating high reproducibility among biological replicates (Fig. S24). Of the 49,786 detected genes, 13,024 exhibited significant differences among regions in Tukey tests (Data S11). For each region we calculated the average TMM per region and performed Tukey tests. The genes showing significant differences and the corresponding Tukey tests are listed in Data S11. Scripts for RNA-seq mapping, principal component plot, and Tukey tests were deposited in Zenodo [99].

4.3. Single-cell RNA sequencing

4.3.1. Protoplast preparation and 10x Genomics library construction and sequencing

We used a cell wall digestion method optimized to preserve meristem cells in maize

inflorescence [66]. Briefly, the inflorescence spike was finely dissected (Fig. 1A) under a

microscope and digested in a MOPS buffer-based enzyme mix (10 mM MOPS, pH 7.5; 10 mM

L-Arginine HCl, pH 7.5; 0.65 M mannitol; 1.5% Cellulase RS; 1.5% Cellulase R10; 1.0%

Macerozyme; 1.0% hemicellulose; 0.1% BSA; 5 mM β-mercaptoethanol; 1 mM $CaCl_2$) for 2

hours. Before adding $CaCl_2$, β-mercaptoethanol, and BSA, the solution was heated to 55°C for

10 minutes.

The digested cells and enzyme mix were passed through a 30 µm pluriStrainer (pluriSelect,

Catalog No. 43-50030-50) for filtration. The filtered protoplasts were pelleted by centrifugation

at 350g for 2 minutes at 4°C and washed with wash buffer (10 mM MOPS, pH 7.5; 10 mM L-

Arginine HCl, pH 7.5; 0.65 M mannitol). The protoplasts were pelleted again by centrifugation

at 350g for 2 minutes at 4°C. The clean protoplast pellets were gently resuspended in wash

buffer, and viability was checked using trypan blue. High-viability protoplasts ($\geq$ 70%) were

immediately loaded into a 10x Genomics Chromium System using Next GEM Single Cell 3' Kit

v3.1 chemistry. The scRNA-seq libraries were sequenced on the Element Bio AVITI platform

using the 10x Genomics recommended sequencing format, with 122 cycles used for Read 2

(Data S7).

4.3.2. Single-cell RNA-seq analysis and clustering

Single-cell RNA-seq reads were mapped to the Kronos genome assembly v1.1. (annotation v1.0)

[67] (https://doi.org/10.5281/zenodo.11106422) using STARsolo [102, 103]. Since scRNA-seq

reads were generated from the 3' end of the transcripts and not all Kronos genes have good

annotations of 3' UTR regions, we extended the 3' end of all gene models by 500 bp.

We used the Seurat V5 pipeline [69] to cluster the cells. Briefly, genes expressed in at least three

cells were retained. All cells from both stages were filtered to include those with 1,500 to 25,000

expressed genes and 8,000 to 300,000 unique molecular identifiers (UMIs).

Reads from individual samples were log-normalized, and 2,000 highly variable genes were

selected for downstream analysis. The integrated matrix was scaled using the 'ScaleData'

function. Principal component analysis (PCA) was performed with the 'RunPCA' function.

Further integrative analysis of all W2.5 and W3.5 stage samples was conducted using the

'IntegrateLayers' function with the anchor-based RPCA integration method. The top 35 principal

components (PCs) were used. A resolution of 1 was applied with the 'FindClusters' function to

identify clusters. Cell cycle clusters were identified (Fig. S23), and the remaining clusters were

re-normalized, scaled, integrated using the RPCA method, and re-clustered. The top 35 PCs were

used to generate the final clusters, at a resolution of 1 (Fig. 5B-D). Scripts used for mapping

reads and clustering analysis and the final Seurat object files were deposited in Zenodo [99].

4.3.3. Correlations among cell clusters

Normalized expression data for the 99 genes in the smFISH study were used to calculate Pearson

correlation coefficients between pairs of spatial clusters (Data S3). Normalized expression data

for the 200 genes used in the spatial transcriptomics study in hexaploid wheat were kindly

provided by Dr. Cristobal Uauy (John Innes Centre, UK) [21]. The hexaploid wheat data was

provided separately for the four different stages included in their study, but we only used the

three early stages (late double ridge, lemma primordia and terminal spikelet) that overlapped

with the stages of spike development used in our scRNAseq study.

The average normalized gene expression across all cells for each single-cell cluster were

obtained with the "AverageExpression" function in Seurat v5 in R after joining the six

replications (three for each stage) of the scRNA-seq data. For this analysis, we used the final

integrated data after removing the initial cell-cycle clusters and filtered the average normalized

data per cluster using two criteria. First, we selected genes with coefficients of variation higher

than 0.5 to eliminate genes with very similar expression levels across all clusters. We also

eliminated genes with a total sum across clusters <0.1 or >20 to minimize the effect of genes

with very high or very low expression levels (and multiple 0s). Using these criteria, we selected

14,903 informative genes that were used to calculate Pearson correlation coefficients between all

possible pairs of single-cell clusters using their average normalized expression across all

common genes (Data S8).

To calculate the correlations between the single-cell clusters and the spatial transcriptomics

clusters from hexaploid wheat, we first identified the overlapping genes for each comparison.

The coding regions of wheat homeologs are on average 97.5% identical, so most of the probes

used in the wheat spatial transcriptomics experiment are expected to hybridize with the different

homeologs. To make the single-cell data comparable, we summed the normalized expression of

the three homeologs in hexaploid wheat, but separate correlation matrices were generated for the

late double ridge (Data S13), lemma primordia (Data S14) and the terminal spikelet stage (Data

S15).

Clustering of the correlation matrices was done by the function "heatmap" from R package [104]

with default parameters (clustering_distance_rows = "euclidean", clustering_distance_cols =

"euclidean", clustering_method = "complete"). Scripts used to extract cluster averages and to

perform the correlation analyses are deposited in Zenodo [99].

4.3.4. **Imputation of scRNA-seq gene expression into spatial transcriptomics cells**

To integrate the smFISH and scRNA-seq data without losing spatial information, we imputed the

expression levels of all genes from the five nearest scRNA-seq cells into each cell from the

spatial transcriptomic experiment using a recently published approach [70]. Briefly, we used

SEURAT v5 [100] to calculate the cosine similarity (1 - cosine distance, varying from 0 for most

dissimilar to 1 for the most similar) among the smFISH and scRNA-seq cells based on the

expression of the 91 genes present in both studies (normalized counts per million). We identified

the five nearest scRNA-seq cell neighbors to each smFISH-labelled cell and calculated a

weighted average expression for each gene, using the cosine similarity as weight. The imputed

gene expression was then incorporated into the smFISH-labelled cells.

We integrated the smFISH and scRNA-seq for each stage (W2.5 and W3.5) separately and then

combined the imputed data. We imputed 70,878 genes for W2.5 and 72,475 for W3.5, resulting

in 74,464 for the combined analysis. We also developed a website for visualization of the

combined data (https://dubcovskylab.ucdavis.edu/imputed-genes). The scripts for data

imputation, the final imputed data, and the source data of the website are included in the Zenodo

depository (folder imputed_data) [99].

Using a similar procedure as in the imputation of scRNA-seq gene expression into spatial

transcriptomics cells, after the integration, we found the five nearest smFISH cells for each

scRNA-seq cell in W2.5 and W3.5 separately with one additional filtering: the nearest cell

should have cosine similarity > 0.3 to exclude very different cells and at the same time keep

enough number of cells for following analysis. We then assigned the scRNA-seq cell the most

frequent smFISH cluster number within the 5 neighbors to the scRNA-seq cell. If there were no

common smFISH cluster numbers (all 5 cluster numbers are unique), then this scRNA-seq cell

had no smFISH cluster projection. After the projection, we calculated the frequency matrix

between scRNA-seq original clusters and the projected smFISH clusters. The count matrix was

normalized by the total number of cells used for each smFISH cluster (the frequency counts /

total number of cells used in that smFISH cluster) to normalize for differences of cluster sizes.

4.3.5. Annotation of scRNA-seq cell clusters

To annotate the scRNA-seq clusters, we used markers known to be expressed in specific cell

types and/or previously associated with known functions in other species (Data S1) and validated

their spatial expression in Kronos developing spikes using the imputation method described

above. We supplemented this information with the correlations calculated between the single-cell

and spatial expression clusters identified in the published spatial transcriptomics study in

hexaploid wheat [21] (Data S13-15), and by the imputed expression of multiple differentially

expressed genes in each scRNA-seq cluster using the methods described in the previous section.

Additional information to annotate the scRNA-seq clusters was extracted from the genes

preferentially expressed in each cluster (Data S9). These genes were selected based on their low

abundance in other clusters (pct.2<0.10) relative to their abundance in the studied cluster

(pct.1>0.10), adjusted *P* values <0.001, and a greater than 2-fold difference in expression levels

(log2Fold>1, Data S9). Clusters sc1 and sc22 had few enriched genes, so we relaxed the filtering

parameters (average log2FC>1, pct.1 no filtering, pct.2<0.2, adjusted *P* value <0.01) (Data S9).

We subsequently determined the pathways enriched in these preferentially expressed genes using

KEGG analyses (Kyoto Encyclopedia of Genes and Genomes, Data S10) through the DAVID

webtool (https://davidbioinformatics.nih.gov/tools.jsp) [105]. We converted the Kronos v1.0

gene IDs into Chinese Spring annotation v1.1 gene IDs, uploaded the CS gene IDs as

"ENSEMBL_GENE_ID" (step 2: Select Identifier) and as "Gene List" (step 3: List Type), used

the "*Triticum aestivum*" as background species, and started the analysis using "Functional

Annotation Tool". In addition, we examined the average log2Fold expression across clusters for

a subset of 41 photosynthetic genes to identify clusters associated with photosynthetic functions,

and for 45 ribosomal protein-encoding genes to identify cells with high protein production,

typically associated with elevated cell division rates. These results were visualized using heat

maps.

To incorporate additional spatial information into our cluster annotation, we used the RNA-seq

data generated from the base, middle and top third of the spike described in section 4.2.

(henceforth described as spike base, middle and tip). To determine if a cell cluster was enriched

in any of these three spike regions, we identified genes with significant differences in Tukey tests

among the preferentially expressed genes for each cluster and quantified the number of genes

expressed at higher levels in each region. To determine if the distribution across the three regions

was significantly different from a random uniform distribution, we performed chi-square tests for

each cluster (Data S16).

4.3.6. **Detection of clusters affected by the protoplasting process**

The protoplasting process may trigger a stress response in some cells, potentially influencing the

clustering process. We employed two independent approaches to assess the potential effects of

protoplasting on gene expression in the different clusters. First, we used a list of 713 maize genes

previously identified as potentially affected by the protoplasting process [84], identified the

closest Kronos homologs, and determined the percentage of preferentially expressed genes in

each cluster overlapping with this list (totaling 215 unique genes). The second approach, which

identified 235 unique genes, involved the identification of preferentially expressed genes in each

single-cell cluster that were not present in two RNA-seq data from Kronos developing spikes

(this study and [68]). A GO-term analysis of the unique genes identified in both analyses

revealed enrichment in stress-related annotations, particularly among the genes not detected in

the two RNA-seq datasets (Data S19). These results suggest that many of these genes may have

been induced by the protoplasting process and are typically not expressed under normal

conditions in the developing wheat spike. The percentages of genes identified by these two

independent methods across clusters were highly correlated (R= 0.813, P< 0.0001) suggesting a

similar underlying phenomenon. Both methods identified clusters sc3 and sc22 as having the

highest risk of being affected by the protoplasting process (Data S16).

4.3.7. **scRNA-seq co-expression analysis**

Using the cell x gene normalized count matrix, we calculated Pearson correlation coefficients (R)

between genes of interest with known spatial distribution (*AGL6, FZP, TCP24, ULT-A1*) and all

other detected genes in the scRNA-seq across all cells (26,009 cells. We focused on co-expressed

genes with correlation coefficients higher than the average plus four standard deviations. The

script used for co-expression analysis was deposited in Zenodo [99].

4.3.8. **Network analysis**

To better understand the interactions between *TCP24* and *FZP*, we performed a gene regulatory

network analysis using the program GENIE3 [96]. The analysis included the 988 genes co-

expressed with *FZP* (Data S20) and/or *TCP24* (Data S21) in the scRNA-seq dataset. We

extracted the top-ranked 500 links to generate and visualize the network using Cytoscape 3.10.3

(https://cytoscape.org/). In Data S22, we summarized the first neighbors of *TCP-A24*, *TCP-B24*,

FZP-A, and *FZP-B* and identified shared and unique transcription factors among them.

4.3.9. Generation of loss-of-function mutants of FZP and SPL14 using CRISPR-Cas9

To validate the specific functions of *FZP* and *SPL14,* we used CRISPR-Cas9 to edit both

homeologs of each target gene in the tetraploid wheat variety Kronos. We designed two guide

RNAs for *FZP* (FZP-gRNA-1 and FZP-gRNA-2, Table S1) and one for *SPL14* (SPL14-gRNA,

Table S1). Guide RNAs for each gene were cloned into a binary vector (Addgene Plasmid

#160393) containing the *Cas9* gene and a *GRF4-GIF1* chimera, which enhances wheat

regeneration efficiency during *Agrobacterium*-mediated transformation [106]. Transgenic plants

were generated at the UC Davis Plant Transformation Facility using methods described

previously [106]. Independent transgenic plants were screened for mutations using next-

generation sequencing (NGS) as previously described [107] using primers described in Table S1.

We identified eight independent T_0 lines with mutations in both *FZP* homeologs and further

confirmed the edits by Sanger sequencing using primers listed in Table S1. For *SPL14*, we

obtained 10 individual T_0 transgenic plants. This line was backcrossed with wildtype Kronos to

segregate out the *Cas9* transgene, and we identified one line in the T_4 generation with successful

edits in both the A and B genomes and without the transgene. We developed CAPS markers to

track the *spl14* mutations across generations using primers described in Table S1.

1353

1354 **ACKNOWLEDGEMENTS**

1355 The authors thank Chaozhong Zhang and Qiujie Liu (UC Davis) for their help with the selection

1356 of the genes included in the smFISH experiment, Samik Bhattacharya from Resolve Biosciences

for his continuous help with the spatial transcriptomics project, Cristobal Uauy from the John Innes Centre UK for his advice with the analysis of the spatial transcriptomics data and early access to his spatial transcriptomics data, Rüdiger Simon and Björn Usadel for their help with the expression imputation method, and Ksenia Krasileva from UC Berkeley for early access to the Kronos genome. JD acknowledges funding from the Howard Hughes Medical institute and the United States Department of Agriculture (USDA) – National Institute of Food and Agriculture (NIFA) grants 2022-68013-36439 (WheatCAP) and 2022-67013-36209. X.X acknowledges USDA-NIFA HATCH project (CA-D-PLB-2850-H) and UC Davis new faculty start-up funding.

AUTHORS CONTRIBUTIONS

XX, HL, and JD conceived the project. Experimental work was performed by HL, XX, FP, KL, CT, and YL. JZ, GB, HL, CL, XX and JD performed data analyses. JD and XX obtained funding and directed the project. HL, XX and JD generated the first draft, and all authors contributed to the revision of the manuscript. XX, CL and JD provided supervision.

CONFLICT OF INTERESTS

The authors declare no conflict of interests.

DATA AVAILABILITY

Data and scripts used in the different analyses are available in Supplemental Data and in Zenodo https://doi.org/10.5281/zenodo.14872904 [99]. The ImageJ version with the Polylux tool plugin, the high-resolution images for the six sections used in the paper (Calcofluor White Figs. S3, S4 and DAPI), the hybridization results for the 99 genes for each image, and instructions for download and installation are also available at our web site (https://dubcovskylab.ucdavis.edu/content/spatial-transcriptomic) and in Zenodo [99]. We also developed a web tool to visualize individual cell clusters and heatmaps of the 99 genes in the six sections https://dubcovskylab.ucdavis.edu/JD99-wheat-spike-smFISH

and a similar tool to visualize the imputed genes https://dubcovskylab.ucdavis.edu/imputed-genes. These tools can be also downloaded from the same Zenodo link and run on individual computers. The raw RNA-seq data of the three regions of the Kronos spike at W3.0 were

deposited in NCBI's Sequence Read Archive (SRA) under BioProject number PRJNA1217025. The processed data, scripts used for mapping and feature counts, and a Readme file with instructions are deposited in Zenodo [99].

The scRNA-seq data were deposited in NCBI's Sequence Read Archive (SRA) under BioProject accession number PRJNA1199502. The final single-cell clusters can be accessed via the Shiny App (https://dubcovskylab.ucdavis.edu/content/kronos-integrated-clusters). Additionally, a 3D-UMAP (https://dubcovskylab.ucdavis.edu/content/kronos-scrna-3d-clusters) was generated to visualize cluster distributions. The Kronos genome assembly v1.1 (annotation v1.0) is available in Zenodo https://doi.org/10.5281/zenodo.11106422. Processed single-cell data, Seurat objects and R scripts are also available in Zenodo https://doi.org/10.5281/zenodo.14872904.

Data and scripts for cluster correlations, the Seurat object produced by Resolve BioSciences "Recognize" online tool, and the R script to replicate the online analysis locally can be also found in Zenodo [99].

REFERENCES

1. FAOSTAT: **Food and Agriculture Organization (FAO) of the United Nations. http://www.fao.org/faostat/en/#data.** 2023.

2. Kellogg EA: **Genetic control of branching patterns in grass inflorescences.** *Plant Cell* 2022, **34:**2518-2533.

3. Li C, Lin H, Chen A, Lau M, Jernstedt J, Dubcovsky J: **Wheat *VRN1, FUL2* and *FUL3* play critical and redundant roles in spikelet development and spike determinacy.** *Development* 2019, **146:**dev175398.

4. Chen H, Zhang X, Xu SH, Song CX, Mao HL: ***TaSPL17*s act redundantly with *TaSPL14*s to control spike development and their elite haplotypes may improve wheat grain yield.** *Frontiers in Plant Science* 2023, **14:**1229827.

5. Chuck G, Whipple C, Jackson D, Hake S: **The maize SBP-box transcription factor encoded by *tasselsheath4* regulates bract development and the establishment of meristem boundaries.** *Development* 2010, **137:**1243-1250.

6. Chuck GS, Brown PJ, Meeley R, Hake S: **Maize SBP-box transcription factors *unbranched2* and *unbranched3* affect yield traits by regulating the rate of lateral primordia initiation.** *Proc Natl Acad Sci U S A* 2014, **111:**18775-18780.

7. Li K, Debernardi JM, Li C, Lin H, Zhang C, Jernstedt J, von Korff M, Zhong J, Dubcovsky J: **Interactions between SQUAMOSA and SVP MADS-box proteins regulate meristem transitions during wheat spike development.** *The Plant Cell* 2021, **33:**3621-3644.

8. Adamski NM, Simmonds J, Brinton JF, Backhaus AE, Chen Y, Smedley M, Hayta S, Florio T, Crane P, Scott P, et al: **Ectopic expression of *Triticum polonicum VRT-A2* underlies elongated glumes and grains in hexaploid wheat in a dosage-dependent manner.** *The Plant Cell* 2021, **33:**2296-2319.

9. Debernardi JM, Lin H, Chuck G, Faris JD, Dubcovsky J: **microRNA172 plays a crucial role in wheat spike morphogenesis and grain threshability.** *Development* 2017, **144:**1966-1975.

10. Greenwood JR, Finnegan EJ, Watanabe N, Trevaskis B, Swain SM: **New alleles of the wheat domestication gene *Q* reveal multiple roles in growth and reproductive development.** *Development* 2017, **144:**1959-1965.

11. Debernardi JM, Greenwood JR, Jean Finnegan E, Jernstedt J, Dubcovsky J: ***APETALA 2*-like genes *AP2L2* and *Q* specify lemma identity and axillary floral meristem development in wheat.** *Plant Journal* 2020, **101:**171-187.

12. Reinheimer R, Kellogg EA: **Evolution of *AGL6-like* MADS box genes in grasses (Poaceae): ovule expression is ancient and palea expression is new.** *Plant Cell* 2009, **21:**2591-2605.

13. Paraiso F, Lin H, Li C, Woods DP, Lan T, Tumelty C, Debernardi JM, Joe A, Dubcovsky J: **LEAFY and WAPO1 jointly regulate spikelet number per spike and floret development in wheat.** *Development* 2024, **151:**dev202803.

14. Kuzay S, Lin H, Li C, Chen S, Woods DP, Zhang J, Lan T, von Korff M, Dubcovsky J: ***WAPO-A1* is the causal gene of the 7AL QTL for spikelet number per spike in wheat.** *PLoS Genetics* 2022, **18:**e1009747.

15. Kuzay S, Xu Y, Zhang J, Katz A, Pearce S, Su Z, Fraser M, Anderson JA, Brown-Guedira G, DeWitt N, et al: **Identification of a candidate gene for a QTL for spikelet number per spike on wheat chromosome arm 7AL by high-resolution genetic mapping.** *Theor Appl Genet* 2019, **132:**2689–2705.

16. Muqaddasi QH, Brassac J, Koppolu R, Plieske J, Ganal MW, Roder MS: ***TaAPO-A1*, an ortholog of rice *ABERRANT PANICLE ORGANIZATION 1*, is associated with total spikelet number per spike in elite European hexaploid winter wheat (*Triticum aestivum* L.) varieties.** *Scientific Reports* 2019, **9:**13853.

17. Voss-Fels KP, Keeble-Gagnere G, Hickey LT, Tibbits J, Nagornyy S, Hayden MJ, Pasam RK, Kant S, Friedt W, Snowdon R, et al: **High-resolution mapping of rachis nodes per rachis, a critical determinant of grain yield components in wheat.** *Theoretical and Applied Genetics* 2019, **132:**2707-2719.

18. Zhang J, Burguener GF, Paraiso F, Dubcovsky J: **Natural alleles of LEAFY and WAPO1 interact to regulate spikelet number per spike in wheat.** *Theor Appl Genet* 2024, **137:**257.

19. Waddington SR, Cartwright PM, Wall PC: **A quantitative scale of spike initial and pistil development in barley and wheat.** *Annals of Botany* 1983, **51:**119-130.

20. Bankhead P, Loughrey MB, Fernandez JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid S, Gray RT, Murray LJ, Coleman HG, et al: **QuPath: Open source software for digital pathology image analysis.** *Sci Rep* 2017, **7:**16878.

21. Long KA, Lister A, Jones MRW, Adamski NM, Ellis RE, Chedid C, Carpenter SJ, Liu X, Backhaus AE, Goldson A, et al: **Spatial transcriptomics reveals expression gradients in developing wheat inflorescences at cellular resolution.** *bioRxiv* 2024, **https://doi.org/10.1101/2024.12.19.629411**.

22. Perico C, Zaidem M, Sedelnikova O, Bhattacharya S, Korfhage C, Langdale JA: **Multiplexed in situ hybridization reveals distinct lineage identities for major and minor vein initiation during maize leaf development.** *Proceedings of the National Academy of Sciences of the United States of America* 2024, **121:**e2402514121.

23. Bonke M, Thitamadee S, Mähönen AP, Hauser MT, Helariutta Y: **APL regulates vascular tissue identity in *Arabidopsis*.** *Nature* 2003, **426:**181-186.

24. Zhang J, Li C, Zhang W, Zhang X, Mo Y, Tranquilli GE, Vanzetti LS, Dubcovsky J: **Wheat plant height locus *RHT25* encodes a PLATZ transcription factor that interacts with DELLA (RHT1).** *Proceedings of the National Academy of Sciences of the United States of America* 2023, **120:**e2300203120.

25. McKim SM: **How plants grow up.** *J Integr Plant Biol* 2019, **61:**257-277.

26.   Wang J, Wang R, Wang Y, Zhang L, Zhang L, Xu Y, Yao S: ***Short and Solid***
***Culm/RFL/APO2* for culm development in rice.** *Plant J* 2017, **91:**85-96.

27.   Ndreca B, Huttly A, Bibi S, Bayon C, Lund G, Ham J, Alarcon-Reverte R, Addy J,
Tarkowska D, Pearce S, et al: **Stacked mutations in wheat homologues of rice *SEMI-***
***DWARF1* confer a novel semi-dwarf phenotype.** *BMC Plant Biol* 2024, **24:**384.

28.   Pearce S, Vanzetti LS, Dubcovsky J: **Exogenous gibberellins induce wheat spike**
development under short days only in the presence of *VERNALIZATION1*. *Plant*
Physiology 2013, **163:**1433-1445.

29.   Sakamoto T, Morinaka Y, Ishiyama K, Kobayashi M, Itoh H, Kayano T, Iwahori S,
Matsuoka M, Tanaka H: **Genetic manipulation of gibberellin metabolism in**
transgenic rice. *Nat Biotechnol* 2003, **21:**909-913.

30.   Wang L, Ming L, Liao K, Xia C, Sun S, Chang Y, Wang H, Fu D, Xu C, Wang Z, et al:
Bract suppression regulated by the *miR156/529-SPLs-NL1-PLA1* module is required
for the transition from vegetative to reproductive branching in rice. *Mol Plant* 2021,
14:1168-1184.

31.   Dong Z, Hu G, Chen Q, Shemyakina EA, Chau G, Whipple CJ, Fletcher JC, Chuck G: **A**
regulatory network controlling developmental boundaries and meristem fates
contributed to maize domestication. *Nat Genet* 2024, **56:**2528-2537.

32.   Lu Z, Yu H, Xiong G, Wang J, Jiao Y, Liu G, Jing Y, Meng X, Hu X, Qian Q, et al:
Genome-wide binding analysis of the transcription activator IDEAL PLANT
ARCHITECTURE1 reveals a complex network regulating rice plant architecture.
Plant Cell 2013, **25:**3743-3759.

33.   Doebley J, Stec A, Hubbard L: **The evolution of apical dominance in maize.** *Nature*
1997, **386:**485-488.

34.   Takeda T, Suwa Y, Suzuki M, Kitano H, Ueguchi-Tanaka M, Ashikari M, Matsuoka M,
Ueguchi C: **The *OsTB1* gene negatively regulates lateral branching in rice.** *Plant J*
2003, **33:**513-520.

35.   Ramsay L, Comadran J, Druka A, Marshall DF, Thomas WT, Macaulay M, MacKenzie
K, Simpson C, Fuller J, Bonar N, et al: ***INTERMEDIUM-C*, a modifier of lateral**
spikelet fertility in barley, is an ortholog of the maize domestication gene
***TEOSINTE BRANCHED 1*.** *Nat Genet* 2011, **43:**169-172.

36. Dixon LE, Greenwood JR, Bencivenga S, Zhang P, Cockram J, Mellers G, Ramm K,
Cavanagh C, Swain SM, Boden SA: *TEOSINTE BRANCHED1* **regulates inflorescence**
architecture and development in bread wheat (*Triticum aestivum***).** *Plant Cell* 2018,
30:563-581.

37. Volkman MM, Martin JM, Hogg AC, Wright L, Hale C, Carr PM, Giroux MJ: **Durum**
wheat *Teosinte Branched1* **null mutations increase tillering.** *Crop Science* 2022,
62:1522-1530.

38. Huijser P, Klein J, Lonnig WE, Meijer H, Saedler H, Sommer H: **Bracteomania, an**
inflorescence anomaly, is caused by the loss of function of the MADS-box gene
squamosa **in** *Antirrhinum majus***.** *Embo Journal* 1992, **11:**1239-1249.

39. Klein J, Saedler H, Huijser P: **A new family of DNA binding proteins includes**
putative transcriptional regulators of the *Antirrhinum majus* **floral meristem**
identity gene *SQUAMOSA***.** *Molecular and General Genetics* 1996, **250:**7-16.

40. Reiser L, Sánchez-Baracaldo P, Hake S: **Knots in the family tree: evolutionary**
relationships and functions of homeobox genes. *Plant Molecular Biology* 2000,
42:151-166.

41. Leiboff S, Li XR, Hu HC, Todt N, Yang JL, Li X, Yu XQ, Muehlbauer GJ, Timmermans
MCP, Yu JM, et al: **Genetic control of morphometric diversity in the maize shoot**
apical meristem. *Nature Communications* 2015, **6:**8974.

42. O'Connor DL, Runions A, Sluis A, Bragg J, Vogel JP, Prusinkiewicz P, Hake S: **A**
division in PIN-mediated auxin patterning during organ initiation in grasses. *Plos*
Computational Biology 2014, **10:**e1003447.

43. Zong J, Wang L, Zhu L, Bian L, Zhang B, Chen X, Huang G, Zhang X, Fan J, Cao L, et
al: **A rice single cell transcriptomic atlas defines the developmental trajectories of**
rice floret and inflorescence meristems. *New Phytol* 2022, **234:**494-512.

44. Patterson EL, Richardson A, Bartlett M: **Pushing the boundaries of organ identity:**
Homology of the grass lemma. *American Journal of Botany* 2023, **110:**e16161.

45. Moraes TD, van Es SW, Hernández-Pinzón I, Kirschner GK, van der Wal F, da Silveira
SR, Busscher-Lange J, Angenent GC, Moscou M, Immink RGH, van Esse GW: **The**
TCP transcription factor HvTB2 heterodimerizes with VRS5 and controls spike
architecture in barley. *Plant Reproduction* 2022, **35:**205-220.

46. Bai F, Reinheimer R, Durantini D, Kellogg EA, Schmidt RJ: **TCP transcription factor,**
BRANCH ANGLE DEFECTIVE 1 (BAD1), is required for normal tassel branch
angle formation in maize. *Proceedings of the National Academy of Sciences of the*
United States of America 2012, **109:**12225-12230.

47. Poursarebani N, Trautewig C, Melzer M, Nussbaumer T, Lundqvist U, Rutten T,
Schmutzer T, Brandt R, Himmelbach A, Altschmied L, et al: **COMPOSITUM 1**
contributes to the architectural simplification of barley inflorescence via meristem
identity signals. *Nat Commun* 2020, **11:**5138.

48. Yuan Z, Gao S, Xue DW, Luo D, Li LT, Ding SY, Yao X, Wilson ZA, Qian Q, Zhang
DB: *RETARDED PALEA1* **controls palea development and floral zygomorphy in**
rice. *Plant Physiol* 2009, **149:**235-244.

49. Komatsu M, Chujo A, Nagato Y, Shimamoto K, Kyozuka J: *FRIZZY PANICLE* **is**
required to prevent the formation of axillary meristems and to establish floral
meristem identity in rice spikelets. *Development* 2003, **130:**3841-3850.

50. Chuck G, Muszynski M, Kellogg E, Hake S, Schmidt RJ: **The control of spikelet**
meristem identity by the *branched silkless1* **gene in maize.** *Science* 2002, **298:**1238-
1241.

51. Dobrovolskaya O, Pont C, Sibout R, Martinek P, Badaeva E, Murat F, Chosson A,
Watanabe N, Prat E, Gautier N, et al: *FRIZZY PANICLE* **drives supernumerary**
spikelets in bread wheat. *Plant Physiology* 2015, **167:**189-199.

52. Poursarebani N, Seidensticker T, Koppolu R, Trautewig C, Gawronski P, Bini F, Govind
G, Rutten T, Sakuma S, Tagiri A, et al: **The genetic basis of composite spike form in**
barley and 'Miracle-Wheat'. *Genetics* 2015, **201:**155-165.

53. Bommert P, Whipple C: **Grass inflorescence architecture and meristem determinacy.**
Seminars in Cell & Developmental Biology 2018, **79:**37-47.

54. Yang Y, Fu D, Zhu C, He Y, Zhang H, Liu T, Li X, Wu C: **The RING-finger ubiquitin**
ligase HAF1 mediates Heading date 1 degradation during photoperiodic flowering
in rice. *Plant Cell* 2015, **27:**2455-2468.

55. Zhu C, Peng Q, Fu D, Zhuang D, Yu Y, Duan M, Xie W, Cai Y, Ouyang Y, Lian X, Wu
C: **The E3 ubiquitin ligase HAF1 modulates circadian accumulation of EARLY**

FLOWERING3 to control heading date in rice under long-day conditions. *Plant Cell* 2018, **30:**2352-2367.

56. McSteen P, Malcomber S, Skirpan A, Lunde C, Wu XT, Kellogg E, Hake S: *barren inflorescence2* **encodes a co-ortholog of the** *PINOID* **serine/threonine kinase and is required for organogenesis during inflorescence and vegetative development in maize.** *Plant Physiology* 2007, **144:**1000-1011.

57. Carles CC, Choffnes-Inada D, Reville K, Lertpiriyapong K, Fletcher JC: *ULTRAPETALA1* **encodes a SAND domain putative transcriptional regulator that controls shoot and floral meristem activity in Arabidopsis.** *Development* 2005, **132:**897-911.

58. Monfared MM, Carles CC, Rossignol P, Pires HR, Fletcher JC: **The** *ULT1* **and** *ULT2* **trxG genes play overlapping roles in** *Arabidopsis* **development and gene regulation.** *Molecular Plant* 2013, **6:**1564-1579.

59. Wang WF, Li G, Zhao J, Chu HW, Lin WH, Zhang DB, Wang ZY, Liang WQ: **DWARF TILLER1, a WUSCHEL-related homeobox transcription factor, is required for tiller growth in rice.** *Plos Genetics* 2014, **10:**e1004154.

60. Bomblies K, Wang RL, Ambrose BA, Schmidt RJ, Meeley RB, Doebley J: **Duplicate** *FLORICAULA/LEAFY* **homologs** *zfl1* **and** *zfl2* **control inflorescence architecture and flower patterning in maize.** *Development* 2003, **130:**2385-2395.

61. Ikeda-Kawakatsu K, Maekawa M, Izawa T, Itoh JI, Nagato Y: *ABERRANT PANICLE ORGANIZATION 2/RFL***, the rice ortholog of Arabidopsis** *LEAFY***, suppresses the transition from inflorescence meristem to floral meristem through interaction with** *APO1***.** *Plant Journal* 2012, **69:**168-180.

62. Selva C, Shirley NJ, Houston K, Whitford R, Baumann U, Li G, Tucker MR: *HvLEAFY* **controls the early stages of floral organ specification and inhibits the formation of multiple ovaries in barley.** *Plant Journal* 2021, **108:**509-527.

63. Glenn P, Woods DP, Zhang J, Gabay G, Odle N, Dubcovsky J: **Wheat bZIPC1 interacts with FT2 and contributes to the regulation of spikelet number per spike.** *Theoretical and Applied Genetics* 2023, **136:**237.

64. Zhu W, Yang L, Wu D, Meng Q, Deng X, Huang G, Zhang J, Chen X, Ferrandiz C, Liang W, et al: **Rice** *SEPALLATA* **genes** *OsMADS5* **and** *OsMADS34* **cooperate to**

limit inflorescence branching by repressing the *TERMINAL FLOWER1-like* gene *RCN4*. *New Phytol* 2022, **233:**1682-1700.

65. Zhang J, Xiong H, Burguener GF, Vasquez-Gross H, Liu Q, Debernardi JM, Akhunova A, Garland-Campbell K, Kianian SF, Brown-Guedira G, et al: **Sequencing 4.3 million mutations in wheat promoters to understand and modify gene expression.** *Proceedings of the National Academy of Sciences of the United States of America* 2023, **120:**e2306494120.

66. Xu X, Passalacqua M, Rice B, Demesa-Arevalo E, Takebayashi MKY, Harris B, Sakakibara H, Gallavotti A, Gillis J, Jackson D: **Large-scale single-cell profiling of stem cells uncovers redundant regulators of shoot development and yield trait variation.** *bioRxiv* 2024, **doi.org/10.1101/2024.03.04.583414**.

67. Seong K, Kumar R, Lunde C, Krasileva K: **Chromosome-level genome assembly of *Triticum turgidum* var 'Kronos' [Data set].** Zenodo. https://doi.org/10.5281/zenodo.11106422 2023.

68. VanGessel C, Hamilton J, Tabbita F, Dubcovsky J, Pearce S: **Transcriptional signatures of wheat inflorescence development.** *Sci Rep* 2022, **12:**17224.

69. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G, Madad S, Fernandez-Granda C, Satija R: **Dictionary learning for integrative, multimodal and scalable single-cell analysis.** *Nat Biotechnol* 2024, **42:**293-304.

70. Demesa-Arevalo E, Dörpholz H, Vardanega I, Maika JE, Pineda-Valentino I, Eggels S, Lautwein T, Köhrer K, Schnurbusch T, von Korff M, et al: **Imputation integrates single-cell and spatial gene expression data to resolve transcriptional networks in barley shoot meristem development.** *bioRxiv* 2025**:**2025.2005.2009.653223.

71. Cho H, Dang TV, Hwang I: **Emergence of plant vascular system: roles of hormonal and non-hormonal regulatory networks.** *Curr Opin Plant Biol* 2017, **35:**91-97.

72. Miyashima S, Sebastian J, Lee JY, Helariutta Y: **Stem cell function during plant vascular development.** *EMBO J* 2013, **32:**178-193.

73. Skipper M: **Genes from the *APETALA3* and *PISTILLATA* lineages are expressed in developing vascular bundles of the tuberous rhizome, flowering stem and flower primordia of *Eranthis hyemalis*.** *Ann Bot* 2002, **89:**83-88.

1631   74.   Houston K, Druka A, Bonar N, Macaulay M, Lundqvist U, Franckowiak J, Morgante M,
1632   Stein N, Waugh R: **Analysis of the barley bract suppression gene Trd1.** *Theor Appl*
1633   *Genet* 2012, **125:**33-45.
1634   75.   Wang L, Yin H, Qian Q, Yang J, Huang C, Hu X, Luo D: ***NECK LEAF 1*, a GATA type**
1635   **transcription factor, modulates organogenesis by regulating the expression of**
1636   **multiple regulatory genes during reproductive development in rice.** *Cell Res* 2009,
1637   **19:**598-611.
1638   76.   Whipple CJ, Hall DH, DeBlasio S, Taguchi-Shiobara F, Schmidt RJ, Jackson DP: **A**
1639   **conserved mechanism of bract suppression in the grass family.** *Plant Cell* 2010,
1640   **22:**565-578.
1641   77.   Wang YG, Du F, Wang J, Wang K, Tian CH, Qi XQ, Lu F, Liu XG, Ye XG, Jiao YL:
1642   **Improving bread wheat yield through modulating an unselected AP2/ERF gene.**
1643   *Nature Plants* 2022, **8:**930-939.
1644   78.   Oikawa T, Kyozuka J: **Two-step regulation of LAX PANICLE1 protein accumulation**
1645   **in axillary meristem formation in rice.** *Plant Cell* 2009, **21:**1095-1108.
1646   79.   Galli M, Liu Q, Moss BL, Malcomber S, Li W, Gaines C, Federici S, Roshkovan J,
1647   Meeley R, Nemhauser JL, Gallavotti A: **Auxin signaling modules regulate maize**
1648   **inflorescence architecture.** *Proc Natl Acad Sci U S A* 2015, **112:**13372-13377.
1649   80.   Jin Y, Luo Q, Tong H, Wang A, Cheng Z, Tang J, Li D, Zhao X, Li X, Wan J, et al: **An**
1650   **AT-hook gene is required for palea formation and floral organ number control in**
1651   **rice.** *Dev Biol* 2011, **359:**277-288.
1652   81.   Gallavotti A, Zhao Q, Kyozuka J, Meeley RB, Ritter MK, Doebley JF, Pe ME, Schmidt
1653   RJ: **The role of *barren stalk1* in the architecture of maize.** *Nature* 2004, **432:**630-635.
1654   82.   Panikashvili D, Shi JX, Bocobza S, Franke RB, Schreiber L, Aharoni A: **The**
1655   ***Arabidopsis* DSO/ABCG11 transporter affects cutin metabolism in reproductive**
1656   **organs and suberin in roots.** *Mol Plant* 2010, **3:**563-575.
1657   83.   Douglas SJ, Li B, Kliebenstein DJ, Nambara E, Riggs CD: **A novel *Filamentous Flower***
1658   **mutant suppresses *brevipedicellus* developmental defects and modulates**
1659   **glucosinolate and auxin levels.** *PLoS One* 2017, **12:**e0177045.
1660   84.   Xu XS, Crow M, Rice BR, Li F, Harris B, Liu L, Demesa-Arevalo E, Lu ZF, Wang LY,
1661   Fox N, et al: **Single-cell RNA sequencing of developing maize ears facilitates**

functional analysis and trait candidate gene discovery. *Developmental Cell* 2021, **56:**557-568.

85. Khanday I, Yadav SR, Vijayraghavan U: **Rice *LHS1/OsMADS1* controls floret meristem specification by coordinated regulation of transcription factors and hormone signaling pathways.** *Plant Physiol* 2013, **161:**1970-1983.

86. Borghi L, Bureau M, Simon R: **Arabidopsis *JAGGED LATERAL ORGANS* is expressed in boundaries and coordinates *KNOX* and *PIN* activity.** *Plant Cell* 2007, **19:**1795-1808.

87. Ohmori Y, Tanaka W, Kojima M, Sakakibara H, Hirano HY: **WUSCHEL-RELATED HOMEOBOX4 is involved in meristem maintenance and is negatively regulated by the CLE gene FCP1 in rice.** *Plant Cell* 2013, **25:**229-241.

88. Wang J, Bao JL, Zhou BB, Li M, Li XZ, Jin J: **The *osa-miR164* target *OsCUC1* functions redundantly with *OsCUC3* in controlling rice meristem/organ boundary specification.** *New Phytologist* 2021, **229:**1566-1581.

89. Hibara K, Karim MR, Takada S, Taoka KI, Furutani M, Aida M, Tasaka M: **Arabidopsis *CUP-SHAPED COTYLEDON3* regulates postembryonic shoot meristem and organ boundary formation.** *Plant Cell* 2006, **18:**2946-2957.

90. Gallavotti A, Malcomber S, Gaines C, Stanfield S, Whipple C, Kellogg E, Schmidt RJ: **BARREN STALK FASTIGIATE1 is an AT-hook protein required for the formation of maize ears.** *Plant Cell* 2011, **23:**1756-1771.

91. Li Z, Liu D, Xia Y, Li Z, Jing D, Du J, Niu N, Ma S, Wang J, Song Y, et al: **Identification of the WUSCHEL-related homeobox (*WOX*) gene family, and interaction and functional analysis of TaWOX9 and TaWUS in wheat.** *Int J Mol Sci* 2020, **21:**1581.

92. Fang F, Ye S, Tang J, Bennett MJ, Liang W: **DWT1/DWL2 act together with OsPIP5K1 to regulate plant uniform growth in rice.** *New Phytol* 2020, **225:**1234-1246.

93. Hendelman A, Zebell S, Rodriguez-Leal D, Dukler N, Robitaille G, Wu XL, Kostyun J, Tal L, Wang PP, Bartlett ME, et al: **Conserved pleiotropy of an ancient plant homeobox gene uncovered by *cis*-regulatory dissection.** *Cell* 2021, **184:**1724-1739.

94.   Bortiri E, Chuck G, Vollbrecht E, Rocheford T, Martienssen R, Hake S: *ramosa2*

encodes a LATERAL ORGAN BOUNDARY domain protein that determines the

fate of stem cells in branch meristems of maize. *Plant Cell* 2006, **18:**574-585.

95.   Whipple CJ: **Grass inflorescence architecture and evolution: the origin of novel**

signaling centers. *New Phytologist* 2017, **216:**367-372.

96.   Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P: **Inferring regulatory networks**

from expression data using tree-based methods. *PLoS One* 2010, **5:**e12776.

97.   Tan F, Zhou C, Zhou Q, Zhou S, Yang W, Zhao Y, Li G, Zhou DX: **Analysis of**

chromatin regulators reveals specific features of rice DNA methylation pathways.

Plant Physiol 2016, **171:**2041-2054.

98.   Baylis T, Cierlik I, Sundberg E, Mattsson J: ***SHORT INTERNODES/STYLISH* genes,**

regulators of auxin biosynthesis, are involved in leaf vein development in

Arabidopsis thaliana. *New Phytol* 2013, **197:**737-750.

99.   Xu X, Lin H, Zhang J, Burguener G, Paraiso F, Li K, Tumelty C, Li C, Liu Y,

Dubcovsky J: **Spatial and single-cell expression analyses reveal complex expression**

domains in early wheat spike development [Data set] In *Zenodo*

https://doi.org/10.5281/zenodo.14872904; 2025.

100.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ, Wilk

AJ, Darby C, Zager M, et al: **Integrated analysis of multimodal single-cell data.** *Cell*

2021, **184:**3573-3587 e3529.

101.   Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S,

Christiansen L, Steemers FJ, et al: **The single-cell transcriptional landscape of**

mammalian organogenesis. *Nature* 2019, **566:**496-502.

102.   Kaminow B, Yunusov D, Dobin A: **STARsolo: accurate, fast and versatile**

mapping/quantification of single-cell and single-nucleus RNA-seq data. *bioRxiv*

2021**:**2021.2005.2005.442755.

103.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

Gingeras TR: **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2013,

29:15-21.

104.   Kolde R: **pheatmap: Pretty Heatmaps. https://CRAN.R-**

project.org/package=pheatmap. 2019.

105.   Huang DW, Sherman BT, Lempicki RA: **Systematic and integrative analysis of large**

gene lists using DAVID bioinformatics resources. *Nature Protocols* 2009, **4:**44-57.

106.   Debernardi JM, Tricoli DM, Ercoli MF, Hayta S, Ronald P, Palatnik JF, Dubcovsky J: **A**

GRF-GIF chimeric protein improves the regeneration efficiency of transgenic

plants. *Nature Biotechnology* 2020, **38:**1274-1279.

107.   Zhang J: *Check CRISPR editing events in transgenic wheat with next-generation*

sequencing. New York, NY: Springer US; 2022.

1730