AMPK- and cGAS-Mediated Dual Sensing of Glucose and dsDNA Synergizes to Enhance Antitumor Immunity

Ruyuan Zhou[1,2,3,#], Jingyan Pei[1,3,#], Shasha Chen[4], Qin Shen[1,3], Qirou Wu[3,5], Shengduo Liu[2,3], Xin-Hua Feng[3,6], Jian Zou[7], Tingbo Liang[1,6], Y. Jessie Zhang[8], Sheng-Cai Lin[9], Penghong Song[1,*], Chen-Song Zhang[8,*], Qian Zhang[3,8,*], and Pinglong Xu[1,2,3,6,*]

[1] Department of Hepatobiliary and Pancreatic Surgery and Zhejiang Provincial Key Laboratory of Pancreatic Disease, The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou, 310058, China;

[2] Institute of Intellectual Medicine, ZJU-Hangzhou Global Scientific and Technological Innovation Center, Hangzhou, 310058, China;

[3] The MOE Key Laboratory of Biosystems Homeostasis & Protection and Zhejiang Provincial Key Laboratory of Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang University, Hangzhou, Zhejiang 310058, China;

[4] Zhejiang Provincial Key Laboratory for Water Environment and Marine Biological Resources Protection, College of Life and Environmental Science, Wenzhou University, Wenzhou 325035, China;

[5] Key Laboratory of Novel Targets and Drug Study for Neural Repair of Zhejiang Province, School of Medicine, Hangzhou City University, Hangzhou 310015, China

[6] Cancer Center, Zhejiang University, Hangzhou, Zhejiang 310058, China;

[7] Eye Center of the Second Affiliated Hospital School of Medicine, Institute of Genetics, School of Medicine, Zhejiang University, Hangzhou 310058, China;

[8] Department of Molecular Biosciences, University of Texas at Austin, Austin, TX 78712, USA;

[9] State Key Laboratory for Cellular Stress Biology, School of Life Sciences, Xiamen University, Xiamen, 361102, China;

[#] These authors contributed equally;

[*] Correspondence should be addressed to P.S., C.-S. Z., Q.Z., or P.X.

Address for correspondence:

Penghong Song

Department of Hepatobiliary and Pancreatic Surgery, The First Affiliated Hospital, School of

Medicine, Zhejiang University

79 Qingchun Road, Hangzhou 310003, China

E-mail: songpenghong@zju.edu.cn

Chen-Song Zhang

State Key Laboratory for Cellular Stress Biology, School of Life Sciences, Xiamen University,

Xiamen, Fujian, China

E-mail: cszhang@xmu.edu.cn

Qian Zhang

Department of Molecular Biosciences, University of Texas at Austin

100 E. 24th Street, Norman Hackerman Building (NHB) 4.130, Austin, TX 78712, USA

Tel: 1-512-471-1029

E-mail: qian.zhang1@austin.utexas.edu

Pinglong Xu

Life Sciences Institute, Zhejiang University

866 Yuhangtang Road, Nano Building, Room A555, Hangzhou, 310058, China

Tel: 86-571-88206078

E-mail: xupl@zju.edu.cn

Running Title: AMPK-TBK1 Integrates Glucose Availability to Antitumor Immunity

ABSTRACT

Intratumoral glucose availability is linked considerably to tumor growth, immune evasion, and metastasis. However, the precise molecular mechanisms by which glucose levels govern cellular and immune responses against tumors remain incompletely understood. Here, we explored the context of tumors of intersections between AMP-activated protein kinase (AMPK) and cGAS-STING signaling, a key pathway initiating antitumor immunity. We unexpectedly found that AMPK activation robustly enhances cGAS-STING signaling within cancer and immune cells, which potently impacts melanoma and colorectal cell senescence, organoid apoptosis, and the infiltration of CD4$^+$ and CD8$^+$ T lymphocytes. Using classic or newly developed AMPK agonist Aldometanib, TBK1$^{S511E/S511E}$ knock-in (KI) mice, glucose metabolic modulator, and AMPKα1/α2 dKO melanoma, we demonstrated that dual sensing of intratumoral glucose and dsDNA integrated by the AMPK-TBK1 cascade is essential for initiating antitumor immunity and suppressing cancer progressions. Notably, novel AMPK agonists or intervening in glucose glycolysis by Lonidamine synergized effectively with STING agonists, substantially inhibiting melanoma growth. Therefore, these findings unravel a concise mechanism integrating glucose metabolism and innate DNA sensing into cancer cell fate and propose an effective therapeutic strategy to enhance antitumor immunity.

Key Words: AMPK; TBK1; cGAS-STING; antitumor immunity; glucose metabolism; synergistic therapy; glucose sensing; nucleic acid sensing; AMPK agonists; senescence; apoptosis; dual sensing; T lymphocytes; melanoma; colorectal cancer; immunotherapy; immune evasion; glycolysis; tumor microenvironment; cancer immunology

HIGHLIGHTS

1. Dual and synergistic sensing of glucose and dsDNA facilitates antitumor immunity in melanoma and colorectal cancer;

2. AMPK substitutes intratumoral cGAS-STING-IRF3 signaling through TBK1 phosphorylation at S511;

3. Glucose deficiency induces cancer cell senescence, apoptosis, and T lymphocyte infiltration via the AMPK-TBK1 cascade;

4. Newly developed AMPK agonists synergize with STING agonists to retrieve striking tumor suppression.

INTRODUCTION

Nucleic acid sensing surveils tissue abnormalities and pathogen invasion and establishes an immune state to restrict microbial infection, modulate adaptive immunity, and guide tissue repair and regeneration [7-11]. Pattern recognition receptors (PRRs) such as cGAS [12,13] monitor microbes and tissue damage by perceiving cytosolic dsDNA derived from pathogens, the mitochondria, or the nucleus [7,8,10,14]. Upon recognition of free dsDNA, cGAS synthesizes 2'3'-cyclic GMP-AMP (cGAMP)[12,15-17], activates STING (also known as MITA or ERIS) [18-20], which triggers the non-canonical STING-PERK pathway in the endoplasmic reticulum (ER) that regulates mRNA translation [21], induces non-canonical autophagy in the ERGIC [22], and facilitates the activation of TBK1 and IRF3 at the Golgi apparatus [23,24]. Phosphorylated IRF3 translocates into nuclear and transcribes type I interferons (IFN-Is) and numerous IFN-stimulated genes (ISGs) [25,26], in coordination with simultaneously activated NF-κB [7,8,10]. Therefore, nucleic acid sensing induces mRNA translation arrest [21], autophagy [22,27,28], phase separation [29], mitochondrial dynamics [30], cell differentiation [31], and triggers cell senescence [32-35] and cell death [36-38].

Effects of cGAS-STING signaling in tumor biology are complex and multifaceted [39]. cGAS-STING signaling is active in cancerous, immune, and stromal cells, shifting their cellular function, metabolism, and fate. The activation of cGAS-STING signaling releases various IFN-I, cytokines, cGAMP, and DNA, which attracts and matures dendritic cells, macrophages, natural killer cells (NKs), and effector T lymphocytes to mount an antitumor immunity [40,41]. STING-induced senescence and immune microenvironment modulation also participate in antitumor immunity regulation, positively or negatively [42,43,44,45,46,47], and STING-dependent residency of tumor-associated macrophages (TAMs) attenuate its antitumor role [48]. Therefore, the effects of cGAS-STING in cancer biology could be context- and stage-dependent and should be carefully evaluated during cancer treatments.

Cells in multicellular organisms depend on balanced nutrient availability for survival, growth, and defense. Despite this, the direct molecular connection between nutrient sensing and tissue damage perception remains less understood. AMP-activated protein kinase (AMPK) plays a central role in detecting glucose deficiency and controlling metabolic pathways [49-51]. AMPK is activated at the lysosomal surface in response to low glucose levels through aldolase

and the v-ATPase-Ragulator complex[52-54], in addition to being induced by AMP/ATP ratios and

kinase such as LKB1 or CaMKII [50,55,56]. AMPK masterly regulates an array of cellular processes

such as glucose and lipid metabolism, protein synthesis, autophagy, mitochondrial biogenesis,

and overall metabolism [50,57], and its dysfunction is broadly accompanied by obesity, type 2

diabetes, and cancers. We recently reported that viral infection causes a rapid and dramatic

decrease in blood glucose levels and substantially activates AMPK in tissues, which directly

phosphorylates TBK1 to trigger IRF3 recruitment and assemble MAVS or STING signalosomes,

thus effectively mounting antiviral immunity[2]. In adipocytes, TBK1 also represses energy

expenditure by inhibiting AMPKα [58]. These observations indicate the presence of the

intersection between metabolic master kinase AMPK and immune key kinase TBK1.

Nevertheless, the roles of metabolism and immunity crosstalk through AMPK-TBK1 signaling

in the context of malignant diseases remain poorly understood.

However, hyperglycemia has been considerably linked to diabetes-related vulnerability to

malignancy [59], and it impacts cancer immunology in distinct ways. For instance, elevated

glucose levels through the hexosamine biosynthesis pathway (HBP) enhance the O-

GlcNAcylation level in TAMs, promoting M2-like polarization and cancer progression [60]. In diet-

induced obesity, more immunosuppressive myeloid-derived suppressor cells (MDSCs) were

observed with higher tumor progression of pancreatic cancer [61,62]. Meanwhile, tumor cells

typically exhibit increased glucose uptake and enhanced glycolysis, which supports key

processes for tumorigenesis, such as the biosynthesis of nucleotides, amino acids, and lipids.

Focusing on the AMPK-TBK1 axis, here we systematically investigated the link between

glucose availability and antitumor immunity. Using the well-documented and newly

developed AMPK agonists and TBK1 S511E knock-in (KI) mice, we found that activated AMPK

directly phosphorylated intratumoral TBK1 at S511, promoting cGAS-STING-initiated

senescence and antitumor immunity, thereby attenuates the progression of melanoma and

colorectal cancer. In summary, we proposed an intratumoral dual and synergistic sensing of

dsDNA and glucose, which forms a "metabolism and immune sensing" bridge through the

concise AMPK-TBK1 signaling and determines tumor progression.

RESULTS

Glucose deficiency promotes intratumoral cGAS-STING signaling

To explore the function of glucose metabolism on antitumor immunity in TME, we investigated the state of glucose deficiency and its impacts on cGAS-STING signaling, an innate immune pathway crucial in antitumor immunity. Intratumoral glucose deficiency was evident, as reported [63-65], and as indicated by AMPK-induced phosphorylation of metabolic regulator ACC1 in melanoma (Fig. 1A-1B). pACC1 signal was particularly robust at tumor edges, where proliferating tumor cells and immune cells need high glucose demand. cGAMP treatment in primary peritoneal macrophages (PMs) triggered robust activation of TBK1 and IRF3, the central kinase and key signaling mediator of cGAS-STING signaling. Simulating an intratumoral glucose-depleted microenvironment using 2-deoxy-D-glucose (2-DG) resulted in an enhanced cGAS-STING-IRF3 pathway when compared to energy-sufficient conditions, accompanied by robust activation of AMPK (Fig. 1C). We observed a similar enhancement of cGAS-STING signaling in colorectal adenocarcinoma cells (DLD1) and mouse embryonic fibroblasts (MEFs) under energy-deficient conditions (Figs. 1D-1E). AICAR, a well-established AMPK agonist [66], expectedly facilitated cGAS-STING signaling in fibroblasts and cancer cells (Figs. 1D-1E). Likewise, glucose depletion amplified mitochondrial DNA (mtDNA)-induced activation of cGAS-STING-IRF3 signaling in colorectal carcinoma cells (Fig. 1F), achieved by combinational treatment of ABT-737, qVD-OPH, and S63845 (AQS) [67-69]. AICAR also potentiated the sensing of cytosolically exposed dsDNA in DLD-1 cells, simulated by poly(dA:dT) transfection (TpdAdT) (Fig. 1G). These observations across immune cells, fibroblasts, and cancer cells suggest that glucose deficiency, frequently seen in tumors, significantly augments cGAS-STING signaling and is likely through AMPK activation.

AMPK facilitates cGAS-STING signaling in cancer cells by TBK1 S511 phosphorylation

To validate that glucose deficiency regulates nucleic acid sensing through AMPK activation, we generated AMPK α1/α2 dKO cells in B16-F10 melanoma cells using a CRISPR-mediated approach (Fig. 2A). Genetic deletion of AMPK catalytic subunits α1/α2 attenuated cGAS-STING-IRF3 signaling in B16 melanoma, as indicated by reduced levels of phospho-STING (S366) and phospho-IRF3 (S396) (Fig. 2A). Likewise, deletion of AMPK α1 and α2 subunits in

colorectal carcinoma HCT116 cells similarly decreased RLR-MAVS-IRF3 signaling (Fig. 2B), the dsRNA pathway with an emerged role in antitumor immunity [70,71]. The role of AMPK on TBK1 activation was also evident as AMPK catalytic subunits strongly potentiated the TBK1-IRF3 interaction (Fig. 2C) and TBK1-stimulated IRF3 transactivation (Fig. 2D). Our previous study and the mass spectrometry analyses of AMPK-coexpressed TBK1 have identified that the S511 residue on TBK1 is abundantly modified by AMPKα1 [2]. We found that AMPK inhibitor dorsomorphin (Compound C) abolished TBK1 S511 phosphorylation (Fig. 2E), and an antibody specifically recognizing phospho-TBK1 S511 demonstrated that AMPKα1 directly phosphorylated TBK1 at this residue (Fig. 2F). We thus evaluated the capability of wild-type, phospho-resistant (S511A), and phosphomimetic TBK1 (S511E) to activate IRF3 by an IRF3-responsive reporter, which indicated that AMPK-mediated TBK1 phosphorylation at S511 was critical for driving IRF3 transactivation (Fig. 2G). Expectedly, a compromised interaction between phospho-resistant TBK1 and IRF3 was seen (Fig. 2H). These findings suggest that AMPK enhances cGAS-STING signaling by phosphorylating TBK1 at S511 to prime TBK1 activation.

To validate this AMPK-TBK1 axis in cancer cells, we generated and examined knock-in (KI) clones of colorectal cells DLD1 with mutations that either prevent (S511A) or mimic (S511D) AMPK-mediated modification by CRISPR-based genome editing [2]. cGAS-STING signaling was attenuated when endogenous TBK1 was mutated into the S511A variant that resists AMPK modification (Fig. 2I). Conversely, STING signaling was robustly activated when endogenous TBK1 was mutated to the S511D variant that mimics AMPK-TBK1 phosphorylation, to a level comparable to those under AICAR treatment (Fig. 2J). Noticeably, the AICAR-enhanced activation of STING signaling was absent in these DLD1 KI cells (Fig. 2J). These observations suggest that AMPK-mediated TBK1 phosphorylation at residue S511 is critical for cancer cells to respond to glucose deficiency.

AMPK drives STING-induced cancer cell senescence and apoptosis

cGAS-STING responds to dsDNA leakage and drives cellular senescence, which could be important in tumor growth arrest [32-34]. We found that upon chemical induction of DNA damage, colorectal carcinoma DLD1 cells displayed strong phenotypes of cellular senescence,

evidenced by SA-β-Gal staining and senescence-associated secretory phenotypes (SASPs) (Fig. 3A-3B). Genetic ablation of AMPK α1/α2 subunits endowed DLD1 with apparent resistance to DNA damage-induced cellular senescence (Fig. 3A). By contrast, AICAR facilitated the progression of these colorectal cancer cells into senescence (Fig. 3B). In agreement with these observations, simulating the glucose-depleted microenvironment using 2-DG greatly enhanced DLD1 senescence (Fig. 3C). We further found that by utilizing a zebrafish model of damage-induced senescence, AMPK activation by AICAR promoted camptothecin (CPT)-induced senescence in zebrafish, indicated by increased numbers of SA-β-Gal-positive cells in the heads and tails of zebrafish (Fig. 3D). Inhibition of AMPK by dorsomorphin strongly attenuated CPT-induced senescence, as expected (Fig. 3D). These observations suggest that AMPK is critical to regulate damage-induced cellular senescence, mediated by cGAS-STING signaling.

The activation of cytosolic dsDNA sensing also mediates a context-dependent apoptotic program [72], an evolved strategy beneficial to tissue homeostasis. We found that activation of cGAS-STING by DNA damage inducer HU [73] triggered significant levels of colorectal cancer cell apoptosis, which was noticeably regulated by AICAR and dorsomorphin (Fig. 3E). Additionally, by utilizing *in vitro*-cultured intestinal organoids, we detected a strong phenotype of HU-induced apoptosis among intestinal cells, which was further exaggerated by AMPK activation (Fig. 3F). These observations suggest that glucose deficiency and resulting AMPK activation controls both senescence and apoptosis of cancer and functional cells through an AMPK-TBK1 axis.

AMPK-TBK1 signaling facilitates STING-mediated antitumor immunity in melanomas

Next, we utilized a naturally occurring and constitutively active STING SAVI mutant (R281Q) to generate B16-F10 melanoma with inducible and constitutively active STING signaling. Upon implanting subcutaneously into wild-type C57BL/6 mice, we found that melanomas harboring STING R281Q SAVI mutation exhibited significantly reduced growth (Figs. 4A-4C), attributed mainly from STING-induced expression of type I interferons (IFNs) [73]. Increased immune cells infiltrated these melanomas with constitutive STING signaling, particularly T cells (Figs. 4D-4E), indicating enhanced antitumor immunity. Administration of AMPK agonist AICAR

significantly intensified T lymphocyte infiltration in melanomas (Figs. 4D-4E) and suppressed tumor growth substantially (Figs. 4A-4C), suggesting that AMPK can amplify STING-mediated antitumor immunity in melanoma.

To determine whether the AMPK-TBK1 axis mediates this enhanced antitumor immunity effects, we generated knock-in (KI) mice harboring the TBK1 S511E mutation using a CRISPR-mediated strategy [2], which mimics AMPK-mediated TBK1 phosphorylation. The TBK1 S511E homozygous mice were viable, fertile, and appeared phenotypically normal. However, melanoma growth was markedly attenuated in these TBK1 S511E mice (Figs. 4F-4H), along with a significantly increased level of T lymphocyte infiltration (Figs. 4I-4J). More elevated antitumor immunity was not seen in TBK1 S511E knock-in mice in the presence of cGAMP (Figs. 4F-4H), further suggesting the importance of AMPK-TBK1 signaling in this process. These consistent observations propose the AMPK-TBK1 axis as a critical factor in facilitating STING-mediated antitumor immunity.

AMPK agonists synergize well with cGAMP to promote antitumor immunity in melanomas

To evaluate the potential for improving intratumoral antitumor immunity through this AMPK-TBK1 axis, we employed in the in vivo model of melanoma a well-characterized AMPK agonist AICAR [66] and a newly developed AMPK agonist Aldometanib [54], which acts through distinct mechanisms. As expected, administration of cGAMP in mice, a natural STING ligand, resulted in significant tumor growth suppression. AICAR (Figs. 5A-5C) and Aldometanib (Figs. 5D-F) markedly enhanced melanoma growth arrest, synergizing well with cGAMP. These findings suggest that maneuvering AMPK activation can effectively strengthen STING-induced antitumor immunity. Intriguingly, we observed that Aldometanib also enhanced antitumor effects in mice bearing AMPK dKO B16-F10 melanomas (Figs. 5G-5I), suggesting an additional cancer cell-nonautonomous effect of Aldometanib in melanoma.

Interrupting aerobic glycolysis potentiates antitumor immunity through the AMPK-TBK1 cascade

Next, we interrupted glucose metabolism by inhibiting aerobic glycolysis in cancer cells using Lonidamine, a classical hexokinase inhibitor [74], to evaluate the potential of glucose

metabolism intervention in antitumor immunity. Lonidamine interferes with energy

metabolism and induces AMPK activation [53,75,76]. Notably, we found that combinational use of

cGAMP and Lonidamine resulted in substantial tumor growth arrest (Figs. 6A-6C), along with

significantly increased infiltration of intratumoral T lymphocytes in melanomas (Figs. 6D-6E).

Furthermore, we found that Lonidamine promoted the robust activation of intratumoral

cGAMP-induced cGAS-STING signaling, evidenced clearly by increased levels of

phosphorylated STING (pSTING S365), STING clustering, and phosphorylated TBK1 (pTBK1

S172) in B16-F10 melanomas (Fig. 6F). These findings suggest that targeting glucose

metabolism can be a promising strategy in antitumor therapeutic, such as in melanomas.

DISCUSSION

Tumor cell metabolism depletes essential nutrients in the tumor microenvironment (TME), leading to significant glucose competition between cancer and immune cells. Meanwhile, the intervention of cGAS-STING signaling is a significant feature of cancers, mainly by genetic deletion of signaling components and suppression of signaling cascades, which attenuates its beneficial roles in antitumor immunity [29,73,77]. However, the exact mechanisms for cGAS-STING activation and its contribution to cellular behaviors in cancer and immune cells, such as cell death, senescence, IFN secretion, and metabolic changes, need further interrogation. This study elucidated how the glucose sensor AMPK senses intratumoral glucose abundance and promotes antitumor immunity through direct phosphorylation and activation of TBK1, an essential kinase during innate immune responses. Given the complex energy metabolism in TME, understanding the AMPK-TBK1 axis opens new avenues for targeting glucose metabolism in both cancer and immune cells. Through activating cGAS-STING signaling and enhancing T-cell-mediated antitumor immunity, these findings provide significant therapeutic targets for drug development in cancer immunotherapy.

Cancer cells primarily rely on aerobic glycolysis for energy metabolism, through which glucose is converted into lactic acid. This metabolic shift exacerbates glucose deficiency stress within the tumor, leading to metabolic competition between tumor cells and immune cells. Notably, these metabolic changes in immune cells contribute to establishing an immunosuppressive microenvironment for accelerating tumor growth and immune evasion. While previous studies have extensively characterized the Warburg effect and its impact on tumor proliferation and immune suppression, this study focuses on how metabolic sensing mechanisms combined with nucleic acid sensing influence the immune landscape of the TME. The cGAS-STING pathway facilitates immune cell recruitment and inhibits tumor progression through cellular senescence and apoptosis [42]. A recent study has highlighted that the AMPK-TBK1 axis plays a crucial role in innate antiviral immunity, extending AMPK's function beyond its classical roles in glucose and lipid metabolism, protein synthesis, autophagy, and mitochondrial function [2]. Changes in glucose metabolism in the TME are closely linked to immune responses. We found that the interaction between AMPK and cGAS-STING forms a "metabolic and immune sensing" bridge, which synergistically senses the metabolic state of

tumor cells and tumor microenvironment to decide the magnitude of immune responses during tumorigenesis. AMPK directly phosphorylates S511 residue on TBK1, enhancing STING-IRF3 signaling activation and promoting T-cell infiltration into the TME, significantly inhibiting melanoma growth. Therefore, these findings expand the understanding of metabolic integration in innate immunity and provide new insights into antitumor immunity.

New therapeutics for tumor treatment increasingly focus on regulating cellular metabolism and immune responses. We propose that modulating glucose levels within tumors, the activation state of AMPK, and the modulation of AMPK-TBK1 signaling could offer novel aspects for antitumor therapeutics. As a critical energy sensor, AMPK is activated in response to glucose deprivation, thereby adjusting the cellular metabolic state. Our study revealed that AMPK activation leads to direct phosphorylation of TBK1 at S511, enhancing cGAS-STING-IRF3 signaling and promoting antitumor immune responses and STING-induced cellular senescence and apoptosis, thereby potently inhibiting tumor growth. Besides, AMPK plays an important role in the adaptive stress response of tumor cells, regulating key metabolic pathways to inhibit tumor cell proliferation [51]. We supposed AMPK also influences the metabolism of immune cells in the TME, particularly macrophage polarization, which impacts tumor immune evasion. By modulating AMPK activity, immune cell surveillance of tumors can be enhanced through cGAS-STING signaling, effectively preventing tumor immune escape.

In conclusion, our study reveals the key role of the concise AMPK-TBK1 axis in regulating intratumoral glucose metabolism and promoting antitumor immune responses, which considerably connects glucose metabolism with innate immunity. We propose that targeting the AMPK-TBK1 axis, the intersection between intratumoral glucose metabolism and antitumor immunity, could reach a substantial synergy for cancer treatments.

ACKNOWLEDGMENTS

This research was sponsored by the NSFC Projects (32321002, 32430028, and 31830052 to

P.X., 82271768 to Q.Z., and 82001668, 32370759 to S.C.) and the National Key Research and

Development Program of China (2021YFA1301401 to P.X. and 2021YFD1801103 to Q.Z.).

Thanks also to technical assistance by the Life Sciences Institute core facilities, Zhejiang

University.

AUTHOR CONTRIBUTIONS

Q.Z. and R.Z. carried out most experiments. C.-S.Z., J.P., S.C., Q.S., Q.W., and S.L. contributed

to several experiments. X.-H.F., J.Z., T.L., Y.J.Z., S.-C.L., P.S., and C.-S.Z. helped with reagents,

data analyses, and discussions. P.X., Q.Z., and R.Z. conceived the study and experimental

design and wrote the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

MATERIALS AND METHODS

Mice

TBK1 S511E transgenic mice were on C57BL/6 background, and both male and female littermates were used in all the experiments. GemPharmatech generated TBK1 Knock-in transgenic C57BL/6 mice. To create point mutations (S511E) at mouse Tbk1 locus by CRISPR/Cas9-mediated genome engineering, Tbk1 S511E (TCT to GAA) specific sgRNA oligos were designed by the CRISPR website (http://crispr.mit.edu/) targeting sequence at S511 locus (5'- TTCTATTGTTCCCTGAGAAC -3') and (5'- ATTTAGCTTTCCAGTTCTCA -3'), with S511E oligo donor sequence (5'-CATTTGGATCTGATCCGTTGTTCTGACCTAACCTAACCCGTTGTATTTAGCTTTCCAGCGAACAGGGAAC AATAGAAAGCAGTCTTCAGGACATCAGCAGCAGGCTGTCTCCAGGGGGCT -3'). Cas9 mRNA and sgRNA generated by in vitro transcription and donor oligo were co-injected into fertilized eggs for knock-in mouse production. The target region of the mouse Tbk1 locus was sequenced to confirm targeting. All the animals were bred and maintained in a pathogen-free animal facility at the laboratory animal center of Zhejiang University. The care of experimental animals was approved by the Zhejiang University committee and followed Zhejiang University guidelines.

Zebrafish

Zebrafish AB wild-type embryos (male/female) were raised at 28.5°C in E3 egg water. The care of experimental animals was approved by the Zhejiang University committee and followed Zhejiang University guidelines.

Peritoneal Macrophages

Peritoneal macrophages were isolated from C57BL/6 mice at 6-8 weeks of age with the Brewer thioglycollate medium (Sigma-Aldrich)-induced approach. Three days after the intraperitoneal injection of 2.5 mL of 3% thioglycollate medium, peritoneal macrophages were isolated and cultured in the RPMI 1640 medium.

Cell Lines

HEK293, HCT116, DLD1, MEFs, and B16-F10 cells were from ATCC. No cell lines used in this study were found in the database of commonly misidentified cell lines maintained by ICLAC

and NCBI Biosample. Cell lines were frequently checked in morphology under microscopy and

tested for mycoplasma contamination but were not authenticated. All cell lines were cultured

in DMEM medium with 10% fetal bovine serum (FBS) at 37°C in 5% CO_2 (v/v), except for

peritoneal macrophages that were maintained in RPMI 1640 medium,

Intestinal Organoids

Isolated small intestines were opened longitudinally, washed three times with cold PBS,

removed the villi by scraping, and incubated twice in cold PBS with 2 mM EDTA for 15 min on

ice. After removing the EDTA medium, the fractions were incubated in 50 mL cold PBS with a

fierce shake and passed through a 70-µm cell strainer (BD Bioscience) to remove the residual

villous material. Isolated crypts were centrifuged at 300 g for 5 min to separate the crypts with

single cells. The centrifuged fractions were suspended and cultured in Matrigel with medium

(Advanced DMEM/F12 supplemented with penicillin/streptomycin, 10 mM HEPES, Glutamax,

13N2, 13B27 (all from Invitrogen), and 1mM N-acetylcysteine (Sigma) containing growth

factors 50 ng mL^{-1} EGF, 100 ng mL^{-1} noggin, 1mg mL^{-1} R-spondin). The organoid formation was

observed and analyzed every day after plating.

METHOD DETAILS

Expression Plasmids, Viruses, Reagents, and Antibodies

Expression plasmids encoding Flag-, Myc-, or HA-tagged wild-type or mutations of human

TBK1, IRF3, and the reporters of IFNβ_Luc and 5xISRE_Luc have been described previously

[78,79]. Flag- or HA-tagged human AMPKα1, AMPKα2, caAMPKα1 and AMPKα1 kinase domains

were constructed using the pRK5 mammalian expression vector. Site-directed mutagenesis

about TBK1 S511A/D/E and AMPKα1 T172D were generated by PCR-based cloning performed

by a kit from Stratagene. All coding sequences were verified by DNA sequencing, and the

detailed information for construction was provided in the attached Supplementary Table 2

and upon requirement.

The pharmacological reagent, Compound C (Selleck), AICAR (Selleck), 991 (Selleck), A-

769662 (Selleck), ABT-737 (Selleck), qVD-OPH (Selleck), S63845 (Selleck), MK2206 (Selleck),

Doxycycline (Sangon Biotech), 2-DG (Sangon Biotech), cGAMP (Invivogen), poly(I:C)

(Invivogen), poly (dA:dT) (Invivogen), puromycin (Yeasen), G418 (Yeasen), Hydroxyurea (HU)

(Sigma), and Lonidamine (Selleck) were purchased and used according to manual instructions.

Aldometanib was generated by Dr. Sheng-Cai Lin (Xiamen University, Xiamen).

The monoclonal anti-TBK1 (3504S, 1:5,000 dilution), anti-pTBK1(S172) (5483S, 1:3,000

dilution), anti-IRF3 (4302S, 1:2,000 dilution), anti-pIRF3 (S396) (4947S, 1:5,000 dilution), anti-

AMPKα (5831S, 1:1,000 dilution), anti-pAMPKα (T172) (50081S, 1:1,000 dilution), anti-pACC

(S79) (3661S, 1:1,000 dilution), anti-ACC (3676S, 1:1,000 dilution), anti-Myc (2276S, 1:3,000

dilution), and anti-HA (3724S, 1:5,000 dilution) antibodies were purchased from Cell Signaling

Technology. The anti-pIRF3 (S386) (ab76493, 1:3,000 dilution), anti-STING (ab181125, 1:100

dilution) were purchased from Abcam, and the anti-α-tubulin (T6199, 1:10,000 dilution), anti-

β-actin (A5441, 1:10,000 dilution), anti-GAPDH (EPR16891, 1:3000), anti-HA (H9658, 1:200

dilution), and anti-Flag (M2) (F3165, 1:5,000 dilution) were purchased from Sigma. The anti-

rabbit IgG and anti-mouse IgG antibodies were purchased from Santa Cruz. The anti-pTBK1

(S511) was generated in collaboration with Abcam, targeting the phospho-TBK1 S511.

Plasmids Transfection and Virus Infection of Cultured Cells

Lipofectamine 3000 (Invitrogen), Polyethyleneimine (PEI, Polysciences), or Lipofectamine

RNAiMAX (Invitrogen) transfection reagents were used for the transfection of plasmids,

poly(dA:dT), and poly(I:C). Digitonin (Sigma) was used to induce cGAMP.

CRISPR/Cas9-mediated Generation of AMPKα1/α2$^{-/-}$ and TBK1 Knock-in Cells

Guide RNA sequences targeting AMPKα1 and α2 mRNA sequence (hAMPKα1, 5'-

AAAGTTTGAGTGCTCAGAAG -3', 5'- GTGATGGAATATGTCTCAGG -3'; hAMPKα2, 5'-

TCAATTAACAGGCCATAAAG -3', 5'- GTTATTTAAGAAGATCCGAG -3'), TBK1 S511A/D knock-in

mRNA sequence (hTBK1, 5'- TATTTAGCTTTCCAGTTCTC -3', 5'- TTCTATTGTTCCCTGAGAAC -3')

were used to clone the genes into the vector pX330, which were transfected into DLD1 cells

by LipofectAmine 3000, together with the pCas9-2A-GFP plasmid, or with pCas9-2A-GFP and

pMD18 in the case of TBK1 S511A/D knock-in generation. 36 hours after transfection, cells

with green fluorescence were sorted with a Flow Cytometer (BD FACS Aria II) and propagated.

Immunoblotting with anti-AMPK(or the sequencing of the genomic PCR products) identified

clones. All gRNAs used in the experiments were also attached in Supplementary Table 1.

Luciferase Reporter Assay

HEK293 cells were transfected with indicated reporters (100 ng) bearing an open read frame

(ORF) coding Firefly luciferase, along with the pRL-Luc with Renilla luciferase coding as the

internal control for transfection and other expression vectors specified in the results section.

In brief, after 24 hours post-transfection, cells were lysed by passive lysis buffer (Promega),

and luciferase assays were performed using a dual luciferase assay kit (Promega), quantified

with POLARstar Omega (BMG Labtech) and normalized to the internal Renilla luciferase

control.

Quantitative RT-PCR Assay

The DLD1 cells stimulated with Hydroxyurea (HU), Camptothecin (CPT), or Etoposide (ETO)

were lysed, and total RNA was extracted using an RNAeasy extraction kit (Axygen). cDNA was

generated by a one-step iScript cDNA synthesis kit (Vazyme), and quantitative real-time PCR

was performed using the EvaGreen Qpcr MasterMix (Abm) and CFX96 real-time PCR system

(Bio-Rad). Relative quantification was expressed as 2-ΔCt, where Ct is the difference between

the primary Ct value of triplicates of the sample and an endogenous L19 mRNA control. The

primer sequences used are listed in the following:

hIL-1α, 5'- AGTGCTGCTGAAGGAGATGCCTGA -3', 5'- CCCCTGCCAAGCACACCCAGTA -3';

hIL-1β, 5'- ATGATGGCTTATTACAGTGGCAA -3', 5'- GTCGGAGATTCGTAGCTGGA -3';

RPL19, 5'- ATGTATCACAGCCTGTACCTG -3', 5'- TTCTTGGTCTCTTCCTCCTTG -3';

All primers used in the qRT-PCR assay were also attached in Supplementary Table 1.

Coimmunoprecipitations and Immunoblottings

HEK293 transfected with specified plasmids encoding Myc-, Flag-, or HA-tagged AMPKα1,

AMPKα2, TBK1s, and IRF3s was lysed using the modified MLB lysis buffer (20 mM Tris-Cl, 200

mM NaCl, 10 mM NaF, 1 mM NaV$_2$O$_4$, 1% NP-40, 20 mM β-glycerophosphate, and protease

inhibitor, pH 7.5) [31]. Cell lysates were then subjected to immunoprecipitation using the

antibodies of anti-Flag (Sigma, F3165-5MG, 1:200 dilution), anti-Myc (CST, 2276S, 1:200

dilution), or anti-HA (Sigma, H9658, 1:200 dilution) for transfected or induced proteins. After

3-4 washes with the MLB, adsorbed proteins were resolved by SDS-PAGE (Bio-Rad) and

immunoblotting with the indicated antibodies. Cell lysates were also analyzed using SDS-PAGE

and immunoblotting to control the protein abundance.

***In vitro* Kinase Assay**

HEK293 cells were transfected with plasmids encoding Flag-, Myc-, or HA-tagged TBK1s or

AMPKα and lysed by the modified MLB lysis buffer after 24 hours of transfection.

Immunoprecipitations were performed by using anti-Flag (Sigma, F3165-5MG, 1:200 dilution),

anti-Myc (2276S, 1:200 dilution), or anti-HA (CST, 3724S, 1:200 dilution) antibodies. With two

washes by the MLB and two washes by the kinase assay buffer (200 μM ATP, 200 μM AMP, 20

mM Tris-HCl, 1 mM EGTA, 5 mM MgCl$_2$, 0.02% 2-mercapto-Ethanol, 0.03% Brij-35, and 0.2 mg

mL^{-1} BSA, PH 7.4), immunoprecipitated TBK1s and AMPKα1 were incubated in the kinase assay

buffer at 30°C for 60 min on THERMO-SHAKER (1400 rpm/min). The reaction was stopped by

adding 2 × SDS loading buffer and subjected to SDS-PAGE and specified immunoblotting.

Cellular senescence assays

To evaluate damage-induced cellular senescence, cells at approximately 60-70% confluence

were treated with Hydroxyurea (HU, 10 mM), Camptothecin (CPT, 1 μM), Etoposide (ETO, 25

μM) or Mitomycin (MMC, 1 μM) for 24 hours, replaced with fresh medium, and cultured for

another 5 days and harvested for SA-β-Gal assay, by using a cellular senescence assay kit

(Yeasen 40754ES60) and according to the manufacturer's manual. Images were acquired

using a converted fluorescence microscope from Nikon. , cells were lysed after 5-8 days

stimulations and subjected to RNA extraction and qRT-PCR assays as described in the previous

section, to detect the senescence-associated secretory phenotype (SASPs), such as the

expression of cytokines including IL-1α and IL-1β.

Senescence assays in zebrafish

Zebrafish embryos were treated with 50 nM CPT with/without 1mM AICAR or 10 μM Compound C from 1.5 days post fertilization (dpf) to 3.5 dpf and then fixed in 4% paraformaldehyde in PBS at 4°C. After 3 washes with PBS (pH 7.4 and pH 6.0) at 4°C, SA-β-Gal staining was performed according to the manual of the senescence assay kit (Yeasen 40754ES60). Animals were photographed under a dissecting microscope.

Murine allograft growth of B16 melanoma

C57BL/6 wild-type mice were maintained under specific-pathogen-free (SPF) conditions and randomly selected for tumor injection. Six- to eight-week-old mice were administered a subcutaneous injection of 1.5×10^6 B16-F10 melanoma cells and treated with or without AICAR (5 mg/kg) or once a day, Aldometanib was supplied either through oral gavaging[54]. From the ninth-day post-tumor inoculation, the injected mice were monitored for tumor growth daily as described in the protocols approved by the IACUC of Zhejiang University. Other treatment regimens included: cGAMP, 2 μg intratumoral injection every 3 days, starting on day 5 post-implantation; Lonidamine, 25 mg/kg intraperitoneal injection every 3 days, starting on day 4 post-implantation. Tumor size is presented as a square caliper measurement and was calculated based on two perpendicular diameters (mm^2). The maximum diameter of tumors is limited up to 10 mm by ethical permission, upon which the mice were sacrificed and considered as dead due to tumor burden.

Murine tumor Immunohistochemistry

For immunohistochemistry assays, tumor samples were dissected, fixed in 4% paraformaldehyde for 12 hours at 4°C, dehydrated with 30% sucrose overnight at 4°C, embedded in optimal cutting temperature compound, and immediately frozen at -80°C. The sectioned samples at 10 μm thickness were washed two times with PBS, permeabilized with 0.5% Triton X-100, blocked in 3% horse serum in PBS for 30 min, and incubated sequentially with primary antibodies, including anti-CD3 (Abcam, 16669, 1:100 dilution), anti-CD4 (eBioscience, 14976680, 1:100 dilution), anti-CD8 (eBioscience, 14080880, 1:100 dilution), and anti-F4/80 (Bio-Rad, MCA497RT, 1:100 dilution). After the sections were incubated with

Alexa-labeled secondary antibodies (Jackson, 111-095-003, 1:500 dilutions, Invitrogen,

A11006, 1:1000 dilutions) and extensively washed, they were stained with DAPI (Santa Cruz

Biotech) and mounted with ProLong^{TM} Gold antifade reagent (Invitrogen).

Immunofluorescence images were obtained and analyzed using a Zeiss LSM710 or LSM 880

confocal microscope.

Nano-liquid Chromatography/Tandem MS (Nano LC-MS/MS) Analysis

Phoenix National Proteomics Core services performed Nano LC/tandem MS analysis for

protein identification and characterization and label-free quantification. Tryptic peptides were

separated on a C18 column and analyzed by LTQ-Orbitrap Velos (Thermo). Proteins were

identified using the National Center for Biotechnology Information search engine against the

human or mouse RefSeq protein databases. The mass spectrometry proteomics data of TBK1

modifications by AMPK have been deposited to the ProteomeXchange Consortium

(http://proteomecentral.proteomexchange.org) via the iProX partner repository with the

dataset identifier PXD033926.

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantitative data are presented as the mean ± standard error of the mean (SEM) from at least

three independent experiments. When appropriate, the statistically significant differences

between multiple comparisons were analyzed using the one-way or two-way ANOVA test with

Bonferroni correction. Differences were considered significant at $p<0.05$. All samples were

included in the analyses if preserved and properly processed, and no samples or animals were

excluded except for zebrafish with conventional injection damage. No statistical method was

used to predetermine sample size, and all experiments except those involving animals were

not randomized. Immunoblotting, reporter assays, and qRT-PCR experiments have been

repeated a minimum of three times independently to ensure reproducibility. The investigators

were not blinded to allocation during experiments and outcome assessment.

REFERENCES

De Martino, M., Rathmell, J. C., Galluzzi, L. & Vanpouille-Box, C. Cancer cell metabolism and antitumour immunity. *Nat Rev Immunol* **24**, 654-669 (2024). https://doi.org/10.1038/s41577-024-01026-4

Zhang, Q. *et al.* AMPK directly phosphorylates TBK1 to integrate glucose sensing into innate immunity. *Mol Cell* **82**, 4519-4536 e4517 (2022). https://doi.org/10.1016/j.molcel.2022.10.026

Kao, K. C., Vilbois, S., Tsai, C. H. & Ho, P. C. Metabolic communication in the tumour-immune microenvironment. *Nat Cell Biol* **24**, 1574-1583 (2022). https://doi.org/10.1038/s41556-022-01002-x

Hamanaka, R. B. & Chandel, N. S. Targeting glucose metabolism for cancer therapy. *J Exp Med* **209**, 211-215 (2012). https://doi.org/10.1084/jem.20120162

Galluzzi, L., Kepp, O., Vander Heiden, M. G. & Kroemer, G. Metabolic targets for cancer therapy. *Nat Rev Drug Discov* **12**, 829-846 (2013). https://doi.org/10.1038/nrd4145

Hay, N. Reprogramming glucose metabolism in cancer: can it be exploited for cancer therapy? *Nat Rev Cancer* **16**, 635-649 (2016). https://doi.org/10.1038/nrc.2016.77

Roers, A., Hiller, B. & Hornung, V. Recognition of Endogenous Nucleic Acids by the Innate Immune System. *Immunity* **44**, 739-754 (2016). https://doi.org/10.1016/j.immuni.2016.04.002

Chen, Q., Sun, L. & Chen, Z. J. Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing. *Nat Immunol* **17**, 1142-1149 (2016). https://doi.org/ni.3558 [pii]
10.1038/ni.3558

9 Chan, Y. K. & Gack, M. U. RIG-I-like receptor regulation in virus infection and immunity. *Curr Opin Virol* **12**, 7-14 (2015). https://doi.org/S1879-6257(15)00005-X [pii]
10.1016/j.coviro.2015.01.004

Takeuchi, O. & Akira, S. Pattern recognition receptors and inflammation. *Cell* **140**, 805-820 (2010). https://doi.org/10.1016/j.cell.2010.01.022

Zhang, Q., Chen, C., Xia, B. & Xu, P. Chemical regulation of the cGAS-STING pathway. *Curr Opin Chem Biol* **69**, 102170 (2022). https://doi.org/10.1016/j.cbpa.2022.102170

Gao, P. *et al.* Cyclic [G(2',5')pA(3',5')p] is the metazoan second messenger produced by DNA-activated cyclic GMP-AMP synthase. *Cell* **153**, 1094-1107 (2013). https://doi.org/10.1016/j.cell.2013.04.046

Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* **339**, 786-791 (2013). https://doi.org/10.1126/science.1232458

Hu, M. M. & Shu, H. B. Innate Immune Response to Cytoplasmic DNA: Mechanisms and Diseases. *Annu Rev Immunol* **38**, 79-98 (2020). https://doi.org/10.1146/annurev-immunol-070119-115052

Ablasser, A. *et al.* cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that activates STING. *Nature* **498**, 380-384 (2013). https://doi.org/nature12306 [pii]
10.1038/nature12306

16 Wu, J. *et al.* Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* **339**, 826-830 (2013). https://doi.org/10.1126/science.1229963

17 Zhang, X. *et al.* Cyclic GMP-AMP containing mixed phosphodiester linkages is an endogenous high-affinity ligand for STING. *Mol Cell* **51**, 226-235 (2013). https://doi.org/10.1016/j.molcel.2013.05.022

18 Ishikawa, H. & Barber, G. N. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* **455**, 674-678 (2008). https://doi.org/10.1038/nature07317

19 Zhong, B. *et al.* The adaptor protein MITA links virus-sensing receptors to IRF3 transcription factor activation. *Immunity* **29**, 538-550 (2008). https://doi.org/10.1016/j.immuni.2008.09.003

20 Sun, W. *et al.* ERIS, an endoplasmic reticulum IFN stimulator, activates innate immune signaling through

595   dimerization. *Proc Natl Acad Sci U S A* **106**, 8653-8658 (2009). https://doi.org/0900850106 [pii]

596 10.1073/pnas.0900850106

597 21 Zhang, D. *et al.* A non-canonical cGAS-STING-PERK pathway facilitates the translational program critical

598   for senescence and organ fibrosis. *Nat Cell Biol* **24**, 766-782 (2022). https://doi.org/10.1038/s41556-

599   022-00894-z

600 22 Gui, X. *et al.* Autophagy induction via STING trafficking is a primordial function of the cGAS pathway.

601   *Nature* **567**, 262-266 (2019). https://doi.org/10.1038/s41586-019-1006-9

602 23 Liu, S. *et al.* Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3

603   activation. *Science* **347**, aaa2630 (2015). https://doi.org/10.1126/science.aaa2630

604 24 Zhang, C. *et al.* Structural basis of STING binding with and phosphorylation by TBK1. *Nature* **567**, 394-

605   398 (2019). https://doi.org/10.1038/s41586-019-1000-2

606 25 Sharma, S. *et al.* Triggering the interferon antiviral response through an IKK-related pathway. *Science*

607   **300**, 1148-1151 (2003). https://doi.org/10.1126/science.1081315

608 26 Fitzgerald, K. A. *et al.* IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nat*

609   *Immunol* **4**, 491-496 (2003). https://doi.org/10.1038/ni921

610 27 Wild, P. *et al.* Phosphorylation of the autophagy receptor optineurin restricts Salmonella growth.

611   *Science* **333**, 228-233 (2011). https://doi.org/10.1126/science.1205405

612 28 Pilli, M. *et al.* TBK-1 promotes autophagy-mediated antimicrobial defense by controlling

613   autophagosome  maturation.  *Immunity*  **37**,  223-234  (2012).

614   https://doi.org/10.1016/j.immuni.2012.04.015

615 29 Meng, F. *et al.* Induced phase separation of mutant NF2 imprisons the cGAS-STING machinery to

616   abrogate  antitumor  immunity.  *Mol*  *Cell*  **81**,  4147-4164  e4147  (2021).

617   https://doi.org/10.1016/j.molcel.2021.07.040

618 30 Chen, S. *et al.* TBK1-Mediated DRP1 Targeting Confers Nucleic Acid Sensing to Reprogram Mitochondrial

619   Dynamics  and  Physiology.  *Mol*  *Cell*  **80**,  810-827  e817  (2020).

620   https://doi.org/10.1016/j.molcel.2020.10.018

621 31 Xu, P. *et al.* Innate antiviral host defense attenuates TGF-beta function through IRF3-mediated

622   suppression  of  Smad  signaling.  *Mol*  *Cell*  **56**,  723-737  (2014).

623   https://doi.org/10.1016/j.molcel.2014.11.027

624 32 Dou, Z. *et al.* Cytoplasmic chromatin triggers inflammation in senescence and cancer. *Nature* **550**, 402-

625   406 (2017). https://doi.org/10.1038/nature24050

626 33 Yang, H., Wang, H., Ren, J., Chen, Q. & Chen, Z. J. cGAS is essential for cellular senescence. *Proc Natl*

627   *Acad Sci U S A* **114**, E4612-E4620 (2017). https://doi.org/10.1073/pnas.1705499114

628 34 Gluck, S. *et al.* Innate immune sensing of cytosolic chromatin fragments through cGAS promotes

629   senescence. *Nat Cell Biol* **19**, 1061-1070 (2017). https://doi.org/10.1038/ncb3586

630 35 Mackenzie, K. J. *et al.* cGAS surveillance of micronuclei links genome instability to innate immunity.

631   *Nature* **548**, 461-465 (2017). https://doi.org/10.1038/nature23449

632 36 Petrasek, J. *et al.* STING-IRF3 pathway links endoplasmic reticulum stress with hepatocyte apoptosis in

633   early alcoholic liver disease. *Proc* *Natl* *Acad* *Sci* *U* *S* *A* **110**, 16544-16549 (2013).

634   https://doi.org/10.1073/pnas.1308331110

635 37 Cerboni, S. *et al.* Intrinsic antiproliferative activity of the innate sensor STING in T lymphocytes. *J Exp*

636   *Med* **214**, 1769-1785 (2017). https://doi.org/jem.20161674 [pii]

637 10.1084/jem.20161674

638 38 Melki, I. *et al.* Disease-associated mutations identify a novel region in human STING necessary for the

control of type I interferon signaling. *J Allergy Clin Immunol* **140**, 543-552 e545 (2017). https://doi.org/S0091-6749(16)31438-5 [pii]

10.1016/j.jaci.2016.10.031

Samson, N. & Ablasser, A. The cGAS-STING pathway and cancer. *Nat Cancer* **3**, 1452-1463 (2022). https://doi.org/10.1038/s43018-022-00468-w

Chin, E. N., Sulpizio, A. & Lairson, L. L. Targeting STING to promote antitumor immunity. *Trends Cell Biol* (2022). https://doi.org/10.1016/j.tcb.2022.06.010

Kwon, J. & Bakhoum, S. F. The Cytosolic DNA-Sensing cGAS-STING Pathway in Cancer. *Cancer Discov* **10**, 26-39 (2020). https://doi.org/10.1158/2159-8290.CD-19-0761

Chen, C. & Xu, P. Cellular functions of cGAS-STING signaling. *Trends Cell Biol* **33**, 630-648 (2023). https://doi.org/10.1016/j.tcb.2022.11.001

Bakhoum, S. F. *et al.* Chromosomal instability drives metastasis through a cytosolic DNA response. *Nature* **553**, 467-472 (2018). https://doi.org/10.1038/nature25432

Chen, Q. *et al.* Carcinoma-astrocyte gap junctions promote brain metastasis by cGAMP transfer. *Nature* **533**, 493-498 (2016). https://doi.org/10.1038/nature18268

Hong, C. *et al.* cGAS-STING drives the IL-6-dependent survival of chromosomally instable cancers. *Nature* **607**, 366-373 (2022). https://doi.org/10.1038/s41586-022-04847-2

Li, S. *et al.* STING-induced regulatory B cells compromise NK function in cancer immunity. *Nature* **610**, 373-380 (2022). https://doi.org/10.1038/s41586-022-05254-3

Concepcion, A. R. *et al.* The volume-regulated anion channel LRRC8C suppresses T cell function by regulating cyclic dinucleotide transport and STING-p53 signaling. *Nat Immunol* **23**, 287-302 (2022). https://doi.org/10.1038/s41590-021-01105-x

Zhou, R. *et al.* TBK1-Zyxin signaling controls tumor-associated macrophage recruitment to mitigate antitumor immunity. *EMBO J* **43**, 4984-5017 (2024). https://doi.org/10.1038/s44318-024-00244-9

Hardie, D. G. & Ashford, M. L. AMPK: regulating energy balance at the cellular and whole body levels. *Physiology (Bethesda)* **29**, 99-107 (2014). https://doi.org/10.1152/physiol.00050.2013

Hardie, D. G., Ross, F. A. & Hawley, S. A. AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat Rev Mol Cell Biol* **13**, 251-262 (2012). https://doi.org/10.1038/nrm3311

Lin, S. C. & Hardie, D. G. AMPK: Sensing Glucose as well as Cellular Energy Status. *Cell Metab* **27**, 299-313 (2018). https://doi.org/10.1016/j.cmet.2017.10.009

Zhang, C. S. *et al.* The lysosomal v-ATPase-Ragulator complex is a common activator for AMPK and mTORC1, acting as a switch between catabolism and anabolism. *Cell Metab* **20**, 526-540 (2014). https://doi.org/10.1016/j.cmet.2014.06.014

Zhang, C. S. *et al.* Fructose-1,6-bisphosphate and aldolase mediate glucose sensing by AMPK. *Nature* **548**, 112-116 (2017). https://doi.org/10.1038/nature23275

Zhang, C. S. *et al.* The aldolase inhibitor aldometanib mimics glucose starvation to activate lysosomal AMPK. *Nat Metab* **4**, 1369-1401 (2022). https://doi.org/10.1038/s42255-022-00640-7

Garcia, D. & Shaw, R. J. AMPK: Mechanisms of Cellular Energy Sensing and Restoration of Metabolic Balance. *Mol Cell* **66**, 789-800 (2017). https://doi.org/10.1016/j.molcel.2017.05.032

Gonzalez, A., Hall, M. N., Lin, S. C. & Hardie, D. G. AMPK and TOR: The Yin and Yang of Cellular Nutrient Sensing and Growth Control. *Cell Metab* **31**, 472-492 (2020). https://doi.org/10.1016/j.cmet.2020.01.015

Mouchiroud, L., Eichner, L. J., Shaw, R. J. & Auwerx, J. Transcriptional coregulators: fine-tuning metabolism. *Cell Metab* **20**, 26-40 (2014). https://doi.org/10.1016/j.cmet.2014.03.027

58   Zhao, P. *et al.* TBK1 at the Crossroads of Inflammation and Energy Homeostasis in Adipose Tissue. *Cell*
172, 731-743 e712 (2018). https://doi.org/10.1016/j.cell.2018.01.007

59   Klil-Drori, A. J., Azoulay, L. & Pollak, M. N. Cancer, obesity, diabetes, and antidiabetic drugs: is the fog
clearing? *Nat Rev Clin Oncol* **14**, 85-99 (2017). https://doi.org/10.1038/nrclinonc.2016.120

60   Rodrigues Mantuano, N. *et al.* Hyperglycemia Enhances Cancer Immune Evasion by Inducing Alternative
Macrophage Polarization through Increased O-GlcNAcylation. *Cancer Immunol Res* **8**, 1262-1272 (2020).
https://doi.org/10.1158/2326-6066.CIR-19-0904

61   Incio, J. *et al.* Obesity-Induced Inflammation and Desmoplasia Promote Pancreatic Cancer Progression
and Resistance to Chemotherapy. *Cancer Discov* **6**, 852-869 (2016). https://doi.org/10.1158/2159-
8290.CD-15-1177

62   Clements, V. K. *et al.* Frontline Science: High fat diet and leptin promote tumor progression by inducing
myeloid-derived   suppressor   cells.   *J   Leukoc   Biol*   **103**,   395-407   (2018).
https://doi.org/10.1002/JLB.4HI0517-210R

63   Keerthana, C. K. *et al.* The role of AMPK in cancer metabolism and its impact on the immunomodulation
of   the   tumor   microenvironment.   *Front   Immunol*   **14**,   1114582   (2023).
https://doi.org/10.3389/fimmu.2023.1114582

64   Reinfeld, B. I. *et al.* Cell-programmed nutrient partitioning in the tumour microenvironment. *Nature* **593**,
282-288 (2021). https://doi.org/10.1038/s41586-021-03442-1

65   Ramakrishnan, G., Terry, A. R., Nogueira, V., Magdy, A. & Hay, N. Deletion of AMP-activated protein
kinase impairs metastasis and is rescued by ROS scavenging or ectopic CD36 expression. *Cell Rep* **44**,
115183 (2025). https://doi.org/10.1016/j.celrep.2024.115183

66   Corton, J. M., Gillespie, J. G., Hawley, S. A. & Hardie, D. G. 5-aminoimidazole-4-carboxamide
ribonucleoside. A specific method for activating AMP-activated protein kinase in intact cells? *Eur J*
Biochem **229**, 558-565 (1995). https://doi.org/10.1111/j.1432-1033.1995.tb20498.x

67   McArthur, K. *et al.* BAK/BAX macropores facilitate mitochondrial herniation and mtDNA efflux during
apoptosis. *Science* **359** (2018). https://doi.org/10.1126/science.aao6047

68   Rongvaux, A. *et al.* Apoptotic caspases prevent the induction of type I interferons by mitochondrial DNA.
Cell **159**, 1563-1577 (2014). https://doi.org/10.1016/j.cell.2014.11.037

69   White, M. J. *et al.* Apoptotic caspases suppress mtDNA-induced STING-mediated type I IFN production.
Cell **159**, 1549-1562 (2014). https://doi.org/10.1016/j.cell.2014.11.036

70   Chiappinelli, K. B. *et al.* Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA
Including   Endogenous   Retroviruses.   *Cell*   **162**,   974-986   (2015).
https://doi.org/10.1016/j.cell.2015.07.011

71   Sasaki, N. *et al.* RNA sensing induced by chromosome missegregation augments antitumor immunity.
Mol Cell (2024). https://doi.org/10.1016/j.molcel.2024.11.025

72   Gulen, M. F. *et al.* Signalling strength determines proapoptotic functions of STING. *Nat Commun* **8**, 427
(2017). https://doi.org/10.1038/s41467-017-00573-w

73   Wu, S. *et al.* HER2 recruits AKT1 to disrupt STING signalling and suppress antiviral defence and
antitumour immunity. *Nat Cell Biol* **21**, 1027-1040 (2019). https://doi.org/10.1038/s41556-019-0352-z

74   Floridi, A. *et al.* Lonidamine, a selective inhibitor of aerobic glycolysis of murine tumor cells. *J Natl*
Cancer Inst **66**, 497-499 (1981).

75   Roberts, D. J., Tan-Sah, V. P., Ding, E. Y., Smith, J. M. & Miyamoto, S. Hexokinase-II positively regulates
glucose starvation-induced autophagy through TORC1 inhibition. *Mol Cell* **53**, 521-533 (2014).
https://doi.org/10.1016/j.molcel.2013.12.019

76 Wu, Y. Q. *et al.* Low glucose metabolite 3-phosphoglycerate switches PHGDH from serine synthesis to
p53 activation to control cell fate. *Cell Res* **33**, 835-850 (2023). https://doi.org/10.1038/s41422-023-
00874-4
77 Ghosh, M. *et al.* Mutant p53 suppresses innate immune signaling to promote tumorigenesis. *Cancer Cell*
39, 494-508 e495 (2021). https://doi.org/10.1016/j.ccell.2021.01.003
78 Zhang, Q. *et al.* Hippo signalling governs cytosolic nucleic acid sensing through YAP/TAZ-mediated TBK1
blockade. *Nat Cell Biol* **19**, 362-374 (2017). https://doi.org/10.1038/ncb3496
79 Liu, S. *et al.* Lck/Hck/Fgr-Mediated Tyrosine Phosphorylation Negatively Regulates TBK1 to Restrain
Innate   Antiviral   Responses.   *Cell   Host   Microbe*   **21**,   754-768   e755   (2017).
https://doi.org/10.1016/j.chom.2017.05.010
737
738

    FIGURE LEGENDS

Figure 1. Glucose deficiency promotes intratumoral cGAS-STING signaling. (A-B), The

intratumoral glucose deficiency state in B16-F10 melanoma was validated through

immunofluorescence of phospho-ACC1 (pACC), a primary substrate of AMPK, by which an

evident signal of pACC was seen in the edge of melanoma (A). AMPK agonist AICAR treatment

was set as the positive control (B). Scale bars=50 μm. **(C-E),** Cytosolic dsDNA sensing was

evaluated in PMs (C), MEFs (D), and DLD1 (E) under 2-DG or AICAR (AMPK agonist)

administration in response to STING agonist cGAMP, as indicated by immunoblotting using

antibodies recognizing active forms of TBK1 and IRF3. **(F-G)**, Sensing of mtDNAs, released

upon the treatment of ABT-737, qVD-OPH, and S63845 (AQS), was similarly augmented by

glucose depletion, revealed by the phosphorylation level of TBK1. **(H)**, The effects of AICAR

administration on TpdAdT-induced DNA sensing were evaluated in DLD1, representing

enhanced dsDNA sensing to double-stranded DNA. ***Applied to figures 1-6***: unless specified,

n=3 independent experiments (mean±SEM); *, P<0.05, **, P<0.01, and ***, P<0.001, by

statistical analysis of the indicated comparison with ANOVA and Bonferroni correction.

Figure 2. AMPK facilitates cGAS-STING signaling in cancer cells by TBK1 S511

phosphorylation. (A), Double knockout (dKO) of AMPK α1 and α2 impaired the activation of

TBK1 and IRF3 following DMXAA stimulation in B16-F10 melanoma cells. **(B)**, The genetic

ablation of AMPK α1 and α2 subunits reduced TpIC-induced cytosolic RNA sensing in HCT116,

revealed by a significant decrease in phosphorylation of TBK1 and IRF3. **(C)**,

Coimmunoprecipitation assay revealed coexpression of AMPKα1 or caAMPKα1 (activated

AMPKα1) enhanced this association of TBK1 and IRF3 2SA. **(D)**, Reporter assay showed that

the expression of AMPK α1 and α2 subunits strongly promoted TBK1-stimulated IRF3

transactivation with IFNβ promoter in HEK293 cells. **(E)**, Mass spectrometry analysis identified

that AMPKα1 phosphorylated the S511 residue of TBK1, and the phosphorylation was

completely inhibited by the AMPK inhibitor dorsomorphin (Compound C). **(F)**, caAMPKα1 and

AMPKa1 kinase domain directly phosphorylated TBK1 at S511 in vitro kinase assay. **(G-H)**,

TBK1 mutants S511A (resistant to AMPK modification) exhibited significantly reduced activity

IRF3-responsive reporter assays (G) and a weaker interaction with IRF3 2SA (H). Meanwhile,

TBK1 S511E (mimics AMPK-phosphorylated TBK1) demonstrated a strong potential to bind

and activate IRF3. **(I-J)**, Generated DLD1 knock-in (KI) clones with mutations that either

prevent (S511A) or mimic (S511D) AMPK-mediated modification using CRISPR-based genome

editing. cGAMP-stimulated cGAS-STING pathway was weakened in TBK1 S511A KI cells (I) but

strengthened in S511D KI cells (J).

Figure 3. AMPK drives STING-induced senescence and apoptosis of cancer cells and

organoids. (A-B), Chemical induction of DNA damage in DLD1 gut epithelial cells by CPT,

mitomycin C (MMC), etoposide (ETO), and hydroxyurea (HU) treatment led to a robust

phenotype of cellular senescence, as assessed by SA-β-Gal staining (A) and SASP features,

including the mRNA expression of IL1α and IL1β (B). Genetic ablation of AMPKα1/α2 in DLD1

cells prevented DNA sensing-initiated cellular senescence (A). Conversely, AICAR treatment

significantly enhanced DNA damage-induced cellular senescence (B). Scale bars = 100 μm. **(C)**,

Energy deficiency stimulation of AMPK by 2-DG or AICAR led to an exaggerated phenotype of

cellular senescence in DLD1 cells, particularly upon HU-induced DNA damage. Scale bars = 100

μm. ***, P<0.001, by ANOVA with Bonferroni correction. **(D)**, Cellular senescence was

assessed in zebrafish embryos (n=30) subjected to chemical-induced damage by incubation

with CPT (50 nM) and to SA-β-Gal staining. A strong phenotype of cellular senescence was

detected in the heads and tails of zebrafish; this phenotype was mildly upregulated by AICAR

treatment but was mitigated by dorsomorphin. Scale bars = 500 μm. **(E)**, FACS analyses of early

and late apoptotic cells among DLD1 cells revealed that activation of AMPK by AICAR

promoted cell apoptosis induced by cytosolic DNA sensing, which triggered by DNA damage

inducer HU, but inhibition of AMPK by dorsomorphin suppressed HU-induced apoptosis. **(F)**,

Intestinal crypts were isolated from mice at 6 weeks of age and cultured *in vitro* in the absence

or presence of the AMPK agonist AICAR. PI staining revealed that cytosolic DNA sensing

stimulated by the DNA damage inducer HU resulted in robust apoptosis of intestinal cells in

organoids, and AICAR exaggerated this effect. Scale bars=50 μm. n=15 intestinal organoids

examined (mean±SEM). *, P=0.002, compared with wild-type organoids (ANOVA with

Bonferroni correction).

Figure 4. AMPK-TBK1 signaling facilitates STING-mediated antitumor immunity in melanomas. (A-C), The STING SAVI mutant (R281Q) was constitutively active (caSTING). The volumes (A), weights (B), and photos (C) of the subcutaneously B16-F10 melanoma growth in wild-type C57BL/6 mice revealed a markedly decreased growth in melanomas expressing the STING SAVI mutant (R281Q). AICAR administration further increased this effect of STING signaling, resulting in robust tumor growth arrest. n=12 inoculations per group (mean ± SEM), measured 7 days post tumor inoculation (dpti). **(D-E)**, The administration of AICAR with caSTING profoundly increased immune cell infiltration, particularly the infiltration of the CD4$^+$ and CD8$^+$ T cells into melanomas, as detected by Immunohistochemistry (IHC) upon STING SAVI mutant expression (D). The statistics for the immunohistochemistry-positive cells were calculated (E). Scale bars=20 μm. **, P<0.01, and ***, P<0.001, by ANOVA with Bonferroni correction. **(F-H)**, B16-F10 melanoma cells were implanted subcutaneously into TBK1$^{WT/WT}$ or TBK1$^{S511E/S511E}$ C57BL/6 mice with intratumoral injection of cGAMP. Volumes (F), weights (G), and photos (H) of melanoma revealed that melanoma growth was markedly attenuated in these TBK1$^{S511E/S511E}$ mice. **(I-J)**, Immunohistochemistry (I), and statistics (J) showed B16-F10 melanoma from TBK1$^{S511E/S511E}$ mice accompanied with the increased immune cell infiltration. Stimulation in TBK1$^{S511E/S511E}$ mice by cGAMP did not further enhance the infiltrated T cells and antitumor immunity effect. Scale bars=20 μm.

Figure 5. AMPK agonists synergize well with cGAMP to promote antitumor immunity in melanomas. (A-C), The weights (A), volumes (B), and photos (C) of the subcutaneously B16-F10 melanoma growth in wild-type C57BL/6 mice with administration of cGAMP or AICAR. AICAR administration further promoted the antitumor immunity effect of cGAMP-induced STING signaling, inhibiting tumor growth. n=6 inoculations per group (mean ± SEM). **(D-F)**, The weights (D), volumes (E), and photos (F) of the subcutaneously B16-F10 melanoma growth in wild-type C57BL/6 mice with administration of Aldometanib and cGAMP. Adding AICAR enhanced the suppression of tumor growth induced by cGAMP. n=4 inoculations per group (mean ± SEM). **(G-I)**, Administration of Aldometanib alone inhibited the growth of AMPK dKO B16-F10 tumors in WT mice. The weights (G), volumes (H), and photos (I) showed similar tumor sizes of AMPK dKO B16-F10 tumors with the treatment of Aldometanib or cGAMP. n=8

inoculations per group (mean ± SEM).

Figure 6. Disrupting aerobic glycolysis potentiates antitumor immunity through the AMPK-
TBK1 cascade. The administration of Lonidamine (Hexokinase inhibitor) with cGAMP
profoundly inhibited the growth of B16-F10 melanoma, revealed by the volumes (D), weights
(E), and photos (F) of B16-F10 melanoma. (D) n=10 inoculations per group (mean ± SEM). (E)
n=8 inoculations per group (mean ± SEM). **(G-H)**, Enhanced infiltration of CD3$^+$, CD4$^+$, and CD8$^+$
T lymphocytes were imaged in B16-F10 tumors with the administration of Lonidamine and
cGAMP; Scale bars, 20 μm. (H) n=3 per group (mean ± SEM). **(I)**, Immunohistochemistry
showed that Lonidamine enhanced cGAMP-induced cGAS-STING signaling in tumors, as
evidenced by increased levels of phosphorylated STING (pSTING S365) and phosphorylated
TBK1 (pTBK1 S172) in B16-F10 melanomas; Scale bars=20 μm.

Figure 1

Figure 2

Figure 4

A

C57BL/6 Mice/8 Inoculations

B

B16-F10 Inoculation

C

C57BL/B16-F10 Inoculation

D

Mice B16-F10 Inoculation

E

Mice B16-F10 Inoculation

F

Mice B16-F10 Inoculation

Figure 1

Figure 1

Figure 1

Figure 1

Figure 1

Figure 2

Figure 2

B

Figure 2

C

IP: Myc-IRF3 2SA
IB: TBK1

Figure 2

F

Figure 2

G

Figure 2

H

Figure 2

I

Figure 2

J

Suppl. Table 1_Oligos Used in Study

qPCR Primers

qPCR Primers	Sequence	Species	Company
IL-1a-Forward Primer	5' - AGTGCTGCTGAAGGAGATGCCTGA - 3'	human	biosune
IL-1a-Reverse Primer	5' - CCCCTGCCAAGCACACCCAGTA - 3'	human	biosune
IL-1b-Forward Primer	5' - ATGATGGCTTATTACAGTGGCAA - 3'	human	biosune
IL-1b-Reverse Primer	5' - GTCGGAGATTCGTAGCTGGA - 3'	human	biosune
RPL19-Forward Primer	5' - ATGTATCACAGCCTGTACCTG - 3'	human	biosune
RPL19-Reverse Primer	5' - TTCTTGGTCTCTTCCTCCTTG - 3'	human	biosune

Primers

Cas9 Primers	Sequence	Species	Company
AMPKα1-Cas9-Forward Primer	5' - CACCGAAAGTTTGAGTGCTCAGAAG - 3'	human	biosune
AMPKα1-Cas9-Reverse Primer	5' - AAACCTTCTGAGCACTCAAACTTTC - 3'	human	biosune
AMPKα2-Cas9-Forward Primer	5' - CACCGTCAATTAACAGGCCATAAAG - 3'	human	biosune
AMPKα2-Cas9-Reverse Primer	5' - AAACCTTTATGGCCTGTTAATTGAC - 3'	human	biosune
TBK1-S511A/D-Knock-In-Forward Primer	5' - CACCGTATTTAGCTTTCCAGTTCTC - 3'	human	biosune
TBK1-S511A/D-Knock-In-Reverse Primer	5' - AAACGAGAACTGGAAAGCTAAATAC - 3'	human	biosune

Suppl. Table 2_ List of Recombinant DNA

DNA	SOURCE	IDENTIFIER
pRK5-HA-TBK1 S511A	This paper	N/A
pRK5-HA-TBK1 S511D	This paper	N/A
pRK5-HA-TBK1 S511E	This paper	N/A
pRK5-Flag-TBK1 S511A	This paper	N/A
pRK5-Flag-TBK1 S511D	This paper	N/A
pRK5-Flag-TBK1 S511E	This paper	N/A
pRK5-HA-AMPKα1	This paper	N/A
pRK5-HA-AMPKα1 T172D	This paper	N/A
pRK5-HA-AMPKα1 kinase domain (1-312)	This paper	N/A
pRK5-Flag-AMPKα1	This paper	N/A
pRK5-Flag-TBK1	Lab stored	N/A
pRK5-HA-TBK1	Lab stored	N/A
pRK5-Myc-TBK1	Lab stored	N/A
pRK5-Myc-IRF3	Lab stored	N/A
pRK5-Myc-IRF3 2SA	Lab stored	N/A
pIFNβ_Luc	Lab stored	N/A
p5xISRE_Luc	Lab stored	N/A
gRNA	Lab stored	N/A
Cas9-2A-GFP	Lab stored	N/A
pMD18	Lab stored	N/A
pX330	Addgene	Cat#42230

Suppl. Table 3_Statistics Source Data

(See details in the following sheets)