Pooled screening identifies combinatorial CAR signaling domains for next-generation CAR-M

immunotherapies

Yuxin Wang[1#], Shiman Zuo[1#], Hanjing Bao[1], Zehui Zhang[1], Yijun Chen[2], Wenlong Zhang[2], Qi Liu[1],

Yan Lu[1], Yahong Huang[1], Wei Zheng[1], Nanfei Yang[1], Lupeng Ye[3], Pingping Shen[1,2,4*]

[1]State Key Laboratory of Pharmaceutical Biotechnology and Department of Urology, Nanjing Drum

Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, School of Life Sciences,

Nanjing University, Nanjing 210023, China.

[2]Macera Therapeutics, Nanjing 210032, China.

[3]Institute of Modern Biology, Nanjing University, Nanjing 210023, China.

[4]Shenzhen Research Institute of Nanjing University, Shenzhen 518000, China.

# Equally Contributing Authors.

* Corresponding author.

E-mail address: ppshen@nju.edu.cn (P. Shen)

Abstract

Chimeric antigen receptor-engineered macrophages (CAR-Ms) hold great promise for solid tumor immunotherapy. The intracellular domains (ICDs) of CARs determine the phenotypic output of therapeutic macrophages but remain largely unexplored. Here, we constructed a CAR library containing 131 unique signaling domains derived from native immune receptors and identified 17 ICDs that enhance macrophage phagocytosis, inflammatory responses, or tumor infiltration *in vitro* and *in vivo*. We further developed a scalable 3' barcode technology, CARode, to uniquely label and trace ICD variants within large-scale combinatorial CAR library and applied it to single-cell RNA sequencing and single-cell CAR analysis to assess the synergetic effects of ICD combinations on macrophage activation. Our approach uncovered a novel CD40-LY9-FCRL1 chimeric receptor that modulates the tumor microenvironment and improves solid tumor clearance. In conclusion, our findings demonstrate that pooled screening can accelerate the discovery of complex ICD constructs, providing a powerful platform for engineering macrophage-based immunotherapies.

Introduction

Chimeric antigen receptor T (CAR-T) cell therapy has demonstrated remarkable efficacy in the treatment of hematological malignancies [1, 2]. However, its application in solid tumors has encountered significant challenges, particularly due to limited tumor infiltration and the presence of immunosuppressive tumor microenvironment (TME) [3, 4]. Recently, CAR-engineered macrophages (CAR-Ms) were introduced as an alternative to CAR-T cell therapy for solid tumors [5, 6]. Macrophages have inherent abilities to penetrate solid tumors and modulate the TME through broad immune responses, such as phagocytosis, antigen presentation, and pro-inflammatory cytokine

secretion [5, 7, 8]. To harness this potential, CAR constructs designed to enhance the phagocytic,

pro-inflammatory, and tumor-infiltrating functions of macrophages are currently being investigated and

have shown promising anti-tumor effects in solid tumors [9-11]. Thus, CAR-engineered macrophages

with enhanced anti-tumor functions hold great potential for revolutionizing immunotherapeutic

approaches for solid tumors.

The anti-tumor efficacy of CAR-Ms is strongly modulated by the intracellular domains (ICDs) of

the CAR constructs [12]. Various engineering strategies have been developed for engineering the

signaling domains of CAR-Ms. One current strategy involves tuning CAR signaling via the Fc receptor

common gamma chain (FcRγ), which facilitates antibody-dependent cellular phagocytosis (ADCP) [13,

14]. The CD3ζ cytosolic domain, a canonical signaling domain for T cell activation, showed functional

similarity with FcRγ signaling in CAR-Ms [10, 15], suggesting the compatibility of receptors that are

not typically expressed by macrophages. The second strategy uses Toll-like receptor signaling to drive

the M1-like phenotype of therapeutic macrophages [11, 16]. This pro-inflammatory signaling has been

combined with the CD3ζ phagocytic domain in CAR-Ms to induce a bifunctional phenotype [11].

Moreover, we previously constructed a CAR bearing the CD147 signaling domain to activate the

expression of matrix metalloproteinases (MMPs) to degrade the extracellular matrix (ECM) of solid

tumors [9]. Taken together, different ICDs induce distinct phenotypes and functions of therapeutic

macrophages. However, the optimal signaling domains required for CAR-M activation and the

potential of combinatorial signaling to elicit multifunctional macrophage responses remain largely

unexplored.

Advances in large-scale pooled screens, such as genome-wide CRISPR screens and

gain-of-function screens, have made it possible to test gene function in a high-throughput manner [17].

Recently, this technology has been applied to CAR engineering, allowing for systematic identification

of ICDs, signaling motifs, and transcription factors (TFs) that increase CAR-T cell efficacy [18-22].

However, such high-throughput approaches have yet to be applied to CAR-M optimization. Several

technical challenges hinder the direct application of large-scale CAR screening to macrophage-based

therapy. Unlike T cells, primary macrophages exhibit limited proliferative capacity *in vitro* [23],

making it difficult to maintain and expand large cell libraries required for pooled screening.

Additionally, macrophages have strong antiviral activity [24], which restricts the efficiency of

viral-based CAR library delivery. The large size and complexity of CAR libraries further complicate

the screening process, as each CAR variant must be effectively expressed and evaluated in parallel.

Therefore, novel screening strategies designed for macrophages are required to address current

limitations and advance the development of CAR constructs for macrophage-based immunotherapy.

Here, we developed a 131-member natural receptor signaling domain CAR (NaSDC) library to

screen the signaling domains of immune response-related receptors as independent ICDs for CAR

constructs to promote macrophage activation and anti-tumor functions. The library was screened in the

human monocytic cell line THP-1 and validated in human monocyte-derived macrophages (hMDMs).

We revealed that 17 ICDs have distinct anti-tumor functions, including phagocytosis, pro-inflammation,

and tumor infiltration. Furthermore, we developed a large-scale combinatorial signaling domain CAR

(CoSDC) library to evaluate the synergistic effects of signaling combinations of the 17 ICDs in

promoting macrophage activation. The CoSDC library was constructed and barcoded via the CARode

(Barcode for CAR) strategy, which facilitates unique labels of combinatorial CAR constructs in a

scalable manner and is compatible with single-cell RNA sequencing (scRNA-seq). The CoSDC library

screens revealed multi-CAR constructs that improved macrophage anti-tumor functions, including a

CD40-LY9-FCRL1 chimeric receptor that modulated the TME and improved solid tumor clearance.

Overall, these studies highlight NaSDC×CoSDC screening as a powerful method to accelerate the

programming of the macrophage phenotype with enhanced anti-tumor functions.

Results

Design of CAR libraries for CAR-M screens

Here, we aimed to develop a platform that is suitable for high-throughput, pooled screening of CAR

signaling domains within macrophages. We developed two libraries to reprogram macrophage function

through single ICD CAR overexpression or combined ICD signaling (Figure 1A). The single ICD CAR

library aims to assess the role of intracellular elements of natural receptors as independent signaling

domains of CAR constructs in promoting macrophage functions, including phagocytosis, tumor

infiltration, and pro-inflammatory responses. The combinatorial signaling domain CAR library is

designed to evaluate the synergetic effects of ICD combinations on macrophage activation.

Human primary macrophages are resistant to virus-based gene transduction. Thus, we first needed

to determine the proper cell types for large-scale CAR variant screens. The human monocytic cell line

THP-1 can be differentiated into macrophages, and this approach has been used previously for

genome-wide screening [25, 26]. The infectivity of lentivirus was greater in THP-1 cells than in human

monocyte-derived   macrophages   (hMDMs)   (Figure   1B).   We   confirmed   that   THP-1-derived

macrophages expressed macrophage markers (Figure S1A) and showed the capacity to polarize to the

M1, M2 and tumor-associated macrophage (TAM) phenotypes (Figure S1B-S1D). We then transduced

CAR variants with the ICD of CD247 (CAR-3ζ) or TLR4 (CAR-t4) into THP-1 cells and observed

increased phagocytosis and pro-inflammatory gene expression in THP-1-derived macrophages in

response to CAR signaling (Figure S1E-S1I), suggesting that THP-1-derived macrophages constitute a proper model for CAR signaling screening studies.

To construct a library for the functional evaluation of natural receptor signaling domains as independent ICDs of CAR constructs within macrophages, we mined macrophage activation-related receptors from various protein families, such as Fc receptors, Toll-like receptors (TLRs), and the TNF receptor superfamily (Figure 1A). Given the homology of receptors across the immune system and the established role of the T cell-related receptor CD3ζ (encoded by *CD247*) as a functional cytosolic domain of CAR-Ms [10], we included receptors that are predominantly expressed in T cells, natural killer (NK) cells, dendritic cells (DCs), and B cells to determine whether subsets of these could engage in distinct and beneficial signaling in macrophages (Figure S2A). Taken together, a total of 131 signaling domains (Table S2) were collected and constructed within the context of a PSMA CAR (containing an anti-PSMA single-chain variable fragment, a CD8α hinge, and a CD8α transmembrane domain) (Figure S2B). The generated library is called the NaSDC library. A truncated CAR without a signaling domain was added to the library as a negative control. Each CAR variant was uniquely barcoded (Figure S2B) and sequenced to confirm the construct. We then generated NaSDC lentiviruses from the plasmid library to transduce THP-1 cells at a multiplicity of infection (MOI) of 0.3 to minimize multiple integrations per cell (Figure S2C). The ability of NaSDC CAR-Ms to bind to soluble PSMA antigen was confirmed by flow cytometry (Figure 1C). We observed no significant correlation between ICD length and abundance in the CAR-M library (Figure S2D). Together, the NaSDC library provides a robust platform for systematically screening diverse natural receptor signaling domains as independent ICD of CAR-Ms.

We next sought to construct a large-scale combinatorial signaling domain CAR (CoSDC) library

to evaluate the synergistic effects of ICD combinations in promoting macrophage activation. A

well-established method for identifying different gene variants in large-scale libraries is to use

randomized 5~20 bp barcode sequences attached to untranslated regions (UTRs) [18, 27, 28], which

require pooled PCR and long-read sequencing to match the barcode sequences with gene variants. Here,

we developed a novel, scalable method to barcode large-scale CAR libraries, namely CARode

(Barcode for CAR). Briefly, we generated constructs with a cloning cassette that was placed between

the ICD sequences and barcode sequences (Figure 1D and S3A). The cassette consisted of customized

cleavage sites and a stop codon and was flanked by linkers that served as homology arms for assembly

(Figure S3B). Each ICD was PCR amplified to combine with the cloning cassette and a unique 6-bp

barcode and then incorporated into a PSMA CAR chassis to generate a single ICD CAR (G1) library

(Figure S3A). The "2nd ICD-cassette-barcode" constructs were subsequently incorporated into the G1

library to replace the cassette of the first ICD, resulting in a second-generation (G2) CAR library with

two fused barcodes (Figure S3A). The third-generation (G3) CARs were generated in a similar way

(Figure S3A). This approach could theoretically enable more than 16 million possible combinatorial

CARodes for the G2 library and over 68 billion CARodes for G3 CARs. Given their proximity to the 3'

UTR, these tandem CARodes allow sequencing of CAR identities in single-cell transcriptomes using

commercially available 3' capture kits (Figure 1D). Additionally, the arrangement of CARodes reflect

the order of ICDs in CAR constructs, and the length of CARodes indicates the generation of CARs

(Figure 1D). In summary, CoSDC screens with CARode technology enable rapid evaluation of CAR

ICD combinations for engineered immunotherapies.

NaSDC allows for the discovery of CAR-M variants with distinct anti-tumor functions

We first aimed to identify signaling domains that trigger the phagocytosis of macrophages in the

context of CARs. We adopted a bead-based selection strategy for the rapid separation of CAR-Ms on

the basis of their ability to phagocytose magnetic substrates (Figure 2A). After incubation with

magnetic particles, NaSDC library cells were passed through a uniform magnetic field that selectively

captured magnetized cells that had ingested magnetic particles, while cells that did not phagocytose

beads passed through. Subsequently, RNA was isolated and reverse transcribed into cDNA, followed

by barcode sequencing to compare the abundance of each CAR variant in the input and output

populations.

      We integrated beta scores from the robust rank aggregation (RRA) model with overall fold

changes to evaluate CAR enrichment in the output populations. The screening was performed in

sextuplicate to enable RRA model and reduce batch effects. The results of the two algorithms were

strongly correlated (Figure 2B). Notably, we found that the previously reported pro-phagocytic

signaling domains FCER1G and CD247 were highly enriched in the output pools (Figure S4A),

suggesting that our screening method yielded reliable results. Signaling by Fc receptor superfamily had

strong effects on macrophage phagocytosis, especially FCRL1 and FCRL6, which exhibited similar

pro-phagocytosis capacity compared with FCER1G signaling (Figure S4B). Among the top positive

hits were HCST (encoding DAP10), which is required for NK cell activation and associated with T cell

responses [29], and TREML1, which can promote the aggregation of platelets [30] (Figure 2B).

Interestingly, the inhibitory signals BTLA and LILRB2, which are expressed primarily in T cells and

NK cells [31, 32] (Figure 2B), also promote macrophage phagocytosis, suggesting distinct functions of

these receptors in macrophages. Overall, eight signaling domains showed an enhanced ability to

phagocytose magnetic substrates compared with FCER1G signaling.

      We next sought to validate the pro-phagocytic function of the CAR variants in human primary

macrophages. Given that macrophages are antiviral-transfected cells, we adopted a lipid nanoparticle

(LNP) delivery system to transfer CAR-encoded mRNAs into primary human macrophages. LNPs

exhibited high gene transduction efficiency and low phenotype perturbance in the context of primary

macrophages (Figure S5A and S5B). The expression level peaked at 12 hours post-transfection and

remained stable for up to 10 days in the cells (Figure S5A). We transduced the mRNAs of the

pro-phagocytic CARs into hMDMs and confirmed their phagocytosis of the antigen-positive tumor

cells (Figure S5D and 2C). Notably, all ICD variants exhibited higher pro-phagocytic capacity

compared with CAR-FCER1G (Figure 2C). Among these variants, the CAR-FCRL6 and CAR-FCRL1

variants exhibited the best performance (Figure 2C).

Therapeutic macrophages must maintain a persistent pro-inflammatory phenotype within tumors

to support their anti-tumor functions [7]. However, macrophages are plastic cells and can shift to an

immunosuppressive phenotype in response to TME signals [7, 8]. To explore CAR variants that can

repolarize immunosuppressive TAMs to a pro-inflammatory phenotype, we performed NFκB pathway

activity screens using a fluorescent reporter (Figure 2D). The reporter was cotransduced with the

NaSDC library into THP-1 cells to generate a reporter CAR-M library. We pretreated the reporter cells

with conditioned media from prostate cancer cell cultures for two days to mimic the

immunosuppressive TME. Downregulation of iRFP fluorescence induced by tumor media was

observed (Figure S4C). The CAR-Ms were subsequently cocultured with antigen-positive target cell

lines, and iRFP670^{high} cells were sorted (Figure S4D) and subjected to barcode sequencing.

Polarization screening revealed eight signaling domains with notable impacts on macrophage

pro-inflammatory polarization (Figure 2E and S4E). We noted that both the TNF receptor superfamily

and TLR family signals promoted NFκB activation (Figure S4F), whereas chimeric receptors with TNF

superfamily intracellular domains, such as CD40, LTBR, TNFRSF1A, and TNFRSF11A, tended to

perform better in this context (Figure 2E). The transduction of TNFRSF1A-CARs and LTBR-CARs

into hMDMs resulted in limited surface expression and decreased cell viability even in the absence of

stimulation (Figure S5C and S5E), suggesting potential stimulation-independent effects. Validation

analyses revealed that the top variants can increase the pro-inflammatory phenotype and cytokine

secretion (Figure 2F and S6A-S6C).

To discover CAR signaling that can promote macrophage infiltration, we next screened the

NaSDC pool in a subcutaneous tumor model (Figure 2G). We subcutaneously (SC) injected

PSMA-positive prostate tumors and peritumorally injected NaSDC macrophages 7 days after tumor

inoculation. Four days later, the tumors were harvested, and the abundance of barcodes was measured

using barcode sequencing (Figure S7A). We computed CAR variant enrichment as described above and

merged the results from all of the mice. We observed high enrichment of CAR-Ms with LY9 and

CD247 ICDs in the tumor tissues (Figure 2H and S4G). The potential of LY9 and CD247 signaling to

increase macrophage abundance in solid tumors was validated in an independent assay in which

hMDMs bearing either CAR-LY9 or CAR-3ζ were injected around the tumor sites, and both of these

CAR-Ms were significantly enriched in target-positive tumors (Figure S7B). Validation analyses

revealed that MMP family gene expression levels increased in these CAR-Ms during coculture with

antigen-positive cells (Figure 2I, S5F, and S7C-S7E), suggesting enhanced matrix degradation capacity.

We noted that these CAR variants lack the ability to induce a pro-inflammatory response in

macrophages (Figure S7D and S7E). RNA-seq analysis revealed a transcriptional similarity of 46% in

upregulated genes between LY9 and CD247 CAR-Ms (Figure S7F). Biological process analysis

showed that CAR-LY9 and CAR-3ζ macrophages behaved more similarly in terms of cell proliferation

(Figure S7G).

In summary, we identified 17 signaling domains, including two previously reported ICDs

(FCER1G and CD247), that can elicit pro-phagocytic, pro-inflammatory, or pro-infiltrating functions in

macrophages through NaSDC screens. These functional signaling domains hold potential for guiding

the design of combinatorial signaling domain CARs that benefit to macrophage-based therapies.

CoSDC enables pooled screening of ICD combinations via scRNA-seq and scCAR-seq

We next explored ICD combinations that could trigger combinatorial functions of macrophages via the

CoSDC library. To construct the CoSDC library, we randomly integrated the 17 ICDs identified from

NaSDC screens to generate second- and third-generation CAR constructs, yielding a theoretical

diversity of 5219 barcoded signaling domain combinations. As described above, each signaling domain

receives a unique barcode sequence to de-multiplex CAR generation, ICD type and position. Barcode

sequencing revealed that the CoSDC plasmid library possessed all possible combinations, with

balanced representation (Figure S8A). We then generated lentiviruses to transduce THP-1 cells at a low

MOI to minimize multiple integrations per cell (Figure S8B). The CAR-M library was PMA-stimulated

and cocultured with antigen-positive target cells. Macrophages bearing truncated CARs were

incorporated into the pooled cocultures as negative controls. Following 24 h of coculture, CAR-Ms

were isolated by FACS (Figure S8C) and processed for scRNA-seq through the 10X Genomics pipeline

(Figure 3A). Barcoded CAR transcripts found in the scRNA-seq cDNA product were selectively

amplified and sequenced.

The transcriptomes of 13,471 CAR-Ms were obtained from scRNA-seq following data

preprocessing and filtering (Figure 3B). Among these cells, 58.7% could be assigned a single specific

CAR variant (Figure 3B), while the remaining cells were excluded because of either the absence of

CAR insertion or the presence of multiple distinct CAR constructs within the same cell. Interestingly,

we noted that CAR variants triggered different functional macrophage states upon encountering target

cells. Seven clusters were identified using unsupervised clustering (Figure 3B). Cluster c6 exhibited a

combinatorial phenotype characterized by M2-like polarization (Figure 3C) and MHC I-mediated

antigen processing and presentation (Figure 3D and S9A). Cluster c5 displayed a similar polarization

pattern to that of c6 but higher levels of MMP expression (Figure 3C and 3E). Both clusters c0 and c3

significantly upregulated the expression of MMP (Figure 3E) and phagocytosis-related genes (Figure

3F), but c3 failed to transition into an M1-like phenotype (Figure 3C) and had diminished antigen

processing and presentation ability (Figure 3D and S9A). Notably, CAR-Ms in cluster c0 exhibited

M1-like polarization, antigen processing and presentation, MMP expression, and phagocytosis (Figure

3C-F and S9A-C), suggesting that the signaling domains within c0 CARs can trigger combinatorial

anti-tumor functions of macrophages.

We next tracked the extent of enrichment for each ICD at each position in the c0 cluster.

Consistent with the standard second- and third-generation of CAR-T cell constructs [1, 33], the CD247

signaling domain tended to perform better at the C-terminal end of the CAR-M constructs (Figure 3G).

In contrast, some ICDs (e.g., TNFRSF1A and LTBR) are suitable for grafting into the

membrane-proximal intracellular space of a CAR (Figure 3G). HCST, TYROBP, and LY9 signaling

can be incorporated into any of the three intracellular positions in a third-generation CAR construct

(Figure 3G). These results demonstrate the positional preferences of different ICDs when incorporated

into a combinatorial signaling CAR.

CoSDC library-guided CAR constructs induce a favorable macrophage phenotype

To validate the CoSDC-guided CAR variants and understand their effects on macrophage activation

and anti-tumor functions, we selected and cloned eight CAR variants from the CoSDC library, based on

the ICD features of cluster c0 (Figure 4A). Each variant consists of a pro-phagocytic module, a

pro-inflammatory module, and a pro-infiltrating module. The ability of CAR-M variants to bind to

PSMA antigen was confirmed by flow cytometry (Figure S10A). The ICD sizes ranged from

~1,000–2,000 bps, and little correlation was observed between the ICD size and surface expression

efficiency (Figure S10B and S10C). Interestingly, we observed significantly different CAR surface

expression levels between constructs with the same ICD elements but different arrangements (COMB1:

HCST-LY9-TNFRSF11A, and COMB4: TNFRSF11A-HCST-LY9) or constructs containing one

different ICD (COMB2: LY9-FCER1G-CD247, and COMB3: LY9-FCER1G-KIR2DS1) (Figure S10D

and S10E), suggesting that the ICD type and position modulate CAR surface expression within

macrophages.

We next evaluated the effect of the combinatorial signaling of CARs on macrophage activation.

We incubated CAR-Ms with mTagBFP2-expressing PC3-PSMA cells at an effector/target (E/T) ratio of

1/1 and observed that COMB3, COMB6, COMB7 and COMB8 CARs led to remarkable phagocytosis

of tumor cells compared to UTD cells over a time course of 4 hours (Figure 4B). Among these

CAR-Ms, COMB6 and COMB7 CAR-Ms showed the best performance in promoting phagocytosis,

suggesting their anti-tumor activity *in vitro*. We noted that all eight CAR variants presented high CD80

and CD86 expression (Figure 4C). Consistent with these findings, the reactive oxygen species (ROS)

levels (Figure 4D) and the *IL6* and *TNF* expression levels were increased in these CAR-Ms (Figure 4E),

suggesting their pro-inflammatory ability. In addition, we noted that all the CAR variants except

COMB5 promoted MMP family gene expression in macrophages (Figure 4F). In summary, the NaSDC

and CoSDC libraries facilitated the rapid identification of CAR ICD combinations that exhibit

synergetic anti-tumor effects in CAR-M immunotherapy. We chose the COMB7 CAR construct for

further development due to its robust pro-phagocytic function and comparable pro-inflammatory and

pro-infiltrating capabilities.

A novel CD40-LY9-FCRL1 CAR-M modifies the TME

Given that the COMB7 construct promoted pro-inflammatory phenotypes of macrophages *in vitro,* we

evaluated the interactions of COMB7 CAR-Ms within the TME in a humanized immune system mouse

model (Figure 5A). Given the variability in the immune system reconstitution rates among individuals,

we adopted a bilateral tumor model to ensure comparability between groups. Briefly, PSMA-positive

prostate tumors were bilaterally inoculated into the subcutaneous flanks of NCG mice. After

confirmation of tumor engraftment, human peripheral blood mononuclear cells (PBMCs) were

transplanted to reconstitute the immune system. UTD or COMB7 CAR-Ms were injected

intratumorally (IT) into either the right or left flank. Four days later, the tumors were harvested, and the

CAR expression levels were measured in matched tumor tissues (Figure S11A). The human

CD45-positive cells were sorted and subjected to scRNA-seq (Figure S11B). The transcriptomes of

17,432 individual cells were obtained from scRNA-seq following data filtering, of which 44.6% were

macrophages and 40.4% were identified as T cells (Figure 5B and S11C).

Using the scRNA-seq data, we analyzed the phenotypic and functional changes of macrophages in

the TME caused by COMB7 CAR-M treatment. We isolated CAR-positive macrophages (Figure 5C

and 5D) from COMB7 CAR-M-treated tumors and observed a strong M1-like polarization phenotype

in these cells (Figure 5E). Notably, CAR-negative macrophages from the COMB7 flank group

presented higher levels of M1 features than those from the UTD group (Figure 5E and 5F), suggesting

that COMB7 CAR-Ms can maintain a pro-inflammatory microenvironment within the human TME.

Although macrophages from UTD-treated tumors expressed several M1-related genes, such as *IL1B*

and *IL6*, these cells simultaneously expressed the M2 markers CD163 and MRC1 (CD206) (Figure 5E

and 5F). Consistent with these findings, UTD cells upregulated the expression of anti-inflammatory

cytokines such as IL-10, which promote an immunosuppressive TME (Figure S11D). In addition,

macrophages from the COMB7 flanks exhibited higher levels of MMP9, suggesting increased ECM

degradation capacity (Figure S11D). Together, these data demonstrate that COMB7 CAR-Ms have the

ability to reduce tumor-induced M2 polarization and maintain an inflammatory TME in solid tumors.

The immunosuppressive environment of solid tumors can drive exhaustion and dysfunction of T

cells [34, 35]. Given the improved TME caused by COMB7 CAR-M treatment and the upregulation of

costimulatory ligands as well as antigen processing and presentation genes in macrophages (Figure 5F

and 5G), we examined whether COMB7 CAR-M therapy could change T cell behavior in solid tumors.

We observed no significant correlations between cytotoxicity scores and COMB7 CAR-M treatment,

but the exhaustion scores were notably decreased in the $CD8^+$ T cells of the COMB7 group (Figure 5H).

Furthermore, we tested the antigen cross-presentation ability of CAR-Ms. We cocultured the CAR-Ms

with NY-ESO-1-expressing PC3-PSMA cells for 24 h and then added the Jurkat T cells expressing

NY-ESO-1-specific T cell receptor (TCR) to the cocultures for 24 h. The Jurkat T cells were activated

by CAR-Ms that ingested NY-ESO-1-expressing PC3-PSMA (Figure 5I). Together, these data suggest

that COMB7 CAR-M treatment maintains T cell function in solid tumors.

CD40-LY9-FCRL1 CAR improves solid tumor clearance

We next evaluated the *in vitro* killing capacity of the COMB7 CAR-induced macrophages. The

COMB7 CARs were transduced to hMDMs using the LNP method. No correlation was observed

between the intracellular levels of CAR-encoded mRNA and the killing efficacy (Figure S12A). We

cocultured PSMA$^+$ cancer cells with COMB7 CAR-Ms and observed that the COMB7 CAR improved

the killing capacity across multiple E/T ratios (Figure 6A). Notably, UTD macrophages had a strong

tumor killing effect at an E/T ratio of 10/1 (Figure 6A), suggesting the intrinsic anti-tumor function of

primary macrophages. Compared with the CAR-3ζ, the COMB7 CAR showed increased cytotoxicity in

macrophages from two human donors (Figure 6B). We next evaluated the killing capacity of the

COMB7 CARs in different tumor models. We generated COMB7 CAR-Ms targeting the solid tumor

antigens HER2 or GPC3 and cocultured these CAR-Ms with antigen-positive tumor cells. We observed

significant killing of HER2-positive SKOV3 cells in COMB7 CAR-M cocultures (Figure 6C). Both

UTD- and COMB7-engineered macrophages exhibited high killing capacity in GPC3-positive HepG2

cells (Figure 6D).

We next evaluated the *in vivo* killing capacity of COMB7 CAR-Ms in NCG mice that were

challenged with intraperitoneal (IP) or subcutaneous prostate tumors (Figure 6E). In the IP model, mice

received a single IP administration of PBS, UTD macrophages, or anti-PSMA CAR-Ms one day

post-tumor challenge. Tumor progression was monitored longitudinally using living imaging. Notably,

mice treated with COMB7 CAR-Ms exhibited significantly enhanced tumor control and improved

survival compared to control groups (Figure 6F, 6G and S12B). In the SC model, CAR-Ms were

administered via in situ injection following confirmation of tumor engraftment (approximately 7 days

post-implantation). Treatment with COMB7 CAR-Ms resulted in sustained suppression of tumor

growth over a 42-day observation period (Figure 6H and 6I). Collectively, these findings demonstrate

that COMB7-based chimeric antigen receptors effectively direct macrophage-mediated antitumor

activity in vivo, supporting the therapeutic potential of this CAR-M platform and providing a

foundation for its clinical translation.

Discussion

The signaling domains of CAR constructs strongly modulate the phenotypic output of CAR-Ms. To discover ICDs that can enhance CAR-M anti-tumor functions, we designed a two-step NaSDC×CoSDC screening platform. The NaSDC library contains CARs with single ICDs derived from immune response-related receptors. NaSDC screening allows us to filter out ICDs with weak anti-tumor effects. CAR variants with different anti-tumor potentials were selected from NaSDC and randomly combined to generate CoSDC. We integrated CoSDC library with scRNA-seq and scCAR-seq to identify synergistic ICD combinations that enhance macrophage anti-tumor functions. As a result, NaSDC screening identified 17 functional CARs with previously unused intracellular signaling domains. Subsequent CoSDC screening using the random combinations of these 17 ICDs identified a novel CD40-LY9-FCRL1 CAR construct with enhanced anti-tumor functions. Together, the CAR candidates identified by the screening platform exhibit functions that are uncommon in current CAR-M designs and may lead to new applications for effective CAR-M therapies.

Unbiased combinatorial CAR signaling domain screens now serve as powerful tools for identifying candidates for improving T cell differentiation, activation, and cytotoxicity in CAR-T therapy [18-20, 22]. However, this screening strategy has not been applied to optimize CAR macrophages. Previous studies have focused primarily on designing CAR-Ms on the basis of known activation and polarization mechanisms of macrophages. Given the conservation of receptor signaling pathways across the immune system [36-38], we hypothesized that certain receptors, which are not naturally expressed in macrophages, could nevertheless activate these cells when incorporated into a CAR construct. Indeed, previous studies have reported T cell-related receptor CD3ζ as a phagocytic

domain of CAR-Ms [10, 15]. Our NaSDC screens demonstrate that lymphatic receptors such as CD3E

and ICOS can function as intracellular signaling domains to promote inflammatory responses or

enhance phagocytosis in macrophages. Together, these results extend the range of signaling domains

from other cell types that can act upon macrophages. Furthermore, macrophages exhibit a wide range

of activation states upon sensing and responding to combined cues in their microenvironment [39-41],

which cannot be accurately predicted at present. Using the CoSDC platform in combination with

scRNA-seq and scCAR-seq, we found that the combinatorial signaling induced at least seven distinct

macrophage activation states, although many of the CAR signaling domains have similar activation

mechanisms or functions. These findings underscore the unpredictable nature of macrophage activation

in response to combinatorial signaling, highlighting the critical importance of pooled screening

strategies in advancing macrophage-based engineered cell therapies.

Gene transduction remains a major challenge in macrophage research. Lentiviral transduction is

the primary method for large-scale library delivery. However, macrophages express retroviral

restriction factors such as SAMHD1 [42-44], which limit the efficiency of lentiviral-based gene

transduction. Previous pooled screening using human monocytic cell line (U937 and THP-1) to

overcome these limitations [25, 26, 45]. In our study, we used PMA-differentiated THP-1 cells as an

alternative model to conduct NaSDC and CoSDC screening. While this model does not fully replicate

the characteristics of activated macrophages [46, 47], we observed similar phenotypic outputs between

THP-1-derived macrophages and hMDMs in response to CAR signaling in NaSDC library screens,

indicating its reliability for CAR identification. To enable direct screening in human primary

macrophages, future research should develop more efficient gene delivery methods, such as improved

viral vectors, or non-viral systems, to overcome retroviral restrictions and support CAR-M

development.

Although we focused on anti-tumor functionality, NaSDC×CoSDC has potential for optimizing

CAR-M therapy in various diseases, such as autoimmune diseases, cardiac injuries, and liver fibrosis.

These conditions involve distinct macrophage phenotypes with diverse functions, such as

anti-inflammatory responses, tissue repair, and fibrosis attenuation. Moving forward, the

NaSDC×CoSDC strategy should not be limited to CAR domain screening but should be expanded to

include the selection of TF combinations, cytokine combinations, and even random combinations of

CARs, TFs and cytokines to further increase macrophage activation and polarization. Moreover, the

integration of CoSDC with scRNA-seq can be used not only to find CAR signaling domains but also as

an optimal approach for investigating the intricate mechanisms of macrophage polarization, making it

possible to obtain snapshots of individual macrophages upon stimulation with different combinations

of signals. Together, the NaSDC×CoSDC strategy will accelerate the process of candidate selection and

design optimization for combinatorial constructs, advancing the discovery of biological mechanisms

and supporting the development of a broad range of cellular therapies.

Methods

Cell lines

HEK293T (referred to as 293T), THP-1, PC-3, K562, Jurkat cell lines were purchased from the cell

bank of the Committee on Type Culture Collection of the Chinese Academy of Sciences. 293T cells

were cultured in DMEM. THP-1, PC-3 cells were maintained in RPMI 1640 medium. To obtain

macrophages, THP-1 cells were differentiated with 1 ng/ml phorbol 12-myristate 13-acetate (PMA) for

48 h. PC-3 cells were transduced with a construct of FFluc-P2A-mTagBFP2 (FFLuc-BFP) to stably

express firefly luciferase (FFLuc) along with a mTagBFP2 fluorescence. PC-3-FFLuc-BFP (referred to

as PC3) cells were then transduced with lentiviral vectors encoding PSMA (*FOLH1*) to generate

PC3-PSMA cells. K562 cells were transduced with lentiviral vectors encoding PSMA to generate

K562-PSMA cells. Cells were maintained in culture medium supplemented with 10% fetal bovine

serum (FBS, GIBCO) and 1% penicillin/streptomycin at 37 °C with 5% CO_2 in a humidified incubator.

Primary cells

De-identified healthy male and female donor blood samples were provided by the SAILYBIO Co., Ltd.

Sample acquisitions were approved and regulated by the Shanghai Liquan Hospital Institutional

Review Board (XF-WBC-220809). PBMCs were isolated from blood samples by density gradient

centrifugation with Lymphocyte separation medium (Yeasen). Monocytes were isolated from PBMCs

from all cell sources by positive selection with anti-CD14 magnetic beads (Miltenyi Biotec). For

macrophage differentiation, isolated CD14[+] monocytes were seeded on plates in X-VIVO 15 medium

(Lonza) supplemented with 10% FBS and 50 ng/ml GM-CSF (Yeasen) and cultured for a further 7 days

at 37 °C with 5% CO_2.

Xenogeneic mouse models

All mouse procedures and experiments for this study were approved by the Institutional Animal Care

and Use Committee at Nanjing University (IACUC-2306005). Animals were housed in a

temperature-controlled specific pathogen-free facilities with a 12 h light/dark cycle with 45-65%

relative humidity. Food and water were available ad libitum. All mice were more than 8 weeks old at

the beginning of the experiments. Animal sample sizes were determined based on previous studies and

literatures of the field using similar experimental paradigms.

Male NOD/ShiLtJGpt-*Prkdc^{em26Cd52}Il2rg^{em26Cd22}*/Gpt (NCG) mice were purchased from

GemPharmatech. For tumor control and survival analyses, mice were injected IP with 1×10^6

PC3-PSMA cells. 24 hours later, 5×10^6 CAR-Ms, UTD macrophages or PBS were injected IP. Tumor

burden was monitored once per week by IP injection of D-Luciferin, Potassium Salt (Yeasen) at a dose

of 150 mg/kg, then measurement of bioluminescence with an IVIS Spectrum imaging system

(PerkinElmer). Tumor burden was quantified by total photon counts using the LivingImage software.

For solid tumor control analyses, mice were injected SC with 1×10^6 PC3-PSMA cells. After

confirmation of tumor engraftment (about 7 days), 5×10^6 CAR-Ms, UTD macrophages or PBS were

injected IT. Tumor volume was monitored once per week. All mice were weighed weekly and were

subject to routine veterinary assessment for signs of overt illness. Animals were killed at experimental

termination or when predetermined IACUC rodent health endpoints were reached. For TME analyses,

4×10^6 PC3-PSMA tumor cells were resuspended in Ceturegel Matrix LDEV-Free Matrigel (Yeasen)

and injected SC into both flanks of NCG mice. After confirmation of tumor engraftment (about seven

days), mice were engrafted intravenously (IV) with 2×10^7 human PBMCs. On day 14, 5×10^6 UTD or

COMB7 CAR-Ms were injected IT into either the right or left flank. Four days later, the tumors were

harvested for further analysis.

NaSDC library construction

The lentiviral vector pHIV-EGFP was gifted by Bryan Welm and Zena Werb (Addgene plasmid

#21373). To generate a backbone vector for ICD insertion, a codon-optimized gene encoding PSMA

CAR composed of CD8α signal peptide, J591 single-chain variable fragments (scFv), CD8α hinge and

transmembrane domain was commercially synthesized by GenScript and cloned into the pHIV-EGFP

vector. An Esp3I cleavable linker and a randomized 18-nucleotide sequence were insert to C-terminal

portion of the CAR construct. The backbone vector was digested with FastDigest Esp3I (Thermo

Fisher), and 131 immune response-related ICDs (Table S2) were individually cloned into the vector

using Gibson Assembly. Then, the bacterial transformation was conducted using the obtained products,

and colonies were randomly selected. Sequences were performed to identify the correct encoding

plasmid with a unique barcode for each CAR variant. The single CAR variant plasmids were pooled at

equimolar ratios, electroporated into Endura Electrocompetent cells (LGC), and purified to generate the

NaSDC plasmid library.

Lentiviral library production

The plasmids pMD2.G and psPAX2 were gifts from Didier Trono (Addgene plasmid #12259 and

#12260). To produce lentivirus, 293T cells were transfected with a plasmid library and the packaging

plasmids psPAX2 and pMD2.G using Lipofectamine 2000 Transfection Reagent (Thermo Fisher) at a

mass ratio of 24:3:1. Medium was changed 12 h after transfection, and the supernatant containing

lentiviruses were collected in 48-72 h after transfection. The supernatant was filtered through 0.45 μm

non-pyrogenic filters. To concentrate lentiviruses, the Lenti-Concentin Virus Precipitation Solution

(ExCell Bio) was added to the supernatant for 12, and centrifuged for 30 min at $2,000 \times g$. The pellet

was then resuspended in OptiMEM medium (Thermo Fisher) and stored at -80 °C. Medium containing

concentrated lentiviral library was used to infect THP-1 cells in the presence of 8 μg/ml polybrene

(Yeasen).

Pro-phagocytic CAR variant screens

Streptavidin-coated magnetic beads (ACROBiosystems) were sterilized for 30 min in 70% isopropanol. Beads were spun down and resuspended in OptiMEM. Biotinylated human PSMA proteins (ACROBiosystems) were added to the beads at a concentration sufficient to occupy 25% of the binding sites. After incubation for 1 h, the beads were washed using PBS and resuspended in OptiMEM. The PSMA-labeled beads were added to macrophages carrying NaSDC library at a bead-to-cell ratio of 5:1. After phagocytosis for 2 h, cells were rinsed eight times by PBS to remove un-ingested particles. Then, the cells were trypsinized, harvested and sorted using a magnetic holder. The captured cells were washed three times by PBS. The input NaSDC CAR-M library and the magnetic sorted cells were store at -80 °C in TRIzol reagent (Thermo Fisher) for subsequent barcode sequencing.

Pro-inflammatory CAR variant screens

To enable screening with fluorescent NFκB reporter system, a construct composed of NFκB response elements (NFRE), a minimal TATA-box promoter, and an iRFP670 fluorescent reporter was cloned into a pLenti6/V5 lentiviral vector to replace the CMV promotor. The fluorescent reporter was then stably co-transduced into THP-1 cells with NaSDC library, and the positive cells were selected using blasticidin (10 μg/ml). The reporter NaSDC THP-1 cells were induced by 1 ng/ml PMA for 48 h. After differentiation, PMA was washed out with PBS, and PC3 conditional media were added to the cells for 48 h to induce TAM polarization. The polarized CAR-M pool was cocultured with PC3-PSMA cells at a ratio of 1:1 for 24 h. Following coculture, cells were trypsinized, and resuspended in FACS buffer (PBS + 1-2% FBS), and iRFP670^{high} cells were sorted by a FACSAria II Cell Sorter (BD Biosciences). The input reporter NaSDC CAR-M library and the sorted cells were store at -80 °C in TRIzol reagent

(Thermo Fisher) for subsequent barcode sequencing.

Pro-infiltrating CAR variant screens

THP-1 cells were lentivirally transduced with a NaSDC library, and the positive cells were selected

using flow cytometry. NCG mice were dosed with 2×10^6 PC3-PSMA tumor cells through subcutaneous

injection. The NaSDC THP-1 cells were induced by 1 ng/ml PMA for 48 h, and 1×10^7 NaSDC

CAR-Ms were peritumorally injected 7 days after tumor inoculation. Four days later, the tumors were

harvested. The input NaSDC CAR-M library and the tumor tissues containing infiltrated CAR-Ms were

store at -80 °C in TRIzol reagent (Thermo Fisher) for subsequent barcode sequencing.

Barcode sequencing and data analysis

RNA was extracted from input and output population using TRIzol reagent, and reversely transcribed

into cDNA using HiScript IV 1st Strand cDNA Synthesis Kit (Vazyme). Subsequently, barcode

sequences were PCR amplified, and Illumina sequencing adaptors were appended using NEBNext

Ultra II Q5 Master Mix (NEB). SYBR Green I (MeilunBio) was incorporated to the rection system to

monitor the progression of PCR. The cDNA was amplified in a C1000 thermocycler (Bio-Rad) until the

fluorescent signal approached the termination of exponential phase. Then, the products were

immediately transferred to another thermocycler, and incubated at 72 °C for 5 min to complete the PCR

program. The PCR products were bead-purified and sequenced by Novogene on an Illumina NovaSeq

X Plus Sequencer. Considering the limited *in vitro* proliferation ability of macrophages, a single round

screening would result in an elevated false positive rate within the output populations. Therefore, we

repeated the screening experiments and sequencing runs to minimize the experimental bias.

DESeq2 (v1.42.1) [48] and RobustRankAggreg (v1.2.1) [49] R packages were integrated to

compute the alterations of barcodes in output vs input populations. DESeq2 was used to merge the

repeated experiments and calculate the overall log2 fold change (log2FC) of barcode representation. In

the RobustRankAggreg analysis, the log2FC was calculated separately for each experimental batch.

The barcodes were then ranked based on their log2FC and aggregated using RRA algorithm. RRA

method uses rho score to retrieve the positive factors from the noise, and represent the significance

probabilities for all the elements. The rho score < 0.05 and overall log2FC > log2FC (CARΔ) was

considered as positive enrichment in output population.

In vitro mRNA transcription

The genes encoding indicated CAR variants or EGFP control were codon-optimized and cloned into a

plasmid carrying a T7 promoter, 5' and 3' UTR elements, and a poly(A) tail. The plasmids were

linearized by FastDigest LguI (Thermo Fisher), and the mRNA was synthesized using T7

Co-transcription RNA Synthesis Kit (Syngene). N1-methyl-pseudouridine-5′-triphosphate

(N1-Me-pUTP) was substituted for uridine triphosphate. Capping of the IVT mRNAs was performed

co-transcriptionally using a trinucleotide cap1 analog CAP GAG (Syngene). The generated mRNAs

were purified using VAHTS RNA Clean Beads (Vazyme). All mRNAs were analyzed by agarose gel

electrophoresis and stored at -80 °C before uses. 10 pmol mRNAs were used to transduce 1×10^6

hMDMs via Lipofectamine MessengerMAX Transfection Reagent (Thermo Fisher) according to the

manufacturer's protocol. The medium was changed 12 h after transduction, and flow cytometry was

performed to detect surface expression of CAR proteins.

Flow cytometry

To block non-specific bindings, FcR Blocking Reagent (Miltenyi Biotech) was always used for staining of monocytes, macrophages or monocytic cell lines expressing Fc receptors. Macrophage purity was tested using the following antibodies: Anti-human CD11b PE (1:100, BioLegend), Anti-human CD14 APC (1:100, BioLegend), Anti-human CD68 APC (1:100, BioLegend). M1 markers on human macrophages were detected with the following antibodies: Anti-human CD80 FITC (1:100, BioLegend), Anti-human CD86 APC (1:100, Thermo Fisher). Surface anti-PSMA CAR expression was detected using PE-Labeled Human PSMA (ACROBiosystems), or Biotinylated Human PSMA (ACROBiosystems) followed by Streptavidin/APC (Solarbio) staining. Flow cytometric analysis was performed on a NovoCyte Flow Cytometer (ACEA Biosciences) or an Attune NxT Flow Cytometer (Thermo Fisher). The data were analyzed by NovoExpress (v1.6.2) and visualized by CNSplot (v1.0.1) R package.

Phagocytosis assay

The PC3-PSMA cells were transduced to stably express mTagBFP2 fluorescence as described above. The mTagBFP2 is a blue fluorescent protein with lower sensitivity to the acidic environment of the lysosome compared to other fluorescence proteins such as EGFP [50]. For each CAR variant, 5×10^5 UTD or CAR-Ms were cocultured with 5×10^5 PC3-PSMA cells for 4-6 h at 37 °C. After coculture, cells were harvested with Accutase (Yeasen) and resuspended in FACS buffer. The CAR-Ms were stained with PE-Labeled Human PSMA (ACROBiosystems), and the UTD cells were stained with Anti-human CD11b PE (BioLegend). The cells were analyzed using an Attune NxT Flow Cytometer (Thermo Fisher). The percent of mTagBFP2$^+$ events within the PE$^+$ population was plotted as

percentage phagocytosis.

Quantitative real-time PCR (qPCR)

Total RNA was isolated from cells or tumor tissue samples with TRIzol reagent. Reverse transcription

was performed with HiScript III RT SuperMix for qPCR (Vazyme). The qPCR assays were performed

on a QuantStudio□3 Real-Time PCR Systems (Thermo Fisher) using AceQ qPCR SYBR Green Master

Mix (Vazyme) according to the manufacturer's protocol. The data were analyzed by QuantStudio

(v1.5.2) and visualized by CNSplot (v1.0.1) R package. The expression of genes was normalized to

housekeeping gene *GAPDH*, and data were normalized to the mean of the control group. Primer

sequences are described in Table S1.

Cytokine assays

For each CAR variant, 1×10^5 UTD or CAR-Ms were cocultured with antigen-positive tumor cells in an

E/T ratio of 1/1 in complete X-VIVO15 medium and incubated at 37 °C for approximately 24 h. After

stimulation, the supernatants were collected and IL-1β, IL-6 or TNFα were measured by ELISA

following the manufacturer's protocol (Thermo Fisher).

RNA sequencing and data analysis

For bulk RNA-sequencing, K562 and K562-PSMA cells were labeled with green fluorescence using

carboxyfluorescein succinimidyl ester (CFSE, Yeasen) method. The hMDMs from independent healthy

donors were transduced with the mRNA of indicated CAR variants, and cocultured with either K562 or

K562-PSMA cells at a E/T ratio of 1/1 for 24 h. Subsequently, the tumor cells were removed from the

cocultures. The absolute removal of tumor cells was confirmed by flow cytometry. Total RNA was

extracted from 5×10^6 CAR-Ms using TRIzol reagent according to the manufacturer's protocol. Next

generation sequencing library preparations were conducted according to the manufacturer's protocol.

The libraries were sequenced by Novogene on an Illumina NovaSeq X Plus Sequencer. Read quality

was assessed with FastQC (v0.12.1) and adapters were trimmed using Cutadapt (v4.4). Reads were

then aligned to the GRCh38 genome using HISAT2 (v2.2.1) [51] and reads in exons were counted

using featureCounts function in Subread (v2.0.3) package [52]. Analysis of differentially expressed

genes was conducted using DESeq2 (v1.40.1) R package. Go analysis was performed using

clusterProfiler (v4.10.1) R package [53]. Count data was transformed to log2 scale and normalized with

respect to library size using rlog transformation produces. Absolute log2 fold change (log2FC) > 1 and

p value < 0.05 was considered statistically significant.

CoSDC library construction using CARode method

The lentiviral vector pLenti6/V5 was used as the backbone vector for CoSDC library construction,

because it uses a shorter 3' LTR region, which is suitable for 3' capture kits of scRNA-seq. A

codon-optimized gene encoding PSMA CAR composed of CD8α signal peptide, J591 single-chain

variable fragments (scFv), CD8α hinge and transmembrane domain was commercially synthesized by

GenScript and cloned into the pLenti6/V5 vector. A linker containing XbaI and XhoI restriction sites

was insert to C-terminal portion of the CAR construct. ICDs were PCR amplified to combine with first

round linkers, a cloning cassette (containing BamHI and XhoI restriction sites) and a unique 6-bp

barcode (a linker-cassette-barcode-linker construct). The PSMA CAR vector was digested with

FastDigest XbaI (Thermo Fisher) and FastDigest XhoI (Thermo Fisher). The ICDs were individually

cloned into the PSMA CAR vector using Gibson Assembly. Then, the bacterial transformation was

conducted using the obtained products. The bacteria containing different CAR variant vectors were

pooled at equal OD ratios, and lysed to extract G1 library plasmid. The single ICD fragments were

PCR amplified again to combine with second round "linker-cassette-barcode-linker". The G1 library

was digested with FastDigest BamHI (Thermo Fisher) and FastDigest XhoI (Thermo Fisher). The 2nd

ICDs were individually cloned into the G1 to generate G2 library. Similarly, ICDs were PCR amplified

to combine with third round "linker-cassette-barcode-linker". The G2 library was digested with BamHI

and XhoI enzyme. The 3rd ICDs were individually cloned into the G2 to generate G3 library. The G1,

G2 and G3 library and a truncated CAR were pooled together, and digested with FastDigest BcuI

(Thermo Fisher). The internal ribosome entry site (IRES) elements and EGFP fluorescence were

inserted into the cloning cassette. The products were electroporated into Endura Electrocompetent cells

(LGC), and purified to generate the CoSDC plasmid library.

Single-cell RNA sequencing and single-cell CAR sequencing

The CoSDC lentiviral library was generated as described above, and transduced to THP-1 cells at a

MOI of 0.3 to minimize multiple integrations per cell. The EGFP$^+$ cells were sorted, PMA-induced and

cocultured with high PSMA-expressing tumor cell PC3-PSMA at an E/T ratio of 1/1 for 24 h.

Immediately after coculture, EGFP$^+$ CAR-Ms were sorted by FACS, and scRNA-seq library was

constructed using the 10X Genomics Chromium system following manufacturer's instructions. In brief,

the suspended cells were loaded into Chromium microfluidic chips and barcoded with a 10× Chromium

Controller (10X Genomics). RNA from the barcoded cells was subsequently reverse-transcribed. The

barcoded cDNA was purified and amplified, followed by fragmenting, A-tailing and ligation with

adaptors. The 10X scRNA-seq libraries were sequenced by Novogene on an Illumina NovaSeq X Plus

Sequencer. For scCAR-seq, CARode sequences were PCR amplified from 100 ng unfragmented 10X

cDNA, and Illumina sequencing adaptors were appended using NEBNext Ultra II Q5 Master Mix

(NEB). The CARode libraries were sequenced by Novogene on an Illumina NovaSeq X Plus

Sequencer.

For scRNA-seq analysis, raw 10X sequencing reads were aligned to the GRCh38 genome using

Cell Ranger (v7.2.0). Gene expression profiles were processed using Seurat (v5.1.0) R package [54].

Briefly, quality control was applied to cells based on the total UMI counts and number of detected

genes. Cells with less than 5,000 UMI count were filtered, as well as cells with more than 10%

mitochondrial gene count. To remove potential doublets, cells with UMI count above 70,000 and

detected genes above 9,000 are filtered out. The gene counts were normalized using SCTransform. For

clustering, top 2,000 variable genes were used, and ribosomal and mitochondrial genes were removed

from the variable gene set before running PCA and identifying clusters using a shared nearest neighbor

(SNN) modularity optimization-based clustering algorithm. The Seurat normalized count matrix was

used for the downstream analysis.

For scCAR-seq analysis, the CARode sequences for each cell were extracted from the raw

sequencing reads and mapped to signal domains. A total of over 21 million CARode reads were

obtained from 16,612 single cells. Cells with more than 10 CARode reads were classified as CAR-Ms.

7911 cells expressing a single specific CAR variant were identified from the scRNA-seq data for

subsequent clustering and functional analyses.

Gene set activity analysis

AUCell (v1.24.0) R package [55] was employed to assess the gene set activities in scRNA-seq

data. AUCell uses the "Area Under the Curve" (AUC) score to quantify the expression level of an input

gene set for each cell. Previously established M1 and M2 signatures [56] were used for macrophage

polarization analysis. MacSpectrum (v0.1.0) R package [57] was also employed to assess macrophage

activation. To evaluate antigen processing and presentation, gene sets from the Reactome (2132295,

983169), KEGG (04612), and GO (0019882) databases were used. Additionally, KEGG (04666) and

GO (0006909) gene sets were used to estimate phagocytic activity. MMP family genes MMP1-28 were

applied to analyze MMP expression levels.

Antigen cross-presentation assay

PC3-PSMA cells were transduced with lentivirus encoding NY-ESO-1 or

HLA-A0201-P2A-NY-ESO-1 to generate PC3-PSMA-NY and PC3-PSMA-HLA-NY cells.

NY-ESO-1-specific T cells were generated by transducing Jurkat cells with lentivirus encoding the 1G4

anti-NYESO-1 specific TCR. CAR-Ms were generated from HLA-A0201$^+$ healthy donors. CAR-Ms

were cocultured with PC3-PSMA-NY cells at an E/T ratio of 1/1. Subsequently, Jurkat cells were

added to the cocultures at an Jurkat/CAR-M ratio of 1/1. Twenty-four hours later, Jurkat cells were

analyzed for activation markers (CD69) by FACS.

Cytotoxicity assay

To measure specific killing, a bioluminescence (BLI)-based cytotoxicity assay was performed.

FFluc-positive PC3-PSMA, SKOV3, and HepG2 tumor cells were used as targets. CAR-Ms were

cocultured with 10,000 target cells at the specified E/T ratios in complete X-VIVO15 medium on black

96-well flat-bottom plates. The BLI was measured using a luminometer (Tecan Spark) in relative light

units (RLU). A spontaneous death control was used including tumor cells alone. The maximal death

control contained 1% Triton X-100 and tumor cells. Specific lysis was calculated using: %Specific

Lysis = 100 × (average spontaneous death RLU − test RLU)/(average spontaneous death RLU −

average maximal death RLU). Negative specific lysis values indicate more signal than in the

spontaneous death control wells.

Statistical analysis

Statistical analysis was performed using rstatix (v0.7.2) R package. Statistical details for all

experiments can be found in the figure legends. Error bars indicated the Standard Error of Mean

(s.e.m.). Differences were analyzed by unpaired Student's t test or one-way ANOVA depending on

experimental conditions. Unless otherwise indicated, the experiments were performed independently in

triplicate, and n is indicated in the figure legends.

Acknowledgements

Thanks to members of Shen's laboratory for helpful discussions, technical assistance, and critical

reading of the manuscript. We thank Fuchou Tang and Lu Wen from Peking University, and Jiaming Su

from Zhejiang University for their help in library construction, high-throughput sequencing and

lentiviral transduction. This work was supported by grants from the Natural Science Foundation of

Jiangsu Province (BK20222009 to P. Shen), the Pivotal Life Sciences Sponsorship Program (1352806

to P. Shen), the Program from National Center of Technology Innovation for Biopharmaceuticals

(NCTIB2023XB01013 to P. Shen), the Program from Proof of Concept Center of Jiangsu Province

(Research, Development and Clinical Study of Chimeric Antigen Receptor Macrophage

Immunotherapy Technology Targeting Malignant Prostate Cancer), and the Guangdong Basic and

Applied Basic Research Foundation (2021B1515120016 to P. Shen).

Author contributions

Y. Wang and S. Zuo conceived the project and performed experiments and data analysis. H. Bao, Z.

Zhang, and Q. Liu performed experiments. Y. Chen, W. Zhang, Y. Lu, Y. Huang, W. Zheng, N. Yang,

and L. Ye provided scientific suggestions and contributed to the manuscript revision. P. Shen

supervised the project and wrote the manuscript. The author(s) read and approved the final manuscript.

Data availability

RNA-seq, scRNA-seq, scCAR-seq data have been deposited in the Gene Expression Omnibus

repository under accession code GSE289202.

References

1.   June, C.H., et al., *CAR T cell immunotherapy for human cancer.* Science, 2018. **359**(6382): p.

1361-1365.

2.   Cappell, K.M. and J.N. Kochenderfer, *Long-term outcomes following CAR T cell therapy:*

what we know so far. Nat Rev Clin Oncol, 2023. **20**(6): p. 359-371.

3.   Maalej, K.M., et al., *CAR-cell therapy in the era of solid tumor treatment: current challenges*

and emerging therapeutic advances. Mol Cancer, 2023. **22**(1): p. 20.

4.   Newick, K., et al., *CAR T Cell Therapy for Solid Tumors.* Annu Rev Med, 2017. **68**: p.

139-152.

5.   Mantovani, A., et al., *Macrophages as tools and targets in cancer therapy.* Nat Rev Drug

Discov, 2022. **21**(11): p. 799-820.

6.   Reiss, K.A., et al., *CAR-macrophage therapy for HER2-overexpressing advanced solid tumors:*

a phase 1 trial. Nat Med, 2025.

7.   Christofides, A., et al., *The complex role of tumor-infiltrating macrophages.* Nat Immunol,

2022. **23**(8): p. 1148-1156.

8.   Cassetta, L. and J.W. Pollard, *A timeline of tumour-associated macrophage biology.* Nat Rev

Cancer, 2023. **23**(4): p. 238-257.

9.   Zhang, W., et al., *Chimeric antigen receptor macrophage therapy for breast tumours mediated*

by targeting the tumour extracellular matrix. Br J Cancer, 2019. **121**(10): p. 837-845.

10.   Klichinsky, M., et al., *Human chimeric antigen receptor macrophages for cancer*

immunotherapy. Nat Biotechnol, 2020. **38**(8): p. 947-953.

11.   Lei, A., et al., *A second-generation M1-polarized CAR macrophage with antitumor efficacy.*

Nat Immunol, 2024. **25**(1): p. 102-116.

12.   Sloas, C., S. Gill, and M. Klichinsky, *Engineered CAR-Macrophages as Adoptive*

Immunotherapies for Solid Tumors. Front Immunol, 2021. **12**: p. 783305.

13.   Dong, X., et al., *Efficacy evaluation of chimeric antigen receptor-modified human peritoneal*

macrophages in the treatment of gastric cancer. Br J Cancer, 2023. **129**(3): p. 551-562.

14.   Morrissey, M.A., et al., *Chimeric antigen receptors that trigger phagocytosis.* Elife, 2018. **7**.

15.   Shah, Z., et al., *Human anti-PSCA CAR macrophages possess potent antitumor activity*

against pancreatic cancer. Cell Stem Cell, 2024. **31**(6): p. 803-817 e6.

16.   Duan, Z., et al., *Chimeric antigen receptor macrophages activated through TLR4 or*

IFN-gamma receptors suppress breast cancer growth by targeting VEGFR2. Cancer Immunol

Immunother, 2023. **72**(10): p. 3243-3257.

17.   Dong, M.B., et al., *Tumor immunology CRISPR screening: present, past, and future.* Trends

Cancer, 2022. **8**(3): p. 210-225.

18.   Gordon, K.S., et al., *Screening for CD19-specific chimaeric antigen receptors with enhanced*

signalling via a barcoded library of intracellular domains. Nat Biomed Eng, 2022. **6**(7): p.

855-866.

19.   Castellanos-Rueda, R., et al., *speedingCARs: accelerating the engineering of CAR T cells by*

signaling domain shuffling and single-cell sequencing. Nat Commun, 2022. **13**(1): p. 6555.

20.   Blaeschke, F., et al., *Modular pooled discovery of synthetic knockin sequences to program*
760   *durable cell therapies.* Cell, 2023. **186**(19): p. 4216-4234 e33.

761   21.   Daniels, K.G., et al., *Decoding CAR T cell phenotype using combinatorial signaling motif*
762   *libraries and machine learning.* Science, 2022. **378**(6625): p. 1194-1200.

763   22.   Goodman, D.B., et al., *Pooled screening of CAR T cells identifies diverse immune signaling*
764   *domains for next-generation immunotherapies.* Science Translational Medicine, 2022. **14**(670):
765   p. eabm1463.

766   23.   Ginhoux, F. and S. Jung, *Monocytes and macrophages: developmental pathways and tissue*
767   *homeostasis.* Nat Rev Immunol, 2014. **14**(6): p. 392-404.

768   24.   Ferreira, I., et al., *Cell Cycle Regulation in Macrophages and Susceptibility to HIV-1.* Viruses,
769   2020. **12**(8).

770   25.   Lindner, B., et al., *A genome-wide CRISPR/Cas9 screen to identify phagocytosis modulators*
771   *in monocytic THP-1 cells.* Sci Rep, 2021. **11**(1): p. 12973.

772   26.   Luteijn, R.D., et al., *SLC19A1 transports immunoreactive cyclic dinucleotides.* Nature, 2019.
773   **573**(7774): p. 434-438.

774   27.   Sack, L.M., et al., *Profound Tissue Specificity in Proliferation Control Underlies Cancer*
775   *Drivers and Aneuploidy Patterns.* Cell, 2018. **173**(2): p. 499-514 e23.

776   28.   Garcia, J., et al., *Naturally occurring T cell mutations enhance engineered T cell therapies.*
777   Nature, 2024. **626**(7999): p. 626-634.

778   29.   Lanier, L.L., *DAP10- and DAP12-associated receptors in innate immunity.* Immunol Rev,
779   2009. **227**(1): p. 150-60.

780   30.   Washington, A.V., et al., *TREM-like transcript-1 protects against inflammation-associated*
781   *hemorrhage by facilitating platelet aggregation in mice and humans.* J Clin Invest, 2009.
782   **119**(6): p. 1489-501.

783   31.   Chen, L. and D.B. Flies, *Molecular mechanisms of T cell co-stimulation and co-inhibition.*
784   Nat Rev Immunol, 2013. **13**(4): p. 227-42.

785   32.   Kaur, G., J. Trowsdale, and L. Fugger, *Natural killer cells and their receptors in multiple*
786   *sclerosis.* Brain, 2013. **136**(Pt 9): p. 2657-76.

33.   Labanieh, L., R.G. Majzner, and C.L. Mackall, *Programming CAR-T cells to kill cancer.* Nat

Biomed Eng, 2018. **2**(6): p. 377-391.

34.   Mellman, I., et al., *The cancer-immunity cycle: Indication, genotype, and immunotype.*
Immunity, 2023. **56**(10): p. 2188-2205.

35.   Zhang, J., et al., *Turning cold tumors hot: from molecular mechanisms to clinical applications.*
Trends Immunol, 2022. **43**(7): p. 523-545.

36.   Dougan, M., G. Dranoff, and S.K. Dougan, *GM-CSF, IL-3, and IL-5 Family of Cytokines:*
Regulators of Inflammation. Immunity, 2019. **50**(4): p. 796-811.

37.   Spangler, J.B., et al., *Insights into cytokine-receptor interactions from cytokine engineering.*
Annu Rev Immunol, 2015. **33**: p. 139-67.

38.   Brandsma, A.M., et al., *Fc receptor inside-out signaling and possible impact on antibody*
therapy. Immunol Rev, 2015. **268**(1): p. 74-87.

39.   Xue, J., et al., *Transcriptome-based network analysis reveals a spectrum model of human*
macrophage activation. Immunity, 2014. **40**(2): p. 274-88.

40.   Pizzurro, G.A. and K. Miller-Jensen, *Reframing macrophage diversity with network motifs.*
Trends Immunol, 2023. **44**(12): p. 965-970.

41.   Locati, M., G. Curtale, and A. Mantovani, *Diversity, Mechanisms, and Significance of*
Macrophage Plasticity. Annu Rev Pathol, 2020. **15**: p. 123-147.

42.   Laguette, N., et al., *SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1 restriction*
factor counteracted by Vpx. Nature, 2011. **474**(7353): p. 654-7.

43.   Berger, A., et al., *SAMHD1-deficient CD14+ cells from individuals with Aicardi-Goutieres*
syndrome are highly susceptible to HIV-1 infection. PLoS Pathog, 2011. **7**(12): p. e1002425.

44.   Hrecka, K., et al., *Vpx relieves inhibition of HIV-1 infection of macrophages mediated by the*
SAMHD1 protein. Nature, 2011. **474**(7353): p. 658-61.

45.   Haney, M.S., et al., *Identification of phagocytosis regulators using magnetic genome-wide*
CRISPR screens. Nat Genet, 2018. **50**(12): p. 1716-1727.

46.   Tedesco, S., et al., *Convenience versus Biological Significance: Are PMA-Differentiated*
THP-1 Cells a Reliable Substitute for Blood-Derived Macrophages When Studying in Vitro
Polarization? Front Pharmacol, 2018. **9**: p. 71.

47.   Mohd Yasin, Z.N., et al., *Macrophage polarization in THP-1 cell line and primary monocytes:*

A systematic review. Differentiation, 2022. **128**: p. 67-82.

48.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

49.   Kolde, R., et al., *Robust rank aggregation for gene list integration and meta-analysis.* Bioinformatics, 2012. **28**(4): p. 573-80.

50.   Subach, O.M., et al., *An enhanced monomeric blue fluorescent protein with the high chemical stability of the chromophore.* PLoS One, 2011. **6**(12): p. e28674.

51.   Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype.* Nat Biotechnol, 2019. **37**(8): p. 907-915.

52.   Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

53.   Wu, T., et al., *clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.* Innovation (Camb), 2021. **2**(3): p. 100141.

54.   Hao, Y., et al., *Dictionary learning for integrative, multimodal and scalable single-cell analysis.* Nat Biotechnol, 2024. **42**(2): p. 293-304.

55.   Aibar, S., et al., *SCENIC: single-cell regulatory network inference and clustering.* Nat Methods, 2017. **14**(11): p. 1083-1086.

56.   Martinez, F.O., et al., *Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression.* J Immunol, 2006. **177**(10): p. 7303-11.

57.   Li, C., et al., *Single cell transcriptomics based-MacSpectrum reveals novel macrophage activation signatures in diseases.* JCI Insight, 2019. **5**(10).

Competing interests

The Shen lab has received research support from Macera therapeutics. P. Shen is a compensated co-founder, member of the scientific advisory board, and works as the CSO of Macera therapeutics. Y. Chen is a compensated co-founder, and works as the CEO of Macera therapeutics. W. Zhang works as the Process Development Director of Macera therapeutics. P. Shen, Y. Wang, S. Zuo, Y. Chen, and W.

Zhang are listed on patent applications related to this work. The other authors declare no competing

interests.

Figure legends

Figure 1. NaSDC×CoSDC screens identify therapeutic CAR-M candidates

(A) Schematic illustration of the NaSDC×CoSDC platform.

(B) Flow cytometric histograms showing EGFP expression in hMDMs and THP-1 cells transduced

with an EGFP reporter lentivirus. Grey peaks indicate protein expression levels in untransduced (UTD)

cells.

(C) Flow cytometric plots illustrating the NaSDC library sorting strategy (left) and CAR expression

levels (right).

(D) Schematic representation of the CoSDC library and CARode. ICD, intracellular domain.

Data in (**B**) and (**C**) are representative of three independent experiments.

Figure 2. NaSDC screens reveal known and novel CAR-M candidates

(A) Schematic representation of the pro-phagocytic module screening strategy.

(B) Correlation between log2 fold change and −log10 rho scores for pro-phagocytic CAR variant

screens. Data represent 18 biological replicates across six NaSDC lentivirus batches.

(C) FACS-based quantification of PC3-PSMA tumor cell phagocytosis by CAR-Ms (n = 4 biological

replicates).

(D) Schematic representation of the pro-inflammatory module screening strategy.

(E) Correlation between log2 fold change and −log10 rho scores for pro-inflammatory CAR variant

screens. Data represent 15 biological replicates across five NaSDC lentivirus batches.

(F) Relative mRNA expression of *TNF*, *IL6* and *IL1B* in CAR-Ms cocultured with PSMA positive

target tumor cells (n = 4 biological replicates).

(G) Schematic representation of the pro-infiltrating module screening strategy.

(H) Correlation between log2 fold change and –log10 rho scores for pro-infiltrating CAR variant

screens. Data represent 9 biological replicates across three NaSDC lentivirus batches.

(I) Relative mRNA expression of MMP family genes in CAR-Ms cocultured with PSMA positive

target tumor cells (n = 4 biological replicates).

Dashed lines in (**C**, **F**, **I**) indicate –log10(0.05) (horizontal) and log2 fold change of truncated CARs

(CARΔ) (vertical). Data in (C) are presented as mean ± s.e.m. and analyzed using one-way ANOVA

followed by Dunnett's multiple-comparison test.

Figure 3. CoSDC screens identify CAR signaling combinations that enhance macrophage

anti-tumor functions

(A) Schematic illustration of the CoSDC library and scCAR-seq screening platform.

(B) Proportion of cells expressing different numbers of CAR variants (left), and tSNE-based

dimensionality reduction and clustering (right) for single-CAR-expressing cells (right).

(C) MacSpectrum plot showing macrophage polarization states of single-cell clusters. MPI,

macrophage polarization index. AMDI, activation-induced macrophage differentiation index. Data are

represented as means of clusters. Black dashed line represents the mean MPI of CARΔ.

(D) Feature plots showing the distribution of AUC scores for antigen processing and

presentation-related gene sets. Reactome gene sets "MHC class II antigen presentation" (2132295) and

"Class I MHC mediated antigen processing & presentation" (983169), GO gene sets "Antigen

processing and presentation" (0019882), and KEGG pathway "Antigen processing and presentation"

(04612) were used.

(E) Normalized AUC scores for MMP-related genes. Dashed line indicates the mean AUC score of

CARΔ.

(F) Dot plot showing the phagocytosis scores for single-cell clusters. KEGG pathway "Fc gamma

R-mediated phagocytosis" (04666) and GO gene set "Phagocytosis" (0006909) were used.

(G) Heatmap showing log2 fold change (c0 vs the other clusters) of each ICD in third-generation

CARs.

Figure 4. Combinatorial CAR constructs enhance macrophage phagocytosis, pro-inflammatory

activation, and infiltration capacity

(A) Description of CAR variants used for individual characterization.

(B) FACS-based quantification of PC3-PSMA tumor cell phagocytosis by UTD and CAR-Ms (n = 4

biological replicates).

(C) Flow cytometric histograms showing surface expression of CD80 and CD86 in UTD and CAR-Ms

cocultured with PSMA positive tumor cells.

(D) Flow cytometric histograms of reactive oxygen species (ROS) levels in UTD and CAR-Ms

cocultured with PSMA positive tumor cells. DCF, 2'-7'dichlorofluorescein.

(E) Relative mRNA expression of *TNF* and *IL6* in UTD and CAR-Ms cocultured with PSMA positive

tumor cells (n = 4 biological replicates).

(F) Relative mRNA expression of *MMP11* and *MMP13* in UTD and CAR-Ms cocultured with PSMA

positive tumor cells (n = 4 biological replicates).

Data in (**B**) are presented as mean ± s.e.m. and analyzed using one-way ANOVA followed by Dunnett's

multiple-comparison. Data in (**C**, **D**) are representative of three independent experiments. Dashed lines

(**C**, **D**) represent the mean fluorescence intensity (MFI) for UTD macrophages.

Figure 5. CD40-LY9-FCRL1 CAR-Ms reshape the TME

(A) Schematic illustration of the humanized immune system TME model. PC3-PSMA tumors were

established bilaterally, followed by PBMC engraftment on day 7. On day 14, 5×10^6 UTD or CAR-Ms

were injected IT. Tumors were harvested on day 18 for scRNA-seq. SC, subcutaneously. IV,

intravenously. IT, intratumorally.

(B) Proportions of cell types from scRNA-seq (left) and tSNE-based clustering of all sequenced cells

(right). B, B cell. T, T cell. Mac, macrophage.

(C) Dimensionality reduction (tSNE) and clustering for macrophages. Data are colored by treatment

conditions.

(D) Density plots within tSNE maps depicting the distribution of CAR-expressing cells.

(E) AUC scores of M1 and M2 gene sets in the indicated macrophage clusters.

(F) Feature plots showing expression levels of selected genes.

(G) MFI of HLA-ABC and HLA-DR in UTD and CAR-Ms after 24 h coculture with PSMA positive

tumor cells (n = 3 biological replicates).

(H) AUC scores of T cell cytotoxicity and exhaustion gene sets in the indicated T cell clusters.

(I) CAR-M cross-presentation assay showing CD69 levels of NY-ESO-1-specific Jurkat cells (T) after

coculturing with NY-ESO-1-expressing PC3-PSMA (PC3-PSMA-NY), HLA-A0201 and NY-ESO-1

co-expressing PC3-PSMA(PC3-PSMA-HLA-NY), COMB7 CAR-Ms (COMB7), and cocultures of

COMB7 CAR-Ms and PC3-PSMA-NY.

Unless specified otherwise, data are presented as mean ± s.e.m. and analyzed using the two-tailed

Student's t-test.

Figure 6. CD40-LY9-FCRL1 CAR-Ms improve solid tumor clearance *in vitro* and *in vivo*

(A) Specific killing of PC3-PSMA cells by UTD and COMB7 CAR-Ms at varying E/T ratios. Data

represent two independent experiments from two human donors.

(B) Specific killing of PC3-PSMA cells by UTD, CD247 CAR-Ms or COMB7 CAR-Ms at an E/T ratio

of 3/1. Data collected from two human donors.

(C-D) Specific killing of HER2$^+$ SKOV3 or GPC3$^+$ HepG2 cells by UTD or COMB7 CAR-Ms at an

E/T ratio of 3/1 (n = 6 biological replicates).

(E) NCG mice were IP injected with 1×10^6 PC3-PSMA cells, followed by treatment with PBS, $5 \times$

10^6 UTD, or 5×10^6 COMB7 CAR-Ms one day later.

(F) Overall survival for mice in each treatment group (n = 6 for PBS, n = 5 for UTD and COMB7).

(G) Tumor burden measured by bioluminescence (total flux). Data shown are individual values (n = 6

for PBS, n = 5 for UTD and COMB7).

(H) Tumor sizes were monitored after tumor injection. Data shown are individual values (n = 5).

(I) Bar plot showing tumor sizes measured at 21, 42, and 63 days after tumor injection. Data are

presented as mean ± s.e.m. and analyzed using one-way ANOVA followed by Dunnett's

multiple-comparison.

Unless specified otherwise, data are presented as mean ± s.e.m. and analyzed using the two-tailed

Student's t-test.

A

Natural receptor Signaling Domain CAR library (NaSDC)

n = 131

Fc receptors: FCER1G, FCGR1A, FCRL1, FCRL2...

Toll-like receptors: TLR1, TLR2, TLR3, TLR4...

TNF receptor superfamily: TNFRSF1A, TNFRSF1B, CD40, LTBR...

Other receptors: CD247, ICOS, HCST...

Functional Modules

Pro-phagocytosis

Pro-inflammation

Pro-infiltration

Combinatorial Signaling Domain CAR library (CoSDC)

Combinatorial CAR Library

ICD i   ICD j   ICD k

CARode: Barcode for CAR ICD

3' LTR

ICD i j k   IRES+EGFP   CARode k j i

Monocytes (THP-1)

Lentiviral CAR Library Transduction

Natural Receptor CAR-M Pool

Combinatorial Signaling CAR-M Pool

scRNA-seq & scCAR-seq

Next-generation CAR-M

Screen for Functional ICD Modules

Screen for Combinatorial CAR Signaling

B

hMDM

EGFP 33.51%

THP−1

EGFP 98.39%

C

Sort 31.01%

0.08 | 98.28
0.25 | 1.39

D

CoSDC Library + CARode

anti-PSMA CAR   ICDs   CARode

IRES+EGFP

scCAR Sequencing

P7   3' LTR   AAAAA   P5   TTTTTT

1. ICD Identity

CARode i j k → CAR1: k+j+i   → CAR2: j+j+i   → CAR3: k+j+k

2. ICDs Order

ICD i j k   k j i CARode   AAAAA

3. CAR Generation

Gen 1 CAR   Gen 2 CAR   Gen 3 CAR

A *Pro-phagocytic Module Screen*

Input CAR-M pool → (Phagocytic cells) → Output (Phagocytic cells)

Barcode Sequencing

B

-log10(rho score)

HCST
TREML1
FCRL1
ICOS
FCRL6
BTLA
LILRB2
TYROBP
FCER1G

-log10(rho)

log2FC
0.0
0.2
0.4

R = 0.628
p < 0.001

log2FoldChange

C Phagocytosis of Target Cells

p < 0.001
p = 0.0031
p = 0.0203
p = 0.0474

Phagocytosis (%)

FCER1G BTLA ICOS TYROBP TREML1 LILRB2 HCST FCRL1 FCRL6

D *Pro-inflammatory Module Screen*

NFRE — iRFP670
NFkB reporter

Input CAR-M pool → Output (iRFP670^high)

Barcode Sequencing

E

-log10(rho score)

CD40
TNFRSF1A
TNFRSF11A
CD3E
LTBR
KIR2DS1
BTLA
FCER1G

log2FC
0.0
0.5
1.0
1.5

-log10(rho)

R = 0.68
p < 0.001

log2FoldChange

F Relative mRNA Expression

TNF
IL6
IL1B

UTD CD40 TNFRSF11A CD3E KIR2DS1 BTLA FCER1G

>6
3
0

1
3
>6

G *Pro-infiltrating Module Screen*

Input CAR-M pool → Output (Intratumoral)

Barcode Sequencing

H

-log10(rho score)

LY9
CD247

log2FC
0.0
0.5
1.0
1.5

-log10(rho)

R = 0.557
p < 0.001

log2FoldChange

I Relative mRNA Expression

MMP13
MMP15
MMP16
MMP17

UTD LY9 CD247

>6
3
0

1
3
>6

A

CoSDC Lentiviral Library → CAR-M Library → Tumor Stimulation → scRNA-seq with Barcode Direct Capture → Decoding CAR-M library phenotype with unique CARode

CARode Sequecing

scRNA Sequecing

B

CAR types per cell

7911 single cells with high−confidence CAR assignment

tSNE2 / tSNE1

Cluster
- c0
- c1
- c2
- c3
- c4
- c5
- c6

C Macrophage Polarization

M2−like / M1−like / Pre-activation / Transitional M1

AMDI / MPI

D Antigen Processing & Presentation

Reactome: 2132295 / Reactome: 983169 / GO: 0019882 / KEGG: 04612

E MMP Levels

Normalized AUC scores

F Phagocytosis Score

KEGG / GO

G

ICD1 / ICD2 / ICD3

HCST, TREML1, ICOS, FCRL1, FCRL6, BTLA, LILRB2, TYROBP, FCER1G, CD40, TNFRSF1A, TNFRSF11A, CD3E, LTBR, KIR2DS1, LY9, CD247

log2FC