Microsporidia infection alters *C. elegans* lipid levels

Hala Tamim El Jarkass[1*], JiHae Jeon[1*], Nick O. Burton[2], Aaron W. Reinke[1#].

[1] Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1A1 Canada

[2] Department of Metabolism and Nutritional Programming, Van Andel Research Institute, Grand Rapids,
MI 49503 USA.

* These authors contributed equally

Corresponding author

aaron.reinke@utoronto.ca

Abstract

Microsporidia are fungal-related obligate intracellular parasites that infect many types of animals.
Microsporidia have exceptionally reduced genomes resulting in limited metabolic capabilities and are
thought to be reliant on host metabolism to fuel their own growth. Here, we investigate the impact of
microsporidia infection on host lipid metabolism using the nematode *Caenorhabditis elegans* along with
its natural microsporidian pathogen *Nematocida parisii*. We show that infection causes an increase in the
level of *C. elegans* lipid droplet associated lipase, ATGL-1, and a decrease in host fat levels. A mutation
that decreases ATGL-1 activity and overexpression of ATGL-1 did not significantly change *N. parisii*
infection levels. Using lipidomics we show that *N. parisii* infection decreases *C. elegans* triglyceride levels
and results in increased ceramides that we speculate are synthesized by *N. parisii*. Mutations in host
genes involved in ceramide synthesis did not significantly change the levels of *N. parisii* infection.
Together these results show that microsporidia can cause changes to lipid metabolism of their hosts, but
some individual mutations of *C. elegans* lipid enzymes do not alter microsporidian growth.

Introduction

Microsporidia are a large group of obligate intracellular parasites with the smallest known eukaryotic genomes [1–3]. Microsporidia infect most groups of animals and many species infect humans [4–7]. Microsporidia genomes are missing many metabolic enzymes, including those involved in lipid metabolism and some species have lost the ability to carry out glycolysis, making these parasites highly dependant upon their hosts for nutrients [8–10]. Microsporidia encode a number of transporters that allow them to acquire nucleotides and energy from their hosts [11–14]. Microsporidia also rely on other types of nutrients from their hosts, including lipids. Several lipids such as phosphatidic acid and linoleic acid were shown to be limiting factors for microsporidia proliferation [15,16]. Microsporidia infection has a profound effect on metabolism in fruit flies, shrimp, crabs, and silkworms, causing a decrease in the levels of many types of lipids [15,17–20].

The free-living nematode *Caenorhabditis elegans* has been developed as a biological model system to study interactions with microsporidia [21]. Several species of microsporidia have been found to infect *C. elegans* either in the wild or in laboratory settings [10,22]. *Nematocida parisii* is the species most frequently found to infect *C. elegans* in the wild and is also the most commonly studied species in laboratory settings. [23]. This parasite has become a model system for microsporidia research and has been used to understand how hosts defend against infection [24–27], mechanisms of invasion [28], proliferation [16,29–31], spore exit [32], and to identify microsporidia inhibitors [16,33,34]. Infection begins when *N. parisii* spores are ingested by the worm, and the spore germinates using its unique infectious apparatus to deposit a sporoplasm inside of an intestinal cell [28,35]. The sporoplasms then divide and proliferate throughout the animal as a form called a meront [36,37]. During this process the 20 intestinal cells of *C. elegans* are fused together into a single cellular syncytium [29]. Sporulation then occurs, the spores exit without lysing the host plasma membrane, and the spores are defecated from the animal [32].

C. elegans fatty acid metabolism is regulated both by environmental stresses such as starvation and by bacteria. The majority of fat in *C. elegans* is stored as lipid droplets in intestinal epithelial cells [38]. Lipid droplets are involved in maintaining energy homeostasis and are important for longevity [39,40]. One of the major enzymes which regulates fat storage is the adipose triglyceride lipase, ATGL-1 [41]. Fasting causes an upregulation of ATGL-1 and a decrease in triglycerides [42,43]. Fatty acid synthesis is regulated in response to pathogen infection [44]. Several types of lipids, including cholesterol, and monosaturated and polyunsaturated fatty acids have been shown to modulate innate immunity in response to pathogens [45–47]. Additionally, pathogenic bacteria and yeast can cause a reduction in lipid droplets [44,48]. Bacteria that are part of the *C. elegans* microbiome can cause changes to fatty acid levels [37,49,50]. Additionally, bacterially produced sphingolipids have been shown to alter sphingolipid biosynthesis in *C. elegans*, providing protection against a pathogenic bacteria [51].

Here, we report the interactions between host lipid metabolism and microsporidia infection. We show that the lipase ATGL-1 is upregulated in response to *N. parisii* infection and that fat stores are depleted. Lipidomic profiling of infected animals reveals large changes in lipid composition including changes consistent with starvation. We also detect many novel ceramide species that appear to be made by *N. parisii*. As *N. parisii* has lost the enzymes to generate ceramides de novo, together these results suggest that *N. parisii* may be modifying host-provided lipids to synthesize these ceramides. We show that mutations in genes in the sphingolipid biosynthesis pathway did not change the amount of *N. parisii* infection. Overall, these experiments highlight changes to host lipid metabolism during microsporidia infection.

Results

ATGL-1 is upregulated and lipid stores are reduced in response to *N. parisii* infection

During fasting-induced nutritional deprivation of *C. elegans*, lipid stores are reduced [43]. This effect is largely mediated by the ATGL-1 lipase which colocalizes with intestinal lipid droplets and hydrolyzes triglycerides to form diglycerides [42]. To determine whether *N. parisii* induces a starvation-like state in *C. elegans*, we tested the effect of microsporidia infection on ATGL-1 expression using strain VS20, which contains an integrated transgenic array that overexpresses GFP-tagged ATGL-1 under its native promoter [43]. We infected the *atgl-1::gfp* strain with *N. parisii* for either 24, 48, or 72 hours. By 72 hours post infection (hpi), we observed a significant increase in ATGL-1 expression in infected animals (**Fig 1A and 1B**). To measure changes in the lipid stores, wild type (N2), *atgl-1::gfp*, and a reduction of function variant of this gene, *atgl-1* (*gk176565*) (ATGL-1 P87S*),* were infected with *N. parisii* for 72 hours*,* fixed, and lipid droplets were stained using Oil Red O dye [43,52]. In wild-type animals there was a significant decrease in lipid staining upon *N. parisii* infection. We also observed a decrease in staining, though not significant, in the two *atgl-1* strains when they were infected with *N. parisii* (**Fig 1C and 1D***).*

ATGL-1 activity impacts the growth of *N. parisii*

To determine the role of ATGL-1 during infection with *N. parisii*, we analyzed the pathogen burden and the effect on the host in *atgl-1* mutant strains. We infected wild type, *atgl-1::gfp, and atgl-1* (*gk176565*) animals for 48 or 72 hours, then fixed each sample and stained with a Fluorescence In Situ Hybridization (FISH) probe specific for *N. parisii* 18S rRNA, which can label the intracellular meront stage [53]. We also stained with Direct Yellow 96 (DY96), which binds to the chitin of worm embryos and microsporidia spores [29,54]. We did not find any significant difference in the growth of *N. parsii* in either *atgl-1* strain at either

time point (**Fig 2A-2C**). To determine how infection impacts the host, we counted the number of embryos as a proxy for estimating the relative fitness of the worm (**Fig 2D**) [26,29]. There was a reduction in embryo number upon infection in all three strains, with infection causing a larger decrease in relative number of embryos in both ATGL-1 mutant strains, compared to N2 (**Fig 2E**).

Lipidomics reveals impact of microsporidia on host lipid metabolism

To understand how *N. parisii* globally alters *C. elegans* lipid metabolism, we measured the lipidome of infected animals. Animals were grown in the presence or absence of *N. parisii* spores, these adults and their embryos were collected, and their lipids were measured using mass spectrometry. In total, 1451 different ionized lipids were reliably detected and 169 exhibited more than two-fold change with a significant p-value ($p < 0.01$) (**Fig 3 and Table S1**). Infected animals displayed decreased levels of triglycerides and increased levels of diglycerides (**Fig 3A**). Long-chain acylcarnitines were also upregulated (**Fig 3B**). Phosphatidylcholines were most impacted by infection with some of these lipids exhibiting more than a 1000-fold change (**Fig 3C**). These highly upregulated phosphatidylcholines were not observed in another study of *C. elegans* lipids, and thus potentially represent metabolites generated by *N. parisii* [55]. Many d17 ceramides are downregulated in response to infection. In contrast, several d18 and d20 ceramides are highly upregulated (**Fig 3D**). These lipids are potentially synthesised by microsporidia as d17 is the only ceramide type known to be synthesized in *C. elegans* [56,57]. In embryos from infected and uninfected parents, 62 significantly regulated lipids were detected. In addition to the regulation of acylcarnitines, phosphatidylcholines, and ceramides, 14 phosphatidic acid molecules were upregulated in embryos from infected parents (**S1 Fig**). Overall, lipid profiling demonstrates that *N. parisii* infection causes a profound alternation to the lipidome of *C. elegans* in both adults and offspring.

Mutations in *C. elegans* sphingolipid biosynthesis impact microsporidian infection

As we observed that putative novel ceramides may be synthesized by *N. parisii*, we sought to understand how host sphingolipid biosynthesis could impact *N. parisii* infection. We tested mutants for 9 out of 11 genes encoded by *C. elegans* which function in the core sphingolipid biosynthetic pathway (**Fig 4**). Wild type and mutants deficient in sphingolipid biosynthesis were infected with *N. parisii* or left uninfected for 72 hours. We observed no significant changes in the levels of meronts or spores, except a small increase in the amount of spores in *asah-2* mutant animals (**Fig 5A and 5B**). To further probe if acid ceramidase mutants increase *N. parisii* infection levels, we tested both *asah-1* and *asah-2* for 48 hours and 72 hours. At the 48-hour time point we observed a slight increase in meronts in both mutants and we again observed a small increase in spores in *asah-2* at 72 hours, though none of these changes were statistically significant (**Fig 5E and 5F**). To determine if a reduction in sphingolipid biosynthesis enzymes affected

the fitness of infected *C. elegans* we measured the number of embryos in sphingolipid mutants in the presence and absence of *N. parisii* spores (**Fig 5C**). We observed that the population of *ttm-5* mutant animals is significantly less gravid than N2, although these mutants have decreased levels of embryos in the absence of infection (**Fig 5C and 5D**).

Sphingosine (d17:1) supplementation does not increase *N. parisii* proliferation

Ceramides are synthesised either through the desaturation of dihydroceramide or through ceramide synthase using sphingosine as a substrate (**Fig 4**). *N. parisii* encodes a ceramide synthase and could be generating ceramides by utilizing host sphingosine. To test if exogenously added sphingosine could play a role during microsporidian infection, we performed infections using media supplemented with sphingosine. Wild-type *C. elegans* were exposed to *N. parisii* for 72 hours on 50 µM sphingosine or control plates containing ethanol. We observed no increase in the amount of spores generated when infected animals were supplemented with sphingosine (**Fig 5G**).

Discussion

To understand if lipids are involved during *N. parisii* infection of *C. elegans*, we determined how infection alters lipid biosynthesis. We show that *N. parisii* infection upregulates ATGL-1, decreases the levels of some lipids, including triglycerides, and increases the levels of many types of lipids, including diglycerides, phosphatidylcholines, and acylcarnitines. We also observed that modifying ATGL-1 activity, reducing the function of some sphingolipid biosynthesis genes, or supplementing with sphingosine did not result in significant changes to *N. parisii* growth. Altogether these results suggest that lipid metabolism is affected by infection, but changes to some lipid levels does not have a large impact on microsporidia growth. One limitation of our results is that there are several steps of sphingolipid biosynthesis where *C. elegans* encodes paralogous enzymes, which could lead to redundancy that prevented us from seeing an effect on microsporidia infection. Additionally, we observed that sphingosine supplementation did not increase *N. parisii* growth and it is possible that using a higher concentration, or using the branched form of sphingosine which is synthesized by *C. elegans* could increase microsporidia proliferation [56].

Large changes to lipid metabolism occur during microsporidia infection. There are several effects of *N. parisii* infection that are similar to starvation. When exposed to fasting conditions, *C. elegans* display upregulated ATGL-1, decreased Oil Red O staining, and decreased triglyceride levels, which are all changes we observed upon *N. parisii* infection [42,43]. We also observed acylcarnitines being upregulated during *N. parisii* infection, which has been shown to occur during starvation [58]. However, there are also differences, such as cardiolipin upregulation during starvation and we observed a downregulation of cardiolipins upon *N. parisii* infection [58]. Infection of *C. elegans* with *N. parisii* and other microsporidia did not show upregulation of *atgl-1* mRNA [59–61]. In fasted worms, *atgl-1* has been

proposed to be post-transcriptionally regulated as there is a larger increase in protein than mRNA, and a
similar regulation may be occurring during *N. parisii* infection [42]. We observed that some lipid mutants
have a decrease in embryo production due to microsporidia infection. We previously showed that vitamin
B12 provides tolerance to *N. parisii* infection and this suggests that depletion of other nutrients can alter
the impact of microsporidia infection on the host [62]. Infection with microsporidia also induces an
intergenerational response which protects progeny from microsporidia infection [26]. The mechanisms of
this intergenerational response are mostly unknown, and it is possible that the changes we see in lipid
levels in offspring from infected animals may play a role in this immunity.

Sphingolipids are an important class of lipids that are components of membranes and used as signaling
molecules [63]. Ceramides are a central type of sphingolipid that is made up of a sphingoid base and a
fatty acyl chain [64]. *C. elegans* generate 17 carbon ceramides (d:17:1) and dihydroceramides (d:17:0)
which contain a branched structure [56]. In several studies profiling lipids in *C. elegans*, d:17:1 and d:17:0
ceramides with different length of fatty acyl chains have been reported, but to the best of our knowledge,
other sphingoid base branch lengths such as d:18 and d:20 have not been observed [55,65–67]. Upon
infection we detected a downregulation of many d:17 ceramides species and upregulation ~30-80 fold of
6 lipid species with either d:18 or d:20 sphingoid bases. Although we have no direct evidence, we
speculate that the most likely explanation for this result is that these longer sphingoid bases are
generated by *N. parisii* enzymes. Other species of fungi have been observed to generate ceramides
containing d:18 and d:20 sphingoid bases [68,69]. Most microsporidia, like *N. parisii*, encode a partial
ceramide synthesis pathway, but do not have the ability to make ceramides de novo, whereas *Edhazardia*
aedis contains the full ceramide synthesis pathway and *Enterocytozoon bieneusi* has lost the entire
pathway.

In conclusion, we provide evidence that *C. elegans* lipid biosynthesis is affected by *N. parisii* infection.
We have previously shown that mutants in the fatty acid desaturase *fat-2*, which is responsible for
synthesising linoleic acid, delays *N. parisii* infection [16]. Future work using other lipid mutants or using a
combination of mutants and RNAi for steps with multiple enzymes is likely to uncover other pathways
necessary for *N. parisii* growth. Additionally, labeling *C. elegans* lipids could be used to determine if and
how *N. parisii* can modify host lipids to generate ceramides [70].

Acknowledgments

We are grateful to Meng Xiao, Yin Chen Wan, and Jonathan Tersigni for providing helpful comments on
the manuscript. *C. elegans* strains were provided by the *Caenorhabditis* Genetics Center, which is
funded by the National Institutes of Health (NIH) Office of Research Infrastructure Programs Grant P40
OD010440 and we thank WormBase. **Funding:** This work was supported by the Canadian Institutes of
Health Research grant no. 400784 (to A. R.) **Competing interests:** The authors declare that they have

no competing interests. **Data availability:** All data needed to evaluate the conclusions in the paper are
present in the paper and supplementary materials. Source Data are provided in **Data S1**.

Contributions: N. B. and A.R. conceived of the project. A. R. performed infections for the lipidomic data
and N. B. generated the lipidomic data. J. J. performed the ATGL-1 fluorescence and infection
experiments as well as preliminary infections on other lipid mutants. H.T.E.J. performed all other
experiments. All authors analyzed the data. J. J. and A. R. co-wrote the paper with edits from the other
authors.

Materials and Methods

Nematode strains and microsporidia culture
All *C. elegans* strains were maintained at 21 °C on Nematode Growth Medium (NGM) plates seeded with
Escherichia coli OP50-1, as previously described [71]. All strains were derived from the Bristol N2 wild-
type animals and obtained from the Caenorhabditis Genetics Centre (CGC). Details of strains used are
listed in Table S2. Synchronized populations of worms were generated by treating mixed stage
populations with NaOH and hypochlorite, washing with M9, and hatching with rotation for 18-24 hours at
21 °C. *Escherichia coli* OP50-1 was grown at 37 °C for 18-24 in LB and then concentrated to make 10x
stocks. *N. parisii* (ERTm1) spores were prepared as previously described [26]. Briefly, infected
populations of *C. elegans* were harvested, disrupted with 2 mm diameter zirconia beads, and filtered (5
μm). Spores were confirmed to be free of bacterial contamination, counted by staining with DY96 and
using a sperm counting slide, and frozen in aliquots at -80 °C.

Infection and fluorescence imaging of ATGL-1::GFP

L1 stage *atgl-1::gfp* worms were divided into three groups and infected or left uninfected for 24, 48, or 72
h. For the 72 h infected group, 1,000 L1 animals were mixed with 1 million spores and 250 μl 10x OP50
and placed onto 3.5-cm NGM plate. For the 24 h and 48 h infected groups, 1,000 L1 stage worms were
first grown on NGM plates seeded with OP50-1 for 48 h and 24 h, respectively. The worms were then
washed in M9 and mixed with 1 million spores and 250 μl 10X OP50 and placed onto 3.5-cm NGM plates.
Uninfected controls were treated the same way as the infected populations except no *N. parisii* spores
were added. Plates were incubated at 21 °C for a total of 72 hours. Worms were then paralyzed in 1 mM
levamisole for 2 minutes and then transferred onto a freshly prepared 3% agarose pad on glass slide.
Imaging was performed using a ZEISS Axio Imager 2 and the level of GFP was quantified for each animal
and normalized by worm size (net fluorescence divided by its total area) using FIJI 2.1.0 [72].

Oil red O staining and imaging

1,000 L1 stage animals, 400 µl of 10x *E. coli* OP50 and 3.5 million spores of *N. parisii* were mixed, plated

onto a 6-cm unseeded NGM plate and incubated at 21 °C for 72 hours. Uninfected controls contained a

volume of M9 equivalent to that used for spores. After 72 hours, animals were washed with PBST, fixed

with 40% isopropanol stained with Oil Red O solution as previously described [73]. Animals were mounted

onto glass slides and imaged using a ZEISS Axio Imager 2 with a mobile phone camera to capture stained

animals. Oil red O levels were quantified using FIJI by first drawing around individual animals to designate

regions of interest, followed by thresholding and measuring stain intensity.

N. parisii infection of lipid mutants

L1 stage animals were mixed with OP50 and *N. parisii* spores and placed onto an NGM plates. Infections

of synchronized *atgl-1::gfp* and *atgl-1 (gk176565)* worms were done with 1,000 L1s, 1.5 million spores,

and 250 µl 10x OP50 on a 3.5-cm plate. Infection of all other lipid mutants was done with 1,000 L1s, 3.5

million spores, and 400 µl 10x OP50 on a 6-cm NGM plate. Infection plates were incubated at 21 °C for

either 48 or 72 hours, depending on the experiment. Worms were washed off plates with M9 + 0.1%

Tween-20 and fixed with acetone.

DY96 and FISH staining

To stain spores and embryos, fixed worms were washed twice with 1 ml PBS + 0.1% Tween-20 then

incubated in 500 µl of DY96 staining solution (1x PBS, 0.1% Tween-20, 20 µg/ml DY96, and 0.1% SDS)

for 30 minutes at 21 °C. To stain microsporidia meronts, fixed worms were washed twice with 1 ml PBS

+ 0.1% Tween-20 and once with 1 ml hybridization buffer (900 mM NaCl, 20 mM Tris [pH 8.0], 0.01%

SDS in dH2O). Supernatant was then removed, 100 µl hybridization buffer containing 5 ng/µl FISH probe

was added and the worms were incubated at about 46 °C overnight. The probe used in the study was

MicroB with Cal Fluor Red 610 attached (CTCTCGGCACTCCTTCCTG) [53]. Samples were washed

once with wash buffer (hybridization buffer containing 5 mM EDTA and 20 µg/mL DY96). Worms were

incubated in 500 µl wash buffer at 46 °C for 30 - 60 minutes. Supernatant was removed and stained

worms were mounted on glass slides using 20 µl Everbrite containing DAPI for imaging with a ZEISS

Axio Imager 2. Spore and meront fluorescence was quantified using FIJI 2.1.0 by thresholding

fluorescence intensity and the remaining fluorescence signal was measured as percentage of the

nematode body area [54,72].

Lipidomics of *N. parisii* infected *C. elegans*

258 Synchronized L1 N2 animals were prepared by sodium hypochlorite treatment of mixed populations of

259 worms. Infections were carried out by adding 10,000 worms, 5 million *N. parisii* spores, and 1 ml 10x

260 OP50-1 per plate onto 13 10-cm NGM plates. For uninfected animals, 2,500 animals and 1 ml 10x OP50-

1 were added per plate onto 10 10-cm NGM plates. Animals were incubated at 21°C for 72 hours. For adult samples, worms from 1 plate of each condition were washed into 1.5 ml tubes and then washed twice with 1 ml M9. To remove embryos and L1s, worms were allowed to settle for 1 minute, supernatant was removed, and 1 ml M9 was added. This was done a total of 4 times. Samples were then frozen in liquid nitrogen.

To prepare embryos, the remaining infected and uninfected plates were treated with sodium hypochlorite in 15 mL conical tubes and washed 3 times with 10 ml M9 and resuspended in 5 ml M9. To remove carcasses, tubes were left to settle for 4 minutes, supernatant removed and washed twice with 10 mL M9. Embryos were then added to a 10-cm NGM plate and incubated for 3 hours at room temperature. Embryos were washed off plates with M9 into 1.5 ml tubes and froze in liquid nitrogen. Experiment was performed in triplicate.

Liquid chromatography separation with mass spectrometry detection (LC–MS) of intact lipid species was performed as described previously [74].

Sphingosine Supplementation

Sphingosine (d17:1, from Cayman Chemicals) was dissolved in 100% ethanol as a vehicle to form a stock of 10 mM. This stock was then used to supplement sphingosine to NGM plates post autoclave to a final concentration of 50 uM sphingosine and 0.5% ethanol. This concentration of sphingosine was chosen based on a previous study performing sphingosine complementation in *C. elegans* [56]. Infections were performed using 400 L1 stage N2 animals, 160 µl of 10x OP50, and 1 million *N. parisii* spores on a 3.5-cm plate. After a 72-hour incubation at 21 °C, worms were harvested, stained with DY96, and imaged as described above.

Fig 1. ATGL-1 levels increase and lipid droplets decrease in response to *N. parisii* infection. (A-B) *atgl-1::gfp* L1 stage *C. elegans* were cultured for a total of 72 hours and were either left uninfected or infected for 24, 48, or 72 hours. **(A)** Representative fluorescence images of live adult *atg-1::gfp* worms. White lines indicate the boundaries of worm bodies. Scale bars are 124 µm. **(B)** Quantification of ATGL-1::GFP. Data is from three independent replicates of 18-20 worms each. **(C-D)**. L1 stage Wildtype (N2), *atgl-1::gfp, atgl-1 (gk176565)* worms of each strain were either infected or left uninfected for 72 hours and stained with Oil Red O. **(C)** Representative images of Oil Red O stained worms. **(D)** Quantification percentage Oil Red O intensity per worm. Data is from three independent replicates of at least 20 worms each. Mean ± SD represented by horizontal bars. P-values were determined using one-way ANOVA with post hoc Šidák correction with tests between infected and uninfected samples of the same strain (A and B) and between uninfected samples of different strains (B). Significance is defined as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, and ns means not significant.

Fig 2. Increased or reduced ATGL-1 activity does not significantly change *N. parisii* growth. L1
stage wild-type (N2), *atgl-1::gfp, atgl-1 (gk176565)* worms were either not infected or infected with *N.
parisii* for 48 **(A)** or 72 hours **(D-E)**. Worms were then fixed and stained with a FISH probe against the *N.
parisii* 18S RNA and DY96. **(A-B)** Quantitation of pathogen burden (meronts) per animal. **(C)**
Quantification of pathogen burden (spores) per animal **(D)** Quantification of number of embryos per
animal. **(E)** Number of embryos per worm, normalized to the uninfected condition for each strain. Data is
from three independent replicates of at least 20 worms each. Mean ± SD represented by horizontal bars.
P-values were determined using one-way ANOVA with post hoc Dunnett's test with comparisons to N2
(A-C), Šidák correction (D and E) with tests between infected and uninfected samples of the same strain
(D and E), and between infected samples of different strains (E). Significance is defined as * $p < 0.05$, **
$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, and ns means not significant.

Fig 3. *N. parisii* induces global changes to *C. elegans* lipidome. (A-D) *C. elegans* were infected or left uninfected for 72 h and lipids measured using LC/MS. Heat map showing lipid metabolites that are upregulated (blue) or downregulated (yellow) upon infection at least 2-fold with a significant p-value (p < 0.01). **(A)** All classes of lipids with 2 or more lipids significantly up or down regulated. **(B)** Acylcarnitines. **(C)** Phosphatidylcholines. **(D)** Ceramides. Those lipids detected in Burton et al. 2017 colored in orange [55].

serine + fatty acyl-CoA

serine palmitoyl transferase
SPTL-1, SPTL-2, SPTL-3

3-ketodihydrosphingosine

3-ketodihydrosphingosine reductase
Y37E11AM.3

Human
C. elegans
Yeast
N. parisii

sphinganine

ceramide synthase
LAGR-1, HYL-1, HYL-2

dihydroceramide

dihydroceramide desaturase
TTM-5, F33D4.4

ceramide

ceramide synthase
LAGR-1, HYL-1, HYL-2

acid ceramidase
ASAH-1, ASAH-2

sphingosine

329 **Fig 4. Sphingolipid biosynthetic pathway.** Schematic of sphingolipid biosynthesis pathway.
330 Metabolites are indicated in italics, enzymes indicated in bold, and *C. elegans* enzyme names are
331 indicated in purple [75]. Presence of enzymes from humans, *C. elegans*, yeast (*Saccharomyces*
332 *cerevisiae*), and *N. parisii* was determined with BLAST, using a threshold E-value of greater than 10^{-5}
333 [76]. The presence of the enzyme is indicated by a colored circle corresponding to the legend at the left.

Fig 5. *N. parisii* infection is not significantly changed in *C. elegans* sphingolipid biosynthesis mutants. (A-D) L1 stage N2 or indicated mutant worms were either not infected or infected with *N. parisii* for 72 hours. Worms were then fixed and stained with a FISH probe against the *N. parisii* 18S and DY96. **(A)** Quantitation of pathogen burden (meronts) per animal. **(B)** Quantification of pathogen burden (spores) per animal. **(C)** Quantification of number of embryos per animal. **(D)** Quantification of population gravidity (gravidity is defined as an animal carrying at least 1 embryo), normalized to the uninfected condition for each strain. **(E-F)** L1 stage wild-type (N2), *asah-1 (ve779), asah-2 (ok564)* worms were infected with *N. parisii* for 48 (E) or 72 hours (F), fixed, and stained with FISH probe and DY96. **(E)** Quantitation of pathogen burden (meronts) per animal. **(F)** Quantification of pathogen burden (spores) per animal. **(G)** L1 stage N2 animals were infected on control or sphingosine containing medium for 72 hours, fixed, and

stained with DY96. Pathogen burden (spores) was quantified. Data is from three independent replicates of at least 20 worms each, except for (G) which had two replicates. Mean ± SD represented by horizontal bars. P-values were determined using one-way ANOVA with post hoc Dunnett's test with comparisons to N2 and those that have a $p > .05$ and are not shown (A-D), Šidák correction with tests between infected and uninfected samples of the same strain (E), and Tukey's multiple comparison between all samples (F). Significance is defined as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, and ns means not significant.

Supplementary materials

S1 Fig. *N. parisii* infection of *C. elegans* parents alters lipids in offspring.

(A-E) *C. elegans* were infected for 72 hours and were harvested using sodium hypochlorite treatment.

After 3 hours embryos were harvested and lipids measured using LC/MS. Heat map showing lipid

metabolites that are upregulated or downregulated in embryos from infected parents by more than 2-fold

with a significant p-value ($p < 0.01$). (A) All classes of lipids significantly upregulated or downregulated.

(B) Acylcarnitines. (C) Phosphatidylcholines. (D) Phosphatidic acids. (E) Ceramides. (B-E) Lipid species

detected in Burton et al. 2017 colored in orange [55].

Table S1. Lipidomic data from infected and uninfected adults and their embryos.

Table S2. List of *C. elegans* strains used in the study.

Strain	Genotype
N2	N2
VS20	*hjIs67 [atgl-1p::atgl-1::GFP + mec-7::RFP]*
VC20458	*atgl-1 (gk176565) [P87S] III*
RG3279	*asah-1 (ve779 [LoxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + LoxP]) I*
RB782	*asah-2 (ok564) II*
RB1465	*sptl-1 (ok1693) II*
RB1579	*sptl-3 (ok1927) V*
RB1036	*hyl-1 (ok976) IV*
VC334	*hyl-1 (gk203) IV*
RB1498	*hyl-2 (ok1766) X*
RB1685	*ttm-5 (ok2095) I*
RB2135	*F33D4.4 (ok2843) IV*
VC507	*lagr-1 (gk263) I*
VC765	*lagr-1 (gk331) I*

391   **Data S1. All infection data from the study.**

References

1.   Wadi L, Reinke AW. Evolution of microsporidia: An extremely successful group of eukaryotic intracellular
parasites. PLoS Pathog. 2020;16: e1008276. doi:10.1371/journal.ppat.1008276

2.   Katinka M, Duprat S, Cornillot E, Méténier G, Thomarat F, Prensier G, et al. Genome sequence and gene
compaction of the eukaryote parasite Encephalitozoon cuniculi. Nature. 2001;414: 450–3.
doi:10.1038/35106579

3.   Vavra J, Lukes J. Microsporidia and "the art of living together". Adv Parasitol. 2013;82: 253–319.
doi:10.1016/B978-0-12-407706-5.00004-6

4.   Murareanu Brandon M., Sukhdeo Ronesh, Qu Rui, Jiang Jason, Reinke Aaron W., Weiss Louis M.
Generation of a Microsporidia Species Attribute Database and Analysis of the Extensive Ecological and
Phenotypic Diversity of Microsporidia. mBio. 12: e01490-21. doi:10.1128/mBio.01490-21

5.   Willis AR, Reinke AW. Factors That Determine Microsporidia Infection and Host Specificity. Exp Suppl.
2022;114: 91–114. doi:10.1007/978-3-030-93306-7_4

6.   Bojko J, Reinke AW, Stentiford GD, Williams B, Rogers MSJ, Bass D. Microsporidia: a new taxonomic,
evolutionary, and ecological synthesis. Trends in Parasitology. 2022;38: 642–659.
doi:10.1016/j.pt.2022.05.007

7.   Han B, Pan G, Weiss LM. Microsporidiosis in Humans. Clinical Microbiology Reviews. 2021;34: e00010-20.
doi:10.1128/CMR.00010-20

8.   Nakjang S, Williams TA, Heinz E, Watson AK, Foster PG, Sendra KM, et al. Reduction and expansion in
microsporidian genome evolution: new insights from comparative genomics. Genome Biol Evol. 2013;5:
2285–2303. doi:10.1093/gbe/evt184

9.   Wiredu Boakye D, Jaroenlak P, Prachumwat A, Williams TA, Bateman KS, Itsathitphaisarn O, et al. Decay of
the glycolytic pathway and adaptation to intranuclear parasitism within Enterocytozoonidae
microsporidia. Environ Microbiol. 2017;19: 2077–2089. doi:10.1111/1462-2920.13734

10.   Wadi L, El Jarkass HT, Tran TD, Islah N, Luallen RJ, Reinke AW. Genomic and phenotypic evolution of
nematode-infecting microsporidia. PLOS Pathogens. 2023;19: e1011510.
doi:10.1371/journal.ppat.1011510

11.   Heinz E, Hacker C, Dean P, Mifsud J, Goldberg AV, Williams TA, et al. Plasma membrane-located purine
nucleotide transport proteins are key components for host exploitation by microsporidian intracellular
parasites. PLoS Pathog. 2014;10: e1004547. doi:10.1371/journal.ppat.1004547

12.   Dean P, Sendra KM, Williams TA, Watson AK, Major P, Nakjang S, et al. Transporter gene acquisition and
innovation in the evolution of Microsporidia intracellular parasites. Nat Commun. 2018;9: 1709.
doi:10.1038/s41467-018-03923-4

13.   Major P, Sendra KM, Dean P, Williams TA, Watson AK, Thwaites DT, et al. A new family of cell surface
located purine transporters in Microsporidia and related fungal endoparasites. Elife. 2019;8.
doi:10.7554/eLife.47037

14. Dean P, Hirt RP, Embley TM. Microsporidia: Why Make Nucleotides if You Can Steal Them? PLoS Pathog.
2016;12: e1005870. doi:10.1371/journal.ppat.1005870

15. Franchet A, Niehus S, Caravello G, Ferrandon D. Phosphatidic acid as a limiting host metabolite for the
proliferation of the microsporidium Tubulinosema ratisbonensis in Drosophila flies. Nature Microbiology.
2019;4: 645–655. doi:10.1038/s41564-018-0344-y

16. Jarkass HTE, Castelblanco S, Kaur M, Wan YC, Ellis AE, Sheldon RD, et al. The Caenorhabditis elegans
bacterial microbiome influences microsporidia infection through nutrient limitation and inhibiting parasite
invasion. bioRxiv; 2024. p. 2024.06.05.597580. doi:10.1101/2024.06.05.597580

17. Ning M, Wei P, Shen H, Wan X, Jin M, Li X, et al. Proteomic and metabolomic responses in hepatopancreas
of whiteleg shrimp Litopenaeus vannamei infected by microsporidian Enterocytozoon hepatopenaei. Fish
Shellfish Immunol. 2019;87: 534–545. doi:10.1016/j.fsi.2019.01.051

18. Wu Y, Chen J, Liao G, Hu M, Zhang Q, Meng X, et al. Down-Regulation of Lipid Metabolism in the
Hepatopancreas of Shrimp Litopenaeus vannamei upon Light and Heavy Infection of Enterocytozoon
hepatopenaei: A Comparative Proteomic Study. International Journal of Molecular Sciences. 2022;23:
11574. doi:10.3390/ijms231911574

19. Ding Z. Lipid metabolism disorders contribute to the pathogenesis of Hepatospora eriocheir in the crab
Eriocheir sinensis. Journal of Fish Diseases. 2021;44: 305–313. doi:10.1111/jfd.13284

20. Su Y, Liu M, Li M, Han Z, Lü D, Zhang Y, et al. Metabolomic analysis of lipid changes in *Bombyx mori*
infected with *Nosema bombycis*. Developmental & Comparative Immunology. 2023;147: 104750.
doi:10.1016/j.dci.2023.104750

21. Tecle E, Troemel ER. Insights from C. elegans into Microsporidia Biology and Host-Pathogen Relationships.
Exp Suppl. 2022;114: 115–136. doi:10.1007/978-3-030-93306-7_5

22. Zhang G, Sachse M, Prevost M-C, Luallen RJ, Troemel ER, Felix M-A. A Large Collection of Novel Nematode-
Infecting Microsporidia and Their Diverse Interactions with Caenorhabditis elegans and Other Related
Nematodes. PLoS Pathog. 2016;12: e1006093. doi:10.1371/journal.ppat.1006093

23. Gang SS, Lažetić V. Microsporidia: Pervasive natural pathogens of Caenorhabditis elegans and related
nematodes. Journal of Eukaryotic Microbiology. 2024;71: e13027. doi:10.1111/jeu.13027

24. Reddy KC, Dror T, Sowa JN, Panek J, Chen K, Lim ES, et al. An Intracellular Pathogen Response Pathway
Promotes Proteostasis in C. elegans. Current Biology. 2017;27: 3544-3553.e5.
doi:10.1016/j.cub.2017.10.009

25. Lažetić V, Wu F, Cohen LB, Reddy KC, Chang Y-T, Gang SS, et al. The transcription factor ZIP-1 promotes
resistance to intracellular infection in Caenorhabditis elegans. Nat Commun. 2022;13: 17.
doi:10.1038/s41467-021-27621-w

26. Willis AR, Zhao W, Sukhdeo R, Wadi L, Jarkass HTE, Claycomb JM, et al. A parental transcriptional response
to microsporidia infection induces inherited immunity in offspring. Science Advances. 2021;7: eabf3114.
doi:10.1126/sciadv.abf3114

27.   Tecle E, Chhan CB, Franklin L, Underwood RS, Hanna-Rose W, Troemel ER. The purine nucleoside
476   phosphorylase pnp-1 regulates epithelial cell resistance to infection in C. elegans. PLOS Pathogens.
477   2021;17: e1009350. doi:10.1371/journal.ppat.1009350

478   28.   Tamim El Jarkass H, Mok C, Schertzberg MR, Fraser AG, Troemel ER, Reinke AW. An intestinally secreted
479   host factor promotes microsporidia invasion of C. elegans. Lourido S, Sengupta P, editors. eLife. 2022;11:
480   e72458. doi:10.7554/eLife.72458

481   29.   Balla KM, Luallen RJ, Bakowski MA, Troemel ER. Cell-to-cell spread of microsporidia causes Caenorhabditis
482   elegans organs to form syncytia. Nature Microbiology. 2016;1: 16144.

483   30.   Luallen RJ, Reinke AW, Tong L, Botts MR, Felix M-A, Troemel ER. Discovery of a Natural Microsporidian
484   Pathogen with a Broad Tissue Tropism in Caenorhabditis elegans. PLoS Pathog. 2016;12: e1005724.
485   doi:10.1371/journal.ppat.1005724

486   31.   Reinke AW, Balla KM, Bennett EJ, Troemel ER. Identification of microsporidia host-exposed proteins
487   reveals a repertoire of rapidly evolving proteins. Nat Commun. 2017;8: 14023. doi:10.1038/ncomms14023

488   32.   Szumowski SC, Botts MR, Popovich JJ, Smelkinson MG, Troemel ER. The small GTPase RAB-11 directs
489   polarized exocytosis of the intracellular pathogen N. parisii for fecal-oral transmission from C. elegans. Proc
490   Natl Acad Sci U S A. 2014;111: 8215–8220. doi:10.1073/pnas.1400696111

491   33.   Murareanu BM, Antao NV, Zhao W, Dubuffet A, El Alaoui H, Knox J, et al. High-throughput small molecule
492   screen identifies inhibitors of microsporidia invasion and proliferation in C. elegans. Nat Commun. 2022;13:
493   5653. doi:10.1038/s41467-022-33400-y

494   34.   Huang Q, Chen J, Pan G, Reinke AW. Screening of the Pandemic Response Box identifies anti-microsporidia
495   compounds. PLOS Neglected Tropical Diseases. 2023;17: e0011806. doi:10.1371/journal.pntd.0011806

496   35.   Han B, Takvorian PM, Weiss LM. Invasion of Host Cells by Microsporidia. Front Microbiol. 2020;11.
497   doi:10.3389/fmicb.2020.00172

498   36.   Troemel ER, Félix M-A, Whiteman NK, Barrière A, Ausubel FM. Microsporidia Are Natural Intracellular
499   Parasites of the Nematode Caenorhabditis elegans. Ward GE, editor. PLoS Biol. 2008;6: e309.
500   doi:10.1371/journal.pbio.0060309

501   37.   Jarkass HTE, Reinke AW. The ins and outs of host-microsporidia interactions during invasion, proliferation
502   and exit. Cellular Microbiology. 2020;22: e13247. doi:10.1111/cmi.13247

503   38.   Mak HY. Lipid droplets as fat storage organelles in Caenorhabditis elegans: Thematic Review Series: Lipid
504   Droplet Synthesis and Metabolism: from Yeast to Man. Journal of Lipid Research. 2012;53: 28–33.
505   doi:10.1194/jlr.R021006

506   39.   Papsdorf K, Miklas JW, Hosseini A, Cabruja M, Morrow CS, Savini M, et al. Lipid droplets and peroxisomes
507   are co-regulated to drive lifespan extension in response to mono-unsaturated fatty acids. Nat Cell Biol.
508   2023;25: 672–684. doi:10.1038/s41556-023-01136-6

509   40.   Watts JL. Fat synthesis and adiposity regulation in Caenorhabditis elegans. Trends in Endocrinology &
510   Metabolism. 2009;20: 58–65. doi:10.1016/j.tem.2008.11.002

511   41.   Zhang SO, Box AC, Xu N, Le Men J, Yu J, Guo F, et al. Genetic and dietary regulation of lipid droplet
512   expansion in Caenorhabditis elegans. Proceedings of the National Academy of Sciences. 2010;107: 4640–
513   4645. doi:10.1073/pnas.0912308107

514   42.   Lee JH, Kong J, Jang JY, Han JS, Ji Y, Lee J, et al. Lipid droplet protein LID-1 mediates ATGL-1-dependent
515   lipolysis during fasting in Caenorhabditis elegans. Mol Cell Biol. 2014/09/08 ed. 2014;34: 4165–4176.
516   doi:10.1128/MCB.00722-14

517   43.   Zaarur N, Desevin K, Mackenzie J, Lord A, Grishok A, Kandror KV. ATGL-1 mediates the effect of dietary
518   restriction and the insulin/IGF-1 signaling pathway on longevity in C. elegans. Molecular Metabolism.
519   2019;27: 75–82. doi:https://doi.org/10.1016/j.molmet.2019.07.001

520   44.   Dasgupta M, Shashikanth M, Gupta A, Sandhu A, De A, Javed S, et al. NHR-49 Transcription Factor
521   Regulates Immunometabolic Response and Survival of Caenorhabditis elegans during Enterococcus faecalis
522   Infection. Infection and Immunity. 2020;88: 10.1128/iai.00130-20. doi:10.1128/iai.00130-20

523   45.   Nandakumar M, Tan M-W. Gamma-linolenic and stearidonic acids are required for basal immunity in
524   Caenorhabditis elegans through their effects on p38 MAP kinase activity. PLoS Genet. 2008/11/21 ed.
525   2008;4: e1000273–e1000273. doi:10.1371/journal.pgen.1000273

526   46.   Anderson SM, Cheesman HK, Peterson ND, Salisbury JE, Soukas AA, Pukkila-Worley R. The fatty acid oleate
527   is required for innate immune activation and pathogen defense in Caenorhabditis elegans. PLOS
528   Pathogens. 2019;15: e1007893. doi:10.1371/journal.ppat.1007893

529   47.   Peterson ND, Icso JD, Salisbury JE, Rodríguez T, Thompson PR, Pukkila-Worley R. Pathogen infection and
530   cholesterol deficiency activate the C. elegans p38 immune pathway through a TIR-1/SARM1 phase
531   transition. Gruber J, Kaeberlein M, Troemel ER, editors. eLife. 2022;11: e74206. doi:10.7554/eLife.74206

532   48.   Nhan JD, Turner CD, Anderson SM, Yen C-A, Dalton HM, Cheesman HK, et al. Redirection of SKN-1 abates
533   the negative metabolic outcomes of a perceived pathogen infection. Proceedings of the National Academy
534   of Sciences. 2019;116: 22322–22330. doi:10.1073/pnas.1909666116

535   49.   Han H-F, Nien S-F, Jiang H-S, Wu J-C, Chiang C-Y, Li M-T, et al. Dietary bacteria control C. elegans fat
536   content through pathways converging at phosphatidylcholine. eLife. 2024;13. doi:10.7554/eLife.96473.1

537   50.   Fox BW, Helf MJ, Burkhardt RN, Artyukhin AB, Curtis BJ, Palomino DF, et al. Evolutionarily related host and
538   microbial pathways regulate fat desaturation in C. elegans. Nat Commun. 2024;15: 1520.
539   doi:10.1038/s41467-024-45782-2

540   51.   Peters L, Drechsler M, Pees B, Angelidou G, Salzer L, Moors KA, et al. Polyketide synthase-derived
541   sphingolipids determine microbiota-mediated protection against pathogens in C. elegans. bioRxiv; 2024. p.
542   2024.02.06.579051. doi:10.1101/2024.02.06.579051

543   52.   O'Rourke EJ, Soukas AA, Carr CE, Ruvkun G. C. elegans Major Fats Are Stored in Vesicles Distinct from
544   Lysosome-Related Organelles. Cell Metabolism. 2009;10: 430–435. doi:10.1016/j.cmet.2009.10.002

545   53.   Troemel ER, Felix M-A, Whiteman NK, Barriere A, Ausubel FM. Microsporidia are natural intracellular
546   parasites of the nematode Caenorhabditis elegans. PLoS Biol. 2008;6: 2736–2752.
547   doi:10.1371/journal.pbio.0060309

54.   Willis AR, Jarkass HTE, Reinke AW. Studying Inherited Immunity in a Caenorhabditis elegans Model of
Microsporidia Infection. JoVE (Journal of Visualized Experiments). 2022; e63636. doi:10.3791/63636

55.   Burton NO, Furuta T, Webster AK, Kaplan REW, Baugh LR, Arur S, et al. Insulin-like signalling to the
maternal germline controls progeny response to osmotic stress. Nat Cell Biol. 2017;19: 252–257.
doi:10.1038/ncb3470

56.   Hannich JT, Mellal D, Feng S, Zumbuehl A, Riezman H. Structure and conserved function of iso-branched
sphingoid bases from the nematode Caenorhabditis elegans. Chem Sci. 2017;8: 3676–3686.
doi:10.1039/C6SC04831E

57.   Cheng X, Jiang X, Tam KY, Li G, Zheng J, Zhang H. Sphingolipidomic Analysis of *C. elegans* reveals
Development- and Environment-dependent Metabolic Features. Int J Biol Sci. 2019;15: 2897–2910.
doi:10.7150/ijbs.30499

58.   Dall KB, Havelund JF, Harvald EB, Witting M, Færgeman NJ. HLH-30-dependent rewiring of metabolism
during starvation in C. elegans. Aging Cell. 2021;20: e13342. doi:10.1111/acel.13342

59.   Bakowski MA, Desjardins CA, Smelkinson MG, Dunbar TA, Lopez-Moyado, IF, Rifkin SA, et al. Ubiquitin-
Mediated Response to Microsporidia and Virus Infection in C. elegans. PLoS Pathog. 2014;10: e1004200.
doi:https://doi.org/10.1371/journal.ppat.1004200

60.   Mok C, Xiao MA, Wan YC, Zhao W, Ahmed SM, Luallen R, et al. High-throughput phenotyping of C. elegans
wild isolates reveals specific resistance and susceptibility traits to infection by distinct microsporidia
species. bioRxiv; 2022. p. 2022.06.20.496912. doi:10.1101/2022.06.20.496912

61.   Wan YC, Troemel ER, Reinke AW. Conservation of Nematocida microsporidia gene expression and host
response in Caenorhabditis nematodes. PLOS ONE. 2022;17: e0279103. doi:10.1371/journal.pone.0279103

62.   Willis AR, Zhao W, Sukhdeo R, Burton NO, Reinke AW. Parental dietary vitamin B12 causes
intergenerational growth acceleration and protects offspring from pathogenic microsporidia and bacteria.
iScience. 2024;27. doi:10.1016/j.isci.2024.110206

63.   Hannun YA, Obeid LM. Sphingolipids and their metabolism in physiology and disease. Nat Rev Mol Cell Biol.
2018;19: 175–191. doi:10.1038/nrm.2017.107

64.   Lee M, Lee SY, Bae Y-S. Functional roles of sphingolipids in immunity and their implication in disease. Exp
Mol Med. 2023;55: 1110–1130. doi:10.1038/s12276-023-01018-9

65.   Zhu H, You Y, Yu B, Deng Z, Liu M, Hu Z, et al. Loss of the ceramide synthase HYL-2 from Caenorhabditis
elegans impairs stress responses and alters sphingolipid composition. Journal of Biological Chemistry.
2024;300. doi:10.1016/j.jbc.2024.107320

66.   Luo H, Zhao X, Wang Z-D, Wu G, Xia Y, Dong M-Q, et al. Sphingolipid profiling reveals differential functions
of sphingolipid biosynthesis isozymes of *Caenorhabditis elegans*. Journal of Lipid Research. 2024;65:
100553. doi:10.1016/j.jlr.2024.100553

67.   Hänel V, Pendleton C, Witting M. The sphingolipidome of the model organism *Caenorhabditis elegans*.
Chemistry and Physics of Lipids. 2019;222: 15–22. doi:10.1016/j.chemphyslip.2019.04.009

68. Usmani SA, Kumar M, Arya K, Ali B, Bhardwaj N, Gaur NA, et al. Beyond membrane components:
uncovering the intriguing world of fungal sphingolipid synthesis and regulation. Research in Microbiology.
2023;174: 104087. doi:10.1016/j.resmic.2023.104087

69. Montefusco DJ, Matmati N, Hannun YA. The yeast sphingolipid signaling landscape. Chemistry and Physics
of Lipids. 2014;177: 26–40. doi:10.1016/j.chemphyslip.2013.10.006

70. Rivera Sánchez R, Bandi S, Scheidt M-D, Laaroussi H, William Fox B, Ishida Y, et al. iso-Fatty Acid
Metabolism in Caenorhabditis elegans' Ceramide Biosynthesis. Helvetica Chimica Acta. 2023;106:
e202300131. doi:10.1002/hlca.202300131

71. Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77: 71–94.

72. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform
for biological-image analysis. Nature Methods. 2012;9: 676–682. doi:10.1038/nmeth.2019

73. Escorcia W, Ruter DL, Nhan J, Curran SP. Quantification of Lipid Abundance and Evaluation of Lipid
Distribution in Caenorhabditis elegans by Nile Red and Oil Red O Staining. Journal of Visualized
Experiments (JoVE). 2018; e57352. doi:10.3791/57352

74. Burton N, Dwivedi V, Burkhart K, Kaplan R, Baugh L, Horvitz H. Neurohormonal signaling via a
sulfotransferase antagonizes insulin-like signaling to regulate a Caenorhabditis elegans stress response.
Nature Communications. 2018;9. doi:10.1038/s41467-018-07640-w

75. Watts J, Ristow M. Lipid and Carbohydrate Metabolism in Caenorhabditis elegans. Genetics. 2017;207:
413–446. doi:10.1534/genetics.117.300106

76. Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. NCBI BLAST: a better web
interface. Nucleic Acids Res. 2008;36: W5–W9. doi:10.1093/nar/gkn201