Chemical diversity promotes ecosystem function

Jeremy A. Fonvielle[1,2*], Sarah R. Sandor[1], Thorsten Dittmar[3,4], Andrew J. Tanentzap[1,3,5*]

[1]Ecosystems and Global Change Group, Department of Plant Sciences, University of
Cambridge, Cambridge CB2 3EA, United Kingdom

[2]Department of Biochemistry, Hopkins Building, University of Cambridge, Tennis Court
Road, Cambridge, CB2 1QW, United Kingdom

[3]Institute for Chemistry and Biology of the Marine Environment, Carl von Ossietzky
University, Oldenburg 26129, Germany

[4]Helmholtz Institute for Functional Marine Biodiversity, Carl von Ossietzky University,
Oldenburg 26129, Germany

[5]Ecosystems and Global Change Group, School of the Environment, Trent University,
Peterborough, Ontario K9L 0G2, Canada

*Authors for correspondence: jaf91@cam.ac.uk; atanentzap@trentu.ca

Abstract

Nature benefits people in many ways that rely on how microorganisms and abiotic processes interact with organic matter to influence ecosystem function (EF). Despite many studies evaluating how biodiversity affects EF, most variation in EF remains unexplained. Here, we tested how the variety of organic compounds in the environment, termed chemodiversity, influences EF. Using a laboratory experiment and 101-lake survey spanning Europe, we discovered that chemodiversity, not biodiversity, consistently explained variation in multiple functions related to carbon cycling and ecosystem metabolism. We attributed positive chemodiversity-ecosystem multifunctionality relationships to the accumulation of compounds that could be biochemically transformed in more diverse ways at higher chemodiversity. We estimated chemodiversity will decline by 2100 due to changes in organic matter sources, causing European lakes to emit, on average, 1.6-times more carbon. These findings reframe our understanding of ecosystem functioning and suggest chemodiversity is key to sustain the benefits that nature delivers to people.

Introduction

Nature benefits society such as by regulating the climate and providing food and clean water[1,2]. These services are in part delivered by the multiple functions undertaken by microorganisms that break down and transform different organic substrates[3]. For example, as microbes assimilate detritus and increase in biomass[4], they support the growth of higher trophic levels[5], as well as modify organic compounds into forms that sequester carbon longer in the environment and thereby contribute to climate regulation[6]. Consequently, ecosystem function is enhanced by both a greater diversity of metabolic capabilities in microbial communities[3] and the environmental conditions that favour this biodiversity[7]. However, large amounts of variation in the capacity of ecosystems to provide multiple functions simultaneously, hereafter termed ecosystem multifunctionality[1], remain unexplained. Many large-scale studies have found that biodiversity only explains between 2 to 20% of the variation in ecosystem multifunctionality[3,8,9], though there are some exceptions[10].

The organic substrates that microbes use to influence biological, geochemical, and physical processes within ecosystems are an overlooked source of variation in past studies that predict ecosystem multifunctionality[6,11,12]. Natural organic matter in aquatic and terrestrial ecosystems contains tens of thousands (and likely millions) of unique molecules of varying origin and composition, making it one of the most complex mixtures on Earth[13]. This variety of molecules within organic matter and their characteristics, such as their bioavailability and stoichiometry, are together referred to as chemodiversity[14]. Chemodiversity can be described by measuring the diversity and abundance of molecular formulae and their characteristics, similar to approaches for studying biological mixtures of species[15]. Despite many surveys of chemodiversity in waters and soils[16,17], there has been no attempt to ask how chemodiversity

influences ecosystem function in nature as compared with biodiversity and other

environmental variables.

We hypothesised that chemodiversity influences ecosystem multifunctionality in two main

ways. First, the chemical properties of organic compounds (e.g. structure, elemental

composition, and biological and photochemical reactivity) determine the biotic and abiotic

pathways by which they are degraded and transformed[18,19]. A greater number of compounds

can therefore directly promote ecosystem multifunctionality by increasing the number of

ways that microbial communities and abiotic processes transform organic matter[13]. For

example, adding compounds to a mixture can increase the variety of unique chemical

properties and thus the number of organic compounds that are degraded, subsequently

increasing microbial biomass. Second, chemodiversity can indirectly influence ecosystem

multifunctionality by changing the physical environment. For example, because compounds

vary in their ability to absorb light[20], the addition of light-absorbing compounds to aquatic

ecosystems can reduce photosynthetic rates and limit heat penetration into deeper waters[21,22].

These changes can result in warmer and more nutrient-rich surface waters that accelerate

microbial activity and carbon cycling between the land surface and atmosphere[6,23]. For these

reasons, the number and characteristics of organic compounds in the environment can

provide an integrative measure across multiple ecosystem functions whose importance can be

compared directly with other variables traditionally used to explain ecosystem

multifunctionality, including biodiversity[9] and climate[3,8]. However, whether chemodiversity

explains a relatively large amount of variation in ecosystem multifunctionality remains

untested.

Here we tested the hypothesis that chemodiversity promotes ecosystem multifunctionality by

combining a manipulative lab experiment with the largest field survey of these variables to

date. We focused on lakes unlike past studies that tested associations between biodiversity

and ecosystem multifunctionality in terrestrial ecosystems[8]. Despite covering only 3.7% of

the Earth's surface, the functions of lakes, including carbon cycling and food web

production[23,24], are disproportionately important to many services valued by society, such as

climate regulation and food provisioning. We first used a lab experiment to test how

changing chemodiversity changes ecosystem multifunctionality, and then tested the

importance of this association in natural lakes that span 30° of latitude as compared with

known drivers of ecosystem multifunctionality, namely biodiversity and physicochemical

variables. We defined multifunctionality from a human perspective that values services like

climate regulation and food web productivity, and measured chemodiversity in surface waters

using ultrahigh-resolution Fourier-transform ion cyclotron resonance mass spectrometry (FT-

ICR MS) to identify the diversity of compounds involved in biotic and abiotic reactions.

Although the analytical window of FT-ICR MS can omit low (<150 m/z) and high-

molecular-mass (>1,000 m/z) compounds that contribute to biotic and abiotic reactions[25], it is

the most sensitive tool available for characterising chemodiversity and captures most of the

organic compounds that interact with microbes[26]. We then used the field survey and current

theories that synthesise how different types of chemodiversity originate[13,27] to develop a

general understanding of how and where chemodiversity matters for ecosystem

multifunctionality. As lakes are more abundant at northern latitudes[28] that are undergoing

large changes in organic matter inputs, we use our results to forecast the consequences of

environmental change for chemodiversity and the potential impacts on the benefits that lakes

provide to society.

Chemodiversity is the main driver of ecosystem multifunctionality

To test if ecosystem multifunctionality increases with increasing chemodiversity, we first

exposed four levels of biodiversity (1, 2, 3, and 6 bacterial species) under controlled

laboratory conditions to three environmental sources of dissolved organic matter (DOM). The

three DOM sources naturally varied in chemodiversity (1090, 1745, and 2564 formulae),

which closely correlated with indices summarising potential biogeochemical functions, such

as defined from the bioavailability, reactivity, and molecular mass of DOM[29] (Fig. S1). The

molecular composition of the three DOM sources was also largely nested (Fig. S2), whereby

increasing chemodiversity primarily consisted of the addition of aromatic-like formulae

(Table S1). Our fully factorial design allowed us to compare the importance of

chemodiversity versus biodiversity for predicting ecosystem multifunctionality from the main

effects of chemodiversity and biodiversity and their interaction. Ecosystem multifunctionality

was defined using changes in cell abundances, dissolved organic carbon (DOC)

concentrations, and total dissolved nitrogen (TDN) concentrations over a 7-day period. We

chose these functions because they are important measures of food web productivity and

nutrient cycling[30]. Because high microbial activity can increase both DOM and TDN

concentration by releasing exudates or transforming particulate organic matter into dissolved

compounds, as well as reduce DOM and TDN concentration by assimilating and respiring

organic compounds[31], we considered the absolute change in DOC and TDN concentration as

an indicator of microbial activity. We found that these three functions were largely

orthogonal to each other in our closed system, thereby representing complementary axes of

ecosystem multifunctionality (Fig. S3). We then summed standardized values (z-scores) of

the change in each of the three functions between day 0 and day 7 into a single index of

ecosystem multifunctionality as is common in ecology[1,3]. Larger values of this index reflect

higher ecosystem multifunctionality. We tested the main and interactive effects of

chemodiversity and biodiversity as factors on multifunctionality using a linear model without

any other variables, i.e. only accounting for the experimental design (see Methods).

We found strong experimental support for our hypothesis that chemodiversity enhances

ecosystem multifunctionality. Increasing chemodiversity led to higher values of both the

multifunctionality index (Fig. 1b) and each individual ecosystem function, with one

exception (Fig. S4). Over the range of observed chemodiversity (1090 to 2563 formulae),

multifunctionality was estimated to increase from a mean (95% confidence interval [CI]) of -

0.90 (-1.49 – -0.32) to a mean of 1.44 (0.10 – 2.77) (Fig. 1a). This mean increase

corresponded to an absolute change of 260% (107 – 966%). In support of these findings,

DOM was increasingly transformed with increasing chemodiversity. We specifically found

ten-times as many compounds were consumed at the highest compared to the lowest level of

chemodiversity, whereas the percentage of compounds that were produced in the incubations,

such as might arise from cell lysis, was relatively stable across chemodiversity treatments

(Fig. S5). Importantly, we found no evidence that chemodiversity was an outcome of greater

multifunctionality. The change in chemodiversity, that is, the difference between

chemodiversity measured at the end of the incubation in control incubations with no microbes

and chemodiversity measured at the end of the incubation in the presence of microbes, was

not correlated with multifunctionality (ρ=-0.22, p=0.203). Chemodiversity alone explained

24% of the variation in multifunctionality, which is greater than the effect of biodiversity in

many other studies of ecosystem multifunctionality[3,8–10].

Neither ecosystem multifunctionality (Fig. 1b) nor any individual function increased

consistently with biodiversity in our experiment (Fig. S4). Although the two species

community had higher multifunctionality, its taxa were included in the six species

community (Fig. 1b). Thus, if increasing biodiversity increased ecosystem function, we

would expect the six species community to have an equal or higher multifunctionality than

the two species community. However, we observed lower multifunctionality with six species

than with two species (Fig. 1b). Adding biodiversity can sometimes have no added benefit for

ecosystem function, potentially because of competition and redundancy in the functions

undertaken by different species[32,33]. In our experiment, we found that the two species

community grew faster and caused larger absolute change in DOC and TDN than when its

members were grown alongside other taxa in the six species community (Fig. S4), suggesting

competitive inhibition in the latter. Most microbial taxa negatively interact when grown

together[34], which reduces their growth relative to expectations from when taxa are grown by

themselves[35].

We also observed an interaction between chemodiversity and biodiversity that further

emphasizes the importance of chemodiversity for ecosystem multifunctionality. Higher

biodiversity increased ecosystem multifunctionality only at the highest chemodiversity level

(Fig. S6). These results suggest that without a sufficient diversity of compounds for microbes

to use, increasing biodiversity alone cannot sustain a higher ecosystem multifunctionality. By

contrast, in environments with low biodiversity and low chemodiversity, increasing

chemodiversity alone can increase ecosystem multifunctionality by enabling the species

already present to perform more functions. Because ecosystem multifunctionality-

biodiversity relationships[3,8–10,32,36–38] are mostly studied in terrestrial environments, where

chemodiversity is far higher than in aquatic environments[39], the importance of

chemodiversity may have been overlooked.

A potential criticism of our experiment is that because we observed an increase in absolute

carbon and nitrogen concentration, changes in ecosystem multifunctionality may have been

due to cell lysis rather than microbes transforming chemodiversity. However, both carbon

and nitrogen could have come from particulate organic matter as well as nitrogen fixation by

some of our study species[40,41]. In support of the former explanation, between 1.6 to 10.6% of

the total organic carbon in each treatment was in particulate rather than dissolved form. We

also found evidence that nitrogen was incorporated into DOM from other pools than cell

lysis. Specifically, we found that a higher proportion of N-containing compounds were

produced at higher chemodiversity, and this result could not have been due simply to cell

lysis because there was a higher overall consumption of compounds (Fig. S5). Taken

together, our results suggest that increasing chemodiversity increases ecosystem

multifunctionality by enabling microbes to use and transform substrates in more ways than is

achieved by increasing biodiversity alone.

To test the generality of our lab experiment, we sampled surface waters from 101 natural

lakes spanning continental Europe (~30° latitude). As in the lab experiment, the proportion of

aromatic compounds increased from 3.1% to 28.4% over the range of chemodiversity

observed in the field from 396 to 3387 molecular formulae (linear model, t=6.54, df=100,

p<0.001). The chemical characteristics of DOM, including its molecular weight, further

suggested that aromatic-like compounds were increasing with chemodiversity like in the lab

experiment (Fig. S7). We also measured different functions in the field than in our

experiment to test if chemodiversity promoted ecosystem multifunctionality irrespective of

the specific functions being measured, that is, if the association between chemodiversity and

ecosystem multifunctionality was generalizable. The functions measured in the field were

related to carbon cycling (CH_4 and CO_2 emissions) and ecosystem metabolism (total

respiration, maximum net primary production, and bacterial protein production). We reversed

the sign of the measured values for CH_4 emissions, CO_2 emissions, and total respiration so

that a higher multifunctionality value corresponded to a greater benefit to society[37] (see

Methods). Thus, larger values of these three functions indicated lakes were both emitting and

mineralizing less carbon. We then summed z-scores of the five functions into a single index

as in the lab experiment. We also compared the importance of chemodiversity with the

importance of microbial (bacteria, fungi, archaea) and eukaryotic biodiversity estimated from

shotgun metagenomic sequencing of environmental DNA in 82 of the study lakes. Our

biodiversity sampling detected more taxa than in other large-scale studies[42,43] with between

97 to 1589 species (median = 610) per lake.

We found a positive association between ecosystem multifunctionality and chemodiversity

but not with biodiversity across 82 of the 101 field sites with data for all these three variables.

Over the range of chemodiversity (396 to 3387 molecular formulae, median = 2445),

multifunctionality was estimated to increase from a mean of -1.27 (95% CI: -2.16 – -0.37) to

0.30 (-0.33 – 0.92) (linear model: t=2.24, p=0.030, df=81; Fig. 1c). This mean increase

corresponded to a change of 124% (43 – 144%). Neither biodiversity nor its interaction with

chemodiversity influenced multifunctionality (t=1.26, p=0.21 and t=-0.02, p=0.99,

respectively, df=81 for both). Much like in our experiment, the effect of biodiversity on

multifunctionality was limited by chemodiversity. We found that the effect of biodiversity on

multifunctionality only became statistically significant at the highest levels of chemodiversity

(Fig. S8). Three of the five measured functions, CH_4 and CO_2 emissions and bacterial protein

production (BPP), also varied individually with chemodiversity whereas only CO_2 emissions

significantly decreased with biodiversity (Fig. S9). At a molecular-level, these functions were

associated with different formulae, potentially reflecting differences in how microbial

functions consumed organic substrates. For example, BPP was negatively correlated with

generally low H:C formulae (Fig. S10), indicative of consumption[4], because these formulae

were relatively more abundant and thus easier for bacteria to find to build cellular structures

than rarer high H:C compounds that may release more energy[4,44]. CH_4 was positively

associated with the proportion of formulae per lake that contained either N or S (ρ=0.24,

p=0.006), which are precursors for oxic methanogenesis[45,46], and CO_2 negatively correlated

with high H:C formulae (Fig. S10), which are preferably respired by bacteria because they

provide the highest energy return[4,44]. Chemodiversity alone explained 5% of the variation in

multifunctionality, which is lower than in our experiment but we considered more functions

than in our experiment and our experiment controlled for environmental noise, e.g. climate.

This result is also comparable to, and in some occasions up to five-times higher than, the

effect of biodiversity reported in influential field studies explaining variation in ecosystem

multifunctionality[8,36]. The choice of functions affects the strength of the relationship between

biodiversity and multifunctionality, so may explain some of these differences among studies.

To test further the importance of chemodiversity relative to published estimates of

biodiversity, we computed multifunctionality based on the exceedance of data-derived

thresholds[37]. This method was specifically designed to facilitate inter-study comparisons[37].

We found that the reduced explanatory power of biodiversity relative to chemodiversity

persisted when we estimated ecosystem multifunctionality with this threshold approach

(Supplementary Note 1). Our estimated effect of chemodiversity on multifunctionality with

this threshold approach was larger than the effect of biodiversity on ecosystem function that

was previously estimated from 47 published studies[9], including freshwaters (Supplementary

Note 1). Unlike for biodiversity-ecosystem multifunctionality, where positive associations

have been reported in studies measuring plant productivity[3,8], and rarely in studies measuring

macronutrient cycling[33], the chemodiversity-ecosystem multifunctionality relationship may

be less sensitive to the specific functions measured. We found consistent positive associations

between chemodiversity and ecosystem multifunctionality across our experimental and field

datasets with fundamentally different functions associated with primary productivity and

macronutrient cycling. Together, our results from both controlled and natural conditions

suggest that the large role of chemodiversity in supporting ecosystem multifunctionality has,

to date, been overlooked and must be considered in efforts aimed at delivering benefits from

nature to society.

Recalcitrance underlies positive chemodiversity-multifunctionality associations

Based on theoretical models that predict that the reactivity of DOM varies with its molecular

composition and local environmental conditions[13,27], we hypothesised that the effect of

chemodiversity on ecosystem multifunctionality depends on the source of chemodiversity

itself. By sampling lakes at similar periods of annual heat accumulation, and with a similar

light environment and DOC concentration (see Methods), we found that spatial variation in

chemodiversity at a continental scale primarily reflected differences in climate and

surrounding land use[16,47] (Fig. S11). Specifically, chemodiversity strongly increased from

southern to northern Europe ($\rho=0.40$, $p<0.001$), suggesting that the environment in which

chemodiversity occurred and/or its source could explain why chemodiversity varied

throughout Europe. Here we extend these two potential explanations (see Supplementary

Note 2 for further discussion) to contend that they differentially influence how

chemodiversity is used by microbes and thus ecosystem multifunctionality. If chemodiversity

was high because of environmental conditions ("environment" hypothesis in Fig. 2a) that

slow microbial transformation of DOM, such as lower mean annual temperature (MAT) or

lower macronutrient concentrations, we expected that ecosystem functions like microbial

respiration and bacterial protein production would also be reduced[48], resulting in a negative

correlation between chemodiversity and multifunctionality. By contrast, if chemodiversity

primarily arose because compounds with greater chemical complexity were imported into

lakes, we expected these compounds to support a greater number of biochemical

transformations and increase ecosystem multifunctionality ("source" hypothesis in Fig. 2a).

Complex compounds, such as those with high aromaticity, are abundant in lakes and their

degradation by microbes involves many biochemical pathways that increase metabolic

activity and biomass formation[4,49,50], and ultimately multifunctionality. By contrast, simpler

compounds are expected to decrease multifunctionality because they are often used to

produce energy, which emits CO_2 as a by-product.

We tested how both the complexity of DOM and the environment in which DOM was

degraded affected microbial transformation of DOM and subsequently the effect of

chemodiversity on ecosystem multifunctionality. We estimated DOM complexity in each

lake based on an aromaticity index[51] (see Methods). We expected this index to reflect

terrestrial vegetation[16], which is a main source of lake DOM[40]. Slower growing vegetation

tends to invest more in protecting biomass, such as with secondary metabolites[52], because it

faces a relatively greater cost to produce biomass. Compounds involved in plant protection

tend to be more complex and have chemical properties, including a higher aromaticity, which

require more biochemical reactions to be transformed[14,27]. We expected complexity to

influence multifunctionality by determining the extent to microbes can transform DOM and

we expected complexity itself to be determined by terrestrial net primary productivity (NPP),

i.e., vegetation growth rates, in surrounding catchments. We also considered nine

environmental variables (MAT, annual precipitation, water temperature, pH, water colour,

lake area, biodiversity, trophic state, and nutrient concentrations) that could influence

chemodiversity by changing the diversity of terrestrial and aquatic organic matter inputs,

controlling the physical energy required to break chemical bonds, and regulating photo-

oxidation[53,54,55]. NPP and the environmental variables were estimated from high-resolution

remote sensing data and in-situ measurements (see Methods). We then averaged the number

of times molecular formulae could be transformed in each lake using a list of 1255 putative

biochemical transformations[19] (see Methods). If the positive association we observed

between chemodiversity and ecosystem multifunctionality arose from the input of complex

compounds rather than the limitation of microbial transformations of DOM, we expected

NPP would be more strongly associated with chemodiversity than the nine other

environmental variables.

Our results indicate that the source of DOM explains how and where a positive

chemodiversity-ecosystem multifunctionality relationship can arise. Specifically, we found

that the production of complex compounds by low terrestrial NPP (ρ=-0.70, p<0.001, Fig.

S12a) correlated with both greater chemodiversity (ρ=0.56, p<0.001; Fig. S12b) and a greater

number of average transformations that formulae could undergo in each lake (ρ=0.45,

p<0.001, Fig. S12c). Consequently, chemodiversity correlated positively with the number of

transformations per formulae (ρ=0.40, p<0.001, Fig. 2b). Correlations between

chemodiversity and each of complexity, NPP, and the number of biochemical transformations

are predicted by the source hypothesis. By contrast, all the environmental variables had

weaker associations with chemodiversity, though some were statistically significant (Table

S2). Together, our results indicate that the production of complex compounds by low NPP

increases both chemodiversity and the number of biochemical transformations that each

compound can support.

We synthesised the observed correlations among chemical complexity, biochemical

transformations, and chemodiversity to propose a new general understanding of ecosystem

multifunctionality. This new understanding was that the positive effect of chemodiversity on

multifunctionality arises because of the production of complex compounds by plants and

environmental conditions that favour those compounds within DOM pools. We found further

support for this framework with a path analysis explicitly testing presumed causal pathways.

The path analysis modified our original hypotheses to incorporate additional paths identified

by tests of model structure (Fig. S13). With this analysis, we found that effects related to the

production of organic compounds (i.e. NPP) influenced chemodiversity by changing DOM

complexity and subsequently the number of biochemical transformations it can receive (Fig.

2a). Although environmental variables (i.e., water colour and MAT) also influenced DOM

complexity, the direction of these effects suggested that they were associated with reduced in-

lake primary productivity and an increase in the relative importance of terrestrial DOM (Fig.

S13). This interpretation was further supported by the direct negative effect of chlorophyll *a*

concentrations on chemodiversity (Fig. S13).

To test further if ecosystem multifunctionality was enhanced by chemodiversity because of

the properties of complex compounds associated with low terrestrial NPP, we analysed the

composition and putative source of all 8456 molecular formulae identified in the field survey.

We first identified formulae whose relative abundances across our study lakes were

correlated with terrestrial NPP of the surrounding catchment. Positive and negative

correlations suggest formulae likely originated from high and low terrestrial NPP,

respectively. We compared these correlations with the average number of biochemical

transformations that each formula was predicted to undergo in each lake[19]. This analysis

allowed us to infer the origins of compounds that can be highly transformed by biotic and

abiotic reactions, which contribute to higher ecosystem multifunctionality. Compounds that

were associated with low NPP appeared more transformed (t=-8.19, df=6639, p<0.001), and

positively correlated with chemodiversity (t=8.50, df=6639, p<0.001; warmer colours in Fig.

2c). These compounds were more complex, on average, with a mean (± standard error)

aromaticity index of 0.52 (±0.17) compared to 0.32 (±0.24) in the entire dataset (Wilcoxon

rank sum test, W=5.4×10^6, p<0.001, Fig. S14). These results further supported the

hypothesis that complex compounds can be degraded in more ways and support higher

multifunctionality. We then selected molecular formulae that most increased in abundance

with decreasing NPP, that is, were in the top 10% of negative associations based on

Spearman rank correlations. We found that these formulae could support, on average, 1.4

times more known putative biochemical transformations than formulae that were weakly

associated with NPP, i.e. lowest 10% of negative associations (t=5.80, df=860, p<0.001), and

1.8 times more transformations than formulae strongly positively associated with NPP, i.e.

top 10% of positive correlations (t=8.00, df=1090, p<0.001). Together, these results

demonstrate that the effect of chemodiversity on multifunctionality emerges from the

properties of individual formulae.

We expect chemodiversity will be less important for ecosystem multifunctionality in

environments where microbes cannot access and transform large quantities of organic matter.

For example, DOM concentrations are typically >5 mg C L^{-1} in rivers, most lakes, wetlands,

or mangroves[19,47,48,56], but <1 mg C L^{-1} in the deep ocean[29] or in large (surface area >500

km^2) deep lakes with centennial residence times[57]. At these concentrations, individual

compounds are more likely to be too dilute to interact with microbes[52], irrespective of the

diversity of compounds in the environment, and may result in a shallower slope between

chemodiversity and ecosystem function. In support of this expectation, we found that

compounds transformed in lakes reacted less frequently when found in the deep ocean. We

identified 2745 formulae shared between our study lakes and a representative deep ocean

sample[18], which is likely an overestimate because we cannot differentiate structural isomers.

We then calculated the number of times that each formula was transformed in each sample

using the established list of putative biochemical transformations[19]. We found that these

formulae underwent, on average, 0.08 (range: 0 to 1.32, standard deviation = 0.17) putative

transformations in lakes but no transformations were detected in the deep ocean. Given that

many of the detected transformations are ubiquitous, such as methylation, a likely

explanation for why we did not find them in the ocean sample is because their substrates

occur at concentrations lower than our femtomole detection limit[58]. Such concentrations

would be several order of magnitude beneath concentrations of the most limiting organic

compounds[59]. This observation further suggests that chemodiversity will have a greater

influence on ecosystem multifunctionality where individual compounds occur at high enough

concentrations to support many biochemical transformations.

Chemodiversity and ecosystem multifunctionality under global change scenarios

NPP is expected to increase because of rising atmospheric CO_2 concentrations[60], and our

results suggest that this increase could reduce the chemodiversity of lakes in ways that are

relevant to global carbon budgets and climate modelling efforts. To predict how the resulting

changes in chemodiversity will influence ecosystem multifunctionality, we first modelled the

value of each ecosystem function that varied with chemodiversity in our field survey (Fig.

S9). By adding chemodiversity to statistical models of CH_4 and CO_2 emissions and BPP that

contained predictors identified as important in previous large-scale studies[23,24], we increased

explained variation of each function by up to 50% as compared with past studies (Table S4).

We achieved the large increase in explained variation despite being unable to measure some

prominent drivers of ecosystem metabolism, such as phosphorus concentrations[61].

Importantly, based on the properties of linear models, additional predictors are unlikely to

change the effect of chemodiversity on individual functions and will only further reduce

unexplained variation. We then projected our models for each ecosystem function to all

European lakes >10 ha (see Methods). For chemodiversity, we generated a European-scale

raster with the model used in the path analysis (Fig. S13). This analysis further validated our

study because we found that the extrapolated estimates were within bounds of global-scale

studies[23,24]; no European-only studies exist for comparison. Overall, we estimated that

European lakes emitted, on average, 22.7 Tg C yr^{-1} as CH_4 (95% CI 11.2 – 32.0; model R^2 =

0.46) and 27.8 Tg C yr^{-1} as CO_2 (95% CI: 9.9 – 37.4; R^2 = 0.22) in 2019. Despite these

emissions potentially being overestimated because they only considered summertime

fluxes[62], they corresponded to 7% and 19% of estimated global lake CH_4 and CO_2

emissions[23,24], respectively, and Europe contains ~20% of the world's total number of lakes[63].

We also estimated that European lakes fixed an average of 1.8 Tg C yr^{-1} (95% CI: 1.0 – 3.3;

R^2 = 0.45) in 2019 into microbial biomass, that is, from BPP.

To forecast the consequences of changes in NPP for European lakes, we input NPP estimated

in the year 2100 by Earth system models[64] into the linear model that predicted

chemodiversity (Table S3) and then used the predictions to forecast each ecosystem function

(Table S4). We focused solely on changes in NPP, holding other variables constant, because

NPP was the best individual predictor of chemodiversity (Table S2) and had a clear

mechanistic link to chemodiversity (Fig. S13). We used two shared socioeconomic pathways

(SSPs) representing likely (SSP2) and worst case (SSP5) scenarios for future changes in

atmospheric CO_2 concentrations. We found that future increases in NPP alone will reduce

chemodiversity in 99.1% of lakes, with the median chemodiversity decreasing by 13% and

24% under the SSP2 and SSP5 scenarios, respectively (Fig. 3a). When summed across lakes,

we estimated that the change in chemodiversity will cause lakes to emit an additional mean

28.4 (95% CI: 3.3 – 166.1) and 32.2 (95% CI: 14.0 – 231.2) Tg C yr^{-1} as CH_4 under SSP2

and SSP5 scenarios, respectively (Figs 3b, S15). Similarly, the changes in chemodiversity

will cause lakes to emit an additional 3.7 (95% CI: 1.3 – 5.1) and 4.0 Tg C yr^{-1} as CO_2 (95%

CI: 1.3 – 5.2 Tg C yr^{-1}) under SSP2 and SSP5 scenarios, respectively (Figs 3c, S15).

Microbes will, however, fix 4.3 (95% CI: 1.7 – 10.2) and 4.4 Tg C yr^{-1} (95% CI: 1.7 – 12.6

Tg C yr^{-1}) more carbon into biomass under SSP2 and SSP5 scenarios, respectively (Figs 3d,

S15). Together, our estimates suggest that European lakes may emit, on average, 1.6-times

more carbon by 2100, mostly as CH_4, because of NPP-associated declines in chemodiversity.

Although our estimates are higher than previous estimates of future CH_4 emissions by lakes

in absolute terms by 50%, they are comparable in terms of relative change, i.e. 73-84% in our

study versus 58-88%[62]. The difference in absolute terms is likely due to the lack of winter

measurements in our dataset as lower microbial activity in colder weather typically reduces

greenhouse gas emissions[62]. The similarity in relative terms instead emphasises that

chemodiversity is a necessary predictor of CH_4 emissions.

We forecasted that lakes in southern Europe, and in the south of the Scandinavian peninsula,

will experience the largest changes in ecosystem function (Fig. 3). These lakes are located

near large population centres and, therefore, provide opportunities to improve management of

their surrounding catchments in ways that can enhance the benefits that European lakes

provide to society. For example, our results suggest that chemodiversity and subsequently

ecosystem multifunctionality can be enhanced by promoting slow growing vegetation, such

as many Fabaceae[65]. Slow growing vegetation produce large numbers of complex polymeric

compounds, which undergo many biochemical transformations when leached into water from

litterfall and soils[17]. They also have higher wood density[66], that can increase long-term

carbon storage on land[67], resulting in a win-win for land and water.

Conclusions

Here, we provided experimental and large-scale observational evidence that increasing

chemodiversity delivers beneficial ecosystem functions, especially a stronger carbon sink

capacity corresponding with greater bacterial biomass and lower greenhouse gas emissions.

We then demonstrated that the ways in which ecosystems deliver these functions depends on

the molecular properties of DOM, namely the capacity of individual compounds to undergo

biochemical transformation. The molecular properties of compounds ultimately depend on

sources of DOM and external environmental conditions, such as their concentrations, as

found in many inland waters, wetlands, and mangroves[19,47,48,56]. Here we add to this

understanding by showing the importance of these properties for ecosystem

multifunctionality and their potential to explain ecosystem-specific differences in

chemodiversity–multifunctionality relationships. Our findings also suggest that

chemodiversity and its effect on multifunctionality can be altered across different aquatic

ecosystems by manipulating sources of DOM within surrounding catchments. Finally, our

study suggests that accounting for the strong association between NPP and chemodiversity

can improve integration between terrestrial and aquatic ecosystem models[5]. For example,

because chemodiversity is strongly predicted by NPP, which itself is predictable at high

temporal and spatial scale from bioclimatic data, we can readily incorporate the effect of

chemodiversity on greenhouse gas emissions into existing carbon budgets. More generally,

our study demonstrates that sustaining the benefits that lakes provide to people will require

management that extends beyond shorelines and considers the chemodiversity exported from

land.

Acknowledgments

We are grateful for help with field work from Samuel Cottingham, Carolyn Ewins, Sophie Guillaume, Eleanor Sheridan, and Samuel Woodman. We thank Yi Zhang and Caroline Kemp for laboratory assistance, Del Hawtin, and Simon Hoer for logistical support, and Katrin Klaproth for assistance with FT-ICR MS measurements. We thank Jim McGeer for providing the DOM sources and Thomas Scheuerl for providing the bacterial strains used in the experiment. This work was funded by H2020 ERC Grant sEEIngDOM #804673 to AJT and the University of Cambridge Harding Distinguished Postgraduate Scholars Programme.

Author contributions

AJT and JAF designed the study. JAF performed the field and associated lab work with input from AJT and TD. SRS and JAF performed the laboratory experiment. JAF analysed the data with input from AJT. JAF and AJT wrote the manuscript with input from TD and SRS.

Data availability

All data are available within the manuscript or upon reasonable request.

Figure 1: Chemodiversity increases ecosystem function. Multifunctionality increased with **(a)** higher chemodiversity but not with **(b)** higher biodiversity under controlled conditions. Multifunctionality was measured as the sum of *z*-scores of changes in each of cell abundances, and transformation of dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) over 7 days in 36 microcosms represented by individual points. Boxes encompass the inter-quartile range with median denoted by solid horizontal line and whiskers equal to 1.5-times the interquartile range. Different letters indicate statistically significant ($p<0.05$) differences between factor levels with Tukey's pairwise comparison tests. Chemodiversity of each experimental source was estimated from the exponential of the Shannon diversity index. In a separate field study, multifunctionality increased with **(c)** higher chemodiversity but not with **(d)** higher biodiversity across 82 European lakes represented by individual points. Multifunctionality was measured in the field study as the sum of *z*-scores of CH_4 and CO_2 emissions, total respiration (Resp), maximal net primary production (PP), and bacteria protein production (BPP). Lines in **(c)** and **(d)** are estimated mean associations based on a linear model ± 95% confidence intervals. For all, darker colours indicate higher chemodiversity (brown points) and biodiversity (blue points).

Figure 2: The origin of chemodiversity influences its effect on ecosystem multifunctionality. (a) We hypothesized that the positive effect of chemodiversity on ecosystem multifunctionality was caused by environmental conditions reducing the transformation of DOM ("Environment" hypothesis) and sources of DOM producing compounds that could be transformed in a greater number of ways ("Source" hypothesis). + and – symbols show hypothesised cause-and-effect of one variable on another, with either positive or negative links between NPP and complexity possible in the Source hypothesis. We tested these hypotheses using path analysis fitted to 101 lake-level observations. To simplify the path analysis, we created a composite variable "environment" that encompassed 9 measured parameters (see Fig. S13 for effect of each parameter), whereas "source" was equal to NPP in the surrounding catchment of each lake. Complexity and transformations corresponded with the intensity-weighted modified aromaticity index and mean number of putative transformations per molecular formula per lake, respectively. Coloured arrows show statistically significant paths with standardized effects. **(b)** The mean number of transformations per formula per lake was closely correlated with chemodiversity (ρ=0.40, p<0.001). Each dot represents a lake (N=101) coloured for chemodiversity. **(c)** We also correlated the number of transformations that each of 8456 molecular formulae underwent across all lakes with the NPP of the lake catchment. Formula (points) associated with low NPP (i.e., negative value along the *y*-axis) generally underwent more transformations (i.e. larger value along the *x*-axis; ρ=-0.10, p<0.001). Numbers of transformations were weighted by the relative abundance of each formula across lakes. Points are coloured according to their correlation with chemodiversity with low (blue) and high (red).

Figure 3: Lakes will emit more carbon by 2100 due to continental-wide declines in chemodiversity. We estimated **(a)** chemodiversity in 2100 while holding all other variables at 2019 levels for all European lakes >10 ha in surface area (R^2 for chemodiversity model = 0.36). Values are the change in chemodiversity compared to 2019. Using future chemodiversity, we then estimated mean changes in **(b)** CO_2 emissions **(c)** bacterial biomass production, and **(d)** CH_4 emissions from pelagic surface waters between 2019 and 2100 using linear models (R^2 = 0.22, 0.45, and 0.46, respectively; Table S4). For all, points indicate N=175,500 individual lakes. Histograms indicate the number of lakes found with corresponding changes. 95% confidence intervals for each function are given in Fig. S15.

Materials and methods

Laboratory experiment

To test if multifunctionality varies with chemodiversity, we incubated bacterial communities of four biodiversity levels with three different levels of chemodiversity. The four biodiversity levels were 1) *Pseudomonas fluorescens* SBW25 2) *Chryseobacterium* sp. and *Sphingobacterium* sp. isolated from natural communities that degrade DOM[68]; 3) *Rahnella* sp., *Raoultella* sp., and *Pantoea* sp. from the same natural communities[68]; 4) all six microbes together. Before starting the experiment, we grew each microbial isolate individually from a frozen archive for five days in R2A media. A control of filtered (0.2 μm, Sterivex-GP, MilliporeSigma, USA) sterile water was included alongside each treatment and indicated no contamination as measured with flow cytometry (see below). For chemodiversity, the three treatment levels were obtained from reverse-osmosis-concentrated water[69] of a temperate wetland (Luther Marsh, Ontario, Canada; 43°57'43"N, 80°23'50"W), boreal lake (Clearwater Lake, Ontario, Canada; 46°22'11"N, 81°03'04"W), and Arctic river (Cameron River, Northwest Territories, Canada; 62°29'33"N, 113°32' 48"W). Using MilliQ water, we diluted each reverse-osmosis concentrate to a final dissolved organic carbon (DOC) concentration of 10 mg C L^{-1}, as measured with a TOC-TNM-L analyser (Shimadzu, Japan), and filled fifteen 250 mL pre-combusted (4 hr at 550°C) Erlenmeyer flasks with 100 mL of each water source (n = 3 replicates for each of 4 biodiversity levels with 3 controls). Before inoculating the DOM sources with each microbial community, we estimated strain-specific cell number with a CLARIOstar fluorometer (BMG Labtech, UK) after 30 min of staining with SYBR Green (10,000× stock solution, Invitrogen, USA) diluted to a 10× final concentration. Based on these data, we diluted an unstained aliquot of each strain to an expected 10,000 fluorescence units and mixed them to form the four communities. This normalisation created a substitutive design that ensured each community had a similar abundance of DNA and thus potential

activity. Then, we added 1 mL of each of the four microbial communities to each

chemodiversity level in triplicate and incubated the flasks at ca. 20°C for seven days in the

dark without shaking.

Measuring chemodiversity

We estimated chemodiversity in the experiment from solid-phase extraction of DOM

following a standard protocol[70]. Briefly, 80 mL of each incubation flask was acidified to a pH

of 2 by adding HCl, passed through 1 g of a polar styrene-divinylbenzene polymer (Bond

Elut PPL, Agilent, USA), and eluted with 3.5 mL of ultrapure methanol (LC-MS LiChrosolv,

Merck, Germany). We diluted the extracts to a final concentration of 2.5 ppm DOC in a 1:1

(v:v) methanol:water solution before injecting 100 µL into a 15 Tesla Solarix FT-ICR MS

(Bruker Daltonic, Germany) with direct infusion electrospray ionization in negative mode.

We collected 200 scans in duplicate for each sample and calibrated the spectra using the

DataAnalysis software (Bruker Daltonic, Germany). Masses ranging from 150 to 1000 m/z

were exported, and we assigned molecular formulas using the online platform ICBM-

OCEAN[71]. We used the N, S, P rule, deleted all singletons, and used isotopic confirmation to

accept/reject a formula. We limited formula attributions to $C_{1-100}H_{2:200}O_{0-70}N_{0-3}P_{0-2}S_{0-2}$ with a

tolerance of 0.2 ppm and retrieved a total of 3873 individual molecular formulas. We

computed chemodiversity as the exponential of the Shannon diversity index using all

assigned molecular formulas[72] and the summed normalised peak intensity per sample. This

index represents the diversity in a community when all formulae are equally common, so

corrects for variation in evenness that can bias raw counts[73] of formulae number. We also

computed the ratio of hydrogen to carbon atoms in each compound as an indicator of

bioavailability, as described elsewhere[44]. The three DOM sources were largely nested with

one another with 93% of formulae present in the Arctic river (low chemodiversity) also

present in the boreal lake (medium chemodiversity), 70% of boreal lake formulae present in

the temperate wetland (high chemodiversity), and 80% of formulae present in the Arctic river

present in the temperate wetland (Fig. S2). The large number of shared formulae among

treatments confirmed that sources with higher chemodiversity contained the vast majority of

formulae present in sources with lower chemodiversity, i.e. were nested. Additionally, the

molecular composition of the DOM used in the experiment differed as much from the lakes

sampled in the field study as the lakes differed among themselves (Fig. S16).

Ecosystem multifunctionality in the laboratory experiment

We estimated the multifunctionality of each replicate community in the experiment by

measuring carbon consumption, nitrogen consumption, and bacterial growth rate. At the start

and end of the incubation, we measured the DOC and TDN concentrations of each flask using

a Shimadzu TOC-TNM-L analyser. Carbon and nitrogen consumption were estimated as the

change in concentration between the start and end of the incubation. To measure growth rate,

we determined the difference in cell abundance between the start and end of the incubation

using flow cytometry. Briefly, we removed 800 µL of water from each flask at the start and

end of the experiment. Samples were incubated with 200 µL of 25% glutaraldehyde in the

dark at 4°C for 30 min before being preserved at -80°C. Within one week, we thawed the

samples, added SYBR Green at a 10× final concentration (10,000× stock solution, Invitrogen,

USA), and incubated the samples overnight at 4°C. Then, we injected 100 µL of each sample

into a flow cytometer (Attune NxT, Thermo Fisher Scientific, USA) at a rate of 250 µL per

minute and used the same instrument settings for all samples. We optimised the gating for

each DOM source based on stained/unstained samples using FlowJo v10.9 (BD Biosciences,

USA) and used the same gating for each community. Each function was standardized to a

mean of zero and standard deviation of one to obtain a *z*-score, which we summed across the

three functions to estimate multifunctionality[8].

Selection of field study sites

We selected lakes to sample across Europe ensuring there were no systematic biases in DOC

concentrations and water chemistry across our latitudinal gradient. Using the EU's Water

Information System for Europe (WISE) database version 4[74], we subsampled all entries

labelled as "lakes" for which DOC was reported in mg C L^{-1}. We also selected entries with a

total organic carbon (TOC) concentration equal or lower than 5 mg C L^{-1} to include more

lakes with low DOC values, which were underrepresented in the WISE database despite

being known to be the most common on Earth[75]. From the 2169 resulting entries, we

removed 256 entries with a DOC/TOC concentration >30 mg C L^{-1}, which is outside the

expected values in lakes[75] and likely represent measurement error or atypical lakes, and 1595

duplicated measurements. We grouped the remaining 318 lakes into five bins in 5 mg C L^{-1}

intervals: <5 mg C L^{-1}, 5-10 mg C L^{-1}, 10-15 mg C L^{-1}, 15-20 mg C L^{-1}, >20 mg C L^{-1} and six

5° latitudinal intervals: <45°N, 45-50°N, 50-55°N, 55-60°N, 60-65°N, >65°N. The last two

DOC/TOC bins (i.e. >15 mg C L^{-1}) were absent from the first two latitudinal bins. Therefore,

we removed all 47 entries with DOC/TOC concentrations >15 mg C L^{-1}. In a global

analysis[75] of lake DOC concentrations, 93% of lakes were predicted to have DOC

concentrations <15 C L^{-1}, so this filtering minimises potential sampling bias while ensuring

we still represent the vast majority of lake DOC concentrations. To standardise the sites

further, we considered total nitrogen (TN) and total phosphorus (TP) concentrations of all

entries. Six entries lacked TP values, so we discarded all but one, which was located between

50-55°N where entries were sparse. Only 10 entries had TP concentrations higher than 0.2

mg L^{-1}, and 21 entries had a TN value higher than 1 mg L^{-1}, so we removed these to avoid

large differences in lake trophic levels.

After reducing the number of entries to lakes with similar water chemistry, entries were

selected based on their distance from one another. Therefore, we discarded 28 entries from

eastern Poland, one in Latvia, and two in Hungary as they were logistically challenging to

reach. We similarly removed 175 entries situated in the United Kingdom and western France.

We then inspected each remaining entry individually using Google Earth satellite imagery,

removing 86 sites that were dammed or visibly artificial. This process removed all entries in

the WISE database situated south of 40° N and north of 69°N. To extend the latitudinal

gradient, we manually added eight sites in southern Italy (<40°N) and seven in northern

Norway (>69°N) without knowledge about their DOC/TOC or trophic state. None of the

manually chosen lakes were dammed or visibly artificial. This selection process resulted in a

total of 109 lakes. During sampling, severe weather conditions, unexpected fences around

lakes, or equipment failure prevented us from sampling eight lakes, and one site in Poland

was substituted for a lake for which no DOC data were available. Ultimately, we sampled

101 lakes spread over 31 latitudinal degrees. There was no statistically significant correlation

between latitude and DOC concentration at the time of sampling (Spearman correlation,

$\rho=0.05$, $p=0.57$, Fig. S17).

Timing of field sampling

The order in which lakes were sampled was chosen to minimise differences in annual heat

accumulation and water temperatures across the latitudinal gradient. Sampling dates were

originally chosen to minimise differences in growing degree days (GDD) among lakes. GDD

represents the difference between a given temperature threshold at which primary production

commences and mean daily temperature[76]. Here we used a threshold of 10°C as is classically

employed[76]. To compute GDD, we used the rnoaa package[77] in R v.4.1.1[78] to retrieve all

weather stations within 50 km of the 101 sampled lakes from the US National Oceanic and

Atmospheric Administration National Climatic Data Center database. We determined the

median GDD for each day of the year based on temperature data collected from the last 10

years. For years with 366 days, the GDD on the 29th of February was added to the GDD from

the 28th of February. Cumulative GDD (cGDD) – representing the sum of GDD from the 1

January until a given date – were obtained by sequential addition for each day of the year for

each lake. The computed cGDD indicated that the growing season commenced (i.e. cGDD

>10) as early as the 1 January at the southernmost site and ended (i.e. cGDD reached its

highest values and no longer increased) as early as the 10 October at the northernmost site.

We then selected sampling dates to survey lakes during similar periods of annual heat

accumulation. To find the optimal sampling day, we calculated the variation in cGDD along

the latitudinal gradient as the ratio between the 3rd and the 1st quartiles of lake-level cGDD,

assuming we started sampling the southernmost site and moved northwards to the closest site

at a pace of one lake per day. By starting sampling on the 8 April 2019, we could minimise

the variation in cGDD across the latitudinal gradient. Overall, the observed maximal

difference between the 5° latitudinal bins was 194.4 GDD, which was much less than the 368

that was originally expected based on the historical data (Fig. S17). This value corresponded

with sampling the warmest site 10 days later than the coldest site, during which temperatures

remained above 20°C. We further measured underwater irradiance at 1 m depth in 60 lakes

across the latitudinal gradient using a HOBO light pendant (UA-002-64, Onset, USA) to test

for additional biases in irradiance. Pictures representing the latitudinal gradient are presented

in Fig. S18.

Dissolved organic matter sampling

At each lake, we collected 10 L of surface water in an acid-washed plastic carboy. We sampled from the point with maximum water column depth or the point equidistant to all shorelines when no bathymetry information was available. We expected these locations would be the best mixed in the epilimnion and most representative of the lake water quality. Directly after sampling, 1 L of lake water was passed through a pre-combusted (450°C, 4 hours) glass fibre filter (0.5 µm, Macherey-Nagel, Germany), and the filtrate dispatched into three 40 mL pre-combusted amber borosilicate vials and one 500 mL pre-combusted amber borosilicate bottle. Once on land, we acidified the samples to a pH of 2 by adding HCl. We kept the filtrates in the dark, and either on ice or at ca. 4°C for up to 15 days before shipping them to Cambridge, UK. In the laboratory, we stored the samples at 4°C for up to a week and determined the DOC in each 40 mL vial using a Shimadzu TOC-TNM-L analyser (Shimadzu, Japan). In line with regulatory standards[79], we averaged DOC concentrations from the three replicates when the standard deviation was within 10% of the mean and averaged the two closest values otherwise. We estimated chemodiversity as described for the laboratory experiment but solid-phase extracted 500 mL of water instead of 80 mL and retrieved a total of 8456 formulae. We also evaporated 10 µL of extract and redissolved it in 10 mL of Milli-Q water acidified to a pH of 2 using HCl to determine the DOC concentration in the extract and corresponding extraction efficiency (mean = 62%, SD =15%).

To determine if chemodiversity promoted multifunctionality because it could undergo more biochemical transformations, we also estimated the number of times that each molecular formula could be biochemically transformed using an established approach[19]. Fudyma *et al.*[19] compiled 1255 pairs of molecular formulae associated with commonly observed biotic and

abiotic transformations, such as enzymatic reactions that add or subtract CH_2 groups and

physical condensation or dehydration, respectively. We considered such a transformation to

have occurred if we detected each pair of formulae from a reaction in a sample.

We estimated the complexity of each compound using an aromaticity index (AI_{mod})[51]. A

molecule is considered more complex if it possesses more three-dimensional structures and

aromatic rings, which are both characterised by the presence of double bonds. However,

atoms such as nitrogen can be linked to carbon with double bonds without necessarily

forming aromatic rings. By using AI_{mod}, we considered both the number of double bound

equivalents (DBE), which approximates the saturation of each molecular formula, and the

number of heteroatoms, which provides insight into how carbon is bound. The number of

DBE and AI_{mod} were computed within the ICBM-OCEAN pipeline using equations described

by Koch & Dittmar[51,71].

Biodiversity sampling

We estimated biodiversity by sampling environmental DNA in 82 of the study lakes. During

sampling, we passed 1L of water through a 0.2 µm filter cartridge (Sterivex, Millipore, USA).

The cartridges were stored at -20°C until DNA was using an established protocol[80]. Briefly,

filter cartridges were opened in a sterile environment, and we placed the filters in a cryotube

containing silica and zirconia beads (3.0, 0.7, and 0.1 mm diameter). We then vortexed the

tubes at 2,850 rpm for 15 min to detach cells from the filter. To ensure DNA was released

into solution from lysed cells, we added 0.6 mL of phenol-chloroform-isopropanol (25:24:1),

0.6 mL of 5% cetrimonium bromide, 60 µL of 10% sodium dodecyl sulfate, and 60 µL 10%

N-lauroylsarcosine before vortexing the tubes a second time at 2,850 rpm for 15 min. We

then centrifuged the tubes at 16,000 g for 15 min at 4°C and collected the supernatant. To

remove phenol residues, we added 0.6 mL of chloroform-isopropanol (24:1) to the

supernatant, gently mixed the samples by inversion, and centrifuged at 16,000 g for 10 min at

4°C. We again collected the supernatant (ca. 0.6 mL) and added 1.2 mL of polyethylene

glycol with 1.6M sodium chlorine to precipitate the DNA overnight at 4°C. The following

morning, we centrifuged the samples at 17,000 g for 90 min at 4°C to obtain a DNA pellet

that we washed with ice-cold (-20°C) 70% ethanol. The pellet was dried at 37°C for 10 min

before being dissolved in 50 µL of ultrapure water (Qiagen, Germany). We used the same

protocol to extract DNA from a ZymoBIOMICS™ Microbial Community Standard II (Log

Distribution) (Zymo Research, USA) and ultrapure water (Qiagen, Germany) that served as

positive and negative controls, respectively. The positive control indicated no external

contamination, and no DNA was retrieved from the negative control.

We prepared shotgun metagenomics sequencing libraries with the NEBNext Ultra II FS kit

(New England Biolabs, USA) following the manufacturer's instructions. Briefly, DNA was

digested for 15 minutes to obtain end polished and A-tailed fragments of ~350bp. Then, we

attached Illumina adaptors by incubating the fragments at 37°C for 15 min in presence of a

ligation enhancer. We cleaned the samples using AMPure XP beads (Beckman Coulter,

USA) and attached Illumina-specific indexes (New England Biosciences, USA). We purified

the product using the AMPure XP beads[81], quantified DNA concentration on a Qubit

fluorometer (ThermoFisher, USA), and pooled libraries at equimolar concentrations before

sequencing on a NovaSeq 6000 PE 250 bp (Illumina, USA) at Novogene (Cambridge, UK),

who performed demultiplexing and checked read quality (Q30 $\geq$ 75% and Q20 = 100%).

We obtained a similar coverage for all samples (1st and 3rd quartiles: 4.35×10^7 and 6.70×10^7

reads). We assigned taxonomy to the reads using Kraken2[82] and a database containing

bacteria, eukaryotes (including fungi), and archaea. We used Bayesian Reestimation of

Abundance with Kraken to obtain read counts at the genus level[83] and estimated biodiversity

as the exponential of the Shannon index as for chemodiversity.

Ecosystem functioning in the field survey

At each field site, we measured three functions related to ecosystem metabolism (maximum

primary production, total respiration, and bacterial protein production) and two functions

related to carbon cycling (CH_4 and CO_2 emissions) from the lake surface.

Primary production and respiration were obtained based on differences in dissolved O_2 as

previously described[84]. First, we filled six glass serum bottles with 20 mL of lake water. All

bottles were crimped gastight without air bubbles. Two bottles were processed directly, two

bottles were incubated in the dark for 24 hours, and the last two bottles faced a cold light

(5000K) lamp and received ca. 34,000 lux for 3 hours. For each bottle, we measured the

dissolved O_2 concentration using optical fibre sensors (Presens, Germany). We recorded

dissolved O_2 concentration every second for approximately five minutes until we reached a

plateau. Then, we averaged the last 30 values of the plateau to infer the dissolved O_2

concentration inside the serum bottles. Primary production was estimated as the difference in

dissolved O_2 between the start of the incubation and after exposure to light. Respiration was

the difference in dissolved O_2 between the start and after 24 hours of dark incubation. We

averaged duplicates to obtain one value per lake.

Bacteria protein production (BPP) was estimated based on ^{3}H-leucine incorporation[85].

Directly after sampling, we filled six tubes (Eppendorf, Germany) containing 17nM of ^{3}H-

leucine with 1.5 mL of lake water. We previously added 300 µL of trichloroacetic acid to

three tubes that served as blanks. All tubes were immersed inside the lake in a dark container

at ca. 0.5 m depth for exactly 1 hour. We terminated the incubation by adding 300 μL of

trichloroacetic acid and preserved the samples on ice until we reached the lake shore – the

journey time varied from 1 to 30 mins. Then, we centrifuged the samples for 10 min at

16,000 g, removed the supernatant, washed the pellets with 1 mL of 5% trichloroacetic acid,

and centrifuged again for 10 min at 16,000 g. We removed the supernatant and let the pellets

air-dry for up to 10 min. We added 1 mL of scintillation cocktail (HiSafe, Perkin Elmer,

USA) and measured counts per minutes with a portable liquid scintillation counter (Triathler,

Hidex, Finland). To obtain the final BPP, we converted counts per minutes into degradation

per minutes, subtracted the value of the mean of the corresponding blanks and expressed the

results in mg C L^{-1} day^{-1} as previously described[86]. We averaged the values from the three

replicates when the standard deviation was within 10% of the mean and averaged the two

closest values otherwise.

CH_4 and CO_2 emissions were measured using the floating chamber technique[87]. We used a

chamber with a 0.07 m^2 perimeter and 0.002 m^3 volume that was covered in aluminium tape

to prevent heating. We connected the chamber to an ultraportable greenhouse gas analyser

(Los Gatos Research, USA) that uses off-axis integrated cavity output spectroscopy using

gastight PVC tubing. Before starting any measurements, we let the chamber equilibrate with

the atmosphere by waiting until the CH_4 and CO_2 curves plateaued. Then, we placed the

chamber on the water with 3 cm of the chamber immersed underwater. We monitored the

concentration of CH_4 and CO_2 inside the chamber every second for 15 mins. We estimated

fluxes as the linear slope between the gas concentration and time[87]. We repeated the process

(atmospheric equilibration and measurement) twice, and a third time if the two fluxes differed

by more than 10%. We then averaged the slopes to estimate CH_4 and CO_2 fluxes.

To calculate a multifunctionality index that reflects human values of nature, we assessed the

societal benefit of each measured function[1]. We considered that society benefits from higher

primary production because it can fuel food web productivity and fix atmospheric carbon into

biomass. Similarly, larger BPP benefits society because it increases the amount of carbon in

the food web and can store carbon for longer inside lakes. However, higher greenhouse gas

(CO_2 and CH_4) emissions negatively impact society due to their effect on climate. Higher

microbial respiration similarly consumes the carbon stored in lakes and emits CO_2 into the

environment, some of which will escape to the atmosphere. We therefore multiplied each of

CO_2 emissions, CH_4 emissions and respiration rate by -1 to ensure that they had the same

directionality of their societal benefits as primary production and BPP, that is, all functions

benefited society more as they increased in value. This decision seemingly contrasted with

the lab experiment where we calculated absolute values because both increases and decreases

in DOC and TDN indicate greater rates of beneficial biogeochemical cycling. However, both

approaches are consistent with the idea that different functions are valued in different ways.

We then calculated multifunctionality as for the laboratory experiment by summing z-scores

of each function.

Predictors of chemodiversity and ecosystem function

Ecosystem functions and chemodiversity are often influenced by physicochemical

variables[16,36] and climate[8,9]. Therefore, during the field study, we recorded water temperature

and pH using a multiprobe (HI-99171, Hanna Instruments, USA), measured water colour as

the absorbance at 254nm using a spectrophotometer (Flame UV-Vis, Ocean Optics, USA),

and measured chlorophyll *a* using a portable fluorometer (FluoroProbe III, bbe Moldaenke,

Germany). In the lab, we measured TDN concentrations concurrent with DOC concentrations

on the Shimadzu TOC-TNM-L analyser. Climate variables were the mean annual temperature

(MAT) and mean annual precipitation (MAP) of each lake at a ca. 31km resolution compiled

by the European Centre for Medium-Range Weather Forecasts ERA5 data product[88]. We

clipped the ERA5 analysis to the area of each catchment retrieved from the Global Lake area,

Climate, and Population dataset[89] and averaged across months for all pixels in a catchment to

obtain MAT and MAP. Because ecosystem function can also depend on the surrounding

catchment, a net primary production (NPP) raster (500m resolution) based on MODIS

satellite images[90] was clipped to each catchment and averaged across all pixels in a catchment

to derive NPP.

Testing the effects of chemodiversity and biodiversity on ecosystem multifunctionality

We first determined which variables influenced multifunctionality in the lab experiment and

field survey by fitting a linear model that only contained biodiversity and chemodiversity. We

then fitted a second linear model that included an interaction between biodiversity and

chemodiversity to test if any effects of chemodiversity on multifunctionality were conditional

on biodiversity. For the lab experiment, chemodiversity and biodiversity were treated as

factors, as we only had ≤4 levels of each variable, whereas we used them as continuous

variables in the field survey. We used the same model for testing the effect of chemodiversity

on multifunctionality as for testing the effect of chemodiversity on each individual function

in the laboratory experiment. However, we added mean annual temperature, water colour,

lake area, and chlorophyll *a* concentration to model each function in the field survey because

previous studies have revealed that these variables also predict ecosystem function[23,24]. All

models were fitted with R version 4.1.1[78].

In addition to the linear models, we tested explicit hypotheses about the drivers of

chemodiversity-ecosystem multifunctionality relationships using a path analysis. Here, we

designed a path analysis explaining variation in multifunctionality, chemodiversity, the

number of biochemical transformations, and DOM complexity. We hypothesized that

variation in multifunctionality was explained by chemodiversity, as well as biodiversity,

MAT, and MAP, which all predict ecosystem multifunctionality in other studies[3,10,38]. We

then hypothesized that chemodiversity was explained by its complexity and potential

transformation consistent with established theories for the reactivity of DOM[27]. Based on the

same theories, we hypothesized that the number of times a molecule was transformed was

also influenced by its complexity and all the environmental variables we measured, including

biodiversity. Finally, we hypothesized that DOM complexity was influenced by terrestrial

primary production that naturally produces many complex compounds that are leached into

downstream waters[65]. Although microbial activity also increases complexity[13], we expected

its effect to be minimal because most DOM in lakes originates from terrestrial production[16].

We then fitted this path analysis to lake-level observations using the piecewiseSEM

package[91] in R, which translates our hypotheses into a set of linear models. After fitting the

initial path analysis, we included missing paths identified by tests of directed separation as

described in Fig. S13.

Identifying the production and consumption of individual compounds

For the laboratory experiment, we first identified individual molecular formulae that were

consumed or produced by microbes. We compared the relative abundance of each molecular

formula at the end of the incubation experiment from all biodiversity levels against the

relative abundance of the same formula in the control with a Welch's t-test. We only

considered the final time point of the control to account for compounds that could be

produced or consumed simply due to abiotic processes. We considered compounds were

produced and consumed if their abundance significantly increased and decreased compared to

the control, respectively. We then summed the total number of produced and consumed

compounds at each chemodiversity level.

For the field survey, we tested how the consumption and production of formulae was

associated with each function that varied with chemodiversity across Europe (CH_4 and CO_2

emissions and bacterial biomass production). We correlated the relative abundance of each

formula in each lake with the value of each function in the corresponding lake using

Spearman rank correlations. We considered statistically significant negative and positive

correlations to indicate that consumption and production of a formula, respectively, was

associated with a given function.

Extrapolating chemodiversity and ecosystem function from our field survey

To extrapolate chemodiversity and each measured function that varied with chemodiversity

across Europe (CH_4 and CO_2 emissions and bacterial biomass production), we first fitted

linear models to these variables. The models included all physio-chemical and environmental

variables measured in the field survey, and we included chemodiversity and biodiversity as

predictors when modelling each function. Then, we simplified the full models with stepwise

variable selection guided by the Akaike information criterion (AIC). We sequentially

removed the least supported variable, that is, most increased AIC when retained. If all

variables reduced AIC when retained, we dropped the variable that least reduced AIC by <2

units[92]. We repeated this selection process until all variables retained in the model reduced

AIC by ≥2 units. The goal of this analysis was to predict each function as accurately as

possible to extrapolate across Europe rather than simply compare the importance of

chemodiversity versus biodiversity as when modelling multifunctionality. Model selection

indicated that chemodiversity was best explained by NPP, MAP, and chlorophyll *a*

concentration (Table S3), which was identical to the variables that explained variation in

chemodiversity within our path analysis (Figs 2, S13).

To extrapolate our linear models across Europe in 2019, we forced them with the

same MAP and NPP rasters described above. The other predictors required in our models

were water colour and chlorophyll *a* concentration (Table S3, S4), which are unavailable at a

European-scale in publicly available datasets. Therefore, we predicted water colour from

elevation, MAP, and lake area using a linear model identical to past global studies predicting

water colour[75]. Similarly, we predicted chlorophyll *a* from MAT, elevation, MAP, and lake

area using another linear model. Past global studies have used all these same variables to

predict chlorophyll *a*, except that they included TP, which was also unavailable from publicly

available datasets at a European-scale[23,24]. Although the models explained 25% and 29% of

the variation in water colour and chlorophyll *a*, respectively, the errors introduced by using

these models are not relevant for our calculations focusing on relative changes over time (see

section below).

Forecasting the future functioning of lakes

We estimated how each function that varied with chemodiversity (CH_4 and CO_2 emissions

and bacterial biomass production) would change across Europe by 2100 solely because of

changes to chemodiversity. To predict chemodiversity in 2100, we first obtained the IPSL-

CM6A[60], HadGEM3-GC31[93], and ACCESS-ESM1.5[94] climate forecasts for 2100. We

selected these forecasts because they emphasise the relevance of terrestrial-atmosphere

exchange and are frequently used in climate studies[95]. We downloaded rasters of MAT and

MAP for likely (SSP2) and worst case (SSP5) scenarios for future changes in atmospheric

CO_2 concentrations for each forecast from the Copernicus Climate Data Store

(https://cds.climate.copernicus.eu/). NPP predictions for each forecast were downloaded from

the Earth System Grid Federation servers (https://esgf-node.llnl.gov/search/cmip6/). Because

no NPP data were available for the HadGEM3-GC31 model, we used the average of the

IPSL-CM6A and ACCESS-ESM1.5 models. We then input these rasters into our linear

model of chemodiversity to predict chemodiversity in 2100 across Europe (Table S3). To

generate the predictor of chlorophyll a, we again used the model described in the previous

section but with MAT and MAP from the climate forecasts, as well as elevation and lake

area. The chemodiversity raster was then input into the linear models of CH_4 and CO_2

emissions and bacterial biomass production (Table S4). All predictors other than

chemodiversity were set at their 2019 values. Rather than focus on absolute values of each

function, we calculated the change in each function between 2019 and 2100 when only

changing chemodiversity, that is, holding all other variables constant. Because errors

associated with models of water colour and chlorophyll a are likely to be systematic, that is,

non-random, the errors should remain similar between time periods with chemodiversity

being the sole difference in the comparison. Furthermore, estimates of the magnitude of

changes of CH_4 and CO_2 for all of Europe were well in line with previous global studies[23,24],

further suggesting that errors in absolute values of water colour and chlorophyll a were of

minor importance.

References

1. Manning, P. *et al.* Redefining ecosystem multifunctionality. *Nature Ecology and*

Evolution **2**, 427–436 (2018).

2. Naeem, S., Duffy, J. E. & Zavaleta, E. The Functions of Biological Diversity in an Age

of Extinction. *Science* **336**, 1401–1406 (2012).

3. Delgado-Baquerizo, M. *et al.* Microbial diversity drives multifunctionality in terrestrial

ecosystems. *Nature Communications* **7**, 10541 (2016).

4. Guillemette, F., Leigh Mccallister, S. & Del Giorgio, P. A. Selective consumption and

metabolic allocation of terrestrial and algal carbon determine allochthony in lake bacteria.

ISME Journal **10**, (2016).

5. Tanentzap, A. J. *et al.* Terrestrial support of lake food webs: Synthesis reveals controls

over cross-ecosystem resource use. *Science Advances* **3**, (2017).

6. Tanentzap, A. J. *et al.* Chemical and microbial diversity covary in fresh water to

influence ecosystem functioning. *Proceedings of the National Academy of Sciences of the*

United States of America **116**, 24689–24695 (2019).

7. Bruelheide, H. *et al.* Global trait–environment relationships of plant communities. *Nature*

Ecology and Evolution **2**, 1906–1917 (2018).

8. Maestre, F. T. *et al.* Plant species richness and ecosystem multifunctionality in global

drylands. *Science* **335**, 214–218 (2012).

9. Lefcheck, J. S. *et al.* Biodiversity enhances ecosystem multifunctionality across trophic

levels and habitats. *Nature Communications* **6**, 6936 (2015).

10. Jing, X. *et al.* The links between ecosystem multifunctionality and above- and

belowground biodiversity are mediated by climate. *Nat Commun* **6**, 8159 (2015).

11. Noriega-Ortega, B. E. *et al.* Does the chemodiversity of bacterial exometabolomes

sustain the chemodiversity of marine dissolved organic matter? *Frontiers in*

Microbiology **10**, 215 (2019).

12. Graham, E. B. *et al.* Multi 'omics comparison reveals metabolome biochemistry, not

microbiome composition or gene expression, corresponds to elevated biogeochemical

function in the hyporheic zone. *Science of the Total Environment* **642**, 742–753 (2018).

13. Dittmar, T. *et al.* Enigmatic persistence of dissolved organic matter in the ocean. *Nature*

Reviews Earth and Environment **2**, 570–583 (2021).

14. Tanentzap, A. J. & Fonvielle, J. A. Chemodiversity in freshwater health. *Science* **383**,

1412–1414 (2024).

15. Testa, B., Vistoli, G., Pedretti, A. & Bojarski, A. J. Atomic Diversity, Molecular

Diversity, and Chemical Diversity: The Concept of Chemodiversity. *Chemistry &*

Biodiversity **6**, 1145–1151 (2009).

16. Kellerman, A. M., Dittmar, T., Kothawala, D. N. & Tranvik, L. J. Chemodiversity of

dissolved organic matter in lakes driven by climate and hydrology. *Nature*

Communications **5**, 3804 (2014).

17. Hall, S. J., Ye, C., Weintraub, S. R. & Hockaday, W. C. Molecular trade-offs in soil

organic carbon composition at continental scale. *Nature Geoscience* **13**, 687–692 (2020).

18. Osterholz, H., Niggemann, J., Giebel, H. A., Simon, M. & Dittmar, T. Inefficient

microbial production of refractory dissolved organic matter in the ocean. *Nature*

Communications **6**, 1–8 (2015).

19. Fudyma, J. D., Chu, R. K., Graf Grachet, N., Stegen, J. C. & Tfaily, M. M. Coupled

Biotic-Abiotic Processes Control Biogeochemical Cycling of Dissolved Organic Matter

in the Columbia River Hyporheic Zone. *Frontiers in Water* **2**, (2021).

20. Helms, J. R. *et al.* Absorption spectral slopes and slope ratios as indicators of molecular

weight, source, and photobleaching of chromophoric dissolved organic matter. *Limnology*

and Oceanography **53**, 955–969 (2008).

21. Rose, K. C., Winslow, L. A., Read, J. S. & Hansen, G. J. A. Climate-induced warming of

lakes can be either amplified or suppressed by trends in water clarity. *Limnology and*

Oceanography Letters **1**, 44–53 (2016).

22. Senar, O. E., Creed, I. F. & Trick, C. G. Lake browning may fuel phytoplankton biomass

and trigger shifts in phytoplankton communities in temperate lakes. *Aquat Sci* **83**, 21

(2021).

23. Rosentreter, J. A. *et al.* Half of global methane emissions come from highly variable

aquatic ecosystem sources. *Nature Geoscience* **14**, 225–230 (2021).

24. DelSontro, T., Beaulieu, J. J. & Downing, J. A. Greenhouse gas emissions from lakes and

impoundments: Upscaling in the face of global change. *Limnology And Oceanography*

Letters **3**, 64–75 (2018).

25. Benner, R. & Amon, R. M. W. The Size-Reactivity Continuum of Major Bioelements in

the Ocean. *Annual Review of Marine Science* **7**, 185–205 (2015).

26. Hutchins, R. H. S. *et al.* The Optical, Chemical, and Molecular Dissolved Organic Matter

Succession Along a Boreal Soil-Stream-River Continuum. *Journal of Geophysical*

Research: Biogeosciences **122**, 2892–2908 (2017).

27. Kothawala, D. N., Kellerman, A. M., Catalán, N. & Tranvik, L. J. Organic Matter

Degradation across Ecosystem Boundaries: The Need for a Unified Conceptualization.

Trends in Ecology & Evolution **36**, 113–122 (2020).

28. Lehner, B. & Döll, P. Development and validation of a global database of lakes,

reservoirs and wetlands. *Journal of Hydrology* **296**, 1–22 (2004).

29. Mentges, A., Feenders, C., Seibt, M., Blasius, B. & Dittmar, T. Functional Molecular

Diversity of Marine Dissolved Organic Matter Is Reduced during Degradation. *Frontiers*

in Marine Science **4**, 194 (2017).

30. Kritzberg, E. S. *et al.* Browning of freshwaters: Consequences to ecosystem services,

underlying drivers, and potential mitigation measures. *Ambio* **49**, 375–390 (2020).

31. Kolmakova, O. V. *et al.* Effects of zooplankton carcasses degradation on freshwater

bacterial community composition and implications for carbon cycling. *Environmental*

Microbiology **21**, 34–49 (2019).

32. Gamfeldt, L. & Roger, F. Revisiting the biodiversity-ecosystem multifunctionality

relationship. *Nature Ecology and Evolution* **1**, (2017).

33. Le Bagousse-Pinguet, Y. *et al.* Phylogenetic, functional, and taxonomic richness have

both positive and negative effects on ecosystem multifunctionality. *Proceedings of the*

National Academy of Sciences **116**, 8419–8424 (2019).

34. Foster, K. R. & Bell, T. Competition, Not Cooperation, Dominates Interactions among

Culturable Microbial Species. *Current Biology* **22**, 1845–1850 (2012).

35. Rivett, D. W. & Bell, T. Abundance determines the functional role of bacterial

phylotypes in complex communities. *Nat Microbiol* **3**, 767–772 (2018).

36. Delgado-Baquerizo, M. *et al.* Multiple elements of soil biodiversity drive ecosystem

functions across biomes. *Nat Ecol Evol* **4**, 210–220 (2020).

37. Byrnes, J. E. K. *et al.* Investigating the relationship between biodiversity and ecosystem

multifunctionality: Challenges and solutions. *Methods in Ecology and Evolution* **5**, 111–

124 (2014).

38. Allan, E. *et al.* Land use intensification alters ecosystem multifunctionality via loss of

biodiversity and changes to functional composition. *Ecology Letters* (2015)

doi:10.1111/ele.12469.

39. Kellerman, A. M. *et al.* Unifying Concepts Linking Dissolved Organic Matter

Composition to Persistence in Aquatic Ecosystems. *Environmental Science and*

Technology **52**, 2538–2548 (2018).

40. Merbach, W., Ruppel, S. & Schulze, J. Dinitrogen Fixation of Microbe-Plant

Associations as Affected by Nitrate and Ammonium Supply. *Isotopes in Environmental*

and Health Studies **34**, (1998).

41. Jung, H. *et al.* Comparative genomic analysis of Chryseobacterium species: deep insights

into plant-growth-promoting and halotolerant capacities. *Microbial Genomics* **9**, 001108

(2023).

42. Kraemer, S. A. *et al.* A large-scale assessment of lakes reveals a pervasive signal of land

use on bacterial communities. *ISME Journal* **14**, 3011–3023 (2020).

43. Thompson, L. R. *et al.* A communal catalogue reveals Earth's multiscale microbial

diversity. *Nature* **551**, 457–463 (2017).

44. D'Andrilli, J., Cooper, W. T., Foreman, C. M. & Marshall, A. G. An ultrahigh-resolution

mass spectrometry index to estimate natural organic matter lability. *Rapid*

Communications in Mass Spectrometry **29**, 2385–2401 (2015).

45. Günthel, M. *et al.* Contribution of oxic methane production to surface methane emission

in lakes and its global importance. *Nat Commun* **10**, 5497 (2019).

46. Ernst, L. *et al.* Methane formation driven by reactive oxygen species across all living

organisms. *Nature* **603**, 482–487 (2022).

47. Mosher, J. J., Kaplan, L. A., Podgorski, D. C., McKenna, A. M. & Marshall, A. G.

Longitudinal shifts in dissolved organic matter chemogeography and chemodiversity

within headwater streams: a river continuum reprise. *Biogeochemistry* **124**, 371–385

(2015).

48. Lapierre, J.-F., Seekell, D. A. & del Giorgio, P. A. Climate and landscape influence on

indicators of lake carbon cycling through spatial patterns in dissolved organic carbon.

Global Change Biology **21**, 4425–4435 (2015).

49. Fuchs, G., Boll, M. & Heider, J. Microbial degradation of aromatic compounds — from

one strategy to four. *Nat Rev Microbiol* **9**, 803–816 (2011).

50. Zhang, R., Wang, J., Milligan, S. & Yan, Y. Microbial utilization of lignin-derived

aromatics: Via a synthetic catechol meta-cleavage pathway. *Green Chemistry* **23**, 8238–

8250 (2021).

51. Koch, B. P. & Dittmar, T. From mass to structure: An aromaticity index for high-

resolution mass data of natural organic matter. *Rapid Communications in Mass*

Spectrometry **20**, 926–932 (2006).

52. Arrieta, J. M. *et al.* Dilution limits dissolved organic carbon utilization in the deep ocean.

Science **348**, 331–333 (2015).

53. Weng, J. K., Philippe, R. N. & Noel, J. P. The rise of chemodiversity in plants. *Science*

336, 1667–1670 (2012).

54. Bertilsson, S. & Tranvik, L. J. Photochemical transformation of dissolved organic matter

in lakes. *Limnology and Oceanography* **45**, 753–762 (2000).

55. Olson, C. R., Solomon, C. T. & Jones, S. E. Shifting limitation of primary production:

experimental support for a new model in lake ecosystems. *Ecology Letters* **23**, 1800–

1808 (2020).

56. Dittmar, T., Hertkorn, N., Kattner, G. & Lara, R. J. Mangroves, a major source of

dissolved organic carbon to the oceans. *Global Biogeochemical Cycles* **20**, (2006).

57. Cuhel, R. L. & Aguilar, C. Ecosystem Transformations of the Laurentian Great Lake

Michigan by Nonindigenous Biological Invaders. *Annual Review of Marine Science* **5**,

289–320 (2013).

58. Waska, H., Koschinsky, A., Ruiz Chancho, M. J. & Dittmar, T. Investigating the

potential of solid-phase extraction and Fourier-transform ion cyclotron resonance mass

spectrometry (FT-ICR-MS) for the isolation and identification of dissolved metal–

organic complexes from natural waters. *Marine Chemistry* **173**, 78–92 (2015).

59. Azam, F. & Malfatti, F. Microbial structuring of marine ecosystems. *Nat Rev Microbiol*

5, 782–791 (2007).

60. Boucher, O. *et al.* Presentation and Evaluation of the IPSL-CM6A-LR Climate Model.

Journal of Advances in Modeling Earth Systems **12**, e2019MS002010 (2020).

61. Zhang, W., Shen, J. & Wang, J. Linking pollution to biodiversity and ecosystem multifunctionality across benthic-pelagic habitats of a large eutrophic lake: A whole-ecosystem perspective. *Environmental Pollution* **285**, 117501 (2021).

62. Zhuang, Q. *et al.* Current and Future Global Lake Methane Emissions: A Process-Based Modeling Analysis. *Journal of Geophysical Research: Biogeosciences* **128**, e2022JG007137 (2023).

63. Messager, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nature Communications* **7**, 13603 (2016).

64. O'Neill, B. C. *et al.* The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model Development* **9**, 3461–3482 (2016).

65. Defossez, E. *et al.* Spatial and evolutionary predictability of phytochemical diversity. *Proceedings of the National Academy of Sciences of the United States of America* **118**, (2021).

66. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).

67. Brienen, R. J. *et al.* Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature communications* **11**, 4241 (2020).

68. Scheuerl, T. *et al.* Bacterial adaptation is constrained in complex communities. *Nat Commun* **11**, 754 (2020).

69. Gheorghiu, C., Smith, D. S., Al-Reasi, H. A., McGeer, J. C. & Wilkie, M. P. Influence of natural organic matter (NOM) quality on Cu–gill binding in the rainbow trout (Oncorhynchus mykiss). *Aquatic Toxicology* **97**, 343–352 (2010).

70. Dittmar, T., Koch, B., Hertkorn, N. & Kattner, G. A simple and efficient method for the

solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater. *Limnology*

and Oceanography: Methods **6**, 230–235 (2008).

71. Merder, J. *et al.* ICBM-OCEAN: Processing Ultrahigh-Resolution Mass Spectrometry

Data of Complex Molecular Mixtures. *Analytical Chemistry* **92**, 6832–6838 (2020).

72. Shannon, C. E. C. & Weaver, W. *The Mathematical Theory of Information. Urbana*

University of Illinois Press (1949). doi:10.1007/978-1-4615-0975-2.

73. Jost, L. Entropy and diversity. *Oikos* **113**, 363–375 (2006).

74. European Environmental Agency. WISE Water Framework Directive Database.

https://www.eea.europa.eu/data-and-maps/data/wise-wfd-4 (2019).

75. Toming, K. *et al.* Predicting lake dissolved organic carbon at a global scale. *Scientific*

Reports **10**, 8471 (2020).

76. McMaster, G. S. & Wilhelm, W. W. Growing degree-days: One equation, two

interpretations. *Agricultural and Forest Meteorology* **87**, 291–300 (1997).

77. Scott Chamberlain. rnoaa: 'NOAA' Weather Data from R. R package version 0.8.4.

https://CRAN.R-project.org/package=rnoaa (2019).

78. R Core team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. Preprint at

https://doi.org/10.2788/95827 (2017).

79. Potter, B. B. & Wimsatt, J. C. USEPA method 415.3: Quantifying TOC, DOC, and

SUVA. *Journal AWWA* **104**, E358–E369 (2012).

80. Nercessian, O., Noyes, E., Kalyuzhnaya, M. G., Lidstrom, M. E. & Chistoserdova, L.

Bacterial populations active in metabolism of C1 compounds in the sediment of Lake

Washington, a freshwater lake. *Applied and Environmental Microbiology* **71**, 6885–6899

(2005).

81. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet. J. **17**, 10–12 (2011).

82. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biology **20**, 1–13 (2019).

83. Lu, J., Breitwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: Estimating species

abundance in metagenomics data. *PeerJ Computer Science* **2017**, e104 (2017).

84. Revsbech, N. P., JØrgensen, B. B. & Brix, O. Primary production of microalgae in

sediments measured by oxygen microprofile, H14CO3 ☐ fixation, and oxygen exchange

methods. *Limnology and Oceanography* **26**, 717–730 (1981).

85. Simon, M. & Azam, F. Protein content and protein synthesis rates of planktonic marine

bacteria. *Marine Ecology Progress Series* **51**, 201–213 (1989).

86. Amand, A. S. *et al.* CASCADING TROPHIC INTERACTIONS PROJECT. (1978).

87. Bastviken, D., Cole, J., Pace, M. & Tranvik, L. Methane emissions from lakes:

Dependence of lake characteristics, two regional assessments, and a global estimate.

Global Biogeochemical Cycles **18**, GB4009 (2004).

88. Copernicus Climate Change Service. C3S ERA5-Land reanalysis. Copernicus Climate

Change Service. (2019) doi:10.24381/cds.68d2bb30.

89. Meyer, M. F., Labou, S. G., Cramer, A. N., Brousil, M. R. & Luff, B. T. The global lake

area, climate, and population dataset. *Scientific Data* **7**, 174 (2020).

90. Running, S. & Zhao, M. MOD17A3HGF MODIS/Terra Net Primary Production Gap-

Filled Yearly L4 Global 500 m SIN Grid V006. 2019, distributed by NASA EOSDIS

Land Processes DAAC. https://doi.org/10.5067/MODIS/MOD17A3HGF.006 (2020).

91. Lefcheck, J. S. piecewiseSEM: Piecewise structural equation modelling in r for ecology,

evolution, and systematics. *Methods in Ecology and Evolution* **7**, 573–579 (2016).

92. Burnham, K. P. & Anderson, D. R. Multimodel Inference: Understanding AIC and BIC

in Model Selection. *Sociological Methods & Research* **33**, 261–304.

93. Williams, K. D. *et al.* The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and

GC3.1) Configurations. *Journal of Advances in Modeling Earth Systems* **10**, 357–380

(2018).

94. Bi, D. *et al.* Configuration and spin-up of ACCESS-CM2, the new generation Australian

Community Climate and Earth System Simulator Coupled Model. *JSHESS* **70**, 225

(2020).

95. Friedlingstein, P. *et al.* Global Carbon Budget 2021. *Earth System Science Data* **14**,

1917–2005 (2022).

1194