Microbial metabolism of food allergens determines the severity of IgE-mediated anaphylaxis

Elisa Sánchez-Martínez[1,x], Liam E. Rondeau[2,x], Manuel Garrido-Romero[1,3], Bruna Barbosa da Luz[2], Dominic A. Haas[2], Gavin Yuen[2], Peter Hall[2], Rebecca Dang[2], Xuan-Yu Wang[2], Lucía Moreno-Serna[1], Celia López-Sanz[1], Emilio Nuñez-Borque[1], Maria Garrido-Arandia[4], Araceli Diaz-Perales[4], Yolanda R. Carrasco[5], Joshua F.E. Koenig[6], Tina D. Walker[6], Manel Jordana[6], Elena F. Verdu[2], Premysl Bercik[2], Giada De Palma[2] Michael G. Surette[2], Pedro Ojeda[7], Francisco Vega[8], Carlos Blanco[8], Lingdi Zhang[9], Supinda Bunyavanich[9], Wayne G. Shreffler[10], Sarita U. Patil[10], F. Javier Moreno[3], Rodrigo Jiménez-Saiz[1,6,11,y,*], Alberto Caminero[2,y,z,*]

[1]Department of Immunology, Instituto de Investigación Sanitaria Hospital Universitario de La Princesa (IIS-Princesa), Universidad Autónoma de Madrid (UAM), Madrid, Spain.

[2]Department of Medicine, Farncombe Family Digestive Health Research Institute, McMaster University, Hamilton, ON, Canada.

[3]Department of Bioactivity and Food Analysis, Instituto de Investigación en Ciencias de la Alimentación (CIAL), CSIC-UAM, CEI, Madrid, Spain.

[4]Centre for Plant Biotechnology and Genomics (CBGP), Universidad Politécnica de Madrid (UPM-INIA), Madrid, Spain.

[5]Department of Immunology and Oncology, Centro Nacional de Biotecnología (CNB)-CSIC, Madrid, Spain.

[6]Department of Medicine, McMaster Immunology Research Centre (MIRC), Schroeder Allergy and Immunology Research Institute (SAIRI), McMaster University, Hamilton, ON, Canada.

[7]Clínica de Asma y Alergia Dres. Ojeda, Madrid, Spain.

[8]Department of Allergy, Hospital Universitario de La Princesa, IIS-Princesa, Madrid, Spain.

[9]Department of Genetics and Genomic Sciences and Department of Pediatrics, Icahn School of Medicine at Mount Sinai, New York, New York, USA.

[10]Food Allergy Center and Center for Immunology and Inflammatory Diseases, Massachusetts General Hospital, Boston, Massachusetts, USA.

[11]Faculty of Experimental Sciences, Universidad Francisco de Vitoria (UFV), Madrid, Spain.

[x]First co-authors who contributed equally to this work.
[y]Senior co-authors.
[z]Lead contact.
*Correspondence: rodrigo.jimenez@ufv.es / @R_JimenezSaiz (R.J.-S); acamine@mcmaster.ca (A.C.) / @LabCaminero

SUMMARY

Anaphylaxis is an acute, potentially life-threatening reaction, often triggered by foods and largely mediated by IgE. Critically important to anaphylaxis are the factors that modulate its severity. The human microbiota is known to influence oral tolerance, but the microbial mechanisms directly involved in IgE-mediated anaphylaxis remain unknown. Here, we demonstrate that human saliva harbors peanut-degrading bacteria that metabolize immunodominant allergens (Ara h 1 and 2) and alter IgE-binding. Additionally, we provide *in vivo* evidence that oral bacteria metabolize peanut allergens, influencing systemic allergen exposure and anaphylaxis severity. Finally, in clinical studies, we observe that common peanut-degrading bacteria, such as *Rothia,* from the oral cavity, are more abundant in peanut-allergic patients who exhibit better tolerance to allergen exposure. Altogether, these results demonstrate that human microbiota modulates IgE-mediated reactions through allergen metabolism. We reveal a novel microbial mechanism with potential to prevent, or reduce, the severity of IgE-mediated anaphylaxis.

KEYWORDS

Allergens, Allergy, Anaphylaxis, IgE, Inflammation, Metabolism, Microbiota, Oral, Peanut, Saliva

INTRODUCTION

Inflammation typically restores homeostasis in response to tissue damage or infection but can cause immunopathology when dysregulated[1]. Anaphylaxis exemplifies this, with acute and potentially fatal outcomes within minutes. The main pathway of anaphylaxis is mediated by immunoglobulin (Ig)E and mast cells: upon allergen binding, mast cell-bound IgE induces the rapid release of inflammatory mediators like tryptase or leukotrienes, which drive the acute clinical manifestations of anaphylaxis[2-4]. Foods are common anaphylaxis triggers, with peanut (PN) being a leading cause of food-induced anaphylaxis and allergy-related deaths among children[5,6]. Food-anaphylaxis, particularly PN-triggered, is most common in North America and Europe and less frequent in the Asia-Pacific[7,8]. The burden of PN allergy is aggravated by its persistence in >70% of individuals, the absence of curative treatments, and the frequent accidental exposures despite patients' avoidance efforts[9-12].

The severity of food-induced anaphylaxis is not solely determined by allergen-specific IgE levels, as clinical reactivity is influenced by a complex interplay of genetic, environmental, dietary, and behavioral factors, as well as co-factors and comorbidities[13,14]. Notably, clinical observations often reveal a disconnect between serum levels of allergen-specific IgE—the key molecule involved in food-induced anaphylaxis—and clinical reactivity[15-17], underscoring the importance of elucidating additional modulators of anaphylaxis[2]. The human microbiota has gained considerable attention in this regard for its capacity to influence both oral tolerance and dietary antigen immunogenicity[18-22]. Indeed, studies have revealed differences in the intestinal microbiota composition of food-allergic patients[23-25]. Nonetheless, the microbial mechanisms involved in food-induced anaphylaxis remain largely unknown.

The oro-gastrointestinal microbiota is often regarded as a "second metabolic organ" capable of breaking down dietary components that are otherwise resistant to human digestive enzymes. Certain food allergens, including the immunodominant PN allergens Ara h 1 and 2[26,27], resist complete digestion by mammalian digestive enzymes[28,29]. Here, we demonstrate that human saliva and the small intestine contain PN-degrading bacteria capable of metabolizing immunodominant allergens and modulating IgE-specific immunity. We also show the *in vivo* capacity of these bacteria to participate in PN-metabolism*,* thereby modulating systemic allergen access and IgE-mediated anaphylaxis. Finally, in two clinical studies, we describe that common PN-degrading bacteria, such as *Rothia* from the oral cavity*,* are more abundant in allergic patients who exhibit higher allergen thresholds to controlled allergen exposure. These results demonstrate the human microbiota's ability to modulate IgE-mediated reactions through allergen metabolism and reveal a novel microbial mechanism with translational potential to prevent or mitigate IgE-mediated anaphylaxis.

RESULTS

Microbiota participates in peanut allergen metabolism

To study the role of microbes in PN metabolism, we used C57BL/6 mice with distinct microbiota compositions: germ-free (GF), specific pathogen-free (SPF; diverse), and altered Schaedler flora (ASF; stable with limited species) (**Figure S1**). Mice were gavaged with PN and sacrificed after 40 minutes (**Figure 1A**). The quantities of PN allergens Ara h 1 and 2 were higher in small intestinal content of GF and ASF mice, compared to mice with diverse SPF microbiota (**Figure 1B**). Moreover, microbial alpha-diversity across groups correlated inversely with Ara h 1 and 2 in the small intestine (**Figure S1D**). We also detected variations in the concentrations of these allergens systemically. Compared to SPF mice, GF and ASF mice presented higher serum levels of Ara h 1 but lower serum levels of Ara h 2 (**Figure 1C**). To determine whether these differences were due to variations in the digestive capacity, intestinal contents of PN-naïve mice were incubated with PN (crude PN extract, CPE) *ex vivo*. The intestinal contents of SPF mice degraded Ara h 1 and 2 more effectively than those of GF and ASF mice (**Figure 1D**). We hypothesized that microbial metabolism influenced the differences observed in PN allergen digestion and attempted to isolate PN-degrading bacteria from SPF and ASF mouse intestinal contents. Notably, PN-degrading bacteria were present in the small and large intestines of SPF mice but not found in ASF mice. While various PN-degrading bacterial species were identified, their capacity to digest Ara h 1 and 2 varied by bacterial strain (**Figure 1E**). Together, these results demonstrate that the intestinal microbiota metabolizes PN allergens and influences their systemic availability.

Figure 1. Microbiota participates in allergen metabolism *in vivo*.
(A) Experimental design. Germ-free (GF), specific pathogen-free (SPF), and altered Schaedler flora (ASF)
C57BL/6 mice received PN intragastrically (i.g.). Peripheral blood and intestinal contents were collected 40
minutes post-PN delivery. n=4–16 mice per group; 2 independent experiments. (B–C) PN allergens (Ara h
1 and 2) in small intestinal content (B) and serum from peripheral blood (C). Each dot=one mouse. *P* values
were calculated using one-way ANOVA with Tukey's post-hoc test. (D) Heatmap showing digestion capacity
of GF, ASF, and SPF microbiota against Ara h 1 and 2. PN-naïve intestinal contents were incubated with
PN allergens (CPE) *in vitro* and remaining allergens quantified. Each row = one mouse. Degradation
capacity is expressed as % degraded relative to control incubations with CPE alone (blue=0%, red=100%).
Statistics: one-way ANOVA with Tukey's post-hoc test. Significance levels: Small intestine Ara h 1: GF vs.
SPF (*P*<0.0001), ASF vs. SPF (*P*=0.0290). Small intestine Ara h 2: GF vs. SPF (*P*<0.0001), ASF vs. SPF
(*P*<0.0001). Large intestine Ara h 1: GF vs. SPF (*P*<0.0001), ASF vs. SPF (*P*=0.0289). Large intestine Ara
h 2: GF vs. SPF (*P*<0.0001), ASF vs. SPF (*P*<0.0024). (E) Heatmap of PN-degrading isolates and
abundance of bacterial isolates from the small and large intestine of SPF mice. Isolates were obtained by
plating SPF intestinal contents on PN-agar and selecting colonies with hydrolytic halos, indicative of PN

degradation. Isolates were incubated in liquid media with CPE to assess Ara h 1 and 2 degradation. Each row=one isolate. Allergen degradation is shown using the same blue-red scale as in (D). Relative abundance is represented on a green scale (low=light, dark=high), based on the frequency of isolation relative to the total number of isolates. See also Figure S1.

Mice with limited peanut-degrading bacteria exhibit stronger mucosal anaphylaxis markers

To investigate the impact of microbial PN metabolism on acute allergic reactions, we used ASF and SPF C3H/HeN mice, a strain more susceptible to type 2 immunity and anaphylaxis upon oral PN challenge[30]. Consistent with our previous findings in C57BL/6 mice, ASF mice exhibited a reduced capacity to metabolize PN compared to SPF C3H/HeN mice (**Figure S2A–C**). ASF and SPF C3H/HeN mice were sensitized to PN with cholera toxin (CT) via intragastric (i.g.) administration once per week for 6 weeks, followed by an i.g. PN-challenge to assess anaphylactic reactions to PN allergens (**Figure 2A**). ASF mice, which present reduced PN-degrading bacteria, had higher serum levels of PN-specific IgE than SPF mice (**Figure 2B**). They also exhibited higher serum levels of mucosal mast cell protease-1 (mMCP-1), indicating local activation of effector cells in the gastrointestinal tract, along with systemic hypothermia following PN challenge (**Figure 2C & Figure S3A**). To bypass the influence of microbial PN metabolism on IgE production, we sensitized SPF and ASF mice with PN + CT via intraperitoneal (i.p.) injection (**Figure 2A**). While both groups exhibited similar serum levels of PN-specific IgE (**Figure 2D**), ASF mice had higher serum mMCP-1 levels and developed hypothermia following PN challenge (**Figure 2E & Figure S3B**). We further bypassed the endogenous capacity to generate allergen-specific IgE in the sensitization process by passively transferring serum from PN-allergic mice into SPF- and ASF-colonized mice (**Figure 2A**). Both mouse strains presented similar levels of PN-specific IgE in the serum after passive sensitization (**Figure 2F**). However, ASF mice again exhibited higher serum levels of mMCP-1, platelet-activating factor acetylhydrolase (PAF-AH), and hypothermia (**Figure 2G & Figure S3C–D**). These results demonstrate that the presence of certain microbes at the site of allergen challenge can protect from anaphylaxis.

To further investigate how microbiota composition influences anaphylactic outcomes, we characterized bacterial community structure and diversity in ASF and SPF mice across all sensitization protocols (**Figure S4A**). As expected, ASF mice exhibited lower alpha-diversity compared to SPF mice, and this reduction in observed species was associated with increased intestinal levels of Ara h 1 and 2 following PN challenge (**Figure S4B**). Moreover, Ara h 1 and 2 levels inversely correlate with alpha-diversity across sensitization methods (**Figure 2H & Figure S4C**), and serum mMCP-1 concentrations showed similar negative correlations with alpha-diversity in i.g., i.p., and passive models (**Figure S4D**). These findings demonstrate that microbial complexity is linked to enhanced PN degradation and reduced effector responses.

Figure 2. Limited microbial peanut metabolism amplifies mucosal anaphylaxis markers.
(A) Experimental design for peanut (PN) sensitization and challenge of C3H-HeN mice colonized with ASF and SPF microbiota. Mice were sensitized (1) intragastrically (i.g.; PN + cholera toxin) once per week for 6 weeks, (2) intraperitoneally (i.p.; PN + cholera toxin) once per week for 4 weeks, and (3) passively (i.p.; serum from PN-allergic mice) once every 48 hours, 3 times. Sensitized mice were challenged with a PN bolus i.g. and sacrificed 40 minutes later. (B–C) Serum PN-specific IgE (B) and serum mucosal mast cell protease 1 (mMCP-1) (C) of i.g. sensitized ASF and SPF mice after PN challenge. (D-E) Serum PN-specific IgE (D) and serum mMCP-1 (E) of i.p. sensitized ASF and SPF mice after PN challenge. (F–G) Serum PN-specific IgE (F), serum mMCP-1 (G) passively sensitized ASF and SPF mice after PN challenge. (H) Correlation plot of Ara h 1 against alpha-diversity in small intestinal content across sensitization methods (y=-0.01375x + 1.095, R^2=0.7053, P<0.0001). n=4–10 mice per group; 2 independent experiments for each sensitization method; each dot=one mouse. P values were calculated using one-way ANOVA with Tukey's post-hoc test (B–E) and Student's t-test (F–G). See also Figures S2-S4.

Human oral cavity and small intestine harbor bacteria with the capacity to digest peanut

Food-induced allergic reactions can occur within minutes of allergen exposure. Given this rapid onset, we investigated whether humans harbor PN-degrading bacteria in the oral cavity and the small intestine. We analyzed saliva from 13 volunteers and jejunum aspirates from 5 volunteers with no reported food allergies (**Table S1**). Analysis of the oral and small intestinal microbiota composition via 16S rRNA sequencing revealed donor-specific profiles, with *Prevotella*, *Rothia,* and *Fusobacterium* as the most abundant genera in the saliva and *Streptococcus* as the dominant genus in the jejunum (**Figure 3A**). Among these taxa, *Rothia* was particularly notable, accounting for up to 43% of the salivary microbiota and as much as 14% of the small intestinal microbiota. We then plated saliva samples on PN-enriched media agar to isolate bacteria with PN-degrading capacity, selecting strains that produced a hydrolytic halo, indicating their ability to degrade PN (**Figure 3B**). PN-degrading bacteria were detected in nearly all donors, with *Rothia, Staphylococcus, Streptococcus,* and *Veillonella* being the most frequently isolated genera. Each isolated bacterial strain was then incubated with CPE to facilitate allergen digestion, and the remaining Ara h 1 and 2 were quantified. Additionally, we screened a well-characterized bacterial collection for PN-degrading potential[31]. While many bacterial strains exhibited PN digestion capabilities on agar plates, their capacity to degrade the allergens varied by taxon and strain (**Figure 3C**). *Rothia*, a dominant oral genus, consistently degraded Ara h 1 and 2 *in vitro*. This activity was observed in all tested *Rothia* strains and a phylogenetically related genus, *Micrococcus*, suggesting a conserved PN-degrading function (**Figure 3C**). In contrast, most strains from other PN-degrading genera such as *Gemella* or *Streptococcus* lacked significant allergen-degrading capacity (**Figure S5A**). While *Staphylococcus* strains showed strong PN-degrading potential, their ability to degrade PN allergens was strain-specific (**Figure 3C**). Like the oral cavity, we identified strains of *Streptococcus*, *Rothia* and *Staphylococcus* in the small intestine of healthy individuals with PN allergen-degrading capacity (**Figure S5C**). Altogether, these findings demonstrate that oral and intestinal bacteria degrade PN allergens with genus- and strain-dependent efficiency.

A   Oral and small intestinal microbial composition (genus)

B   PN-degrading *Rothia mucilaginosa*
on PN-enriched agar

C   PN-degrading bacteria

Figure 3. Human saliva harbors peanut-degrading bacteria.
(A) Genus-level relative abundance of bacteria in saliva (n=13) and jejunal aspirates (n=5) from non-peanut (PN)-allergic individuals. Taxa <1% relative abundance not shown. (B) PN-degrading bacteria (*Rothia mucilaginosa*) from human saliva on PN-agar showing zone of clearance. (C) PN-degrading bacteria isolated from human saliva, identified for whole PN-degradation by growth on PN-agar and assessed for Ara h 1 and 2 degradation. Heatmaps show allergen degradation (blue=0%, red=100% relative to crude peanut extract control) and isolate abundance (green scale, light=low, dark=high). Data presented at the genus and species levels for *Actinomycetota* (pink), *Bacillota* (grey) and *Pseudomonota* (yellow). See also Figure S5.

Peanut allergens are degraded by bacteria

To further study the microbial capacity for PN allergen digestion, we conducted proteomic analyses focusing on key bacterial taxa. We selected two primary genera: *Rothia*, a dominant oral taxon with efficient allergen-digestion capacity, and *Staphylococcus,* which has been previously associated with allergic diseases (**Figure S5B**)[32]. First, we confirmed the capacity of these taxa to degrade the main PN allergens Ara h 1 (64 kDa) and Ara h 2 (17 kDa) using SDS-PAGE (**Figure 4A–B**). To strengthen these data, we performed tandem mass spectrometry analysis on the fraction below 3 kDa, screening for PN-derived peptides after bacterial digestion. Given the low relative abundance of Ara h 2 in the PN matrix (5.9-9.3% of total protein) and the noise introduced by bacterial proteins in the samples, we centered the proteomic analyses on Ara h 1 (12-16% of total protein)[33]. We identified several Ara h 1-derived peptides following PN digestion by strains of *Rothia* and *Staphylococcus*. As expected, we quantified more peptides from bacterial strains that degraded PN more extensively (*Rothia* R3, 9 peptides; S1, 11 peptides; S3, 13 peptides), according to the electrophoretic analysis. Using molecular modelling, we visualized these peptides, marking in yellow peptides digested by bacteria (**Figure 4C–F**). Interestingly, the identified peptides shown in purple have been reported as clinically relevant IgE epitopes recognized by PN-allergic patients[34-45]. These epitopes are distributed across various regions of the 3D structure of Ara h 1, highlighting the lack of localization of the identified peptides to a single structural domain (**Figure 4C–F**). Considering the ability of these *Rothia* and *Staphylococcus* species to degrade PN, we evaluated their impact on human IgE-binding to PN after bacterial digestion by Western blotting. We used a pool of sera from PN-allergic patients with elevated levels of Ara h 1- and Ara h 2-specific IgE (**Table S2**). The analysis showed a substantial reduction in IgE-binding to PN proteins overall, and to Ara h 1 and 2 in particular (**Figure 4G–H**). To precisely define the impact of microbial metabolism on main PN allergens, we mono-sensitized mice to native Ara h 1 or recombinant Ara h 2 and used their sera to analyze IgE recognition of the digested PN fragments via Western blotting (**Figure 4I–J**). *Rothia* digestion of PN reduced IgE-binding to Ara h 1, most notably by strain R3. Similarly, *Staphylococcus* strains (S1 and S3) completely degraded and reduced IgE-binding to Ara h 1, though some residual fragments retained IgE-binding capacity. *Staphylococcus* strains with lower PN-degrading capacity, such as S2, showed less efficacy in reducing binding to Ara h 1 (**Figure 4I**). On the other hand, sera from mice sensitized with recombinant Ara h 2

predominantly recognized a band around ~37 kDa, likely representing its oligomerized

forms. Nevertheless, PN-digestion by *Rothia* strains impaired Ara h 2 recognition by IgE,

particularly with strain R3. *Staphylococcus* strains S1 and S3 also reduced IgE-binding to

Ara h 2, while strain S2 showed little effect (**Figure 4J)**. Altogether, these findings

demonstrate the potential of *Rothia* and *Staphylococcus* strains to modify the structure of

key PN allergens, including IgE-binding epitopes, thereby altering their recognition by

both murine and human IgE.

Figure 4. Peanut allergens are degraded by *Rothia* and *Staphylococcus*.
Selected *Rothia* (R1- *R. aeria,* R2- *R. dentacariosa,* and R3- *R. mucilaginosa*) and *Staphylococcus* (S1- *S. epidermidis,* S2- *S. aureus*, and S3- *S. aureus*) strains were incubated with crude peanut extract (CPE) in

liquid media to assess Ara h 1 and 2 degradation. (A–B) SDS-PAGE of Ara h 1 (64 kDa) and Ara h 2 (17
kDa) degradation by *Rothia* (A) and *Staphylococcus* (B) vs. controls. (C–F) Molecular modeling of Ara h 1
(Alphafold database code: AF-P43238-F1): peptides <3 kDa identified by mass spectrometry after digestion
by R1 (C), R3 (D), S1 (E), and S3 (F) in yellow; human IgE-recognized epitopes in purple. (G–H) Western
blot showing peanut (PN)-allergic human IgE-binding to PN allergens after *Rothia* (G) and *Staphylococcus*
(H) digestion relative to controls. (I–J) Western blot showing IgE-binding from Ara h 1 (I) or Ara h 2-
sensitized (J) mice after bacterial digestion. Bar charts show allergen quantification after bacterial digestion
(mean ± SEM). *P* values calculated using unpaired *t*-test between undigested and bacterially-digested PN.
286

Microbial metabolism mitigates peanut allergenicity

The structural modifications of Ara h 1 and 2 caused by microbial metabolism affected
IgE recognition, prompting us to assess the functional consequences using a mast cell
activation assay[46,47]. We sensitized bone marrow-derived mast cells (BMMCs) with sera
from mice allergic to either native Ara h 1 or recombinant Ara h 2 and assessed IgE-
mediated mast cell activation after challenge with bacterial PN digests. This was done by
challenging the BMMCs with either bacterially digested PN or undigested PN, and
measuring CD63 and CD107a surface expression, along with β-hexosaminidase activity
in the cell supernatants (**Figure 5 & Figure S6A**). Our initial controls confirmed that serum
from Ara h 1-allergic mice induced BMMC activation following Ara h 1 or CPE challenge,
but not upon Ara h 2 stimulation, and vice versa (**Figure S6B–C**). We then observed that,
among the three *Rothia* strains, PN-digestion by strain R2 consistently reduced BMMC
activation for both Ara h 1 and 2 (**Figure 5A–B**); while PN-digestion by strains R1 and R3
significantly decreased BMMC responses in Ara h 2-sensitized cells (**Figure 5A–B**). On
the other hand, among the three *Staphylococcus* strains, PN-digestion by strain S1
significantly reduced BMMC activation in Ara h 1-sensitized cells **(Figure 5C)**, while
BMMC responses to Ara h 2 were impaired by the digestion of the three strains (**Figure
5D**). In sum, these results demonstrate that the metabolism of key PN allergens by human
oral bacteria can modulate the ability of these allergens to trigger IgE-mediated mast cell
activation.

To further characterize the mechanisms underlying microbial allergen degradation, we
performed whole-genome sequencing of the different *Rothia* and *Staphylococcus* species
and identified candidate genes encoding protease repertoires (**Figure S7 & Figure S8**).
Notably, conserved proteases in *Rothia* include members of the peptidase S8 family of
subtilisin-like serine proteases that are secreted extracellularly, which have been
previously attributed to degrading recalcitrant dietary proteins such as gluten[48-50], and
may likewise contribute to PN allergen degradation. Protease-encoding genes were more
variable in *Staphylococcus*, with distinct, non-conserved extracellular proteases identified
in each strain. When comparing PN-degrading functionality with annotated protease
genes, we observed that potential extracellular proteases including the cysteine protease
EcpA, were present in the most efficient degraders against immunodominant allergens.
These results confirm the presence of proteolytic machinery potentially involved in PN
degradation by bacteria.

A   Ara h 1 sensitized

B   Ara h 2 sensitized

C   Ara h 1 sensitized

D   Ara h 2 sensitized

Figure 5. Challenge with bacterially digested peanut impairs mast cell activation. Mast cells (MCs) were sensitized with pooled sera from mice allergic to Ara h 1 or 2 and challenged with undigested peanut (PN; gray, positive control) or PN digested by *Rothia* (orange; R1–*R. aeria*, R2–*R. dentocariosa*, R3–*R. mucilaginosa*; A–B) or *Staphylococcus* (blue; S1–*S. epidermidis,* S2–*S. aureus,* S3–*S. aureus*; C–D). Negative controls (white) were MCs sensitized with sera from non-allergic mice and challenged with pooled bacterially-digested PN from all three *Rothia* or *Staphylococcus* species. Bar graphs indicate CD63 and CD107a expression, and β-hexosaminidase activity, normalized to 100%. Data=mean ± SEM. All three *Rothia* and *Staphylococcus* strains are displayed in the same graphs, but statistical analyses were performed separately for each strain vs. undigested PN. *P* values calculated using one-way ANOVA with Dunnett's post-hoc test, except for specific cases R1-CD63-Ara h 2 (B), S2-CD107a-Ara h 1 (C), and S1-CD107a-Ara h 2 (D), where the Kruskal-Wallis test with Dunn's correction was used. Comparisons with undigested PN used one-way ANOVA with Dunnett's test for β-hexosaminidase activity (D), CD63 (A), and CD107a (A–C), or the Kruskal-Wallis with Dunn's correction for β-hexosaminidase activity (A–C), CD63 (B–D), and CD107a (D). See also Figures S6-S8.

Microbial peanut metabolism alters systemic allergen access

To study the *in vivo* capacity of human oral bacteria to degrade PN allergens, we colonized GF C57BL/6 mice with bacterial strains exhibiting varying allergen-degrading capacities. We selected three *Staphylococcus* strains based on their *in vitro* PN-degrading capacities: one with strong capacity to degrade both Ara h 1 and 2 (S1), one that degrades only Ara h 1 (S3), and one with no detectable allergen-degrading capacity (S2). Additionally, we included a *Rothia* strain (R3) capable of degrading both Ara h 1 and 2. GF mice were used as controls. After colonization, mice were gavaged with PN (**Figure 6A**), as previously described for SPF and ASF mice. Successful transfer of PN-degrading capacity was confirmed in colonized mice (**Figure 6B**). Notably, we observed a reduction in Ara h 1 and 2 levels in the small intestinal content of mice colonized with allergen-degrading bacteria, such as S1 and R3 (**Figure 6C**). Additionally, systemic levels of Ara h 1 and 2 were reduced in these mice (**Figure 6D**). In contrast, Ara h 1—and particularly Ara h 2—showed increased systemic levels in mice colonized with PN-degrading bacteria that had impaired allergen-degrading capacity (S2) (**Figure 6D**). To study whether bacterial metabolism alters allergen passage through the mucosa and subsequent systemic access, we assessed allergen translocation across *ex vivo* mouse small intestinal tissue using Ussing chambers. CPE, pre-incubated with or without the different bacterial strains was applied to the mucosal side of the chamber, and Ara h 1 and 2 levels were quantified on the serosal side of the tissue after 2 hours. We found that PN-degrading bacteria with impaired allergen-degrading capacity facilitated the passage of Ara h 1 and 2 through the intestinal mucosa (**Figure 6E**). These findings suggest that bacteria metabolize PN *in vivo* and control allergen passage through the intestinal mucosal barrier.

Figure 6. Bacteria dictate systemic access of allergens *in vivo*.
(A) Experimental design. GF C57BL/6 mice were mono-colonized with bacteria (R3–*Rothia mucilaginosa*, S1–*Staphylococcus epidermidis*, S2–*Staphylococcus aureus*, and S3–*Staphylococcus aureus*) for 3 weeks before peanut (PN) gavage; GF mice served as controls. Mice were sacrificed 40 min after PN gavage. (B) PN-degrading capacity of small intestinal contents measured by hydrolytic halo diameter on PN-agar, and degradation of Ara h 1 and 2 (heatmap, blue=0%, red=100% relative to crude peanut extract). (C–D) Ara h 1 and 2 in small intestinal content (C) and serum (D). (E) Translocation of PN allergens digested by bacteria from the mucosal to serosal side of small intestinal tissue in Ussing chambers (R3, orange; S1, light blue; S3, dark blue, S2, medium blue; non-growth control, black). Data are shown as mean + SD (B) and mean where each dot represents an individual mouse (C, D, E). n=4–8 mice/group. *P* values calculated using a one-way ANOVA with Tukey's post-hoc test.

Microbial peanut metabolism dictates anaphylactic reactions *in vivo*

To evaluate the impact of microbial PN metabolism on anaphylactic responses, ASF C3H/HeN mice—which have impaired microbial PN metabolism and are susceptible to anaphylactic reactions upon oral exposure—were sensitized to PN via i.p. injection and subsequently challenged with bacterially pre-digested PN (**Figure 7A**). Four groups of mice received an i.g. PN challenge with either untreated PN (control) or PN pre-digested by one of the following bacteria: a) *Rothia* R3, b) *Staphylococcus* S1, or c) *Staphylococcus* S2, each exhibiting distinct allergen-degrading capacities. While all groups showed similar PN-specific IgE levels after sensitization (**Figure 7B**), serum mMCP-1 levels were lower in mice challenged with microbially pre-digested PN (**Figure 7C**). Notably, mice challenged with PN pre-digested by *Staphylococcus* S2, which has limited allergen-reducing capacity, exhibited higher serum levels of mMCP-1. To evaluate the protective capacity of *Rothia* colonization against food-induced anaphylaxis, ASF C3H/HeN mice were sensitized to PN (i.g.) and then colonized with *Rothia* R3 with efficient PN-degrading capacity (**Figure 7D**). After sensitization, ASF mice with and without PN-degrading bacteria were challenged to PN i.g. Despite having similar PN-specific IgE levels in serum and small intestinal mucosal mast cells (**Figure 7E & Figure S9A–E**), mice colonized with *Rothia* R3 (capable of digesting PN allergens) demonstrated significantly reduced serum levels of mMCP-1 (**Figure 7F**) and PAF-AH (**Figure S9B**). Furthermore, PN challenge in ASF mice colonized with or without *Rothia* after i.p. sensitization led to reduced mast cell degranulation in *Rothia*-colonized mice despite similar PN-specific IgE levels and mucosal mast cells (**Figure S9F–J**). Overall, these findings highlight the critical role of microbial allergen metabolism in modulating allergic responses *in vivo*.

Peanut-degrading bacteria are increased in allergic patients better tolerating peanut

To investigate the role of the oral microbiota in allergic reactions, we collected saliva samples from a small well-defined cohort of PN-allergic patients (**Figure 7G & Table S1**). Prior to initiating oral immunotherapy (OIT), allergic patients underwent a PN challenge to determine their threshold for reactivity (PN threshold). We analyzed serum levels of IgE and the composition of the oral microbiota. Notably, serum levels of IgE specific to Ara h 1 and 2 showed a poor association with PN threshold (**Figure S10A**), suggesting the role of additional factors. Examining the oral microbiota, we observed non-significant clustering based on PN threshold (**Figure S10B**). While PN-degrading taxa were present across patients with varying PN thresholds, individuals with higher PN thresholds exhibited a greater abundance of taxa capable of degrading Ara h 2 (**Figure 7H**). Notably, *Micrococcales*—a taxon that includes *Rothia* and *Micrococcus*, both efficient at degrading immunodominant Ara h 1 and 2—was reduced in PN-allergic patients with low PN threshold (0–40 mg), independent of serum Ara h 2-specific IgE levels (**Figure 7I**). Interestingly, some high-threshold patients with elevated Ara h 2-specific IgE had

increased *Micrococcales*, potentially implicating non-IgE factors in reactivity. To validate
our clinical findings, we examined the relative abundance of *Rothia* in saliva of PN-allergic
and non-allergic children in an external cohort of 120 children who underwent double-
blind, placebo-controlled food challenges[51]. The participants included 23 children with no
allergy (controls), 74 with high-threshold PN allergy (reacting to ≥ 443 mg cumulative PN
protein), and 23 with low-threshold PN allergy (reacting to <443 mg cumulative PN
protein). Notably, *Rothia aeria* ASV14171 was significantly more abundant in the saliva
of non-PN-allergic and high PN threshold allergic individuals compared to allergic
individuals with low PN thresholds (**Figure 7J**). Other *Rothia* species also showed a trend
toward enrichment in the groups (**Figure S10C**). These findings suggest that the oral
microbiota could serve as a predictive marker of threshold reactivity to PN, highlighting
the potential importance of microbial allergen metabolism in IgE-mediated reactions.

A PN challenge with bacterially digested PN

PN sensitization (i.p.)

PN challenge (i.g.) Digestions or Control

4 weeks | 2 weeks | 40 min | Endpoint

SPF C3H-HeN (n=5)

__Bacterial CPE digestions for challenge__
- **1. Control:** Non-digested CPE
- **2. *Rothia* R3:** Degrades Ara h 1 & 2
- **3. *Staphylococcus* S1:** Degrades Ara h 1 & 2
- **4. *Staphylococcus* S2:** Degrades PN but not Ara h 1 & 2

B PN-specific IgE

PN-IgE (Relative Binding, OD)

15 — 10 — 5 — 0

C mMCP-1

0.025
0.0007
0.0012

mMCP-1 (ng/mL)

200 — 150 — 100 — 50 — 0

D PN challenge in mice colonized with PN-degrading bacteria

PN sensitization (i.g.)

PN challenge (i.g.)

6 weeks | 3 weeks | 40 min | Endpoint

Colonization

ASF C3H-HeN (n=7)

__Colonization__
1. Control (vehicle)
2. *Rothia* R3

E PN-specific IgE

PN-IgE (Relative Binding, OD)

80 — 60 — 40 — 20 — 0

ASF Control | ASF *Rothia*

F mMCP-1

0.0037

mMCP-1 (ng/mL)

800 — 600 — 400 — 200 — 0

ASF Control | ASF *Rothia*

G PN threshold assessment

PN dose escalation

Baseline threshold tolerance
Low (0–40 mg)
High (>40 mg)

PN-allergic patients (n=19)

H PN-degrading taxa abundance based on PN threshold

PN-degrading

% Relative abundance

60 — 50 — 40 — 30 — 20 — 10 — 0

Low (0–40) | High (>40)
PN threshold (mg)

Ara h 2-degrading

% Relative abundance

6 — 5 — 4 — 3 — 2 — 1 — 0

Low (0–40) | High (>40)
PN threshold (mg)

- *Micrococcales*
- *Actinomyces*
- *Staphylococcus*
- *Veillonella*
- *Gemella*
- *Granulicatella*
- *Streptococcus*

I PN-degrading taxa in PN-allergic patients

Micrococcales

% Relative abundance

12 — 10 — 8 — 6 — 2.5 — 2.0 — 1.5 — 1.0 — 0.5 — 0.0

High Ara h 2 IgE
Low Ara h 2 IgE

Low (0–40) | High (>40)
PN threshold (mg)

J External cohort: *Rothia aeria* abundance in PN-allergic and non-allergic individuals

Not PN allergic (NPA) n=23

High PN threshold (HT) n=74

Low PN threshold (LT) n=23

Rothia aeria
ASV 14171

% Relative abundance

1.5 — 1.0 — 0.5 — 0.0

0.0031
0.0132

NPA | HT | LT

DISCUSSION

The human microbiome is essential for maintaining homeostasis, and disruptions in its
composition are associated with immune disorders. The oro-gastrointestinal microbiota
has gained attention for its role in influencing oral tolerance and sensitization to food
antigens[18-20,22,52]. While studies have reported differences in the intestinal microbiota
composition of food-allergic patients[23-25], the microbial mechanisms underlying food-
induced anaphylaxis remain largely unknown. In this study, we show that human saliva
and small intestinal content harbor allergen-degrading bacteria capable of metabolizing
immunodominant PN allergens, thereby modulating IgE-mediated reactions to foods.
Identifying the microbes involved in PN metabolism in humans and characterizing the
related microbially-mediated IgE-specific immune responses could have implications for
reducing the severity of allergic reactions.

The antigenic properties of many foods stem from their resistance to gastrointestinal
digestion[28,29]. PN allergens, for example, are poorly metabolized by human digestive
enzymes[26,27], whereas the intestinal microbiota encodes diverse metabolic pathways
absent in the host that participate in food digestion[53]. For example, the capacity of the
intestinal microbiota to degrade other recalcitrant antigenic proteins, such as gluten[54], has
been demonstrated[55,56]. Using gnotobiotic mouse models, we demonstrate that
microbiota composition and complexity contribute to PN allergen metabolism *in vivo* and
the amount of allergen reaching the intestine and circulation. While the intestinal
microbiota affects host metabolism and food-dependent immune responses through
mechanisms such as altering the intestinal barrier, motility, or immune activation[18,57-59],
we confirmed the presence of PN-degrading bacteria capable of digesting Ara h 1 and 2
along the gastrointestinal tract in wild-type SPF mice. These findings reveal the role of

the intestinal microbiota in allergen metabolism and raise important questions about how diet-microbiota interactions shape food allergy.

Food allergies are primarily IgE-mediated[60], and even minute allergen doses can trigger life-threatening reactions [61]. We hypothesized that microbial metabolism modulates IgE-dependent responses to PN. First, we demonstrated that mice with limited PN-degrading capacity (ASF) developed enhanced allergic reactions to PN than those with efficient PN metabolism (SPF) following i.g. sensitization and challenge. ASF mice exhibited greater mucosal mast cell activation, likely increasing intestinal permeability and facilitating systemic allergen access[62,63]—a critical step in IgE-mediated anaphylaxis[64]. Intriguingly, ASF mice also developed higher titers of antigen-specific IgE than SPF controls, potentially due to their reduced ability to metabolize PN allergens. Similar increases in antigen-specific IgE have been reported in GF mice and mice colonized with limited microbial communities following sensitization[65-67]. Given that food-induced anaphylaxis is largely IgE-dependent[68,69], and the microbiota may influence IgE-mediated food sensitization independently of allergen degradation[18], differences in sensitization between mouse strains could explain the severity of allergic reactions. To bypass local microbial factors, we evaluated the role of microbial metabolism in IgE-mediated food responses using additional models of systemic[70] and passive[47,71] sensitization. Following PN challenge, ASF mice with limited PN-degrading capacity developed enhanced allergic reactions, characterized by increased mast cell degranulation and hypothermia compared to conventional mice, despite similar PN-IgE titers. Altogether, these results indicate that microbial metabolism is a key determinant of IgE-mediated food allergy in mice.

Food-induced anaphylaxis usually results from accidental allergen exposure, posing a persistent threat that significantly impacts quality of life[72,73]. Although co-factors such as exercise, estrogen levels, alcohol, and drugs can heighten anaphylaxis risk[14,74], variations in allergen thresholds and susceptibility remain poorly understood. A recent study demonstrated that anaphylaxis severity is independent of the amount of allergen ingested, emphasizing individual sensitivity and the potential role of microbiota in modulating IgE-responses[75]. To explore the potential role of microbial allergen metabolism in humans, we searched for PN-degrading bacteria in healthy donors, focusing on the oral cavity and small intestine where ingested food first interacts with the microbiota. While most microbiota research has centered on the large intestinal lumen due to its ease of sampling and high bacterial density, recent studies have highlighted the stability and complexity of the oral and small intestinal microbiota, as well as their relevance in health and disease[76]. Moreover, the oral microbiota serves as a continuous source of bacteria for the intestinal tract, seeding the small intestine with taxa such as *Rothia* and *Staphylococcus*[77], which may persist and contribute to allergen metabolism as food moves through the upper gut[78]. Using SDS-PAGE, Western blotting, and proteomics, we confirmed the presence of PN-degrading bacteria in the oral cavity with strain-specific capacities to degrade immunodominant PN allergens, disrupt human IgE epitopes, and alter IgE-binding of sera from PN-allergic patients and sensitized mice.

Overall, these results confirm the capacity of bacteria to potentially alter allergenicity in the oral cavity and small intestine of humans.

Dominant taxa in the oral cavity and small intestine such as *Rothia*, and related genera like *Micrococcus*, consistently digested both Ara h 1 and 2 *in vitro,* altering IgE recognition. The capacity of *Rothia* to degrade other dietary antigens such as gluten proteins has been also suggested[50,79]. Remarkably, impaired IgE recognition of PN upon *Rothia* digestion hampered the activation of the classical pathway of anaphylaxis both *in vitro* (mast cell assays) and *in vivo* (PN-allergic mice challenged with digested PN). In addition, the PN-degrading capacity of *Rothia* was validated *in vivo*, reducing the concentration of major PN allergens reaching the small intestine and circulation. We also tested the capacity of *Rothia* strains to alter IgE-mediated reactions following colonization in a competitive environment. To avoid changes in the adaptive immune response, ASF mice were colonized following either i.g. or i.p. sensitization. ASF mice colonized with *Rothia* had reduced IgE-mediated allergic responses and mucosal mast cell degranulation following challenge compared to ASF controls. These results confirm that the protective effects of allergen metabolism arise by reducing allergen availability in the intestinal environment rather than by altering systemic sensitization. Altogether, our data demonstrate that PN-degrading bacteria can modulate acute allergic responses *in vivo*.

Microbial PN metabolism does not always protect against IgE-mediated responses. Certain PN-degrading genera, such as *Gemella* or *Streptococcus,* lacked allergen-degrading capacity *in vitro*, while *Staphylococcus*, a main PN-degrading taxon in the oral cavity*,* displayed strain-specific effects in our work. Others have reported that skin colonization by *Staphylococcus aureus* induces keratinocyte release of IL-36α[80], a pro-inflammatory cytokine linked to allergic sensitization to foods[32]. Additionally, *Staphylococcus* and other microbial proteases such as EcpA can induce immune activation and low-grade inflammation in the colon[81], and promote inflammatory responses to innocuous antigens[82,83]. Our findings revealed that *Staphylococcus* strains vary in their capacity to remove PN allergens and alter IgE-mediated recognition and mast cell activation *in vitro*. In gnotobiotic mice colonized with *Staphylococcus* strains exhibiting differing PN-degrading capacities, efficient strains reduced Ara h 1 and 2 levels in the small intestine and serum. However, these immunodominant peptides increased systemically in mice colonized with a strain lacking efficient PN-degrading capacity. Enhanced passage of Ara h 1 and 2 through the intestinal mucosa after partial digestion of PN by this *Staphylococcus* strain was also confirmed with Ussing chambers, highlighting the role of bacterial activity in systemic allergen access, a critical step in anaphylactic reactions[64]. These results indicate that *Staphylococcus* species, including *Staphylococcus aureus,* may facilitate allergen passage through the mucosal barrier and promote allergic inflammation when allergen degradation is incomplete[84,85]. Some *Staphylococcus* strains can also compromise the epithelial barrier and increase permeability directly[86,87]. Therefore, microbes may have a dual impact on IgE-mediated immune responses, depending on the bacteria's ability to degrade PN allergens, regardless of its taxonomy. Complete microbial degradation of PN allergens reduces IgE

responses, while inefficient metabolism may promote anaphylaxis by enhancing the passage of allergens through the mucosa. Altogether, the ability of the microbiota to completely degrade immunodominant allergens and reduce IgE recognition of PN significantly influences subsequent mast cell activation and anaphylactic reactions.

Despite these findings, the precise microbial proteases responsible for PN allergen metabolism remain to be fully identified. Whole-genome sequencing of different *Rothia* species revealed diverse protease repertoires, including subtilisin-like serine proteases. Members of this family have been shown to degrade other recalcitrant dietary proteins[48-50], such as gluten, suggesting they may contribute to PN allergen metabolism. Efficient PN-degrading *Staphylococcus* possessed annotated extracellular proteases, including the virulence-associated cysteine protease EcpA[88,89]. However, given the broad proteolytic capacity of the oral microbiota, multiple enzymes could be involved in modulating allergenicity. Future studies integrating genetic modulation will be required to pinpoint the specific microbial enzymes mediating these effects.

While our findings consistently support the role of microbial metabolism in modulating PN allergenicity, variability was observed in our results across models and microbial communities. Both technical (*e.g.*, antibody and *ex vivo* assay variability) and biological factors (*e.g.*, postprandial state, vendor-specific SPF microbiota) likely contributed. We used i.g. challenges to focus on mucosal mast cell activation, though these induce weaker clinical anaphylaxis than the i.p. route and add variability. Specifically, immune responses upon i.g. challenge are influenced by factors including allergen delivery, intestinal permeability, gastric emptying, and genetic background. This variability is further amplified after i.g. sensitization that also enhances mast cell recruitment to the small intestinal mucosa and intestinal permeability[90]. GF mice, the gold standard for studying microbially-mediated phenotypes, were used to investigate PN metabolism, but they exhibit altered intestinal barrier function, impaired motility, and increased proteolytic activity, and their underdeveloped immunity can exaggerate sensitization and anaphylaxis[65,67,91-93]. Importantly, sensitization was not the primary focus of this study and allergic responses may be influenced by factors beyond microbial[22], such as the extent of mast cell recruitment to the intestine[90]. To better isolate microbial effects, we used ASF mice, which provide minimal but stable microbiota with limited endogenous PN-degrading activity. ASF mice serve as a stronger comparator for high PN-degrading capacity microbiota or as a model for supplementation with PN-degrading taxa. Colonization of ASF mice with PN-degrading *Rothia* after i.g. or i.p. sensitization to PN reduced anaphylaxis markers and mast cell degranulation. Microbial composition and complexity were key determinants of allergen degradation and IgE-mediated responses, as shown by a negative correlation between alpha-diversity and both intestinal allergen levels and serum mMCP-1 in ASF and SPF mice. This suggests that variability in PN degradation and anaphylaxis observed in our study may be partially attributed to individual differences in microbiota composition and complexity, with some SPF mice potentially harboring fewer PN-degrading bacteria.

The primary recommendation for food allergic patients is strict avoidance of allergens, yet this is a challenge for ubiquitous foods such as egg, wheat, milk or PN, leading to frequent accidental exposures[10,12,94,95]. Allergen thresholds differ among PN-allergic patients, with some individuals experiencing anaphylaxis to trace amounts[96,97]. To investigate the role of microbial allergen metabolism in food-induced anaphylaxis, we characterized the oral microbiota of PN-allergic patients initiating OIT with varying allergen thresholds. Allergen-specific IgE did not correlate well with allergen thresholds in our study, consistent with previous findings[98,99]. However, patients with higher PN thresholds showed greater abundance of *Micrococcales* (an order including PN-degrading *Rothia* and *Micrococcus*), than highly sensitive allergic patients to PN, independent of IgE titers. Consistent with this observation, our analysis of an external pediatric cohort revealed that *Rothia aeria* was significantly more abundant in the saliva of non-PN-allergic and high-threshold PN-allergic individuals compared to allergic individuals with low thresholds, with other *Rothia* species also showing a trend toward enrichment[51]. A previous study also reported an increased relative abundance of *Rothia* in the saliva of PN-allergic children after undergoing OIT[100], and *Veillonella*, an oral taxon with PN-degrading capacity, is increased in those with greater reaction thresholds[101]. Together, these findings suggest that allergen threshold associates with the metabolic activity of oral microbiota against food allergens, and may serve as a marker of reactivity. Still, the mechanistic link between oral microbiota and allergen threshold in humans remains speculative, requiring preclinical and longitudinal validation. Longitudinal assessment of oral microbiota and allergen reactivity during PN OIT in future studies will further inform how microbiota may influence OIT outcomes. Although OIT remains the only allergen-specific treatment with disease-modifying potential, it is associated with limitations such as events that compromise patient adherence, quality of life, and overall effectiveness[73,102,103]. Functional analysis of the oral microbiota may help identify patients with low allergen thresholds less suitable for OIT, while harnessing PN-degrading bacteria may raise allergen thresholds. Characterizing PN-degrading bacteria could also be applied to reduce cross-contamination risks—a leading cause of unintentional allergen exposure and a significant concern in food allergy management[95,104-106]—and improve the formulation of hypoallergenic products. In summary, our findings reveal a novel microbial mechanism of relevance in food-induced anaphylaxis. Our results underscore the role of the human microbiota in dictating the severity of IgE-mediated reactions through allergen metabolism and highlight the therapeutic potential of harnessing bacterial allergen-degrading capabilities for managing food allergies.

RESOURCE AVAILABILITY

Lead contact: Further information and requests for resources and reagents should be directed to the lead contact Alberto Caminero (acamine@mcmaster.ca).

Materials availability: This study did not generate new unique reagents.

Data and code availability: The microbiome sequencing data generated in this study have
been deposited in the NCBI BioProject database under the accession number
PRJNA1223483. The mass spectrometry proteomics data have been deposited to the
ProteomeXchange Consortium via the PRIDE partner repository with the dataset
identifier PXD060888 and 10.6019/PXD060888. Any additional information required to
reanalyze the data reported in this paper is available from the lead contact upon
reasonable request.

ACKNOWLEDGMENTS

We thank the McMaster University AGU staff, Dr. Heather G. Galipeau, Joe Noratangelo,
Michael Rosati, and Sarah Armstrong for their assistance with mouse care. We thank the
McMaster Genome Facility, Laura Rosati, and Michelle Shah for technical support with
16S rRNA gene sequencing. We thank Gaston Rueda for assistance with clinical
samples. We thank the staff of the Proteomics Unit at CNB-CSIC, particularly Dr. Alberto
Paradela, for analysis and technical support. We are grateful to Drs. Pablo Rodríguez del
Río, Salvador Iborra and Francisco Sánchez-Madrid for their critical review of the
manuscript. This work was supported by grants from The Nutricia Research Foundation
(NRF-2021-13) and the New Frontiers in Research Fund (NFRFE-2019-00083) to R.J.-
S. and A.C.; and by a grant from the European Food Safety Authority
(GP/EFSA/ENCO/2020/02-1) to F.J.M. R.J.-S.´s laboratory is supported by the Instituto
de Salud Carlos III (ISCIII; CP20/00043; PI22/00236; PI25/00477; RD24/0007/0037) and
by the Ministerio de Ciencia, Innovación y Universidades (CNS2024-154194). A.C. holds
a Family Douglas Research Chair in Intestinal Research and is funded by the Canadian
Institutes of Health Research (CIHR; Project Grant 202010PJT), NSERC (DGECR-2022-
00221) and by Crohn's and Colitis Canada (CCC, Grants-In-Aid grant). L.E.R. is
supported by a Frederick Banting and Charles Best Canada Graduate Scholarship-
Doctoral award from CIHR and a graduate scholarship from the Farncombe Institute.
E.S.-M. is supported by a Formación de Profesorado Universitario grant (FPU23/03341)
from Ministerio de Universidades. S.U.P. is supported by NIAID NIH R01 grants
1R01AI155630 and 1R01AI182001. L.M.-S. is supported by the Investigo program
through the Ministerio de Trabajo y Economía Social, Servicio Público de Empleo
(SEPE), which is funded by "Plan de Recuperación, Transformación y Resiliencia" and
"NextGenerationEU" of the European Union (2022- C23.I01). C.L.S and E.N.-B are
supported by a Río Hortega (CM23/0019) and a Sara Borrell grant (CD23/00125),
respectively, from ISCIII. E.F.V. is supported by a CIHR Project Grant (PJT-183881) and
holds a Tier 1 Canada Research Chair in Microbial Therapeutics and Nutrition in
Gastroenterology. L.Z. and S.B. received funding from NIAID NIH R01 AI147028 and
U196053.

AUTHOR CONTRIBUTIONS

Conceptualization, R.J.-S., A.C.; Methodology, R.J.-S., A.C.; Investigation, E.S.-M., L.E.R., M.G.-R., D.H., P.H., B.B., G.Y., R.D., J.F.E.K., T.D.W., L.M.-S., C.L.S., E.N.-B., X.Y.W., G.D.P., L.Z., F.J.M., R.J.-S., A.C.; Resources, M.G.-A., A.D.-P., E.F.V., Y.R.C., P.O., F.V., C.B., J.F.E.K, M.J., W.G.S., S.U.P., F.J.M., M.G.S., S.B., P.B.; Writing – Original Draft Preparation, E.S.-M., L.E.R., R.J.-S., A.C.; Writing – Review & Editing Preparation, E.S.-M., L.E.R., R.J.-S., A.C.; Visualization Preparation, E.S.-M., L.E.R.; Supervision, R.J.-S., A.C.; funding Acquisition, R.J.-S., A.C.

DECLARATION OF INTERESTS

R.J.-S. receives research funds from a collaboration agreement between FIB-Hospital Universitario de La Princesa and Inmunotek S.L., which does not present any conflicts with the data presented. Provisional patent (63/670,312) was filed in the United States Patent and Trademark office based on data presented in this manuscript ("Microbial metabolism of peanut allergens and uses thereof).

SUPPLEMENTAL INFORMATION INDEX

Figures S1-S10 and their legends, as well as Tables S1 and S2 are in a PDF.

References

1. Meizlish, M.L., Franklin, R.A., Zhou, X., and Medzhitov, R. (2021). Tissue Homeostasis and Inflammation. Annu Rev Immunol *39*, 557-581. 10.1146/annurev-immunol-061020-053734.

2. Nunez-Borque, E., Dribin, T.E., Rodriguez Del Rio, P., Camargo, C.A., Jr., Esteban, V., du Toit, G., Jimenez-Saiz, R., and Giovannini, M. (2025). Anaphylaxis: Spotlight on Inflammation. Clin Exp Allergy *55*, 8-10. 10.1111/cea.14610.

3. Bachtel, N.D., Cullen, J.L., Liu, M., Erickson, S.A., Kutyavin, V.I., El-Naccache, D.W., Florsheim, E.B., Lim, J., Sullivan, Z.A., Imaeda, R., et al. (2025). Intestinal mast cell-derived leukotrienes mediate the anaphylactic response to ingested antigens. Science *389*, eadp0246. 10.1126/science.adp0246.

4. Hoyt, L.R., Liu, E., Olson, E.C., Jacobsen, D.R., Siniscalco, E.R., Krier-Burris, R.A., Greenfield, K.G., McBride, C.D., Alfajaro, M.M., Amat, J.A.R., et al. (2025). Cysteinyl leukotrienes stimulate gut absorption of food allergens to promote anaphylaxis in mice. Science *389*, eadp0240. 10.1126/science.adp0240.

5. Dyer, A.A., Rivkina, V., Perumal, D., Smeltzer, B.M., Smith, B.M., and Gupta, R.S. (2015). Epidemiology of childhood peanut allergy. Allergy Asthma Proc *36*, 58-64. 10.2500/aap.2015.36.3819.

6. Cianferoni, A., and Muraro, A. (2012). Food-induced anaphylaxis. Immunol Allergy Clin North Am *32*, 165-195. 10.1016/j.iac.2011.10.002.

7. Muraro, A., Sublett, J.W., Haselkorn, T., Nilsson, C., and Casale, T.B. (2021). Incidence of anaphylaxis and accidental peanut exposure: A systematic review. Clin Transl Allergy *11*, e12064. 10.1002/clt2.12064.

8. Leung, A.S.Y., Tham, E.H., Pacharn, P., Xing, Y., Trinh, H.K.T., Lee, S., Ahn, K., Chatchatee, P., Sato, S., Ebisawa, M., et al. (2024). Disparities in pediatric anaphylaxis triggers and management across Asia. Allergy *79*, 1317-1328. 10.1111/all.16098.

9. Neuman-Sunshine, D.L., Eckman, J.A., Keet, C.A., Matsui, E.C., Peng, R.D., Lenehan, P.J., and Wood, R.A. (2012). The natural history of persistent peanut allergy. Ann Allergy Asthma Immunol *108*, 326-331 e323. 10.1016/j.anai.2011.11.010.

10. Cherkaoui, S., Ben-Shoshan, M., Alizadehfar, R., Asai, Y., Chan, E., Cheuk, S., Shand, G., St-Pierre, Y., Harada, L., Allen, M., and Clarke, A. (2015). Accidental exposures to peanut in a large cohort of Canadian children with peanut allergy. Clin Transl Allergy *5*, 16. 10.1186/s13601-015-0055-x.

11. Peters, R.L., Guarnieri, I., Tang, M.L.K., Lowe, A.J., Dharmage, S.C., Perrett, K.P., Gurrin, L.C., and Koplin, J.J. (2022). The natural history of peanut and egg allergy in children up to age 6 years in the HealthNuts population-based longitudinal study. J Allergy Clin Immunol *150*, 657-665 e613. 10.1016/j.jaci.2022.04.008.

12. Vasse, M., Moyart, M., Cisterne, C., Thumerelle, C., Drumez, E., Behal, H., Pouessel, G., Lejeune, S., and Deschildre, A. (2025). Children Allergic to Peanut and/or Tree Nuts Are at Risk of Accidental Allergic Reactions. Clin Exp Allergy *n/a*. 10.1111/cea.70132.

13. Benede, S., Garrido-Arandia, M., Martin-Pedraza, L., Bueno, C., Diaz-Perales, A., and Villalba, M. (2017). Multifactorial Modulation of Food-Induced Anaphylaxis. Front Immunol *8*, 552. 10.3389/fimmu.2017.00552.

14. Munoz-Cano, R., Pascal, M., Araujo, G., Goikoetxea, M.J., Valero, A.L., Picado, C., and Bartra, J. (2017). Mechanisms, Cofactors, and Augmenting Factors Involved in Anaphylaxis. Front Immunol *8*, 1193. 10.3389/fimmu.2017.01193.

15. Flinn, A., and Hourihane, J.O. (2013). Allergic reaction to peanuts: can we predict reaction severity in the wild? Curr Allergy Asthma Rep *13*, 645-650. 10.1007/s11882-013-0369-5.

16. Petek, T., Lajhar, M., Krasovec, B., Homsak, M., Kavalar, M., Korosec, P., Koren, B., Tomazin, M., Hojnik, T., and Berce, V. (2023). Risk Factors for Anaphylaxis in Children Allergic to Peanuts. Medicina (Kaunas) *59*. 10.3390/medicina59061037.

17. Turner, P.J., Baumert, J.L., Beyer, K., Boyle, R.J., Chan, C.H., Clark, A.T., Crevel, R.W., DunnGalvin, A., Fernandez-Rivas, M., Gowland, M.H., et al. (2016). Can we identify patients at risk of life-threatening allergic reactions to food? Allergy *71*, 1241-1255. 10.1111/all.12924.

18. Caminero, A., Meisel, M., Jabri, B., and Verdu, E.F. (2019). Mechanisms by which gut microorganisms influence food sensitivities. Nat Rev Gastroenterol Hepatol *16*, 7-18. 10.1038/s41575-018-0064-z.

19. Stephen-Victor, E., Crestani, E., and Chatila, T.A. (2020). Dietary and Microbial Determinants in Food Allergy. Immunity *53*, 277-289. 10.1016/j.immuni.2020.07.025.

20. Cerovic, V., Pabst, O., and Mowat, A.M. (2025). The renaissance of oral tolerance: merging tradition and new insights. Nat Rev Immunol *25*, 42-56. 10.1038/s41577-024-01077-7.

21. Stephen-Victor, E., Kuziel, G.A., Martinez-Blanco, M., Jugder, B.E., Benamar, M., Wang, Z., Chen, Q., Lozano, G.L., Abdel-Gadir, A., Cui, Y., et al. (2025). RELMbeta sets the threshold for microbiome-dependent oral tolerance. Nature *638*, 760-768. 10.1038/s41586-024-08440-7.

22. Caminero, A., Sanchez-Martinez, E., Wulczynski, M., Moreno-Serna, L., Verdu, E.F., and Jimenez-Saiz, R. (2025). Potential role of microbes in antigen-based food sensitivities. Semin Immunol *79*, 101982. 10.1016/j.smim.2025.101982.

23. Azad, M.B., Konya, T., Guttman, D.S., Field, C.J., Sears, M.R., HayGlass, K.T., Mandhane, P.J., Turvey, S.E., Subbarao, P., Becker, A.B., et al. (2015). Infant gut microbiota and food sensitization: associations in the first year of life. Clin Exp Allergy *45*, 632-643. 10.1111/cea.12487.

24. De Filippis, F., Paparo, L., Nocerino, R., Della Gatta, G., Carucci, L., Russo, R., Pasolli, E., Ercolini, D., and Berni Canani, R. (2021). Specific gut microbiome signatures and the associated pro-inflamatory functions are linked to pediatric allergy and acquisition of immune tolerance. Nature Communications *12*. 10.1038/s41467-021-26266-z.

25. Bao, R., Hesser, L.A., He, Z., Zhou, X., Nadeau, K.C., and Nagler, C.R. (2021). Fecal microbiome and metabolome differ in healthy and food-allergic twins. J Clin Invest *131*. 10.1172/JCI141935.

26. Koppelman, S.J., Hefle, S.L., Taylor, S.L., and de Jong, G.A. (2010). Digestion of peanut allergens Ara h 1, Ara h 2, Ara h 3, and Ara h 6: a comparative in vitro

study and partial characterization of digestion-resistant peptides. Mol Nutr Food Res *54*, 1711-1721. 10.1002/mnfr.201000011.

27. Kopper, R.A., Odum, N.J., Sen, M., Helm, R.M., Steve Stanley, J., and Wesley Burks, A. (2004). Peanut protein allergens: gastric digestion is carried out exclusively by pepsin. J Allergy Clin Immunol *114*, 614-618. 10.1016/j.jaci.2004.05.012.

28. Jimenez-Saiz, R., Benede, S., Molina, E., and Lopez-Exposito, I. (2015). Effect of processing technologies on the allergenicity of food products. Critical reviews in food science and nutrition *55*, 1902-1917. 10.1080/10408398.2012.736435.

29. Moreno, F.J. (2007). Gastrointestinal digestion of food allergens: effect on their allergenicity. Biomed Pharmacother *61*, 50-60. 10.1016/j.biopha.2006.10.005.

30. Gertie, J.A., Zhang, B., Liu, E.G., Hoyt, L.R., Yin, X., Xu, L., Long, L.L., Soldatenko, A., Gowthaman, U., Williams, A., and Eisenbarth, S.C. (2022). Oral anaphylaxis to peanut in a mouse model is associated with gut permeability but not with Tlr4 or Dock8 mutations. J Allergy Clin Immunol *149*, 262-274. 10.1016/j.jaci.2021.05.015.

31. Whelan, F.J., Waddell, B., Syed, S.A., Shekarriz, S., Rabin, H.R., Parkins, M.D., and Surette, M.G. (2020). Culture-enriched metagenomic sequencing enables in-depth profiling of the cystic fibrosis lung microbiota. Nat Microbiol *5*, 379-390. 10.1038/s41564-019-0643-y.

32. Tsilochristou, O., du Toit, G., Sayre, P.H., Roberts, G., Lawson, K., Sever, M.L., Bahnson, H.T., Radulovic, S., Basting, M., Plaut, M., et al. (2019). Association of Staphylococcus aureus colonization with food allergy occurs independently of eczema severity. J Allergy Clin Immunol *144*, 494-503. 10.1016/j.jaci.2019.04.025.

33. Koppelman, S.J., Vlooswijk, R.A., Knippels, L.M., Hessing, M., Knol, E.F., van Reijsen, F.C., and Bruijnzeel-Koomen, C.A. (2001). Quantification of major peanut allergens Ara h 1 and Ara h 2 in the peanut varieties Runner, Spanish, Virginia, and Valencia, bred in different parts of the world. Allergy *56*, 132-137. 10.1034/j.1398-9995.2001.056002132.x.

34. Burks, A.W., Shin, D., Cockrell, G., Stanley, J.S., Helm, R.M., and Bannon, G.A. (1997). Mapping and mutational analysis of the IgE-binding epitopes on Ara h 1, a legume vicilin protein and a major allergen in peanut hypersensitivity. Eur J Biochem *245*, 334-339. 10.1111/j.1432-1033.1997.t01-1-00334.x.

35. Shin, D.S., Compadre, C.M., Maleki, S.J., Kopper, R.A., Sampson, H., Huang, S.K., Burks, A.W., and Bannon, G.A. (1998). Biochemical and structural analysis of the IgE binding sites on ara h1, an abundant and highly allergenic peanut protein. J Biol Chem *273*, 13753-13759. 10.1074/jbc.273.22.13753.

36. Jiang, S., Wang, S., Sun, Y., Zhou, Z., and Wang, G. (2011). Molecular characterization of major allergens Ara h 1, 2, 3 in peanut seed. Plant Cell Rep *30*, 1135-1143. 10.1007/s00299-011-1022-1.

37. Cong†, Y., Lou†, F., Xue, W., Li, L., and Chena, M. (2008). Characterisation of the IgE-binding immunodominant epitopes on Ara h1. Food and Agricultural Immunology *19*, 175-185. 10.1080/09540100802172599.

38. Bogh, K.L., Nielsen, H., Madsen, C.B., Mills, E.N., Rigby, N., Eiwegger, T., Szepfalusi, Z., and Roggen, E.L. (2012). IgE epitopes of intact and digested Ara

h 1: a comparative study in humans and rats. Molecular immunology *51*, 337-346. 10.1016/j.molimm.2012.04.002.

39. Bogh, K.L., Nielsen, H., Eiwegger, T., Madsen, C.B., Mills, E.N., Rigby, N.M., Szepfalusi, Z., and Roggen, E.L. (2014). IgE versus IgG4 epitopes of the peanut allergen Ara h 1 in patients with severe allergy. Molecular immunology *58*, 169-176. 10.1016/j.molimm.2013.11.014.

40. Hansen, C.S., Dufva, M., Bogh, K.L., Sullivan, E., Patel, J., Eiwegger, T., Szepfalusi, Z., Nielsen, M., and Christiansen, A. (2016). Linear epitope mapping of peanut allergens demonstrates individualized and persistent antibody-binding patterns. J Allergy Clin Immunol *138*, 1728-1730. 10.1016/j.jaci.2016.06.019.

41. Shreffler, W.G., Beyer, K., Chu, T.H., Burks, A.W., and Sampson, H.A. (2004). Microarray immunoassay: association of clinical history, in vitro IgE function, and heterogeneity of allergenic peanut epitopes. J Allergy Clin Immunol *113*, 776-782. 10.1016/j.jaci.2003.12.588.

42. Rambo, I.M., Kronfel, C.M., Rivers, A.R., Swientoniewski, L.T., McBride, J.K., Cheng, H., Simon, R.J., Ryan, R., Tilles, S.A., Nesbit, J.B., et al. (2023). IgE and IgG4 epitopes of the peanut allergens shift following oral immunotherapy. Front Allergy *4*, 1279290. 10.3389/falgy.2023.1279290.

43. Suarez-Farinas, M., Suprun, M., Bahnson, H.T., Raghunathan, R., Getts, R., duToit, G., Lack, G., and Sampson, H.A. (2021). Evolution of epitope-specific IgE and IgG(4) antibodies in children enrolled in the LEAP trial. J Allergy Clin Immunol *148*, 835-842. 10.1016/j.jaci.2021.01.030.

44. Chen, G., Shrock, E.L., Li, M.Z., Spergel, J.M., Nadeau, K.C., Pongracic, J.A., Umetsu, D.T., Rachid, R., MacGinnitie, A.J., Phipatanakul, W., et al. (2021). High-resolution epitope mapping by AllerScan reveals relationships between IgE and IgG repertoires during peanut oral immunotherapy. Cell Rep Med *2*, 100410. 10.1016/j.xcrm.2021.100410.

45. Suarez-Farinas, M., Lee, K.W., Suprun, M., Bahnson, H.T., Foong, R.X., Du Toit, G., Lack, G., and Sampson, H.A. (2025). Kinetics of epitope-specific IgE and IgG(4) in early peanut allergy development and resolution. J Allergy Clin Immunol. 10.1016/j.jaci.2025.06.022.

46. Lopez-Sanz, C., Sanchez-Martinez, E., and Jimenez-Saiz, R. (2022). Protocol to desensitize human and murine mast cells after polyclonal IgE sensitization. STAR Protoc *3*, 101755. 10.1016/j.xpro.2022.101755.

47. Fernandez-Gallego, N., Castillo-Gonzalez, R., Moreno-Serna, L., Garcia-Civico, A.J., Sanchez-Martinez, E., Lopez-Sanz, C., Fontes, A.L., Pimentel, L.L., Gradillas, A., Obeso, D., et al. (2024). Allergic inflammation triggers dyslipidemia via IgG signalling. Allergy *79*, 2680-2699. 10.1111/all.16187.

48. Helmerhorst, E.J., Zamakhchari, M., Schuppan, D., and Oppenheim, F.G. (2010). Discovery of a novel and rich source of gluten-degrading microbial enzymes in the oral cavity. PLoS One *5*, e13264. 10.1371/journal.pone.0013264.

49. Wei, G., Darwish, G., Oppenheim, F.G., Schuppan, D., and Helmerhorst, E.J. (2020). Commensal Bacterium Rothia aeria Degrades and Detoxifies Gluten via a Highly Effective Subtilisin Enzyme. Nutrients *12*. 10.3390/nu12123724.

50. Wei, G., Tian, N., Siezen, R., Schuppan, D., and Helmerhorst, E.J. (2016). Identification of food-grade subtilisins as gluten-degrading enzymes to treat

celiac disease. Am J Physiol Gastrointest Liver Physiol *311*, G571-580. 10.1152/ajpgi.00185.2016.

51. Zhang, L., Chun, Y., Grishina, G., Lo, T., Reed, K., Wang, J., Sicherer, S., Berin, M.C., and Bunyavanich, S. (2025). Oral and Gut Microbial Hubs Associated With Reaction Threshold Interact With Circulating Immune Factors in Peanut Allergy. Allergy. 10.1111/all.16481.

52. Gomez de Aguero, M., Ganal-Vonarburg, S.C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., Steinert, A., Heikenwalder, M., Hapfelmeier, S., Sauer, U., et al. (2016). The maternal microbiota drives early postnatal innate immune development. Science *351*, 1296-1302. 10.1126/science.aad2571.

53. Caminero, A., Guzman, M., Libertucci, J., and Lomax, A.E. (2023). The emerging roles of bacterial proteases in intestinal diseases. Gut Microbes *15*, 2181922. 10.1080/19490976.2023.2181922.

54. Shan, L., Molberg, O., Parrot, I., Hausch, F., Filiz, F., Gray, G.M., Sollid, L.M., and Khosla, C. (2002). Structural basis for gluten intolerance in celiac sprue. Science *297*, 2275-2279. 10.1126/science.1074129.

55. Caminero, A., Galipeau, H.J., McCarville, J.L., Johnston, C.W., Bernier, S.P., Russell, A.K., Jury, J., Herran, A.R., Casqueiro, J., Tye-Din, J.A., et al. (2016). Duodenal Bacteria From Patients With Celiac Disease and Healthy Subjects Distinctly Affect Gluten Breakdown and Immunogenicity. Gastroenterology *151*, 670-683. 10.1053/j.gastro.2016.06.041.

56. Caminero, A., McCarville, J.L., Zevallos, V.F., Pigrau, M., Yu, X.B., Jury, J., Galipeau, H.J., Clarizio, A.V., Casqueiro, J., Murray, J.A., et al. (2019). Lactobacilli Degrade Wheat Amylase Trypsin Inhibitors to Reduce Intestinal Dysfunction Induced by Immunogenic Wheat Proteins. Gastroenterology *156*, 2266-2280. 10.1053/j.gastro.2019.02.028.

57. Tremaroli, V., and Backhed, F. (2012). Functional interactions between the gut microbiota and host metabolism. Nature *489*, 242-249. 10.1038/nature11552.

58. Thomson, C.A., Morgan, S.C., Ohland, C., and McCoy, K.D. (2022). From germ-free to wild: modulating microbiome complexity to understand mucosal immunology. Mucosal Immunol *15*, 1085-1094. 10.1038/s41385-022-00562-3.

59. Jimenez-Saiz, R., Anipindi, V.C., Galipeau, H., Ellenbogen, Y., Chaudhary, R., Koenig, J.F., Gordon, M.E., Walker, T.D., Mandur, T.S., Abed, S., et al. (2020). Microbial Regulation of Enteric Eosinophils and Its Impact on Tissue Remodeling and Th2 Immunity. Front Immunol *11*, 155. 10.3389/fimmu.2020.00155.

60. Jutel, M., Agache, I., Zemelka-Wiacek, M., Akdis, M., Chivato, T., Del Giacco, S., Gajdanowicz, P., Gracia, I.E., Klimek, L., Lauerma, A., et al. (2023). Nomenclature of allergic diseases and hypersensitivity reactions: Adapted to modern needs: An EAACI position paper. Allergy *78*, 2851-2874. 10.1111/all.15889.

61. Galli, S.J., and Tsai, M. (2012). IgE and mast cells in allergic disease. Nat Med *18*, 693-704. 10.1038/nm.2755.

62. Leyva-Castillo, J.M., Galand, C., Kam, C., Burton, O., Gurish, M., Musser, M.A., Goldsmith, J.D., Hait, E., Nurko, S., Brombacher, F., et al. (2019). Mechanical Skin Injury Promotes Food Anaphylaxis by Driving Intestinal Mast Cell Expansion. Immunity *50*, 1262-1275 e1264. 10.1016/j.immuni.2019.03.023.

63. Jacob, C., Yang, P.C., Darmoul, D., Amadesi, S., Saito, T., Cottrell, G.S.,
Coelho, A.M., Singh, P., Grady, E.F., Perdue, M., and Bunnett, N.W. (2005).
Mast cell tryptase controls paracellular permeability of the intestine. Role of
protease-activated receptor 2 and beta-arrestins. J Biol Chem *280*, 31936-31948.
10.1074/jbc.M506338200.
64. Strait, R.T., Mahler, A., Hogan, S., Khodoun, M., Shibuya, A., and Finkelman,
F.D. (2011). Ingested allergens must be absorbed systemically to induce
systemic anaphylaxis. J Allergy Clin Immunol *127*, 982-989 e981.
10.1016/j.jaci.2011.01.034.
65. Cahenzli, J., Koller, Y., Wyss, M., Geuking, M.B., and McCoy, K.D. (2013).
Intestinal microbial diversity during early-life colonization shapes long-term IgE
levels. Cell Host Microbe *14*, 559-570. 10.1016/j.chom.2013.10.004.
66. Wyss, M., Brown, K., Thomson, C.A., Koegler, M., Terra, F., Fan, V., Ronchi, F.,
Bihan, D., Lewis, I., Geuking, M.B., and McCoy, K.D. (2019). Using Precisely
Defined in vivo Microbiotas to Understand Microbial Regulation of IgE. Front
Immunol *10*, 3107. 10.3389/fimmu.2019.03107.
67. Stefka, A.T., Feehley, T., Tripathi, P., Qiu, J., McCoy, K., Mazmanian, S.K.,
Tjota, M.Y., Seo, G.Y., Cao, S., Theriault, B.R., et al. (2014). Commensal
bacteria protect against food allergen sensitization. Proc Natl Acad Sci U S A
111, 13145-13150. 10.1073/pnas.1412008111.
68. Arias, K., Chu, D.K., Flader, K., Botelho, F., Walker, T., Arias, N., Humbles, A.A.,
Coyle, A.J., Oettgen, H.C., Chang, H.D., et al. (2011). Distinct immune effector
pathways contribute to the full expression of peanut-induced anaphylactic
reactions in mice. J Allergy Clin Immunol *127*, 1552-1561 e1551.
10.1016/j.jaci.2011.03.044.
69. Gowthaman, U., Chen, J.S., Zhang, B., Flynn, W.F., Lu, Y., Song, W., Joseph, J.,
Gertie, J.A., Xu, L., Collet, M.A., et al. (2019). Identification of a T follicular helper
cell subset that drives anaphylactic IgE. Science *365*, eaaw6433.
10.1126/science.aaw6433.
70. Chu, D.K., Jimenez-Saiz, R., Verschoor, C.P., Walker, T.D., Goncharova, S.,
Llop-Guevara, A., Shen, P., Gordon, M.E., Barra, N.G., Bassett, J.D., et al.
(2014). Indigenous enteric eosinophils control DCs to initiate a primary Th2
immune response in vivo. J Exp Med *211*, 1657-1672. 10.1084/jem.20131800.
71. Jimenez-Saiz, R., Chu, D.K., Mandur, T.S., Walker, T.D., Gordon, M.E.,
Chaudhary, R., Koenig, J., Saliba, S., Galipeau, H.J., Utley, A., et al. (2017).
Lifelong memory responses perpetuate humoral T(H)2 immunity and anaphylaxis
in food allergy. J Allergy Clin Immunol *140*, 1604-1615 e1605.
10.1016/j.jaci.2017.01.018.
72. Knibb, R.C., Jones, C.J., Herbert, L.J., and Screti, C. (2024). Psychological
support needs for children with food allergy and their families: A systematic
review. Pediatr Allergy Immunol *35*, e14108. 10.1111/pai.14108.
73. Mori, F., Giovannini, M., Barni, S., Jimenez-Saiz, R., Munblit, D., Biagioni, B.,
Liccioli, G., Sarti, L., Liotti, L., Ricci, S., et al. (2021). Oral Immunotherapy for
Food-Allergic Children: A Pro-Con Debate. Front Immunol *12*, 636612.
10.3389/fimmu.2021.636612.

74. Niggemann, B., and Beyer, K. (2014). Factors augmenting allergic reactions. Allergy *69*, 1582-1587. 10.1111/all.12532.

75. Piletta-Zanin, A., Scherl, A., Benhamou, A., Braendle, G., Caubet, J.C., Graham, F., Grosgurin, O., Harr, T., Manzano, S., Nigolian, H., et al. (2025). The Severity of Allergic Reactions in a Real-World Environment Is Independent of the Eliciting Amounts of Foods. Allergy *80*, 238-247. 10.1111/all.16413.

76. Baker, J.L., Mark Welch, J.L., Kauffman, K.M., McLean, J.S., and He, X. (2024). The oral microbiome: diversity, biogeography and human health. Nat Rev Microbiol *22*, 89-104. 10.1038/s41579-023-00963-6.

77. Constante, M., Libertucci, J., Galipeau, H.J., Szamosi, J.C., Rueda, G., Miranda, P.M., Pinto-Sanchez, M.I., Southward, C.M., Rossi, L., Fontes, M.E., et al. (2022). Biogeographic Variation and Functional Pathways of the Gut Microbiota in Celiac Disease. Gastroenterology *163*, 1351-1363 e1315. 10.1053/j.gastro.2022.06.088.

78. Kunath, B.J., De Rudder, C., Laczny, C.C., Letellier, E., and Wilmes, P. (2024). The oral-gut microbiome axis in health and disease. Nat Rev Microbiol *22*, 791-805. 10.1038/s41579-024-01075-5.

79. Zamakhchari, M., Wei, G., Dewhirst, F., Lee, J., Schuppan, D., Oppenheim, F.G., and Helmerhorst, E.J. (2011). Identification of Rothia bacteria as gluten-degrading natural colonizers of the upper gastro-intestinal tract. PLoS One *6*, e24455. 10.1371/journal.pone.0024455.

80. Patrick, G.J., Liu, H., Alphonse, M.P., Dikeman, D.A., Youn, C., Otterson, J.C., Wang, Y., Ravipati, A., Mazhar, M., Denny, G., et al. (2021). Epicutaneous Staphylococcus aureus induces IL-36 to enhance IgE production and ensuing allergic disease. J Clin Invest *131*. 10.1172/JCI143334.

81. Rondeau, L.E., Da Luz, B.B., Santiago, A., Bermudez-Brito, M., Hann, A., De Palma, G., Jury, J., Wang, X., Verdu, E.F., Galipeau, H.J., et al. (2024). Proteolytic bacteria expansion during colitis amplifies inflammation through cleavage of the external domain of PAR2. Gut Microbes *16*, 2387857. 10.1080/19490976.2024.2387857.

82. Agaronyan, K., Sharma, L., Vaidyanathan, B., Glenn, K., Yu, S., Annicelli, C., Wiggen, T.D., Penningroth, M.R., Hunter, R.C., Dela Cruz, C.S., and Medzhitov, R. (2022). Tissue remodeling by an opportunistic pathogen triggers allergic inflammation. Immunity *55*, 895-911 e810. 10.1016/j.immuni.2022.04.001.

83. Caminero, A., McCarville, J.L., Galipeau, H.J., Deraison, C., Bernier, S.P., Constante, M., Rolland, C., Meisel, M., Murray, J.A., Yu, X.B., et al. (2019). Duodenal bacterial proteolytic activity determines sensitivity to dietary antigen through protease-activated receptor-2. Nat Commun *10*, 1198. 10.1038/s41467-019-09037-9.

84. Jimenez-Saiz, R., Ellenbogen, Y., Koenig, J.F.E., Gordon, M.E., Walker, T.D., Rosace, D., Spill, P., Bruton, K., Kong, J., Monteiro, K., et al. (2019). IgG1(+) B-cell immunity predates IgE responses in epicutaneous sensitization to foods. Allergy *74*, 165-175. 10.1111/all.13481.

85. Kong, J., Chalcraft, K., Mandur, T.S., Jimenez-Saiz, R., Walker, T.D., Goncharova, S., Gordon, M.E., Naji, L., Flader, K., Larche, M., et al. (2015).

Comprehensive metabolomics identifies the alarmin uric acid as a critical signal for the induction of peanut allergy. Allergy *70*, 495-505. 10.1111/all.12579.

86. Kwak, Y.K., Vikstrom, E., Magnusson, K.E., Vecsey-Semjen, B., Colque-Navarro, P., and Mollby, R. (2012). The Staphylococcus aureus alpha-toxin perturbs the barrier function in Caco-2 epithelial cell monolayers by altering junctional integrity. Infect Immun *80*, 1670-1680. 10.1128/IAI.00001-12.

87. Moran, M.C., Brewer, M.G., Schlievert, P.M., and Beck, L.A. (2023). S. aureus virulence factors decrease epithelial barrier function and increase susceptibility to viral infection. Microbiol Spectr *11*, e0168423. 10.1128/spectrum.01684-23.

88. Martinez-Garcia, S., Rodriguez-Martinez, S., Cancino-Diaz, M.E., and Cancino-Diaz, J.C. (2018). Extracellular proteases of Staphylococcus epidermidis: roles as virulence factors and their participation in biofilm. APMIS : acta pathologica, microbiologica, et immunologica Scandinavica *126*, 177-185. 10.1111/apm.12805.

89. Cau, L., Williams, M.R., Butcher, A.M., Nakatsuji, T., Kavanaugh, J.S., Cheng, J.Y., Shafiq, F., Higbee, K., Hata, T.R., Horswill, A.R., and Gallo, R.L. (2021). Staphylococcus epidermidis protease EcpA can be a deleterious component of the skin microbiome in atopic dermatitis. J Allergy Clin Immunol *147*, 955-966 e916. 10.1016/j.jaci.2020.06.024.

90. Forbes, E.E., Groschwitz, K., Abonia, J.P., Brandt, E.B., Cohen, E., Blanchard, C., Ahrens, R., Seidu, L., McKenzie, A., Strait, R., et al. (2008). IL-9- and mast cell-mediated intestinal permeability predisposes to oral antigen hypersensitivity. J Exp Med *205*, 897-913. 10.1084/jem.20071046.

91. Hayes, C.L., Dong, J., Galipeau, H.J., Jury, J., McCarville, J., Huang, X., Wang, X.Y., Naidoo, A., Anbazhagan, A.N., Libertucci, J., et al. (2018). Commensal microbiota induces colonic barrier structure and functions that contribute to homeostasis. Scientific reports *8*, 14184. 10.1038/s41598-018-32366-6.

92. Li, Y., Watanabe, E., Kawashima, Y., Plichta, D.R., Wang, Z., Ujike, M., Ang, Q.Y., Wu, R., Furuichi, M., Takeshita, K., et al. (2022). Identification of trypsin-degrading commensals in the large intestine. Nature *609*, 582-589. 10.1038/s41586-022-05181-3.

93. Santiago, A., Hann, A., Constante, M., Rahmani, S., Libertucci, J., Jackson, K., Rueda, G., Rossi, L., Rithwick, R., Ruf, W., et al. (2023). Crohn's disease proteolytic microbiota enhances inflammation through PAR2 pathway in gnotobiotic mice. Gut Microbes *15*, 2205425. 10.1080/19490976.2023.2205425.

94. Rotella, K., and Oriel, R.C. (2022). Accidental Reactions to Foods: Frequency, Causes, and Severity. Curr Treat Options Allergy *9*, 157-168. 10.1007/s40521-022-00314-5.

95. Hsu, C., Hosakoppal, S., Yong, M., Gupta, R., Makhija, M., and Singh, A.M. (2024). Prevalence and Characteristics of Accidental Ingestions Among Pediatric Food Allergy Patients. J Allergy Clin Immunol Pract *12*, 3089-3095 e3082. 10.1016/j.jaip.2024.08.006.

96. Li, J.C., Rotter, N.S., Stieb, E.S., Stockbridge, J.L., Theodorakakis, M.D., and Shreffler, W.G. (2024). Utility of food allergy thresholds. Ann Allergy Asthma Immunol *132*, 321-327. 10.1016/j.anai.2023.12.012.

97. Patel, N., Adelman, D.C., Anagnostou, K., Baumert, J.L., Blom, W.M., Campbell, D.E., Chinthrajah, R.S., Mills, E.N.C., Javed, B., Purington, N., et al. (2021). Using data from food challenges to inform management of consumers with food allergy: A systematic review with individual participant data meta-analysis. J Allergy Clin Immunol *147*, 2249-2262 e2247. 10.1016/j.jaci.2021.01.025.

98. Santos, A.F., Barbosa-Morais, N.L., Hurlburt, B.K., Ramaswamy, S., Hemmings, O., Kwok, M., O'Rourke, C., Bahnson, H.T., Cheng, H., James, L., et al. (2020). IgE to epitopes of Ara h 2 enhance the diagnostic accuracy of Ara h 2-specific IgE. Allergy *75*, 2309-2318. 10.1111/all.14301.

99. Patil, S.U., Steinbrecher, J., Calatroni, A., Smith, N., Ma, A., Ruiter, B., Virkud, Y., Schneider, M., and Shreffler, W.G. (2019). Early decrease in basophil sensitivity to Ara h 2 precedes sustained unresponsiveness after peanut oral immunotherapy. J Allergy Clin Immunol *144*, 1310-1319 e1314. 10.1016/j.jaci.2019.07.028.

100. Blackman, A.C., Thapa, S., Venkatachalam, A., Horvath, T.D., Runge, J.K., Haidacher, S.J., Hoch, K.M., Haag, A.M., Luna, R.A., and Anagnostou, A. (2022). Insights into Microbiome and Metabolic Signatures of Children Undergoing Peanut Oral Immunotherapy. Children (Basel) *9*. 10.3390/children9081192.

101. Zhang, L., Chun, Y., Ho, H.E., Arditi, Z., Lo, T., Sajja, S., Rose, R., Jones, D., Wang, J., Sicherer, S., and Bunyavanich, S. (2022). Multiscale study of the oral and gut environments in children with high- and low-threshold peanut allergy. J Allergy Clin Immunol *150*, 714-720 e712. 10.1016/j.jaci.2022.04.026.

102. Chu, D.K., Wood, R.A., French, S., Fiocchi, A., Jordana, M., Waserman, S., Brozek, J.L., and Schunemann, H.J. (2019). Oral immunotherapy for peanut allergy (PACE): a systematic review and meta-analysis of efficacy and safety. Lancet *393*, 2222-2232. 10.1016/S0140-6736(19)30420-9.

103. Barshow, S., Tirumalasetty, J., Sampath, V., Zhou, X., Seastedt, H., Schuetz, J., and Nadeau, K. (2024). The Immunobiology and Treatment of Food Allergy. Annu Rev Immunol *42*, 401-425. 10.1146/annurev-immunol-090122-043501.

104. Fierstein, J.L., Brown, D., Gupta, R., and Bilaver, L. (2021). Understanding Food-Related Allergic Reactions Through a US National Patient Registry. J Allergy Clin Immunol Pract *9*, 206-215 e201. 10.1016/j.jaip.2020.08.011.

105. Taylor, S.L., and Baumert, J.L. (2010). Cross-contamination of foods and implications for food allergic patients. Curr Allergy Asthma Rep *10*, 265-270. 10.1007/s11882-010-0112-4.

106. Miller, T.A., Koppelman, S.J., Bird, J.A., Hernandez-Trujillo, V., Thyagarajan, A., Mack, D., Chalil, J.M., Green, T.D., and Baumert, J.L. (2022). Peanut cross-contamination in randomly selected baked goods. Ann Allergy Asthma Immunol *128*, 439-442. 10.1016/j.anai.2022.01.037.

107. Dewhirst, F.E., Chien, C.C., Paster, B.J., Ericson, R.L., Orcutt, R.P., Schauer, D.B., and Fox, J.G. (1999). Phylogeny of the defined murine microbiota: altered Schaedler flora. Appl Environ Microbiol *65*, 3287-3292. 10.1128/AEM.65.8.3287-3292.1999.

108. Burton, O.T., Stranks, A.J., Tamayo, J.M., Koleoglou, K.J., Schwartz, L.B., and Oettgen, H.C. (2017). A humanized mouse model of anaphylactic peanut allergy. J Allergy Clin Immunol *139*, 314-322 e319. 10.1016/j.jaci.2016.04.034.

109. Whelan, F.J., and Surette, M.G. (2017). A comprehensive evaluation of the sl1p pipeline for 16S rRNA gene sequencing analysis. Microbiome *5*, 100. 10.1186/s40168-017-0314-2.

110. Pazos-Castro, D., Gonzalez-Klein, Z., Montalvo, A.Y., Hernandez-Ramirez, G., Romero-Sahagun, A., Esteban, V., Garrido-Arandia, M., Tome-Amat, J., and Diaz-Perales, A. (2022). NLRP3 priming due to skin damage precedes LTP allergic sensitization in a mouse model. Sci Rep *12*, 3329. 10.1038/s41598-022-07421-y.

111. Wang, Y., Matsushita, K., Jackson, J., Numata, T., Zhang, Y., Zhou, G., Tsai, M., and Galli, S.J. (2021). Transcriptome programming of IL-3-dependent bone marrow-derived cultured mast cells by stem cell factor (SCF). Allergy *76*, 2288-2291. 10.1111/all.14808.

112. Kechin, A., Boyarskikh, U., Kel, A., and Filipenko, M. (2017). cutPrimers: A New Tool for Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing. J Comput Biol *24*, 1138-1143. 10.1089/cmb.2017.0096.

113. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal *17*, 10-12. https://doi.org/10.14806/ej.17.1.200.

114. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., and Holmes, S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods *13*, 581-583. 10.1038/nmeth.3869.

115. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glockner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res *41*, D590-596. 10.1093/nar/gks1219.

116. Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS One *5*, e9490. 10.1371/journal.pone.0009490.

117. McMurdie, P.J., and Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One *8*, e61217. 10.1371/journal.pone.0061217.

118. Derakhshani, H., Bernier, S.P., Marko, V.A., and Surette, M.G. (2020). Completion of draft bacterial genomes by long-read sequencing of synthetic genomic pools. BMC genomics *21*, 519. 10.1186/s12864-020-06910-6.

119. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics *34*, i884-i890. 10.1093/bioinformatics/bty560.

120. Chen, S. (2023). Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp. Imeta *2*, e107. 10.1002/imt2.107.

121. Wick, R.R., Judd, L.M., Gorrie, C.L., and Holt, K.E. (2017). Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. PLoS Comput Biol *13*, e1005595. 10.1371/journal.pcbi.1005595.

122. Schwengers, O., Jelonek, L., Dieckmann, M.A., Beyvers, S., Blom, J., and Goesmann, A. (2021). Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. Microb Genom *7*. 10.1099/mgen.0.000685.

1152   123.   Page, A.J., Cummins, C.A., Hunt, M., Wong, V.K., Reuter, S., Holden, M.T.,
1153   Fookes, M., Falush, D., Keane, J.A., and Parkhill, J. (2015). Roary: rapid large-
1154   scale prokaryote pan genome analysis. Bioinformatics *31*, 3691-3693.
1155   10.1093/bioinformatics/btv421.

1156

1157

1158

1159

1160

1161

STAR PROTOCOLS

RESOURCE AVAILABILITY

Lead contact

Requests for further information or resources should be directed to the lead contact Alberto Caminero (acamine@mcmaster.ca).

Materials availability

This study did not generate new unique reagents.

Data and code availability

The microbiome sequencing data generated in this study have been deposited in the NCBI BioProject database under the accession number PRJNA1223483. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD060888 and 10.6019/PXD060888.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Human samples

Human saliva from peanut (PN) allergic patients entering oral immunotherapy (OIT; n=19) were collected. These samples were obtained from children aged 1-14 years who participated in a standard clinical oral food challenge to establish PN threshold prior to clinical management with PN OIT at Massachusetts General Hospital. Saliva was collected with the SalivaBio Children's Swab (Salimetrics) following the manufacturer's protocol, and samples were stored at -80°C until further use. PN- and antigen-specific Ig levels in plasma from patients undergoing OIT were measured using a Phadia ImmunoCAP 1000 instrument (ThermoFisher) according to the manufacturer's instructions. Saliva samples from healthy volunteers (n=13) were collected at McMaster University under BUP#423. All patients and healthy volunteers provided written informed consent as per local institutional review board guidelines. Supportive and demographics data can be found in Table S1.

To complement oral sampling, small intestinal aspirate samples (mid-jejunal washes) were obtained from non-allergic control individuals without any chronic gastrointestinal or metabolic conditions (n=5). These samples were collected under an ongoing clinical protocol led by Dr. Premysl Bercik at McMaster University (HIREB #15311). Push enteroscopy was performed following an overnight fast, and jejunal washes (15–20 mL) were collected as previously described[77]. Briefly, sterile saline was applied using a water jet to dislodge mucus-associated and luminal bacteria from the mid-jejunum and samples were frozen with or without 20% glycerol for further use. Small intestinal microbiota was

analyzed by 16S rRNA sequencing, and samples were cultured for isolation and identification of PN-degrading bacteria. Supportive and demographics data can be found in Table S1.

Human sera for Western blotting were provided by the Allergology Service at Hospital Universitario de La Princesa (Madrid, Spain). The Research Ethics Committee of Hospital Universitario de La Princesa approved the study protocol (reference 4460). All the donors provided written informed consent with no conflict of interest. Donor sera characterization is detailed in Table S2.

Mice

Age-, sex-, and strain-matched controls were used in all experiments. Specific pathogen-free (SPF) C57BL/6 mice were purchased from Charles River and Taconic, while SPF C3H-HeN mice were purchased from Taconic. Germ-free (GF) C57BL/6 mice used for colonization experiments were originally purchased from Taconic and derived GF by two-stage embryo transfer. They were bred in flexible-film gnotobiotic isolators at McMaster University's Farncombe Family Digestive Health Research Institute Axenic Gnotobiotic Unit (AGU). C3H-HeN mice with altered Schaedler flora (ASF) microbiota, a simple and stable microbiota[81,107], were bred and housed in GF conditions at the AGU. The C3H-HeN ASF mice were originally purchased from Taconic and have been maintained in the AGU for over 20 generations. All mice were maintained on a 12-h light-dark cycle and fed an autoclaved chow diet and sterile water *ad libitum*. Eight-to-12-week-old mice were used throughout. All procedures were approved by the Environmental Council of the Community of Madrid with PROEX references 45.2/20 and 9.7/23, and the McMaster University Animal Care Committee and McMaster University Animal Research Ethics Board in accordance with the Animal Utilization Protocol #22-26. Ethical regulations for animal research were strictly followed.

Food allergy and anaphylaxis model

To study the impact of microbes on PN metabolism and anaphylaxis, C3H-HeN mice with different microbiota were sensitized to PN using distinct techniques. 1. Intra-gastric sensitization: mice were provided 3.25 mg of crude PN extract (CPE; Greer Laboratories Inc) and 0.625 µg of cholera toxin (List Biological Laboratories) in 0.1 mL of PBS by oral gavage once a week for 6 weeks; 2. Intra-peritoneal sensitization: mice were provided 3.25 mg of CPE and 6.25 µg of cholera toxin in 0.1 mL of PBS intra-peritoneally once a week for 4 weeks; 3. Passive sensitization- mice were subjected to 3 intra-peritoneal injections (0.1, 0.2, and 0.4 mL) of serum obtained from intra-peritoneally sensitized PN-allergic mice as described[47]. After sensitization, mice were challenged with a 24 mg bolus of CPE in 0.8 mL of PBS by oral gavage. In some experiments, CPE was pre-digested with PN allergen-degrading bacteria (*Rothia* R3, *Staphylococcus* S1, or S2), or mice were colonized with PN-degrading bacteria (*Rothia* R3), prior to the oral challenge to evaluate

the impact of bacterial PN metabolism on allergenicity and anaphylaxis. Serum was
collected 48 h before challenge. Mice were monitored for 40 min after challenge for
changes in rectal (core) body temperature. Peripheral blood was collected, and mice were
euthanized 40 min after the oral challenge. Mucosal mast cell protease 1 (mMCP-1) was
quantified in serum after the PN challenge, as it is a key biomarker of anaphylaxis in mice,
serving a similar role to serum tryptase for indicating mast cell activation in humans[108].
Platelet-activating factor acetylhydrolase (PAF-AH) was quantified in serum as a marker
of mast cell activation.
1245
<u>Microbiota model for studying microbial PN metabolism</u>
To investigate the role of microbiota in PN metabolism, we used two colonization
approaches: 1. We studied PN metabolism in mice colonized with ASF and SPF
microbiota, and GF controls; 2. GF C57BL/6 mice were mono-colonized with selected
PN-degrading bacteria (*Rothia* R3 or *Staphylococcus* S1, S2, or S3). After colonization,
mice were administered 24 mg of CPE in 0.8 mL of PBS by oral gavage. Small and large
intestinal contents and serum from peripheral blood were collected 40 minutes post-
administration for quantification of PN allergens Ara h 1 and 2 in biological samples by
ELISA (Indoor Biotechnologies) as described below. Additionally, small intestinal
contents were incubated *ex vivo* with CPE overnight (o/n), and remaining Ara h 1 and 2
levels were quantified by ELISA (Indoor Biotechnologies). C57BL/6 mice were used in
these experiments because of difficulties generating C3H-HeN mice under GF conditions.
1258
<u>Characterization of bacterial PN-degradation and immune activation</u>
To assess bacterial degradation of PN allergens and its impact on immune activation, we
performed *in vitro* digestions, proteomic characterization, and mast cell activation assays.
Bacterial isolates were incubated with CPE and degradation of Ara h 1 and Ara h 2 was
quantified by ELISA. Then, *Rothia* (R1, R2, and R3) and *Staphylococcus* (S1, S2, and
S3) candidates were selected for further characterization by SDS-PAGE, and Western
blotting. Western blotting was performed using sera from PN-allergic patients as well as
sera from mice allergic to Ara h 1 or to Ara h 2, to identify IgE-binding epitopes in the
digested PN allergens. Proteomic analysis using nano liquid chromatography coupled to
mass spectroscopy was used to characterize the degradation products and PN-specific
epitopes. For immune activation studies, bone marrow-derived mast cells (BMMCs) were
cultured and sensitized to PN with sera from PN-allergic mice. These cells were
challenged with bacterially digested PN to evaluate BMMC activation by β-
hexosaminidase release and flow cytometry analysis of degranulation markers (CD63,
CD107a)[46].
1274
1275

METHOD DETAILS

Mouse colonization procedures

For colonization of GF C57BL/6 mice with ASF and SPF microbiota, fresh cecal and colon contents were harvested from SPF C57BL/6 mice (Taconic) or ASF C3H-HeN mice, diluted 1:10 in sterile PBS in anaerobic conditions, and 0.2 mL of each cecal/fecal suspension was orally gavaged to mice.

For mono-colonization of GF C57BL/6 mice with PN-degrading bacteria, *Rothia* R3 or *Staphylococcus* S1, S2, or S3 were grown o/n in brain heart infusion broth (BHI; Research products International Corp.) and 10^9 colony forming units (CFU) from the culture were provided by oral gavage with PBS as a vehicle. GF controls were provided PBS by oral gavage.

For the PN anaphylaxis model, ASF C3H-HeNTac mice received 10^9 CFU total of *Rothia* R3, by oral gavage, administered once per week for three weeks after sensitization and before PN challenge. Sterile PBS was used as a vehicle and served as a non-colonization control.

Colonization was confirmed by culturing intestinal content combined with PCR of the 16S rRNA gene and Sanger sequencing to verify bacterial identity (methods detailed below). These checks were performed throughout the experiments and at endpoint. Mice with evidence of contamination or failed colonization were excluded from the analysis. For selected experiments, 16S rRNA microbiota analysis was also performed to confirm overall microbial community composition.

PN-specific IgE determination

Peripheral blood was collected by retro-orbital bleeding 48 h before the PN challenge. Serum PN-specific IgE was measured using a sandwich ELISA adapted from previously described methods[47]. Briefly, 96-well plates (high-binding, flat-bottom, polystyrene; Corning, USA) were coated o/n at 4°C with 2 µg/mL rat anti-mouse IgE antibody diluted in PBS. Plates were washed with PBS/0.05% Tween-20 (PBS-T; Sigma Aldrich), then blocked with 1% BSA in PBS-T for 2 h at room temperature (RT). Plates were washed thoroughly with PBS-T, then serum samples were incubated on the plate at 1/2 dilution o/n at 4°C. To detect PN-specific IgE, plates were incubated with biotinylated-CPE (biot-CPE; 0.1518 µg/mL in blocking solution) for 90 min at RT. Biot-CPE was generated using the EZ-Link Sulfo-NHS-LC-Biotinylation kit (Thermo Scientific). A standard curve, to which no capture antibody or sera was added, was generated using biot-CPE, serially diluted from 75 ng/mL to 0.6 ng/mL in PBS. Plates were washed thoroughly, then incubated with streptavidin-HRP (1:250 dilution in blocking solution; Thermo Scientific) and tetramethylbenzidine substrate (TMB; Thermo Scientific or Biolegend). The reaction

1316 was stopped with 1 N H_2SO_4. OD was measured at 450 nm, and the values were
1317 interpolated against the biot-CPE standard curve to determine relative PN-specific IgE-
1318 binding. Results are expressed as PN-IgE (Relative Binding, OD), reflecting the ability of
1319 serum PN-specific IgE to bind PN allergens.
1320

<u>mMCP-1 quantification</u>
Peripheral blood was collected by retro-orbital bleeding 40 min after the PN challenge.
The concentration of mMCP-1 was quantified using a Mouse mMCPT-1 ELISA kit
(Thermo Scientific) following the manufacturer's instructions. Briefly, serum samples were
diluted by 1/25 and incubated on the capture antibody (anti-mouse mMCPT-1)-coated
plate for 2 h at RT. After washing, a biotin-conjugated anti-mouse mMCPT-1 antibody
was added to the plate and incubated for 1 h at RT. The reaction was developed with
avidin-HRP and TMB, stopped with 1N H_2SO_4 and OD was measured at 450 nm. The
results are expressed as ng of mMCP-1 per mL of serum and multiplied by the dilution
factor of the sample.
1331

<u>ELISA detection of PN allergens in biological samples</u>
Ara h 1 and 2 were detected in samples harvested 40 min following intra-gastric
administration of 24 mg of CPE in 0.8 mL of PBS. Serum from peripheral blood was
collected by retro-orbital bleeding and analyzed undiluted. Whole small and large
intestinal content was analyzed after dilution by 1/5 and 1/10 in PBS. The quantification
of PN allergens was performed using proprietary ELISA kits for Ara h 1 and 2 following
the manufacturer's instructions (Indoor Biotechnologies). Briefly, for Ara h 1 detection by
ELISA (Indoor Biotechnologies): Plates pre-coated with a monoclonal anti-Ara h 1
antibody were washed with wash buffer, and biological samples and Ara h 1 standards
were added and incubated for 1 h. The plates were washed, and a second monoclonal
anti-Ara h 1 antibody conjugated to peroxidase was added and incubated for 1 h. After a
final wash, TMB substrate was added, and the reaction was stopped with 0.5 N H_2SO_4.
OD was measured at 450 nm, and results were expressed as the concentration of Ara h
1 in biological samples, multiplied by the dilution factor. For Ara h 2 detection by ELISA
(Indoor Biotechnologies): Plates pre-coated with a monoclonal anti-Ara h 2 antibody were
washed with wash buffer, and biological samples and Ara h 2 standards were added and
incubated for 1 h. After washing, a rabbit anti-Ara h 2 polyclonal detection antibody was
added and incubated for 1 h. This was followed by the addition of a peroxidase-
conjugated anti-rabbit IgG for 1 h. After the final wash, TMB substrate was added, and
the reaction was stopped with 0.5 N H_2SO_4. OD was measured at 450 nm, and results
were expressed as the concentration of Ara h 2 in biological samples, multiplied by the
dilution factor.
1354

1355

 Detection of PN degradation by microbes isolated from human saliva and mouse and
human intestinal content
Whole human saliva samples from healthy individuals (n=13), human small intestinal
aspirates (jejunum) (n=5), and mouse small and large intestinal contents (10 mg/mL in
PBS) were plated on agar media plates designed to isolate PN-degrading bacteria. BHI
media enriched with powdered PN (organic partially defatted PN; PB&Me, Sahah
Naturals) or agar supplemented with powered PN were used. Plates were incubated with
samples for 48 h in aerobic and anaerobic conditions (Bactron IV anaerobic chamber).
The bacteria were isolated based on their capacity to generate a visible hydrolytic halo in
the media, which is indicative of the breakdown of PN components in the media. Isolated
bacteria demonstrating PN hydrolytic activity were incubated in liquid BHI with CPE (0.5
mg/mL). After 48 h of incubation under aerobic or anaerobic conditions, the amount of
Ara h 1 and 2 in media was quantified by ELISA as described above. Allergen degradation
was calculated as the percentage of allergen degraded relative to the initial concentration
in non-growth controls (CPE). These values were visualized in heatmaps using a blue-
red color scale, where blue represents no degradation (0%) and red indicates complete
degradation (100%).

Relative abundance of each bacterial isolate was determined by calculating the proportion
of times each taxon was isolated across all samples. These values were expressed as
percentages and displayed using a green color scale, ranging from light green (low
relative abundance) to dark green (high relative abundance).

Salivary and intestinal microbes were identified using Sanger sequencing technology.
Briefly, DNA from isolates was extracted by picking single bacterial colonies into water,
boiling, and centrifuging at 2,000 RCF for 2 min. The 8F-926R region of the 16S rRNA
gene was amplified by PCR using the extracted DNA and sequences were determined
using Sanger sequencing (FW primer: 5'- AGAGTTTGATCCTGGCTCAG-3', RV primer:
5'-CCGTCAATTCCT-TTRAGTTT-3'). The resulting sequences for the isolates were
taxonomically assigned using the NCBI nucleotide collection database[109].
1385

PAF-AH quantification
Serum levels of platelet-activating factor acetylhydrolase (PAF-AH) were quantified using
a mouse PAF-AH/PLA2G7/Lp-PLA2 ELISA kit (Novus Biologicals) according to the
manufacturer's instructions. Briefly, serum samples were diluted 1:10 in PBS and
incubated capture antibody coated plates. After washing, a biotin-conjugated detection
antibody was added for 1 h at RT, followed by HRP conjugate for 30 min. The reaction
was developed with TMB, stopped with 1N H_2SO_4, and OD was measured at 450nm.
Results were expressed as ng/mL serum and corrected for the dilution factor.
1394

 Quantification of mucosal tryptase-positive mast cells

Paraffin-embedded small intestinal sections were deparaffinized in xylene and
redhydrated through a graded ethanol series. Antigen retrieval was performed by heat in
citric acid buffer, and non-specific binding was blocked with 3% BSA in PBS for 45 min.
Sections were incubated o/n at 4°C with a rabbit monoclonal anti-mast cell tryptase
antibody (1:200, Abcam). Immunoreactive cells were visualized using 3,3-
diaminobenzidiene (DAB; Dako). Negative controls were performed by omitting the
primary antibody. Tryptase positive mast cells were quantified in the mucosa and
normalized to mucosal area using an Axio Imager Z2 microscope (Carl Zeiss, Jena,
Germany). The data are expressed as tryptase-positive cells/μm^2.

1406

Microbial digestion of PN for challenge

Selected bacteria (*Rothia aeria* (R3), *Staphylococcus epidermidis* (S1), and
Staphylococcus aureus (S3)) were grown o/n on BHI agar plates enriched with 1%
powdered PN to confirm PN degradation capability (visible PN hydrolytic halo). For each
strain, a 10 mL pre-inoculum was prepared by o/n culture in liquid BHI. Then, 10 µL of
pre-inoculum was transferred into 5 mL of OptiMEM medium containing 30 mg/mL CPE
and incubated for 6 h to complete the PN digestion process. The resulting digests were
boiled for 5 min to kill the bacteria, and 800 µL of the digests, containing 24 mg of digested
PN, was administered by oral gavage at the time of challenge as described above.
OptiMEM medium served as the vehicle for non-digested CPE challenge.

1417

Measurement of *ex vivo* intestinal PN-degrading capacity

The ability of the intestinal content from mice colonized with PN-degrading bacteria to
degrade PN was assessed using ELISA and a bioassay with agar media enriched with
PN. Solid BHI agar enriched with 1% powdered PN was prepared, and a 50 µL aliquot of
diluted (1/2 in PBS) small intestinal content was placed into wells created in the agar. The
plates were then incubated for 24 h. The PN-degrading capacity was determined by
measuring the diameter of the clear halo around the inoculation site, indicating
degradation, and the results were expressed in millimeters. In addition, the intestinal
contents were also incubated with CPE (0.5 mg/mL). After 48 h of incubation, the quantity
of Ara h 1 and 2 in media was analyzed by ELISA as described above. The data for Ara
h 1 and 2 degradation are represented as blue-red color scales showing the percentage
of allergen degradation relative to the control incubations (CPE), with blue indicating no
degradation (0%) and red indicating complete degradation (100%).

1431

1432 Ussing chamber detection of PN translocation after bacterial digestion

1433 Translocation of PN allergens after digestion by bacteria was evaluated *ex vivo* using
1434 small intestinal sections of SPF C3H-HeN mice and the Ussing chamber technique. For

preparation of bacterial PN allergen digestions, *Rothia mucilaginosa* (R3), *Staphylococcus epidermidis* (S1), and *Staphylococcus aureus* (S2 & S3) were grown for 16 h in BHI liquid medium. Bacteria were then incubated in BHI with 5 mg/mL CPE for 4 h to allow for bacterial digestion of the PN allergens. After incubation, samples were boiled for 15 min. Bacterial digestions were performed in triplicate, with a negative control containing no bacterial inoculum. For the Ussing chamber technique, 1.5 cm sections of proximal small intestine were collected, opened along the mesenteric border, and mounted on the sliders of Ussing chambers. Each chamber exposed 0.25 cm² of tissue surface area to 4 mL of circulating oxygenated Krebs buffer containing 10 mM glucose (serosal side) and 10 mM mannitol (mucosal side), maintained at 37°C and aerated with 95% O_2 and 5% CO_2. Then, 0.2 mL of each bacterial digestion was placed into 4 mL of Krebs buffer on the mucosal side of the chamber. The experiment was run for 2 h, after which the total mucosal and serosal volumes were collected. The concentrations of PN allergens Ara h 1 and Ara h 2 on the serosal side were quantified by ELISA and expressed as intestinal translocation of allergens. The Ussing chamber system was from Physiologic Instruments.

Microbial digestion of PN for proteomic and immune activation assays

Rothia (R1- *Rothia aeria* R2- *Rothia dentacariosa* R3- *Rothia mucilaginosa*) and *Staphylococcus* (S1- *Staphylococcus epidermidis,* S2- *Staphylococcus aureus,* S3- *Staphylococcus aureus*) species were selected for characterization of PN allergen degradation and allergenicity. Bacteria were grown o/n in BHI (10 mL), then 10 µL of bacterial culture was taken and used to inoculate 1 mL of OptiMem with 0.5 mg/mL CPE. The bacteria were incubated in the media with CPE for 4 h at 37°C. After the incubation, the digestion was stopped by boiling for 5 min. Subsequently, the digestion of Ara h 1 and 2 were detected using ELISA, SDS-PAGE and Western blotting, and BMMC activation assays.

SDS-PAGE and Western blotting

To assess PN degradation profiles produced by degradation by *Rothia* and *Staphylococcus* species we performed SDS-PAGE and Western blotting. Bacteria were grown in 0.5 mg/mL as mentioned above. Undigested PN and Ara h 1 and 2 were used as controls. Then Laemmli sample buffer (Bio-Rad) with β-mercaptoethanol (Sigma-Aldrich) at 1/10 was added at 1/4 in a final volume of 50 µL. Before loading the gel (Any kD™ Mini-PROTEAN® TGX™ Precast Protein Gels, Bio Rad), samples were heat-denatured at 100ºC for 5 min. Moreover, a protein molecular weight marker was added (ThermoFischer). The gel was run in 1X running buffer (*i.e.*, SDS/Tris/Glycine 10X, Bio-Rad) for 1-1.5 h at 100 V using an electrophoretic cell and a power supply Mini-Protean System Bio-Rad (Bio-Rad).

For SDS-PAGE, after running, the gel was placed in staining solution (45% of methanol and 10% of acetic acid in distilled water, plus 0.2% w/v of Coomassie blue, Bioscience) for 1 h in agitation. Then, the staining solution was removed using destaining solution (20% methanol and 10% acetic acid, Sigma-Aldrich, in distilled water) in constant agitation, the process was repeated every 15 min until the background was eliminated. Then, the gel was photographed using a molecular imaging system (Amersham ImageQuant™ 800).

For Western blotting, after running the gel, the bands of the gel were transferred to a nitrocellulose membrane (Bio-Rad) using a transfer system (fiber pad, filter paper, membrane, gel, filter paper and fiber pad) and transfer buffer (20% methanol and 10% of Tris-Glycine 1X in distilled water). The transference was run for 1 h at 400 mA. Then, to check that bands were transferred correctly, the membrane was stained with Ponceau S Staining solution (ThermoScientific). After cleaning with water and 3 washes with TBS-T solution, the membrane was blocked in constant agitation for at least 1 h with 3% BSA (NZYtech) in TBS-T. Then, the membrane was incubated with sera from PN-allergic donors, or sera from mice allergic to Ara h 1 or to Ara h 2 (diluted in 3% BSA at different dilutions depending on the amount of IgE) as primary antibody, at 4ºC o/n with smooth agitation. Sera from non PN-allergic donors (Table S2) and non PN-allergic mice were used as negative control. During day 2, after performing 6 washes with TBS-T every 10 min with agitation, the membrane was incubated with anti-human IgE-HRP (Invitrogen) 1:10,000 or anti-mouse IgE-HRP (SouthernBiotech) 1:1,000 as secondary antibody for 1 h at RT with agitation. Then, 6 washes with TBS-T every 10 min with agitation were performed before revealing the membrane According to the instructions of the manufacturer (Cytiva-Merck), 1:1 of peroxide and luminol and 1 mL of the mixture was added to the membrane. The chemiluminescent signal of the gel was captured in a chemiluminescent imaging system (Amersham ImageQuant™ 800). SDS-PAGE and Western blots were analyzed using Fiji ImageJ.

Proteomics

Proteomics was performed to identify bacterial degradation products of PN allergens and to map PN-specific epitopes. Microbial PN digestions (above) by *Rothia* (R1, R2, and R3) and *Staphyloccocus* (S1, S2, and S3) were ultra-filtered using microcon filter units with a 3 kD cutoff that were previously equilibrated and washed with water. The peptides of the flow-through were selected (<3 kD) and they were cleaned with ZipTip with 0.6 µL C18 resin (Merck Millipore) to avoid the presence of elements that could interfere with the mass spectrometry, and they were drained in a SpeedVac Vacuum Concentrator. The samples, resuspended in a volume of 14 µL and 2 µL LC-MS grade water containing 2% (v/v) acetonitrile and 0.1% (v/v) formic acid, were used for the quantification with the QBIT method, analyzing around 0.8 µg per sample. The peptides were analyzed by nano liquid

chromatography coupled to mass spectrometry in data-dependent-acquisition mode using an UltiMate 3000 High Pressure Liquid Chromatograph (Fisher Scientific) and a Orbitrap Exploris™ 240 Mass Spectrometer (Fisher Scientific). The flow of the chromatograph was 250 nL/min for 60 min, and an Easy-spray PepMap C18 analytical column 50 cm × 75 μm (Fisher Scientific) was used.

The obtained MS1 and MS2 spectra were analyzed using the Peaks (https://www.bioinfor.com/peaksdb/) search engine against an *Arachis hypogaea* proteome database obtained from UniprotKB (https://www.uniprot.org/) combined with a list of typical laboratory contaminants, using a tolerance of 10 ppm and 0.02 Da for the precursor ions and fragments, respectively. Carbamidomethylation in cysteines was used as fixed modification and acetyl in the N-terminal end of the protein and methionine oxidation were used as variable modifications. The search was performed without restrictions to any proteases and the results were shown as proteins identified with at least a unique peptide with a False Discovery Rate ≤1%.

The unique peptides obtained after the analysis were located and identified in the sequences of Ara h 1, which were downloaded from the Protein Data Bank (https://www.rcsb.org/). The most common human IgE-binding epitopes of Ara h 1 were found in the following allergen databases: Allergen Nomenclature (https://allergen.org/); Allergen Online (http://allergenonline.org/); Compare Database (https://db.comparedatabase.org/); and The Immune Epitope Database (https://www.iedb.org/). The following studies were considered to specify immunodominant IgE-binding epitopes[34-45]. Ara h 1 peptides that were contiguous or overlapping were considered distinct if their cut-off points differed. Peptides not fully aligned with the canonical sequence were required to have at least 75% alignment and a minimum of four consecutive identical amino acids Molecular structure analysis was performed with Chimera v1.17.3 for tridimensional visualization of allergens and relevant epitopes before and after bacterial digestion.

Mice sensitization to Ara h 1 and Ara h 2

Mice were sensitized to immunodominant PN allergens to generate sera specific to Ara h 1 or Ara h 2 specific IgE to be used in mast cell degranulation assays (above). To sensitize mice against native Ara h 1 (InBio) or recombinant Ara h 2 (prepared as reported[110] using Uniprot reference Q6PSU2-1 without the signal peptide), 3 i.p. injections with 10 μg of the allergen plus 1 mg aluminum hydroxide (Alhydrogel® adjuvant 2%, InvivoGen) were performed. Previously, aluminum hydroxide and allergen were mixed at 4ºC during 30 min in a ferris wheel. Between the first and second injection, 2 weeks were left; and 1 week between the second and third injection. Blood samples were collected and centrifuged at 16,000 RCF to obtain sera for mast cell activation assays (see below) and Western blot analysis. Control sera from PN-allergic mice were generated following immunization with a classical model of food allergy.[47,71]

1555

Bone marrow-derived mast cell culture

BMMCs were cultured to assess mast cell activation by bacterially digested PN allergens. Bone marrow from femur and tibia of C57BL/6 (Charles River) mice were used to obtain BMMCs as described[111]. In short, bone marrow was flushed out inserting a 23-gauge needle (BD Microlance) attached to the 10 mL syringe filled with PBS (Gibco) at the knee side of both types of bone. Then, the cell suspension collected was filtered with a 40 µm filter (Falcon). After centrifugating 5 min at 286 RCF and RT, the pellet was lysed with ACK lysing buffer (Lonza). Erythrocyte-lysed bone marrow cells were cultured for 4 weeks in Petri dishes at 37ºC and 5% CO_2 in Iscoves´s Modified Dulbecco´s Medium (IMDM, Gibco) supplemented with 10% fetal bovine serum (FBS, Cytiva), 1% minimal essential medium (MEM, Gibco), 1% sodium pyruvate (Biowest), 1% penicillin/streptomycin (P/S, Gibco), 1% non-essential amino acids (Biowest), 0.01% recombinant murine stem cell factor (SCF, PeproTech) and 0.05% recombinant murine Interleukin (IL-3, PeproTech), until they differentiated into mature BMMCs. Cell growth and viability were monitored using the trypan blue exclusion test. The culture media was changed weekly, and the cellularity was kept below 0.5×10^6 cells/mL.

BMMC activation

BMMC activation was conducted as reported[46,47]. In brief, BMMCs were sensitized o/n at 37ºC and 5% CO_2 with sera at a concentration range of 15–40 ng/mL from Ara h 1- or Ara h 2-sensitized mice. Culture media was changed 1-2 days previous sensitization. PN-allergic mice sera were used as a positive control. BMMC sensitization was done at a cell density of 1×10^6 cells/mL. The next day, BMMCs were washed to eliminate unbound Igs (cells were spun at 265 RCF for 5 min and low break) and resuspended in supplemented IMDM without cytokines, nor FBS, at a cell density of 1×10^6 cells/mL. For BMMC activation, bacteria with PN-degrading potential isolated from saliva and gut of healthy donors were selected. Then, 0.1×10^6 BMMCs per condition were placed in a U/V bottom 96-well culture plate (Falcon) and challenged with bacterially digested PN (Stallergenes Greer) at 25 µg/mL and 100 µg/mL in Ara h 1-sensitized BMMCs, or at 50 µg/mL and 200 µg/mL in Ara h 2-sensitized BMMCs. Media and undigested PN were used as negative and positive controls, respectively. After 20 min at 37ºC and 5% CO_2, the reaction was stopped on ice and BMMC activation was determined via β-hexosaminidase activity in the supernatants, and phenotypically (CD63 and CD107a expression) by flow cytometry[46].

β-hexosaminidase activity assay

BMMCs were centrifugated as above described to recover cell-free supernatants. A volume of supernatant of 45 µL was added in duplicates to 45 µL of β-hexosaminidase substrate solution (2 mM p nitrophenyl N-acetyl β-D-glucosamine, Sigma-Aldrich) diluted

in 0.1 M citrate buffer (45 mM dehydrated sodium citrate, 55 Mm citric acid, Sigma-Aldrich,
in distilled H_2O) in a flat-bottom 96-well plate. After incubation at 37ºC for 45 min in the
dark, 45 µL NaOH 1 M (1M sodium hydroxide, Sigma-Aldrich, in distilled H_2O) were added
to stop the reaction. OD was measured at 405 nm with a microplate reader. Buffers alone
and BMMC lysates (0.5% Triton X-100, Sigma Aldrich, was used for cell lysis) were used
as control, absorbance background (B) and total lysis, respectively. The percentage of
BMMC degranulation, based on β-hexosaminidase activity, was calculated as follows:

1602
$$\text{Degranulation (\%)}=\frac{\text{OD supernatant-B}}{\text{OD total lysis-B}}\times100$$

Flow cytometry
BMMCs were resuspended in ice-cold FACS buffer (2.5 mM EDTA, PanReacAppliChem,
0.5% BSA in PBS). BMMCs were blocked with 1:50 Fc block (purified anti-mouse
CD16/32, BioLegend) for 15 min on ice to prevent non-specific antibody binding. Then,
BMMCs were incubated with BV421 rat anti-CD117 (c-kit) (BioLegend) 1/200; APC rat
anti-CD63 (BioLegend) 1:200; PerCP-Cy5.5 rat anti-CD107a (LAMP-1) (BioLegend)
1:200; PE rat anti-IgE (BioLegend) 1:200; PE-Cy7 rat anti-FcεR1α (BioLegend) 1:200; on
ice for 30 min covered from light. Viability was assessed with efluor780 dye (eBioscience)
1:4,000. After washing with FACS buffer to eliminate unbound antibodies, cells were
analyzed on a BD FACSCantoll flow cytometer. On average, 10,000 live and single cells
were recorded. Data were analyzed with FlowJo v10 software; dead cells and aggregates
were excluded and fluorescence minus one (FMO) controls were used for gating.
1615
Microbiota analysis
DNA was extracted from mouse fecal pellets, human saliva, and human jejunal aspirates,
and the hypervariable V3-V4 regions of the 16S rRNA gene were amplified with
polymerase chain reaction (PCR) using Taq polymerase (Life Technologies), as
previously described[81]. Forward barcoded primers targeting the V3 region (v3f_341f-
CCTACGGGNGGCWGCAG) and reverse primers targeting the V4 region (v4r_806r-
GGACTACNVGGGTWTCTAAT) were used. Forward primers included six-base pair
barcodes to allow multiplexing samples. Purified PCR products were sequenced using
the Illumina MiSeq platform by the McMaster Genomics facility. Primers were trimmed
from the obtained sequences with Cutadapt software[112,113], and processed with Divisive
Amplicon Denoising Algorithm 2 (DADA2; version 1.14.0) using the trained SILVA
reference database (version 138.1)[114,115]. A phylogenetic tree of the sequences was
calculated using FastTree 2[116], and data was explored using the phyloseq package
(version 1.30.0) in R (version 3.6.2)[117]. After data cleanup, a total of 292,618 reads were
obtained with a minimum of 11,651 and maximum of 60,790 with an average of 32,513
reads per sample for mouse microbiota. For human healthy control salivary microbiota, a
total of 273,686 reads were obtained with a minimum of 1,881 and maximum of 48,620
with an average of 16,099 reads per sample. For PN allergic patient salivary microbiota,

a total of 844,167 reads were obtained with a minimum of 3,236 and maximum of 145,324
with an average of 36,702 reads per sample. Alpha-diversity was measured using
observed species and Chao1 indices. Beta-diversity was calculated on normalized data
and the originated matrices were ordinated using principal coordinate analysis based on
Jaccard distance (mouse microbiota) and Bray-Curtis dissimilarity (human microbiota).

For whole genome sequencing of bacterial isolates, genomic DNA was extracted as
previously described[81]. Illumina short-read libraries were prepared using the NEBNext
Ultra II FS DNA Library Prep Kit for Illumina (NEB, Ipswich, MA, USA). Final libraries were
sequenced on an Illumina NextSeq 2000 platform using a P1 flow cell at the McMaster
Metagenomics Facility (Hamilton, ON, Canada). The run yielded 3.7 million paired-end (2
× 150 bp) reads per genome. Short reads were processed as previously described[118],
with few modifications. Briefly, fastp was used for quality trimming and removal of
adapters from Illumina reads[119,120]. Unicycler (v.0.5.1) was then used for de novo
assembly[121], followed by Bakta for annotation of the genomes[122]. Genomes were then
compared using Roary[123]. The data was further analysed using the MEROPs database
and R (4.4.3) to search for proteases. The data are presented as heatmaps where light
blue=gene not present and dark blue=present.

Rothia abundance (16S rRNA) data in saliva of non-PN allergic and PN-allergic (high
threshold and low threshold) individuals were obtained from a previously published PN
allergy cohort described by Zhang *et al*[51]. PN threshold was determined as described in
the primary study[51]. In this cohort, amplicon sequence variants (ASVs) belonging to
Rothia were identified after filtering for low prevalence and low abundance, and relative
abundances were quantified following taxonomic assignment against NCBI databases.
We used the *Rothia* ASV abundance values as provided by the authors for secondary
analysis in the present study.

STATISTICAL ANALYSIS
All variables were analyzed using GraphPad Prism 9 and 10 software (GraphPad
Software, USA). Parametric data are depicted as dot plots with each dot representing an
individual mouse or biological replicate. Normal distribution was determined by
D'Agostino-Pearson omnibus normality test, Shapiro-Wilk test, and Kolmogorov-Smirnov
test with Dallal-Wilkinson-Lillie correction. One way analysis of variance (ANOVA) was
used to evaluate differences between more than two groups with a parametric distribution
and Tukey's or Dunnet's post-hoc corrections were applied. Student's t-test (two-tailed)
was performed to evaluate the differences between two independent groups as
appropriate. Data with non-normal distribution were evaluated with Kruskal-Wallis test
with Dunn's post-hoc test for more than two groups. Correlation analyses were performed
using Pearson correlation for parametric data and Spearman correlation for non-

parametric data. A P value of ≤ 0.05 was selected to reject the null hypothesis. Information

regarding specific P values, value of n, and how data are presented can be found in figure

legends.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rat anti-mouse IgE (clone R35-72)	BD Pharmingen	Cat# 553413
Anti-mouse IgE-HRP (clone: 23G3)	SouthernBiotech	Cat#1130-05
Anti-human IgE-HRP	Invitrogen	Cat#A18793
Sera from Ara h 1- or Ara h 2-sensitized mice	This paper	N/A
Purified anti-mouse CD16/32 (clone: 93)	BioLegend	Cat#101302
BV421 rat anti-CD117 (c-kit) (clone: ACK2)	BioLegend	Cat#135124
APC rat anti-CD63 (clone: NVG-2)	BioLegend	Cat#143906
PerCP-Cy5.5 rat anti-CD107a (LAMP-1) (clone: 1D4B)	BioLegend	Cat#121626
PE rat anti-IgE (clone: RME-1)	BioLegend	Cat#406908
PE-Cy7 rat anti-FcεR1α (clone: MAR 1)	BioLegend	Cat#134318
Rabbit anti-mast cell tryptase	Abcam	Cat#ab151757
Bacterial and virus strains		
Rothia aeria	This paper	R1
Rothia dentacariosa	This paper	R2
Rothia mucilagenosa	This paper	R3
Staphylococcus epidermidis	This paper	S1
Staphylococcus aureus	This paper	S2
Staphylococcus aureus	This paper	S3
Biological samples		
Serum from peanut-allergic donors	Hospital Universitario de la Princesa	N/A
Saliva from Healthy controls	McMaster University	N/A
Saliva from peanut-allergic patients	Mass General Hospital	N/A
Chemicals, peptides, and recombinant proteins		
ACK Lysing Buffer	Lonza	Cat#BP10-548E
Iscoves´s Modified Dulbecco´s Medium (IMDM)	Gibco	Cat#12440-053
Fetal Bovine Serum (FBS)	Cytiva	Cat#SV30160.03
Minimal essential medium (MEM)	Gibco	Cat#11120-037
Penicillin/Streptomycin	Gibco	Cat#15140122
Non-essential amino acids	Biowest	Cat#X0557-100
Sodium Pyruvate	Biowest	Cat#L0642
Recombinant Murine Stem Cell Factor (r-mSCF)	PeproTech	Cat#250-03
Recombinant Murine Interleukin 3 (r-mIL-3)	PeproTech	Cat#213-13
Native Ara h 1	InBio	Cat#NA-AH1-1
Native Ara h 2	InBio	Cat#NA-AH2-1
Recombinant Ara h 2	Centre for Plant Biotechnology and Genomics	N/A
Aluminum hydroxide, Alhydrogel® adjuvant 2%	InvivoGen	Cat#21645-51-2
Bovine Serum Albumin (BSA)	NZYtech	Cat#9048-46-8
Tetramethylbenzidine (TMB)	Biolegend	Cat#421101
Peanut (PN; CPE)	Stallergenes Greer	Cat#XPF171D3A25
p nitrophenyl N-acetyl β-D-glucosamine	Sigma-Aldrich	Cat#N9376-100MG
0.5% Triton X-100	Sigma Aldrich	Cat#9036-19-5
Ethylenediaminetetracetic acid (EDTA)	PanReacAppliChem	Cat#131026

Efluor780 viability	eBioscience	Cat#65-0865-14
Any kD™ Mini-PROTEAN® TGX™ Precast Protein Gels	Bio Rad	Cat#4569034
Nitrocelulose membrane	Bio Rad	Cat#1620115
Protein molecular weight marker	ThermoFischer	Cat#26619
Microcon units	Sigma	Cat#UFC5003
Cholera toxin from *Vibrio cholerae* Inaba 5698	List Biological Laboratories	Cat#100B, 9100B
Brain-heart infusion (BHI) broth	Research Products International Corp.	Cat#B11000-1000
Powdered PN	PB & Me, Sahah Naturals	SKU#62845116652
Opti-MEM™ Reduced Serum Medium	Gibco, ThermoScientific	Cat#31985070
3,3-diaminobenzidiene (DAB)	DAKO	Cat#K4065
Critical commercial assays		
Mouse MCpT-1 Uncoated ELISA Kit with Plates	Thermo Scientific	Cat#88-7503-22
Ara h 1 ELISA 2.0	Indoor Biotechnologies	Cat#EPD-AH1-5
Ara h 2 ELISA 2.0	Indoor Biotechnologies	Cat#EPD-AH2-5
EZ-Link Sulfo-NHS-LC-Biotinylation Kit	ThermoScientific	Cat#21435
Mouse PLA2G7/PAF-AH/Lp-PLA2 ELISA Kit	Novus Biologicals	NBP2-81152
Deposited data		
Microbiome and proteomics data (see Star Protocols)		
Experimental models: Cell lines		
Experimental models: Organisms/strains		
Mouse:C57BL/6 (6–12 weeks of age, male or female)	Charles River Laboratory	Strain No.: 027
Mouse:C3H-HeN	Taconic Biosciences	Strain No.: C3H/HeNTac
Mouse:C57BL/6N (germ-free)	Taconic Biosciences	Strain No.: C57BL/6NTac
Oligonucleotides		
Microbiota primers F: (v3f_341f-CCTACGGGNGGCWGCAG) R: (v4r_806r-GGACTACNVGGGTWTCTAAT)	IDT DNA	N/A
Sanger sequencing primers F: (5'- AGAGTTTGATCCTGGCTCAG-3') R: (5'-CCGTCAATTCCT-TTRAGTTT-3')	IDT DNA	N/A
Recombinant DNA		
Software and algorithms		
Graph Pad Prism v 9.0	GraphPad	https://www.graphpad.com/demos/
FlowJo v10	BD	https://www.flowjo.com/solutions/flowjo
Peaks	Bioinformatics Solutions Inc.	https://www.bioinfor.com/peaksdb/
UniprotKB	Uniprot	https://www.uniprot.org/

Protein Data Bank	RCSB PDB	https://www.rcsb.org/
Allergen Nomenclature	Allergen Nomenclature. Financial contributions from IUIS, EAACI, and AAAAI	https://allergen.org/
Allergen Online	AllergenOnline. University of Nebraska-Lincoln	http://allergenonline.org/
Compare Database	Compare	https://db.comparedatabase.org/
Chimera v1.17.3	University of California San Francisco (UCSF)	https://www.cgl.ucsf.edu/chimera/
ImageJ 1.53q	Fiji	https://fiji.sc/
Illumina MiSeq	Illumina Inc.	https://www.illumina.com/
Cutadapt	Marcel Martin[1]	https://cutadapt.readthedocs.io/
DADA2 (version 1.14.0)	Benjamin Callahan[2]	https://benjjneb.github.io/dada2/
SILVA Database (version 138.1)	SILVA Consortium[3]	https://www.arb-silva.de/
FastTree 2	Morgan N. Price et al.[4]	https://www.microbesonline.org/fasttree/
Phyloseq (version1.30.0)	Paul J. McMurdie & Susan Holmes[5]	https://joey711.github.io/phyloseq/
R (version 3.6.2)	R Foundation for Statistical Computing	https://www.r-project.org/
fastp	Chen et al. [6,7]	
Unicycler (v.0.5.1)	Wick RR et al.[8]	
Bakta	Schwengers et al.[9]	
Roary	Page A et al.[10]	
Other		
Sterile Dish 100 mm x 20 mm CC-treated	Corning	Cat#430167
U/V-bottom 96 well TC-treated microplates	Falcon	Cat#353910
Flat-bottom 96 well high binding microplates	Costar	Cat#3590
23-gauge needle	BD Microlance	Cat#300800
40 µm filter	Falcon	Cat#352340
Microplate reader	Promega	GloMax Discover
SalivaBio Children's Swab	Salimetrics	Cat#5001.06
Flow cytometer	BD	FACSCantoII
Ussing chamber system	Physiologic Instruments	Cat#P2400
Imaging system	Amersham	ImageQuant™ 800
ZipTip with 0.6 µL C_{18} resin	Merck Millipore	Cat#ZTC18S008
UltiMate 3000 High Pressure Liquid Chromatograph	Fisher Scientific	Cat# IQLAAAGABHFAPBMBFB
Orbitrap Exploris™ 240 Mass Spectrometer	Fisher Scientific	Cat#BRE725535
Easy-spray PepMap C18 analytical column 50 cm × 75 µm	Fisher Scientific	Cat#ES903
Non-heparinizing microcapillaries (Fisher Scientific),	Fisher Scientific	Cat#11884040

References

1. Martin, M. (2011). Cutadapt removes adapter sequences from high-througput sequencing reads. EMBnet.journal *17*, 10-12. https://doi.org/10.14806/ej.17.1.200.
2. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., and Holmes, S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods *13*, 581-583. 10.1038/nmeth.3869.
3. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glockner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res *41*, D590-596. 10.1093/nar/gks1219.
4. Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS One *5*, e9490. 10.1371/journal.pone.0009490.
5. McMurdie, P.J., and Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One *8*, e61217. 10.1371/journal.pone.0061217.
6. Chen, S. (2023). Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp. Imeta *2*, e107. 10.1002/imt2.107.
7. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics *34*, i884-i890. 10.1093/bioinformatics/bty560.
8. Wick, R.R., Judd, L.M., Gorrie, C.L., and Holt, K.E. (2017). Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. PLoS Comput Biol *13*, e1005595. 10.1371/journal.pcbi.1005595.
9. Schwengers, O., Jelonek, L., Dieckmann, M.A., Beyvers, S., Blom, J., and Goesmann, A. (2021). Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. Microb Genom *7*. 10.1099/mgen.0.000685.
10. Page, A.J., Cummins, C.A., Hunt, M., Wong, V.K., Reuter, S., Holden, M.T., Fookes, M., Falush, D., Keane, J.A., and Parkhill, J. (2015). Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics *31*, 3691-3693. 10.1093/bioinformatics/btv421.

Figure S1. Intestinal microbiota of Altered Schaedler Flora (ASF) and specific pathogen free (SPF) C57BL/6 mice. (A) Beta-diversity (Jaccard index) plot of fecal microbiota of ASF and SPF mice where each dot represents one mouse. (B) Alpha-diversity metrics based on observed species and Chao1 index of ASF and SPF mice where each dot represents one mouse. (C) Order-level relative abundance of the microbial composition of ASF and SPF mice where each bar represents one mouse. (D) Correlation plot of Ara

h 1 (y=-0.01467x + 2.231, R^2=0.5678, P=0.0002) and Ara h 2 (y=-0.0004257x + 0.07693, R^2=0.0137, P=0.0137) against alpha-diversity in small intestinal content of germ free, ASF, and SPF mice. n=4–5 mice per group.

Figure S2. Peanut (PN) digestion by Altered Schaedler Flora (ASF) and specific pathogen-free (SPF) C3H-HeN mice. ASF and SPF C3H-HeN mice were provided a 24 mg bolus of PN i.g. Peripheral blood and small intestinal contents were collected 40 minutes post PN delivery for analysis. n=5 mice per group. PN allergens (Ara h 1 and 2) in small intestinal content (A) and in serum from peripheral blood (B). (C) Heatmap showing digestion capacity of ASF and SPF microbiota of C3H-HeN mice against Ara h 1 and 2. Small intestinal contents were subsequently incubated with PN allergens *in vitro* and remaining allergens were quantified after digestion. Allergen degradation capacity is expressed as the percentage of degraded allergen relative to control non-growth incubations (containing crude peanut extract (CPE)), with the color scale ranging from blue (0% degradation; minimum) to red (100% degradation; maximum). Each row represents one mouse. Significance levels: Ara h 1 SPF vs. ASF (*P*<0.0001), Ara h 2 SPF vs. ASF (*P*= 0.0001). Data are presented as mean where each dot represents an individual mouse (A-B), or each row represents one mouse (C). Displayed *P* values were calculated using an unpaired Student's *t*-test.

Figure S3. Body temperature reduction of Altered Schaedler Flora (ASF) and specific pathogen-free (SPF) mice in response to peanut (PN) challenge in different sensitization models, and characterization of allergic donor serum used for passive sensitization corresponding to Figure 2. (A-C) Core temperature reduction after PN challenge for intragastrically (i.g.), intraperitoneally (i.p.), and passively sensitized mice, respectively. (D) Serum platelet-activating factor-acetylhydrolase (PAF-AH) after PN challenge for passively sensitized mice. (G) PN-specific IgE and IgG1 in allergic donor serum used for passive sensitization. Data are presented as mean ± SD (A-C, E) and mean where each dot represents one mouse (D). Displayed P values were calculated using an unpaired Student's t-test.

A Gut microbiota composition and α-diversity in ASF vs. SPF mice across sensitization methods

B PN-allergen levels in ASF and SPF mice

C Relationship between Ara h 2 in intestinal content and α-diversity

D Relationship between mMCP-1 and α-diversity across sensitization methods

Figure S4. Gut microbiota composition and diversity of ASF and SPF C3H-HeN mice across sensitization methods, and their relationship with peanut (PN)-challenge outcomes (corresponding to Figure 2). (A) Relative abundance of bacterial orders and α-diversity (observed species) in the small intestine across intragastric (i.g.), intraperitoneal (i.p.), and passive sensitization, including sensitized non-sensitized mice. (B) Levels of Ara h 1 and 2 in intestinal content across sensitization methods, including sensitized and non-sensitized mice. (C) Correlation plot of Ara h 1 against α-diversity in small intestinal content (y=-0.001039x + 0.07793, R^2=0.6736, P<0.0001). (D) Relationship between

serum mucosal mast cell protease 1 (mMCP-1) concentrations and α-diversity across sensitization methods.

Figure S5. Characterization of peanut (PN)-degrading bacteria isolated from human saliva and small intestine. (A) Heatmap of PN allergen degradation capacity of bacterial strains isolated from saliva. (B) Degradation of Ara h 1 and 2 by bacterial strains that were further characterized in Figures 4 and 5 for their proteomic profiles and effects on mast cell activation. (C) Heatmap of PN allergen degradation capacity of bacterial strains isolated from the small intestine (jejunal aspirates). Color scale (blue-red) represents the percentage of allergen degradation relative to control non-growth incubations, with blue indicating no degradation and red indicating complete degradation. For (C), the relative abundance of each isolated strain is shown using a green scale, ranging from light green (low abundance) to dark green (high abundance), calculated as a percentage of total isolates from saliva.

Figure S6. Characterization of sera specificity from mono-sensitized mice. (A) Gating strategy. Bone marrow-derived mast cells (MCs) co-express FcεRI and CD117. MC activation was detected by the expression of CD63 and CD107a. Gates of positive and negative controls are shown. Controls of sera specificity to (B) Ara h 1 and (C) Ara h 2. MCs were sensitized with a pool of sera from Ara h 1- or Ara h 2-allergic mice, respectively, and then challenged with undigested PN (blue), native Ara h 1 (white) or recombinant Ara h 2 (grey). Data are presented as the mean +/- SEM of each group. Displayed *P* values were calculated using a one-way ANOVA with Tukey's post-hoc test in panels B and C (CD107a and β-hexosaminidase activity) or using Kruskal-Wallis test with Dunn´s post hoc test in panel C (CD63).

Protease repertoire of *Rothia* species identified by whole genome sequencing

Species columns (left to right): *Rothia muciaginosa* **R3**, *Rothia muciaginosa*, *Rothia dentocariosa* **R2**, *Rothia dentocariosa*, *Rothia aeria* **R1**, *Rothia kristinae*

Intracellular

Gene	Annotation
pepB	leucyl aminopeptidase
pepN	aminopeptidase N
pepP	Xaa-Pro aminopeptidase
pepO	peptidase M13
map	type I methionyl aminopeptidase
pqqL	peptidase M16
dacB	D-alanyl-D-alanine carboxypeptidase/D-alanyl-D-alanine-endopeptidase
htpX	Zn-dependent protease with chaperone function
argE	dipeptidase
degQ	MarP family serine protease
lepB	signal peptidase I
lspA	signal peptidase II
IAP4	M18 family aminopeptidase
CPBP family intramembrane metalloprotease domain-containing protein	CPBP family intramembrane metalloprotease domain-containing protein
ptrB	oligopeptidase B
pcp	peptidase C15
peptidase E	peptidase E
peptidase A24	peptidase A24
glpG	rhomboid family intramembrane serine protease
ATP-dependent protease Clp, ATPase subunit	ATP-dependent protease Clp, ATPase subunit
clpS	ATP-dependent Clp protease adapter ClpS
clpX	ATP-dependent Clp protease ATP-binding subunit ClpX
clpP	ATP-dependent Clp protease proteolytic subunit
ftsH	ATP-dependent zinc metalloprotease FtsH
protease/lipase ABC transporter permease/ATP-binding protein	protease/lipase ABC transporter permease/ATP-binding protein
immA	ImmA/IrrE family metallo-endopeptidase
peptidase M48	peptidase M48
M23 family metallopeptidase	M23 family metallopeptidase
prepilin peptidase	prepilin peptidase
Membrane protein related to metalloendopeptidase	Membrane protein related to metalloendopeptidase
Periplasmic protease	Periplasmic protease
nlpD	Murein DD-endopeptidase MepM and murein hydrolase activator NlpD, contains LysM domain
pepE	peptidase S51
tonB	serine protease
rseP	zinc metalloprotease
Peptidase C39 domain-containing protein	Peptidase C39 domain-containing protein
CAAX amino terminal protease family protein	CAAX amino terminal protease family protein
CAAX protease self-immunity	CAAX protease self-immunity
M48 family peptidase	M48 family peptidase
Peptidase M20 domain-containing protein 2	Peptidase M20 domain-containing protein 2
Endopeptidase La	Endopeptidase La
Site-2 protease family protein	Site-2 protease family protein
Peptidase family M50	Peptidase family M50
Peptidase-M3 domain-containing protein	Peptidase-M3 domain-containing protein
L,D-transpeptidase	L,D-transpeptidase
Peptidase-A24 domain-containing protein	Peptidase-A24 domain-containing protein
lspA	Lipoprotein signal peptidase

Extracellular

Gene	Annotation
peptidase S8	peptidase S8
group_294	peptidase S8
group_1522	peptidase S8
group_2124	Peptidase S8
group_2125	peptidase S8
Peptidase S8	Peptidase S8
aprE	peptidase S8
acm	serine protease like S8
group_5120	peptidase S8
Subtilisin	peptide S8
Peptidase-S8 domain-containing protein	Peptidase-S8 domain-containing protein
eMpr	V8-like Glu-specific endopeptidase
Minor extracellular protease vpr	Minor extracellular protease vpr
Extracellular basic protease	Extracellular basic protease

Both/Unknown

Gene	Annotation
Di- and tripeptidase	Di- and tripeptidase
prolyl aminopeptidase	prolyl aminopeptidase
serine protease	serine protease
Peptidase S1 domain-containing protein	Peptidase S1 domain-containing protein
Peptidase-M78 domain-containing protein	Peptidase-M78 domain-containing protein
group_2044	peptidase
group_2073	peptidase
zinc metalloprotease	zinc metalloprotease
Peptidase C51 domain-containing protein	Peptidase C51 domain-containing protein
Peptidase	Peptidase
group_2882	Peptidase
group_2894	peptidase
group_2945	peptidase
peptidase	peptidase
group_2952	peptidase
group_2963	peptidase
group_2964	peptidase
Zn-dependent proteases	Zn-dependent proteases
group_5239	zinc metalloprotease
group_6644	serine protease
X-Pro dipeptidyl-peptidase (S15 family) domain protein	X-Pro dipeptidyl-peptidase (S15 family) domain protein
Peptidase-M23 domain-containing protein	Peptidase-M23 domain-containing protein
Peptidase C56	Peptidase C56
Peptidase-S9 domain-containing protein	Peptidase-S9 domain-containing protein
Pyroglutamyl-peptidase I	Pyroglutamyl-peptidase I
Zinc-dependent metalloprotease	Zinc-dependent metalloprotease

Gene presence

No   Yes

Degradation scale

Min   Max

Ara h 1

Ara h 2

Allergen degradation capacity
(western blotting)

Figure S7. Protease repertoire of *Rothia* species identified by whole-genome sequencing. Heatmap showing predicted proteases encoded in *Rothia* genomes, including *Rothia mucilaginosa* (R3), *Rothia dentocariosa* (R2), and *Rothia aeria* (R1), as well as additional species without specific isolate names. Rows represent protease genes grouped by subcellular localization (intracellular, extracellular, or both/unknown) with functional annotations, and columns correspond to individual *Rothia* genomes. Blue shading indicates gene presence (light blue: not present, dark blue: present). Green-highlighted labels denote duplicate recognition of the same protease across different sequence annotations. Allergen degrading capacity against Ara h 1 and Ara h 2, determined by Western blotting, is shown at the bottom right for reference and the heat map colour scale (blue-red) represents the percentage of allergen degradation relative to control non-growth incubations (crude peanut extract).

Figure S8. Protease repertoire of *Staphylococcus* species identified by whole-genome sequencing. Heatmap showing predicted proteases encoded in *Rothia* genomes, including *Staphylococcus epidermidis* (S1), *Staphylococcus aureus* (S2), and *Staphylococcus aureus* (S3). Rows represent protease genes grouped by subcellular localization (intracellular, extracellular, or both/unknown) with functional annotations, and columns correspond to individual *Staphylococcus* genomes. Blue shading indicates gene presence (light blue: not present, dark blue: present). Green-highlighted labels denote duplicate recognition of the same protease across different sequence annotations. Allergen degrading capacity against Ara h 1 and Ara h 2, determined by Western blotting,

is shown at the bottom right for reference and the heat map colour scale (blue=0%, red=100%) represents the percentage of allergen degradation relative to control non-growth incubations (crude peanut extract).

Figure S9. (A-D) Peanut (PN) intragastric sensitization and challenge in ASF-microbiota mice colonized with or without *Rothia* R3. (A) Experimental design. C3H-HeN mice with altered Schaedler flora (ASF) microbiota sensitized i.g. to PN once per week for 6 weeks.

One week after the last sensitization, mice were co-colonized with *Rothia* R3 once per week for three weeks. Mice were provided a PN bolus i.g. three weeks after the first colonization and sacrificed 40 minutes later. n=7 mice per group. (B) Serum platelet activating factor-acetylhydrolase (PAF-AH). (C) Tryptase-positive mast cells quantified in small intestinal sections. (D) Representative images of small intestinal mucosal tryptase-positive mast cells. (E) Small intestinal microbiota composition (order). (F-H) PN i.p. sensitization and challenge in ASF-microbiota mice colonized with or without *Rothia* strains. (F) Experimental design. C3H-HeN mice with ASF microbiota sensitized i.p. to PN once per week for 4 weeks. One week after the last sensitization, mice were co-colonized with *Rothia* strains once per week for three weeks. Mice were provided a PN bolus i.g. three weeks after the first colonization and sacrificed 40 minutes later. n=7 mice per group. (G) Serum PN-specific IgE before PN challenge. (H) Serum mMCP-1 after PN challenge. (I) Tryptase-positive mast cells quantified in small intestinal sections. (J) Representative images of small intestinal mucosal tryptase-positive mast cells.Displayed *P* values were calculated using an unpaired Student's *t*-test.

Figure S10. (A-B) Peanut (PN)-allergic patients (n=19) were tested for PN threshold using increasing PN concentrations before enrolment in oral immunotherapy. (A) Correlation plot of Ara h 1-specific IgE (y= -0.09x + 42.45, R^2 = 0.15, P=0.11) and Ara h 2-specific IgE (y= -0.11x + 58, R^2= 0.21, P=0.06) against PN threshold. (B) Bray-Curtis-tomrepresents serum Ara h 2-specific IgE levels, while dot color corresponds to the PN threshold (scales shown). (C) Abundance of *Rothia* species in saliva from analysis of 16s rRNA data from an external cohort including 120 children (23 non-PN-allergic controls (NPA), 74 PN-allergic with high PN threshold (HT; >443 mg), and 23 PN-allergic with low PN threshold (LT; <443 mg)) who underwent double-blind, placebo-controlled PN challenges. Data are presented as interquartile range (IQR) with whiskers extending to the min and max data points.

Table S1: Detailed subject demographics. Peanut, PN; Male, M; Female; F.

Status	ID	Sex	Age	Race	PN threshold (mg)	Ara h 1 IgE (kUA/mL)	Ara h 2 IgE (kUA/mL)
PN-allergic saliva	PN1	F	10	Unknown	300	0.10	1.19
	PN2	F	7	Asian	300	0.10	2.44
	PN3	F	1	White	300	0.10	0.10
	PN4	F	27	White	440	15.2	13.8
	PN5	F	6	White	375	20.3	25.7
	PN6	M	6	White	40	11.6	0.41
	PN7	M	7	White	300	N/A	N/A
	PN8	F	15	White	440	17.3	12.1
	PN9	M	9	White	12	100	25.8
	PN10	M	13	White	6	9.11	0.19
	PN11	M	10	White	43	18	88.5
	PN12	F	13	White	13	100	58.2
	PN13	F	13	White	43	100	100
	PN14	F	14	White	12	100	96.7
	PN15	M	9	White	6	5.95	20.7
	PN16	M	11	Asian	160	7.62	9.92
	PN17	F	2	Asian	80	1.41	0.84
	PN18	F	2	Black/African American	15.5	0.18	40.1
	PN19	M	4	Unknown	12	16.7	100
Non-PN-allergic saliva	HV1	M	35	White	N/A	N/A	N/A
	HV2	M	22	White			
	HV3	M	20	White			
	HV4	F	26	White			
	HV5	F	33	White			
	HV6	F	24	White			
	HV7	F	18	White			
	HV8	M	30	White			
	HV9	F	38	White			
	HV10	F	20	Black/African American			
	HV11	M	40	Asian			
	HV12	F	66	White			
	HV13	M	33	White			
Non-PN-allergic jejunal washes	HV14	M	57	White	N/A	N/A	N/A
	HV15	M	63	White			
	HV16	F	57	White			
	HV17	F	35	White			
	HV18	M	53	White			

Table S2: Serum characterization for western blotting. Peanut, PN.

Serum	Total IgE levels (kU/L)	PN IgE (kUA/mL)	rAra h 1 IgE (kUA/mL)	rAra h 2 IgE (kUA/mL)	rAra h 3 IgE (kUA/mL)	rAra h 8 IgE (kUA/mL)	rAra h 9 IgE (kUA/mL)	Sex	Age	Positive Skin Prick Test	Other hypersensitivities
A	12.49	Unknown	52.9	>100	26.6	0.02	0.02	F	30	PN, pollen of Cupressaceae, Acer pseudoplatanus, olive, grasses, Salsola, dog, cat, horse and rabbit. Minimal reaction to mites and aspergillus	Persistent moderate asthma and intermittent rhinoconjunctivitis
B	176	54.9	20.4	24.5	0	0	0	F	19	PN, chickpea, arizonica pollen, minimal reaction to pine nuts	Atopic dermatitis