1 **Warning signals for the long-term innovative potential of**
2 **genetic research based on 20 million papers and patents**
3

4 Junghun Chae[1,16], WooJoong Kim[1,16], Woochul Jung[2,16], Dawoon Jeong[3–6,16],
5 Roktaek Lim[1,7,16], Manoj Chamlagain[7], Giju Jung[2], Juneil Jang[2,8], Jae Won Lee[9,10],
6 Nam Kyu Kang[10], Kwangryul Baek[10], Jonghyeok Shin[10], Ye-Gi Lee[9,10], Hyun Gi Koh[10],
7 Chanwoo Kim[9,10], Sangdo Yook[9,10], Allen Ka Loon Cheung[7], Yong-Su Jin[9,10],
8 Hyejin Youn[5,6,11–13,*], Pan-Jun Kim[7,14,15,*], and Cheol-Min Ghim[1,2,15,*]
9

[1]Department of Physics, Ulsan National Institute of Science and Technology, Ulsan 44919,
Republic of Korea
[2]Department of Biomedical Engineering, Ulsan National Institute of Science and Technology,
Ulsan 44919, Republic of Korea
[3]Technology Management, Economics and Policy Program, Seoul National University, Seoul
08826, Republic of Korea
[4]Institute of Engineering Research, Seoul National University, Seoul 08826, Republic of Korea
[5]Kellogg School of Management, Northwestern University, Evanston, IL 60208, USA
[6]Northwestern Institute on Complex Systems, Evanston, IL 60208, USA
[7]Department of Biology, Hong Kong Baptist University, Kowloon, Hong Kong
[8]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA
[9]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign,
Urbana, IL 61801, USA
[10]Department of Food Science and Human Nutrition, University of Illinois at Urbana−Champaign,
Urbana, IL 61801, USA
[11]Department of Civil and Environmental Engineering, Northwestern University, Evanston, IL
60208, USA
[12]London Mathematical Lab, London WC2N 6DF, UK
[13]Santa Fe Institute, Santa Fe, NM 87501, USA
[14]Center for Quantitative Systems Biology & Institute of Computational and Theoretical Studies,
Hong Kong Baptist University, Kowloon, Hong Kong
[15]Asia Pacific Center for Theoretical Physics, Pohang, Gyeongbuk 37673, Republic of Korea
[16]J.C., W.J.K., W.J., D.J., and R.L. contributed equally to this work.
33
34 *Correspondence:
35   hyejin.youn@kellogg.northwestern.edu (H.Y.); extutor@gmail.com (P.-J.K.);
36   cmghim@unist.ac.kr (C.-M.G.)

Abstract

Current technological revolutions, involving artificial intelligence, mRNA vaccines, and quantum computing, are largely driven by industry. Despite the existing perception that commercial motives promote cutting-edge innovation, concerns may arise about their risk of limiting scientific exploration from diverse perspectives, which nurtures long-term innovative potential. Here, we investigate the interplay between scientific exploration and industrial influence by analyzing about 20 million papers and US, Chinese, and European patents in genetic research, a domain of far-reaching societal importance. We observe that research on new genes has declined since the early 2000s, but the exploration of novel gene combinations still underpins biotechnology innovation. Fields of highly practical or commercial focus are less likely to adopt the innovative approaches, exhibiting lower research vitality. Additionally, continuous scientific research creates exploratory opportunities for innovation, while industry's R&D efforts are typically short-lived. Alarmingly, up to 42.2–74.4% of these exploratory opportunities could be lost if scientific research is restrained by industry interests, highlighting the cost of over-reliance on commercially-driven research. Given the industry's dominance in recent technologies, our work calls for a balanced approach with long-term scientific exploration to preserve innovation vitality, unlock the full potential of genetic research and biotechnology, and address complex global challenges.

Introduction

Groundbreaking innovation emerges as the resonance of science and technology [1–3]. From the landmark developments at the Palo Alto Research Center (PARC) and Bell Labs to today's artificial intelligence (AI), robotics, mRNA vaccines, and quantum computing [4–8], industrial entrepreneurship has powered technology innovation. Given the particular dominance of industry in the ongoing technological shifts, it is imperative to discuss the positive and negative impacts of this dominance on the course of research and development (R&D) [6–8]. Ethical and regulatory aspects aside, different perspectives may exist for the lasting effects on the innovation itself [9–16], in relation to the following stories.

Recalling the history of biotechnology, the Human Genome Project decades ago was primarily led by government agencies with widespread support from academia, but Celera Genomics from the private sector played a pivotal role in accelerating the process [16]. On the other hand, commercially-supported research is vulnerable to cuts or compromise upon changes in management conditions. The storied Bell Labs, an industrial laboratory once

known as an innovation powerhouse across fundamental and applied sciences, could not

maintain its standing amid company downturns [5].

More broadly, innovative attempts can face innovation barriers such as cost-

effectiveness, particularly under commercial focus in favor of early profitability [6, 11–13].

This situation reflects the conflict between entrepreneurial risk-taking and economic

rationality, and the latter can exert pressure on innovators to deliver quick and profitable

marketable outcomes. This pressure, which we refer to as commercial pressure here, tends

to discourage long-term exploration efforts and promote risk-averse approaches against

uncertain returns and tight regulations [9–13]. It may in turn limit the spectrum of approaches

from diverse perspectives and ultimately harm the long-term vitality of innovation itself, as

our analysis will later indicate. In parallel to this issue, concerns have already been raised

about the decline of innovative research output across academic disciplines and industrial

sectors [17, 18]. Notably, it is the risk-averse nature of pharmaceutical firms that fosters

underinvestment in groundbreaking innovation and thereby suppresses novelty in drug

development [10, 19]. Besides, a recent COVID-19 R&D study suggests that non-monetary

incentives can actually be more common drivers of innovation [9].

As a crucial step to facilitate further discussions, we here examine a gap between the

activities of scientific exploration and industry-driven technology development from the

aspect of commercial pressure and innovation vitality, unlike the previous data-driven

studies that often focused on the flow of scientific or technological knowledge into future

innovation [20–23]. We target the empirically observed patterns in the realm of

biotechnology—specifically, genetic research. The recent year (2023) marked both the 70th

anniversary of the discovery of the DNA double helix and the 20th anniversary of the

completion of the Human Genome Project. Biotechnology has extraordinary impacts on

modern society, with an undeniable role in overcoming the COVID-19 pandemic. Especially,

research on genes and gene products forms the foundation of modern biotechnology and is

recognized for its applicability in medicine, agriculture, food industry, energy supply,

environmental remediation, and grand challenges such as climate change [16, 19, 24–26].

For the sake of simplicity in this study, both genes and gene products (proteins) will

henceforth be referred to as genes, unless specified.

103
104   **Results**
105
106   **Popular genes in scientific research and technology inventions**
107
We traced research of 53,484 different genes for protein production through the systematic

survey of 16 million papers and 2 million patent families, the latter from the U.S. Patent and

Trademark Office (USPTO), the China National Intellectual Property Administration (CNIPA),

and the European Patent Office (EPO). The genes were curated from the UniProt

Knowledgebase (UniProtKB)/Swiss-Prot (Methods) [27]. Our analysis focuses on the periods

of 1990–2021 for paper publications (i.e., until the COVID-19 pandemic era) and 1990–2020

for patent applications. These periods are chosen because the Human Genome Project was

launched in 1990 as a milestone in the history of genetic research, whereas the patent

application records after about 2020 did not seem fully accessible at the time of this

manuscript preparation (Methods).

Like previous studies, we view papers as carriers of scientific research and patents as

carriers of technology inventions [20–22, 28–31]. Accordingly, we obtained the annual

counts of papers and patents for each gene as the activity levels of scientific research and

technology inventions with that gene, respectively (Methods). Because not all genes in the

full text of a given document might directly contribute to its subject, we only considered

genes in the title and abstract. According to our analysis, a patent is typically company-

assigned with a 76.0% chance for each gene, whereas a paper rarely includes company-

funded research, with only a 5.2% chance for each gene (Methods). In other words, the

patent-filed inventions can be viewed primarily as industry-driven, but the paper-published

research is not.

One example of our analysis is that the research on hemoglobin grew until 2021 with five

times as many paper publications as in 1990. Similarly, hemoglobin-involving inventive

activities grew with ten times as many patent applications as in 1990 (figure 1A). The

examples of CRISPR associated protein 9 (Cas9), angiotensin-converting enzyme 2 (ACE2),

and cyclooxygenase-2 (COX-2) are presented in figure 1B–D. Intriguingly, ACE2 is a binding

target of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) to enter host cells

and cause COVID-19 [32, 33], and as expected, its research and inventive activities

skyrocketed from 2020 (figure 1C). On the other hand, COX-2 showed the decline of the

inventive activities in the 2000s (figure 1D), and this case will be elaborated on later. As an

alternative way, among the multiple genes in a given paper or patent, we identified only the

genes of the main focus in that paper or patent (Methods). In this study, when analyzing only

those main genes, we will explicitly state it.

We found that interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF-α) are the top-

ranked genes in the cumulative volumes of scientific research (figure 1E). The top ten genes

for research included many immune genes, several cancer-related genes such as tumor

suppressor p53 (TP53) and epidermal growth factor receptor (EGFR), and others (figure

1E). In the case of genes in inventive activities, serum albumin and EGFR were the top-

ranked, but their positions were overtaken by Cas9 and programmed cell death protein 1

(PD-1) during 2016–2020 (figure 1*F*). PD-1 and programmed cell death ligand 1 (PD-L1)

together came to the front in this period and they are both immunotherapy-related genes.

Similarly, the top genes among only the genes of the main focus in papers or patents are

presented in electronic supplementary material, figure S1 (see also Methods).

150
151
152

153
154

Figure 1. Genetic research trend since the launch of the Human Genome Project (1990). *A*–*D*,

Annual counts of papers (blue) or patents (red) whose titles/abstracts include a given gene. *E*, *F*, Ten

most frequent genes in papers or patents of all time (1990–2021 for papers and 1990–2020 for

patents) and latest five years (2017–2021 for papers and 2016–2020 for patents). The reason for

selecting different periods for paper and patent counting is explained in Methods. The vertical width of

each ribbon is proportional to the paper or patent number of the respective gene in that period.
161
162

Beyond those prominent genes, our analysis expands to all other genes for a

comprehensive view of genetic research, as will soon be unfolded here.

Research on new genes slows down

One may suppose that the vitality in genetic research can be partially related to the pace of

studying new genes. We observe that the total number of new genes in papers and patents

each year, which were not homologous to any genes in prior studies, grew until 2002 and

then turned to robust decline (figure 2*A* and Methods). A similar tendency has also been

observed in a previous study with only human protein-coding genes [26]. Consequently, the

exploration of new genes in research and inventive activities continued to slow down soon

after the turn of the millennium, which saw the announcement of the first draft human

genome and the near-end of the Human Genome Project. Compared to 2002, the new

genes fell into one-third in 2021 (figure 2*A*). Despite this overall tendency, an exception is

the continuous increase of new genes in Chinese patents each year, unlike US and

European patents (electronic supplementary material, figure S2*A–D*). This exception

resulted from the unusual growth of the total annual patents themselves from China. Still,

whichever US, Chinese, or European patents, the *proportion* of patents with new genes (i.e.,

the probability of new gene occurrence per patent) began to decline just before 2000 and

finally recorded 3.3–5.4% (electronic supplementary material, figure S3). Such a declining

trend and the other key results throughout this study remain unchanged when we only

consider the genes of the main focus in papers or patents (Methods). The overall decline of

newly researched genes calls to mind the pre-existing concerns about conventionally-biased

human gene studies and drug developments [10, 19].

For further analysis of this trend, we categorized genes according to their frequent

research themes. Mainly based on the expert-examined classifications of journals and

patents as the venues of genetic research, each gene was assigned to an appropriate

thematic category (electronic supplementary material, data S1–S3 and Methods). As will be

clear, commercially-oriented categorical genes exhibit some unique patterns amid the overall

new gene decline. While newly researched genes in medical science mirror the

aforementioned post-millennium decline trend, new genes in the field of metabolic

engineering underwent an earlier and severer decline, even dating back to 1992 (figure

2*B*,*C*). Of note, metabolic engineering is based on commercial and economic motives to

enhance the desirable performance of living organisms by altering their biochemical

pathways [34, 35]. Likewise, its parent field, applied biotechnology shows an early decline of

newly researched genes (see electronic supplementary material, figure S4 for all the gene

categories). We will revisit these issues later. On the other hand, those in plant science just

started to decline after 2008 (figure 2*D*). Interestingly, 2008 was close to the end of the NSF-

funded 2010 Project that aimed to determine the function of as many genes in model plant

Arabidopsis thaliana as possible [36]. From 2017, the further decline of new genes in plant

science stopped, mainly by the contributions of the genes related to *Arabidopsis* (48.5%)

and rice (16.6%) (figure 2*D* and Methods).

Figure 2. Research on new genes. *A*, Annual counts of new genes in papers and patents. *B–D*,

Annual counts of new genes in each thematic gene category (Methods). *E*, Output of our

mathematical model for the annual counts of newly researched genes. A heatmap shows the relative

errors of the annual counts of the new genes between the model output and actual data over

parameters τ (horizontal) and β (vertical), while the model assumes that fraction β of newly reported

amino acid (AA) sequences would be studied sometime later with characteristic latent period τ

(Methods). For particular τ and β with the small relative error, the heatmap is projected to the graph of

the cumulative number of the studied genes as a function of the number of their non-redundant AA

sequences, i.e., individual gene copies (left) and the graph of the annual counts of newly studied

genes (right), which is relevant to *A*: the linear interpolation of the model output (solid line) and the

real data points (circle) are shown together in these graphs. Over a broad range of τ and β, the model

produces the quantitative pattern in *A*.

To account for the overall decline of new genes from 2002 in figure 2*A*, we developed a

mathematical model that describes the cumulative number of distinct genes in papers and

patents over time (Methods). In this model, an effectively finite pool of genes and their

copies across species becomes a target of current-day genetic research. For these gene

copies, we considered all the possible orders of their initial studies with approximately equal

probabilities, and then calculated the expected cumulative number of the distinct studied

genes each year in combination with annually-updated gene copy numbers from the

UniProtKB/Swiss-Prot data. This model accurately reproduces figure 2*A* and gives other

results compatible with empirical data over a broad range of conditions (figure 2*E*, electronic

supplementary material, figure S5, and Methods; $P < 10^{-4}$). Our model suggests that the vast

majority (71.3 ± 5.4%) of taxonomically-dispersed genes studied as of 2021 had already

been studied at least once before 2002 (avg. ± s.d. and Methods). In the case of

taxonomically-restricted genes [37, 38] as of 2021, only their 8.0 ± 0.8% had been studied

before 2002, though (avg. ± s.d.). Hence, the time around the completion of the Human

Genome Project was a turning point with the substantial accumulation of the researched,

taxonomically-dispersed genes, which then rarified the exploration of the new ones.

238
239 **Novel gene combinations keep actively explored, but not in every field**
240
Despite the significance of new genes, a more common mode of novel genetic research

could be the use of genes in new combinations. In stark contrast to the slowdown of

exploring new genes from 2002, new combinations of genes tended to increase until recently

(figure 3*A* and Methods). There exist some variations between different patent sources: from

the 2000s, the numbers of novel gene combinations in Chinese and US patents were

accelerating and running steadily, respectively. Still, none of US, Chinese, and European

patents entered a decline phase in novel gene combinations and they all remained active for

those combinations (electronic supplementary material, figure S2*E–P*).

Regarding different thematic categories of genes, we observe an increase in novel gene

combinations in medical science and a similar but recently weaker trend in plant science.

About commercially-oriented categories, we observe a contrast in metabolic engineering and

applied biotechnology with drop-offs and stagnancy in the novel gene combinations from

2017, possibly warning for their low innovation vitality (figure 3*B–D*, and electronic

supplementary material, figure S6 for all the gene categories; see also Methods).

Back to the overall trend beyond regional and categorical patterns, this "innovation by

combination" [22, 39–41] involves the succession of highly frequent genes in those

combinations over time: the most frequent in the newly paired genes were β-galactosidase

in 1990–1997, green fluorescent protein (GFP) in 1998–2015, and Cas9 in 2016–2021,

along with less yet highly frequent genes such as mammalian target of rapamycin (mTOR)

and PD-L1 (figure 3*E–H* and electronic supplementary material, table S1). For example,

Cas9 newly teamed with 4,141 different genes between 2016 and 2021. We view the genes

of such frequent fresh partnership as have been the workhorses of biotechnology innovation,

recapitulating the history of landmark ideas and technologies from DNA cloning (β-

galactosidase) and live cell imaging (GFP) to genome editing (Cas9) (figure 3*E–G*) [42, 43].

Figure 3. Research on novel gene combinations. *A*, Annual counts of new pairs of genes in papers

and patents. For separate counting of paper and regional patent cases, and for larger gene

combinations beyond a pair, see electronic supplementary material, figure S2*E–P*. ***B–D*,** Annual

counts of new pairs of genes in each thematic gene category (Methods). ***E–G*,** For each gene that

once had the highest share (relative frequency) in new pairs of genes, its share is presented over

time. Shaded is the period when this gene had the highest shares. ***H*,** The shares of PD-L1 and

mTOR in new pairs of genes over time. They were not the highest but still ranked high. ***I*,** The

proportion of genes in each thematic gene category, paired with Cas9 or GFP until the peak time of its

annual paper and patent counts. In this pairing, we considered genes that debuted in papers/patents

in 1990–2000 (left) and 2001–2011 (right), separately (Methods). Abbreviations: med. sci., medical

science; pharma., pharmaceutical; metab. eng., metabolic engineering; plant sci., plant science.

Among the above prominent genes of biotechnology innovation, Cas9 and GFP are
recognized for their utility across disparate domains such as medical and plant sciences.
Therefore, we examined how many genes in each category teamed with Cas9 or GFP
across papers and patents, especially during the early stages of Cas9 and GFP. We found
that, in their infancy to peaks, Cas9 and GFP respectively teamed with 6.4–13.1% and 6.7–
23.4% of medical science genes that debuted in papers or patents during similar periods
(figure 3*I* and Methods). Interestingly, if we limit our focus to pharmaceutical genes, they fall
short of this gross medical science result, with the particularly lower proportion for Cas9
(figure 3*I*; $P \leq 2 \times 10^{-4}$ and Methods). Of note, pharmaceutical research within medical
science is known to prioritize marketable outcomes. Another observation from our data is
that both metabolic engineering and (its parent) applied biotechnology rank among the
lowest in the proportions of the genes teaming early with Cas9 or GFP (figure 3*I*, and
electronic supplementary material, figure S7 for all the gene categories; $P \leq 10^{-4}$ and
Methods). These patterns do not largely depend on specific periods of the early teaming
(electronic supplementary material, figure S7). The patterns of pharmaceutical, metabolic
engineering, and applied biotechnology fields indicate that the highly practical focus of these
fields may be in fact limiting the test and adoption of newcomer technologies. There exist
further field-specific issues, such as the emphasis on product price competitiveness in the
cases of metabolic engineering and applied biotechnology, requiring the mindset of rational
managers rather than risk-taking innovators. Such commercial pressure restraining the
innovative potentials appears to be consistent with the previous, long-alarming trend in figure
2*C*, 3*C* and electronic supplementary material, figure S4*D,E*, S6*D,E*, which exhibited severe
decline of both newly researched genes and novel gene combinations in metabolic
engineering and applied biotechnology. In a later part of this study, we will further elaborate
on the downside of commercially-controlled research activities.
304
305   **Short-lived industry interests and long-lived scientific research**
306
Recall the very initial finding in this study that patent-filed technology inventions are typically
industry-driven, while paper-published scientific research is not. Because industry R&D is
closely aligned with market needs and demands, commercial pressure would influence
patent trajectories more than paper trajectories. Our subsequent results do remain valid
when considering only the company-assigned patents among all the patents.
Figure 4*A* shows the paper and patent time-series of COX-2, which has been studied for
anti-inflammatory treatment and pain relief [44–48]. Notably, its annual patent applications
started to decline in 2004 while the paper publications still persisted (figure 4*A*). This patent
falloff, in fact, coincided with the withdrawal of a COX-2 inhibitor drug for arthritis patients

(Vioxx) from the market, due to the risk of heart attack and stroke [44]. Phosphodiesterase 5

(PDE5) from our data, well known for its inhibitor sildenafil to treat erectile dysfunction

(Viagra), is another gene with a patent falloff but persistent paper publications. The use of

sildenafil was approved for pulmonary arterial hypertension in 2005, following its previous

approval for erectile dysfunction [49, 50]. Upon sildenafil's dominance in the market, patent

applications with PDE5 started to fall in 2007. Therefore, the cases of COX-2 and PDE5

presumably exemplify the industry's R&D depression due to risk management and

anticipated lower returns in a saturated market, respectively [10, 12, 13].

324
325

326
327

Figure 4. Paper–patent falloff relationship and research continuation. *A*, Annual paper (blue) or

patent (red) counts of COX-2. ***B***, The proportion of genes whose paper falloffs precede the patent

falloffs (blue) and the proportion of genes whose patent falloffs precede the paper falloffs (red). Here,

we considered genes that debuted in papers/patents in 1990–2000 (left) and 2001–2011 (right),

separately. For a fair comparison between paper and patent falloffs, we set the endpoint of the paper

time-series to the same year (2020) as the patent time-series. ***C***, On the left side, the proportion of

genes with clinical trial papers before ("Lively") or after/at ("Low") their patent falloffs, among the

clinical-trial genes that debuted in papers/patents in 2002–2011. These debut years were chosen

regarding the first year of the mandated trial posting to ClinicalTrials.gov (Methods). On the right side,

the probability of industry or other sponsorship for a clinical trial with each gene in a given case

("Lively" or "Low") from the left side. ***D***, The proportion of genes with papers before ("Lively") or

after/at ("Low") their patent falloffs, if these papers received ≥3 patent citations each within the first

three years of publications. We considered the genes that debuted in papers/patents in 1990–2000

(left) and 2001–2011 (right), separately. Alternative criteria for high patent citations did not

qualitatively change the observed result here.

From now on, we define the paper or patent falloff as a decrease in its number by at least a

quarter of the peak value, without a later full rebound (Methods). Alternations of this

definition did not change our main results. Remarkably, the preceding patent falloff above is

not limited to COX-2 or PDE5 but characterizes the general course of genetic research. For

two-thirds of genes that debuted in papers or patents during similar periods, their patent

falloffs preceded the paper falloffs (figure *4B*; $P < 10^{-4}$ and Methods). For example, genes

that debuted in 1990–2000 show their patent falloffs 6.5 ± 4.9 years (avg. ± s.d.) ahead of

the paper falloffs, and a quarter of these genes even show ≥10-years earlier patent falloffs.

Our further analysis excludes the possibility that the policy requirement of non-overlap of a

patent content with prior art might be responsible for the rapider patent falloff (electronic

supplementary material, figure S8 and Methods). We hence conclude that commercial

pressure tends to disrupt the industry R&D on a given gene first, despite the ongoing

scientific interest in that gene. One natural consequence of this relatively early patent falloff

is that the total patent volume of each gene becomes capped by the total paper volume with

some scaling (electronic supplementary material, figure S9 and Methods).

During the period of continuous scientific research without much of the industry's drive,

the pursuit of breakthroughs and impactful works did not stop. Figure *4C* shows that even

~60% of genes with clinical trials underwent these trials during the times of their limited

patent levels (Methods). Besides, a trial with each gene was usually not sponsored by the

industry, with a ~70% probability whether it was the limited patent case or not (figure *4C*; $P \leq$

0.05 and Methods). This fact reminds us of the previous report that 70% of COVID-19

clinical trials were driven by public research institutions [9]. To evaluate the technological

impact of scientific research, we measured it by the patent citations of each paper within a

defined period [20–22, 28, 30, 41]. We found that a substantial body of the genes (37.0–

50.7%) with the highly impactful papers did appear in those papers during the times of the

limited patent levels (figure *4D* and Methods).

These general tendencies are repeated in the cases of COX-2 and PDE5. After the

patent falloff, COX-2 continued to be studied for its function in various research papers,

along with many clinical trials [45–48, 51]. They include the studies of COX-2, EGFR, and

mTOR relations in cancer, later followed by a clinical trial [45, 46], the relationship between

ω−3 fatty acids and COX-2, with many patent citations [47], and the relief of

hyperinflammatory responses in COVID-19 [48]. In the case of PDE5 after the patent falloff,

its inhibition continued to be tested in paper-published research and clinical trials from

various aspects, such as β-catenin-involving anti-cancer mechanism and muscle ischemia

alleviation, both followed by patent citations [52–54], cardiovascular risk reduction [55], and

ventilation improvement for single-ventricle patients [56]. In other words, the industry's

interest in these genes turned cold first, but the innovative scientific research persevered.

380
381   **Commercial pressure can even disrupt long-term innovative opportunities**
382

Our results indicate that genes with the fading industry's interest still retain the opportunities

for innovative approach and possible breakthrough explorations. To further examine this

issue, we preselected eight workhorse genes in biotechnology innovation, based on their

shares in novel gene combinations explained above: Cas9 for genome editing [43], PD-L1

for immunomodulation and immunotherapy [57], mTOR and TP53 mainly for cancer

research [58, 59], GFP and luciferase for live cell imaging [60, 61], β-catenin mainly for

regenerative medicine and cancer research [62, 63], and β-galactosidase for DNA cloning

[42] (Methods). We will henceforth refer to these genes as *select genes*. Interestingly, the

aforementioned COX-2 teamed with two select genes—PD-L1 and Cas9—for the first time

after its patent falloff, although the patent applications reduced to one-third and one-fourth of

the peak, respectively (figure 5*A*). In these cases, PD-L1 and Cas9 had not yet reached their

own peaks, but they respectively served for the investigations of the suppression of

transplant rejection and the mechanism of gastric cancer [64, 65]. Likewise, the

aforementioned PDE5 newly teamed with three early-stage select genes—GFP, TP53, and

mTOR—when its patent applications reduced to one-third to one-fourth of the peak. GFP,

TP53, and mTOR here served for the monitoring of cyclic guanosine monophosphate

(cGMP) levels inside living cells and the investigations of kidney injury protection and liver

cancer treatment, respectively [66–68].

Indeed, we found that a substantial portion of genes teamed with new select genes after

their patent falloffs, although these select genes were still in the early stages of work. Of

genes that teamed with select genes in their infancy to peaks, the majority (54.5%) initially

teamed with at least one of these select genes after the patent falloffs (figure 5*B* and

Methods). About a half of these genes, including COX-2 and PDE5, initially teamed with two

or even more of those select genes after the patent falloffs (figure 5*B* and Methods). In

addition, for one-third to a half of the genes that teamed with Cas9, PD-L1, or mTOR in its

infancy to peak, these teams were initially formed after the patent falloffs, and this fraction

had already been appreciable during an earlier period of the teaming (figure 5*B,* electronic

supplementary material, figure S10, and Methods).

These results suggest that the loss of the industry's active interest does not necessarily

close the exploratory opportunities of genetic research. Further analysis reveals that

sustained scientific research is a key to bring this possibility to life, despite the fading

industry's interest. Specifically, the teaming of each gene with a new, early-stage select

gene after the patent falloff predominantly occurred (≥72.9%) when the active paper

publications still continued (figure 5*C*, electronic supplementary material, figure S11, and

Methods; $P < 10^{-4}$).

418
419

420
421
Figure 5. Teaming with new select genes in their early stages. *A*, Annual paper (blue) or patent

(red) counts of COX-2 (dashed line for a period before its patent falloff). An arrow indicates the initial

time of the teaming with each select gene if this teaming comes after or coincides with the patent

falloff. ***B*,** The proportion of genes teaming with ≥1, ≥2, or ≥3 new select genes after (or at) the patent

falloffs, and the proportion of genes initially teaming with a given select gene—Cas9, PD-L1, mTOR,

or GFP—after (or at) their patent falloffs (Methods). We only considered the teaming with the select

genes until the peak times of their annual paper and patent counts. ***C*,** The proportion of genes initially

teaming with a given select gene before ("Lively") or after/at ("Low") their paper falloffs, among the

genes with this teaming after/at the patent falloffs (Methods). We only considered the teaming with the

select gene until the peak time of its annual paper and patent counts. ***D*,** The estimated proportion of

genes that would lose the teaming with a given select gene, among the genes that initially teamed

with this select gene amid their limited patent applications ("Low") or among all the genes that teamed

with this select gene ("All") (Methods). We only considered the teaming with the select gene until the

peak time of its annual paper and patent counts. In ***B*–*D*,** we considered genes that debuted in

papers/patents in 1990–2000. Genes with post-2000 debuts were not considered due to the

fundamentally low overlap of their research periods with the early stages of the select genes.

438
439

These findings raise a fundamental question: if commercial pressure propagates from

industry to scientific activities even outside the industry with financial or other impacts [12,

69], to what extent would this pressure disrupt the exploratory opportunities of genetic

research? One instance of such propagation is the blow to biofuel research at leading

universities in the mid-2010s due to the oil price drop and subsequent investment cuts [70–

72]. To answer our question, we conducted a "thought experiment" that assumed the

hindering of scientific research on a given gene due to a lack of the industry's active interest:

specifically, we altered the temporal trend of the paper counts to match that of the patent

counts, which had come to a low level (Methods). The lack of the industry's visible interest in

a particular research area may not only affect its own investment but also possibly diminish

the public and governmental support for that research [73]. Our thought experiment gives

rise to a remarkable estimation that the vast majority (≥69.7%) of the genes, which initially

teamed with a given early select gene amid the limited industry's interest and patent

applications, would now lose this teaming (figure 5*D*, electronic supplementary material,

figure S12, and Methods). Among all the genes whether affected or not by our experimental

scenario, still 42.2–74.4% of the genes that once teamed with a given early select gene

would lose this teaming, with the exception of PD-L1 although it would also lose one-fourth

of the teams (figure 5*D,* electronic supplementary material, figure S12, and Methods). In the

example of COX-2 and PDE5 whose annual patents fell in 2005 and 2007, the hindering of

their scientific research would drastically reduce the chances of the teaming with each early

select gene by ≥83.5% at most (Methods). Hence, these estimations suggest the cost of

over-reliance on industry-dependent research, which would miss the long-term exploratory

and innovative opportunities once the industry's drive becomes limited.

463
464 **Discussion**
465
Through the macroscopic overview of decades-long genetic research courses, our work

challenges the existing perception that commercial motives promote cutting-edge innovation.

In fact, excessive commercial pressure can distort research trajectories and stifle the full

exploration of the innovation landscape in genetic research. Consistently, the fields of highly

practical or commercial focus, such as metabolic engineering, are less likely to embrace

innovative approaches and exhibit lower research vitality (figure 3*C*,*I* and electronic

supplementary material, figure S6, S7). Our analysis also shows that steady scientific

activities create the long-term opportunities for exploration and innovation, while industry's

R&D efforts are typically short-lived (figure 4*B*–*D*, 5*C* and electronic supplementary material,

figure S11). Of note, prioritizing early returns and risk aversion has been recognized as

common features of industry [6, 9–13]. We estimate that up to 42.2–74.4% of the exploratory

opportunities with notable gene combinations would be lost if the scientific research

becomes hindered due to the industry's limited interest (figure 5*D* and electronic

supplementary material, figure S12). In other words, the propagation of the commercial

pressure from the industry to the external scientific research can endanger long-term

innovation efforts, suggesting the cost of over-reliance on industry-driven research.

Regarding such propagation, we envisage that the lack of the industry's active interest may

not only suppress its own investment but also diminish the societal support for that research

[73]. Although our present data analysis takes various controls and null models to minimize

the possibility of alternative data interpretations (Methods), a future work beyond the current

scope should identify more specific causal relationships and mechanisms. This identification

may benefit from the use of highly refined data [74–76] and is a prerequisite for setting

relevant policies. Consideration of hidden biases in patent data due to commercial interests

is also warranted [23].

Our work points to the downside of commercial pressure (or its analogous effects at

large) on research activities, which can narrow the spectrum of approaches by restricting an

intellectual scope and may ultimately harm the vitality of innovation in an entire field itself

(e.g., figure 2*C*, 3*C*). Regarding the issue of the industry that is supposed to spur cutting-

edge innovation but can act contrary through the commercial pressure, a proper innovation

policy might aim to preserve the entrepreneurship of innovators. One way could be

governmental partnership to reduce innovation risks. For example, the government may offer

the support for the commercialization of new technologies by providing firms with

opportunities to test, apply, and publicize these technologies [77, 78]. Furthermore, the

government may help establish the communication channels with consumers to identify the

social needs for new technologies and thereby lower the uncertainty about market response

[79, 80]. These policies would encourage the industry to remain risk-taking innovators rather

than risk-averse rational managers, and thus contribute positively to the innovation

ecosystem.

To summarize, despite the perceived positive role of commercial pressure in technology

progress, it risks undermining long-term innovative potential. Our findings call for a balanced

approach with stable support for exploratory research [1, 5, 73]. This balance will be

essential to unlock the full potential of biotechnology and to address complex challenges

such as climate change, pandemics, major technological shifts, and social inequality [6, 81–

84], for which the need for innovative and creative solutions has never been greater.

Methods

Listing of gene names

Each entry of the UniProtKB/Swiss-Prot database presents a curated, non-redundant protein sequence mainly at the species level [27, 85]. For example, IL-6 protein has an entry for each single species such as human (*Homo sapiens*), mouse (*Mus musculus*), rat (*Rattus norvegicus*), or pig (*Sus scrofa*). Across all entries, we collected gene and protein names and their synonyms from the attributes in electronic supplementary material, table S2 (UniProtKB/Swiss-Prot XML file, Release 2022_04). These names and synonyms were later used for their search in paper and patent texts. For simplicity, we here refer to a set of homologous genes and their proteins as a gene, and refer to their names and synonyms as synonyms unless specified. The comprehensive synonym set of each gene was identified by the grouping of the entries with highly shared synonyms, as these entries often corresponded to homologous genes. Specifically, we excluded improper synonyms such as the names of cleaved chains or subdomains of main peptide bodies, and constructed the graph where each entry is a node and two nodes are linked if their entries share a substantial portion of the synonyms to satisfy $|A \cap B|/\min(|A|, |B|) \geq 0.75$ (A and B are the synonym sets of these entries). By applying the generalized Louvain method to this graph (v0.3; https://sourceforge.net/projects/louvain) [86], we detected densely-knitted subgraphs, i.e., network communities. For synonyms still shared by multiple network communities, each synonym was assigned only to the community with the largest share of the entries for that synonym. The whole synonyms in each community were then considered the gene's tentative synonym set. After the test run of synonym matches with paper and patent texts (described below), we identified some homonyms irrelevant to genes. To reduce these homonyms, we removed non-biological phrases in common use based on their frequent occurrences in published books in society, available from *Google Books Ngram Corpus* (version 3, February 2020) [87]: the phrases with higher counts than an empirically-chosen threshold (1.8×10^7 counts in the books published by 2020) were automatically removed from our gene synonyms. The remaining synonyms underwent extensive manual inspection and curation, and we then finalized our gene synonym sets.

Collection of paper and patent data

To survey research activities in genetic subjects, we targeted the papers in the journals of biology, applied biology, medicine, and other related disciplines. The categories of those disciplines and the associated journal lists were obtained from the Web of Science (WoS) Journal Citation Reports (JCR) (https://jcr.clarivate.com). We manually curated these journal

lists for better association with the journal categories and removed the journals irrelevant to

contemporary genetic research. The resulting journal lists are provided in electronic

supplementary material, data S1 with our expert-curated journal categories JG0, JG1, ⋯,

and JG8: JG1 includes biomedical journals, along with its subsets JG2 and JG3 for

veterinary medicine and pharmaceutics, respectively. JG4 includes journals in applied

biotechnology, along with its subset JG5 for metabolic engineering. JG6 and JG7 include

plant science and general microbiology journals, respectively (we here use the term "general

microbiology" to distinguish it from applied microbiology covered by JG4). JG0 includes all

the journals in JG1 to JG7, with additional journals for animal, algal, and aquatic research.

JG8 includes broad-scope journals and other journals not fit to JG0.

At the same time, we downloaded the annual releases of the baseline data files from the

PubMed FTP server (April 22, 2022; https://ftp.ncbi.nlm.nih.gov/pubmed/baseline) [88],

which contained the titles and abstracts of the articles serviced by PubMed. Other than the

titles and abstracts, we did not pursue the full text analysis because all the genes in the full

texts were not necessarily relevant to the very subjects of the articles. We considered only

the articles in the above JG0 and JG8 journals and published in 1990–2021 because the

Human Genome Project was launched in 1990 and our gene synonyms derived from the

UniProtKB/Swiss-Prot data dated May 2022 as noted above.

To survey inventive activities in genetic subjects, we collected the USPTO, CNIPA, and

EPO patent application data (US, Chinese, and European, respectively) in the PATSTAT

Global database (version 5.22, 2023 Autumn Edition) provided by the EPO. For each patent

family or any set of patents with duplicated titles and abstracts, we selected only the earliest-

dated patent to avoid data redundancy. Among the Cooperative Patent Classification (CPC)

codes jointly managed by the EPO and the USPTO (dated May 2, 2022) and the

International Patent Classification (IPC) codes managed by the World Intellectual Property

Organization (WIPO) (dated October 1, 2023), we selected possibly gene-related codes

based on expert opinions (electronic supplementary material, data S2) and considered the

patents with these codes. There is a known delay between patent application and

publication, and we did observe a roughly two-year delay with a spurious decrease in recent

patent application records. Therefore, taking this delay and our data access in Fall 2023 into

consideration, the patents applied by the end of 2020 were included in our further analysis.

Throughout this study, we considered only the patent titles and abstracts because the genes

there mostly overlapped with those in the claims and the rest included genes not necessarily

relevant to the very subjects of the patents.

Search of genes in papers and patents

To find the synonyms of each gene in the paper or patent titles and abstracts, we used the string-matching algorithm that directly compares the phrases in the target literature to the strings from the gene synonyms, while incorporating the indexed tree and tokenizing scheme for efficient computation [89]. Several rules were added to the search algorithms to consider common variations in the synonyms, such as the use of Greek letters, plural forms, and different letter cases. For each gene, papers and patents were counted when matching at least one of its synonyms. However, papers and patents with too many genes (>10 genes each) were excluded from our analysis, regarding their ambiguous research targets among those genes. The precision and recall of the gene identification were 0.90 and 0.71, respectively, according to the comparison with manually identified genes from 100 randomly selected papers (electronic supplementary material, data S4). Although the existing resource, PubTator3 also provides a list of genes in each article [90], the precision and recall (0.96 and 0.79) were only marginally better than our result above (electronic supplementary material, data S4). Critically, PubTator3 serves only PubMed papers, and therefore we excluded its use for consistency of our paper and patent analyses.

Throughout this study, we performed additional analyses that involve only the genes of the main focus in papers or patents. We identified these genes with the following heuristic rules: in the case of a paper, we considered a gene as of its main focus if this gene appears in the title or appears ≥3 times in the abstract. In the case of a patent, we considered a gene as of its main focus if this gene appears in the title or appears ≥2 times in the abstract, or if this gene is the only gene in the title and abstract. We validated these rules by the manual interpretation of 100 randomly chosen papers and patents. This manual inspection showed the precision of 0.86 and the recall of 0.90 for papers and the precision of 0.88 and the recall of 0.85 for patents (electronic supplementary material, data S5).

Paper funding source and patent assignee analyses

We manually identified the funding sources of 300 different papers in genetic research (electronic supplementary material, data S6), selected evenly as will be explained below. If the funding sources of a given paper included at least one company, the paper was considered company-funded. In the case of other funding sources from this paper set, we confirmed their public nature by reviewing the specific funding sources. For the selection of these 300 papers, we first randomly selected 50 different genes each (total 200), whether they debuted in papers/patents in 1990–2000 or 2001–2011, and whether their patent time-series include falloff years or are fully spanned by states II and III, which will be defined later.

These four cases were considered due to their high relevance to many analyses in this
study. In the cases of the patent falloffs, one paper before the patent falloff and the other
paper after (or at) this falloff were randomly selected for each gene; otherwise, any single
paper was randomly selected for each gene. The probability of a paper being company-
funded for each gene was obtained by the weighted average of the proportion of company-
funded papers over all those six cases, reflecting two from each patent-falloff case (one
before the falloff and the other afterward). In this calculation, we assigned a weight to each
case using the actual proportions of genes and their papers from the entire dataset.
For patent assignees of unambiguous characters, we considered "COMPANY", "GOV",
"NON-PROFIT", "UNIVERSITY", and "HOSPITAL" among the patent applicant sectors from
the PATSTAT database. For the genes with patents assigned to these sectors, we
calculated the average proportion of patents, each with at least one company assignee. This
quantity represents the probability of a patent being company-assigned for each gene.

632
633   **Thematic categorization of genes**
634
The common categorization of genes based on molecular or biochemical contexts rarely
allowed the best interpretation of our paper or patent counts with the genes. Therefore, we
categorized the genes according to their frequent research themes. Mainly based on the
journal categories and CPC/IPC codes of the papers and patents with each gene, the gene
was assigned to one of the following thematic categories (electronic supplementary material,
data S3). The methods described here were heuristically developed by the examination of
the categorization results.
"Medical science": for each gene, we calculated the ratio of its papers in the above JG1
journals to those in the JG0 journals (electronic supplementary material, data S1). Genes
with these ratios greater than or equal to an empirically-chosen threshold (0.7) were
assigned to category "medical science". In addition, for each gene, we calculated the fraction
of its patents with CPC or IPC codes "A61K38", "A61K31", "A61K39", "A61K48", and their
child codes (electronic supplementary material, data S2). Genes with these fractions greater
than or equal to an empirically-chosen threshold (0.5) were also assigned to the category
"medical science".
"Genetic tools": the genes utilized as the tools for a range of molecular biology and
genetic experiments were assigned to category "genetic tools" based on manual literature
search and expert opinions, in the absence of their specific journals and patent classification
codes. This category itself was not used for our analysis, but for excluding its genes from
other categories below.

"Applied biotechnology": this category was intended to include the genes mainly in non-
medical applied biotechnology. For each gene, we calculated the fraction of its patents with
CPC or IPC code "A61K" and its child codes (electronic supplementary material, data S2).
For each of the genes with these fractions lower than an empirically-chosen threshold (0.5),
we counted its papers in the JG4 journals (electronic supplementary material, data S1).
Each gene with this count ≥ 40% of the total number of its papers in the JG0 journals was
assigned to category "applied biotechnology". To avoid the overlap with the other categories,
the genes in the previous categories "medical science" and "genetic tools" were here
excluded from the outset.
"Plant science" and *"general microbiology"*: as noted above, we use the term "general
microbiology" to distinguish it from applied microbiology covered by "applied biotechnology".
For each gene, we calculated the ratio of its papers in the JG6 (JG7) journals to those in the
JG0 journals, and the genes with these ratios ≥ 0.5 were assigned to category "plant
science" ("general microbiology"). To avoid the overlap with the other categories, the genes
in "medical science", "genetic tools", and "applied biotechnology" were here excluded from
the outset. The overlap between "plant science" and "general microbiology" was avoided by
manual recategorization of their shared genes.
A part of genes in "medical science" were subcategorized as follows: for each gene in
"medical science", we calculated the ratio of its papers in the JG3 journals to those in the
JG1 journals. The genes with these ratios ≥ 0.25 were assigned to subcategory
"pharmaceutical". Another subcategory *"immune"*, given its definite biological function and
existing database, was assigned the genes of Gene Ontology (GO) term "immune response"
(GO:0006955) and its child terms, but "hemocyte proliferation" (GO:0035172) or its child
terms [91]. Additionally, the genes associated with Anatomical Therapeutic Chemical (ATC)
codes "immunostimulants" (L03) and "immunosuppressants" (L04) in the DrugBank
database [92] were manually reviewed and thereby assigned to the subcategory "immune".
Note that the above subcategory "pharmaceutical" allows the overlap with this subcategory
"immune".
Each gene in "applied biotechnology" was assigned to subcategory *"metabolic*
engineering", when the ratio of its papers in the JG5 journals to those in the JG4 journals
was ≥ 0.4.
The genes in the above categories and subcategories, particularly those near their
thresholds, underwent further manual reviews to adjust and finalize the categorization.
Naturally, there remain genes without any category assignment.

Annual count of newly researched genes

We counted new genes in papers and patents each year in the following ways: (*i*) based on the debut year of each gene in papers or patents, whichever earlier for that gene (figure 2*A*); (*ii*) based on the debut year of each gene in papers (electronic supplementary material, figure S2*A*); and (*iii*) based on the debut year of each gene in US, Chinese, or European patents, whichever the earliest for that gene (electronic supplementary material, figure S2*B*– *D*). Because of the low availability of the patent data after 2020 as noted above, we only considered the paper data for 2021 in the case of (*i*). Nevertheless, because 2021 is just one year and the paper volume far exceeds the patent volume, the result is unlikely to change much even when the 2021 patent data are considered.

To count the genes related to *Arabidopsis thaliana* or rice among new genes in the category of plant science, we considered the new genes whose paper or patent titles/abstracts matched one of words "Arabidopsis", "A. thaliana", and "Arabidopsis thaliana" for *A. thaliana*, and "rice", "Oryza", "O. sativa", and "Oryza sativa" for rice, in a case- insensitive way.

Mathematical modeling of new gene research over time

In our mathematical model, we consider a total of *L* genes and their *M* non-redundant amino acid (AA) sequences across species (i.e., *M* gene copies across species). These sequences can correspond to the aforementioned UniProtKB/Swiss-Prot entries and are assumed to have the potential for prospective research. We label these *M* sequences by $1, 2, \cdots, M$ in the temporal order of their debuts in research in papers and patents, and consider all the possible configurations of this labeling with approximately equal *a priori* probabilities. If *X* and *Y* are the number of the member sequences of a given gene and the earliest label of these member sequences, respectively, then the probability of $Y \leq y$ for the gene with $X = x$ is given by

$$\Pr(Y \leq y | X = x) = \begin{cases} 1 - \binom{M-y}{x}\binom{M}{x}^{-1} & \text{if } 1 \leq y \leq M - x \\ 1 & \text{if } y > M - x \end{cases}, \qquad (1)$$

where the binomial coefficient is given by $\binom{a}{b} = \frac{a!}{b!(a-b)!}$. If *f*(*x*) is the number of genes with *x* member sequences each, it satisfies the following relations:

$$\sum_{x=1}^{M} f(x) = L \; ; \quad \sum_{x=1}^{M} xf(x) = M \, . \tag{2}$$

719   Let $b(y)$ be the expected number of genes that debuted in research by the time of the y^{th}-

720   labeled sequence's debut in research. $b(y)$ is then expressed as

$$b(y) = \sum_{x=1}^{M} f(x)\,\mathrm{Pr}(Y \le y | X = x) = \sum_{x=1}^{M} f(x) - \sum_{x=1}^{M-y} f(x) \binom{M-y}{x} \binom{M}{x}^{-1} . \tag{3}$$

721   $b(y)$ in Eq. (3) can be further expressed as the function of time t, i.e., $b(y(t))$ if $y(t)$ is the total

722   number of the researched sequences by that time t. Note that the reporting of a given AA

723   sequence to the scientific community does not necessarily coincide with the research of this

724   sequence. After the reporting of the sequence, the event of its first research may be

725   modeled as a Poisson process with the following $y(t)$:

$$y(t) = \sum_{t''=t_0}^{t} \sum_{t'=t_0}^{t''} \beta N(t') e^{-(t''-t')/\tau} \left(1 - e^{-1/\tau}\right) . \tag{4}$$

Here, $N(t')$ is the number of the newly reported, non-redundant sequences in year t'

obtainable from the UniProtKB/Swiss-Prot entries, β is the fraction of the sequences for

prospective research ($0 \le \beta \le 1$), t'' denotes the debut year of each sequence in research,

τ is the characteristic latent period between the sequence's first report and research, t_0 is the

earliest of year t' for the available $N(t')$ data (i.e., 1939 in the case of the

UniProtKB/Swiss-Prot data), and $\left(1 - e^{-1/\tau}\right)$ serves as the normalization factor.

In practice, the reporting years of the sequences in the UniProtKB/Swiss-Prot entries

often indicate their first sequencing years, which may not be the same as their actual earliest

reporting years, particularly in the case of the early-reported genes before the wide

popularization of sequencing technologies. In this regard, we examined the first reporting

years in the UniProtKB/Swiss-Prot entries and found some cases before 1997 where each

gene's reporting was spuriously preceded by its debut in research. Therefore, we considered

only $t \ge 1997$ (year) in our modeling.

If there exists x' to satisfy $\sum_{x=x'}^{M} f(x) \ll \sum_{x=1}^{M} f(x)$ and $x' \ll M - y$, $b(y)$ in Eq. (3) is

simplified as follows:

$$b(y) \approx \sum_{x=1}^{M} f(x) - \sum_{x=1}^{M} f(x)\left(1 - \frac{y}{M}\right)^x. \tag{5}$$

741 To avoid the overfitting of our model to empirical data, we lump $f(x)$'s and reduce their
742 number, as follows:

$$f(x)\begin{cases} \geq 0 & \text{if } x = 1 \text{ or } x = k \ (k > 1) \\ = 0 & \text{otherwise} \end{cases}. \tag{6}$$

As noted previously, the non-redundant sequence in each UniProtKB/Swiss-Prot entry is
mainly valid at the species level, and therefore genes with $x = 1$ in Eq. (6) correspond to
taxonomically-restricted genes (TRGs) or orphans that lack recognizable homologs in other
species [37, 38]. Here, we call the other genes taxonomically-dispersed genes (TDGs), and
they are lumped to $x = k$ in Eq. (6) where k denotes the effective number of the species for a
TDG. Eq. (5) is now converted to

$$\begin{aligned} b(y) &\approx \sum_{x=1}^{M} f(x) - f(1)\left(1 - \frac{y}{M}\right)^1 - f(k)\left(1 - \frac{y}{M}\right)^k \\ &\approx \sum_{x=1}^{M} f(x) - f(1)\left(1 - \frac{y}{M}\right)^1 - f(k)e^{-ky/M} \\ &= \frac{f(1)}{M}y + f(k)\left(1 - e^{-ky/M}\right) = sy + i\left[1 - e^{-(1-s)y/i}\right], \end{aligned} \tag{7}$$

749 where $s \equiv f(1)/M$, $i \equiv f(k)$, and we used this relation:

$$\frac{k}{M} = \frac{1}{f(k)}\frac{kf(k)}{M} = \frac{1-s}{i}. \tag{8}$$

750 Because the values of $b(y(t)) - b(y(t - 1 \text{ year}))$ are available from figure 2A, their summation
751 over time leads to $b(y(t))$ with some constant shift. If we reinterpret $b(y)$ as this summation of
752 the empirical values, then it is rewritten as

$$b(y) \approx sy + i\left[1 - e^{-(1-s)y/i}\right] + c, \tag{9}$$

where c is a constant.
Next, we seek the parameters in Eq. (9). For each pair of τ and β in Eq. (4) with $(\tau, \beta) \in$
$\{0, 1, 2, \cdots, 19\} \otimes \{0.1, 0.2, 0.3, \cdots, 1.0\}$ (τ takes the unit of year), we randomly selected 10^6
sets of the initial parameters of (s, i, c) in the ranges of $(0,1) \otimes [[, b(t = 2021)] \otimes [[, b(t =$
1997)] ($b(t)$ denotes the above-mentioned empirical data of $b(y(t))$, t takes the unit of year,
and we consider only $t \geq 1997$ as previously discussed). This random initial parameter

selection was conducted by the PCG algorithm in NumPy v1.19.1. Plugging the empirical

values of $N(t')$ in Eq. (4) for $y(t)$, we minimized the following error functions by the

sequential least squares programming (SLSQP in SciPy v1.6.2):

$$\chi_1 \equiv \frac{\sum_{t=1997}^{2021}|\Delta b(t) - \Delta b_{\text{est}}(t)|}{\sum_{t=1997}^{2021}\Delta b(t)}; \tag{10}$$

$$\chi_2 \equiv \frac{\sum_{t=1997}^{2021}|b(t) - b_{\text{est}}(t)|}{\sum_{t=1997}^{2021}b(t)}, \tag{11}$$

where $\Delta b(t) \equiv b(t) - b(t - 1 \text{ year})$, $\Delta b_{\text{est}}(t) \equiv b_{\text{est}}(t) - b_{\text{est}}(t - 1 \text{ year})$, and $b_{\text{est}}(t)$ is the

right-hand side of Eq. (9) with $y(t)$ in Eq. (4). The summation in the error functions runs

from the year 1997 for the aforementioned reason. Given τ and β, the minimization of χ_1

gives rise to the best-fit *s* and *i*, and the subsequent minimization of χ_2 with these *s* and *i*

gives rise to the best-fit *c*. Figure 2*E* presents the heatmap of the resulting χ_1 and the

corresponding graphs of $b_{\text{est}}(t)$ and $\Delta b_{\text{est}}(t)$ as the functions of $y(t)$ and *t*, respectively.

The comparison of $b_{\text{est}}(t)$ and $b(t)$ shows their excellent agreement ($\chi_1, \chi_2 \leq 0.07$) over a

broad range of τ and β: P <10^{-4} for either the resulting χ_1 or χ_2 (one-tailed test) from the

null distribution of χ_1 or χ_2 with the direct use of 10^4 randomly selected sets of (*s*, *i*, *c*) for

each (τ, β) in the same parameter ranges as above (PCG algorithm in NumPy v1.19.1). We

stress that the observed match between $b_{\text{est}}(t)$ and $b(t)$ is not a mere result of parameter

overfitting but stands on the genuine agreement of the empirical data with our model output:

$b(t)$ itself initially curves and approaches a straight line as a function of $y(t)$ (figure 2*E*) as

exactly predicted by Eq. (9) in the model, and thereby always results in $\chi_2 < 0.01$ whenever

$\chi_1 \leq 0.07$ is satisfied.

Our model output involves the quantities about TRGs and TDGs mentioned above. First,

the fraction of TRG sequences among all the non-redundant sequences of research merit in

the model is *f*(1)/*M* = *s*. Across all the τ and β pairs resulting in $\chi_1, \chi_2 \leq 0.07$, we obtained

the probability distribution of *s* in electronic supplementary material, figure S5*A*. The

expected fraction of the TRGs among all the researched genes as of year *t* is *f*(1)·[*y*(*t*)/*M*]/

$b_{\text{est}}(t)$ = *sy*(*t*)/$b_{\text{est}}(t)$. In the same χ_1 and χ_2 conditions as above, we obtained the

probability distribution of the latest available *sy*(*t*)/$b_{\text{est}}(t)$ (i.e., *t* = 2021 (year)) as in

electronic supplementary material, figure S5*B*.

Because the expected number of TDGs researched as of year *t* is $\sim f(k)\{1 -$

$[1 - y(t)/M]^k\} \approx i[1 - e^{-(1-s)y(t)/i}]$, the expected fraction of the TDGs researched as of

year t_1 among the TDGs as of year t_2 ($t_1 < t_2$) is

$F_{\mathrm{TDG}}(t_1, t_2) \equiv \left[1 - e^{-(1-s)y(t_1)/i}\right]/\left[1 - e^{-(1-s)y(t_2)/i}\right]$. Because the expected number of TRGs

researched as of year t is $f(1) \cdot [y(t)/M] = sy(t)$, the expected fraction of the TRGs researched

as of year t_1 among the TRGs as of year t_2 ($t_1 < t_2$) is $F_{\mathrm{TRG}}(t_1, t_2) \equiv y(t_1)/y(t_2)$. In the same χ_1

and χ_2 conditions as above, we computed $F_{\mathrm{TDG}}(t_1, t_2)$ and $F_{\mathrm{TRG}}(t_1, t_2)$ when $t_1 < 2002$ (year)

and $t_2 = 2021$ (year), as presented in the main text for the discussion of the result in figure

2*A*.

794
795   **Identification of gene combinations**
796
To identify the combination of genes that are meaningfully linked in a given paper or patent,

we considered the pair, triplet, or entire combination (*n*-tuple with $n \geq 2$) of the genes that

include at least one gene of the main focus in that paper or patent (see above for the

identification of the genes of the main focus in a paper or patent). Because of the low

availability of the patent data after 2020 as noted above, we considered only the paper data

in the case of gene combinations in 2021. Nevertheless, because 2021 is just one year and

the paper volume far exceeds the patent volume, the result is unlikely to change much even

when the 2021 patent data are considered.

805   If any gene in a given pair belongs to a certain thematic gene category, we treated this

806   pair as affiliating with this thematic category (figure 3*B–D* and electronic supplementary

807   material, figure S6).

808   For each year or period, we defined the share of a given gene in new pairs as the

number of new pairs with this gene divided by the total number of new pairs in that year or

period (figure 3*E–H* and electronic supplementary material, table S1). Among the genes with

the highest annual shares, Cas9 and GFP are recognized for their utility across disparate

domains such as medical and plant sciences. Therefore, for each thematic gene category,

we computed the proportion of genes paired with Cas9 or GFP. In this calculation, we

focused on the pairings during the early stages of Cas9 and GFP, as these early stages

indicate the novelty and innovativeness of the pairings. Specifically, for Cas9 or GFP, we

considered the period up to the peak time of its annual paper and patent number.

Additionally, we even considered an earlier stage, the period up to the midpoint of the

cumulative paper and patent number at the peak time. As will be seen, the same definitions

of the early pairing can apply to other genes of interest beyond Cas9 and GFP. If the annual

paper and patent number of such a gene continued to grow until 2021, then 2021 is treated

as the peak time in these definitions. Back to the proportion of genes paired with Cas9 or

GFP, we considered genes that debuted in papers or patents during similar periods, in order

to control for variations from different periods of debut years: one group of genes with debut
years of 1990–2000 (figure 3*I*, left and electronic supplementary material, figure S7*A*,*C*) and
the other with 2001–2011 (figure 3*I*, right and electronic supplementary material, figure
S7*B*,*D*). More recent debut years (2012–) were not considered because the number of these
genes was relatively small and they may not yet have reached the levels of research stages
for the full analysis.

For the statistical significance test of the lower proportion of pharmaceutical genes paired
with Cas9 or GFP than that of the whole medical science genes (figure 3*I*), we generated
10^4 random subsets of the medical science genes and each subset included the same
number of genes as pharmaceutical genes. Using these subsets, we obtained the P value of
the observed proportion of pharmaceutical genes paired with Cas9 or GFP in its above-
mentioned early stage (one-tailed test). In addition, we performed the statistical significance
test of the relatively low proportion of metabolic engineering (applied biotechnology) genes
paired with Cas9 or GFP, compared to those in other categories (figure 3*I* and electronic
supplementary material, figure S7). Specifically, we randomly regrouped genes in major
thematic categories—medical science, plant science, metabolic engineering (applied
biotechnology), and general microbiology—while keeping the original sizes of these
categories. In this way, we generated 10^4 null configurations of the category membership
and obtained the P value of the observed proportion of metabolic engineering (applied
biotechnology) genes paired with Cas9 or GFP in its above-mentioned early stage and the
rank of this proportion among the categories (one-tailed test). All these statistical tests were
performed for the genes that debuted in papers or patents during similar periods, as
described above.

846
847 **Paper and patent falloff patterns**
848
If annual paper numbers of a given gene are decent enough (≥ 5 papers/year on average for
certain three consecutive years), we smoothened the paper time-series with a moving
window average (3-year window) to reduce its fluctuations. If the papers in the last window
do not exceed a certain threshold (75% of the peak), we considered the latest year of
meeting or crossing this threshold after the peak time as the paper falloff year. We defined
the patent falloff year in a similar way. Alternations of our methods, however, did not much
change our main results.

If annual papers of a given gene are scarce (without any three consecutive years of ≥ 5
papers/year on average), we averaged the paper numbers in year t–1, t, and t +1, and
termed this average X_t here. To each year t, we assigned one of the following states based
on the majority of X_{t-1}, X_t, and X_{t+1}: state I if the majority are ≥1 each; state II if the majority

are 0 each; and otherwise, state III. In the case that $t-1$ is the debut year of the gene in

papers or patents, or $t+1$ is the last time point, only two of X_{t-1}, X_t, and X_{t+1} are defined and

we assigned one of the following states to year t: state I if any of two is ≥1 and both are >0;

state II if any of two is 0 and both are <1; and otherwise, state III. In the case that t is the

debut year of the gene in papers or patents, or t is the last time point, only either X_{t-1} or X_{t+1}

is defined and we assigned one of the following states to year t: state I if the value is ≥1;

state II if the value is 0; and otherwise, state III. If the paper time-series includes state I

sometime before the last point and state II or III at the last point, we considered the year next

to the latest time of state I as the paper falloff year. We defined the patent falloff year in a

similar way. Alternations of our methods, however, did not much change our main results.

To investigate paper and patent falloff patterns, we obtained the proportion of genes

whose paper falloffs preceded the patent falloffs and vice versa (figure 4*B*). In this

calculation, we used the same period (–2020) for both paper and patent time-series for their

fair comparison and considered only the genes that satisfy the following conditions: (*i*) any of

the paper and patent time-series includes the falloff year defined above; and (*ii*) neither the

paper nor patent time-series is fully spanned by states II and III defined above. In addition,

we considered the genes that debuted in papers or patents during similar periods (1990–

2000 and 2001–2011, but not 2012–) for the same reasons explained earlier.

To examine whether the policy requirement of non-overlap of a patent content with prior

art is responsible for the observed tendency of the paper and patent falloffs (figure 4*B*), we

filtered out papers and patents with highly similar contents to their prior works and re-

analyzed the data (electronic supplementary material, figure S8). This test ensures a fair

paper and patent treatment, with the analyzed papers being as dissimilar to their prior works

as the patents are. Specifically, we used PaECTER, a citation-informed sentence

transformer [93] and obtained the cosine similarity of every pair of papers (patents) for each

gene, based on the paper (patent) titles and abstracts. If the total number of papers (patents)

exceeded 5,000 for a given gene, we only used 5,000 randomly selected papers (patents)

for the similarity calculation, which would otherwise become computationally impractical. For

the papers (patents) of each gene in a particular year, we filtered out the later papers

(patents) with the similarity at or above a certain threshold, and repeated this process across

all the considered years in chronological order. The similarity threshold here (0.97) was

determined by the manual inspection of the contents of paper or patent pairs with varying

threshold values. After the filtering, we obtained the proportion of genes whose paper falloffs

preceded the patent falloffs and vice versa. Regarding the control group without the filtering,

we used the aforementioned 5,000 papers (patents) for each gene if this gene occurred in

more than 5,000 papers (patents), and otherwise, all its papers (patents).

For the statistical significance test of the dominance of genes whose patent falloffs

preceded the paper falloffs (figure 4*B* and electronic supplementary material, figure S8), we

randomly swapped the paper and patent time-series of each gene and generated 10^4 null

configurations. Using these null configurations, we obtained the *P* value of the observed

proportion of genes whose patent falloffs preceded the paper falloffs (one-tailed test). This

statistical test was performed for the genes that debuted in papers or patents during similar

periods, as described above.

The total paper versus patent numbers of each gene

The practical upper limit of the patent number of a gene can be expressed as a function of

its paper number, $y = \alpha x^\gamma$ where *x* and *y* are the paper and patent numbers, respectively

(electronic supplementary material, figure S9). The parameters *γ* and *α* were initially found

by the fitting of the function with the least square method (polyfit in NumPy v1.19.1) and then

adjusted manually. To test the statistical significance of the observed fraction of the genes

below or on the upper limit, we randomly permuted the paper numbers of the genes (or

equivalently, their patent numbers) and obtained the probability of the fraction of the genes

(below or on the same above-mentioned upper limit) being greater than or equal to the

observed fraction. This probability serves as the *P* value of the observed fraction (one-tailed

test) and was computed with 10^4 trials of the randomization by the PCG algorithm in NumPy

v1.19.1.

Selection of genes that signify biotechnology innovation

We first calculated the share of each gene in new annual pairs of genes, as defined above.

We then identified the genes of the highest shares for ≥5 consecutive years or among the

top five share genes for ≥10 years in 1990–2021. They were a total of nine genes that signify

biotechnology innovation: Cas9 for genome editing [43], PD-L1 for immunomodulation and

immunotherapy [57], mTOR, phosphatidylinositol 3-kinase (PI3-K), and TP53 mainly for

cancer research [58, 59, 94], GFP and luciferase for live cell imaging [60, 61], β-catenin

mainly for regenerative medicine and cancer research [62, 63], and β-galactosidase for DNA

cloning [42]. We excluded PI3-K from further analysis due to its strong contextual overlap

with mTOR through the biochemical pathway sharing. The remaining eight genes are

referred to as *select genes* in this study.

Clinical trial analysis

We sought the papers with a given gene as the main focus in the above-mentioned way,

while each paper contains the titles/abstracts with "clinicaltrials.gov" (case-insensitive) and

"NCT" followed by an eight-digit number. Using this National Clinical Trials (NCT) number,

we identified the type of the sponsor for that clinical trial from ClinicalTrials.gov (June 6,

2024), managed by the U.S. National Library of Medicine as a registry with information on

global clinical trial activities. Among the sponsor types, we only considered "INDUSTRY",

"NIH", "OTHER_GOV", "FED", and "OTHER". In contrast to "INDUSTRY", we confirmed the

public nature of those other sponsor types by reviewing their specific sponsors.

Since September 2007, the Food and Drug Administration Amendments Act of 2007 has

mandated the posting of trials to ClinicalTrials.gov. According to our observation, it typically

took ~6 years between the debut of a gene in papers/patents and its first trial, suggesting

2002– as the appropriate debut years of genes for the analysis. In addition, the genes of

relatively recent debut years (2012–) were not considered because the number of these

genes was relatively small and they may not yet have reached the levels of research stages

for the full analysis. As a result, we focused on the genes of the debut years 2002–2011.

Among these genes in clinical trial papers as their main focus, we obtained the proportion

of the genes in those papers before the patent falloffs (figure 4C, left "Lively"), and the

proportion after (or at) the patent falloffs or with the patent time-series fully spanned by

states II and III defined above (figure 4C, left "Low"). To calculate the probability of industry

(or other) sponsorship for a clinical trial with each gene before its patent falloff, we obtained

the average proportion of clinical trials with industry (or other) sponsorship described above,

over all the genes in the clinical trial papers before the patent falloffs (figure 4C, right

"Lively"). Similarly, we calculated the probability of industry (or other) sponsorship for a

clinical trial with each gene after (or at) its patent falloff or of the patent time-series fully

spanned by states II and III defined above (figure 4C, right "Low").

For the statistical significance test of the high probability of non-industry sponsorship for

a clinical trial with each gene compared to industry sponsorship, we randomly but equally

reassigned industry or other sponsorship to each clinical trial and generated the 10^4 null

configurations. From these null configurations, we obtained the P value of the observed

probability of non-industry sponsorship in the case before the patent falloff, or in the case

after (or at) the patent falloff or of the patent time-series fully spanned by states II and III

(one-tailed test).

965
966 **Patent citation analysis**
967
We collected the data of patent citations of papers from SciSciNet [76]. For compatibility with

our paper data, the PubMed reference numbers (PMIDs) of papers in SciSciNet were

retrieved from OpenAlex (May 25, 2024) [75]. We focused on highly patent-cited works, the

papers that received ≥3 patent citations each within the first three years of publications.

However, alternative criteria of high patent citations did not qualitatively change our analysis
result. Among the genes of the main focus in the highly patent-cited works, we obtained the
proportion of genes in those works before the patent falloffs (figure 4*D*, "Lively"), and the
proportion after (or at) the patent falloffs or with the patent time-series fully spanned by
states II and III defined above (figure 4*D*, "Low"). To control for variations from different
debut years of genes, we considered the genes of similar debut periods, 1990–2000 and
2001–2011 separately. More recent debut years (2012–) were not considered for the same
reason as the previous analysis.
980
981 **Analysis of the teaming with new select genes**
982
We considered genes that had the patent falloff years defined above and were paired with
any select genes until the aforementioned peak times of these select genes. Among them,
we obtained the proportion paired with ≥1, ≥2, or ≥3 new select genes after (or at) the patent
falloffs, until the peak times of the select genes (figure 5*B*). In addition, for a given select
gene, we considered genes paired with this select gene until its peak time. Among them, we
obtained the proportion of genes initially paired with that early select gene before/after the
patent falloffs (figure 5*B* and electronic supplementary material, figure S10*A*). Similarly, we
obtained the proportion for the above-defined midpoint of the peak time, instead of the peak
time (electronic supplementary material, figure S10*B*). All these analyses were performed for
the genes that debuted in papers or patents during a similar period, 1990–2000. Genes with
post-2000 debuts were not considered due to the fundamentally low overlap of their active
research periods with the early stages of the select genes.
For a given select gene, we next focused on the genes that satisfy the following
conditions: (*i*) they were initially paired with this select gene until its peak time or the
midpoint of the peak time, after (or at) the patent falloffs; (*ii*) their paper time-series are not
fully spanned by states II and III defined above; (*iii*) the patent falloffs preceded or coincided
with the paper falloffs. Among these genes, we obtained the proportion of genes initially
paired with that early select gene before/after the paper falloffs (figure 5*C* and electronic
supplementary material, figure S11). In addition, we performed the statistical significance
test of the higher proportion of those genes before the paper falloffs than after (or at) the
paper falloffs, observed for most select genes. Specifically, we randomly reassigned the
initial pairing times of each select gene with equal *a priori* probabilities over the range from
the select gene's debut year or its paired gene's patent falloff year (whichever later) to the
latest point of at least one paper or patent of this paired gene. Across all the relevant select
genes and their paired genes, we generated 10^4 null configurations of the initial pairing times
and obtained the *P* value of the observed, joint proportions of genes initially paired with the

select genes before the paper falloffs. All the analyses were performed for the genes that

debuted in papers or patents during a similar period (1990–2000), as discussed earlier.

1011
1012   **Missing exploratory opportunities under commercial pressure**
1013
To examine the effect of commercial pressure on the exploratory opportunities of genetic

research, we conducted the "thought experiment" for each select gene, as follows.

Define set A as the genes that teamed with this select gene until its peak time and satisfy

the following condition (i) or (ii): (i) they were initially paired with that early select gene after

(or at) the patent falloffs and the patent falloffs preceded or coincided with the paper falloffs;

or (ii) the patent time-series are fully spanned by states II and III defined above but the paper

time-series are not. In addition, define set B as a subset of A, where genes were initially

paired with that early select gene before the paper falloffs. Also, define set C as a subset of

A, where genes were initially paired with that early select gene after (or at) the paper falloffs.

For the genes initially paired with a given early select gene with their limited patent

volumes (i.e., set A), we consider a hypothetical scenario that the papers undergo immediate

falloffs due to the limited patent volumes. This scenario allows us to estimate the maximum

extent to which the early teaming with this select gene would be lost if the research activities

become hindered by the commercial pressure from the industry's limited interest. The

estimated proportion of genes that would lose this early teaming is given by $(|A| - x_0)/|A|$

(figure 5D and electronic supplementary material, figure S12), where x_0 is the number of

genes that would still maintain the teaming despite the paper falloffs. Here, $x_0 \equiv P|B| + |C|$

with $P \equiv |C|/|A|$. P denotes the conditional probability that each gene in B would still

maintain the teaming. By definition, all genes in C would clearly maintain the teaming.

We further define set D as all genes that teamed with a given early select gene. We also

define set E as $E \equiv D - B$. Among all the genes in D whether affected or not by the above

hypothetical scenario, the estimated proportion of genes that would lose the early teaming

with the select gene is given by $(|D| - x_1)/|D|$ (figure 5D and electronic supplementary

material, figure S12), where $x_1 \equiv P|B| + |E|$ with P defined previously. This derivation is

similar to the above formulation.

Additionally, we conducted the same thought experiment, but for the teaming with each

select gene until the above-defined midpoint of its peak time. All these analyses were

performed for the genes that debuted in papers or patents during a similar period (1990–

2000), as discussed previously. In the example of COX-2 or PDE5, the estimated proportion

of the chances to lose the early teaming with a particular select gene is given by $1 - P$,

calculated for this select gene.

Acknowledgments

We thank Taekyun Kim, Hong Lok Lung, Seolmin Yang, Daehyun Kim, Liming Xiong, Yiji Xia, Adam G. Craig, and Anna Oi Wah Leung for useful discussions.

References

1. Bush V. Science the Endless Frontier: A Report to the President. Washington, DC: United States Office of Scientific Research and Development; 1945.

2. Stokes DE. Pasteur's Quadrant: Basic Science and Technological Innovation. Washington, DC: Brookings Institution; 1997.

3. Mokyr J. The Gifts of Athena: The Origin of the Knowledge Economy. Princeton, NJ: Princeton University; 2002.

4. Hiltzik MA. Dealers of Lightning: Xerox PARC and the Dawn of the Computer Age. New York, NY: HarperCollins; 1999.

5. Georgescu I. Bringing back the golden days of Bell Labs. Nat Rev Phys. 2022;4(2):76-8.

6. Ahmed N, Wahed M, Thompson NC. The growing influence of industry in AI research. Science. 2023;379(6635): 884-6.

7. Kumar A, Blum J, Le TT, Havelange N, Magini D, Yoon IK. The mRNA vaccine development landscape for infectious diseases. Nat Rev Drug Discov. 2022;21(5): 333-4.

8. Gibney E. Quantum gold rush: the private funding pouring into quantum start-ups. Nature. 2019;574: 22-4.

9. Agarwal R, Gaule P. What drives innovation? Lessons from COVID-19 R&D. J Health Econ. 2022;82: 102591.

10. Krieger J, Li D, Papanikolaou D. Missing novelty in drug development. Rev Financ Stud. 2022;35: 636-79.

11. Teece DJ. Profiting from technological innovation: Implications for integration, collaboration, licensing and public policy. Res Policy. 1986;15: 285-305.

12. Blumenthal D, Causino N, Campbell E, Louis KS. Relationships between academic institutions and industry in the life sciences—an industry survey. N Engl J Med. 1996;334(6): 368-73.

13. Pellegrino G. Barriers to innovation in young and mature firms. J Evol Econ. 2018;28: 181-206.

14. Azoulay P, Ding W, Stuart T. The impact of academic patenting on the rate, quality, and direction of (public) research output. J Ind Econ. 2009;57(4): 637-76.

15. Lee ED, Kempes CP, West GB. Idea engines: Unifying innovation & obsolescence from
markets & genetic evolution to science. Proc Natl Acad Sci U S A. 2024;121(6):
e2312468120.

16. McElheny VK. Drawing the Map of Life: Inside the Human Genome Project. New York,
NY: Basic Books; 2010.

17. Park M, Leahey E, Funk RJ. Papers and patents are becoming less disruptive over time.
Nature. 2023;613(7942): 138-44.

18. Bloom N, Jones CI, Van Reenen J, Webb M. Are ideas getting harder to find? Am Econ
Rev. 2020;110: 1104-44.

19. Gates AJ, Gysi DM, Kellis M, Barabási A-L. A wealth of discovery built on the Human
Genome Project - by the numbers. Nature. 2021;590(7845): 212-5.

20. Ahmadpoor M, Jones BF. The dual frontier: Patented inventions and prior scientific
advance. Science. 2017;357(6351): 583-7.

21. Poege F, Harhoff D, Gaessler F, Baruffaldi S. Science quality and the value of
inventions. Sci Adv. 2019;5(12): eaay7323.

22. Shi F, Evans J. Surprising combinations of research contents and contexts are related to
impact and emerge with scientific outsiders from distant disciplines. Nat Commun.
2023;14(1): 1641.

23. Kwon S. Underappreciated government research support in patents. Science. 2024;
385(6712): 936-8.

24. McGuire AL, Gabriel S, Tishkoff SA, Wonkam A, Chakravarti A, Furlong EEM, et al. The
road ahead in genetics and genomics. Nat Rev Genet. 2020;21(10): 581-96.

25. Wu W, Zhang HH, J. WM, B KP. Gene Biotechnology. 2nd ed. Boca Raton, FL: CRC
Press; 2003.

26. Stoeger T, Amaral LAN. The characteristics of early-stage research into human genes
are substantially different from subsequent research. PLOS Biol. 2022;20(1): e3001520.

27. The UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. Nucleic
Acids Res. 2021;49(D1): D480-9.

28. Narin F, Noma E. Is technology becoming science? Scientometrics. 1985;7: 369-81.

29. Narin F, Noma E, Perry R. Patents as indicators of corporate technological strength. Res
Policy. 1987;16: 143-55.

30. Meyer M. Does science push technology? Patents citing scientific literature. Res Policy.
2000;29: 409-34.

31. Sampat B, Williams HL. How do patents affect follow-on innovation? Evidence from the
human genome. Am Econ Rev. 2019;109(1): 203-36.

32. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al.
1116   SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically
1117   proven protease inhibitor. Cell. 2020;181(2): 271-80.

1118   33. Stoeger T, Amaral LAN. Meta-Research: COVID-19 research risks ignoring important host
1119   genes due to pre-established research patterns. eLife. 2020;9: e61981.

1120   34. Stephanopoulos G, Aristidou AA, Nielsen JH. Metabolic Engineering : Principles and
1121   Methodologies. San Diego: Academic Press; 1998.

1122   35. Sun L, Lee JW, Yook S, Lane S, Sun ZQ, Kim SR, et al. Complete and efficient
1123   conversion of plant cell wall hemicellulose into high-value bioproducts by engineered
1124   yeast. Nat Commun. 2021;12(1): 4975.

1125   36. Provart NJ, Alonso J, Assmann SM, Bergmann D, Brady SM, Brkljacic J, et al. 50 years
1126   of *Arabidopsis* research: highlights and future directions. New Phytol. 2016;209(3): 921-
1127   44.

1128   37. Wilson GA, Bertrand N, Patel Y, Hughes JB, Feil EJ, Field D. Orphans as taxonomically
1129   restricted and ecologically important genes. Microbiol-Sgm. 2005;151: 2499-501.

1130   38. Khalturin K, Hemmrich G, Fraune S, Augustin R, Bosch TCG. More than just orphans:
1131   are taxonomically-restricted genes important in evolution? Trends Genet. 2009;25(9):
1132   404-13.

1133   39. Youn H, Strumsky D, Bettencourt LMA, Lobo J. Invention as a combinatorial process:
1134   evidence from US patents. J R Soc Interface. 2015;12(106): 20150272.

1135   40. Fink TMA, Reeves M. How much can we influence the rate of innovation? Sci Adv.
1136   2019;5(1): eaat6107.

1137   41. Ke Q. Technological impact of biomedical research: The role of basicness and novelty.
1138   Res Policy. 2020;49: 104071.

1139   42. Ullmann A, Jacob F, Monod J. Characterization by *in vitro* complementation of a peptide
1140   corresponding to an operator-proximal segment of beta-galactosidase structural gene of
1141   *Escherichia coli*. J Mol Biol. 1967;24(2): 339-43.

1142   43. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable
1143   dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science.
1144   2012;337(6096): 816-21.

1145   44. Topol EJ. Failing the public health—rofecoxib, Merck, and the FDA. N Engl J Med.
1146   2004;351: 1707-9.

1147   45. Shin DM, et al. Chemoprevention of head and neck cancer by simultaneous blocking of
1148   epidermal growth factor receptor and cyclooxygenase-2 signaling pathways: preclinical
1149   and clinical studies. Clin Cancer Res. 2013;19(5): 1244-56.

46. Saba NF, et al. Chemoprevention of head and neck cancer with celecoxib and erlotinib:
results of a phase Ib and pharmacokinetic study. Cancer Prev Res. 2014;7(3): 283-91.

47. Groeger AL, et al. Cyclooxygenase-2 generates anti-inflammatory mediators from
omega-3 fatty acids. Nat Chem Biol. 2010;6(6): 433-41.

48. Chen JS, et al. Nonsteroidal anti-inflammatory drugs dampen the cytokine and antibody
response to SARS-CoV-2 infection. J virol. 2021;95(7): e00014-21.

49. Jackson G, Gillies H, Osterloh I. Past, present, and future: a 7-year update of Viagra®
(sildenafil citrate). Int J Clin Pract 2005;59(6): 680-91.

50. Bhogal S, Khraisha O, Al Madani M, Treece J, Baumrucker SJ, Paul TK. Sildenafil for
pulmonary arterial hypertension. Am J Ther. 2019;26(4): e520-6.

51. U.S. National Institutes of Health. ClinicalTrials.gov. Available from:
http://www.clinicaltrials.gov.

52. Tinsley HN, et al. Colon tumor cell growth-inhibitory activity of sulindac sulfide and other
nonsteroidal anti-inflammatory drugs is associated with phosphodiesterase 5 inhibition."
Cancer Prev Res. 2010;3(10): 1303-13.

53. Tinsley HN, et al. Inhibition of PDE5 by sulindac sulfide selectively induces apoptosis and
attenuates oncogenic Wnt/β-catenin–mediated transcription in human breast tumor cells.
Cancer Prev Res. 2011;4(8): 1275-84.

54. Nelson SF, Miceli MC, Elashoff RM, Sweeney HL, Victor RG. PDE5 inhibition alleviates
functional muscle ischemia in boys with Duchenne muscular dystrophy. Neurology.
2014;82(23): 2085-91.

55. Wang R, Lei X, Rakesh CK. PDE5 inhibitor tadalafil and hydroxychloroquine cotreatment
provides synergistic protection against type 2 diabetes and myocardial infarction in mice.
J Pharmacol Exp Ther. 2017;361(1): 29-38.

56. Amedro P, et al. Efficacy of phosphodiesterase type 5 inhibitors in univentricular
congenital heart disease: the SV-INHIBITION study design. ESC Heart Fail. 2020;7(2):
747-56.

57. Ohaegbulam KC, Assal A, Lazar-Molnar E, Yao Y, Zang XX. Human cancer
immunotherapy with antibodies to the PD-1 and PD-L1 pathway. Trends Mol Med.
2015;21(1): 24-33.

58. Zou ZL, Tao T, Li HM, Zhu X. mTOR signaling pathway and mTOR inhibitors in cancer:
progress and challenges. Cell Biosci. 2020;10(1): 31.

59. Mantovani F, Collavin L, Del Sal G. Mutant p53 as a guardian of the cancer cell. Cell
Death Differ. 2019;26(2): 199-212.

60. Lippincott-Schwartz J, Patterson GH. Development and use of fluorescent protein
markers in living cells. Science. 2003;300(5616): 87-91.

61. Liu S, Su YC, Lin MZ, Ronald JA. Brightening up biology: advances in luciferase systems
for *in vivo* Imaging. Acs Chem Biol. 2021;16(12): 2707-18.

62. Liu JQ, Xiao Q, Xiao JN, Niu CX, Li YY, Zhang XJ, et al. Wnt/beta-catenin signalling:
function, biological mechanisms, and therapeutic opportunities. Signal Transduct Tar.
2022;7(1): 3.

63. Zhang Y, Wang X. Targeting the Wnt/beta-catenin signaling pathway in cancer. J
Hematol Oncol. 2020;13(1): 165.

64. English K, Barry FP, Field-Corbett CP, Mahon BP. IFN-γ and TNF-α differentially
regulate immunomodulation by murine mesenchymal stem cells. Immunol Lett.
2007;110(2): 91-100.

65. Liu X, Ji Q, Zhang C, Liu X, Liu Y, Liu N, et al. miR-30a acts as a tumor suppressor by
double-targeting COX-2 and BCL9 in H. pylori gastric cancer models. Sci Rep. 2017;7(1):
7113.

66. Biswas KH, Sopory S, Visweswariah SS. The GAF domain of the cGMP-binding, cGMP-
specific phosphodiesterase (PDE5) is a sensor and a sink for cGMP. Biochemistry.
2008;47(11): 3534-43.

67. Küçük A, Yucel M, Erkasap N, Tosun M, Koken T, Ozkurt M, et al. The effects of PDE5
inhibitory drugs on renal ischemia/reperfusion injury in rats. Mol Biol Rep. 2012;(39):
9775-82.

68. Tavallai M, Hamed HA, Roberts JL, Cruickshanks N, Chuckalovcak J, Poklepovic A, et
al. Nexavar/Stivarga and viagra interact to kill tumor cells. J Cell Physiol. 2015;230(9):
2281-98.

69. Krimsky S. Science in the Private Interest: Has the Lure of Profits Corrupted Biomedical
Research? Lanham, MD: Rowman & Littlefield; 2003.

70. BP cuts funding for 'most promising' biofuel. Investigate Midwest. August 20, 2015;
https://investigatemidwest.org/2015/08/20/bp-cuts-funding-for-most-promising-biofuel-2.

71. Stocker M, Baffes J, Vorisek D. What triggered the oil price plunge of 2014–2016 and
why it failed to deliver an economic impetus in eight charts. World Bank Blogs. January
18, 2018; https://blogs.worldbank.org/en/developmenttalk/what-triggered-oil-price-plunge-
2014-2016-and-why-it-failed-deliver-economic-impetus-eight-charts.

72. Asase RV, Okechukwu QN, Ivantsova MN. Biofuels: present and future. Environ Dev
Sustain. 2024; https://doi.org/10.1007/s10668-024-04992-w.

73. Mandt R, Seetharam K, Cheng CHM. Federal R&D funding: the bedrock of national
innovation. MIT Sci Policy Rev. 2020;1: 44-54.

74. Marx M, Fuegi A. Reliance on science by inventors: Hybrid extraction of in-text patent-to-
article citations. J Econ Manage Strat. 2022;31(2): 369-92.

75. Priem J, Piwowar H, Orr R. OpenAlex: A fully-open index of scholarly works, authors,
venues, institutions, and concepts. arXiv:2205.01833 [Preprint]. 2022. Available from:
https://arxiv.org/abs/2205.01833

76. Lin Z, Yin Y, Liu L, Wang D. SciSciNet: A large-scale open data lake for the science of
science research. Sci Data. 2023;10(1): 315.

77. Haessler P, Giones F, Brem A. The who and how of commercializing emerging
technologies: A technology-focused review. Technovation. 2023;121: 102637.

78. Levinthal DA. The slow pace of rapid technological change: Gradualism and punctuation
in technological change. Ind Corp Change. 1998;7(2): 217-47.

79. Scaringella L, Miles RE, Truong Y. Customers involvement and firm absorptive capacity
in radical innovation: The case of technological spin-offs. Technol Forecast Soc Change.
2017;120: 144-62.

80. Arthur WB. The structure of invention. Res Policy 2007;36: 274-87.

81. Arnold C. Death by climate change. Nat Clim Change. 2022;12(7):607-9.

82. Perry BL, Aronson B, Pescosolido BA. Pandemic precarity: COVID-19 is exposing and
exacerbating inequalities in the American heartland. Proc Natl Acad Sci U S A.
2021;118(8): e2020685118.

83. Yun J, Kim P-J, Jeong H. Anatomy of scientific evolution. PLOS ONE. 2015;10(2):
e0117388.

84. Chen CY, Kahanamoku SS, Tripati A, Alegado RA, Morris VR, Andrade K, et al.
Systemic racial disparities in funding rates at the National Science Foundation. eLife.
2022;11: e83071.

85. Bairoch A, Apweiler R, Wu CH, Barker WC, Boeckmann B, Ferro S, et al. The universal
protein resource (UniProt). Nucleic Acids Res. 2005;33: D154-D9.

86. Browet A, Absil P-A, Dooren PV. Fast community detection using local neighbourhood
search. arXiv:1308.6276 [Preprint]. 2013. Available from: https://arxiv.org/abs/1308.6276

87. Lin Y, Michel J-B, Aiden EL, Orwant J, Brockman W, Petrov S. Syntactic annotations for
the Google Books Ngram Corpus. Proceedings of the 50th Annual Meeting of the
Association for Computational Linguistics. 2012: 169-74.

88. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database
resources of the National Center for Biotechnology Information in 2023. Nucleic Acids
Res. 2022;50: D20-6.

89. Fredkin E. Trie Memory. Commun Acm. 1960;3(9): 490-9.

90. Wei CH, Allot A, Lai PT, Leaman R, Tian S, Luo L, et al. PubTator 3.0: an AI-powered
literature resource for unlocking biomedical knowledge. Nucleic Acids Res, 2024;52(W1):
W540-6.

91. The Gene Ontology Consortium. The Gene Ontology resource: enriching a GOld mine.
Nucleic Acids Res. 2021;49(D1): D325-34.
92. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a
major update to the DrugBank database for 2018. Nucleic Acids Res. 2018;46(D1):
D1074-82.
93. Ghosh M, Erhardt S, Rose ME, Buunk E, Harhoff D. PaECTER: patent-level
representation learning using citation-informed transformers. arXiv:2402.19411 [Preprint].
2024. Available from: https://arxiv.org/abs/2402.19411
94. Fruman DA, Christian R. PI3K and cancer: lessons, challenges and opportunities. Nat
Rev Drug Discov. 2014;13(2): 140-56.