Environmental and geographic drivers of global bat phylogenetic diversity

Abigail E. Green[1], Camilo A. Calderón-Acevedo[3], J. Angel Soto-Centeno[4], Tara A. Pelletier[2]

1. Department of Mathematics & Statistics, Radford University, Radford, VA 24142
2. Department of Biology, Radford University, Radford, VA 24142
3. Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá
4. Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural History,
New York, NY 10024

Biosketch

Our team is interested in exploring biodiversity patterns in a wide-range of taxa using methods from
population genetics, phylogeography, and systematics.

AEG: www.linkedin.com/in/abigail-abbi-green

TAP: https://sites.google.com/site/taraapelletier/

CCA: https://mammalbiogeo.wordpress.com/

ASC: https://www.mormoops.com/

Abstract
Aim
Understanding the patterns and factors that shape biodiversity is vital to conserving species. We
combined open-source genetic, environmental, and geographic information to analyze bat phylogenetic
diversity (PD) patterns in continuous ecoregions across the globe. This information is important for
developing bat conservation strategies, and our methodology can work for any taxa with sufficient
georeferenced genetic data available.

Location
Global.

Methods
After curating a global dataset containing 14,037 COI DNA sequences from 343 described species of
bats, we calculated PD for continuous ecoregions at different spatial scales. To avoid the difficulties of
using current species names, we used genetic OTUs identified by a single-locus species delimitation
method to reconstruct and date a phylogeny. We then calculated PD, estimated a lineage through time
plot, and used random forest predictive modeling to identify environmental and geographic predictors
of PD.

Results
In addition to current temperature, temperature during the last glacial maximum and temperature
changes between the last glacial maximum and last interglacial were most closely associated with PD.
However, at different spatial scales, the top variables differed slightly. When using smaller ecoregions,

latitude and population density were also identified as important, though not significant. Southeast Asia
and South America had the highest levels of PD, along with parts of Africa and the Himalayas. We
demonstrate that, regardless of spatial scale and uneven sampling across the globe, single-locus genetic
data can reflect species diversity gradients and identify predictors of PD.
46
Main conclusions
We show that publicly-available, single-locus data can be used to analyze large-scale evolutionary
patterns and inform conservation efforts. Additionally, choice of biodiversity measure and spatial scale
matter when assessing species patterns. When looking at bats, temperature variables with a historical
component are most important for predicting PD broadly, but latitude and population density could also
be important on smaller scales.
53
54 **Keywords**
55 Chiroptera, phylogenetic diversity, COI, species delimitation, ecoregions, random forest
56
57
58 **Introduction**
59
Considering multiple components of biodiversity is necessary for understanding the relative roles of
evolutionary processes in shaping communities at different spatial scales (Stevens & Gavilanez, 2015),
and different measures of biodiversity may vary across different spatial scales. For example, species
richness and within-species genetic diversity are not always evenly or similarly distributed across
geographic space and can be influenced by different evolutionary processes (Amador, et al., 2024;
Brown et al., 2020; Lino, et al., 2021). Within-species genetic diversity and phylogeographic breaks
follow a similar pattern to the latitudinal species diversity gradient (da Silva Fonseca, et al., 2024; Smith
et al., 2014) while patterns of isolation-by-distance or -environment are better explained by range size
(Pelletier & Carstens, 2018; Amador et al. 2025). Documenting these differing biodiversity components
is important for structuring conservation efforts that are effective and efficient (Harper et al., 2021;
Hosegood et al., 2020). Phylogenetic diversity accounts for branch lengths on a phylogenetic tree (Faith,
1992) or genetic distance (i.e., divergence) among species and is therefore a useful measure of
biodiversity to consider for conservation purposes. Assemblages of species that contain more distantly
related species have been shown to consist of and produce higher levels of biomass (Cadotte, 2013).
More distantly related groups also capture more diversity in species traits, which increases future
potential for adaptation (Forest et al., 2007) and may better preserve natural processes. Phylogenetic
diversity is also listed as an important consideration by the International Union for Conservation of
Nature (www.pdtf.org) and globally can contribute to the Convention on Biological Diversity's global
conservation efforts (www.cbd.int).
Most data available in open-source repositories can lead to new ways of documenting and exploring
biodiversity, especially by integrating environmental, geographic, and genetic data. For instance, efforts
to create global maps of biodiversity can be useful for tracking changes over time (Perreira 2016),
identifying areas of high diversity in need of conservation (Pelletier, et al., 2018), and understanding the
evolutionary processes promoting biodiversity (Guralnick & Hill, 2009; Theissinger et al., 2023). Recent

studies using open-source data to document biodiversity provide information to facilitate conservation
action and may help unravel the processes that shape highly speciose regions (e.g., phylogeographic
breaks in birds: Smith, et al., 2017; response to Pleistocene glaciations in bats: Carstens, et al., 2018;
hidden diversity in mammals: Parsons, et al., 2022). Conservation initiatives that include more complex
levels of biodiversity, incorporate evolutionary history, and capture functional diversity can help
maintain the stability of a wide range of taxonomic groups and minimize species loss. For example,
when space for protected areas is limited, targeting phylogenetic diversity can more effectively conserve
a larger proportion of the mammal tree of life (Rosauer, et al., 2017; Rosauer et al., 2018).
92
Increased anthropogenic land use and habitat fragmentation can negatively affect the composition of
biodiverse communities (Newbold et al, 2015; López￼Baucells et al., 2022). Habitat degradation reduces
ecosystem stability and productivity which endangers important ecosystem services, like food
production, at local and regional scales (Cardinale et al., 2012; Moreira-Hernández, et al., 2021).
Additionally, the increased rate of the rise in global temperatures caused by human emissions of
greenhouse gases and pollution is making many habitats inhospitable by humans, wildlife, and
vegetation alike (Grimm et al., 2013; Weiskopf et al., 2020). A better understanding of the key
environmental and geographic factors that both foster and discourage long-term biodiversity trends can
help improve conservation decision making and land use management. Strong conservation initiatives
are now more pertinent than ever given the rapid increase in biodiversity and habitat loss that has led to
the endangerment and extinction of many species in recent millennia (Ceballos, et al., 2020; Newbold et
al, 2015). Furthermore, climate change is inevitable and understanding how biodiversity responds to the
climate in its environment can have useful conservation implications.
106
Many studies have examined the effects of environmental factors on phylogenetic diversity (PD) across
multiple taxonomic and geographic scales (Grimshaw & Higgins, 2017; Carvalho, et al., 2021; Paz et al.,
2021; Montaño-Centellas, et al., 2023; Hu et al., 2021). The importance of different environmental
factors may change with the measure of phylogenetic diversity used, taxonomic group, or spatial scale
(Grimshaw & Higgins, 2017; Paz et al., 2021; Montaño-Centellas et al., 2023). For example, Paz et al.
(2021) found that, in plants, mean nearest taxon distance (MNTD) and mean pairwise distance (MPD)
were more closely associated with temperature while Faith's index was more closely associated with
precipitation. Nonetheless, there is limited information about the effects of environmental factors on
biodiversity globally because most studies are conducted at regional geographic scales (Hu et al., 2021).
PD in the bats of Mexico (Grimshaw & Higgins, 2017) and the Atlantic Forest of Brazil (Stevens and
Gavilanez 2015) was best explained by temperature, fluctuations in temperature, and latitude. Elevation
predicted phylogenetic turnover in phyllostomid bats in the Amazon and Atlantic Forest of Brazil (de
Carvalho et al., 2019; Carvalho et al., 2023). For all mammals in western Mexico, however, precipitation
was a more important factor for taxonomic diversity (Mason-Romo et al., 2017). On a global scale,
raptor PD was lower in areas with historically high temperatures and higher human disturbance
(Montaño-Centellas et al., 2023).

Bats (order Chiroptera) comprise 1500 species, representing about 22% of crown mammal diversity
(Mammal Diversity Database, 2024; Simmons & Cirranello, 2025). Two suborders and 21 families of bats
are currently recognized (Teeling et al., 2005). The phylogenetic history of bats reflects a deep
diversification event where two suborders diverged ~63mya (Foley et al. 2023; Hao et al. 2024). The
suborder Yinpterochiroptera is composed of five families of megabats and smaller insectivore and
carnivore species. The suborder Yangochiroptera contains the remaining families and exhibits the largest
ecological variation. This ecological variation is evident when comparing the most diverse families:
Vespertilionidae, a largely insectivorous lineage with one carnivorous representative; Phyllostomidae in
the Neotropics with less than half the bat species (230), but with the largest Eltonian and Grinnelian
niche amplitudes of any mammal clade (Fleming et al., 2020); and Pteropodidae, the flying foxes, which
are primarily nectarivorous and frigivorous. Given their unique ability to fly, many bats may be able to
disperse across large areas providing important ecosystem services like insect pest control, pollination of
over 500 species of angiosperms by the three aforementioned mentioned families (Muchhala &
Tschapka 2020; Moréira-Hernández et al., 2021), and seed dispersal (Kunz, et al., 2011; Moréira-
Hernández et al., 2021; Ramírez-Fráncel et al., 2022). Over one-third of bat species assessed by the IUCN
are considered threatened or data deficient, and 80% have unknown or decreasing population trends
(Frick, et al., 2020). Human derived activities at local, regional, and global scales are the main source of
bat species decline and population fragmentation (Frick et al., 2020; Soto-Centeno & Calderón-Acevedo,
2023); however, we have limited knowledge on how phylogenetic diversity relates to habitat and
environmental change across space and time in most taxonomic groups, especially at a global scale.
144
We explored how environmental and geographic factors affect bat phylogenetic diversity on a global
scale. First, we estimated phylogenetic diversity using georeferenced DNA sequences and species
delimitation methods across ecoregions. Then, we used a random forest predictive algorithm to
determine which geographic and environmental variables had the largest impact on phylogenetic
diversity measures in continuous ecoregions across the globe. We also included historical climatic data
starting in the Pliocene because paleoclimate leaves impressions on species ranges and diversity
(Svenning et al. 2015), so it will be important to consider future climate change in conservation planning
on a global scale (i.e., ecoregions; Dobrowski et al. 2021). We expect largely heterogeneous ecoregions
with more stable, warmer temperatures to contain the highest levels of PD given that landscape barriers
to dispersal and time should promote speciation, particularly in tropical areas (Smith et al. 2014).
Furthermore, given regional studies conducted in bats, climate variables pertaining to temperature and
historical climate could also have an impact on PD on larger scales. Our study serves as a first step in
defining areas that contain high levels of phylogenetic diversity and the conditions under which
biodiversity could be maintained and promoted moving forward through estimates of PD across
ecoregions with heterogeneous habitats that should capture broad-scale biodiversity and evolutionary
potential.
161

Methods

DNA sequence data and OTUs

We obtained data from public databases, such as Genbank and the Global Biodiversity Information
Facility (GBIF), through the *phylogatR* platform (Pelletier et al. 2022; https://phylogatr.org/) using the
search term 'Chiroptera' on February 22, 2022. The total dataset represented 17 out of 21 (80%)
families, 429 out of 1482 (29%) species, and 18 loci with a total of 20,020 DNA sequences. *PhylogatR*
provides DNA sequence alignments on a species-by-species basis that are almost analysis ready. In this
study, we focused on the mitochondrial gene *Cytochrome oxidase I* (*COI*) as this locus was the best
represented in the data. We first checked for potentially misidentified sequences. Each species
sequence alignment was run through BLAST via the NCBI database. The top hit for each DNA accession
was compared to the species name associated with the DNA sequence in *phylogatR*. Five hundred thirty
sequences that were considered not plausible data (i.e., too short) and two that did not BLAST to bats
(i.e. *Mus musculus* and *Apodemus draco*) were removed from the alignments. Cleaned alignments from
this step were merged for all species and re-aligned using MAFFT v7.5 (Kotah & Standley, 2013) with the
default settings and –adjustdirection. We visually inspected the concatenated *COI* sequence alignment
and trimmed the ends of the sequences to remove large areas with missing data resulting in a 658 base-
pair region of this barcoding gene. Our final dataset consisted of 14,037 *COI* DNA sequences found
globally (Fig. 1; Table S1). Of these, 1215 (8%) had species names in GenBank that did not match records
existing in the GBIF or the Barcode of Life Data System (BOLD). Because we used OTUs as units of
analysis, we chose to retain these taxa without adjusting taxonomy (details below).
183
Mammal taxonomy is rapidly changing (Burgin, et al., 2018; Parsons et al., 2022; Reeder, et al., 2007;
Solari & Martínez-Arias, 2014; Simmons & Cirranello, 2025; Mammal Diversity Database, 2024) and
species names are often incorrect in online databases (Mulcahy et al., 2022). Therefore, we used OTUs
estimated from the DNA sequence data itself, rather than species names, as terminal taxa in our
phylogeny. By using OTUs, we captured biodiversity above and below currently named species diversity,
which can have useful conservation implications (Rosauer et al., 2018). This is crucial, particularly for
groups with complicated taxonomy, such as bats (Calderón-Acevedo, et al., 2021; Calderón-Acevedo, et
al., 2022; da Silva Fonseca, 2024; Mônico & Soto-Centeno, 2024; Rodríguez-Posada, et al., 2021; Yi et al.,
2024) and small mammals in general (Parsons et al. 2022). To estimate genetic lineages and define
OTUs, we used the distance-based species delimitation method Assemble Species by Automatic
Partitioning (ASAP; Puillandre, et al., 2020), as tree-based methods (e.g., GMYC [Pons et al., 2006] or PTP
[Zhang, et al., 2013]) would be prohibitive given the size of our dataset. All default settings in ASAP were
used. This method establishes a set of ten different partitions ranked by an ASAP score. Each partition is
a list of genetically distinct groups defining which individuals belong to each group. The ASAP score is a
combination of the probability that each ASAP group is a single species along with the width of the
barcode gap between the current and previous partition. Generally, the smaller the ASAP score, the
better the partition is (Puillandre et al., 2020), so we used the partition with the lowest score (i.e., best
ranked).
202
Phylogeny reconstruction and estimates of divergence
One DNA sequence was randomly sampled from each ASAP genetic group (i.e., OTU) to estimate a
phylogenetic tree representing each *COI* lineage present in our dataset. This phylogeny (Supporting files
S1 & S2) was then used in subsequent estimates of divergence times, lineage through time plots, and

phylogenetic diversity (see below). We first used ModelFinder (Kalyaanamoorthy et al., 2017) to select
the best model of nucleotide evolution. Then, we implemented the resulting model (GTR+F+R8) in
RAxML-NG (Kozlov, et al., 2019). This analysis ran for an initial search of 10 Maximum Likelihood (ML)
trees followed by a 1000 bootstrap tree search (see Supporting Information). In this analysis, we used
Yinpterochiroptera as an outgroup to all other bats following well-supported previously published bat
relationships that combined mitochondrial and nuclear sequences (Shi & Raboski, 2015; Teeling et al.,
2005). Given the limited information in a single-locus dataset, we wanted to ensure that we followed
major accepted bat relationships (Teeling et al., 2005; Shi & Raboski, 2015; Baker, et al., 2016;
Cirranello, et al., 2016; Rojas, et al., 2016). A configuration file (Supporting file S3) for the RaxML-NG
analysis included constrained relationships at the level of superfamily and family, but allowing the
inference of subfamily, genera, and species/OTU (all intermediate tree construction outputs can be
found in Supporting files S1-S12).
219
We used the rooted constrained RaxML-NG phylogeny (Supporting files S2 and S9) to compute
divergence time estimates with the penalized likelihood semi-parametric method available in TreePL
(Smith & O'Meara, 2012). This method allows for the estimation of different rates on different branches
on a previously estimated phylogeny with branch lengths. To estimate divergences, we used the node
ages of Yingochiroptera and Yangochiroptera (71–58 my), Noctilionoidea (37–47 my), the node of
Pteropodidae and Rhinolophoidea (53–63 my), the node between *Pteronotus* and *Mormoops* (36.2–30.8
my), the internal node of Stenodermatinae and Phyllostominae within Phyllostomidae (36.2–27.33), and
the first node within the subfamily Desmodontinae (23.9–16.7 my) as calibration points previously
available from the literature (see Teeling et al. 2005; Shi & Raboski, 2015; Rojas et al., 2016). The TreePL
analysis then followed an initial priming run where optimization parameters were obtained, followed by
a cross-validation step to select the best smoothing parameter for the penalized likelihood estimates
(Supporting file S10, Smith & O'Meara, 2012). The resulting divergence time estimates (Supporting file
S11) were used to construct a lineage through time (LTT) plot with the R packages 'ape' and 'phytools'
(Paradis & Schliep, 2019; Revell, 2024). The LTT plot was produced using the family Phyllostomidae as a
reference because this clade of bats shows a remarkable rate of species diversification in the Neotropics
(Shi & Raboski, 2015; Rojas et al., 2016).
236
Phylogenetic diversity and geographic units, ecoregions
The most common ways to measure phylogenetic diversity include Faith's index, mean pairwise distance
(MPD), and mean nearest taxon distance (MNTD) (Faith, 1992; Warwick & Clarke, 1995; Webb, et al.,
2002; Webb, 2000). Faith's index sums of all the branch lengths in a phylogenetic tree. Mean pairwise
distance calculates average distance among all species in a phylogenetic tree. Mean nearest taxon
distance calculates the average distance between a species and its closest evolutionary neighbor.
Because MPD and MNTD include measurements of branch lengths based on nearest neighbors and not
total branch lengths, these measures, MPD in particular, should be somewhat protected from missing
species, as they focus on relationships between tips of the phylogeny (Cadotte et al. 2012; Li et al.
2012); however, we used the standardized effect size (SES) approach for MPD and MNTD to better
control for missing taxa. OTU richness was calculated for each ecoregion (see below), and all three
measures of PD were estimated using the R package 'picante' (Kembel et al., 2010).

To estimate levels of PD across the globe, we chose to analyze data by ecoregion rather than using grid
cells that may be arbitrary from a biological perspective (Paz-Vinas et al., 2021). From a conservation
perspective, an ecoregion approach can yield higher representation of novel habitats, natural
communities, and species-level diversity than other units of analysis such as protected areas or political
boundaries; therefore, more adequately protecting species, their natural processes, connectivity, and
the species they interact with (Funk and Fa 2010; Giakoumi et al. 2013; Hanson et al. 2022; Zoderer et
al. 2024). Our goal was not to focus on single species or endemics, but to use the presence or absence of
a lineage in each ecoregion to identify the ecoregions that may foster the persistence of biodiversity in
bats, as ecoregions are ecologically cohesive but span a suite of novel habitats and specialized niches
(Zoderer et al. 2024).

We used the 846 RESOLVE Ecoregions 2017 dataset (Olson et al., 2001; Dinerstein et al., 2017). There
were 161 ecoregions containing DNA sequences in our dataset. However, many of those had very few
species, limiting our ability to make comparisons of OTU level diversity. For example, the 161 ecoregions
ranged from 2 – 85 OTUs per ecoregion and had an average of 8.96 OTUs per region. There were 109
out of 161 ecoregions that had more than two OTUs represented and could be used to estimate PD. We
also created a second set of larger continuous ecoregions by grouping RESOLVE Ecoregions that
contained DNA sequences from our dataset, were adjacent to each other, and belonged to the same
biome. This dataset contained 25 continuous ecoregions, from 185 RESOLVE Ecoregions, as some
regions contained small fragmenting ecoregions that were included to create a continuous space (Table
S2). There were 3 – 117 OTUs per ecoregion with an average of 22.44 OTUs. We used both datasets, a
109-eco set that used the smaller single ecoregions and a 25-eco set with combined ecoregions, to
conduct subsequent analyses. Investigating the same sets of predictor variables at different spatial
scales allowed us to determine whether spatial scale can influence trends in bat diversity.
We also conducted linear regressions to test whether OTU richness explained the different measures of
PD to determine whether PD captures a different feature of biodiversity other than simply OTU richness,
especially given that our bat dataset is not complete (i.e., open-source databases rarely contain all
species). OTU richness explained almost all of the variation in Faith's index (109-eco: R^2 = 0.9767, p <
0.001; 25-eco: R^2 = 0.9815, p < 0.001), so we moved forward with predictive modeling on PD
measurements that were not significantly correlated with OTU richness: MNTD (109-eco: R^2 = 0.04689, p
= 0.02372; 25-eco: R^2 = 0.03237, p = 0.1925) and MPD (109-eco: R^2 = 0.1139, p < 0.001; 25-eco: R^2 =
0.1218, p = 0.0488). PD estimates were used as the response variable in the predictive modeling
framework described below.

Predictor variables

We curated a set of predictor variables using 115 environmental data layers (Table S3) and geographic
information associated with each ecoregion. We took the median and standard deviation of each data
layer in each region by randomly sampling 1000 points per region. This was done to capture information
about the ecoregion as a whole and the habitat heterogeneity that contributes to or inhibits
connectivity, which might promote or hinder speciation and be important under future climate change

scenarios (Dobrowski et al. 2021). Identifying the environmental factors that contribute to biodiversity
may help predict areas that could harbor the most biodiversity. The data layers included 19 BIOCLIM
layers for present time from the Chelsa database (Karger et al., 2017; Karger et al., 2018; Karger, et al.,
2021) at 1km resolution, elevation (Tachikawa et al., 2011), population density (CIESIN, 2017), terrestrial
habitat heterogeneity (Tuanmu & Jetz, 2015), gross domestic product (GIS processing World Bank
DECRG, 2010), global land cover classification (European Space Agency, 2009), global river classification
(Dallaire, et al., 2019), disaster risk (Peduzzi, 2019), anthropogenic biome (Ellis, et al., 2010), global lakes
and wetlands database (Lehner & Döll, 2004), and various indicators of seasonal growth (Karger et al.,
2021). We also included BIOCLIM layers for four different historical time periods: Mid-Holocene (ca.
8.326-4.2 ka); Last Glacial Maximum (LGM, ca. 21 ka), Last Interglacial (LIG, ca. 130 ka), and Pliocene (ca.
3.3 ma) (Fordham et al., 2017; Karger et al., 2021; Otto-Bliesner, et al., 2006; Dolan, et al., 2015)**.** We
used the R packages 'raster' (v3.6-14; Hijmans, 2023), 'tools' (R Core Team, 2022)**,** and 'sf' (Pebesma,
2018) to extract ecoregion specific information for each of these layers. We also calculated the median,
standard deviation, maximum, and minimum values for latitude and longitude of each ecoregion and
used the st_area function to calculate the area of each ecoregion (Pebesma, 2018).

To capture change of climate variables over time, we calculated the difference in the median values
between time periods: Pliocene—LIG, LIG—LGM, LGM—Holocene, and Holocene—present in addition
to the median and standard deviation of each layer. This resulted in 310 variables that represent a
variety of environmental and geographic traits for each ecoregion (Tables S4 & S5). Twelve of the 310
variables for the 25-eco dataset contained NA values and were removed from the dataset, as the
random forest algorithm does not handle missing data. These twelve variables included the median and
standard deviation for three growing season temperature variables, snow water equivalent, and two
global lakes and wetlands classifications. There were six additional variables that contained NA values
and were removed for the 109-eco dataset: the mean and standard deviation for the last day of the
growing season, the first day of the growing season, and global drought events. The resulting 25-eco
dataset contained 298 variables and the 109-eco dataset contained 292 variables for further
exploration.
318
Predictive modeling
To reduce the number of variables in our predictive models, we used different sets of predictor variables
based on correlation coefficients and variable categories. We first calculated pairwise Pearson
correlation coefficients for all environmental and geographic variables using the R package 'Hmisc'
(Harrell Jr., 2023) and removed predictor variables with a correlation coefficient greater than or equal to
0.75 (Tables S6 & S7). We then split the variables into seven groups based on category of variable and
ran a Random Forest (RF) for each group: 1) current climate, 2) geography (including latitude, longitude,
flood risk, growing season, anthropogenic biomes, etc.), 3) Holocene climate, 4) LGM climate, 5) LIG
climate, 6) Pliocene climate, and 7) climate change (the difference across time periods). We ran these
category analyses using all variables and with the correlation cutoff of 0.75. In total, we ran 14 RF
analyses with different sets of predictor variables for both MPD and MNTD as response variables with
the R package 'randomForest' (Liaw & Wiener, 2002) using the default settings, 1000 trees, and
retaining the importance values. While random forest handles a high variable to observation ratio

relatively well (Genuer et al. 2008), we conducted 100 iterations for each RF to assess stability of the
model.

Based on the predictive power of the RF models (Table 1), we focus on MPD as our phylogenetic
diversity response variable. Percent increase in mean squared error (MSE) was calculated for each
variable when removed from the model to determine which variables contribute the most to the
prediction. We used this value to demonstrate the relative importance of each variable. To further
reduce the variables in our data and assess which were most important, we took the top 2-4 predictors
from the seven random forest models (each category listed above with 0.75 or higher correlated
variables removed) to run a final RF. Variables were chosen for the final RF if they were ranked in the
top five most important predictors for their model in 100% of the RF iterations . There were 19 top
predictors for the 109-eco set and 23 top predictors for the 25-eco set across the seven different
categories of variables (Tables S8 & 9). For the top predictors in the final RF models, we used the R
package 'nlme' (Pinheiro & Bates, 2000; Pinheiro et al., 2025) to conduct generalized least squares (GLS)
linear regressions and test the overall significance of the predictor variables while accounting for the
spatial autocorrelation that commonly arises in geographical analyses like ours.

Finally, we conducted another round of RF with the 109-eco set regions that had significant values for
the SES MPD randomization procedure (Table S5). Given that the original dataset contained missing
taxa, we wanted to ensure that any signals in the data were consistent and reliable. This dataset
contained 41 ecoregions. We only used predictor variables sets with highly correlated variables removed
because we were starting with a smaller number of observations. We also only conducted these
analyses with MPD as the prior models with MNTD as the response variable did not demonstrate any
predictive power. We followed the same procedure as above and retained only the top variables from
each category for a final RF analysis (Table S10), followed by GLS to assess significance.

Results

Genetic data and OTUs
After data cleaning and trimming the sequence alignment, there were 658 base-pairs of *COI* from 343
species and 14,037 individual sequences (Fig. 1; Table S1). ASAP identified 372 genetic partitions or
OTUs. With >14,000 sequences, the ASAP analysis took less than 20 minutes on iMac (3.6 GHz 8-Core
Intel Core i9 16 GB 2667 MHz DDR4). The ASAP results largely matched species names but there were
cases where nominal species were split, merged, and sometimes both (Fig. S1; Table S11). More
specifically, 155 nominal bat species matched the ASAP groups exactly, where all individual sequences
from each species only belonged to one ASAP group, 81 nominal species were split into multiple ASAP
groups, and 161 nominal species were combined in 69 ASAP groups. A total of 54 nominal species that
were split into multiple ASAP groups also had individuals merged with other nominal species. The
average value for OTU richness across the 25-eco set was 21.48 with East Asia having the highest (116)
and four different regions (Mountainous region of Asia (Himalayas), Northwest Asia, Northwest North
America, and the Philippines) having the lowest (3) (Table S4). The average value for OTU richness across

the 109-eco set was 12.28 with the Guianan lowland moist forests having the highest (85), followed by
the Northern Annamites rain forests (53), and 23 regions with the lowest (3) (Table S5).
375
Bat phylogeny reconstruction and lineages through time
The estimated phylogeny (Fig. 2 & S2; Supporting file S2) accurately inferred most of the currently
accepted relationships of bats both at the levels of subfamily and genus. Estimates of lineages through
time based on the reconstructed phylogeny demonstrated a pattern congruent with rapid species
diversification in Phyllostomidae, the most speciose Neotropical bat family, compared to more gradual
lineage accumulation for other more deeply or recently divergent families (e.g., Molossidae and
Hipposideridae; Fig. 3 & S3). However, we noted that some deep nodes (i.e., the sister relationship of
Emballonuroidea with and Noctilionoidea) in the phylogeny were recovered with short branches (Fig. 2).
Furthermore, the divergence time estimates produced by TreePL show deeper (i.e., older) divergences
than expected based on previously published studies (e.g., Teeling et al., 2005; Rojas et al., 2016). These
observations were likely due to our use of a single mitochondrial marker with a fast mutation rate
compared to other studies that use a combination of nuclear and mitochondrial data. Despite that, our
choice of using a single mitochondrial marker allowed us to include a wide sampling breadth across 70%
of all bat families worldwide, which facilitated the analysis of drivers of diversity globally (read below).
390
Phylogenetic diversity and predictive modeling
Average MNTD and MPD values for the 25-eco dataset across the ecoregions were 0.38 and 0.62,
respectively. There were no obvious global geographic patterns in PD as measured by MPD, except that
the southern hemisphere contained higher levels of MPD (Fig. S4). Mideast Africa and Southeast Asia
had the highest levels of MPD, followed by regions in Southern North America, Central America,
Northern South America, and Southern Africa. Western North America had the lowest levels of MPD.
Average PD values for the 109-eco dataset for MNTD and MPD across the ecoregions were 0.42 and
0.60, respectively. Ecoregions in Asia had the highest MPD values, followed by some areas in Africa (Fig.
1). Similarly, the southern, more tropical regions of the globe had higher levels of MPD.
400
MNTD as the response variable had low predictive power. For the 25-eco set, at most, 11.2% of the
variation was explained by LGM variables. For the 109-eco set, at most, 19.9% of the variation was
explained by geographic variables. Geography, the LGM, and climate change were the only groups of
variables that produced models with predictive power for MNTD, though still relatively low. The models
using MPD as a response variable explained anywhere from 47.4-67.8% [25-eco] or 33.5-58.4% [109-
eco] of the variation depending on the model. Thus, we focus our attention on the results using MPD as
our our response variable for predictive modeling (Table 1). While the percent variation explained
differed among our MPD RF models, the geographic (25-eco: 67.8%; 109-eco: 57.4%) and climate change
(25-eco: 65.3%; 109-eco: 42.2%) variables had the most explanatory power for both sets of ecoregions,
though all categories had moderate predictive power. For each set of ecoregions, we conducted an RF
analysis with the top 2-4 predictors from each category chosen based on the number of times the
variable was in the top five predictors of the categorical RF analyses (Tables S8 & S9).
413
For the 25-eco set, there were 23 top variables in the final RF (Fig. 4) which included:

Current BIOCLIM: median temperature of the wettest quarter (bio8_med), median maximum
temperature of the warmest month (bio5_med), and standard deviation of temperature seasonality
(bio4_sd)
Geography: median longitude (lon_med) and standard deviation of flood risk(fl_risk_sd)
Holocene BIOCLIM: median temperature of the wettest quarter (bio8_Hol_med), median
temperature of the coldest quarter (bio11_Hol_med), median temperature annual range
(bio7_Hol_med), and median temperature of the warmest quarter (bio10_Hol_med)
LGM BIOCLIM: median temperature of the wettest quarter (bio8_LGM_med), median
temperature of the coldest quarter (bio11_LGM_med), and standard deviation and median temperature
of the warmest quarter (bio10_LGM_sd and bio10_LGM_sd)
LIG BIOCLIM: median minimum temperature of the coldest month (bio6_LIG_med), median
temperature of the coldest quarter (bio11_LIG_med), median temperature of the wettest quarter
(bio8_LIG_med), and median precipitation of the wettest month (bio13_LIG_med)
Pliocene BIOCLIM: median temperature of the wettest quarter (bio8_Plio_med), median
temperature of the coldest quarter (bio11_Plio_med), and standard deviation of the mean temperature
of the driest quarter (bio9_Plio_sd)
Climate Change: median maximum temperature of the warmest month between the LIG and
LGM (bio5_LGM_LIG_meddiff), median temperature of the warmest quarter between the LIG and LGM
(bio10_LGM_LIG_meddiff), and median temperature of the warmest quarter between the Holocene and
LGM (bio10_Hol_LGM_meddiff).
435
This top variable RF model had high predictive power, explaining 78.91% of the variation in MPD. The
top three variables from this analysis were current median temperature of the wettest quarter
(bio8_med), LGM median temperature of the wettest quarter (bio8_LGM_med), and median maximum
temperature of the warmest month between the LIG and LGM (bio5_LGM_LIG_meddiff), all showing up
as top five predictors 100% of the time (Fig. 4). All three variables were significant after implementing a
Bonferroni correction (0.05/3=0.0166) and accounting for spatial autocorrelation (Table 2).
442
For the 109-eco set, there were 19 top variables in the final RF (Fig. 5) which included:
Current BIOCLIM: median temperature of the wettest quarter (bio8_med), median diurnal
temperature range (bio2_med: mean of monthly [max temp – min temp]), and standard deviation of
annual precipitation (bio12_sd)
Geography: median latitude (lat_med), median longitude (lon_med), median anthropogenic
biomes (anth_med), and median SEDAC world population density (gpw_med)
Holocene BIOCLIM: median temperature of the wettest quarter (bio8_Hol_med) and median
temperature of the coldest quarter (bio11_Hol_med)
LGM BIOCLIM: median temperature of the wettest quarter (bio8_LGM_med)
LIG BIOCLIM: standard deviation of the mean precipitation of the wettest month (bio13_LIG_sd),
median temperature of the coldest quarter (bio11_LIG_med), and median temperature of the wettest
quarter (bio8_LIG_med)

Pliocene BIOCLIM: standard deviation of mean precipitation of warmest quarter (bio18_Plio_sd),
median temperature of the wettest quarter (bio8_Plio_med), and median temperature of the coldest
quarter (bio11_Plio_med)
Climate Change: median maximum temperature of the warmest month between the Holocene
and LGM (bio5_Hol_LGM_meddiff), median precipitation of wettest quarter between the present and
Holocene (bio16_curr_Hol_meddif), and median maximum temperature of the warmest month (BIO5)
between the LGM and LIG (bio5_LGM_LIG_meddiff).

This top variable RF model had high predictive power, explaining 56.78% of the variation in MPD. The
top three variables from this analysis were median latitude (lat_med), median world population density
(gpw_med), and current median temperature of the wettest quarter (bio8_med), all showing up as top
five predictors 100% of the time (Fig. 5). Only current median temperature of the wettest quarter was
statistically significant after implementing a Bonferroni correction (0.05/3=0.0166) and accounting for
spatial autocorrelation (Table 2).

Finally, for the 109-eco set reduced to 41 ecoregions after removing PD estimates that were not
significant after randomization, there were 18 top variables in the final RF (Fig. 6) which included:
Current BIOCLIM: median precipitation seasonality (bio15_med) and median temperature of the
wettest quarter (bio8_med)
Geography: standard deviation of the Köpppen-Geiger climate classification (kg2_sd) and
standard deviation of snow cover days (scd_sd)
Holocene BIOCLIM: median temperature of the driest quarter (bio9_Hol_med), median
temperature of the coldest quarter (bio11_Hol_med), median precipitation seasonality
(bio15_Hol_med), and median temperature of the wettest quarter (bio8_Hol_med)
LGM BIOCLIM: median temperature of the coldest quarter (bio11_LGM_med)
LIG BIOCLIM: median temperature of the coldest quarter (bio11_LIG_med), median temperature
of the driest quarter (bio9_LIG_med), and standard deviation of the mean precipitation of the wettest
month (bio13_LIG_sd)
Pliocene BIOCLIM: median temperature of the driest quarter (bio9_Plio_med), median
temperature of the coldest quarter (bio11_Plio_med), and standard deviation of the mean temperature
of the driest quarter (bio9_Plio_sd)
Climate Change: median maximum temperature of the warmest month between the present
and Holocene (bio5_curr_Hol_meddiff), median temperature seasonality between the present and
Holocene (bio4_curr_Hol_meddiff), and median temperature annual range between the present and
Holocene (bio7_curr_Hol_meddiff)

This top variable RF model had similar predictive power to the full 109-eco set, explaining 47.03% of the
variation in MPD. The top three variables from this model were standard deviation of the Köpppen-
Geiger climate classification, Pliocene standard deviation of the mean temperature of the driest quarter
(bio9_Plio_sd), and LIG median temperature of the coldest quarter (bio11_LIG_med). While only
Köpppen-Geiger climate classification showed up as a top five predictor 100% of the time, the other two

were in the top five 82% and 79% of the time. All were significant after implementing a Bonferroni
correction (0.05/3=0.0166) and accounting for spatial autocorrelation (Table 2).
498
499 **Discussion**
500
Terrestrial ecosystems have been rapidly altered by anthropogenic pressures (Ellis et al., 2021). The
degradation and fragmentation of habitats and overexploitation of species globally have culminated in
the current biodiversity crisis. Various conservation initiatives have proposed targets of 30 to 50%
protection of the terrestrial biosphere to mitigate species loss (Dinerstein et al., 2019; Watson & Venter,
2017). Deciphering the environmental and spatial factors that shape different aspects of biodiversity is
imperative to resolve regional and global patterns of diversity and can support conservation practices by
identifying high levels of biodiversity that are less likely to be lost given future climate change. We
leveraged publicly available, single-locus genetic data and spatial layers to estimate OTUs and examine
the distribution of bat phylogenetic diversity (PD) at a global scale and its driving factors using random
forest predictive modeling with a large set of starting predictor variables.
511
We produced a global bat phylogeny and defined OTUs using the Assemble Species by Automatic
Partitioning (ASAP) method. While ASAP genetic groups generally matched current bat taxonomy, we
encountered various groups in the dataset where merging and/or splitting of nominal species was
common. These groups fit into different categories. For example, some merged groups included OTUs
within genera with cryptic species that have undergone recent taxonomic revisions (e.g., Jarrín-V &
Kunz, 2008; Parlos, et al., 2014; Velazco, et al., 2017; Calderón-Acevedo & Muchhala, 2018; Calderón-
Acevedo & Muchhala, 2020; Calahorra-Oliart, et al., 2021; Calderón-Acevedo et al., 2022; Calahorra-
Oliart et al., 2021; da Silva Fonseca et al. 2024; Mônico & Soto-Centeno 2024), particularly small
Stenodermatine frugivores (Rodríguez-Posada, et al., 2018; Morales-Martínez, et al., 2021). Other
mismatched groups included widely distributed insectivores that occur from Africa to Southeast Asia
(Foley et al., 2015; Foley, et al., 2017; Patterson et al., 2019; Taylor et al., 2012; Csorba, et al., 2011).
These examples highlight some of the many recent efforts to provide more stable taxonomy based on
species delimitation (Calahorra-Oliart et al., 2021; da Silva Fonseca et al. 2024; Mônico & Soto-Centeno
2024). For example, the *Vampyressa melissa* species complex was recently split into multiple species
with the use of *Cytochrome b* sequence data (Morales-Martínez et al., 2021). Three of these species,
that are now considered at-risk due to their small ranges and rare occurrences, would have otherwise
gone undetected without the use of genetic data, increasing their risk of extinction. Our ASAP
delimitation results (Table S11) provide a guide for future similar taxonomic work in multiple clades that
might contain cryptic species whose identification could have downstream conservation implications
(Hending, 2024).
532
Patterns in OTU PD match those found in species richness gradients and global phylogenetic diversity
patterns in birds (Udy et al., 2021; Barrios & Martinez-Nuñez, 2024). Southeast Asia is known for its high
levels of bat species richness, and our analysis revealed that it also exhibits some of the highest levels of
PD. Despite having the second highest bat species richness, the Neotropics showed lower phylogenetic
diversity than parts of Africa and the Himalayas, both of which had the next highest levels of PD in our

analysis. This matches previously found patterns in species diversity and other measures of PD in bats
(López-Aguirre, et al., 2018). The fact that the Himalayas had relatively higher levels of PD but lower
species richness, a pattern observed previously, might be particularly important because of this region's
high sensitivity to climate change (Chakravarty, et al., 2021). Furthermore, in birds, the only other group
of flying vertebrates, PD was higher in mountainous regions while being lower in overall species richness
(Jarzyna, et al., 2021; Montaño-Centellas et al., 2023). Similar to the latitudinal gradient of species
diversity, we detected relatively high PD around the tropics compared to temperate areas (Fig. 1 & S4),
and latitude was identified as an important predictor. Eastern South America, central and southern
Africa, central Asia, northern Mexico, and southern Australia have been identified as places to expand
protected areas as they are locations that have intact landscapes and transition potential (Dobrowski et
al. 2021). Some of these regions overlap with the same areas where we detected high levels of PD in
bats and could serve as important hotspots for future biodiversity conservation. The ecoregions we have
identified with high PD using *COI* can be used as a starting point for local conservation efforts that want
to incorporate PD and climate indicators in their decision making. However, it is important to note that
overall sampling was lower in the global south, including large areas of South America and parts of
central Africa, where more of our ecoregions did not have robust estimates of PD (according to SES
resampling). With this caveat, the distribution of PD does not follow the same pattern as sampling (Fig.
1) and temperature variables and bioclimatic history consistently come out as top predictors even when
ecoregions with weak PD estimates were removed from the analysis.
557
At both spatial scales, the 25- and 109-ecoregion sets, median temperature of the wettest quarter was a
significant contributor to PD in bats and was moderately important when using the 41-eco set. Median
temperature of the wettest quarter during the LGM and the change in median temperature of the
warmest month between the LGM and LIG were also significant when looking at larger spatial units (25-
eco set). Both the standard deviation of temperature of the driest quarter during the Pliocene and
median temperature of coldest quarter during the LIG were significant when looking at smaller spatial
units (41-eco set). These results are congruent with similar studies analyzing the relationship between
temperature and various measures of phylogenetic diversity in bats and birds. Phylogenetic diversity
measures like MNTD and MPD have been associated with temperature when looking at both Mexican
bat communities and global bird diversity (Grimshaw & Higgins, 2017; Montaño-Centellas et al., 2023;
Barrios & Martinez-Nuñez, 2024). Warm climates will result in more habitat stability and heterogeneity
allowing for time and space to fill a diverse set of evolutionary niches (Kozak and Wiens 2010). Overall,
temperature is a consistent factor that contributes to phylogenetic diversity. Additionally, the standard
deviation of the Köppen-Geiger climate classification was important and significant. This further
supports that habitat heterogeneity plays an important role in PD and should be considered when
identifying areas for protection (Udy et al. 2021).
574
There are also indications that considering historical components is important for explaining PD
(Montaño-Centellas et al., 2023; Paz et al., 2021; Svenning et al., 2015), and these variables are often
not included in studies of biodiversity. Many of the temperature variables identified as top predictors in
our analyses have a historical component (e.g. Mid-Holocene, LGM, LIG, and Pliocene). This could be key
to understanding patterns of diversity given that climate oscillations during the Quaternary (i.e.,
Pleistocene and Holocene) have been associated with extinction, shaping genetic structure, and

distribution of many small mammals in temperate regions (Blois, et al., 2010; Shafer, et al., 2010).
Changes in climate during this time likely also played a significant role in the historical demography of
bats (Carstens et al., 2018), conceivably contributing to niche evolution. We did detect the change in
temperature from the LIG to LGM as important, potentially suggesting that while climate stability
increases species diversity, change over time might influence PD differently. Braga et al. (2023) found
that climate stability since the LGM increases relatedness in bat communities. It is possible that we did
not identify climate change as a stronger predictor because its impact may vary and could be difficult to
detect at a global scale (Kissling et al., 2021). Despite that, both deeper and recent climate seem to
shape PD. The warmer Pliocene climate was a driver of speciation in African birds (Voelker, et al., 2010).
However, we observed a negative relationship between temperature during the Pliocene and PD,
suggesting that extremely hot and arid climates can limit speciation and niche evolution in bats. This
information can be considered when making future predictions for the stability of species richness in
bats over long timeframes, especially when the goal is to preserve PD.

We showed a slowdown in speciation rates over time in bat families globally. This pattern has been
previously observed in studies sampling all bat families (Shi & Rabosky 2015) and in the superfamily
Noctilionoidea (Rojas et al. 2016). These studies attributed speciation slowdowns to a combination of
macroecological and biogeographic processes with protracted speciation after a diversification event
related to trophic niche availability (Shi & Raboski 2015; Rojas et al. 2016). Generally, bats depict a
saturated expansion scenario where species diversification plateaus towards the present, with
Phyllostomidae showing a group-specific diversification pattern where species accumulation slowdown
follows a more recent rapid rate of expansion. While we used a narrower genetic dataset compared to
other studies, it still captured the signature of diversification scenarios observed across all bats.
Importantly, the association between PD and climate that we estimated likely describes the observed
patterns of bat evolutionary history. In our analysis, the apparent decoupling of PD estimates from the
timing of increased speciation rate, for example in phyllostomids, highlights the importance of lineage
specific factors that account for patterns at the ecoregion level and are independent of bursts in
speciation rate. For example, Neotropical plants show a pattern of more frequent diversity expansions
compared to tetrapods (Meseguer et al., 2022). These increases in diversification rate were stimulated
by climatic cycles and the expansion of different biomes (e.g. Dick & Pennington, 2019). The increased
plant diversification and resulting habitat heterogeneity associated with climate may explain the
maintenance of higher regional PD in other organisms such as bats.

The measure of phylogenetic diversity being used also seems to influence which predictors are
important. In contrast to analyses using MNTD and MPD, studies looking at Faith's index saw
precipitation variables as top predictors (Grimshaw & Higgins, 2017; Paz et al., 2021). We did not
consider Faith's index in our predictive modeling because it was highly correlated with OTU richness in
our dataset, and we did not want our results to be driven by OTU richness. Although, we did detect
some variation in predicting PD at different spatial scales. While the 25-eco worldwide dataset had top
variables that were primarily temperature related, similar to analyses from smaller regions like the
Atlantic Forest of Brazil and Mexico (Grimshaw & Higgins, 2017; Paz et al., 2021), the 109-eco data also
identified median latitude and population density as important, though these were not significant when

accounting for spatial autocorrelation. We believe it could be worthwhile to more systematically explore
if the same patterns apply within and between ecoregions and what differences arise. For example, like
our results, López-Baucells et al. (2022) found that at small spatial scales environmental variables do not
seem to explain PD in aerial insectivorous bats, but PD was higher in larger forest fragments. This could
influence how we think about conservation practices for species at different spatial scales in conjunction
with evolutionary history, as it is not unusual for some patterns, like latitudinal diversity gradients or
human impact, to be more or less strong at different spatial scales (Hillebrand, 2004; Barrios et al.
2024). Furthermore, protected areas do not always maintain within-species genetic diversity
(Thompson, et al., 2021) or fully capture phylogenetic diversity (Saraiva, et al., 2018). Future studies at
local scales should examine the usefulness of protected areas for preserving phylogenetic diversity in
bats (e.g., Leon-Alvarado & Miranda-Esquivel, 2023). Specifically, protected areas could be evaluated on
how much PD they currently have and what the future projections of their climate would look like for
variables that have been deemed important (i.e., temperature).
636
Monitoring biodiversity is essential for conserving current and future levels of biodiversity. In addition to
the thousands of phylogenetic and phylogeographic studies that produce mtDNA sequence data,
environmental DNA metabarcoding and barcoding efforts produce a large amount of sequence data that
can be used for these purposes (Rodríguez-Ezpeleta et al. 2021). When it comes to biodiversity
assessments needed to justify and inform conservation efforts, we have shown that open-source, single-
locus data can be a valuable tool. While our study analyzed environmental and geographic predictors of
bats, this computationally efficient method could be applied to any taxa of interest with sufficient
georeferenced data and used to inform conservation efforts that prioritize the evolutionary history of
species. Both phylogenetic and functional diversity have been found to predict biodiversity effects like
biomass production (Flynn, et al., 2011). Since collecting functional diversity data is more resource
intensive, focusing on PD may be a more efficient and effective way of documenting biodiversity and
ecosystem services that go beyond species richness (Molina-Venegas, et al., 2021; Forest, et al., 2007;
Pio, et al., 2011). PD has also been shown to be connected to the stability of vegetation production,
which is an important factor in assessing an environment's ability to withstand climatic changes
(Mazzochini, et al., 2019).
652
Global bat phylogenetic diversity is highest around the tropics of Asia, Central America, South America,
and Africa which aligns well with patterns of species richness, but assessments at different spatial scales
may provide slightly different results. However, using a broad-scale ecoregion approach, studies of
phylogenetic diversity in other taxa can provide areas of overlap among organisms that could make for
the most valuable targets for conservation. In bats, phylogenetic diversity is best explained by
temperature variables while also considering a historical component. By understanding what
environmental and geographic factors foster higher levels of phylogenetic diversity, we can better
determine what environments contain the diversity necessary to withstand climate change and where
conservation resources will have the largest impact. PD also has more evolutionary potential and can be
useful to consider for conservation purposes long-term. While this information would never be a sole
determinant of conservation focus areas, the analysis is easy to run and can be an additional decision-
making tool when it comes to allocation of conservation resources.

Acknowledgements

Funding provided by National Science Foundation DBI-1911293 to TAP. We thank Radford University's
Office of Undergraduate Research for supporting AEG through the Highlander Research Rookies
program.

References

Amador, L., Arroyo-Torres, I., & Barrow, L. N. (2024). Machine learning and phylogenetic models identify
predictors of genetic variation in Neotropical amphibians. *Journal of Biogeography, 51*(5), 909-
923. https://doi.org/10.1111/jbi.14795
Amador, L., Wiley, D. L., Arroyo-Torres, I., McDaniels, C. X., Rosario-Sanchez, E. O., Farmer, H., … Barrow,
L. N. (2025). Caudata macrogenetics: Species range size predicts intraspecific genetic variation in
global salamanders. *bioRxiv*, 2025-01.
Baker, R. J., Solari, S., Cirranello, A., & Simmons, N. B. (2016). Higher level classification of phyllostomid
bats with a summary of DNA synapomorphies. *Acta Chiropterologica, 18*(1), 1–38.
https://doi.org/10.3161/15081109acc2016.18.1.001
Barrios, P. and Martinez-Nuñez, C. (2024), Bird Phylogenetic Diversity Increases With Temperature
Worldwide. *Diversity and Distribution, 30*, e13930. https://doi.org/10.1111/ddi.13930
Blois, J. L., McGuire, J. L., & Hadly, E. A. (2010). Small mammal diversity loss in response to late-
Pleistocene climatic change. *Nature, 465*(7299), 771-774.
Braga, P. H. P., Kembel, S., & Peres-Neto, P. (2023). Historical and contemporary processes drive global
phylogenetic structure across geographical scales: Insights from bat communities. *Global
Ecology and Biogeography*, 32, 747–765. https://doi.org/10.1111/geb.13650
Brown, J. L., Paz, A., Reginato, M., Renata, C. A., Assis, C., Lyra, M., … Carnaval, A. C. (2020). Seeing the
forest through many trees: Multi-taxon patterns of phylogenetic diversity in the Atlantic Forest
hotspot. *Diversity and Distributions, 26*(9), 1160-1176. https://doi.org/10.1111/ddi.13116
Burgin, C. J., Colella, J. P., Kahn, P. L., & Upham, N. S. (2018). How many species of mammals are there?
Journal of Mammalogy, 99(1), 1–14. https://doi.org/10.1093/jmammal/gyx147
Cadotte, M. W. (2013). Experimental evidence that evolutionarily diverse assemblages result in higher
productivity. *Proceedings of the National Academy of Sciences, 110*(22), 8996-9000.
https://doi.org/10.1073/pnas.1301685110
Cadotte, M. W., Dinnage, R., & Tilman, D. (2012). Phylogenetic diversity promotes ecosystem stability.
Ecology, 93(8), S223-S233. https://doi.org/10.1890/11-0426.1
Calahorra-Oliart, A., Ospina-Garcés, S.M. & León-Paniagua, L. (2021) Cryptic species in *Glossophaga
soricina* (Chiroptera: Phyllostomidae): do morphological data support molecular evidence?
Journal of Mammalogy, 102, 54–68.
Calderón-Acevedo, C. A., Bagley, J. C., & Muchhala, N. (2022). Genome-wide ultraconserved elements
resolve phylogenetic relationships and biogeographic history among Neotropical leaf-nosed bats
in the genus Anoura (Phyllostomidae). *Molecular Phylogenetics and Evolution*, 167, e107356.
https://doi.org/10.1016/j.ympev.2021.107356
Calderón-Acevedo, C.A. & Muchhala, N. (2020) First report of the Broad-toothed Tailless Bat, *Anoura
latidens* Handley, 1984 (Chiroptera, Phyllostomidae), in Bolivia. *Check List*, 16, 1545–1550.

Calderón-Acevedo, C.A. & Muchhala, N.C. (2018) Identification and diagnosis of *Anoura fistulata* with
remarks on its presumed presence in Bolivia. *Journal of Mammalogy*, 99, 131–137.
Calderón-Acevedo, C. A., Rodríguez-Posada, M. E., & Muchhala, N. (2021). Morphology and genetics
concur that *Anoura carishina* is a synonym of *Anoura latidens* (Chiroptera, Glossophaginae).
Mammalia, 85, 471–481. https://doi.org/10.1515/mammalia-2020-0183
Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., … Naeem, S. (2012).
Biodiversity loss and its impact on humanity. *Nature, 486*(7401), 59-67.
https://doi.org/10.1038/nature11148
Carstens, B. C., Morales, A. E., Field, K., & Pelletier, T. A. (2018). A global analysis of bats using
automated comparative phylogeography uncovers a surprising impact of Pleistocene glaciation.
Journal of Biogeography, 45(8), 1795-1805. https://doi.org/10.1111/jbi.13382
Carvalho, W. D., Fluck, I. E., de Castro, I. J., Hilário, R. R., Martins, A. C. M., de Toledo, J. J., da Silva
Xavier, B., Dambros, C., & Bobrowiec, P. E. D. (2023). Elevation drives taxonomic, functional and
phylogenetic β-diversity of phyllostomid bats in the Amazon biome. *Journal of Biogeography*,
50, 70–85. https://doi.org/10.1111/jbi.14533
Carvalho, W. D., Lourenço, E. C., Costa, L. M., Bergallo, H. G., & Esbérard, C. E. L. (2021). Patterns and
drivers determining phyllostomid bat diversity in land-bridge islands off the south-east coast of
Brazil, *Biological Journal of the Linnean Society*, 134(3), 604-619.
https://doi.org/10.1093/biolinnean/blab112
Ceballos, G., Ehrlich, P. R., & Raven, P. H. (2020). Vertebrates on the brink as indicators of biological
annihilation and the sixth mass extinction. *Proceedings of the National Academy of Sciences,
117*(24), 13596-13602. https://doi.org/10.1073/pnas.1922686117
Chakravarty, R., Mohan, R., Voigt, C. C., Krishnan, A., & Radchuk, V. (2021). Functional diversity of
Himalayan bat communities declines at high elevation without the loss of phylogenetic diversity.
Scientific Reports, 11(1), 22556.
CIESIN - Center for International Earth Science Information Network, Columbia University. (2017).
Gridded population of the world, version 4 (GPWv4): Population density, revision 11. *NASA
Socioeconomic Data and Applications Center (SEDAC)*. https://doi.org/10.7927/H49C6VHW
Cirranello, A., Simmons, N. B., Solari, S., & Baker, R. J. (2016). Morphological diagnoses of higher-level
phyllostomid taxa (Chiroptera: Phyllostomidae). *Acta Chiropterologica, 18*(1), 39–71.
https://doi.org/10.3161/15081109acc2016.18.1.002
Csorba, G., Son, N. T., Saveng, I., & Furey, N. M. (2011). Revealing cryptic bat diversity: three new *Murina*
and redescription of *M. tubinaris* from Southeast Asia. *Journal of Mammalogy, 92*(4), 891-904.
da Silva Fonseca, B., Soto-Centeno, J. A., Simmons, N. B., Ditchfield, A. D., & Leite, Y. L. R. (2024). A
species complex in the iconic frog eating bat *Trachops cirrhosus* (Chiroptera, Phyllostomidae)
with high variation in the heart of the Neotropics. *American Museum Novitates, 4021*, 1–27.
Dallaire, C. O., Lehner, B., Sayre, R., & Thieme, M. (2019). A multidisciplinary framework to derive global
river reach classifications at high spatial resolution. *Environmental Research Letters, 14*(2),
024003.
de Carvalho, W. D., Martins, M. A., Esbérard, C. E. L., & Palmeirim, J. M. (2019). Traits that allow bats of
tropical lowland origin to conquer mountains: Bat assemblages along elevational gradients in
the South American Atlantic Forest. *Journal of Biogeography, 46*(2), 316–331.
https://doi.org/10.1111/jbi.13506

Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., … Noss, R. (2017). An ecoregion-based approach to protecting half the terrestrial realm. *BioScience*, *67*(6), 534–545. https://doi.org/10.1093/biosci/bix014

Dinerstein, E., Vynne, C., Sala, E., Joshi, A. R., Fernando, S., Lovejoy, T. E., … Wikramanayake, E. (2019). A global deal for nature: guiding principles, milestones, and targets. *Science advances, 5*(4), eaaw2869.

Dobrowski, S. Z., Littlefield, C. E., Lyons, D. S., Hollenberg, C., Carroll, C. Parks, S. A., Abatzoglou, J. T., Hegewisch, K., & Gage, J. (2021). Protected-area targets could be undermined by climate change-driven shifts in ecoregions and biomes. *Communications Earth & Environment, 2*(198). https://doi.org/10.1038/s43247-021-00270-z

Dolan, A. M., Haywood, A. M., Hunter, S. J., Tindall, J. C., Dowsett, H. J., Hill, D. J., & Pickering, S. J. (2015). Modelling the enigmatic late Pliocene glacial event—Marine Isotope Stage M2. *Global and Planetary Change, 128*, 47-60. https://doi.org/10.1016/j.gloplacha.2015.02.001

Ellis, E. C., Klein Goldewijk, K., Siebert, S., Lightman, D., & Ramankutty, N. (2010). Anthropogenic transformation of the biomes, 1700 to 2000. *Global ecology and biogeography, 19*(5), 589-606. https://doi.org/10.1111/j.1466-8238.2010.00540.x

Ellis, E.C., Gauthier, N., Klein Goldewijk, K., Bliege Bird, R., Boivin, N., Díaz, S., … Watson, J.E.M. (2021). People have shaped most of terrestrial nature for at least 12,000 years. *Proceedings of the National Academy of Sciences, 118*, e2023483118.

European Space Agency (ESA). 2009. ESA GlobCover project. https://due.esrin.esa.int/page_globcover.php

Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological conservation, 61*(1), 1-10. https://doi.org/10.1016/0006-3207(92)91201-3

Fleming, T. H., Dávalos, L. M., & A. R. Mello, M. (2020). *Phyllostomid Bats: A Unique Mammalian Radiation.* University of Chicago Press. https://doi.org/10.7208/chicago/9780226696263.001.0001

Flynn, D. F., Mirotchnick, N., Jain, M., Palmer, M. I., & Naeem, S. (2011). Functional and phylogenetic diversity as predictors of biodiversity–ecosystem-function relationships. *Ecology, 92*(8), 1573-1581. https://doi.org/10.1890/10-1245.1

Foley, N.M., Goodman, S.M., Whelan, C. V., Puechmaille, S.J. & Teeling, E. (2017). Towards navigating the minotaur's labyrinth: Cryptic diversity and taxonomic revision within the speciose genus hipposideros (Hipposideridae). *Acta Chiropterologica, 19*, 1–18. https://doi.org/10.3161/15081109ACC2017.19.1.001

Foley, N.M., Mason, V.C., Harris, A.J., Bredemeyer, K.R., Damas, J., Lewin, H.A., Eizirik, E., Gatesy, J., Karlsson, E.K., Lindblad-Toh, K., Zoonomia Consortium‡, Springer, M.S., Murphy, W.J., Andrews, G., Armstrong, J.C., Bianchi, M., Birren, B.W., Bredemeyer, K.R., Breit, A.M., Christmas, M.J., Clawson, H., Damas, J., Di Palma, F., Diekhans, M., Dong, M.X., Eizirik, E., Fan, K., Fanter, C., Foley, N.M., Forsberg-Nilsson, K., Garcia, C.J., Gatesy, J., Gazal, S., Genereux, D.P., Goodman, L., Grimshaw, J., Halsey, M.K., Harris, A.J., Hickey, G., Hiller, M., Hindle, A.G., Hubley, R.M., Hughes, G.M., Johnson, J., Juan, D., Kaplow, I.M., Karlsson, E.K., Keough, K.C., Kirilenko, B., Koepfli, K.-P., Korstian, J.M., Kowalczyk, A., Kozyrev, S.V., Lawler, A.J., Lawless, C., Lehmann, T., Levesque, D.L., Lewin, H.A., Li, X., Lind, A., Lindblad-Toh, K., Mackay-Smith, A., Marinescu, V.D., Marques-Bonet, T., Mason, V.C., Meadows, J.R.S., Meyer, W.K., Moore, J.E., Moreira, L.R., Moreno-Santillan, D.D., Morrill, K.M., Muntané, G., Murphy, W.J., Navarro, A., Nweeia, M., Ortmann, S., Osmanski,

A., Paten, B., Paulat, N.S., Pfenning, A.R., Phan, B.N., Pollard, K.S., Pratt, H.E., Ray, D.A., Reilly,
S.K., Rosen, J.R., Ruf, I., Ryan, L., Ryder, O.A., Sabeti, P.C., Schäffer, D.E., Serres, A., Shapiro, B.,
Smit, A.F.A., Springer, M., Srinivasan, C., Steiner, C., Storer, J.M., Sullivan, K.A.M., Sullivan, P.F.,
Sundström, E., Supple, M.A., Swofford, R., Talbot, J.-E., Teeling, E., Turner-Maier, J., Valenzuela,
A., Wagner, F., Wallerman, O., Wang, C., Wang, J., Weng, Z., Wilder, A.P., Wirthlin, M.E., Xue,
J.R. & Zhang, X. (2023) A genomic timescale for placental mammal evolution. *Science, 380*,
eabl8189. https://www.science.org/doi/10.1126/science.abl8189

Foley, N.M., Thong, V.D., Soisook, P., Goodman, S.M., Armstrong, K.N., Jacobs, D.S., … Teeling, E.C.
(2015). How and why overcome the impediments to resolution: Lessons from rhinolophid and
hipposiderid bats. *Molecular Biology and Evolution*, *32*, 313–333.
https://doi.org/10.1093/molbev/msu329

Fordham, D. A., Saltré, F., Haythorne, S., Wigley, T. M., Otto-Bliesner, B. L., Chan, K. C., & Brook, B. W.
(2017). PaleoView: a tool for generating continuous climate projections spanning the last 21 000
years at regional and global scales. *Ecography, 40*(11), 1348-1358.
https://doi.org/10.1111/ecog.03031

Forest, F., Grenyer, R., Rouget, M., Davies, T. J., Cowling, R. M., Faith, D. P., … Savolainen, V. (2007).
Preserving the evolutionary potential of floras in biodiversity hotspots. *Nature, 445*, 757-760.
https://doi.org/10.1038/nature05587

Frick, W. F., Kingston, T., & Flanders, J. (2020). A review of the major threats and challenges to global bat
conservation. *Annals of the New York Academy of Sciences, 1469*(1), 5–25.
https://doi.org/10.1111/nyas.14045

Funk S. M., & Fa, J. E. (2010). Ecoregion prioritization suggests an armoury not a silver bullet for
conservation planning. *PLOS ONE 5*(1), e8923. https://doi.org/10.1371/journal.pone.0008923

Genuer, R., Poggi, J. M., & Tuleau, C. (2008). Random Forests: some methodological insights. arXiv
preprint arXiv:0811.3619.

Giakoumi S, Sini M, Gerovasileiou V, Mazor T, Beher J, et al. (2013). Ecoregion-Based Conservation
Planning in the Mediterranean: Dealing with Large-Scale Heterogeneity. *PLOS ONE 8*(10):
e76449. https://doi.org/10.1371/journal.pone.0076449

GIS processing World Bank DECRG. (2010). Washington, DC, extrapolation UNEP/GRID-Geneva.
https://doi.org/10.57966/edbt-r040

Grimm, N. B., Chapin III, F. S., Bierwagen, B., Gonzalez, P., Groffman, P. M., Luo, Y., … Williamson, C. E.
(2013). The impacts of climate change on ecosystem structure and function. *Frontiers in Ecology*
and the Environment, 11(9), 474-482. https://doi.org/10.1890/120282

Grimshaw, J. R., & Higgins, C. L. (2017). Environmental correlates of phylogenetic structure in Mexican
bat communities. *Journal of Mammalogy, 98*(6), 1657-1666.
https://doi.org/10.1093/jmammal/gyx101

Guralnick, R., & Hill, A. (2009). Biodiversity informatics: Automated approaches for documenting global
biodiversity patterns and processes. *Bioinformatics, 25*(4), 421-428.
https://doi.org/10.1093/bioinformatics/btn659

Hanson, J. O., Vincent, J., Schuster, R., Fahrig, L., Brennan, A., Martin, A. E., Hughes, J. S., Pither, R., &
Bennett, J. R. (2022). A comparison of approaches for including connectivity in systematic
conservation planning. *Journal of Applied Ecology, 59*, 2507-2519.
https://doi.org/10.1111/1365-2664.14251

Hao, X., Lu, Q., & Zhao, H. (2024). A molecular phylogeny for all 21 families within Chiroptera (bats).
Integrative Zoology 19, 989–998. https://doi.org/10.1111/1749-4877.12772

Harper, M., Mejbel, H. S., Longert, D., Abell, R., Beard, T. D., Bennett, J. R., … Cooke, S. J. (2021).
Twenty-five essential research questions to inform the protection and restoration of freshwater
biodiversity. *Aquatic Conservation: Marine and Freshwater Ecosystems, 31*(9), 2632-2653.
https://doi.org/10.1002/aqc.3634

Harrell Jr., F. (2023). Hmisc: Harrell Miscellaneous. R package version 5.0-1, https://CRAN.R-
project.org/package=Hmisc

Hending, D. (2024). Cryptic species conservation: a review. *Biological Reviews, 100*(1), 258-274.

Hijmans R (2023). _raster: Geographic Data Analysis and Modeling_. R package version 3.6-14,
https://CRAN.R-project.org/package=raster

Hillebrand, H. (2004). On the Generality of the Latitudinal Diversity Gradient. T*he American Naturalist,*
163(2), 192-211. https://doi.org/10.1086/381004

Hosegood, J., Humble, E., Ogden, R., De Bruyn, M., Creer, S., Stevens, G. M., … Carvalho, G. (2020).
Phylogenomics and species delimitation for effective conservation of manta and devil rays.
Molecular ecology, 29(24), 4783-4796. https://doi.org/10.1111/mec.15683

Hu, Y., Fan, H., Chen, Y., Chang, J., Zhan, X., Wu, H., … Wei, F. (2021). Spatial patterns and conservation
of genetic and phylogenetic diversity of wildlife in China. *Science Advances, 7*(4), eabd5725.
https://doi.org10.1126/sciadv.abd5725

Jarrín-V., P. & Kunz, T.H. (2008). Taxonomic history of the genus Anoura (Chiroptera: Phyllostomidae)
with insights into the challenges of morphological species delimitation. *Acta Chiropterologica,*
10, 257–269.

Jarzyna, M. A., Quintero, I., & Jetz, W. (2021). Global functional and phylogenetic structure of avian
assemblages across elevation and latitude. *Ecology Letters, 24*(2), 196-207.
https://doi.org/10.1111/ele.13631

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. (2017). ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589.
https://doi.org/10.1038/nmeth.4285

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., … Kessler, M. (2017).
Climatologies at high resolution for the Earth land surface areas. *Scientific Data, 4.* 170122.
https://doi.org/10.1038/sdata.2017.122

Karger D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., … Kessler, M. (2018). Data
from: Climatologies at high resolution for the earth's land surface areas. *Dryad Digital*
Repository. http://dx.doi.org/doi:10.5061/dryad.kd1d4

Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H., & Zimmermann, N. E. (2021). CHELSA-TraCE21k
v1. 0. Downscaled transient temperature and precipitation data since the last glacial maximum.
Climate of the Past Discussions, 1-27. https://doi.org/10.5194/cp-19-439-2023

Katoh, K. & Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and evolution, 30*, 772–780.
https://doi.org/10.1093/molbev/mst010

Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., … Webb, C. O.
(2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics, 26*(11), 1463-
1464. https://doi.org/10.1093/bioinformatics/btq166

Kissling, W. D., Eiserhardt, W. L., Baker, W.J., Borchsenius, F., Couvreur, T. L. P., Balslev, H. & Svenning, J.
(2012). Cenozoic imprints on the phylogenetic structure of palm species assemblages
worldwide, Proceedings of the National Academy of Sciences, 109(19), 7379-7384.
https://doi.org/10.1073/pnas.1120467109
Kozak, K. H., & Wiens, J. J. (2010). Accelerated rates of climate-niche evolution underlie rapid species
diversification. Ecology Letters, 13, 1378-1389. https://doi.org/10.1111/j.1461-
0248.2010.01530.x
Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. (2019). RAxML-NG: a fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35(21)**, 4453–
4455. https://doi.org/10.1093/bioinformatics/btz305
Kunz, T.H., Braun de Torrez, E., Bauer, D., Lobova, T. & Fleming, T.H. (2011). Ecosystem services provided
by bats. *Annals of the New York Academy of Sciences, 1223*, 1–38.
https://doi.org/10.1111/j.1749-6632.2011.06004.x
Lehner, B., & Döll, P. (2004). Development and validation of a global database of lakes, reservoirs and
wetlands. *Journal of hydrology, 296*(1-4), 1-22. https://doi.org/10.1016/j.jhydrol.2004.03.028
Leon-Alvarado, O.D. & Miranda-Esquivel, D.R. (2023) Phylogenetic diversity and North Andean block
conservation. *PeerJ, 11*, e16565.
Li, D., L. Trotta, H. E. Marx, J. M. Allen, M. Sun, D. E. Soltis, P. S. Soltis, R. P. Guralnick, & B. Baiser. (2019). For
common community phylogenetic analyses, go ahead and use synthesis phylogenies. *Ecology*
100(9):e02788. https://doi.org/10.1002/ecy.2788
Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News, 2*(3), 18-22.
https://CRAN.R-project.org/doc/Rnews/
Lino, A., Ferreira, E., Fonseca, C., Fischer, E. & Ramos Pereira, M.J. (2021). Species–genetic diversity
correlation in phyllostomid bats of the Bodoquena plateau, Brazil. *Biodiversity and Conservation*,
30, 403–429. https://doi.org/10.1007/s10531-020-02097-0
López-Aguirre, C., Hand, S. J., Laffan, S. W., & Archer, M. (2018). Phylogenetic diversity, types of
endemism and the evolutionary history of New World bats. *Ecography, 41*(12), 1955-1966.
https://doi.org/10.1111/ecog.03260
López-Baucells, A., Rowley, S., Rocha, R., Bobrowiec, P. E., Palmeirim, J. M., Farneda, F. Z., & Meyer, C. F.
(2022). Interplay between local and landscape-scale effects on the taxonomic, functional and
phylogenetic diversity of aerial insectivorous neotropical bats. *Landscape Ecology, 37*(11), 2861-
2875. https://doi.org/10.1007/s10980-022-01493-x
Mammal Diversity Database. (2024). Mammal Diversity Database (Version 1.11) [Data set]. Zenodo.
https://doi.org/https://doi.org/10.5281/zenodo.7830771
Mason-Romo, E. D., Farías, A. A., Ceballos, G. (2017) Two decades of climate driving the dynamics of
functional and taxonomic diversity of a tropical small mammal community in western Mexico.
PLOS ONE 12(12),e0189104. https://doi.org/10.1371/journal.pone.0189104
Mazzochini, G. G., Fonseca, C. R., Costa, G. C., Santos, R. M., Oliveira-Filho, A. T., & Ganade, G. (2019).
Plant phylogenetic diversity stabilizes large-scale ecosystem productivity. *Global Ecology and
Biogeography, 28*(10), 1430-1439. https://doi.org/10.1111/geb.12963
Molina-Venegas, R., Rodríguez, M. Á., Pardo-de-Santayana, M., Ronquillo, C., & Mabberley, D. J. (2021).
Maximum levels of global phylogenetic diversity efficiently capture plant services for
humankind. *Nature Ecology & Evolution, 5*(5), 583-588.

Mônico, P. I. & Soto-Centeno, J. A. (2024). Phylogenetic, morphological and niche differentiation unveil
new species limits for the big brown bat (*Eptesicus fuscus*). *Royal Society Open Science*, *11* (2),
231384.
Montaño-Centellas, F., Baiser, B., McGrew, A., Trotta, L., & Li, D. (2023). Global patterns and drivers of
raptor phylogenetic and functional diversity. Global Ecology and Biogeography, 32(2), 281-294.
https://doi.org/10.1111/geb.13619
Morales-Martínez, D.M., Rodríguez-Posada, M.E. & Ramírez-Chaves, H.E. (2021). A new cryptic species
of yellow-eared bat *Vampyressa melissa* species complex (Chiroptera: Phyllostomidae) from
Colombia. *Journal of Mammalogy, 102*, 90–100.
Moreira-Hernández, J.I., Calderón-Acevedo, C.A. & Muchhala, N. (2021). Fur, wings, and flowers:
Development and progress on nectarivorous bat research in the last 50 years. 50 Years of Bat
Research. In B.K. Lim, M.B. Fenton, R.M. Brigham, S. Mistry, A. Kurta, E.H. Gillam, A. Russell, and
J. Ortega, & (Eds.), pp. 135–149. Springer International Publishing, Cham.
https://doi.org/10.1007/978-3-030-54727-1_9
Muchhala, N., & Tschapka, M. (2020). The Ecology and Evolution of Nectar Feeders. In T. H. Fleming,
Dávalos, & M. A. R. Mello (Ed.), *Phyllostomid Bats* (pp. 273–294). University of Chicago Press.
https://doi.org/10.7208/chicago/9780226696263.003.0016
Mulcahy, D. G., Ibáñez, R., Jaramillo, C. A., Crawford, A. J., Ray, J. M., Gotte, S. W., ... de Queiroz, K.
(2022). DNA barcoding of the National Museum of Natural History reptile tissue holdings raises
concerns about the use of natural history collections and the responsibilities of scientists in the
molecular age. *Plos one, 17*(3), e0264930. https://doi.org/10.1371/journal.pone.0264930
Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., ... Purvis, A. (2015). Global
effects of land use on local terrestrial biodiversity. *Nature, 520*(7545), 45-50.
https://doi.org/10.1038/nature14324
Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., ...
Kassem, K. R. (2001). Terrestrial ecoregions of the world: A new map of life on Earth: A new
global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity.
BioScience, *51*(11), 933–938. https://doi.org/10.1641/0006-
3568(2001)051[0933:TEOTWA]2.0.CO;2
Otto-Bliesner, B. L., Marshall, S. J., Overpeck, J. T., Miller, G. H., Hu, A., & CAPE Last Interglacial Project
members. (2006). Simulating arctic climate warmth and icefield retreat in the last
interglaciation. *Science, 311*(5768), 1751-1753.
Paradis, E. & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*, 35, 526-528. doi:10.1093/bioinformatics/bty633
Parlos, J.A., Timm, R.M., Swier, V.J., Zeballos, H. & Baker, R.J. (2014) Evaluation of the paraphyletic
assemblages within Lonchophyllinae, with description of a new tribe and genus. *Occasional
Papers, Museum of Texas Tech University, 320*, 1–23.
Parsons, D. J., Pelletier, T. A., Wieringa, J. G., Duckett, D. J., & Carstens, B. C. (2022). Analysis of
biodiversity data suggests that mammal species are hidden in predictable places. *Proceedings of
the National Academy of Sciences, 119*(14), e2103400119.
https://doi.org/10.1073/pnas.2103400119
Patterson, B.D., Webala, P.W., Kerbis Peterhans, J.C., Goodman, S.M., Bartonjo, M. & Demos, T.C.
(2019). Genetic variation and relationships among Afrotropical species of *Myotis* (Chiroptera:
Vespertilionidae). *Journal of Mammalogy, 100,* 1130–1143.

Paz, A., Brown, J.L., Cordeiro, C.L.O., Aguirre-Santoro, J., Assis, C., Amaro, R.C., … Carnaval, A.C. (2021). Environmental correlates of taxonomic and phylogenetic diversity in the Atlantic Forest. *Journal of Biogeography, 48*, 1377–1391. https://doi.org/10.1111/jbi.14083

Paz-Vinas, I., Jensen, E. L., Bertola, L. D., Breed, M. F., Hand, B. K., Hunter, M. E., … & Hoban, S. (2021). Macrogenetic studies must not ignore limitations of genetic markers and scale. *Ecology Letters, 24*(6), 1282-1284. https://doi.org/10.1111/ele.13732

Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal 10*(1), 439-446. https://doi.org/10.32614/RJ-2018-009

Peduzzi, P. (2019). The disaster risk, global change, and sustainability nexus. *Sustainability, 11*(4), 957. https://doi.org/10.3390/su11040957

Pelletier, T. A., & Carstens, B. C. (2018). Geographical range size and latitude predict population genetic structure in a global survey. *Biology Letters, 14*(1), 20170566. https://doi.org/10.1098/rsbl.2017.0566

Pelletier, T. A., Carstens, B. C., Tank, D. C., Sullivan, J., & Espíndola, A. (2018). Predicting plant conservation priorities on a global scale. Proceedings of the National Academy of Sciences, 115(51), 13027-13032. https://doi.org/10.1073/pnas.1804098115

Pelletier, T. A., Parsons, D. J., Decker, S. K., Crouch, S., Franz, E., Ohrstrom, J., & Carstens, B. C. (2022). phylogatR: Phylogeographic data aggregation and repurposing. *Molecular Ecology Resources*, *22*(8), 2830– 2842. https://doi.org/10.1111/1755-0998.13673

Pereira, H. M. (2016). A latitudinal gradient for genetic diversity. *Science, 353*(6307), 1494-1495. https://doi/10.1126/science.aah6730

Pinheiro, J.C. & Bates, D.M. (2000). Mixed-Effects Models in S and S-PLUS. *Springer*, *New York*. doi:10.1007/b98882 https://doi.org/10.1007/b98882.

Pinheiro, J., Bates, D., & R Core Team (2025). nlme: Linear and Nonlinear Mixed Effects Models. doi:10.32614/CRAN.package.nlme https://doi.org/10.32614/CRAN.package.nlme, R package version 3.1-168, https://CRAN.R-project.org/package=nlme.

Pio, D. V., Broennimann, O., Barraclough, T. G., Reeves, G., Rebelo, A. G., Thuiller, W., … Salamin, N. (2011). Spatial predictions of phylogenetic diversity in conservation decision making. *Conservation Biology, 25*(6), 1229-1239.

Pons, J., Barraclough, T. G., Gomez-Zurita, J., Cardoso, A., Duran, D. P., Hazell, S., … Vogler, A. P. (2006). Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology, 55*(4), 595–609. https://doi.org/10.1080/10635150600852011

Puillandre, N., Brouillet, S., & Achaz, G. (2020). ASAP: assemble species by automatic partitioning. *Molecular Ecology Resources, 21*(2), 609– 620. https://doi.org/10.1111/1755-0998.13281

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Ramírez-Fráncel, L.A., García-Herrera, L.V., Losada-Prado, S., Reinoso-Flórez, G., SÁánchez-Hernández, A., Estrada-Villegas, S., … Guevara, G. (2022). Bats and their vital ecosystem services: a global review. *Integrative Zoology, 17*, 2–23. https://doi.org/10.1111/1749-4877.12552

Reeder, D. M., Helgen, K. M., & Wilson, D. E. (2007). Global trends and biases in new mammal species discoveries. Occasional Papers, Museum of Texas Tech University.

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). *PeerJ*, 12, e16505.

Rodríguez-Ezpeleta, N., Zinger, L., Kinziger, A., Bik, H.M., Bonin, A., Coissac, E., Emerson, B.C., Lopes,
C.M., Pelletier, T.A., Taberlet, P. and Narum, S. (2021), Biodiversity monitoring using
environmental DNA. *Molecular Ecology Resources, 21*, 1405-1409.
https://doi.org/10.1111/1755-0998.13399
Rodríguez-Posada, M.E., Morales-Martínez, D.M., Gutiérrez-Sanabria, D.R., Calderón-Capote, M.C. &
Fernández-Rodríguez, C. (2018). New records of the rare bat *Vampyressa melissa* Thomas, 1926
(Chiroptera: Phyllostomidae: Stenodermatinae) from the eastern slope of the Andes in the
Colombian Orinoco basin. *Mammalia, 82*, 298–302.
Rodríguez-Posada, M. E., Morales-Martínez, D. M., Ramírez-Chaves, H. E., Martínez-Medina, D., &
Calderón-Acevedo, C. A. (2021). A new species of Long-eared Brown Bat of the genus *Histiotus*
(Chiroptera) and the revalidation of *Histiotus colombiae*. *Caldasia*, *43*(2), 221–234.
https://doi.org/10.15446/caldasia.v43n2.85424
Rojas, D., Warsi, O. M. & Dávalos, L. M. Bats (Chiroptera: Noctilionoidea) challenge a recent origin of
extant neotropical diversity. (2016). *Syst. Biol. 65*, 432–448.
https://doi.org/10.1093/sysbio/syw011
Rosauer, D. F., Byrne, M., Blom, M. P., Coates, D. J., Donnellan, S., Doughty, P., … Moritz, C. (2018).
Real-world conservation planning for evolutionary diversity in the Kimberley, Australia,
sidesteps uncertain taxonomy. *Conservation Letters, 11*(4), e12438.
https://doi.org/10.1111/conl.12438
Rosauer, D. F., Pollock, L. J., Linke, S., & Jetz, W. (2017). Phylogenetically informed spatial planning is
required to conserve the mammalian tree of life. *Proceedings of the Royal Society B: Biological*
Sciences, 284(1865), 20170627. https://doi.org/10.1098/rspb.2017.0627
Saraiva, D. D., dos Santos, A. S., Overbeck, G. E., Giehl, E. L. H., & Jarenkow, J. A. (2018). How effective
are protected areas in conserving tree taxonomic and phylogenetic diversity in subtropical
Brazilian Atlantic Forests? *Journal for Nature Conservation, 42*, 28-35.
https://doi.org/10.1016/j.jnc.2018.02.002
Shafer, A. B., Cullingham, C. I., Côté, S. D., & Coltman, D. W. (2010). Of glaciers and refugia: a decade of
study sheds new light on the phylogeography of northwestern North America. *Molecular*
ecology, 19(21), 4589-4621.
Shi, J. J., & D. L. Rabosky. (2015). Speciation dynamics during the global radiation of extant bats.
Evolution, *69*:1528–1545.
Simmons, N. B., & Cirranello, A. L. (2025). *Batnames.org Species List Version 1.8.1*. Accessed on
07/15/2025. https://zenodo.org/records/16937886
Smith, B. T., McCormack, J. E., Cuervo, A. M., Hickerson, M. J., Aleixo, A., Cadena, C. D., … Brumfield, R.
T. (2014). The drivers of tropical speciation. Nature, 515(7527), 406-409.
Smith, B. T., Seeholzer, G. F., Harvey, M. G., Cuervo, A. M., & Brumfield, R. T. (2017). A latitudinal
phylogeographic diversity gradient in birds. *PLoS biology, 15*(4), e2001073.
https://doi.org/10.1371/journal.pbio.2001073
Smith, S.A. & O'Meara, B.C. (2012). treePL: divergence time estimation using penalized likelihood for
large phylogenies. *Bioinformatics*, **28**, 2689–2690.
Solari, S., & Martínez-Arias, V. (2014). Cambios recientes en la sistemática y taxonomía de murciélagos
Neotropicales (Mammalia: Chiroptera). *Therya*, *5*(1), 167–196. https://doi.org/10.12933/therya-
14-180

Soto-Centeno, J. A., & Calderón-Acevedo, C. A. (2023). Global change and the conservation of Caribbean
1054   bat communities. In A. Kurta & A. Rodríguez-Durán (Eds.), *Bats of the West Indies, A Natural*
1055   *History and Field Guide* (pp. 72–86). Cornell University Press.

1056   Stevens, R. D., & Gavilanez, M. M. (2015). Dimensionality of community structure: phylogenetic,
1057   morphological and functional perspectives along biodiversity and environmental gradients.
1058   *Ecography, 38*(9), 861-875. https://doi.org/10.1111/ecog.00847

1059   Svenning, J. C., Eiserhardt, W. L., Normand, S., Ordonez, A, & Sandel, B. (2015). The influence of
1060   paleoclimate on present-day patterns in biodiversity and ecosystems. *Annual Review of Ecology,*
1061   *Evolution, and Systematics, 46*, 551-572. https://doi.org/10.1146/annurev-ecolsys-112414-
1062   054314

1063   Tachikawa, T., Kaku, M., Iwasaki, A., Gesch, D. B., Oimoen, M. J., Zhang, Z., … Carabajal, C. (2011). ASTER
1064   global digital elevation model version 2-summary of validation results. *NASA*.

1065   Taylor, P. J., Stoffberg, S., Monadjem, A., Schoeman, M. C., Bayliss, J., & Cotterill, F. P. (2012). Four new
1066   bat species (*Rhinolophus hildebrandtii* complex) reflect Plio-Pleistocene divergence of dwarfs
1067   and giants across an Afromontane archipelago. *Plos One, 7*(9): e41744.

1068   Teeling, E. C., Springer, M. S., Madsen, O., Bates, P., O'Brien, S. J., & Murphy, M. J. (2005). A molecular
1069   phylogeny for bats illuminates biogeography and the fossil record. *Science, 307*, 580–584.
1070   https://doi.org/10.1126/science.1105113

1071   Theissinger, K., Fernandes, C., Formenti, G., Bista, I., Berg, P. R., Bleidorn, C., … Zammit, G. (2023). How
1072   genomics can help biodiversity conservation. *Trends in Genetics, 39*(7), 545-559.
1073   https://doi.org/10.1016/j.tig.2023.01.005

1074   Thompson, C. E., Pelletier, T. A., & Carstens, B. C. (2021). Genetic diversity of North American
1075   vertebrates in protected areas. *Biological Journal of the Linnean Society, 132*(2), 388-399.
1076   https://doi.org/10.1093/biolinnean/blaa195

1077   Tuanmu, M. N., & Jetz, W. (2015). A global, remote sensing-based characterization of terrestrial habitat
1078   heterogeneity for biodiversity and ecosystem modelling. *Global Ecology and Biogeography,*
1079   *24*(11), 1329-1339. https://doi.org/10.1111/geb.12365

1080   Udy K, Fritsch M, Meyer KM, Grass, I., Hanß, S., Hartig, F., Kneib, T., Kreft, H., Kukunda, C.B., Pe'er, G.,
1081   Reininghaus, H., Tietjen, B., Tscharntke, T., van Waveren, C-S., & Wiegand, K. (2021).
1082   Environmental heterogeneity predicts global species richness patterns better than area. *Global*
1083   *Ecology and Biogeography, 30*, 842–851. https://doi.org/10.1111/geb.13261

1084   Velazco, P.M., Soto-Centeno, J.A., Fleck, D.W., Voss, R.S. & Simmons, N.B. (2017). A new species of
1085   nectar-feeding bat of the genus *Hsunycteris*. *American Museum Novitates*, 1–26.

1086   Voelker, G., Outlaw, R. K., & Bowie, R. C. (2010). Pliocene forest dynamics as a primary driver of African
1087   bird speciation. *Global Ecology and Biogeography, 19*(1), 111-121.
1088   https://doi.org/10.1111/j.1466-8238.2009.00500.x

1089   Warwick, R. M., & Clarke, K. R. (1995). New 'biodiversity' measures reveal a decrease in taxonomic
1090   distinctness with increasing stress. *Marine ecology progress series, 129*, 301-305.
1091   https://doi.org/10.3354/meps129301

1092   Watson, J. E., & Venter, O. (2017). A global plan for nature conservation. *Nature, 550*(7674), 48-49.
1093   https://doi.org/10.1038/nature24144

1094   Webb, C. O. (2000). Exploring the phylogenetic structure of ecological communities: an example for rain
1095   forest trees. *The American Naturalist, 156*(2), 145-155. https://doi.org/10.1086/303378

Webb, C. O., Ackerly, D. D., McPeek, M. A., & Donoghue, M. J. (2002). Phylogenies and community
ecology. *Annual review of ecology and systematics, 33*(1), 475-505.
https://doi.org/10.1146/annurev.ecolsys.33.010802.150448
Weiskopf, S. R., Rubenstein, M. A., Crozier, L. G., Gaichas, S., Griffis, R., Halofsky, J. E., … Whyte, K. P.
(2020). Climate change effects on biodiversity, ecosystems, ecosystem services, and natural
resource management in the United States. *Science of the Total Environment, 733*, 137782.
https://doi.org/10.1016/j.scitotenv.2020.137782
Yi, X., Latch, E. K., Lim, B. K., Koroiva, R., Da Rocha, P. A., & Feijó, A. (2024). UCE-derived mitochondrial
phylogeny reveals pervasive mito-nuclear discordances in serotine bats (genus *Eptesicus*) and
complex evolutionary history in Eptesicus (Histiotus). *Mammalian Biology*, *104*(4), 417–430.
https://doi.org/10.1007/s42991-024-00418-2
Zhang, J., Kapli, P., Pavlidis, P., & Stamatakis, A. (2013). A general species delimitation method with
applications to phylogenetic placements. *Bioinformatics, 29*(22), 2869-2876.
https://doi.org/10.1093/bioinformatics/btt499
Zoderer, B. M., Thomas Marsoner, T., & Tasser, E. (2024). Protecting wilderness or rewilding? An
ecoregion-based approach to identifying priority areas for the protection and restoration of
natural processes for biodiversity conservation. *Journal for Nature Conservation, 81*, 126661.
https://doi.org/10.1016/j.jnc.2024.126661

1114
1115
1116

1117 **Data availability statement**

1118 All intermediate datafiles and list of DNA sequence accessions are available as supplemental tables
1119 (Table S1-8). All data analysis scripts and supplemental files 1-12 are on Dryad: Reviewer URL:
1120 http://datadryad.org/stash/share/Wrp3Bem5LbdGpxC8bdzp8h0f3eyMJW0zoc6hNW11y8A.

1121
1122

Table 1. Percent variation explained for all random forest models and the number of predictor variables
used in each model. 0.75 cut-off refers to the variable set where those correlated higher than 0.75 were
removed. 0.05 cut-off refers to the set where ecoregions were removed if their PD values were not
significant after randomization. BIOCLIM variables for present day (Current), the Holocene, the Last
Glacial Maxima (LGM), the Last Interglacial (LIG), and the Pliocene (Plio) were grouped into their own
categories. The final two categories were geographic variables (Geography), which also contained other
data layers, and Climate change which included differences between each of time periods for all
BIOCLIM variables.

Group	MPD % Var		MNTD % Var		# variables	
	25-eco	109-eco	25-eco	109-eco	25-eco	109-eco
Current bioclim	56.84	40.28	4.09	7.81	38	38
Geography	67.81	57.35	-2.98	19.92	47	47
Holocene	52.72	38.51	-4.21	5.29	38	38
Last glacial maximum	60.41	37.58	11.18	3.92	38	38

Last interglacial	47.78	37.35	1.72	4.85	38	38
Pliocene	50.51	36.55	-8.85	3.42	28	28
Climate change	65.26	42.19	-6.63	14.72	71	71
0.75 cut-off						
Current bioclim	52.55	42.98	0.85	1.35	12	12
Geography	53.16	58.39	-3.94	18.99	29	28
Holocene	49.94	39.08	-3.41	6.49	27	24
Last glacial maximum	56.13	38.24	11.07	3.25	28	27
Last interglacial	48.89	33.54	-2.38	5.91	29	26
Pliocene	47.41	35.73	-10.22	2.27	24	24
Climate change	65.26	42.11	-6.31	14.59	71	71
0.75 cut-off & 0.05 cut-off						
Current bioclim	-	21.13	-	-4.94	-	12
Geography	-	24.03	-	-6.56	-	28
Holocene	-	24.98	-	11.12	-	24
Last glacial maximum	-	24.78	-	-12.40	-	27
Last interglacial	-	23.26	-	-2.48	-	26
Pliocene	-	19.20	-	0.94	-	24
Climate change	-	16.29	-	-6.80	-	71

1131

1132

1133

1134

1135   **Table 2**. P-values for the top three variables in each of the three RF final models.

Variable	eco-set	p-value
median LGM BIO8	25	<0.0001
median present BIO8	25	<0.0001
median LGM-LIG BIO5	25	<0.0001
absolute latitude	109	0.2654
median gpw	109	0.9779
median BIO8	109	<0.0001
standard deviation kg2	41	0.0142
standard deviation Pliocene BIO9	41	0.0138
median LIG BIO11	41	<0.0001

1136

1137   **Figures and figure legends**

1138

1139
1140 **Figure 1**: 109-ecoregions. (a) Phylogenetic diversity of bats Worldwide and number of DNA sequences
1141   (given in natural logarithm of actual sequence numbers in a 2x2 degree grid, for better
1142   visualization). (b) The Neotropics (c) East Africa (d) Southeast Asia
1143

1144
1145
Figure 2: Time calibrated phylogeny of bats worldwide representing 17 families and all extant
superfamilies. Estimates of divergence were performed in TreePL using the RaxML-NG COI
phylogeny. Color coded nodes represent the superfamily and family taxonomic designations at
tips of the phylogeny. Time scale represented in millions of years and following the geological
time designations Pal= Paleogene, Eo = Eocene, Ol = Oligocene, Mio = Miocene, and Pli =
Pliocene (the Pleistocene and Holocene not labeled).
1152

1153
1154
1155   **Figure 3**: Lineage through time (LTT) plot based on the dated maximum likelihood RaxML-NG COI tree
1156   showing lineage accumulation in the last 30 million years for seven bat families. Plot scaled
1157   using the family Phyllostomidae (New World leaf-nosed bats, black line) as reference.
1158

1159

Figure 4. (a) Bar plot of % increase in MSE for all variables in the top random forest for the 25-ecoregion
dataset. Red bars indicate those variables that were in the top five predictors 100% of the time
for 100 random forest iterations. (b) Scatterplots with linear regression lines for the three top
variables. All three were significant.

1165
1166 **Figure 5.** (a) Bar plot of % increase in MSE for all variables in the top random forest for the 109-
1167   ecoregion dataset. Red bars indicate those variables that were in the top five predictors 100% of
1168   the time for 100 random forest iterations. (b) Scatterplots with linear regression lines for the
1169   three top variables. Only median temperature of the wettest quarter was significant.
1170

1171
1172 **Figure 6.** (a) Bar plot of % increase in MSE for all variables in the top random forest for the ecoregion
1173 dataset with only significant mpd values. Red bars indicate those variables that were in the top
1174 five predictors 100% of the time for 100 random forest iterations, and orange bars indicate
1175 those that were in the top five 75% or more of the time for 100 random forest iterations. (b)
1176 Scatterplots with linear regression lines for the three top variables. All three were significant.