Tracing the function expansion for a primordial protein fold in the era of fold-

based function prediction: beta-trefoil

Moushmi Goswami[1] and Subhashini Srinivasan[1, 2]

1. Institute of Bioinformatics and Applied Biotechnology, Biotech Park, Electronic City Phase I,

Bangalore 560100, India

2. Corresponding author

ABSTRACT

The incredibly narrow protein fold bottleneck, which separates the billions of unique

proteins on one side to deliver diverse biological functions on the other, arises from folds tolerating

mutations during evolution. One such fold, called beta-trefoil, is present in functionally diverse

proteins including cytokines involved in the immune system. The unrecognizable sequence-level

diversity even among paralogs of interleukin-1, one of the cytokines within the same chromosomal

locus, suggests the resilience of this fold to mutational on-slaught. Furthermore, beta-trefoil

containing-proteins are known to exist along with other domains allowing them to bind to diverse

macromolecular types in achieving the observed functional diversity. In this study, we use the

beta-trefoil, a fold prevalent across kingdoms, to assess the reach and limitations of function

prediction using fold-fold comparison. We found thirty-two distinct functions for proteins

containing at least one trefoil domain by mining both the PDB and AlphaFold databases. Among

the novel domain architecture including trefoil domain, we find domains such as chitinase, lipase,

beta-glucosidase, protein kinase, peptidoglycan-binding + peptidase matrixin, glycosyl hydrolases

family 3 + PA14 + fibronectin type- III, alpha galactosidase A, PhoD-like phosphatase, insecticidal

crystal toxin, trypsin, alginate lyase and two unknown domains. The limitations of structure-

structure comparison tools may arise from converging evolution in folds. This was addressed by

creating phylogenetic tree from structure-based multiple sequence alignment of trefoil domains

from diverse kingdoms and functions. The clades by kingdoms as opposed to functions may

support the hypothesis that beta-trefoil evolved independently, perhaps creating a twilight zone in

function prediction using fold-fold comparison.

INTRODUCTION

Nobel Laureate Francois Jacob, in his essay on Evolution and Tinkering, stated that nature

is a better tinkerer than an inventor. As evolution proceeds to create complex organisms from a

small repertoire of protein folds, one can expect accumulating diversity in the amino acid sequences of proteins sharing the same fold. Considering that protein functions are dictated by their unique three-dimensional structures, the accepted mutations during evolution are expected to preserve the overall structural fold of the original protein. While, in some functional families, one can recognize the remnant sequence motifs of the ancient protein, there are protein fold-families with no recognizable sequence signatures, yet sharing folds and, perhaps, function.

In 1992, Cyrus Homi Chothia estimated the number of unique folds to be as low as one thousand[3]. As of 2023, the protein data bank (PDB) has 3D structures of 227,933 proteins with ~7000 clusters less than 30% identity representing only 1,257 unique folds[1]. The five-fold reduction in the number of folds from sequence-based clustering reflects both i) the diversity in sequences within a family below the 30% identity cutoff, and ii) the differing domain architecture within a protein family limiting homology over good lengths of the proteins. More recently, the number of sequence clusters in the BFD (big fantastic database), using a collation of 2 billion proteins, is as large as 65,983,866 using a 30% identity cutoff and 90% coverage over the length of proteins[2].

The exponential lag in solving 3D structures of proteins and, until recently, our inability to predict structures of divergent and novel proteins from primary sequences, made sequence homology based-function prediction a norm for the last several decades. We have come a long way since the efforts by Dayhoff in collating thousands of published protein sequences[3]. The development of tools for sequence homology-based function prediction kept pace with growing sequence databases, bridging the lag in structure determination. For example, the early 1990's saw a boom in DNA sequence databases from EST-sequencing efforts, triggering an unprecedented race to discover novel paralogs of genes of therapeutic importance. Function prediction from

fragmented DNAs was the need of the time, which was met wholesomely by the timely

development of the most popular molecular biology tool in 1990 called BLAST[4]. To improve the

sensitivity in sequence-based structure prediction, among divergent paralogs/homologs, hidden

markov models (HMM) for individual fold families were developed[5].

Theoretically one should be able to predict functions of divergent proteins using the shared

folds. Parallel efforts continued to predict structures of proteins based on homology at the

sequence-level. However, the progression of homology-based structure prediction can be likened

to decades where nothing happened. Slowly but surely, novel strategies including the incorporation

of multiple templates, the refinement of secondary structure prediction, the optimization of *ab

initio* loop modeling, and the use of backbone-dependent rotamer libraries for side-chain

placement continued to improve structure prediction methods over decades[6]. That the native fold

is present among the thousands of simulated structures kept the lights-on for researchers through

community-based challenge[7]. Wodak et al. has provided a detailed review of progress in

methodologies achieved via collective intelligence through community experiments like CASP

(critical assessment of structure prediction) and other efforts[8]. Significant improvements in

methods were observed by CASP10 by introducing co-evolution to guide the folding process[9].

Parallel to this effort, the deluge in protein sequences from advances in next generation sequencing

revealed more and more coevolving pairs within folding domains adding additional distance

constraints between distal amino acids in the primary sequence that are likely to be spatially

proximal — advancing structure prediction by leaps and bounds. By 2020, the convergence of

collective and artificial intelligence along with advances in high-throughput sequencing

technologies, we entered an era when decades happen in weeks. By the time of CASP14,

AlphaFold2 predicted 3D structures of test proteins to unprecedented accuracy[10].

Currently, AlphaFold-predicted structure database (AFdb) contains hundreds of millions

of predicted structures providing the basis for functional genomics using fold-fold comparison.

Methods to predict functions using AFdb are on the rise. In one such effort, using DeepFRI, the

contribution of AFdb in enhancing the accuracy of protein function prediction achieved previously

by just using PDB was demonstrated[11]. Their results highlighted the importance of AFdb structures

among all three Gene Ontology (GO) categories.

The success of AlphaFold in predicting high-quality tertiary structures inspired

development of PANDA-3D[12], which is a novel computational tool designed to predict protein

functions by leveraging the high-quality 3D structural models in AFdb. Developed as a deep-

learning-based method, PANDA-3D focuses on combining the geometric vector perceptron graph

neural networks and transformer decoders for multi-label classification of GO over the traditional

function prediction tools using sequence-based annotation such as in PANDA[13] .

The research by Durairaj et al. explores the diversity of protein structures and functions

through advanced computational tools, leveraging the AFdb[14]. By integrating structure, sequence

similarity and machine learning tools like DeepFRI, the study developed an annotated sequence

similarity network to explore protein functions and evolutionary relationships. The study analyzed

over 350 million protein sequences from UniProt, revealing that many proteins remain

unannotated/uncharacterized due to limitations in sequence-based annotation methods. Their

efforts significantly enhance understanding of these "dark" proteins by the identification of new

protein families, such as TumE-TumA, a superfamily of toxin-antitoxin systems and discovery of

the beta-flower fold, a unique structural motif.

In yet another effort the authors attempted to cluster the 214 million structures in AFdb

using a novel structure comparison tool[15]. The authors use Foldseek, a scalable structural-

alignment-based method, to identify ~18 million clusters with 2.3 million non-singletons. Out of

the non-singleton clusters, 31% was unannotated representing potentially novel structures. Their

key findings include species-specific clusters comprising 4% of the total clusters, hinting at *de*

novo gene birth. Evolutionary insights suggest most clusters are ancient, with some human

immune-related proteins showing remote homology to prokaryotic species, suggesting ancient

systems may have been co-opted into specialized functions in higher organisms. Also, they report

new domain families, expanding the understanding of protein evolution and function.

Building on these advances, we have used both PDB and AFdb to explore the functional

diversity achieved by a single fold such as a primordial beta-trefoil fold[16]. This approach could be

considered a bottom-up approach to the work just described above, which can be considered top-

down. There are 1,387 protein containing trefoil domain in the current version of the PDB.

According to SCOP classification, beta-trefoil is implicated in eight distinct functional families

with capacity to bind to DNA (3brf), to carbohydrate (1jly, 1jlx), to chlorophyll (2dre), to

immunoglobulin-like receptor domain (1ira), to actin in a pH dependent fashion (1hcd), proteases

(6dwu) and AbfB (1wd3 and 2wd4).

RESULTS

In this work, the beta-trefoil fold was chosen to demonstrate the reach and limitations of

functional classification of a primordial fold using fold-fold comparison. Trefoil fold not only

displays functional diversity by promiscuously binding to every class of macromolecules, but is

the most divergent fold family challenging the sequence-based function prediction even among

paralogs of the same functional class within the same locus in human[17].

Functional diversity of trefoil fold in PDB

As representative of the trefoil fold, we selected accessions for 3D structures of five known proteins including 1i1b, 5uc6, 4p0j, 5bow and 6ncu from PDB. During the first iteration, each of these structures were searched for potential homologs against the PDB using DALI[18], a structure comparison tool. The r.m.s.d. cutoff for homolog selection was set at 2 Å and the query coverage greater than 80%. This process revealed nineteen distinct functions among proteins containing the trefoil fold. This is however an underestimate, since we missed other functions present in PDB because of the stringent cutoff. For example, we missed proteins containing the trefoil fold in the PDB such as LAG-1 (3brf), AbfB (1wd3) and beta-L-arabinopyranosidase (3a21). As shown in Figure 1, we found 51 entries containing trefoil domain spanning diverse functions in the PDB including GalNAc, Hemagglutinin, Lectin, Mosquitocidal toxin, Carbohydrate-Binding Family, Fibroblast Growth Factor, Ubiquitin, Neurotoxin, Ebulin, Cytolysin, Mycocypin/Clitocypin, Ricin, Lysenin, Kunitz Inhibitor, Inositol-3 Phosphate receptor, Fascin, Spindlin, Sevil and IL-1.

Figure 1: The core of the sundial represents the trefoil fold, a primordial recurring structural motif

in various proteins. The second ring classifies the proteins into functional families. The third layer

shows the distribution of trefoil fold across biological kingdoms, color-coded as blue for bacteria,

gray for fungi, green for plants, red for Metazoa, and yellow for protozoa. Pie charts within

functional families depict kingdom-level distribution of multidomain trefoil proteins in AFdb. The

outermost layer illustrates domain architecture, with colored blocks indicating different protein

domains, where purple represents the trefoil domain and others denote additional domains that

contribute to the function within the spoke such as Agglutinin (orange), Glycosyl transferase

family (dark-yellow), Glycosyl hydrolases family (dark-blue), Clostridium neurotoxin receptor-

binding domain (light-blue), RIH domain (brown), Ryanodine receptor-associated domain (dark-

green), Unknown domain 1 (light-pink), Unknown domain 2 (dark-pink), Unknown domain 3

(lemon-yellow), Unknown domain 4 (grey), Unknown domain 5 (pastel-green), Spindlin/Ssty

family domain (light-green), Cellulase (red), Carbohydrate binding family domain (forest-green).

The domain architecture of every protein identified containing the trefoil fold from the

PDB was characterized using HMM and their domain architecture is shown in the outermost ring

within each of the respective functional spokes. Table 1 lists 18 representative protein structures

containing distinct domain architecture that were selected for our second iteration. These

representative protein structures were queried against the AFdb v2.0 using the 3Di option in

Foldseek[19]. During this iteration, the search results were filtered to include only those with an

r.m.s.d. below 3 Å and covering more than 30% of the query structure. The 3 Å increase in cutoff

is based on our experience with PDB search, where we missed some functions because of the 2 Å

cutoff. The stringency of 30% query coverage for this iteration was essential to identify multi-

domain proteins containing trefoil from AFdb.

Details of the search results from mining AFdb are elaborated in Table 1. The second and

third column contain the query IDs for the representative query structures and their respective

kingdoms. The fourth column reports the number of hits with 80% versus 30% coverage from

AFdb and the last column highlights the kingdom diversity among the 30% hits. With a few

exceptions, hits with 80% and more coverage show less diversity at the kingdom level. We find

considerable kingdom-level diversity from AFdb with respect to the queries in column 3. For

example, with the query 3vsz belonging to bacteria, we find AFdb hits covering all four kingdoms.

On the other hand, 7kdu belonging to plants does not yield a single hit from plants. Similarly, 1ba7

from plants yields the structures only from bacteria.

Table 1: Validation of the approach taken here

Protein function	Representative Query PDB ID	Query Kingdom	Total number of Hits (>80% /30% coverage)	Number of Hits from a different kingdom among 30% coverage and less
Ricin	7KDU	Plantae	221/2250	

Ricin	3VSZ	Bacteria	2/368	
Ricin	3PHZ	Fungi	30/2038	
Hemagglutinin	3AH2	Bacteria	14/721	
Lectin	2IHO	Fungi	37/2271	
Lectin	4HR6	Plantae	152/553	
Carbohydrate binding family (CBF)	5G56	Bacteria	5/159	
Neurotoxin	5TPB	Bacteria	39/42	-
Ebulin	1HWM	Plantae	175/622	
Lysenin	3ZXD	Metazoa	5-Mar	

Lysenin	5EC5	Metazoa	12-Jan	
Kunitz inhibitor	1BA7	Plantae	631/634	
Inositol 3-P receptor	1N4K	Metazoa	Jun-34	
Fascin	4GOV	Metazoa	1049/1203	
Spindlin	5Y5W	Metazoa	16/55	
Spindlin	4UY4	Metazoa	76/188	
Spindlin	5A1H	Metazoa	1069/1223	
Galnac	5FV9	Metazoa	102/104	-

In column 4 of Table 1, the number of hits with 80% and 30% query coverage is shown.

For some of the queries, the number of hits are large for 80% cutoff allowing us to interrogate the

accuracy of predicted structures with respect to sequence identity. For the eleven query structures

with sufficient number of hits (purple, Table 1: column 2), the r.m.s.d. versus sequence identity

are plotted in Figure 2. As expected, there is a clear inverse correlation between r.m.s.d. and

sequence identity among these hits, validating that AlphaFold is a mega homology modeling tool.

Figure 2: For the eleven query PDB structures with accession mentioned inside, each plot shows

r.m.s.d. (x-axis: 0-3 Å) versus percent identities (y-axis: 30-100) for hits against AFdb with more

than 80% query coverage.

Figure 2 shows the plot of r.m.s.d on the x-axis verses sequence identity on the y-axis for

hits covering 80% of the query sequence against AFdb, for the eleven query structures inset within

the respective plots. The hits from AFdb for the eleven selected query PDB structures shown in

Figure 2, vary in pattern widely. A few follow a step pattern with each step representing distinct

classes or orders. For example, each step in the hits from 4gov, from metazoa, represent homologs

from birds at 78%, to fish at 66%, whale at 55% and Arthropoda at 42% identity. Similarly, hits

from 5a1h, from metazoa, have a step at 78% representing birds and bats. Other predominant

patterns in Figure 2 include those with missing structures above ~60% identity cutoff and below

~1.5 r.m.s.d. cutoffs, such as hits from 4hr6, 2iho and 5tpb. For example, the query 4hr6, is a lectin

from snake gourd- *Trichosanthes anguina*, a plant with hits between 60% - 50% belong to the

order Cucurbitales, hits between 50% - 40% mostly belong to the order Dipsacales and

Malpighiales, and a few from Rosales and Fabales, hits between 40% - 30% mostly belong to the

order Malvales and very few from Malpighiales, Cucurbitales and Caryophyllales. The query 2iho

is a lectin, which is from mushroom, and finds hits that are also from mushrooms (Basidiomycetes)

with a few exceptions from Ascomycetes, which is a sac fungus. The query 5tpb is a toxin from

bacterium *Clostridium botulinum* with most hits belonging to the same species and a few from the

same genus. In order to explain the gap in the region spanning low r.m.s.d. (<1.5 Å) and high

sequence identity (> 60%) in hits using query structure 4hr6, 2iho and 5tpb in Figure 2, we blasted

the sequence for the query protein against NR (Supplementary Figure 2 1C, 2B and 3C) and

UniProt databases (Supplementary Figure 2 1B and 3B) and find that this is resulting from

incompleteness in AFdb resulting from missing UniProt entries.

The third pattern is a sudden drop to 60% identity still showing low r.m.s.d. such as in

7kdu, a Rosid and in 1hwm, an Asterid. The 3-4 layers of hits against 7kdu are all plants from

different orders. For example, 98% identity is for another Rosid, 66% for plant species under

magnolids, and species under monocots show up at 43% identity. Similarly, hits against 1hwm at

81% come under Asterids and those at 43% belong to Rosids.

Functional annotation

Functional annotations of UniProt proteins available online are by homology at the

sequence level. These will include partial domain-level homology and may miss domain

architecture-based functional annotation. Hereafter, even for characterized entries in UniProt,

independent annotation was attempted using fold-fold comparison. In order to discover novel

domain architecture containing the trefoil fold, we interrogated hits with query coverage greater

than 30% for multi-domain query structures with the assumption that trefoil fold covers at least

30% of the multi-domain proteins. Kingdom-level diversity in functions for the query functions in

Table 1 is shown in pie-chart within the respective functional spokes in Figure 1 and Table 1. This

makes it evident that the fold-fold comparison strategy has the reach to identify homologs across

kingdoms. For example, the CBF (carbohydrate binding family) extends to metazoans and is not

limited to microbes as observed from PDB.

In Figure 3, we report thirteen novel UniProt proteins from AFdb containing a trefoil

domain along with an additional domain, which are not represented together in the PDB. These

proteins were categorized based on the presence or absence of these additional domains in the PDB

with or without trefoil domain. i) Category-1: Proteins that share their trefoil fold, but the structure

of the other domain is not present in the PDB database, suggesting novel domain architecture. ii)

Category-2: Proteins whose additional domain is found in the PDB without the trefoil fold. In these

cases, it was verified that the trefoil domain is absent even in the full length protein of the

corresponding PDB structures, indicating that while the additional domain is well represented, its

association with the trefoil fold is unique to the proteins identified. iii) Category-3: Proteins that

are represented in the PDB with both the trefoil fold and the extra domain as the hits, perhaps

missed during our first iteration because of stringent r.m.s.d. cutoff.

Figure 3: Novel domain architecture of UniProt proteins containing trefoil domains predicted by

AlphaFold that are not yet represented in the PDB. A tabular format is included in Supplementary

Table 1.

Among the 13 hits from AFdb, we found four category-1 and eight category-2 hits as shown

in Figure 3 and listed in Supplementary Table 1. Figure 4a, 4c, 4e, 4f, 5c, 5d, 5e and 5f shows hits

belonging to category-2. A0A6I5GSK5, which is annotated as chitinase sharing its chitinase

domain with 2dsk (r.m.s.d 1.342 Å), lacks trefoil domain even in the corresponding full-length

proteins (WP_317259669.1) with 742 amino acids. A0A4Q2KX23, sharing its beta-glucosidase

domain with 4i3g (r.m.s.d 1.016 Å) lacks trefoil domain in the corresponding full-length protein

(WP_230883105.1) with 832 amino acids. A0A3D9WMQ9 shares its protein kinase domain with

3f61 (1.95 Å), which lacks trefoil domain in the corresponding full-length protein

(WP_057333683.1) of length 626 amino acids. A0A820M8C8 shares its metallopeptidase domain

with 1slm (0.824 Å) lacking trefoil domain in the corresponding full-length protein (NP_002413.1)

with 477 amino acids. A0A191US67 shares its Beta-L- arabinofuranosidase domain with 4qjy

(2.83 Å) lacking trefoil domain in the corresponding full-length protein (WP_371878443.1) with

642 amino acids. A0A844M7R0 shares its phoD domain with 2yeq (0.741 Å) lacking trefoil

domain in the corresponding full-length protein (WP_009969242.1) with 583 amino acids.

A0A820T4Y8 shares the trypsin domain with 7n7x (0.628 Å) lacking trefoil domain in the

corresponding full-length protein (XP_054205941.1) with 436 amino acids. A0A3A9W2R1

shares the alginate lyase domain with 3zpy (0.976 Å) lacking trefoil domain in the corresponding

full-length protein (WP_303396801.1) with 446 amino acids.

Figures 4b, 4d, 5a and 5b show category-1 hits. The trefoil domains of A0A318NTU0,

A0A1I6E245, A0A3L7BAS7 and A0A0U3MKC8 gives an r.m.s.d. of 0.795 Å, 1.867 Å, 0.686

Å and 0.888 Å against a representative trefoil domain to confirm the presence of this domain in

these novel hits. The non-trefoil domains of these proteins there is no representation in the PDB.

Perhaps the annotation for these domains in UniProt is based on the presence of the annotated

domain in PFam.

Figure 4: Stereo diagram of the six characterized UniProt predicted structures not represented in

PDB. PDB queries in B and D representing category-1 are in green, AFdb structures in pink, PDB

hits for categories-2 and 3 in A, C, E and F are in blue.

Figure 5g shows the structure of R8CKC2, category-3, superimposed on the full structure

of 8bad annotated as a toxin from *Bacillus thuringiensis* with an r.m.s.d. of 0.976 Å. An enlarged

view of all proteins in Figure 4 and Figure 4 are shown individually in stereo figures in

Supplementary Figure 3a-f and Supplementary Figure 4a-g.

Figure 5: Stereo diagram of the seven uncharacterized proteins from AFdb containing trefoil domain not part of the functions and domain architecture presented in Figure 1. PDB queries in A and B representing category-1 are in green, AFdb structures in orange, PDB hits for categories-2 and 3 in C through G in blue.

DISCUSSION

The incomprehensible convergence in fold for billions of proteins, divergent at the

sequence-level, can now be used in functional characterization of thousands of proteins under each

organism using AFdb. Here, by identifying novel domain architecture in the AFdb structures

containing the trefoil fold, one of the primordial protein folds, we demonstrate the usefulness of

fold prediction in exploring functional diversity.

While the SCOP database represents eight functional diversities among proteins containing

the trefoil domain, our search in the PDB revealed 19 distinct functions for proteins containing the

trefoil fold shown in Figure 1. Except for a few, these functions span all kingdoms including plants,

bacteria, fungi and Metazoans. This could be an underestimate because of the 2 Å r.m.s.d. cutoff

used to limit false positives.

The discovery of the functional diversity from PDB is likely underestimated, as the PDB

is highly redundant and/or biased towards proteins of therapeutic or economic importance to

humans. Therefore, extending our analysis further, we searched the AFdb of over 200 million

predicted protein structures using representative PDB structures for each functional class

containing trefoil fold (inner ring, Figure 1). Not surprisingly, as shown in Figure 2, among the

AFdb structures from a given PDB query with 80% coverage, there is a clear inverse correlation

between sequence identity and r.m.s.d. for the predicted structures. Interestingly, the steps in the

plots of r.m.s.d. against sequence identity in Figure 2, represent a distinct class or order or

kingdom. In that, AlphaFold is a mega homology modeling tool on the one end and a novel fold

prediction tool on the other.

In order to look for novel functional architecture among the trefoil containing proteins in

AFdb, we interrogated hits with greater than 30% query coverage. Note that this cutoff leads to

false negatives by eliminating large proteins in which trefoil domains may span less than 30%. For

example, proteins with domain structures like 6mu1 (Supplementary Figure 6) where the trefoil

region spans only 15% of the entire structure. As mentioned in the results section, the hits are

binned into various categories depending on the presence of trefoil and/or other domains in PDB.

Among the proteins from AFdb containing trefoil fold, we discovered novel domain

architectures where the trefoil fold is found along with an additional domain that is either not yet

represented in the PDB or the complete domain architecture represents a novel domain

arrangement that is yet to be structurally characterized. With the exception of A0A820T4Y8, all

the novel proteins containing a trefoil domain identified here are from culturable bacteria and are

amenable to functional characterization using experimental techniques. For example,

A0A3L7BAS7 is a protein from *Micromonospora sp.* CV4, A0A191US67 is a protein from

Streptomyces parvulus, A0A844M7R0 is a protein from *Scytonema sp.* UIC 10036

(cyanobacteria), R8CKC2 from *Bacillus cereus*, A0A820T4Y8 from *Rotaria sp. Silwood2*

(Metazoa) and A0A0U3MKC8 from *Roseateles depolymerans*.

It should be mentioned that FoldSeek is able to find homologs of trefoil across diverse

kingdoms and functions, which is not possible using sequence-based annotation especially of folds

like trefoil (work not published). On the other hand, the extra reliance on Foldseek algorithm for

this project may also have limited the discovery of novel proteins containing trefoil fold, because

Foldseek reduces the structural elements into alphabets, which may still be biased toward sequence

similarity for short range interaction. For example, the performance of Foldseek to find structural

homologs of helical cytokines (results not shared) was poor. On the other hand, Foldseek performs

better with folds containing beta sheets, such as the TNF fold family (manuscript in preparation).

We believe that this is because helical cytokines are overrepresented on short-range interactions

compared to long-range interactions, which is the opposite for beta-sheet proteins.

Trefoil folds appear to be highly tolerant of amino acid changing mutations even within

the core. For example, paralogs within the same genome loci binding to the same receptor show

no significant sequence homology. For example, the function of IL1RA, an interleukin-1 receptor

antagonist, could only be predicted using 1D-3D threading approaches[17]. The work reported here

shows that the trefoil fold is present in all kingdoms with diverse domain architecture.

Furthermore, the diversity in functions of trefoil domain containing proteins is by binding to

diverse macromolecules including, DNA, carbohydrate, ig-like receptor domains, proteases,

chlorophyll, actin and by large oligomerization forming pores. In order to find out if beta-trefoil is

a converging or diverging evolution, we generated a phylogenetic tree using structure-based

sequence alignment for all proteins shown in Supplementary Figure 7. In divergent evolution one

would expect beta-trefoil containing proteins within the same functional family to cluster together

irrespective of kingdom. As shown in the phylogenetic tree (Supplementary Figure 7) these

proteins actually cluster by kingdom and not by function, suggesting convergent evolution. The

clustering by kingdom along with the observed diversity in delivering function via binding to

diverse macromolecules, it is tempting to hypothesize that this fold may have evolved

independently several times. If true, fold-based function prediction will hit a twilight zone just like

sequence-based function prediction. However, based on the amino acids at the equivalent positions

forming the core, shown in Supplementary Figure 8, the jury is still out on this hypothesis

METHOD

In Figure 6 we show the flowchart of the approach taken here. The r.m.s.d. cutoff of 2 Å

was used during the first round of searching PDB database for trefoil containing proteins. The

cutoff was relaxed to 3 Å while searching the AFdb with a query containing multiple domains.

Nevertheless, these cutoffs are stringent and may have led to false negatives.

Structure-based Functional Genomics

369 As shown in Figure 6a a few sample structures containing trefoil domain from the PDB database

370 were searched against PDB using DALI for all known structures containing trefoil fold. A filter of

371 2 Å was used for functional classification.

373 Figure 6: Flowchart of various steps along with filtering criteria

Validation of our approach

376   As shown in Figure 6b, the accepted hits for a given query against AFdb with a query coverage of

377   80% and more is plotted using r.m.s.d. against sequence identity ranging from 30% to 100%.

Identification of novel protein containing trefoil fold from PDB

As shown in Figure 6c, an expanded list of representative structures were selected from each

kingdom representing a given function including those containing multiple domains. These were

used as query sequences to search AFdb using Foldseek with 3Di option and a cut of 3 Å was used

to select hits for downstream analysis. These hits included characterized and uncharacterized

proteins based on existing annotation and were classified as various categories based on the

existence of 3D structure of individual domains in PDB.

ACKNOWLEDGEMENT

387   The authors thank GoK for funding for data analysis personnel via BioIT grant and computing

388   infrastructure via the Department of IT, BT and ST.

CONFLICT OF INTEREST

390   Authors have no conflict of interest.

AUTHOR CONTRIBUTION

392   MG: The entire bioinformatics work, SS: for conceptualization, guiding bioinformatics work and

393   writing of the manuscript

DATA AVAILABILITY

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

BIBLIOGRAPHY

1. Chothia, C. Proteins. One thousand families for the molecular biologist. *Nature* **357**, 543–544 (1992).

2. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

3. Dayhoff, M., Schwartz, R. & Orcutt, B. *Atlas of Protein Sequence Nd Structure*. vol. 5 (1978).

4. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

5. Eddy, S. R. Hidden Markov models. *Curr. Opin. Struct. Biol.* **6**, 361–365 (1996).

6. Hameduh, T., Haddad, Y., Adam, V. & Heger, Z. Homology modeling in the time of collective and artificial intelligence. *Comput. Struct. Biotechnol. J.* **18**, 3494–3506 (2020).

7. Simons, K. T., Strauss, C. & Baker, D. Prospects for ab initio protein structural genomics. *J. Mol. Biol.* **306**, 1191–1199 (2001).

8. Wodak, S. J., Vajda, S., Lensink, M. F., Kozakov, D. & Bates, P. A. Critical Assessment of Methods for Predicting the 3D Structure of Proteins and Protein Complexes. *Annu. Rev.*

Biophys. **52**, 183–206 (2023).

9. Kamisetty, H., Ovchinnikov, S. & Baker, D. Assessing the utility of coevolution-based

residue-residue contact predictions in a sequence- and structure-rich era. *Proc. Natl. Acad.*

Sci. U. S. A. **110**, 15674–15679 (2013).

10. Kryshtafovych, A., Schwede, T., Topf, M., Fidelis, K. & Moult, J. Critical assessment of

methods of protein structure prediction (CASP)-Round XIV. *Proteins* **89**, 1607–1617 (2021).

11. Ma, W. *et al.* Enhancing Protein Function Prediction Performance by Utilizing AlphaFold-

Predicted Protein Structures. *J. Chem. Inf. Model.* **62**, 4008–4017 (2022).

12. Zhao, C., Liu, T. & Wang, Z. PANDA-3D: protein function prediction based on AlphaFold

models. *NAR Genomics Bioinforma.* **6**, lqae094 (2024).

13. Wang, Z., Zhao, C., Wang, Y., Sun, Z. & Wang, N. PANDA: Protein function prediction

using domain architecture and affinity propagation. *Sci. Rep.* **8**, 3484 (2018).

14. Durairaj, J. *et al.* Uncovering new families and folds in the natural protein universe. *Nature*

622, 646–653 (2023).

15. Barrio-Hernandez, I. *et al.* Clustering predicted structures at the scale of the known protein

universe. *Nature* **622**, 637–645 (2023).

16. Tenorio, C. A., Longo, L. M., Parker, J. B., Lee, J. & Blaber, M. Ab initio folding of a

trefoil-fold motif reveals structural similarity with a β-propeller blade motif. *Protein Sci.*

Publ. Protein Soc. **29**, 1172–1185 (2020).

17. Bazan, J. F., Timans, J. C. & Kastelein, R. A. A newly defined interleukin-1? *Nature* **379**,

591 (1996).

18. Holm, L. & Sander, C. Dali: a network tool for protein structure comparison. *Trends*

Biochem. Sci. **20**, 478–480 (1995).

19. Fast and accurate protein structure search with Foldseek - PubMed.

https://pubmed.ncbi.nlm.nih.gov/37156916/.