'Caroten-omics' of pigments synthesised by *Gordonia rubropertincta* through

High Resolution Mass Spectrometry and *in-silico* programming approach

Gayathri Vemparala[a,c] , Gangagni Rao Anupoju[a,c], Thenkrishnan Kumaraguru[b,c,*]

[a]Bioengineering and Environmental Sciences (BEES) group, Department of Energy and

Environmental Engineering (DEEE),

CSIR-Indian Institute of Chemical Technology (IICT), Tarnaka, Hyderabad – 500007, India

[b]Department of Organic Synthesis and Process Chemistry (OSPC), CSIR-IICT

[c]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India - 201002

*Corresponding Author Email: kumaraguru@iict.res.in; kumaraguruchem@gmail.com

ABSTRACT

Carotenoids are tetraterpenoid pigments synthesised by microorganisms, plants and present in some animals. They are metabolites of importance in research and industry. The present study aimed to analyse the High Resolution Mass Spectrometry (HRMS) experimental data through R programming and identify all the carotenoids that *Gordonia rubropertincta* produces. The growth rate of the bacterium was 0.24 ± 0.132 h^{-}1, produced about 140 ± 20 mg carotenoids/ g wet cell weight (WCW) on extraction with methanol and the extract was subjected to HRMS. The number of carotenoids the bacterium synthesizes was identified to be 93 and the concentration of few carotenoids at particular time was pinpointed using R. The distinctive features of the bacterium like high number of carotenoids synthesis through various pathways, their fluctuation with respect to time, storage of carotenoids in cell cytoplasm, and its auto-fluorescent property were also discovered.

Keywords: Carotenoids, Growth rate, HRMS, R programming, Microscopy, Auto-fluorescent

1.0 INTRODUCTION

Nutraceuticals are potential solutions to challenges such as malnutrition (Katata-Seru et al., 2019), disease prevention (Daliu et al., 2019) and an alternate to chemical food additives and enhancers (Munekata et al., 2020). Molecules like amino acids, vitamins, fatty acids, along with metabolites like carotenoids, antioxidants, polyphenols, fall under the nutraceutical umbrella (GNK et al., 2018; Padmavathi, 2018). Carotenoids are a diverse group of pigments that are popular for their therapeutic properties (Cvetković et al., 2022) and thus are important for production at industrial scale. Carotenoid pigments are produced by wide range of organisms starting from bacteria through plants and also found in animals (De Miguel et al., 2001).

Carotenoids are red-yellow coloured, secondary metabolites made up of isoprenoid units (López et al., 2021). They are tetraterpenoids that contain eight isoprene units (Kumari et al., 2020; López et al., 2021). Most carotenoids are hydrocarbons and are highly non-polar compounds while some carotenoids contain a few oxygen atoms that contribute to their partial polarity (Giuffrida et al., 2018). More than 1200 varieties of carotenoid pigments have been identified to date from various sources and documented (Nupur et al., 2016; Yabuzaki, 2017). Carotenoids are microbial and plant secondary metabolites of high importance in various industries and research purposes. They are well-known for their applications as nutrient supplements, precursors, anti-microbial, as well as cell and DNA protecting agents (Kłodawska et al., 2019). Some carotenoids such as, staphyloxanthin, are known to confer virulence to the host organism producing it. Therefore it can be used as a target for such pathogenic treatments (Ye et al., 2022). According to Business Communications Company (BCC) Research, the current market value of carotenoids is estimated to be $2 billion in 2022 which could rise up to $2.7 billion by 2027 (Bcc and Staff, 2022).

Among the different sources of carotenoids, microorganisms such as bacteria (including

Actinobacteria) and microalgae are the best sources for industrial scale production when

compared to plants and animals (Loh et al., 2020). Microbes grow quickly in large numbers, are

easy to handle at both laboratory and commercial scales, and, simple to study and rewire

genetically for higher yield. Microorganisms belonging to genera, *Dunaliella, Haematococcus,*

Chlorella, Paracoccus, Dietzia, Rhodotorula, and *Brevundimonas,* are known for their

carotenoid producing capacity (Lee et al., 2022; Mussagy et al., 2021; Ram et al., 2020; Zhao et

al., 2023). Actinobacteria contain bacterial species that are mostly pigmented and thus appear

coloured. Genera such as *Micrococcus, Arthrobacter, Curtobacterium, Corynebacterium,* and

Rhodococcus are actinobacteria that are known to produce carotenoids (Sandmann, 2021).

Gordonia is an actinobacterial genera that were reported to synthesise new carotenoid types that

were not known previously (Sowani et al., 2018). The *Gordonia* species like *G. alkanivornas*

(Silva et al., 2016)*, G. ajoucoccus* (Kim et al., 2014)*, G. tearrae* (Elfalah et al., 2013)*,* and *G.*

jacobaea (De Miguel et al., 2001)*,* are known for their ability to synthesize carotenoids. *G.*

rubropertincta is another species that was not studied with respect to the carotenoid pigments it

synthesises. Moreover, to study carotenoids synthesised by any bacterium, elaborate extraction

procedures and analysis such as High Performance Liquid Chromatography (HPLC) that require

standardizations with controls is necessary which might make the process expensive and limited

to certain carotenoid metabolites. Therefore, the intention behind current work is to identify the

carotenoids that *G. rubropertincta*, produces by combining the methods of untargeted metabolite

analysis through High Resolution Mass Spectrometry (HRMS) and the data analysis using R

programming to quickly identify all carotenoids in the bacterium.

Initially, the growth pattern of the bacterium was studied which gave us a brief insight into how

the pigment synthesis might proceed with respect to the incubation time as microbes synthesize

carotenoids during the stationary phase of growth.. Then a solvent extraction method was applied

to extract the pigments from the bacterial cell. Then, extracted pigments were identified by

HRMS which revealed various types of carotenoids pigments the bacterium synthesises. The

data obtained through HRMS were sorted and interpreted. Finally, the morphology of the

microbe was studied initially by Scanning Electron Microscopy (SEM) followed by fluorescence

and confocal microscopy to locate the carotenoids in the cell. Transmission Electron

Microscopiy (TEM) technique was employed to confirm the location of the pigment in the cells

which was expected to aid in the carotenoid extraction process.

2.0 MATERIALS AND METHODS

2.1 Bacterium acquisition, maintenance, and morphological studies

The current study was carried out using *G. rubropertincta* collected from Microbial Type Culture

Collection and Gene Bank (MTCC), CSIR-Institute of Microbial Technology, Chandigarh, India,

(Accession No. 289). The microbial culture was maintained in the Nutrient Broth medium

(Himedia). For all the experiments of the present study, 1000 mL of bacterial culture was grown

at 25 ℃ for 48 h and stored at 2-4 ℃.

2.2 Bacterial growth curve and growth rate analysis

To perform this experiment, bacterial culture was inoculated in nutrient broth medium, and 1 mL

of liquid sample was collected every 2 h up to 48 h and later, samples were collected every 24 h.

Based on the results the experiment was terminated after 15 d. The culture was centrifuged at

7000 rpm for 2 min. The supernatant was discarded, and the pellet weight was recorded to

calculate growth rate of the microbe. The pellet was then suspended in distilled water and

Optical Density (OD) was taken at 600nm (Hach-DR6000 UV-VIS spectrophotometer,

Loveland, CO, USA). Graph has been plotted for growth curve and growth rate was calculated

using the Equation 1 (Fernández-naveira et al., 2016). This experiment was performed in

triplicates and the average of these triplicates were used to plot the graphs.

111
$$\mu = \frac{[Ln(N_t) - Ln(N_0)]}{(t - t_0)} \quad \square^{-1} \qquad \text{Eq. 1}$$

Here, μ is the growth rate, N_t is the cell density in g/L at time t (hours), N_0 is the

cell density at time 0 h (t_0).

2.3 Extraction and quantification of pigments

The carotenoid pigments were extracted from the bacterial cell at the end of 15 d of incubation

through a modified solvent extractions process (Bisht et al., 2020; Poddar et al., 2021). The

bacterial culture was pelleted by centrifuging at 7000 rpm for 5 min. The supernatant was

removed, and pellet was washed thrice with distilled water. 50 mg of wet pellet was taken in five

centrifuge tubes each for four different solvents extraction and a combined solvents extraction.

The solvents used for extraction were acetone, methanol, 1-butanol, and ethyl acetate (Sisco

Research Laboratories Pvt. Ltd., Mumbai, India). Their combination in equal proportion was also

used for extraction. 2 mL of solvent respectively was added to each of the tubes and vortexed for

1-2 min. It was then centrifuged at 7000 rpm for 5 min. The supernatant solvent containing the

pigments was collected into separate centrifuge tubes and the procedure was repeated thrice. To

quantify the pigments, the solvent was dried by a rotary evaporator (Heidolph Schwabach,

Germany) and; remaining pigments were weighed and recorded. After weighing, the pigments

were again dissolved in 2 mL respective solvents. All the solvents were mixed and lyophilised.

(Labconco-FreeZone Plus 6 lyophiliser). The lyophilised pigments were used for further

experiments.

2.4 Liquid Chromatography/High Resolution Tandem Mass Spectrometry (LC/HRMS)

The lyophilized pigments were dissolved in methanol and the types of carotenoids present in the

sample were analysed by LC/HRMS (Waters-Xevo G2-XS QTof, Milford, MA, USA). This

method was developed for the identification of carotenoid pigments and was done by using the

data about every carotenoid such as molecular weight and structure available in the Carotenoid

Database (López et al., 2021).

2.5 Scanning Electron Microscopy (SEM)

The bacterial culture was centrifuged as above to get a pellet and supernatant was discarded. The

pellet was washed thrice with distilled water and then lyophilized. The lyophilized bacterial

culture was placed on a conductive carbon double-sided adhesive tape (Nisshin EM Co., Ltd.,

Tokyo, Japan) attached on one side to a metal stub. The stub was left to rest to facilitate the

sample sticking to the carbon tape. The stub was placed in the sample chamber of the SEM and

observed (Jeol-JSM-7610F Schottky Field Emission Scanning Electron Microscope, Tokyo,

Japan).

2.6 Fluorescence and Confocal microscopy

The localisation of the pigments synthesised by the bacterium was done using various

microscopic techniques. To find out the wavelength range of the pigments produced by the

bacteria under study, initially the bacteria were observed under three filters of a fluorescent

microscope namely, DAPI, GFP and TRITC. This is a simple procedure designed to fix bacterial

cells between a slide and coverslip for making the observation under the microscope easy. The

slide was prepared by pelleting the cells by centrifugation, followed by washing the pellet with

1X Phosphate-Bufferd Saline (PBS) for 3 times. The cells were then suspended in 1X PBS and

200 µL was layered onto a sterile 22 mm^2 cover slip which was placed in a sterile Petri dish. The

setup was incubated overnight under 30 ℃ temperature. Then, the cover slip was washed thrice

with 1X PBS and observed under bright-field microscope to confirm the cellular attachment to

the cover slip. The cover slip was fixed on to a microscopic glass slide using Eukitt® Quick-

hardening mounting medium (Sigma-Aldrich, Burlington, MA, USA). The slide was observed

under a fluorescence microscope (Olympus-IX73P1F, Tokyo, Japan) and then under confocal

laser scanning microscope (Olympus-FluoView FV10i, Tokyo, Japan).

2.7 Transmission Electron Microscopy (TEM)

To confirm the location of the pigments inside the bacterial cell, TEM was employed. The

bacterium was grown as described above and pelleted by centrifugation. The pellet was

preserved in 2.5% glutaraldehyde (Sigma Aldrich, 0.1M phosphate buffer – pH 7.3) and stored at

4 ℃ for 24 h and processed for the TEM study as per standard protocol(Lakshman, 2019). The

samples were washed in buffer and post fixed in 1% aqueous Osmium tetroxide (Sigma Aldrich)

later dehydrated in ascending grades of acetone (Qualigens fine chemicals, Mumbai, India),

embedded in Spurr's resin (SPI supplies, araldite 6005 Embedding kit, USA) and were incubated

(Universal incubator-NSW-151) for 24 h at 70 ℃ for complete polymerization of tissues. The

blocks were trimmed manually and semi thin (700-800 nm) sections were made with ultra-

microtome (Leica ultra-cut UCT-GAD/E-100, Germany), stained with 1% toludine blue

(Qualigens fine chemicals) and observed under a light microscope (Olympus-AX 70) to locate

exact area and to remove extra resin, if any for making ultra-thin sections (50nm thickness).

Suitable ultra-thin sections were mounted on copper grids (SPI supplies, USA) allowed to air dry

overnight and were stained with Urenyl Acetate (UA) and 1% Reynold's Lead Citrate (LC) as

previously reported (Bozzola and Russell, 1998). All grids were dried at room temperature and

observed under TEM (Talos F200X TEM) at 120kV.

3.0 RESULTS

3.1 Bacterial growth curve and growth rate analysis

The bacterial growth curve shows the transition of growth phases of the bacterium under study

with respect to time. This information should aid in studying the pigment production as

carotenoids are secondary metabolites synthesised by the bacterium during its stationary phase

(Narsing Rao et al., 2017). The bacteria, as can be seen in **Fig. 1(a),** remain in the lag phase for

about 4 h after inoculation and then enters the logarithmic/exponential phase. The log phase

initiates and rapid exponential growth occurs between 6-12 h of incubation and hence a higher

deviation in the OD could be seen. After 12 h, although not exponential, the bacterium remains

in log phase for about 3 d as can be seen from **Fig. 1(b)**. The stationary phase then starts and

continues to persist for a very long time, i.e., up to 15 d which can be noticed from **Fig. 1(b),** and

then starts descending gradually**.** Hence, the bacterium had the potential to synthesize pigments

for about 15 d after inoculation. Also, the maximal growth rate of the bacterium was found to be

$0.24 \pm 0.132 \, h^{-1}$.

Fig. 1(a): *Gordonia rubropertincta* growth curve for 48h. The dots represent the mean of triplicate experiments and the error bars represent standard deviation from mean.

Fig. 1(b): *Gordonia rubropertincta* **growth curve for 15d. The dots represent the mean of**

triplicate experiments and the error bars represent standard deviation from mean.

3.2 Extraction and quantification of pigments

The pigment extractions were performed by solvent extraction method at the end of 15 d of incubation. Four solvents were chosen based on their polarity, boiling points, pigment solubility, ease of availability and solvent recovery. A combination of all the solvents in equal proportion was also used. Acetone, methanol, 1-butanol, ethyl acetate, and the combination yielded an average of 100 ± 20mg, 140 ± 20mg, 66.67 ± 16mg, 126.67 ± 12.5mg and 90.67 ± 19mg carotenoids/g wet cell weight (WCW) respectively. Fig. 2 is a graphical portrayal of the pigment yield with various solvents. Of all the solvents, it is evident that methanol could extract the highest amount of pigment compared to the other solvents making it the most suitable solvent for carotenoid pigment extraction from the bacterium, *G. rubropertincta*.

Fig. 2: Carotenoids extracted from *G. rubropertincta* using various solvents and their quantification.

3.3 Liquid Chromatography/High Resolution Tandem Mass Spectrometry (LC/HRMS)

As the types of carotenoid pigments present in the bacterium are unknown, a whole metabolite analysis was performed using the HRMS technique. The analysis revealed that the bacterium synthesises 93 types of carotenoid pigments whose quantity in terms of percentage can be seen in **Fig. 3(a)** and out of them, 16 accounted for highest quantity than others. The obtained results were compared with the KEGG carotenoid biosynthesis pathway database (https://www.genome.jp/kegg/pathway.html) and it was determined that the high number of carotenoids identification through HRMS is due to the fact that the intermediates of the pigments synthesis pathways are also carotenoid type. Therefore, the intermediates and the end products together resulted in a very high number. All the identified pigments and their abundance could be seen in **Figure S1**. To confirm this interpretation, same analysis was performed with pigments extracted at specified time intervals of 3 days up to 15 days to check the progression of carotenoid synthesis. The obtained HRMS data were sorted based on factors like carotenoids present throughout 15 d, carotenoids specific to a particular time interval but not present on other days, the quantities of each carotenoid and their fluctuations throughout 15 d. The detailed information on the carotenoids and their abundance on each day are given in the **Figure S2(a)-(e)** in the **supplementary material**. **Fig. 3(b)** shows only those carotenoids that are present continuously throughout all the 15 days and how their quantities fluctuate with time. The presence of varieties of carotenoids varied in quantity at different times of bacterial incubation shows that this is a simple and effective combination of methods to determine the concentration

241 of any metabolite. Also, the bacterium under study is a store house of wide range of carotenoids

242 that could have multiple applications in research and industry.

246 **Fig. 3(a): Carotenoid pigments synthesized by *G. rubropertincta* in highest quantity**

247 **identified by High Resolution Mass Spectrometry**

Fig. 3(b): Carotenoid pigments synthesized by *G. rubropertincta* that are present throughout the 15 d and their varying concentrations at regular time intervals identified by High Resolution Mass Spectrometry

3.4 Scanning Electron Microscopy (SEM)

Fig. 4 shows microscopic observation of *G. rubropertincta* which revealed the surface

morphology to be long, rod-shaped, bacilli-like structures although cells slightly vary in length.

The bacterial cells grow at one or either ends attached to another cell creating a film or a layer in

the broth medium. This might be due to the formation of non-septate hyphae. The width and

depth differ at each point of the microbe resembling a rather non-uniform bacillus. Some cells

show a characteristic structure of one end branching into two and giving it the shape of the

alphabet letter Y. Evidently, the microbe shows features peculiar to actively growing

Actinobacteria (Hopwood, 2007)*,* and further experiments in this study uncover some particular

cellular details about *G. rubropertincta*.

273   **Fig. 4: Scanning Electron Microscopy of *Gordonia rubropertincta***

(a)-(d) micrographs of increasing magnification. (a) & (b) long, slender and connected

bacterial cells (c) cells resemble rods with varying measurements (d) blue arrows indicate

cell branching at one end

3.5 Fluorescence and Confocal microscopy

The bacterium was observed under three filters, DAPI, GFP and TRITC of a fluorescence

microscope to find out the fluorescing capability of the pigments and also to locate them in the

cell. The whole cell appeared blue in colour under DAPI filter indicating that the pigments could

fluoresce and are carotenoid type. The location of the pigments could not be pinpointed and

hence a confocal microscopic technique was employed. **Fig. 5** shows the bacterial cell under

confocal microscope with carotenoids fluorescing in blue. Two distinct layers could be seen on

the exterior of the cell which probably is the cell wall and plasma membrane where no

fluorescence could be seen. As the bacterium under study is a gram positive (Nouioui et al.,

2018) one the outermost thick layer of the cell can be attributed as the cell wall with a cell

membrane inner to it. The fluorescence emission occurs from the interior of the cell and that

indicates the pigment location to be the cytoplasm. Further experiments were carried out to

confirm this interpretation.

3.6 Transmission Electron Microscopy (TEM)

The TEM image, **Fig. 6,** gave a clear picture of the bacterial structure and the pigment location.

The exterior black line in **Fig. 6(b),** is the stained cell wall and plasma membrane of the

bacterium. The genetic material appears as darkly stained body inside the cell as it gets stained

by UA (Kabiri et al., 2019) and the whole cell appears dark indicating the presence of

carotenoids inside the cell being distributed all over in the cytoplasm. **Fig. 6(a)** and **(c)** also

shows dark cells with two distinct layers outside the cell and the genetic material inside as a

black dot. TEM provides evidence that the bacterium under study stores the synthesised

carotenoids in the cell cytoplasm. Till date carotenoids were known to be present in bacterial

cells bound to their plasma membranes (Hagaggi and Abdul-Raouf, 2023; Kirti et al., 2014).

Determining the location of carotenoid storage inside the bacterial cell helps in developing

efficient methods to extract the carotenoid pigments from the bacterium.

(a)   (b)   (c)

312   **Fig. 6: Transmission Electron Microscopy of *Gordonia rubropertincta***

313   **(a) & (c) Individual bacterial cells. (b) One bacterial cell with dark cytoplasm. Blue**

314   **arrow indicates the plasma membrane and red arrow shows the genetic material**

315   **of the bacterium**

4.0 DISCUSSION

The carotenoid pigments produced are secondary metabolites and therefore it would be useful to precisely know the stationary phase of the bacterium for extraction of pigments. Of the four growth phases, the bacterium enters the stationary phase between 6-12 h of incubation and remains in it for about 15 d. The growth rate of the bacterium was 0.24 ± 0.132 h^{-1}.

Four solvents, acetone, methanol, 1-butanol, ethyl acetate and an equal mixture of all four were used for the extraction process. Carotenoids are C_{20}-C_{60} hydrocarbon in nature with some containing a few oxygen atoms. They are highly non-polar molecules with the oxygen contributing to a little polarity to some carotenoids. Hence, solvents of increasing relative polarity (Miller's Home Solvent polarity table) were chosen for extraction so that carotenoids of various polarities get dissolved in the solvents and are extracted easily. Also, the availability of solvents, their ability to dissolve cell wall and plasma membrane to release cellular contents including carotenoids, solubility of pigments and ease of solvent recovery were other solvent properties taken into consideration. The carotenoids could be easily extracted from the cells and separated from cell debris through centrifugation. Of the solvents, methanol was found to be most suitable for extracting high amount of carotenoids i.e., 140 ± 20 mg carotenoids/g WCW. Thus extracted carotenoids were then lyophilised and used for the identification of types of carotenoids by HRMS method.

A mixture of carotenoids were observed that were 93 in total. Each carotenoid showed a particular response that corresponds to the quantity present in the whole mixture. So, HRMS is both a qualitative and quantitative estimation of whole carotenoids present in the extract. When compared with the KEGG (https://www.genome.jp/kegg/pathway.html) and Carotenoid database

(Yabuzaki, 2017) the 93 carotenoids identified were found to be the intermediates and end

products of carotenoid synthesis pathways. It was also found that *G. rubropertincta* is a

bacterium capable of synthesising varieties of carotenoids some of which are not known till date

to be synthesised by bacterium but produced by higher organisms such as algae, fungi and plants.

For example, neurosporaxanthin methyl ester (Parra-Rivero et al., 2020) and crocetin (Guo et al.,

2022) that have pharmacological applications are generally found in fungi and plants

respectively (Yabuzaki, 2017). Synthesis of such important carotenoids by a bacterium could be

highly beneficial to produce them at commercial scales.

The HRMS analysis was performed on carotenoids extracted at regular intervals of time for 15 d

as the bacterium stays in stationary phase for about 15 days as found through growth curve

analysis. Extraction was performed every 3 d up to 15 d. Intermediates and end products were

seen in all 5 samples with the types and abundances varying on each day. This information aids

in identifying the pathway(s) through which the organism is actively synthesising carotenoids,

the concentration of each carotenoid at any point of time, so, targeting and harvesting any

particular carotenoid is possible. *G. rubropertincta* synthesises carotenoids through diverse

pathways such as Neurosporaxanthin, bacterioruberin, spirilloxanthin, okenone. While the types

and concentrations of carotenoids on each are varying, 23 carotenoids were found to be present

throughout the 15 d. The time point at which a carotenoid is present at maximum quantity can be

identified through this method which helps in deciding the incubation or fermentation time to

obtain the maximum amount of product.

SEM revealed the bacteria to be rod-shaped, bacilli-like but cells connected to each other due to

the formation of non-septate hyphae. Carotenoid pigments such as, β-carotene, rhodopin,

spheroidenone, fucoxanthin, lycopene (Gillbro and Cogdell, 1989; Katoh et al., 1991) are all

fluorescent carotenoids whose fluorescence spectra lies in the UV-VIS region (Miller et al.,

1935). *G. rubropertincta,* when observed under three different filters i.e, DAPI (4',6-diamidino-

2-phenylindole), TRITC (Tetramethylrhodamine), GFP (Green Fluorescent Protein), of a

fluorescence microscope, it fluoresced under the DAPI region showing the carotenoids it

synthesises are also auto-fluorescent. Taking advantage of the auto-fluorescence, the bacterium

was then observed under a confocal microscope to precisely locate the carotenoids inside the cell

as that information can be useful in developing an efficient extraction strategy. When observed

using a confocal microscope, a much clearer and sharper image of the bacterium was obtained

showing some fine details on the cellular characteristics. The bacterial cell consists of a thick cell

wall as it is gram-positive in nature, a lipid bilayer inside the cell wall, and the fluorescence

emerges only from the inside of the plasma membrane indicating the location of carotenoids in

the cell cytoplasm. It has been previously reported that carotenoids are present in the plasma

membrane bound to one of the leaflets of the lipid bilayer or embedded inside the cell membrane

with hydrophobic ends tucked inside and exposed hydrophilic ends (Anwar et al., 1977; Chia et

al., 2021). Bacterial carotenoids are also found bound to the bacteriochlorophyll pigments

present in the plasma membrane (Yurkov et al., 1993). The carotenoids synthesised by *G.*

rubropertincta are revealed through confocal microscopy to be stored completely in the

cytoplasm and are not bound to the cell membrane. The carotenoids' location was further

confirmed by TEM. The phospholipid bilayer present on the exterior of the bacterial cell, genetic

material that can be seen as a darkly stained body inside the cell, and the darkly stained

cytoplasm could be seen through TEM. The cytoplasm getting stained in black indicates the

presence of carotenoids in the cytoplasm of the cell. Being present in the cytoplasm makes it

easier to bring the carotenoids out of the cell and separate it from other cell debris compared to

those that are membrane bound. Therefore, extracting the carotenoids and purifying them should

be easier when *G. rubropertincta* is used as the host organism.

5.0 CONCLUSION

To summarize, *Gordonia rubropertincta* is a gram-positive, actinobacteria that synthesises a

broad range of carotenoids through various pathways. The carotenoids it synthesises are auto-

fluorescent and present in the cell cytoplasm, so are easy to extract from the cell. It is a store

house of different carotenoids. Each carotenoid can be used for different applications, and

bacteria are easy to handle, study and maintain at lab to industrial scale compared to higher

organisms like fungi, algae, and yeasts. Hence, this bacterium can be used to commercially

produce carotenoids. The diversity of carotenoids it synthesises makes it an ideal organism to

study carotenoids, their pathway(s), properties, applications, to metabolically engineer and

increase or decrease production of a particular carotenoid, and channelise carotenoid synthesis

into one or more particular pathway(s). This organism can be of high importance in industries to

produce carotenoids commercially.

ACKNOWLEDGEMENTS

The authors are thankful to The Director, CSIR-Indian Institute of Chemical Technology (IICT)

for providing the facilities required for carrying out this research work (**IICT/Pubs./2023/097**).

The authors are grateful to the Academy of Scientific and Innovative Research (AcSIR) for their

immense support during this work. Also, sincere thanks to the Indian Council of Medical

Research (ICMR) for providing Ms. Gayathri Vemparala with research fellowship (JRF-2019,

HRD-22). Thanks to Dr. Mohana Krishna Reddy Mudiam, Dr. B. Sreedhar, Dr. Rajkumar

Banerjee, Dr. Sumana Chakravarty, Narendra Kumar Nagendla, and Sriravali Pulipaka for their

help in analytical techniques. Thank you Dr. M. Srinivasa Rao and Dr. Gurrala Sheelu for

reviewing and enhancing the article. Thank you Bhavya Karumanchi, Abhishiktha Chiliveru and

all other dear friends, family and colleagues for their moral support which led to the successful

completion of this work.

REFERENCES

Anwar, M., Hasan Khan, T., Prebble, J., Zagalsky, P.F., 1977. Membrane-bound carotenoid in

Micrcoccus luteus proects naphthoquinone from photodynamic action. Nature 270, 538–

540.

Bcc, M.F., Staff, P., 2022. The Global Market for Carotenoids.

Bisht, G., Srivastava, S., Kulshreshtha, R., Sourirajan, A., Baumler, D.J., Dev, K., 2020.

Applications of red pigments from psychrophilic Rhodonellum psychrophilum GL8 in

health, food and antimicrobial finishes on textiles. Process Biochem. 94, 15–29.

https://doi.org/10.1016/j.procbio.2020.03.021

Bozzola, J.J., Russell, L.D., 1998. Electron Microscopy Principles and Techniques for

Biologists.

Chia, G.W.N., Seviour, T., Kjelleberg, S., Hinks, J., 2021. Carotenoids improve bacterial

tolerance towards biobutanol through membrane stabilization. Environ. Sci. Nano 8, 328–

341. https://doi.org/10.1039/d0en00983k

Cvetković, B., Kolarov-Bjelobrk, I., Radić, J., Bosanac, M., Janičić, S., Stupar, A., Samardžić-

Lukić, M., Ćuk, D., Tovilović, V., Andrejić-Višnjić, B., 2022. Can carotenoids cure or

protect us? Hosp. Pharmacol. - Int. Multidiscip. J. 9, 1183–1188.

https://doi.org/10.5937/hpimj2202183c

Daliu, P., Santini, A., Novellino, E., 2019. From pharmaceuticals to nutraceuticals: bridging

disease prevention and management. Expert Rev. Clin. Pharmacol. 12, 1–7.

https://doi.org/10.1080/17512433.2019.1552135

De Miguel, T., Sieiro, C., Poza, M., Villa, T.G., 2001. Analysis of canthaxanthin and related

pigments from Gordonia jacobaea mutants. J. Agric. Food Chem. 49, 1200–1202.

https://doi.org/10.1021/jf001169z

Elfalah, H.W.A., Ahmad, A., Usup, G., 2013. Anti-Microbial Properties of Secondary

Metabolites of Marine Gordonia tearrae Extract. J. Agric. Sci. 5, 94–101.

https://doi.org/10.5539/jas.v5n6p94

Fernández-naveira, Á., Abubackar, H.N., Veiga, M.C., 2016. Carbon monoxide bioconversion to

butanol-ethanol by Clostridium carboxidivorans □: kinetics and toxicity of alcohols 4231–

4240. https://doi.org/10.1007/s00253-016-7389-8

Gillbro, T., Cogdell, R.J., 1989. Carotenoid fluorescence. Chem. Phys. Lett. 158, 312–316.

https://doi.org/10.1016/0009-2614(89)87342-7

Giuffrida, D., Donato, P., Dugo, P., Mondello, L., 2018. Recent Analytical Techniques Advances

in the Carotenoids and Their Derivatives Determination in Various Matrixes. J. Agric. Food

Chem. 66, 3302–3307. https://doi.org/10.1021/acs.jafc.8b00309

GNK, G., A, R., V, Suresh, V, Senthil, Baviya, P.R., 2018. Nutraceuticals - a Regulatory

Review. Int. J. Drug Regul. Aff. 3, 22–29. https://doi.org/10.22270/ijdra.v3i2.165

Guo, Z.L., Li, M.X., Li, X.L., Wang, P., Wang, W.G., Du, W.Z., Yang, Z.Q., Chen, S.F., Wu,

D., Tian, X.Y., 2022. Crocetin: A Systematic Review. Front. Pharmacol. 12, 1–28.

https://doi.org/10.3389/fphar.2021.745683

Hagaggi, N.S.A., Abdul-Raouf, U.M., 2023. Production of bioactive β-carotene by the

endophytic bacterium Citricoccus parietis AUCs with multiple in vitro biological potentials.

Microb. Cell Fact. 22, 1–9. https://doi.org/10.1186/s12934-023-02108-z

Hopwood, D., 2007. An introduction to the actinobacteria. Microbiol. Today 34, 60–62.

https://www.genome.jp/kegg/pathway.html, n.d. Kyoto Encyclopedia of Genes and Genomes

[WWW Document].

Kabiri, Y., Angelin, A., Ahmed, I., Mutlu, H., Bauer, J., Niemeyer, C.M., Zandbergen, H.,

Dekker, C., 2019. Intercalating Electron Dyes for TEM Visualization of DNA at the Single-

Molecule Level. ChemBioChem. https://doi.org/10.1002/cbic.201800638

Katata-Seru, L., Ramalapa, B., Tshweu, L., 2019. Nanoformulated Delivery Systems of Essential

Nutraceuticals and Their Applications. Nanoemulsions - Prop. Fabr. Appl. 1–12.

https://doi.org/10.5772/intechopen.86170

Katoh, T., Nagashima, U., Mimuro, M., 1991. Fluorescence properties of the allenic carotenoid

fucoxanthin: Implication for energy transfer in photosynthetic pigment systems. Photosynth.

Res. 27, 221–226. https://doi.org/10.1007/BF00035843

Kim, J.H., Kim, S.H., Yoon, J.H., Lee, P.C., 2014. Carotenoid production from n-alkanes with a

broad range of chain lengths by the novel species Gordonia ajoucoccus A2T. Appl.

Microbiol. Biotechnol. 98, 3759–3768. https://doi.org/10.1007/s00253-014-5516-y

Kirti, K., Amita, S., Priti, S., Mukesh Kumar, A., Jyoti, S., 2014. Colorful World of Microbes:

Carotenoids and Their Applications. Adv. Biol. https://doi.org/10.1155/2014/837891

Kłodawska, K., Bujas, A., Turos-Cabal, M., Żbik, P., Fu, P., Malec, P., 2019. Effect of growth

temperature on biosynthesis and accumulation of carotenoids in cyanobacterium Anabaena

sp. PCC 7120 under diazotrophic conditions. Microbiol. Res. 226, 34–40.

https://doi.org/10.1016/j.micres.2019.05.003

Kumari, S., Goyal, A., Garg, M., 2020. Phytochemistry and Pharmacological Update on

Tetraterpenoids. Nat. Prod. J. 11, 617–628.

https://doi.org/10.2174/2210315510999200806154609

Lakshman, M., 2019. Application of conventional electron microscopy in aquatic animal disease

diagnosis: A review. J. Entomol. Zool. Stud. 7, 470–475.

Lee, S.-A., Kim, M., Esterhuizen, M., Le, V. Van, Kang, M., Ko, S.-R., Oh, H.-M., Kim, Y.J.,

Ahn, C.-Y., 2022. An acceleration of carotenoid production and growth of Haematococcus

lacustris induced by host-microbiota network interaction. Microbiol. Res. 262.

Loh, W.L.C., Huang, K.C., Ng, H.S., Lan, J.C.W., 2020. Exploring the fermentation

characteristics of a newly isolated marine bacteria strain, Gordonia terrae TWRH01 for

carotenoids production. J. Biosci. Bioeng. 130, 187–194.

https://doi.org/10.1016/j.jbiosc.2020.03.007

López, G., Álvarez-rivera, G., Carazzone, C., Ibáñez, E., Leidy, C., Cifuentes, A., 2021.

Carotenoids in Bacteria◻: Biosynthesis , Extraction , Characterization and Applications.

https://doi.org/10.20944/preprints202108.0383.v1

Miller's Home Solvent polarity table [WWW Document], n.d. URL

http://johnhogan.info/hogan/CHEM_1201/Images/SolventPolarityTables.pdf

Miller, E.S., Mackinney, G., Zscheile, Jr., F.P., 1935. ABSORPTION SPECTRA OF ALPHA

AND BETA CAROTENES AND LYCOPENE. Plant phisiology 375–381.

Munekata, P.E.S., Pateiro, M., Barba, F.J., Dominguéz, R., Gagaoua, M., Lorenzo, J.M., 2020.

Development of new food and pharmaceutical products: Nutraceuticals and food additives.

Adv. Food Nutr. Res. 92, 53–96. https://doi.org/10.1016/bs.afnr.2019.12.001

Mussagy, C.U., Khan, S., Kot, A.M., 2021. Current developments on the application of

microbial carotenoids as an alternative to synthetic pigments. Crit. Rev. Food Sci. Nutr. 0,

1–15. https://doi.org/10.1080/10408398.2021.1908222

Narsing Rao, M.P., Xiao, M., Li, W.J., 2017. Fungal and bacterial pigments: Secondary

metabolites with wide applications. Front. Microbiol. 8, 1–13.

https://doi.org/10.3389/fmicb.2017.01113

Nouioui, I., Carro, L., García-López, M., Meier-Kolthoff, J.P., Woyke, T., Kyrpides, N.C.,

Pukall, R., Klenk, H.P., Goodfellow, M., Göker, M., 2018. Genome-based taxonomic classification of the phylum actinobacteria. Front. Microbiol. 9, 1–119. https://doi.org/10.3389/fmicb.2018.02007

Nupur, L.N.U., Vats, A., Dhanda, S.K., Raghava, G.P.S., Pinnaka, A.K., Kumar, A., 2016. ProCarDB: A database of bacterial carotenoids. BMC Microbiol. 16, 1–8. https://doi.org/10.1186/s12866-016-0715-6

Padmavathi, D., 2018. A general review on " Nutraceuticals ": Its golden health impact over human community. Int. J. Food Sci. Nutr. 3, 214-217A.

Parra-Rivero, O., de Barros, M.P., Prado, M. del M., Gil, J.V., Hornero-Méndez, D., Zacarías, L., Rodrigo, M.J., Limón, M.C., Avalos, J., 2020. Neurosporaxanthin overproduction by Fusarium fujikuroi and evaluation of its antioxidant properties. Antioxidants 9, 1–18. https://doi.org/10.3390/antiox9060528

Poddar, K., Padhan, B., Sarkar, D., Sarkar, A., 2021. Purification and optimization of pink pigment produced by newly isolated bacterial strain Enterobacter sp. PWN1.

Ram, S., Mitra, M., Shah, F., Tirkey, S.R., Mishra, S., 2020. Bacteria as an alternate biofactory for carotenoid production: A review of its applications, opportunities and challenges. J. Funct. Foods 67, 103867. https://doi.org/10.1016/j.jff.2020.103867

Sandmann, G., 2021. Carotenoid Biosynthesis in the Phylum Actinobacteria, in: Carotenoids: Biosynthetic and Biofunctional Approaches. pp. 175–181.

Silva, T.P., Paixão, S.M., Alves, L., 2016. Ability of: Gordonia alkanivorans strain 1B for high added value carotenoids production. RSC Adv. 6, 58055–58063.

https://doi.org/10.1039/c6ra08126f

Sowani, H., Kulkarni, M., Zinjarde, S., 2018. An insight into the ecology, diversity and

adaptations of Gordonia species. Crit. Rev. Microbiol. 44, 393–413.

https://doi.org/10.1080/1040841X.2017.1418286

Yabuzaki, J., 2017. Carotenoids Database: Structures, chemical fingerprints and distribution

among organisms. Database 2017, 1–11. https://doi.org/10.1093/database/bax004

Ye, X., Huang, Y., Zhou, Chenyu, Liu, X., Zhao, W., Zhao, X., Xie, X., Wang, L., Bai, Z., Zhou,

Changlin, Ma, L., 2022. MSI-1 combats drug-resistant S. aureus by affecting bacterial

viability and inhibiting carotenoid pigment production. Microbiol. Res. 255, 126909.

https://doi.org/10.1016/j.micres.2021.126909

Yurkov, V., Gad'on, N., Drews, G., 1993. The major part of polar carotenoids of the aerobic

bacteria Roseococcus thiosulfatophilus RB3 and Erythromicrobium ramosum E5 is not

bound to the bacteriochlorophyll a-complexes of the photosynthetic apparatus. Arch.

Microbiol. 160, 372–376. https://doi.org/10.1007/BF00252223

Zhao, D., Li, C., Zhang, N., Li, B., 2023. Integrative analysis of genomic and metabolomic data

reveals key metabolic pathways involved in lipid and carotenoid biosynthesis in oleaginous

red yeast Rhodosporidiobolus odoratus XQR. Microbiol. Res. 270.