Artificial intelligence-enabled automated analysis of transmission electron micrographs to evaluate chemotherapy impact on mitochondrial morphology in triple negative breast cancer

Argenis Arriojas[1,*], Lily M. Baek[2,*], Mariah J. Berner[2], Jiaqi Wang[3], Joseph Duraisingh[4], Alexander Zhurkevich[5], Antentor Othrell Hinton Jr.[6,7], Matthew D. Meyer[8], Lacey E. Dobrolecki[2], Michael T. Lewis[2,9,11], Kourosh Zarringhalam[1,3,#], Gloria V. Echeverria[2,9-11,#]

[1] Center for Personalized Cancer Therapies, University of Massachusetts Boston, Boston, MA, USA

[2] Lester and Sue Smith Breast Center and Department of Medicine, Baylor College of Medicine, Houston, TX, USA

[3] Department of Mathematics, University of Massachusetts Boston, Boston, MA, USA

[4] Department of Cell and Systems Biology, University of Toronto, Toronto, Ontario, Canada

[5] Department of Computer Science, University of Massachusetts Boston, Boston, MA, USA

[6] Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, USA

[7] Microscopy and Cell Analysis Core Facility, Mayo Clinic, Rochester, MN, USA

[8] Shared Equipment Authority, Rice University, Houston, TX, USA

[9] Dan L Duncan Comprehensive Cancer Center, Baylor College of Medicine. Houston, TX, USA

[10] Department of Medicine, Baylor College of Medicine, Houston, TX, USA.

[11] Departments of Molecular and Cellular Biology and Radiology. Baylor College of Medicine. Houston, TX, USA

* These authors contributed equally to this work

co-corresponding authors

DISCLOSURES

MTL is a founder and limited partner in StemMed Ltd. and a manager in StemMed Holdings, its general partner. He is a founder and equity stakeholder in Tvardi Therapeutics Inc. Some PDXs are exclusively licensed to StemMed Ltd., resulting in royalty income to MTL. LED is a compensated employee of StemMed Ltd. Some PDXs are exclusively licensed to StemMed Ltd. resulting in royalty income to LED. G.V.E. previously recieved sponsored research funding from Chimerix, Inc. G.V.E. receives

28 experimental compounds from the Lead Discovery Center of Germany and Jazz Pharmaceuticals, Inc.

29 All other authors have nothing to disclose.

ABSTRACT

Advancements in transmission electron microscopy (TEM) have enabled in-depth studies of

biological specimens, offering new avenues to large-scale imaging experiments with subcellular

resolution. Mitochondrial structure is of growing interest in cancer biology due to its crucial role in

regulating the multi-faceted functions of mitochondria. We and others have established the crucial role

of mitochondria in triple-negative breast cancer (TNBC), an aggressive subtype of breast cancer with

limited therapeutic options. Building upon our previous work demonstrating the regulatory role of

mitochondrial structure dynamics in the metabolic adaptations and survival of chemotherapy-refractory

TNBC cells, we sought to extend those findings to a large-scale analysis of transmission electron

micrographs. Here we present a U-Net artificial intelligence (AI) model for automatic annotation and

assessment of mitochondrial morphology and feature quantification. Our model is trained on 11,039

manually annotated mitochondria across 125 micrographs derived from a variety of orthotopic patient-

derived xenograft (PDX) mouse model tumors and adherent cell cultures. The model achieves an F1

score of 0.85 on test micrographs at the pixel level. To validate the ability of our model to detect

expected mitochondrial structural changes, we utilized micrographs from mouse primary skeletal

muscle cells genetically modified to lack Dynamin-related protein 1 (Drp1). The algorithm successfully

detected a significant increase in mitochondrial elongation, in alignment with the well-established role

of Drp1 as a driver of mitochondrial fission. Further, we subjected *in vitro* and *in vivo* TNBC models to

conventional chemotherapy treatments commonly used for clinical management of TNBC, including

doxorubicin, carboplatin, paclitaxel, and docetaxel (DTX). We found substantial within-sample

heterogeneity of mitochondrial structure in both *in vitro* and *in vivo* TNBC models and observed a

consistent reduction in mitochondrial elongation in DTX-treated specimens. We went on to compare

mammary tumors and matched lung metastases in a highly metastatic PDX model of TNBC, uncovering

significant increase in mitochondrial length in metastatic lesions compared to their cognate mammary tumor. This dataset provides high statistical power to detect frequent chemotherapy-induced shifts in mitochondrial shapes and sizes in residual cells left behind after treatment. The successful application of our AI model to capture mitochondrial structure marks a step forward in high-throughput analysis of mitochondrial structures, enhancing our understanding of how morphological changes may relate to chemotherapy efficacy and mechanism of action. Our large, manually curated electron micrograph dataset - now publicly available - serves as a unique gold-standard resource for developing, benchmarking, and applying computational models, while further advancing investigations into mitochondrial morphology and its impact on breast cancer biology.

INTRODUCTION

Mitochondria are multifaceted organelles responsible for energy production through oxidative phosphorylation (OXPHOS). This process generates ATP, the cell's primary energy currency, by utilizing electrons derived from nutrients[1]. Beyond energy production, mitochondria play critical roles in regulating various cellular processes, including apoptosis, cell cycle control, and cell signaling, all of which are dependent on their structural integrity[2-5].

Mitochondrial structure, regulated by their fusion and fission, is highly dynamic. These organelles are enclosed by a double membrane, with the inner membrane folded into cristae that house the OXPHOS machinery[6]. The balance between mitochondrial fusion and fission, governed by key proteins such as dynamin-related protein 1 (DRP1), optic atrophy 1 (OPA1), and mitofusins (MFNs), is crucial for maintaining mitochondrial function and cellular health[7]. In several diseases including cancer, alterations in mitochondrial structure, including those that favor either elongation or fragmentation, are common, contributing to dynamic metabolic reprogramming that supports tumor growth and survival[8-10]. Alterations in mitochondrial fusion and fission can lead to enhanced proliferation, metastasis, or

other hallmarks of cancer[9]. Thus, delineating these alterations is critical for gaining a deeper understanding of cancer biology.

Triple-negative breast cancer (TNBC) presents significant treatment challenges due to the absence of estrogen and progesterone receptors and human epidermal growth factor receptor two (HER2) amplification or overexpression[11]. Although neoadjuvant (a.k.a pre-surgical) chemotherapy (NACT) with PD-1 immune checkpoint blockers is the standard treatment, approximately half of patients do not achieve complete tumor eradication, leading to significantly elevated risk of metastasis and mortality[12-14]. In our previous studies, we found that after chemotherapy, residual patient-derived xenograft (PDX) tumors which eventually recurred underwent reversible transcriptional and mitochondrial adaptations, with notable increases in OXPHOS-related gene expression and activity relative to treatment-naive tumors[15]. This mitochondrial adaptation was mediated by elevation of the mitochondrial fusion protein OPA1. We also demonstrated that inhibiting OPA1 with the small molecule MYLS22[16] abated mitochondrial fusion, OXPHOS, and chemoresistance[8]. Targeting proteins like DRP1, OPA1, or MFNs could enhance the efficacy of existing treatments and overcome resistance mechanisms, offering new hope for patients with difficult-to-treat cancers[17-20].

Electron microscopy is the gold standard method to gain subcellular resolution of organelles. Due to the complexity of mitochondrial structure, accurately quantifying features in imaging data is crucial for understanding the connection between mitochondrial structure and function. Traditionally, analyzing mitochondrial structures involves manually tracing and counting individual mitochondria in electron micrographs to measure morphological parameters such as length, area, circularity, and linearity. This process becomes challenging when the contrast between mitochondria and other organelles or the cytosol is low. In addition, during manual inspection of mitochondria, the examiner often considers only a subset of mitochondria in each image, leading to a loss of information. Individual tumor cells can contain anywhere from a few dozen to several hundred mitochondria, making this manual procedure ineffective for statistically powered high-throughput analysis, as it is extremely time-

consuming and may be heavily dependent on examiner bias. These issues can lead to unreliable and inconsistent metrics for quantification, especially considering the extensive intra-tumoral heterogeneity characteristic of most cancers.

In recent years, advancements in artificial intelligence (AI), particularly deep machine learning, have transformed image analysis across various medical domains[21-23], including those focused on mitochondrial inner and outer membrane structure[24]. Deep learning models, especially convolutional neural networks, have shown significant success in tasks such as tumor classification, cancer subtype identification, mutation prediction, and treatment outcome prediction using histological images like hematoxylin and eosin (H&E) stained slides[25-27]. These approaches utilize either transfer learning, where pre-trained models are fine-tuned for specific tasks, or full-training strategies that train models from scratch using large domain-specific datasets. Such innovations have enabled the identification of subtle morphological and molecular patterns within images that were previously unnoticed through traditional methods. Similarly, AI-driven approaches have advanced mitochondrial structure studies, enabling precise quantification of mitochondrial dynamics and revealing intricate features like cristae topologies that were difficult to detect. Additionally, pixel-level segmentation of cancer images using autoencoder-based architectures has been adapted for high-resolution segmentation tasks across various image types[28-30]. Among these, U-Net architectures[31] have demonstrated particular efficacy in pixel-level cancer image analysis[25-27,32]. These methods offer automated segmentation and detection capabilities, facilitating more accurate, efficient, and unbiased quantification with exceptional statistical power.

In this study, we utilized *in vitro* (adherent cultures of human TNBC cell lines) and *in vivo* (orthotopic PDX mouse models of TNBC) models, in which we previously delineated mitochondrial features and chemotherapeutic responses[15], to conduct a large-scale assessment of mitochondrial morphology. Building on our studies that demonstrated a link between mitochondrial morphology, OXPHOS, and survival in post-chemotherapy residual TNBC cells[8,15], we sought to develop and

validate an AI-based approach capable of high-throughput segmentation and quantitative analysis of mitochondrial structures in TNBC. To this end, we first generated a large set of transmission electron micrographs from our models, then curated a high-quality dataset by manually annotating 11,039 mitochondria across 125 micrographs from the aforementioned models. These manual annotations served as training data for a U-Net image segmentation model, which was then applied to detect mitochondria at scale. Using computer-vision approaches, we systematically characterized multiple mitochondrial morphological features. This extensive analysis allowed us to statistically quantify sometimes subtle mitochondrial structural upon chemotherapy treatment with exceptional statistical rigor. Here, we describe the extensive training and evaluation of our model, demonstrate its robustness and versatility, and present findings that illuminate how mitochondrial structure adapts in TNBC.

RESULTS

Diverse training datasets to enable robust mitochondrial segmentation

Using our previously established methods to examine residual cells surviving exposure to conventional chemotherapies with *in vivo* and *in vitro* models of TNBC, we systematically collected cell and tumor specimens then subjected them to TEM. We then assembled a diverse training dataset by manually annotating mitochondria from 125 electron micrographs using QuPath[33]. These images ranged in calibrated pixel size from 0.36 to 23 nm/pixel and were collected from two cell culture models – primary mouse skeletal myotubes genetically engineered to lack the mitochondrial fission-driving gene *Drp1* (*Drp1* KO) and human TNBC cells (MDA-MB-231) as well as seven orthotopic patient-derived xenograft (PDX) TNBC models (PIM001-P, PIM005[15], PIM056[34], WHIM-14[35], BCM-2147[36], BCM-15116 (unpublished), and HCI-010[35], each derived from a unique TNBC patient. At an intermediate resolution of 7.5 nm/pixel, slides typically contained between 100 - 500 mitochondria. Data acquisition involved three separate microscopists, each employing unique microscope settings and

capturing various specimens. This diversity in magnification levels, microscope configurations, image

formats (including presence of scale bars), and biological systems, helped us build a model that learned

to recognize mitochondrial structures under varied conditions. This adaptability is critical for its intended

application, as it may allow for accurate predictions irrespective of resolution or experimental setup.

Model development overview

Figure 1 shows a schematic of our pipeline. Briefly, we began with preparation of training data

by manually annotated micrographs, followed by tiling at a common resolution (7.5 nm/pixel), then data

augmentation to enhance model robustness. A U-Net was trained, allowing the model to learn complex

features associated with mitochondrial morphology. The trained model was then utilized to make

predictions on new images, detecting and outlining mitochondrial structures. Finally, we performed

downstream analyses in two major steps: (1) computer vision tasks (e.g., contour detection, feature

characterization such as maximum Feret's diameter and area) and (2) statistical assessments (*e.g.*,

distribution measurements) to compare control versus experimental conditions. This streamlined

workflow laid the foundation for high-throughput, quantitative evaluation of mitochondrial morphology.

Model performance evaluation

We employed a 5-fold cross-validation approach, using images from each individual dataset as

well as an aggregate dataset consisting of mixed images across datasets ("Mixture") (Fig. 2). We

assessed F1 scores to determine the consistency and robustness of mitochondrial segmentation

performance (Fig. 3). The heatmap reveals variations in F1 scores for each dataset, with the PIM001-

P PDX model achieving the highest score on its corresponding test dataset (F1 = 0.86), while *Drp1* KO

and HCI-010 reached 0.84 and 0.72, respectively, indicating strong alignment between model

predictions and manual annotations (Fig. 3A). By contrast, the *Drp1* KO model shows lower

performance on other datasets, with scores of 0.53 on HCI-010 and 0.19 on PIM001-P, indicating that

dataset-specific characteristics can strongly influence robustness. Training on the Mixture attenuates

these differences and yields 0.80 on *Drp1* KO, 0.74 on HCI-010, and 0.84 on PIM001-P, providing

competitive performance with improved consistency across datasets. This trend is observed in the

evaluation of recall, where the Mixture model performs better than single models trained on individual

datasets. Given the class imbalance in this segmentation task, precision–recall (PR) curves are the

most appropriate evaluation. The PR AUC values are 0.92, 0.80, and 0.94 for the *Drp1* KO, HCI-010,

and PIM001-P models, respectively, which align closely with the F1 scores (Fig. 3B).

To further evaluate the model's predictive accuracy, we visually compared the model's

segmentation output with manual annotations for sample images across the datasets. Representative

images are shown where mitochondria identified by manual annotation (blue) and model prediction

(red) are overlaid (magenta) in the "composed" panels, highlighting regions of concordance and minor

discrepancies (Fig. 3C). Overall, the model successfully identified the majority of mitochondria, aligning

with quantitative results, and demonstrating its ability to reliably detect mitochondrial structure under

varied experimental conditions. Despite some challenges in specific datasets, these findings

underscore the model's broad applicability for large-scale mitochondrial morphology analysis,

particularly when dealing with mixed image sets.

Evaluating alternative model architectures

To explore whether other recently developed segmentation architectures could enhance

performance, we trained U-Net++[37] (Mixture++) and Mask R-CNN[38] (Mixture-RCNN) models using the

Mixture training dataset and compared it directly with the corresponding U-Net model. As shown in

Fig.3A, U-Net++ achieved similar F1 scores but consistently improved recall across all three datasets,

indicating a modest advantage in capturing mitochondria pixels that may be missed by the standard U-

Net. In contrast, Mask R-CNN shows substantially lower F1 on all test sets and mixed recall. Although

U-Net++ offered modest improvements in recall, these gains were not accompanied by meaningful

changes in F1 score. Given that our primary analyses were completed using the U-Net architecture

with consistent training, benchmarking, and validation across datasets, we opted to retain the U-Net

model for subsequent analyses.

Model benchmarking against MitoNet

We compared our dataset-specific U-Net models to the general purpose MitoNet-v1 model [39]

using pixel-level F1 score and recall metrics (Fig. 3A). Across all test datasets, our specialized models

outperformed MitoNet-v1, achieving higher F1 scores and recall in most cases. Performance was

especially improved for PIM001-P and HCI-010 datasets, where the score gap was the largest. These

results demonstrate that domain-specific training on TEM images from TNBC models substantially

enhanced segmentation accuracy over a general tool.

In vitro **cell models provide biological validations of the algorithm**

We sought biological validation of our model with two approaches. First, we genetically

manipulated a canonical mitochondrial structure regulator to test the AI's ability to discern an expected

structural difference. Specifically, we conducted TEM of murine skeletal primary cells that were either

wild type (WT) or engineered to lack the mitochondrial fission protein Drp1 (*Drp1* KO). Micrographs

were analyzed at a resolution of 7.5 nm/pixel. Our data demonstrated a statistically significant increase

in elongation (where a value of 1 represents a perfect circle and higher values indicate more elongated

shapes) in *Drp1* KO cells compared to WT (Fig. 4B), consistent with the well-established role of Drp1

in causing mitochondrial fission. Histograms (Fig. 4C) show a clear rightward shift in elongation for *Drp1*

KO relative to WT, with a broader high-elongation tail, consistent with increased mitochondrial

elongation. In contrast, the area distributions are largely overlapping and remain skewed toward small

objects, in line with the non-significant difference seen in the Fig. 4B. These findings highlight the ability

of our model to effectively capture and quantify expected mitochondrial morphological differences.

As a second biological validation, we examined micrographs from drug-treated samples for

which we had previously established mitochondrial structure alterations using a variety of methods and

functional characterizations[8]. We previously reported that different types of standard chemotherapies

impacted mitochondrial morphology of TNBC cells in distinct ways: DNA-damaging chemotherapies

induced mitochondrial elongation, while taxanes, which disrupt microtubule dynamics, usually reduced

mitochondrial elongation (Fig. 5A)[8]. Consistent with those findings, our model detected a significant

increase in elongation in MDA-MB-231 TNBC cells treated with DNA-damaging chemotherapies

doxorubicin (DXR) or carboplatin (CRB) compared to vehicle (VEH) (Fig. 5B). In contrast, treatment

with the taxanes paclitaxel (PTX) or docetaxel (DTX) resulted in decreased mitochondrial elongation

and area compared to VEH (Fig.5B). Together, these findings validate our model's ability to detect

expected biological changes and highlight its potential for broader discovery applications.

Mitochondrial structure changes upon chemotherapy treatment in PDX models

We next sought to investigate whether our approach could detect changes in mitochondrial

features, for which we did not have a priori knowledge, during clinically relevant biological transitions in

TNBC using orthotopic PDX models. These models are well-known to recapitulate metastatic[40] and

therapeutic response[8,15,41] phenotypes that match the originating patient tumors. We treated five unique

PDX models, representing five TNBC patients, with the conventional taxane agent DTX, often included

as a standard treatment for patients with TNBC. Tumors in the five models exhibited varying degrees

of responses to DTX (Fig. 6A), as expected based on heterogeneity of chemotherapy efficacy that is

common amongst TNBC patients. We collected mammary tumors from mice that were either untreated

(n=1-3 per group) or treated with DTX (n=1-3 per group). Residual DTX tumors were harvested 20-28

days after treatment initiation. Tumors were fixed and processed for TEM followed by automated

mitochondrial analysis using our algorithm. In three out of five PDX models, we observed decreased

mitochondrial elongation following DTX treatment, with BCM-2147 showing a significant reduction

(p<0.0001). This was accompanied by a decrease in mitochondrial area. Representative micrographs

illustrate these findings (Fig. 6C). Additionally, in tumors with greater baseline mitochondrial elongation,

we detected a more pronounced reduction in elongation after DTX treatment, suggesting that initial mitochondrial morphology may influence the extent of structural and functional adaptations they undergo (Fig.6D). Indeed, our prior study provided functional evidence that a transition towards mitochondrial fragmentation reduced OXPHOS in TNBC cells[8]. We did not observe any clear association between mitochondrial features and degree of DTX-induced change in tumor volume (Fig.6A-C).

Reproducible mitochondrial elongation in a PDX model of spontaneous lung metastasis

We used a highly metastatic PDX model, PIM056, derived from a treatment-naïve TNBC patient. This model develops multifocal macro-metastatic lung lesions within 16 weeks of mammary fat pad (MFP) engraftment without the need for survival tumor resection surgery. This rapidly aggressive metastatic phenotype is exceedingly rare amongst PDX models. Notably, a prior study using H&E and human-specific mitochondrial immunostaining confirmed that the lung metastases in the PIM056 model are predominantly composed of human tumor cells, rather than mouse stromal tissue, thereby validating the cellular identity of the regions analyzed by TEM[42]. Approximately 16 weeks post engraftment, when the mammary tumors measured on average 1000 mm^3, we collected matched MFP tumors and macro-dissected lung metastases from four individual replicate mice (Fig.7A). TEM followed by automated mitochondria detection revealed significantly longer mitochondria in lung metastases compared to mammary tumors (Fig. 7B-D). In HER2$^+$ breast cancer, brain metastases exhibit increased mitochondrial fission relative to MFP tumors[43]. In TNBC, several studies suggest a greater reliance on mitochondrial OXPHOS in lung metastases than at other metastatic sites[44] ; however, the contribution of mitochondrial dynamics to these differences remains unresolved. The functional relevance of mitochondrial length in TNBC metastasis will be an important topic for future investigations, and our findings highlight the importance of broadening this analysis to additional models of TNBC metastasis.

DISCUSSION

The application of AI is advancing the field of biomedical image analysis[45]. Manual annotation of

cancer images is labor-intensive, prone to human bias, and often inconsistent between analysts,

platforms, and institutions, making large-scale studies impractical. The AI-driven model presented here

addresses these limitations by automating the detection and characterization of mitochondrial

morphology with high accuracy and consistency. A key contribution of our AI model is its capacity to

capture and quantify the heterogeneity and dynamics of mitochondrial morphology in biologically

important transitions within cancer cells.

Chemotherapy-induced changes in mitochondrial structure are crucial indicators of cellular

adaptive mechanisms, including metabolic reprogramming and resistance to cell death. Notably, our

prior work demonstrated that genetically or pharmacologically inducing mitochondrial fission or fusion

was sufficient to modulate OXPHOS and chemotherapeutic responses[8]. By detecting subtle changes

in structural features across large datasets, our approach provides new insights into how mitochondria

may contribute to the survival of residual cancer cells post-treatment. We expect our model and highly

curated annotations may not only accelerate the analysis of large datasets but may also enable a level

of statistical rigor that is difficult to achieve with manual methods. Its robustness, stemming from training

on a diverse set of images, underscores its potential for broad applicability in various biological contexts.

To benefit the research community, we have made both our algorithm code and our extensive collection

of TEM images publicly available. This dataset includes 125 micrographs with a total of 11,039

mitochondrial annotations, which we anticipate will serve as a valuable resource for developing further

AI-driven analysis tools.

We observed substantial intra-tumor heterogeneity of mitochondrial shapes and sizes within

individual micrographs from both cell lines and PDX models (Supplementary Fig. 1). Although genomic,

transcriptomic, epigenomic, and proteomic heterogeneity are well-established phenomena across

diverse cancers, including within the cell models in this study, diversity of mitochondrial structure has

only recently begun to gain attention in cancer[46,47]. Our algorithm's ability to capture this heterogeneity

with high resolution, even within a decades-old, well-established human cell line MDA-MB-231, suggests an important, previously underappreciated layer of complexity in cancer cell biology. These observations raise compelling questions about how variations in mitochondrial morphology may correlate with functional differences or therapy responses, underscoring the need for deeper investigation into the potential role of mitochondrial heterogeneity in tumor progression.

Our study has several limitations. First, PDX models lack a fully intact immune system, which may be important for mitochondrial features of tumor cells. Consequently, extending our algorithm to immune-competent mouse tumor models will be crucial. Second, the generalizability of our observed chemotherapy-induced mitochondrial structure changes should be further assessed in additional *in vitro* and *in vivo* models with additional conventional chemotherapies, and ideally in human tumor specimens. This is especially important given the well-established inter-patient heterogeneity of TNBC[48-50]. Our algorithm is precisely designed to enable such high-throughput analyses on large sample sets. Third, TEM provides only a two-dimensional cross section of mitochondria and is unable to capture their full three-dimensional complexity[47]. We hypothesize that analyzing high numbers of mitochondria, representing a random sampling of cells' cross-sections, as presented in this study, may mitigate this limitation by approximating the "average" morphological properties within each sample.

In our metastasis PDX study, we observed altered mitochondrial elongation reproducibly across four replicate mice when comparing their lung metastases to their cognate MFP tumor. These findings are in line with our prior report of the reproducible clonal bottleneck during the process of metastasis in similar TNBC PDX models, and we conducted analogous experiments in biological replicate mice from that model to uncover reproducible alterations in clonal architecture[40]. The fact that mitochondrial changes arose repeatedly in independent mice suggests potential functional relevance, but that must be rigorously tested in future perturbation experiments in PIM056 and in additional models of metastatic TNBC.

We acknowledge that TEM, while providing the gold standard for subcellular resolution of

mitochondrial structure, involves intrinsically complex sampling and processing procedures. These

include specialized aldehyde fixation, osmium tetroxide post-fixation, resin embedding, ultramicrotomy,

and heavy metal counterstaining; procedures that require significantly longer processing times, higher

costs, and more advanced technical expertise compared to conventional light microscopy. Despite

these practical constraints, TEM remains essential for assessing mitochondrial morphological features

that cannot be resolved with light microscopy. While emerging technologies like focused ion beam

scanning electron microscopy (FIB-SEM) enable three-dimensional reconstruction of mitochondrial

cristae at nanometer resolution, their application remains even more limited, with single-cell

acquisitions requiring days to weeks of imaging and processing[51-53], restricting throughput to dozens of

mitochondria across only a few cells per experiment. This is almost certainly insufficient for capturing

the heterogeneity of cancer cell populations. Our AI-driven approach maximizes the value extracted

from TEM images by enabling rapid, unbiased analysis of thousands of mitochondria across our

manually curated dataset of 125 micrographs, thereby overcoming the traditional bottleneck of manual

annotation while maintaining the resolution necessary to detect morphological changes.

While our current analysis focused exclusively on mitochondrial morphology, a more holistic

understanding of the chemotherapy-induced cellular changes will require analyzing additional

organelles, such as the nucleus, cell membrane, lipid droplets, and endoplasmic reticulum, that can

have important interactions with mitochondria. These organelles play significant roles in cellular

metabolism, signaling, and homeostasis in concert with mitochondria and may be important

contributors to therapeutic resistance in cancer.

Several high-quality datasets and segmentation frameworks have advanced the field of

mitochondrial and organelle image analysis, including MitoNet, MitoEM, OpenOrganelle, CREMI, and

Lucchi[39,54-58]. Each of these resources serves unique purposes. MitoNet provides a generalizable deep-

learning framework for mitochondria segmentation across diverse EM datasets. MitoEM and

OpenOrganelle focus on large-scale 3D connectomics data, enabling volumetric reconstruction and detailed mitochondrial network analysis. Meanwhile, the CREMI and Lucchi datasets, though smaller, have served as important benchmarks for EM segmentation methods, often targeting synaptic or nuclear structures alongside mitochondria. Although these tools are invaluable for general benchmarking, they differ substantially from our target application. Most are designed for broad organelle segmentation or 3D volume analysis, typically using isotropic serial-section or FIB-SEM data from non-cancer tissues. By contrast, our pipeline is purpose-built for 2D TEM of TNBC models, with an emphasis on pixel-level morphological quantification of thousands of mitochondria per experiment, enabling the detection of subtle but potentially relevant structural shifts, such as chemotherapy-induced fragmentation or elongation, across large and heterogeneous datasets.

Our rationale for developing a dedicated TNBC-TEM segmentation framework rests on two key points. First, domain-specific optimization was essential, as benchmarking revealed that general-purpose models often underperform on TNBC TEM data due to differences in tissue type, imaging conditions, and structural complexity. Second, our workflow integrates segmentation with downstream morphometric analysis, allowing rigorous statistical comparisons across biological conditions rather than producing masks alone. In this way, our approach complements rather than replaces generalist models, offering a specialized, high-throughput solution for quantitative mitochondrial morphology analysis in clinically relevant cancer models.

We must note that our models were trained exclusively on 2D TEM images from TNBC PDX samples and cell lines, which inherently limits their generalizability. The models are not necessarily expected to produce accurate segmentation predictions for images outside this training domain, such as different cancer types, tissue types, or imaging modalities. Therefore, researchers applying our model to new biological contexts should carefully validate the predictions through manual review, and may need to perform additional fine-tuning or retraining with domain-specific annotations to achieve optimal performance.

FIGURE LEGENDS

Figure 1. Schematic overview of the U-Net AI pipeline for mitochondrial segmentation and analysis. The workflow begins with the preparation of training data, including transmission electron micrographs and manually annotated masks of mitochondria. Images are then rescaled to a common resolution (7.5nm/pixel) and divided into smaller tiles for input into the neural network. Data augmentation, such as rotations, flips, and brightness adjustments, enhances model robustness to image variability. The U-Net architecture, consisting of contracting and expanding paths, is trained to generate segmentation maps from the input image tiles. Following training, the model is used to make predictions on new images, with segmentation masks stitched together for full-image analysis. Downstream analysis includes computer vision tasks for feature quantification and statistical evaluations of mitochondrial morphology across experimental conditions.

Figure 2. Evaluation of model performance using 5-fold cross-validation across datasets. The datasets include *Drp1* KO, HCI-010, PIM001-P, and a mixed dataset combining samples from all groups. The cross-validation approach divided each dataset into five groups, with one group used for testing and the remaining groups for training in each fold. Line plots on the right display F1 scores for each fold during training, demonstrating convergence and model performance across datasets. Consistent F1 score trends across folds indicate reliable segmentation performance, with variations reflecting dataset-specific characteristics. The mixed dataset exhibited robust generalizability, capturing diverse image features and achieving high performance.

Figure 3. Evaluation of segmentation performance and visual comparison of model predictions. (A) Heatmaps showing F1 scores (left) and recall scores (right) for mitochondrial segmentation models trained on individual datasets (*Drp1* KO, HCI-010, PIM001-P), a pooled dataset (Mixture), or enhanced training strategies (Mixture++, Mixture-RCNN), compared against the benchmark model MitoNet-v1.

Each row represents the test dataset, and each column represents the training dataset or model. Higher values (darker blue) indicate better performance. (B) Precision–recall curves for models trained on individual datasets (*Drp1* KO, HCI-010, PIM001-P) when tested on their respective datasets. The area under the curve (AUC) is indicated for each model. (C) Representative TEM images showing raw micrographs (left), manual annotations, and mitochondrial segmentation outputs from models trained on individual datasets or mixture-based approaches. Each row represents a different magnification (top: whole cell, middle: subcellular region, bottom: field with multiple cells). Segmented mitochondria are pseudocolored for visualization: manual annotations (blue), model predictions (red), and overlapping regions (magenta).

Figure 4. Genetic validation of model performance.

(A) Representative micrographs from wild-type (WT) and *Drp1* knockout (KO) murine skeletal muscle cells. Images are shown as raw (left) or with model-predicted mitochondrial contours overlaid in red (right). Scale bars, 0.2 µm. (B) Quantification of mitochondrial elongation (left) and area (μm^2) (right) in WT and *Drp1* KO cells. Boxplots display the mean (black dot), median (horizontal line), interquartile range (box), and whiskers extending to 15–85% anchors, ±1.2×IQR. Sample sizes are indicated inside boxes. Data points outside these bounds were excluded as outliers. ****$P < 0.0001$, ns= not significant by pairwise Wilcoxon test. (C) Distribution histograms of mitochondrial elongation (left) and area (right) for WT and *Drp1* KO groups, shown as percent of group.

Figure 5. Pharmacologic validation of model performance. (A) Representative micrographs from MDA-MB-231 cells were treated with VEH, DXR, CRB, PTX, or DTX. Images are shown as raw (left) or with model-predicted mitochondrial contours overlaid in red (right). (B) Quantification of mitochondrial elongation (left) and area (μm^2) (right). Boxplots display the mean (black dot), median (horizontal line), interquartile range (box), and whiskers extending to 15–85% anchors, ±1.2×IQR. Sample sizes are indicated inside boxes. Data points outside these bounds were excluded as outliers. ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$, *$P < 0.05$, ns= not significant by pairwise Wilcoxon test.

Figure 6. Analysis of mitochondrial changes in PDX models treated with DTX. (A) Line plots showing normalized tumor volume over time in five PDX models treated with DTX (orange) (starting on Day 0) compared to untreated controls (UNT, blue). Red arrows on the x-axis indicate each DTX administration (*i.p.*, 20mg/kg). (B-C) Box plots summarize mitochondrial elongation (B) and area (C) across the five PDX models. Boxplots display the mean (black dot), median (horizontal line), interquartile range (box), and whiskers extending to 15–85% anchors, ±1.2×IQR. Sample sizes are indicated inside boxes. Data points outside these bounds were excluded as outliers. ****P < 0.0001, ***P < 0.001, **P < 0.01, ns= not significant by pairwise Wilcoxon test. (C) Representative micrographs from BCM-2147 untreated (top) or treated with DTX (bottom). Images are shown as raw (left) or with model-predicted mitochondrial contours overlaid in red (right). Scale bars, 0.2 μm. (E) Scatter plot showing baseline elongation in the untreated (UNT) group (x-axis) vs the percent change (Δ%) in elongation after DTX (y-axis), defined as $(DTX_{mean}-UNT_{mean})/UNT_{mean}$ x 100. Each point represents one model. Colors correspond to the legend. The red dashed line indicates the linear regression fit (P=0.14), and the black dashed line marks no change.

Figure 7. Mitochondrial morphology differences between mammary tumors and lung metastases (A) Schematic representation of the PIM056 PDX model, where tumors were allowed to grow for 16 weeks before isolation of both primary mammary fat pad (MFP) tumors and lung metastases (Lung) for analysis. (B) Representative TEM images show mitochondrial structures in MFP (top) and lung metastases (bottom). Images are shown as raw (left) or with model-predicted mitochondrial contours overlaid in red (right). Scale bars=0.2 μm. (C) Quantification of mitochondrial elongation (top) and area (μm^2) (bottom) in MFP and lung metastases. Boxplots display the mean (black dot), median (horizontal line), interquartile range (box), and whiskers extending to 15–85% anchors, ±1.2×IQR with sample sizes indicated for each group. Data points outside these bounds were excluded as outliers. ****P < 0.0001 by pairwise Wilcoxon test. (D) Distribution histograms of mitochondrial elongation (top) and area (bottom) for MFP and lung metastases, shown as percent of group.

Supplementary Figure 1. Mitochondrial elongation and area distributions reveal model-specific differences in morphology and heterogeneity under control conditions.

(A) Histograms of mitochondrial elongation and area across control samples (untreated conditions) show distinct distribution patterns across TNBC models. (B) Density ridge plots provide an overview of distribution shapes. (C) Heterogeneity in mitochondrial morphology was quantified using the coefficient of variation (CV) and interquartile range (IQR) for elongation and area.

METHODS

TNBC cell culture and drug treatment

The MDA-MB-231 cell line was purchased directly from ATCC (American Type Culture Collection) and cultured in RPMI-1640 (Gibco, 11879020) supplemented with 5 mM glucose (Gibco, A2494001), 10% fetal bovine serum (FBS) (R&D, S11550), and 1× antibiotic-antimycotic solution (Corning, 30-004-CI). Except for carboplatin (solutions were freshly prepared from powder), we prepared stock solutions of 10 mM in DMSO, and further dilutions were prepared in a proper growth medium. Cells were seeded at ~20% confluence and maintained for 48 h before treatment. Cells were then treated with: doxorubicin (final conc. at 100 nM, Sigma-Aldrich, 44583, protected from light), carboplatin (final conc. at 100 μM, Selleck Chemicals, S1215), paclitaxel (final conc. at 10 nM, Selleck Chemicals, Houston TX, S1150) or docetaxel (final conc. at 5 nM, Selleck Chemicals, Houston TX, S1148) for 48 hours before fixation.

Authentication of cell lines and PDX models

Cell cultures were tested for mycoplasma contamination each quarter using PCR with the Universal Mycoplasma Detection Kit (ATCC, 30-1012K), and were confirmed to be mycoplasma-free throughout the duration of this study. Short-tandem repeat (STR) DNA fingerprinting was performed on

DNA extracted from both cell lines and PDX model tumors by the Cytogenetics and Cell Authentication Core (CCAC) at M.D. Anderson Cancer Center. The Promega 16 High Sensitivity STR Kit (Catalog # DC2100) was used for the fingerprinting analysis, and the resulting profiles were compared against online search databases (DSMZ, ATCC, JCRB, RIKEN) for authentication.

Animal studies

This study was carried out in accordance with the *Guide for the Care and Use of Laboratory Animals* from the National Institutes of Health (NIH) IACUC. The protocol was approved by the IACUC at BCM (protocol AN-8243 and AN-2289). Mice were euthanized when they reached defined study or ethical end points. Euthanasia was conducted as recommended by the Association for Assessment and Accreditation of Laboratory Animal Care International. PIM001-P, PIM005[15] and PIM056[34] were propagated as previously described[1]. Briefly, cryo-preserved PDX cell suspensions were quickly thawed, then washed with Dulbecco's modified Eagle's medium (DMEM):F12 (Cytiva HyClone, SH30023.01) supplemented with 5% FBS. Viable cells were counted by staining with AOPI dye (Nexcelom Bioscience, CS2-0106) on a Cellometer K2 (Nexcelom Bioscience). For injection into mammary glands, 0.5-1.0 million viable tumor cells were suspended in a total of 20 µl (1:1 volume mixture of medium and Matrigel (Corning, 354234). Suspensions were then immediately injected unilaterally into the fourth mammary fat pads of 5 to 8-week-old female NOD/SCID [NOD.CB17-Prkdc^{scid}/NcrCrl, Charles River, National Cancer Institute (NCI) Colony] or NRG [NOD-Rag1null IL2rgnull] mice (Envigo). For HCI-010, BCM-2147, BCM-15116, and WHIM-14, cryo-preserved tumor fragments (1-2mm^3) were quickly thawed and washed with DMEM. The tumor chunks were orthotopically xenografted to 4th mammary gland of 6-8-week-old SCID/beige female mice.

Chemotherapy treatment of PDX models was conducted as previously described. Docetaxel solutions were administered by intraperitoneal injection at 20 mg/kg.

For PIM056 metastasis studies, lung metastases and mammary tumors were collected

immediately following euthanasia when primary tumors reached approximately 1200 mm³. Macro-

metastatic lesions visual to the naked eye were isolated with scalpels then preserved for TEM as below.

To generate *Drp1* KO skeletal muscle murine myotubes, mouse husbandry was carried out in

accordance with standard protocols[59] approved by the University of Iowa IACUC. Male C57Bl/6J mice

were housed at 22 °C with a 12 h light, 12 h dark cycle and free access to water and standard chow.

Controls represent pure C57Bl/6J genetic background mice. In accordance with previous studies[60-62],

Tamoxifen-inducible, skeletal muscle-specific *Drp1* KO mice were generated by crossing mice carrying

a homozygous floxed allele of *Drp1* with mice carrying a tamoxifen-inducible Cre recombinase under

the control of the skeletal muscle-specific myogenin promoter (Jackson Lab).

Isolation of satellite cells and differentiation

To collect tissue samples from wild-type or *Drp1* KO mice, animals were anesthetized using

isoflurane at 8–10 weeks old. Skeletal muscles from the gastrocnemius and quadriceps were excised

and washed twice with 1x phosphate-buffered saline (PBS) supplemented with 1% penicillin-

streptomycin and 0.3% fungizone (300 µL/100 mL The muscles were then incubated in Dulbecco's

Modified Eagle's Medium (DMEM)-F12 containing 0.2% collagenase II (2 mg/mL), 1% penicillin-

streptomycin, and 0.3% fungizone (300 µL/100 mL). Samples were shaken for 90 minutes at 37 °C.

Following incubation, the media was removed, and the muscle tissue was washed four times with PBS.

The media was then replaced with DMEM-F12 containing 0.05% collagenase II (0.5 mg/mL), 1%

penicillin-streptomycin, and 0.3% fungizone (300 µL/100 mL), followed by shaking for 30 minutes at

37 °C. Post-digestion, the tissue was ground until cells were dislodged from the tissue matrix and then

passed through a 70 µm cell strainer. The isolated cells were centrifuged, resuspended, and plated on

BD Matrigel-coated dishes. Adherent cells were differentiated into myotubes by adding DMEM-F12, 20%

FBS, 0.004% (40 ng mL-1) basic fibroblast growth factor (R&D Systems, 233-FB/CF), 1× non-essential amino acids, 0.14×10^3 m β-mercaptoethanol, 1x penicillin/streptomycin, and 0.3% fungizone (300 µL/100 mL). Myotubes were maintained in a medium containing 0.001% (10 ng/mL) growth factor until reaching 85% confluency, then were differentiated in DMEM-F12, 2% FBS, and 1× insulin-transferrin-selenium.

Transmission electron microscopy (TEM)

Tumor cells were fixed, processed, and embedded in situ to prevent morphological damage, as follows. Cells were fixed for 24 hours in Karnovsky's fixative, post-fixed in 1% osmium tetroxide, dehydrated in graded ethanol, and embedded in EMbed 812 resin. Ultra-thin sections were cut using a Leica UC7 ultramicrotome, stained with saturated methanolic uranyl acetate and Reynold's lead citrate, and imaged using a JEOL JEM-1230 TEM equipped with an AMT NanoSprint15 sCMOS camera. For tumor tissues, immediately following ethical animal euthanasia, mammary tumors were resected from mice and a ~1 mm-tick tumor slice was placed in Trump's fixation solution and fixed for 24 hours at room temperature with rocking. The necrotic areas that usually found in the core region of solid tumors were avoided. The initial fixed samples then were post-fixed in 1% buffered osmium tetroxide, dehydrated in graded ethanol, and embedded in Polybed 812 resin. Ultra-thin sections were obtained using a Leica UC 7 and sections were stained in 1% aqueous uranyl acetate and Reynold's lead citrate. Images were obtained using a FEI Tecnai Spirit TEM equipped with an Eagle camera and FEI Tia image acquisition software. Approximately 65 images at magnifications of 400-5000x were taken per tumor sample.

Myotubes were processed for TEM in accordance with previous studies[63,64]. For 1 h, cells were fixed by incubating at 37 °C with 2.5% glutaraldehyde in 0.1 m sodium cacodylate buffer. After rinsing

twice with 0.1 m sodium cacodylate buffer, samples were fixed again at room temperature for 30 min

to 1 h using 1% osmium tetroxide and 1.5% potassium ferrocyanide in 0.1 m sodium cacodylate buffer.

After secondary fixation, samples were washed for 5 min with 0.1 m sodium cacodylate buffer (7.3 pH).

From there, two washings of 5 min with diH_2O ensured the plates were cleaned. While keeping all

solutions and plates at room temperature, the samples were incubated with 2.5% uranyl acetate, diluted

with H_2O, at 4 °C overnight. Following this, samples were dehydrated using an ethanol gradient series.

After dehydration, the ethanol was replaced with Eponate 12 mixed in 100% ethanol in a 1:1 solution,

then incubated at room temperature for 30 min. This was repeated three times for 1 h using 100%

Eponate 12. The plates were finally placed in new media and cured in an oven at 70 °C overnight.

Plates were cracked upon hardening, and the cells were separated by submerging the plate in

liquid nitrogen. An 80 nm thickness jeweler's saw was used to cut the block to fit in a Leica UC6

ultramicrotome sample holder. From there, the section was placed on formvar-coated copper grids.

These grids were counterstained in 2% uranyl acetate for 2 min. Then these grids were counterstained

by Reynold's lead citrate for 2 min. Images were acquired by TEM on either a JEOL JEM-1230,

operating at 120 kV, or a JEOL 1400, operating at 80 kV.

Manual annotation of micrographs

For each tumor or cell line, 50-60 transmission electron micrographs were obtained at different

magnifications, ranging from 1000X to 5000X magnification. These micrographs were then loaded into

the QuPath software[33], a versatile tool for bioimage analysis. Each mitochondrion within the

micrographs was manually annotated using the drawing tool provided by QuPath. This process involved

carefully outlining the boundaries of individual mitochondria to ensure precise and accurate

characterization of their morphological features. The manual annotations were subsequently used for

further analysis, providing a detailed dataset for evaluating mitochondrial structure across various tumor

samples.

Model

A U-Net model was utilized to perform semantic segmentation on the TEM images. This deep-

learning model is an autoencoder that consists of a feature extraction stage (contracting path, encoder)

and a reconstruction stage (expansive path, decoder). The model uses a series of convolutional and

max pooling layers to transform the input image into abstract features in a latent space, that will then

be decoded in the reconstruction stage to match the model output with the corresponding training mask.

A defining characteristic of a U-Net model is the use of feature maps from the contracting path, as input

features in the expansive path. This choice helps to localize and assemble a more precise output

mask[31]. The network architecture is described in the following. The 2D input layer is followed by a

Gaussian Noise layer with standard deviation of 0.01. Each convolutional layer has kernel size 3 and

is followed by a leaky ReLu activation layer, a 2D spatial dropout layer (rates 0.2 and 0.1 in the

contracting and expansive paths respectively), and a batch normalization layer. A convolutional block

is composed of two such convolutional layers, followed by either a 2 by 2 max pooling layer in the

contracting path, or a 2 by 2 up-convolution layer with stride 2 in the expanding path. The model is

composed of 9 convolutional blocks in total: 4 in the contracting path, 1 in the bottleneck, and 4 in the

expansive path. The first convolutional block uses 16 filters in each of its convolutional layers. The

number of filters used in the convolutional layers doubles after each max pooling layer, and halves after

each up-convolution layer. The full model has just under 2 million trainable parameters.

Datasets

Three different datasets were used for training four U-Net models. The datasets are labeled as *Drp1* KO, HCI-010, and PIM001-P, with 26, 21, and 37 images respectively. The *Drp1* KO dataset contains images from the control set of an *in vitro* genetically engineered mouse myotube model. HCI-010 and PIM001-P contain images from two corresponding *in vivo* PDX mouse models. All images in the training datasets correspond to non-perturbed (control/vehicle) samples. For each dataset, a model was trained. A mixed model was also trained using images from these three datasets, along with control images of 3 other datasets (BCM-2147, MDA-MB-231, and PIM005), using a total of 125 images for training.

Preprocessing training data

The training dataset consists of electron micrographs, each paired with a manually annotated segmentation mask indicating the mitochondria. Magnification information was parsed from the image to obtain the calibrated pixel size. Prior to training or inference, images were scaled to match a common pixel size of 7.5 nm/pixel. Each image-mask pair was tiled to appropriately sized input samples that fit the model architecture shape (444 by 444 in the input layer, and 260 by 260 pixels at the output). The tiled process was performed in an overlapping fashion with a stride of 130 pixels in both horizontal and vertical directions. To avoid losing the pixels at the border, the images were zero-padded. The pixel values are rescaled to lie in the range [0,1]. The tile sets were converted into TF records to improve training computational performance. The final output was assembled by stitching the overlapping predicted mask tiles and taking the average for the overlapping regions.

Data augmentation

During model training, the input tiles were randomly transformed by rotating and flipping the images. This step was performed to increase the number of training samples and to avoid orientation

bias. Moreover, the brightness, contrast and gamma image parameters were randomly adjusted to

account for variation of these parameters across the input images. Additionally, images were randomly

magnified to boost the robustness of the model to input image resolution.

Model training

The model was trained on Nvidia DGX-A100 GPUs, for a total of 1200 epochs and batch size

128. The Adam optimizer was used with the default parameters, and a weighted binary cross entropy

loss function, assigning double weight to positive mask pixels. Model checkpoints were saved for the

best achieved loss on the validation sets. The convergence of the model was monitored using the

corresponding loss metric, as well as the IOU and DICE coefficients.

Performance evaluation

The data was split into a training set and a test set (85% and 15% approximately). A 5-fold cross

validation was used in the training set, where each fold uses 4 subsets for training and the remaining

5th set as validation. Model performance was assessed using the left-out test set.

Model predictions

Each of the 5 resulting models from the cross-validation procedure were used to make

predictions on new images. An ensemble approach was used to aggregate the results in a single

prediction, by taking the average of all predictions. Since the semantic segmentation models are

optimized for analysis at a calibrated pixel size of 7.5 nm/pixel, input images are scaled to this

magnification prior inference, and predictions are scaled back to the original size.

Assesing alternative model architectures

Our U-Net implementation was extended to the more recent U-Net++ architecture to evaluate whether its enhanced skip connections could improve segmentation performance. The model was constructed using the same parameters as the standard U-Net, resulting in approximately 2.4 million parameters in total, a ~25% increase compared to the original architecture.

We also evaluated a general-purpose instance segmentation framework, Detectron2's COCO-pretrained Mask R-CNN (mask_rcnn_R_50_FPN_3x, Meta AI), to assess whether this architecture could offer improvements over our U-Net-based models. For compatibility, the ground truth annotations from our Mixture training dataset were prepared following Detectron2's requirements, converted from semantic masks into COCO-style JSON format containing per-instance metadata such as contour coordinates and instance IDs, and registered in Detectron2 as a new local dataset. Model inference was performed via a tile-wise pipeline (512 by 512 px tiles with 20% overlap), followed by stitching to full resolution and rasterizing instance predictions into semantic masks. Despite its versatility, the Mask R-CNN baseline did not outperform our domain-specific U-Net models on the test datasets.

Benchmarks

To benchmark our model's performance, we compared it against MitoNet_v1[39], a widely-used mitochondria segmentation tool. The comparison was performed using the Empanada wrapper (v1.2) integrated with Napari (v0.4.18) in a Python 3.9 conda environment. We applied MitoNet's 2D inference tool with default parameters to 13 test images from our dataset, generating instance segmentation masks for mitochondria in each image. To enable direct comparison with our U-Net predictions, the instance segmentation masks were converted to semantic segmentation format by flattening individual mitochondria masks using Napari's "convert to image" feature. Model performance was quantified using pixel-level F1 score and recall metrics.

Postprocessing for morphometric analysis

For morphological measurements, predicted binary masks were first smoothed to reduce single-pixel noise and produce cleaner object boundaries. This was achieved by repeated convolution with a normalized 2D Gaussian kernel (sigma = 5 nm in image space) for 24 iterations, followed by by re-thresholding to recover a binary mask. This smoothing step was applied uniformly to all predictions prior to morphometric quantification, and was not used in the calculation of segmentation performance metrics.

Statistical analysis

For each group, data are summarized as the median, mean (black dot in boxplots), interquartile range (IQR), and whiskers spanning the 15th–85th percentiles with ±1.2×IQR, unless otherwise noted. Differences between groups were assessed using the Wilcoxon rank-sum test (two-sided) because the distributions of elongation and area are non-Gaussian, highly skewed, and contain unequal variances and outlier structures.

Data and code availability

Cell line models of TNBC are available for purchase from ATCC. PDX models can be made available to investigators through a material transfer agreement with the University of Texas MD Anderson Cancer Center (PIM001-P, PIM005 and PIM056) or Baylor College of Medicine (HCI-010, BCM-2147, BCM-15116, and WHIM-14). The Python code will be made available on GitHub (DOI pending). Electron micrographs and annotated mitochondrial masks are deposited in Bioimage Archive[65] under accession # S-BIAD2271.

AUTHOR CONTRIBUTIONS

A.A., L.M.B., K.Z., and G.V.E were responsible for overall completion of the study, writing the manuscript, and preparation of figures.

L.M.B., K.Z., and G.V.E were responsible for project conceptualization.

L.M.B. conducted PDX and cell line experiments, generated electron micrographs, conducted manual annotations of mitochondria, and data analyses under the supervision of G.V.E.

A.A. conducted computational algorithm development, optimization, and testing under the supervision of K.Z.

M.J.B. conducted PDX chemotherapy trials and generated electron micrographs analyzed in this study under the supervision of G.V.E.

A.Z. conducted initial algorithm development under the supervision of K.Z.

J.W. and J.D. contributed to the benchmarking efforts. J.W. performed training and evaluation for the Mixture-RCNN model. J.D. performed model evaluations for the MitoNet-v1 segmentation model.

A.O.H. provided electron micrographs of WT and *Drp1*KO myotubes.

Jiaqi Wang[3], Joseph Duraisingh

JW implemented the Mask-RCNN model and performed benchmark analysis.

JD implemented MitoNet and Empanada benchmakrs.

M.D.M. aided in design of electron microscopy imaging and generated most images in this study.

L.E.D. conducted PDX chemotherapy trials and provided tissues for electron microscopy under the supervision of M.T.L.

ACKNOWLEDGEMENTS

We are grateful to the breast cancer patients who donated their biopsies for cell line and PDX model generation. We are grateful to Mrs. Janice Cowden for providing advocacy support for our research. Dr. James P. Barrish assisted with TEM. Dr. Junegoo Lee assisted with mouse experiments. Dr. Helen Piwnica-Worms provided the PIM001-P and PIM056 PDXs through MTA with the University of Texas MD Anderson Cancer Center, and model generation was supported by a generous gift from the Cazalot Foundation through the MD Anderson Women's Cancer Moonshot Program. STR DNA fingerprinting was done by the Cytogenetics and Cell Authentication Core at M.D. Anderson Cancer Center.

FUNDING

G.V.E. is a Cancer Prevention Research Institute of Texas (CPRIT) Scholar in Cancer Research. The authors are supported by CPRIT RR200009 to G.V.E., National Institutes of Health (NIH) R37CA269783-01A1 to G.V.E., R37CA269783-01A1 to K.Z., R01AI167570 to K.Z., American Cancer Society RSG-22-093-01-CCB to G.V.E, National Science Foundation 2140736 to M.J.B., and Baylor Research Advocates for Student Scholars (BRASS) Myra Branum Wilson Scholarship to M.J.B. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH, NSF, CPRIT, ACS, or BRASS. The BCM PDX core is supported by P30 Cancer Center Support Grant NCI-CA125123, CPRIT Core Facilities Support Grant RP220646.

REFERENCES CITED

1   Anderson, A. J., Jackson, T. D., Stroud, D. A. & Stojanovski, D. Mitochondria-hubs for regulating
cellular biochemistry: emerging concepts and networks. *Open Biol* **9**, 190126,
doi:10.1098/rsob.190126 (2019).

2   Rossmann, M. P., Dubois, S. M., Agarwal, S. & Zon, L. I. Mitochondrial function in development
and disease. *Dis Model Mech* **14**, doi:10.1242/dmm.048912 (2021).

3   Nguyen, T. T. *et al.* Mitochondria-associated programmed cell death as a therapeutic target for
age-related disease. *Exp Mol Med* **55**, 1595-1619, doi:10.1038/s12276-023-01046-5 (2023).

4   Mitra, K., Wunder, C., Roysam, B., Lin, G. & Lippincott-Schwartz, J. A hyperfused mitochondrial
state achieved at G1-S regulates cyclin E buildup and entry into S phase. *Proc Natl Acad Sci U
S A* **106**, 11960-11965, doi:10.1073/pnas.0904875106 (2009).

5   Egan, G. *et al.* Mitochondrial and Metabolic Pathways Regulate Nuclear Gene Expression to
Control Differentiation, Stem Cell Function, and Immune Response in Leukemia. *Cancer Discov*
11, 1052-1066, doi:10.1158/2159-8290.CD-20-1227 (2021).

6   Chen, W., Zhao, H. & Li, Y. Mitochondrial dynamics in health and disease: mechanisms and
potential targets. *Signal Transduct Target Ther* **8**, 333, doi:10.1038/s41392-023-01547-9 (2023).

7   Raven, K. D. & Kapetanovic, R. Mitochondrial dynamics: Regulating cell metabolism,
homoeostasis, health and disease. *Semin Cell Dev Biol* **161-162**, 20-21,
doi:10.1016/j.semcdb.2024.02.002 (2024).

8   Baek, M. L. *et al.* Mitochondrial structure and function adaptation in residual triple negative breast
cancer cells surviving chemotherapy treatment. *Oncogene* **42**, 1117-1131, doi:10.1038/s41388-
023-02596-8 (2023).

9   Boulton, D. P. & Caino, M. C. Mitochondrial Fission and Fusion in Tumor Progression to
Metastasis. *Front Cell Dev Biol* **10**, 849962, doi:10.3389/fcell.2022.849962 (2022).

10 Sessions, D. T. *et al.* Opa1 and Drp1 reciprocally regulate cristae morphology, ETC function, and
NAD(+) regeneration in KRas-mutant lung adenocarcinoma. *Cell Rep* **41**, 111818,
doi:10.1016/j.celrep.2022.111818 (2022).

11 Perou, C. M. Molecular stratification of triple-negative breast cancers. *Oncologist* **16 Suppl 1**,
61-70, doi:10.1634/theoncologist.2011-S1-61 (2011).

12 Symmans, W. F. *et al.* Long-Term Prognostic Risk After Neoadjuvant Chemotherapy Associated
With Residual Cancer Burden and Breast Cancer Subtype. *J Clin Oncol* **35**, 1049-1060,
doi:10.1200/JCO.2015.63.1010 (2017).

13 Yau, C. *et al.* Residual cancer burden after neoadjuvant chemotherapy and long-term survival
outcomes in breast cancer: a multicentre pooled analysis of 5161 patients. *Lancet Oncol* **23**,
149-160, doi:10.1016/S1470-2045(21)00589-1 (2022).

14 Hamy, A. S. *et al.* Prognostic value of the Residual Cancer Burden index according to breast
cancer subtype: Validation on a cohort of BC patients treated by neoadjuvant chemotherapy.
PLoS One **15**, e0234191, doi:10.1371/journal.pone.0234191 (2020).

15　Echeverria, G. V. *et al.* Resistance to neoadjuvant chemotherapy in triple-negative breast cancer
mediated by a reversible drug-tolerant state. *Sci Transl Med* **11**,
doi:10.1126/scitranslmed.aav0936 (2019).

16　Herkenne, S. & Scorrano, L. OPA1, a new mitochondrial target in cancer therapy. *Aging (Albany*
NY) **12**, 20931-20933, doi:10.18632/aging.104207 (2020).

17　Xie, L. *et al.* Drp1-dependent remodeling of mitochondrial morphology triggered by EBV-LMP1
increases cisplatin resistance. *Signal Transduct Target Ther* **5**, 56, doi:10.1038/s41392-020-
0151-9 (2020).

18　Alalawy, A. I. *et al.* Inhibition of Drp1 orchestrates the responsiveness of breast cancer cells to
paclitaxel but insignificantly relieves paclitaxel-related ovarian damage in mice. *Sci Rep* **13**,
22782, doi:10.1038/s41598-023-49578-0 (2023).

19　Noguchi, M. *et al.* Inhibition of the mitochondria-shaping protein Opa1 restores sensitivity to
Gefitinib in a lung adenocarcinomaresistant cell line. *Cell Death Dis* **14**, 241,
doi:10.1038/s41419-023-05768-2 (2023).

20　Decker, C. W. *et al.* Mitofusin-2 mediates doxorubicin sensitivity and acute resistance in Jurkat
leukemia cells. *Biochem Biophys Rep* **24**, 100824, doi:10.1016/j.bbrep.2020.100824 (2020).

21　Rub, J. *et al.* A deep-learning tool for species-agnostic integration of cancer cell states. *bioRxiv*,
2024.2012.2020.629285, doi:10.1101/2024.12.20.629285 (2024).

22　Chaudhry, F., Kim, T. W., Elemento, O. & Betel, D. Machine learning analysis of population-wide
plasma proteins identifies hormonal biomarkers of Parkinson's Disease. *medRxiv*,
doi:10.1101/2024.12.21.24313256 (2024).

23　Wang, J. M. *et al.* Deep learning integrates histopathology and proteogenomics at a pan-cancer
level. *Cell Rep Med* **4**, 101173, doi:10.1016/j.xcrm.2023.101173 (2023).

24　Suga, S. *et al.* An interactive deep learning-based approach reveals mitochondrial cristae
topologies. *PLoS Biol* **21**, e3002246, doi:10.1371/journal.pbio.3002246 (2023).

25　Jiang, Y., Yang, M., Wang, S., Li, X. & Sun, Y. Emerging role of deep learning-based artificial
intelligence in tumor pathology. *Cancer Commun (Lond)* **40**, 154-166, doi:10.1002/cac2.12012
(2020).

26　Noorbakhsh, J. *et al.* Deep learning-based cross-classifications reveal conserved spatial
behaviors within tumor histological images. *Nat Commun* **11**, 6367, doi:10.1038/s41467-020-
20030-5 (2020).

27　Yang, Y., Fang, Q. & Shen, H. B. Predicting gene regulatory interactions based on spatial gene
expression data and deep learning. *PLoS Comput Biol* **15**, e1007324,
doi:10.1371/journal.pcbi.1007324 (2019).

28　Rich, J. M. *et al.* Deep learning image segmentation approaches for malignant bone lesions: a
systematic review and meta-analysis. *Front Radiol* **3**, 1241651, doi:10.3389/fradi.2023.1241651
(2023).

29   Xu, Y. *et al.* Advances in Medical Image Segmentation: A Comprehensive Review of Traditional,
Deep Learning and Hybrid Approaches. *Bioengineering (Basel)* **11**,
doi:10.3390/bioengineering11101034 (2024).

30   Yuan, J. Brain tumor image segmentation method using hybrid attention module and improved
mask RCNN. *Sci Rep* **14**, 20615, doi:10.1038/s41598-024-71250-4 (2024).

31   Ronneberger, O., Fischer, P. & Brox, T. 234-241 (Springer International Publishing).

32   Farahmand, S. *et al.* Deep learning trained on hematoxylin and eosin tumor region of Interest
predicts HER2 status and trastuzumab treatment response in HER2+ breast cancer. *Mod Pathol*
35, 44-51, doi:10.1038/s41379-021-00911-w (2022).

33   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci Rep*
7, 16878, doi:10.1038/s41598-017-17204-5 (2017).

34   Echeverria, G. V. *et al.* Predictors of success in establishing orthotopic patient-derived xenograft
models of triple negative breast cancer. *NPJ Breast Cancer* **9**, 2, doi:10.1038/s41523-022-
00502-1 (2023).

35   Guillen, K. P. *et al.* A human breast cancer-derived xenograft and organoid platform for drug
discovery and precision oncology. *Nat Cancer* **3**, 232-250, doi:10.1038/s43018-022-00337-6
(2022).

36   Zhang, X. *et al.* A renewable tissue resource of phenotypically stable, biologically and ethnically
diverse, patient-derived human breast cancer xenograft models. *Cancer Res* **73**, 4885-4897,
doi:10.1158/0008-5472.CAN-12-4081 (2013).

37   Zhou, Z., Siddiquee, M. M. R., Tajbakhsh, N. & Liang, J. UNet++: A Nested U-Net Architecture
for Medical Image Segmentation. *Deep Learn Med Image Anal Multimodal Learn Clin Decis*
Support (2018) **11045**, 3-11, doi:10.1007/978-3-030-00889-5_1 (2018).

38   Kaiming He, G. G., Piotr Dollár, Ross Girshick. Mask R-CNN. *arXiv*
doi:https://doi.org/10.48550/arXiv.1703.06870 (2017).

39   Conrad, R. & Narayan, K. Instance segmentation of mitochondria in electron microscopy images
with a generalist deep learning model trained on a diverse dataset. *Cell Syst* **14**, 58-71 e55,
doi:10.1016/j.cels.2022.12.006 (2023).

40   Echeverria, G. V. *et al.* High-resolution clonal mapping of multi-organ metastasis in triple
negative breast cancer. *Nat Commun* **9**, 5079, doi:10.1038/s41467-018-07406-4 (2018).

41   Lei, J. T. *et al.* Patient-Derived Xenografts of Triple-Negative Breast Cancer Enable
Deconvolution and Prediction of Chemotherapy Responses. *bioRxiv*,
doi:10.1101/2024.12.09.627518 (2025).

42   Rinkenbaugh, A. L. *et al.* An Annotated Biobank of Triple-Negative Breast Cancer Patient-
Derived Xenografts Features Treatment-Naive and Longitudinal Samples During Neoadjuvant
Chemotherapy. *Cancer Res*, doi:10.1158/0008-5472.CAN-24-5023 (2025).

43 Parida, P. K. *et al.* Limiting mitochondrial plasticity by targeting DRP1 induces metabolic
reprogramming and reduces breast cancer brain metastases. *Nat Cancer* **4**, 893-907,
doi:10.1038/s43018-023-00563-6 (2023).

44 Cheng, S. *et al.* RGCC-mediated PLK1 activity drives breast cancer lung metastasis by
phosphorylating AMPKalpha2 to activate oxidative phosphorylation and fatty acid oxidation. *J
Exp Clin Cancer Res* **42**, 342, doi:10.1186/s13046-023-02928-2 (2023).

45 Pinto-Coelho, L. How Artificial Intelligence Is Shaping Medical Imaging Technology: A Survey of
Innovations and Applications. *Bioengineering (Basel)* **10**, doi:10.3390/bioengineering10121435
(2023).

46 Begum, H. M. *et al.* Spatial Regulation of Mitochondrial Heterogeneity by Stromal Confinement
in Micropatterned Tumor Models. *Sci Rep* **9**, 11187, doi:10.1038/s41598-019-47593-8 (2019).

47 Han, M. *et al.* Spatial mapping of mitochondrial networks and bioenergetics in lung cancer.
Nature **615**, 712-719, doi:10.1038/s41586-023-05793-3 (2023).

48 Fischer, J. R. *et al.* Multiplex imaging of breast cancer lymph node metastases identifies
prognostic single-cell populations independent of clinical classifiers. *Cell Rep Med* **4**, 100977,
doi:10.1016/j.xcrm.2023.100977 (2023).

49 Jovanovic, B. *et al.* Heterogeneity and transcriptional drivers of triple-negative breast cancer.
Cell Rep **42**, 113564, doi:10.1016/j.celrep.2023.113564 (2023).

50 Lehmann, B. D. *et al.* Identification of human triple-negative breast cancer subtypes and
preclinical models for selection of targeted therapies. *J Clin Invest* **121**, 2750-2767,
doi:10.1172/JCI45014 (2011).

51 Dietlmeier, J. *et al.* Few-shot hypercolumn-based mitochondria segmentation in cardiac and
outer hair cells in focused ion beam-scanning electron microscopy (FIB-SEM) data. *Pattern
Recognit Lett* **128**, 521-528, doi:10.1016/j.patrec.2019.10.031 (2019).

52 Gallusser, B. *et al.* Deep neural network automated segmentation of cellular structures in volume
electron microscopy. *J Cell Biol* **222**, doi:10.1083/jcb.202208005 (2023).

53 Gupta, Y., Heintzmann, R., Costa, C., Jesus, R. & Pinho, E. Deep learning-enhanced automated
mitochondrial segmentation in FIB-SEM images using an entropy-weighted ensemble approach.
PLoS One **19**, e0313000, doi:10.1371/journal.pone.0313000 (2024).

54 Casser, V., Kang, K., Pfister, H. & Haehn, D. in *Proceedings of the Third Conference on Medical
Imaging with Deep Learning* Vol. 121 (eds Arbel Tal *et al.*) 111--120 (PMLR, Proceedings of
Machine Learning Research, 2020).

55 Gao, H. & Ji, S. in *2017 IEEE International Conference on Data Mining (ICDM)*   871-876 (IEEE
Computer Society, 2017).

56 Wei, D. *et al.* MitoEM Dataset: Large-scale 3D Mitochondria Instance Segmentation from EM
Images. *Med Image Comput Comput Assist Interv* **12265**, 66-76, doi:10.1007/978-3-030-59722-
1_7 (2020).

57 Xu, C. S. *et al.* An open-access volume electron microscopy atlas of whole cells and tissues.
Nature **599**, 147-151, doi:10.1038/s41586-021-03992-4 (2021).

58 Heinrich, L. *et al.* Whole-cell organelle segmentation in volume electron microscopy. *Nature* **599**,
141-146, doi:10.1038/s41586-021-03977-3 (2021).

59 Pereira, R. O. *et al.* OPA1 deficiency promotes secretion of FGF21 from muscle that prevents
obesity and insulin resistance. *EMBO J* **36**, 2126-2145, doi:10.15252/embj.201696179 (2017).

60 Favaro, G. *et al.* DRP1-mediated mitochondrial shape controls calcium homeostasis and muscle
mass. *Nat Commun* **10**, 2576, doi:10.1038/s41467-019-10226-9 (2019).

61 Neikirk, K. *et al.* Systematic Transmission Electron Microscopy-Based Identification and 3D
Reconstruction of Cellular Degradation Machinery. *Adv Biol (Weinh)* **7**, e2200221,
doi:10.1002/adbi.202200221 (2023).

62 Southard, S. *et al.* A series of Cre-ER(T2) drivers for manipulation of the skeletal muscle lineage.
Genesis **52**, 759-770, doi:10.1002/dvg.22792 (2014).

63 Lam, J. *et al.* A Universal Approach to Analyzing Transmission Electron Microscopy with ImageJ.
Cells **10**, doi:10.3390/cells10092177 (2021).

64 Han, J. *et al.* ER-stress-induced transcriptional regulation increases protein synthesis leading to
cell death. *Nat Cell Biol* **15**, 481-490, doi:10.1038/ncb2738 (2013).

65 Sarkans, U. *et al.* The BioStudies database-one stop shop for all data supporting a life sciences
study. *Nucleic Acids Res* **46**, D1266-D1270, doi:10.1093/nar/gkx965 (2018).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1