A Syntenic Pangenome of *Gardnerella* Reveals Novel Plasmids and Phage, Taxonomic Boundaries, and Species-Level Stratification of Metabolic and Virulence Potential

Heather K. Bouzek*[1,2], Martha A. Zepeda-Rivera*[1,2], Sujatha Srinivasan[2], Elliot M. Lee[2], Susan M. Strenk[2], Dakota S. Jones[2], Elsa F. McMahon[2], Tina L. Fiedler[2], Marko Kostovski[3], Michael T. France[4,5], Jacques Ravel[4,5], David N. Fredricks [‡2], and Christopher D. Johnston[‡1,2]

[1] Department of Genomic Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas, USA

[2] Vaccine and Infectious Disease Division, Fred Hutch Cancer Center, Seattle, Washington, USA

[3] Institute of Microbiology and Parasitology, Faculty of Medicine, University Ss Cyril and Methodius in Skopje, Skopje, Macedonia

[4] Center for Advanced Microbiome Research and Innovation, Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, Maryland, USA

[5] Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, Maryland, USA

[‡]Address correspondence to:

David N. Fredricks, MD, dfredric@fredhutch.org

Christopher D. Johnston, PhD, cdjohnston@mdanderson.org

* These authors contributed equally to this work.

Abstract

Gardnerella species are central to bacterial vaginosis (BV), a condition affecting nearly one in three women of reproductive age and associated with preterm birth and increased susceptibility to sexually transmitted infections. Despite decades of study, progress in defining *Gardnerella* diversity has been hindered by inconsistent taxonomy and poor-quality genomic resources. Here, we sequenced 392 new *Gardnerella* isolates from asymptomatic and BV-associated microbiota and integrated this collection with all publicly available genomes. After stringent curation, we generated a high-quality reference set of 313 distinct genomes that underpins a comprehensive taxonomic framework. Using average nucleotide identity (ANI), digital DNA–DNA hybridization (dDDH), and phylogenomics, we resolved 21 genomic lineages encompassing 11 species and 15 subspecies, each assigned a provisional formal name. Integration of complete long-read assemblies enabled construction of the first syntenic *Gardnerella* pangenome, revealing lineage-specific repertoires of virulence, metabolic, and defense systems - including variable sialidases (NanH), vaginolysin, and amino-acid biosynthetic pathways - and defining conserved genomic architecture across species. Comparative methylome profiling further highlighted restriction-modification system diversity that may influence genetic exchange. Finally, we identified the first native cryptic plasmids in *Gardnerella*, overturning the assumption that the genus lacks plasmids, and demonstrated their use in generating a replicative *E. coli-Gardnerella* shuttle vector. Together, these results establish a complete genomic and functional framework for *Gardnerella*, providing a reproducible foundation for mechanistic and translational studies of BV and a model for resolving taxonomy and functional stratification in other urogenital-associated bacteria.

Main

Bacterial vaginosis (BV) is a common and global condition characterized by an imbalance in the vaginal microbiota[1-3], associated with significant health risks, including increased susceptibility to sexually transmitted infections[4,5] and adverse pregnancy outcomes such as preterm birth and low birth weight[6-8]. While molecular methods have associated many bacterial species with BV, *Gardnerella vaginalis* was the first bacterium isolated from individuals affected with BV[9], and was long thought to be the sole causative agent of this condition[1,10,11] due to its high virulence potential[3,12-20], its presence in 95–100% of BV cases[11,21,22] and its ability to thrive in the vaginal environment[10,19,20,23-25]. Our previous research revealed that while *Gardnerella vaginalis* is correlated with a greater risk ratio for BV than other validly named *Gardnerella* species, no single *Gardnerella* species is specific for BV based on a *cpn60* qPCR. In fact, the presence of a high *Gardnerella* species diversity was identified as a risk factor for the development of BV in our longitudinal study[22]. The presence of *Gardnerella* species in both BV-affected and asymptomatic individuals suggests the potential for both commensal and pathogenic lineages or states[3]. However, the genetic capabilities of individual *Gardnerella* lineages remain unclear, partially due to inconsistent taxonomic assignments and overly complex *Gardnerella* nomenclature[19], despite laudable efforts from multiple research groups[26-35]. Further, many of the public *Gardnerella* genome assemblies are duplicated, contaminated, or incompletely assembled and did not pass our quality metrics for pangenome analysis. Thus, we sequenced over 392 *Gardnerella* isolates genomes from both asymptomatic individuals or those with BV and combined these with all available public genomes to generate a highly curated and deduplicated collection for analyses. We applied comprehensive comparative genomic analysis of this curated collection to taxonomically identify *Gardnerella* lineages and propose a single naming scheme that is reflective of their

phylogenetic placement and historical context. Syntenic pangenome partitioning revealed lineage-

specific repertoires of metabolic pathways, virulence-associated genes, and defense systems.

Collectively, this study integrates pangenome analyses, genome-based taxonomy, and experimental

validation to define species boundaries within *Gardnerella*, characterize lineage-specific metabolic

and virulence-associated potential, and identify mobile genetic elements, including a cryptic

plasmid that enables the development of a shuttle vector for the genetic engineering of *Gardnerella*.

Results

***Gardnerella* and *Bifidobacterium* are distinct genera**

We obtained genomes for 392 *Gardnerella* isolates (Fredricks and Ravel bacterial isolate

collections) from asymptomatic individuals and those with BV. We curated these and additionally

available public *Gardnerella* genomes (n=271 within the NCBI database as of May 2025) using

stringent quality metrics to remove duplicate and low-quality genomes (Methods, Supplementary

Table 1). This resulted in a final curated set of 312 genomes for comparative genomic analyses

(Supplementary Table 2). Analysis of the *cpn60* gene (which encodes a 60 kDa chaperonin)

currently provides the most reliable method for *Gardnerella* species typing, as these species are

indiscernible by traditional 16S rRNA gene sequence typing[36,37]. BLASTn analysis of the *cpn60*

gene of these isolates supported their identification as *Gardnerella* species (Supplementary Table

2).

Despite the classification of these genomes as *Gardnerella* against and within the NCBI

database, initial speciation against the Genome Taxonomy Database (GTDB) assigned many of

these genomes to *Bifidobacterium* (Supplementary Table 2). However, *Bifidobacterium* is known

to be a polyphyletic group, consisting of at least six distinct clades[38,39], whose members exhibit

substantial genetic diversity. The separation of *Gardnerella* from *Bifidobacterium* has previously been demonstrated through comparisons of genome length, G+C percentage, and overall genome composition[31,39]. Consistent with theses prior observations, comparisons of genome length and G+C percentage between our *Gardnerella* genome set, *Bifidobacterium,* and closely related scardovial genera (e.g., *Alloscardovia[40], Neoscardovia[41], Aeriscardovia[42], Scardovia[43]*) (n=229 genomes; Supplementary Table 3) indicate that *Gardnerella* genomes are significantly shorter ($p < 0.001$, Wilcoxon *post hoc* test) and have lower G+C content ($p < 0.001$, Wilcoxon *post hoc* test) than other *Bifidobacterium* members (Extended Data Fig. 1a-b). Notably, *Gardnerella* shares a human-focused and pathogenic niche with *Alloscardovia, Aeriscardovia* and *Scardovia* and clusters more closely with the scardovial genera across these genomic features than with *Bifidobacterium.* Additionally, average amino acid identity (AAI) values between *Gardnerella* and *Bifidobacterium* fell below genus thresholds of 65%[44] (Extended Data Fig. 1c). Together, these results confirmed that our final curated genome set comprises only *Gardnerella* species, further supporting previous evidence that *Gardnerella* and *Bifidobacterium* represent distinct genera within the *Bifidobacteriaceae* family.

Identification and proposed revised nomenclature for *Gardnerella* lineages

Given the importance of *Gardnerella* species diversity in BV[21], we first assessed the number of lineages present in our genome collection. *Gardnerella* has been classified using various approaches[30-33,45,46], but consistent naming has been a challenge for *Gardnerella* despite extensive efforts[26-28,32] (Fig. 1). Early subtyping by Piot *et al.*[26] relied on sugar hydrolysis phenotypes and later expanded to 17 biotypes[27]. Genomic methods such as restriction fragment length polymorphism (RFLP)[28] and amplified ribosomal DNA restriction analysis (ARDRA)[29] identified

several distinct genotypes. PCR and whole-genome sequence comparisons introduced two new

naming schemes including designations "A - D"[37] or "1 - 4"[30] which led to hybrid labels ("C/1",

"B/2", "A/4", and "D/3")[47]. These species were then reframed as reassorted ecotypes[31]

incorporating the genomic context of the distinct types. With widespread whole-genome

sequencing, additional standards based on ANI emerged, describing thirteen genomic species

(Gsp)[32], nine genomospecies (GS)[33], or nine different genotypes (GGtypes)[35]. Subsequent formal

naming of a subset of these species included *G. vaginalis*[9], *G. pickettii*[48], *G. piotii*[32], *G. leopoldii*[32],

G. swidsinskii[32], *G. greenwoodii*[48], *G. lacydonensis*[49], *G. bretellae*[49], *G. massiliensis*[49], and *G.*

phocaeensis[49] (Fig. 1). Despite the use of diverse classification schemes, phylogenetic analyses

across studies consistently recover a deep bifurcation within *Gardnerella*, with the same two major

lineages forming the basal split of the genus. While this division has been explicitly designated as

"Set A" and "Set B" only by Ahmed *et al.*, analogous groupings confirm the primary branching

structure in multiple independent phylogenetic frameworks[30-35,37,46,50,51].

Consistent with this, *cpn60* gene sequences and reference-free whole-genome phylogenies

separated all genomes in our collection into two deeply divergent lineages (Fig. 2, Extended Data

Fig. 2, Supplementary Table 4) here maintaining the Set A and Set B designation. To determine the

number of *Gardnerella* species represented, we calculated pairwise genome-wide average

nucleotide identity (ANI) values across all genomes and examined their distribution (Extended

Data Fig. 3a). In accordance with prior large-scale analyses demonstrating a pronounced

discontinuity in ANI values around the canonical prokaryotic boundary, most inter-lineage

comparisons fell well below 95%, whereas closely related genomes clustered above 95% with an

exceedingly small fraction of the sequence comparisons falling above 94% (Extended Data Fig.

3a). Because ANI and digital DNA–DNA hybridization (dDDH) values were strongly colinear

across the dataset, dDDH did not provide additional resolution but independently supported the

same groupings: all genomes that clustered together by ANI also exceeded the 70% dDDH species

threshold (Extended Data Fig. 3b–c). Thus, species boundaries inferred from ANI were fully

congruent with the universally accepted 70% dDDH. Integrating whole-genome phylogenies with

ANI/dDDH clustering, we identified eleven *Gardnerella* species. Within five of these species,

additional phylogenomic structure supported sub-species level differentiation, resulting in twenty-

one distinct *Gardnerella* lineages (Fig. 2, Supplementary Table 4). These species-level assignments

were congruent with the classification scheme proposed by Potter[33]. The *Gardnerella* species

bretellae, lacydonensis, leopoldii, phocaeensis, *pickettii, piotii*, and *swidsinskii* were previously

described as distinct species when integrating phenotypic characterization with genome-based

analyses. In our dataset, these lineages likewise form well-supported monophyletic clades in whole-

genome phylogenies and remain clearly differentiated from other *Gardnerella* taxa. However, when

evaluated across the expanded genome collection as analyzed here, pairwise ANI and dDDH

comparisons reveal that these lineages consistently fall within the genomic coherence of broader

species-level clusters defined by ≥95% ANI and ≥70% dDDH thresholds. While they are

phylogenetically structured and phenotypically distinguishable, they do not cross the currently

accepted genomic discontinuity thresholds that would necessitate recognition as separate species

under genome-based standards.

Given the extensive history of overlapping and partially incompatible classification

schemes for *Gardnerella* (Fig. 1), we next harmonized existing species and subspecies names and

established a new nomenclature across all identified lineages (Fig. 1, Fig. 2, Supplementary Tables

2 and 4). Details on the rationale for each proposed nomenclatural revision, including species-

specific functional attributes and ecological isolation sources, are provided in the Protologues. Our aim was to integrate previously used species and subspecies names to maintain a consistent framework. The historically recognized "*G. vaginalis*" corresponds to a subspecies-level clade within a broader species and is designated here as *G. vaginalis* subspecies *vaginalis*, preserving historical continuity while reflecting its phylogenetic placement (Fig. 1, Supplementary Table 1, Supplementary Table 4). Similarly, we propose distinct species names for lineages with well-supported genomic and phenotypic divergence, including *G. bivia*, *G. hutchinsoni*, and *G. washingtoni*, among others (Extended Fig. 3g). To minimize ambiguity in referring to *Gardnerella* species in this paper and future efforts, we intentionally propose species names that can be abbreviated with unique two letter abbreviations (*Gb*, *Gg, Gh, Gi*, *Gk*, *Gl*, *Gm, Gp, Gr*, *Gv* and *Gw*), while individual subspecies were designated with discernable three-letter abbreviations (except for *Gardnerella piotii* subsp. *pickettii* and *Gardnerella piotii* subsp. *piotii,* which each require four letters, *Gppk* and *Gppt*) (Fig. 1, Supplementary Table 4).

Figure 1: Historical context of *Gardnerella* nomenclature. a, From top to bottom, the schematic shows the identification method by type, publication first author, names given to *Gardnerella* or naming framework, date of publication, and resulting conclusions from naming schemes. **b**, *Gardnerella* subtyping resulting from the publication in (**a**), where each column represents a subset of *Gardnerella*. Arrows indicate consistent lineages between studies without implying a definitive link. The two right-most columns contain updated lineage assignments with corresponding species and subspecies nomenclature proposed within this study.

Figure 2: Average nucleotide identity values between *Gardnerella* genomes. Average nucleotide

identity values between *Gardnerella* genomes in **a**, Set A and **b**, Set B. Key for each panel indicates

corresponding values to each color. White represents values below 90%. Genomes are organized

based on a kSNP4-generated whole-genome reference-free phylogeny. Figure is a composite of the

kSNP4 tree created in iTOL and the map sorted by the values in the kSNP4 tree as created using

ComplexHeatmap in R.

A syntenic pangenome for *Gardnerella*

Having stringently identified *Gardnerella* species and subspecies lineages in our genome

set and assigning names that adhered as closely to precedent, where possible, we were next able to

investigate the genetic features associated with each lineage. Pangenome analysis provides a

comprehensive view of the genetic repertoire within a group of bacteria by identifying genetic

features that are well-conserved across most (>95%) genomes ("persistent" or "core" genome)

versus those that are only present in a subset of genomes ("cloud" or "accessory" genome). While

the core genome often contains genes involved in essential cellular processes[52,53], analysis of the

accessory genomes can provide key insights about genetic diversity, adaptation, and evolutionary

relationships. For example, previous pangenome efforts in *Gardnerella* have expanded taxonomic

groupings, identified metabolically essential genes, and uncovered virulence and niche adaptation

features[34,50,54]. However, bacterial genes often function in co-regulated units or clusters involved

in specific biological processes (e.g., operons). To identify putative operons within the *Gardnerella*

pangenome, we applied a syntenic approach that accounts for gene order by using the Partitioned

Pangenome Graph of Linked Neighbors (PPanGGOLiN) tool[52] (Figs. 3-4, Supplementary Table

5). Briefly, in these analyses, the coding sequences of genes are grouped into "nodes" based on

amino acid similarity and visualized as circular nodes, with node size reflecting the relative

presence of that gene group across the genome set (Fig. 4a-c). Nodes representing neighboring

genes are linked into syntenic units called "modules" (Extended Data Fig. 4). This analysis clearly

distinguished between the core ("persistent" partition) and accessory genomes, and enabled

identification of accessory genetic elements present at intermediate frequency ("shell" partitions)

or low frequency ("cloud" partition) (Fig. 3). By leveraging modules and pangenome partitions,

conserved arrangements of genes can be traced across multiple genomes and enables insight into

the functional and evolutionary dynamics driving diversity at the genus, species, and strain levels.

In our collection, the core genome of *Gardnerella* reflected approximately 16% of identified

nodes, enriched in essential cellular functions including translation and ribosomal structure,

transcription, RNA processing, signal transduction, energy production, and metabolism of

nucleotides, amino acids, inorganic ions, and lipids (Fig. 3, Supplementary Table 5). Periodically,

these conserved nodes are interrupted by shell-associated nodes (Fig. 4a; nodes in distinct colors,

branching from orange nodes), which mark points of evolutionary divergence. These shell nodes

often represent horizontal gene transfer (HGT) events and are commonly associated with alternate

evolutionary pathways. We noted node differences between shell partitions that were consistent

with virulence, metabolism, and defense systems, warranting further investigation.

Figure 3: *Gardnerella* pangenome partitions. The tile plot displays the presence and absence of nodes in each partition where each row represents a node, and each column corresponds to a *Gardnerella* genome. Colors on the side correspond to core/persistent (orange), phage (red), shells (green), and cloud (blue) partitions. Genomes are ordered by kSNP4 phylogenies of Set A and Set B, with colors on dendrogram indicating lineage.

Figure 4: Syntenic maps of the *Gardnerella* pangenome. a–c, Visualizations of the PPanGGOLiN-generated *Gardnerella* pangenome. Each node represents a gene group, with edges connecting syntenic neighboring genes. Node size reflects the relative frequency of the gene group across *Gardnerella* genomes. Colors indicate core (orange) or shell (other colors) partitions. **b** and

c, The core and shell partitions are shown individually. **d**, map of the presence and absence of the

sialidase genes (*nanH1-nanH4*), type II α-amylase-pullulanase and vaginolysin as colored by the

species and subspecies of *Gardnerella*. **e**, Presence–absence bar chart showing the percentage of

conserved *Gardnerella* vaginolysin and the sialidase genes across each *Gardnerella* set (gray). Raw

presence/absence counts are shown above each bar. **f**, Gene context maps for the sialidase loci.

Virulence factors and predicted pathogenicity across *Gardnerella* lineages

Colonization of the vaginal niche by *Gardnerella*, and its transition to a pathogenic state,

depend on mucin degradation, adherence to the vaginal epithelium, and the formation of

structurally stable biofilms[13,55]. Canonical *Gardnerella* virulence factors implicated in these

processes include exported sialidases and vaginolysin[56], a cholesterol-dependent cytolysin. Among

Gardnerella-associated sialidase isoforms, NanH2 and NanH3 account for the majority of

extracellular sialidase activity, whereas NanH1 lacks a signal peptide and is therefore predicted to

function intracellularly for sugar catabolism[57]. Consistent with this role, *nanH1*[58] is embedded

within a conserved operon across genomes (Extended Figure 5e). In contrast, *nanH2* and *nanH4*[59]

are frequently located in proximity to this operon but are associated with mobile genetic elements,

suggesting more recent paralogous diversification and horizontal acquisition. Notably, *nanH3* is

uniquely not operon-associated; instead, it is found as a solitary gene within defense-associated

genomic regions, often adjacent to prophage elements, consistent with a distinct evolutionary

trajectory and potential phage-linked mobility. Within our pangenome dataset, we identified

lineage-specific nodes encoding vaginolysin, which contributes to host interaction by binding to

and lysing epithelial cells of the vaginal mucosa[55], and multiple enzymes involved in the stepwise

degradation of mucin-associated glycans (Fig. 4d-e, Extended Data Fig. 5, Supplementary Table

5). These include secreted sialidases and additional carbohydrate-active enzymes, such as L-

fucosidase[60-62] and galactosidase[27] that facilitate mucin catabolism when exported[63].

A Type II α-amylase-pullulanase protein was well conserved (present in 100% of

Gardnerella genomes), in agreement with previous reports[64] (Fig. 4d). Comparisons between

Gardnerella Set A and Set B showed that the four sialidase genes (*nanH1-nanH4*) are significantly

enriched in Set A genomes, while vaginolysin (*vly*) is significantly enriched in Set B genomes (Fig.

4e, Extended Data Fig. 5), highlighting lineage-dependent differences in metabolic capacity and

the distribution of putative virulence-associated genes across *Gardnerella*.

Predicted metabolic capabilities across *Gardnerella* lineages

Given the observed difference in metabolism-related nodes, we bioinformatically predicted

the biosynthesis of amino acids and the catabolism of small carbon sources across our *Gardnerella*

genome dataset (Fig. 5a, Fig. 5f, Supplementary Table 6). The biosynthesis of glutamine, glycine,

and proline was well conserved (>99% of genomes) while a high proportion of genomes (89%)

harbored the capacity for cysteine biosynthesis (Fig. 5a). Asparagine biosynthesis was enriched for

overall for Set B versus Set A, but within Set A, 100% of the *Gr* strains carried the genetic pathway.

Within Set B, asparagine biosynthesis was enriched for *Gg* (although depleted for *Ggi*) and the *Gll*

lineage alone amongst other *Gl* (Fig. 5a, b). Methionine biosynthesis is also enriched throughout

Set B, but absent in all of Set A. Within Set B methionine biosynthesis is notably lacking for strains

of *Gb* and *Ggg* (Fig. 5a, Fig. 5d). Threonine biosynthesis is much more sporadically throughout Set

A as compared to Set B (Fig. 5a, Fig. 5e). Additionally, threonine synthesis is more likely to be

found in *Gv* compared to other Set A (Fig. 5a, Fig. 5e). Although cysteine biosynthesis is well

conserved this pathway is entirely absent in *Gh* genomes (Fig. 5a) and depleted in *Gps* as

compared to other *Gp* genomes (Fig. 5a, Fig. 5e) .

247   Genes for the catabolism of ethanol appear to be a well-conserved feature of *Gardnerella*

genomes (Fig. 5f). Genomes across Set A were additionally significantly enriched for the

catabolism of alanine, gluconate, and serine (Fig. 5f, Fig. 5h, Fig. 5j, Fig. 5k). Notably, lineage-

specific associations within Set A were observed for the catabolism of each of these small carbon

sources. While gluconate catabolism is significantly enriched in *Gv*, alanine catabolism is

significantly depleted in *Gvl* as compared to other Set A members (Fig. 5f, Fig. 5h, Fig. 5j). Serine

catabolism was completely absent in *Gr* and (Fig. 5f) and was significantly depleted in *Ggb* and

Gls compared to other lineages within their respective species (Fig. 5f, Fig. 5k). Interestingly, the

catabolism of acetate and deoxyribose was lineage-specific and unique to *Gb* and *Gv*, respectively

(Fig. 5f). Deoxyribose catabolism was enriched in the *Gvv* compared to other *Gv* subspecies (Fig.

5f, Fig. 5i). Our results indicate that Set A and Set B genomes are distinguished by their relative

enrichment of pathways for catabolism of small carbon sources or synthesis of distinct amino acids

with lineage-specific associations.

Figure 5: Predictions for metabolic capabilities across *Gardnerella* lineages. Presence-absence map for GapMind-predicted pathways to **a,** synthesize amino acids (AA) or **f,** catabolize small carbon sources (SCS) where each row represents a *Gardnerella* genome. Genomes are ordered by

kSNP4 phylogenies of Set A and Set B. Presence by proportion comparison of genomes for the **b-**

e, biosynthesis of a subset of amino acids or **g-k**, catabolism of a subset of small carbon sources.

Colors represent each lineage.

Phage and antiphage defense mechanisms across *Gardnerella* genomes

In our pangenome data set, we noted that multiple nodes were related to phage-associated

proteins that encompassed two distinct *Siphoviridae* family *Gardnerella* prophage, including the

previously described vB_Gva_AB1[51,65] and a novel phage we have tentatively designated as

"pGamma" (Extended Data Fig. 5, Supplementary Table 7). This newly recognized phage is from

a distinct family from vB_Gva_AB1 within the Class *Caudoviricetes* in whole genome

comparisons[66]. vB_Gva_AB1 is sometimes reported as several different prophages[46,51], the discrete

synteny within our PPanGGOLiN pangenome suggests that these variants represent at least one

discrete taxon (Extended Data Fig. 5). It is notable that some strains appear to have remnant phage

genes, although these appear to be non-syntenic and scattered throughout the genome

(Supplementary Table 5). While both vB_Gva_AB1 and pGamma were identified in Set A and Set

B genomes, vB_Gva_AB1 was significantly enriched in the Set B species *G. leopoldii* (Bonferroni

adj. $p < 0.01$, Fisher's Exact test) and *G. greenwoodii* (Bonferroni adj. $p = 0.4$, Fisher's Exact test)

(Extended Data Fig. 6).

To counteract phages, bacteria evolved an array of defense systems[67], with the most

widespread and well-characterized being restriction-modification (RM)[68,69] and CRISPR-Cas

mechanisms[70]. Across our *Gardnerella* genome dataset we identified the presence of sixty different

systems, with an average of twelve defenses per genome (Fig. 6a-b, Extended Data Fig. 7,

Supplementary Table 8). Of the defense systems with described mechanisms, dXTPase systems[71]

were well conserved (99% genomes). Type I RM systems were prevalent (56% of genomes) and

did not differ significantly between *Gardnerella* sets. Set A genomes were significantly enriched

for some Abi proteins (AbiD[72] and AbiL[73]), CBASS defenses[74,75], and DRT mechanisms[76,77], while

Set B showed enrichment for a larger number of defenses, including those with Abi-mechanisms

(AbiE[69], Gabija[75], and Lamassu[75,78]), CRISPR-Cas (CRISPR arrays and Cas Type IE proteins), and

Type II RM defenses (Extended Data Fig. 8). Across *Gardnerella* lineages, *Gls* showed an

enrichment for DNA-modification system (DMS) proteins, while *Gvv* showed an enrichment for

AbiL and DRT defenses (Extended Data Fig. 8).

Although defense mechanisms evolved primarily to protect against bacteriophages, their

activity extends against other horizontally transferred genetic elements, as well as plasmid vectors

used during *in vitro* transformation. RM defenses, which are dependent upon detection and

destruction of inappropriately methylated foreign DNA, are one of the most prevalent across

bacteria. Further, it is notable that approaches to overcome these RM systems during genetic

engineering have been shown to significantly improve transformation efficiency across multiple

bacterial species[79-81]. Thus, we sought to characterize RM defenses across a subset of our

Gardnerella genome dataset. We obtained and analyzed methylomes for 42 *Gardnerella* isolates

across thirteen lineages within the Fredricks isolate collection (Supplementary Table 2). A total of

39 methylated motifs were detected and assigned to Type I (25 motifs), Type II (11 motifs), or Type

III (3 motifs) RM systems (Fig. 6c, Supplementary Table 9).

Figure 6: Defense systems, methylation motifs, and plasmid architectures across *Gardnerella* lineages

a-d, Presence-absence maps for PADLOC-predicted defense mechanisms. Genomes are ordered

by kSNP4 phylogenies of **a,** Set A and **b,** Set B. **c,** Sunburst plot shows the sequence of each

methyl-modified motif detected within each *Gardnerella* lineage. Size of section represents

prevalence within each lineage. **d,** Schematics depict plasmid annotations for pDNF01199S-01,

p0756_546_1_1_BHK10-01, and pNR047-01. Full sequences for the plasmids are available in

Supplementary Table 10.

Identification of a native *Gardnerella* cryptic plasmid

There is renewed interest in developing genetic tools for engineering *Gardnerell*a

species[82,83]. Because *Gardnerella* was long thought to lack mobile genetic elements, such as native

plasmids, most efforts have focused on adapting non-*Gardnerella* genetic tools for use in this

genus. Recent studies have been extensive and show promising results, achieving targeted

chromosomal integrations and deletions using *E. coli*-based suicide vectors[82], as well as

reproducible transformation with a *Bifidobacterium-E. coli* shuttle vector[83]. During genome

sequencing of the Fredricks clinical isolate collection, we observed an independent circular contig,

with high coverage, forming a second replicon within the otherwise closed genome sequence of

Gppk strain DNF01199S (Fig. 6d, Supplementary Table 10). Annotation of this contig showed

three coding sequences for hypothetical proteins and a single putative replication initiation protein

(Supplementary Table 11). Subsequent plasmid isolation and validation confirmed the episomal

nature of this contig, marking pDNF01199S-01 as the first cryptic plasmid identified within the

Gardnerella genus.

Motivated by this discovery, we assessed whether additional cryptic plasmids were present

in our *Gardnerella* pangenome. We reasoned that the current limited knowledge of such plasmids

reflects their rarity. As such, we focused on the cloud partition of the accessory genome, which

highlights strain-specific genetic factors and collectively comprises 47.23% of the *Gardnerella*

pangenome. Using structural sequence insights from pDNF01199S-01, we identified two

additional putative plasmid sequences within the cloud partition: one in *Gvv* strain

0756_546_1_1_BHK10[84] (p0756_546_1_1_BHK10-01), which was isolated from cervicovaginal

mucus in South Africa, and another in *Gk* strain NR047[23] (pNR047-01), isolated from a vaginal

swab in Nairobi, Kenya. All three plasmids are approximately 4kb in size, contain coding

sequences for putative hypothetical proteins, and share a highly similar replication initiation

protein (84.48%-95.26% similar), suggesting a potentially conserved replication mechanism for

cryptic plasmids within the *Gardnerella* genus (Fig. 6d, Supplementary Tables 10-11). Given that

pDNF01199S-01 replicates stably in *Gardnerella*, we next sought to engineer a *Gardnerella-E.*

coli shuttle vector suitable for transformation of additional *Gardnerella* isolates. Analysis of our

pangenome revealed that 22% of *Gardnerella* genomes encode *tetM*, including *Gppk* DNF01199S,

consistent with previous reports on variable tetracycline resistance phenotypes across *Gardnerella*

strains[85]. We therefore assessed the growth of both *tetM+* and *tetM-* isolates in the presence of

supplemented tetracycline. As predicted, *tetM+* isolates were resistant to tetracycline, whereas

tetM- isolates were susceptible, confirming that the *tetM* cassette confers tetracycline resistance *in*

vitro. This supports the use of *tetM* as a selectable marker for this vector. To enable the propagation

in *E. coli* strains, we combined pDNF01199S-01, the *tetM* cloned from DNF01199S, and an *E. coli*

minicircle plasmid backbone (pMC) into a single parental plasmid (herein designated pG01-pp)

(Extended Data Fig. 9a-c). As previously reported[86], we induced and purified the pDNF01199S-

tetM-MC <u>minic</u>ircle vector (pG01-MC). We initially attempted to transform this minicircle into

the *tetM*-strain, ATCC 14018[T] (*Gvv*), but no transformants were recovered (Extended Data Fig.

9d).

The strain ATCC 14018[T] encodes a restriction-modification (RM) system which modifies

the "GGCC" motif[84] and has been shown to limit transformation efficiency[82]. Thus, we used

commercially available *HaeIII* methyltransferase (MTase) to methyl-modify these sites in pG01-

MC prior to transformation. Electroporation with methylated pG01-MC successfully transformed

Gvv ATCC 14018[T], but at a very low efficiency (two transformants per µg of DNA). Further

optimization of the protocol increased efficiency to 685 transformants per µg of DNA by

transforming with 0.5 µg total *HaeIII*-treated vector on competent cells grown with 100 mM NaCl

to an OD_{600} of 0.32 and washed with 10% glycerol (Extended Data Fig. 9d). A detailed, step-by-

step   version   of   the   optimized   protocol   is   available   online   at

(https://www.protocols.io/private/1A47BAD2F1AC11F09AA30A58A9FEAC02).

Collectively, these results show that pG01-pp is the first replicative shuttle vector derived

from a native *Gardnerella* cryptic plasmid. Together with recent work by the Dillard and Hill

groups[82,83], these efforts should establish a robust platform for more mechanistic investigations of

Gardnerella biology and pathogenesis.

Discussion

Taxonomic inconsistencies and fluctuating nomenclature have made it inherently

challenging to delineate the roles of *Gardnerella* lineages in health and disease. In this study, we

integrate whole-genome analyses to establish a refined and internally consistent taxonomic

framework for the *Gardnerella* genus. Using a curated set of 313 distinct, high-quality genomes,

we define clear lineage boundaries and propose an updated *Gardnerella* nomenclature that reflects

genome-scale divergence while preserving continuity with existing naming conventions. Although

nomenclatural revisions, particularly for well-established organisms such as '*G. vaginalis,*'

(updated here to *G. vaginalis* subsp. *vaginalis*), can often be contentious, applying established

genomic criteria for species and subspecies delineation is warranted to prevent further taxonomic

ambiguity and this misinterpretation of functional studies.

While single-locus markers such as *cpn60*, which are widely used for *Gardnerella* typing,

can distinguish between *Gardnerella* species, they lack sufficient resolution for subspecies

assignments. Accurate subspecies assignments require whole-genome approaches including

phylogenomic inference and average nucleotide identity or digital DNA-DNA comparisons. As

additional *Gardnerella* lineages are isolated, whole-genome sequencing efforts should allow for

continuous integration into this framework, supporting a stable and extensible taxonomy.

Our genome-based analyses also intersect with a broader, long-standing debate regarding

the placement of *Gardnerella* within the family *Bifidobacteriaceae*. Although, marker gene

frameworks such as the GTDB place *Gardnerella* within *Bifidobacterium*, the two groups differ

markedly in genome size, G+C content, metabolic repertoire, virulence-associated gene content,

mobile genetic element composition, and ecological specialization. Notably, *Gardnerella*

genomes are substantially smaller and lack many biosynthetic and metabolic pathways retained in

larger *Bifidobacterium* genomes. Such genome reduction is widely interpreted as a hallmark of

specialization, reflecting adaptation to host-associated niches where broad metabolic versatility is

unnecessary. In this context, reduced genome size represents ecological specialization rather than

diminished biological significance.

Together, these genome-scale differences suggest that conserved marker phylogeny alone

may be insufficient to define biologically meaningful genus boundaries within this family. A

parsimonious interpretation is that *Bifidobacterium*, as currently circumscribed, encompasses

multiple genomically and ecologically distinct genera. While a formal genus-level revision of

Bifidobacteriaceae lies beyond the scope of this study, our data provide additional genome-scale

evidence supporting the continued recognition of *Gardnerella* as a distinct genus and contribute to

an evidence base that may inform future ICNP-compliant taxonomic restructuring. With this

refined taxonomic and nomenclature framework, we probed for genetic features and putative

operons associated with each lineage by applying a syntenic pangenome approach. These analyses

linked *Gardnerella* lineages to distinct repertoires of virulence-associated factors, metabolic

pathways, and genetic defense systems, highlighting functional diversity across the genus. Notably,

most *Gardnerella* gene content (84%) is species- or strain-variable with many differences related

to virulence-associated functions, metabolism, and genetic defense.

Gardnerella Set A genomes (*Gr, Gp,* and *Gv*) were enriched for sialidases and exhibited

increased potential for the catabolism of small carbon sources. We further noted lineage-specific

enrichment of established virulence-associated factors, including exported sialidases (*nanH2* in

Gr and *Gppt*; *nanH3* in *Gppk* and *Gppt*) and vaginolysin (enriched in *Gvv, Gvl, Gps* and *Gppk*).

The restricted distribution of these factors suggests functional specialization among lineages that

may contribute to mucin degradation and epithelial barrier disruption within the vaginal microbiome. In contrast, Set B genomes (*Gb, Gc, Gg, Gi, Gl, Gk, Gm,* and *Gw*) were enriched in putative virulence-associated proteins (e.g. predicted surface-associated proteins, secretion system components and factors implicated in host interaction), exhibited greater capacity for amino acid biosynthesis, showed a higher propensity to harbor the vG_gva_AB1 prophage, and encoded a broader repertoire of genetic defense mechanisms. Collectively, these features are consistent with distinct colonization strategies among Set B *Gardnerella* lineages, including enhanced persistence on epithelial surfaces, flexibility under nutrient-limited or variable resource conditions, and adaptation to competitive, phage-rich, polymicrobial niches.

Mechanistic studies to assess the roles of *Gardnerella* genetic factors, including those identified in this study, have been hindered by limited genetic tools available for *Gardnerella*. In fact, it has previously been reported that *Gardnerella* species lack mobile genetic elements, such as plasmids[30]. However, our extensive genome sequencing efforts reveal the first *Gardnerella* cryptic plasmid naturally occurring within a single cultivated strain, along with additional homologous sequences in publicly available genomes. Through the identification of this cryptic plasmid (pDNF01199S), we developed a replicative *Gardnerella-E. coli* shuttle vector (pG01-pp). Coupled with recent advances in *Gardnerella* species engineering[82,83], this discovery of extrachromosomal and self-replicative genetic elements within the *Gardnerella* genus should facilitate the development of advanced genetic tools. Combined, these foundational studies seek to advance genetic tools for mechanistic studies of *Gardnerella* species. Collectively, overcoming genetic intractability within *Gardnerella* will greatly advance our ability to investigate the role of this genus in health and disease states.

Although these interpretations are based on comparative genomic potential and do not

directly infer *in vivo* activity or clinical outcomes, taken together they support the view of

Gardnerella as an ecologically diverse genus composed of lineages with distinct metabolic

capacities, host-interaction mechanisms, and mobile genetic repertoires. Different *Gardnerella*

seem to occupy overlapping but not identical niches within their human hosts. Some lineages may

be better adapted to persistence and competitive growth in the vaginal microbiome and urinary

tract, whereas others may be more consistent with transient or context-dependent colonization.

Methods

<u>Study design and participants</u>

Vaginal bacterial isolates were collected from human study participants both with and without BV

in Seattle, Washington, USA[22,87] or from the Ravel isolate collection, harboring strains from

multiple geographic locations, including USA and Africa (Supplementary Table 2). Bacterial

isolates consistent with *Gardnerella* morphotypes on Gram stain were subjected to 16S rRNA

and/or *cpn60* gene amplicon sequencing to confirm phylogenetic placement in the *Gardnerella*

genus[24]. Phylogenetically diverse *Gardnerella* species were selected for whole-genome

sequencing. High molecular weight DNA was extracted from each isolate using the MasterPure

DNA purification kit (Lucigen; Middleton, WI), with two phenol/chloroform cleanups prior to

DNA precipitation where needed. DNA extraction was quantified on a TapeStation 2200

instrument run with a Genomic DNA tape (Agilent Technologies; Santa Clara, CA).

<u>Genome sequencing and assembly</u>

Sequencing on the Pacific Biosciences SeqI/II Platforms

A total of forty-two strains from the Fredricks collection, and seventeen from the Ravel collection, were selected for single-molecule real-time (SMRT) long-read sequencing using PacBio (SeqI or SeqII, Pacific Biosciences; Menlo Park, CA) at the Fred Hutch Cancer Center or the Genomic Resource Center of the University Maryland School of Medicine. Sequencing libraries were prepared using the SMRTBell Template Prep Kit 1.0/2.0 and were selected for a 3-5kb size cutoff using a BluePippin (Sage Science; Beverly, MA). Sequencing reads were processed via Microbial Assembly software within the Pacific Biosciences' SMRTAnalysis pipeline (v.9.0.0.92188).

Sequencing on the Illumina HiSeq 4000 Platform

332 isolates were selected for Illumina sequencing on the HiSeq4000 platform. Short read libraries were prepared using the KAPA HyperPlus Kit (Kapa Biosystems/Roche; Willmington, MA) with KAPA Single-Indexed Adapter Kit Set B. 35μL of genomic DNA was used as input, and libraries were prepared following the manufacturer's protocol with modifications based on concentration. For samples with 0.5 or 0.2 ng input DNA, the fragmentation enzyme was diluted 1:2 or 1:5 with water. All samples were fragmented at 37°C for 5 minutes. Adapter concentrations varied according to the input, and the adapter ligation was carried out overnight at 4°C for all samples. The post-ligation cleanup was performed with 0.8x Ampure XP beads (Beckman Coulter; Indianapolis IN) and 20 μL of sample was used in library amplification. Post-amplification cleanup was performed with 1x Ampure XP beads; libraries with remaining adapter dimer peaks were cleaned a second time. The final elution was in 25 μL of nuclease-free water. Libraries were run on a TapeStation instrument with a D1000 tape (Agilent Technologies; Santa Clara, CA) to assess quality and concentration. Libraries were sequenced on an Illumina HiSeq 4000 instrument using the 150 bp paired-end protocol. Reads were assembled using Unicycler.

<u>Publicly available genomes from the National Center for Biotechnology Information (NCBI)</u>

We used publicly available genomes both to provide a direct reference for previous studies and to

supplement our own sequencing efforts. We searched GenBank[88] assembly databases for the genus

Gardnerella and family *Bifidobacteraceae* (without genus assignment). Using the Type (Strain)

Genome Server (TYGS)[89] to confirm the genus of these genomes, we discovered that several of

the unassigned *Bifidobacteraceae* bacteria in NCBI were *Gardnerella* and included them in our

analysis. *Bifidobacteraceae* (without genus assignment) were included if they were collected as

part of the same NCBI BioProject which added 20 *Gardnerella* that were not typed further than

the *Bifidobacteraceae* family. Additionally, we included three sequences (350_GVAG,

476_GVAG, and 842_GVAG) that were initially assigned to *Gardnerella* but were later renamed

to *Alloscardovia* to verify this placement. The initial number of potential genomes was n = 649.

We removed genomes that were assigned outside of *Gardnerella*. Draft genomes with more than

250 contigs, modified in the laboratory, and those with truncated or missing cpn60 sequences were

removed due to inherent unreliability.

Bifidobacterium strains were collected and curated in much the same manner as the *Gardnerella*

strains. A total of 98 strains from all known species of the *Bifidobacterium* genus and the closely

related scardovial genera (e.g., *Alloscardovia, Aeriscardovia*, *Bombiscardovia*, *Pseudoscardovia*,

and *Scardovia*) were downloaded from NCBI. Whenever possible, the type strains were collected.

Sequences were assessed using CheckM, TYGS, and ANI via cutoffs similar to those used for the

Gardnerella strains. *Bifidobacteriaceae* groups were assigned according to Alessandri *et al*[90].

<u>Genome quality and comparison assessments</u>

Genome completeness and contamination for all newly sequenced and publicly available

Gardnerella genomes retrieved from NCBI were assessed using CheckM (v 1.0.18)[91]. All genomes

were annotated via RAST (v 1.073)[92]. Genomes with contamination greater than 1.5% or

completeness below 90% were excluded from downstream analyses.

Redundant genome sequences were identified based on pairwise average nucleotide identity (ANI).

Genomes sharing > 99.5% ANI with other genomes in the collection were considered duplicates

and a single representative genome was retained for each duplicate group (Supplementary Table

1). Pairwise average nucleotide identity (ANI) metrics were calculated separately for Set A and Set

B genomes using FastANI (v1.34)[93] and visualized using the R packages ComplexHeatmap (v

2.27.0)[94]. Digital DNA-DNA hybridization (dDDH)[95] values were calculated via the Genome-to-

Genome Distance Calculator (v 3.0)[96] and visualized in R (v 4.5.1 -- 2025-06-13 -- "Great Square

Root"). Amino Acid Identity (AAI) values for *Bifidobacteriaceae* type strains were determined

using pairwise comparisons of FastAAI (version 2)[44].

PPanGGOLiN *Gardnerella* pangenome

We used Partitioned PanGenome Graph of Linked Neighbors (PPanGGOLiN v 2.2.0)[52] on our

curated genome set to graph and cluster the genes from all strains, find regions of genomic

plasticity and synteny[75], and separate the pangenome gene clusters into core, shell, and cloud

partitions, with clustering set to 50% coverage and 50% identity. EggNOG[96], InterPro (accessed

October 2025)[97], and KEGG Mapper (v 5.2)[98], were run separately on each genome to gather and

confirm the CAZyme[99], Pfam[100], and KEGG (all accessed October 2025), designations of the gene

clusters. COG categories[101] were assigned based on the consensus of these designations. We added

the annotation "Unclassified Metabolism" to capture gene clusters that are clearly related to

metabolism, but the exact metabolite is not defined or could be of multiple types.

<u>Phylogenomic analyses</u>

Barrnap was used to find and extract 16S rRNA sequences for submission to NCBI[98]. The

sequences in the PPanGGOLiN for the persistent *cpn60* node were MUSCLE[99]-aligned to the same

gene from *Bifidobacterium bifidum* (ATCC 29521) to generate maximum-likelihood distances, and

a neighbor-joined dendrogram using the R package phangorn[100]. The resulting tree was visualized

and annotated using iTOL (v 7.3)[101]. kSNP4 (v 4.1)[102] was used to construct a reference-free whole-

genome phylogenetic tree for *Gardnerella* Set A (kmer=17, FCK=0.167) and *Gardnerella* Set B

(kmer=17, FCK=0.797) genomes, separately. *Bombiscardovia bombi* strain DSM19703 was

included to root both dendrograms. The resulting parsimony trees were visualized in

iTOL[101].GapMind (accessed September 2025)[103-105] metabolism predictions and PADLOC (v

2.0.0)[106,107] defense mechanism predictions were annotated where appropriate. Bacteriophage

detection was performed by VIBRANT (v. 1.2.1)[108], PHageR (accessed Feb 2026)[109], and the use

of phage-annotated nodes to find syntenic modules of phages in PPanGGOLiN. These methods

were combined as no one method found all bacteriophage. Phage sequences are mostly as found

by VIBRANT, when VIBRANT failed to find the phage sequence they were determined from

PPanGGOLiN. Phages were typed by aligning the large terminase sequences from each intact with

MAFFT (v7.526) and then creating a maximum likelihood tree with IQTree2 (v. 2.3.6)[110].

Assessment of greater taxonomic grouping was determined using VICTOR

(https://ggdc.dsmz.de/victor.php accessed March 2026)[66].

Isolation of pDNF01199S-01 plasmid

Gardnerella piotii subsp. *pickettii* strain DNF01199S was grown anaerobically in New York City

III (NYCIII) liquid media in an AS-580 anaerobic chamber (Anaerobe Systems, Morgan Hill, CA)

in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 at 37°C. A QIAamp DNA Miniprep kit (Qiagen,

Hilden, Germany) was used to isolate pDNF01199S-01 from a 5 mL culture of the bacteria.

Isolated DNA was digested with *XhoI* (New England Biolabs, Ipswich, MA) and visualized by gel

electrophoresis to confirm plasmid presence.

Gardnerella tetracycline resistance testing

Gardnerella strains were grown anaerobically in an AS-580 anaerobic chamber (Anaerobe

Systems, Morgan Hill, CA) in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 at 37°C. Bacteria

were recovered from frozen stocks on NYCIII agar plates and subcultured twice in 2 mL of NYCIII

liquid media. Culture purity was verified by Gram stain. The final cultures were normalized to 0.5

McFarland turbidity using fresh NYCIII broth and 25 µL was spotted onto an NYCIII agar plate

containing 16 µg/mL tetracycline and an antibiotic-free NYCIII agar plate. 25 µL of *Lactobacillus*

reuteri CF48-3A was spotted onto each tetracycline plate as a tetracycline-resistant control[102].

Tetracycline resistance was determined according to Clinical and Laboratory Standards Institute

for antimicrobial susceptibility testing protocols, and any growth on tetracycline NYCIII agar

plates was considered resistant to tetracycline.

pMC-pDNF01199S-*tetM* construction and minicircle preparation

pDNF01199S-01 and *tetM* were both cloned from gDNA isolated from strain DNF01199S. Gene

fragments were assembled with pMC by Gibson assembly[103]. The parental construct was

transformed into *E. coli* ZYCY10P3S2T (System Biosciences, Palo Alto, CA) by heat shocking

according to the SBI Minicircle DNA Technology user manual. Transformant colonies were

selected by growing on Luria-Burtani (LB) agar plates containing 50 µg/mL Kanamycin. A

successfully transformed colony was chosen and stocked, then induction of minicircles was

performed according to the SBI Minicircle DNA Technology user manual by growing the bacteria

in induction media containing arabinose. pDNF01199S-*tetM* minicircle DNA was then purified

from the induced *E. coli* using a Qiagen Plasmid Midi Kit (Qiagen, Hilden, Germany). The

sequence of purified plasmid was confirmed against the construct sequence by whole-plasmid

sequencing performed by Plasmidsaurus (Plasmidsaurus, Eugene, OR). "GGCC" sites on the

plasmid were protected by incubating 1mg of the DNA with 1µL of *HaeIII* MTase (New England

Biolabs, Ipswich, MA) at 37°C for 4hrs in the presence of 160 µM S-Adenosyl methionine (New

England Biolabs, Ipswich, MA). Following enzyme treatment, DNA was purified with Monarch

PCR & DNA Cleanup Kit (New England Biolabs, Ipswich, MA).

Preparation of competent *Gardnerella* cells

Gardnerella vaginalis subsp. *vaginalis* ATCC 14018[T] was grown anaerobically in an AS-580

anaerobic chamber (Anaerobe Systems, Morgan Hill, CA) in a gas mixture of 5% CO_2, 5% H_2,

and 90% N_2 at 37°C. Cells were recovered from frozen stocks on NYCIII agar plates and

subcultured into NYCIII liquid media. 500 µL of culture was subcultured into 5 mL NYCIII media

and grown overnight. 4mL of this culture was inoculated into 40mL of modified de Man-Rogosa-

Sharpe media[104] supplemented with 5 g/L glucose, 0.4 g/L L-cysteine HCl, 100 mL/L filter-

sterilized horse serum, and 100 mM NaCl when stated. Once the target OD_{600} was reached, cells

were harvested by centrifugation at 4°C at 4,052 x *g* for 10 minutes. Cells were washed in one of

the following three ways: 1) Twice in 25 mL ice-cold maltose citrate buffer (0.2625 g/L citric acid

and 0.5 M maltose at pH 5.8), 2) Twice in 25 mL ice-cold, autoclaved, 10% glycerol, or 3) Once

in 25 mL ice-cold autoclaved water from a Milli-Q IQ7000 pure water system (Millipore Sigma,

St. Louis, MO) followed by a second wash in 25 mL ice-cold maltose citrate buffer. Washed cells

were then resuspended in 200 µL of maltose citrate buffer, mixed with 200 µL 80% glycerol, and

stored at -80°C in aliquots.

Gardnerella electrotransformation and verification

Competent cells were mixed with plasmid DNA on ice and transferred into pre-chilled 0.1 cm gap

Gene Pulser/MicroPulser Electroporation Cuvettes (Bio-Rad, Hercules, CA). Electroporation was

then performed on a Gene Pulser Xcell (Bio-Rad, Hercules, CA) at 25 µF, 200 Ω, and 2,000 V.

Immediately following electroporation, 950 µL of 37°C pre-warmed NYCIII broth was added to

the cells and they were transferred to an AS-580 anaerobic chamber (Anaerobe Systems, Morgan

Hill, CA) in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 to recover at 37°C for 1 hour. 100 µL

– 200 µL of transformed cells were then plated on NYCIII agar plates with and without 16 µg/mL

tetracycline. Plates were incubated anaerobically at 37°C for 48 hours and transformed colonies

on the tetracycline plates were counted to determine transformation efficiency. After the first

successful transformation, transformant colonies were grown in 5 mL NYCIII media with 16

µg/mL tetracycline and pDNF01199S-*tetM* was recovered from them using a QIAamp DNA

Miniprep kit (Qiagen, Hilden, Germany). The sequence of the purified plasmid was confirmed by

sequencing performed by Plasmidsaurus (Plasmidsaurus, Eugene, OR). Following transformation

of *Gardnerella vaginalis* subspecies *regina* strain DNF00354 with pDNF01199S-*tetM*, the

transformation reaction was plated onto NYCIII agar plates with and without 16 µg/mL

tetracycline. A small amount of growth from a transformant colony on the tetracycline plate or

lawn growth from the control plate was resuspended in sterile water and colony PCR was

performed to amplify and detect the *tetM* gene.

One microliter of resuspended bacteria was added to a reaction mix of 12.5 µL GoTaq Green master

mix (Promega, Madison, WI), 0.5 µL of each primer, and 6.5 µL H_2O. PCR was carried out for 28

cycles with an annealing temperature of 60°C and an extension time of 2.5 min. A detailed version

of this protocol is available online at:

https://www.protocols.io/private/1A47BAD2F1AC11F09AA30A58A9FEAC02.

Primer sequences were as follows (5' to 3'):

Forward: CGCGACCAAATATTGGTACATTATTACAGCTATTTTGTAATCACGTACTC

Reverse: GAGACTGTCGTATTGATGTCACGGGCTTAATAGGTCTTATGTCGG

Protologues

Description of *Gardnerella vaginalis* subsp. *vaginalis* Gardner and Dukes 1955 (Approved

Lists 1980) Bouzek *et al*. comb. nov.

Basonym: *Haemophilus vaginalis* Gardner and Dukes 1955 (Approved Lists 1980)[9]. Basonym:

Haemophilus vaginalis Gardner and Dukes 1955 (Approved Lists 1980)[9].

Earlier homotypic synonym: *Gardnerella vaginalis* (Gardner and Dukes 1955) Greenwood and

Pickett 1980[114]. Earlier homotypic synonym: *Gardnerella vaginalis* (Gardner and Dukes 1955)

Greenwood and Pickett 1980[111].

Gardnerella vaginalis subsp. *vaginalis* (va.gi.na'lis. L. fem. n. *vagina*, sheath, vagina; L. masc.

adj. suff. -*alis,* suffix denoting pertaining to; N.L. fem. adj. *vaginalis*, pertaining to vagina, of the

vagina).

The description of this subspecies remains as given to *Gardnerella vaginalis* by Gardner and Dukes

1955[9] and Greenwood and Pickett 1980[111] with the following additional information. Curated

genomes of 79 *Gardnerella vaginalis* subsp. *vaginalis* cultivated isolates were used in this analysis,

isolated from the vagina, endometrium, cervix, catheter urine, and placenta of humans from

Bangladesh, Belgium, China, France, Russia, the United Kingdom, the United States, and Zambia

(Supplementary Table 2). The mean genome length of *Gardnerella vaginalis* subsp. *vaginalis* is

1.68 Mb [range1.58 – 1.78] and the median G+C content is 41.26% [range 40.99% – 41.82%]. The

type strain for *Gardnerella vaginalis* subsp. *vaginalis* remains 594^T of Gardner and Dukes (=ATCC

14018^T=DSM 4944^T=CCUG 3717^T=JCM 11026^T=NCTC 10915^T=NCTC 10287^T).

Strains used in previous classifying publications: ATCC14019, ATCC49145, DNF01149,

FDAARGOS296, FDAARGOS568, GH015, GH021, HMP9231, JCM11026, JCP7276, JCP7672,

NR001, NR037, NR038, NR039, UGent09.01, UGent09.07, UGent25.49, UMB0032A,

UMB0032B, UMB0061, UMB0233, UMB0298, UMB0386, UMB0768, UMB0770, UMB0775,

WP023, 287, 377, 653, 1492, 3549624, 0284V, 0286E, 0288E, 1-010-6, 1-014-4, 104-V2-10,

1388E, 18-4, 203-8, 23-12, 302000480-V1-10, 315-A, 7571-2, 772-1, and 825-1.

Description of *Gardnerella vaginalis* subsp. *lacydonensis* (Allini Ntiguemassa *et al.* 2026) Bouzek *et al.* comb. nov.

Basonym: *Gardnerella lacydonensis* Allini Ntiguemassa *et al*. 2026[49] Basonym: *Gardnerella lacydonensis* Allini Ntiguemassa *et al*. 2026[49]

Gardnerella vaginalis subsp. *lacydonensis* (la.cy.do.nen'sis. N.L. fem. adj. *lacydonensis*, from Lacydon, the name of the ancient port of Marseille, the French city where the strain was first described).

The description of *Gardnerella vaginalis* subsp. *lacydonensis* remains as given to *Gardnerella lacydonensis* by Allini Ntiguemassa *et al*.[49] with the following additional information. Our collection of curated genomes contains 15 *Gardnerella vaginalis* subsp. *lacydonensis* cultivated strains, isolated from the vagina, endometrium, and ectocervical mucosa of humans from France, Kenya, South Africa, the United States, and Zambia (Supplementary Table 2). The median genome length of these isolates is 1.66 Mb [range 1.62 – 1.72] and the median G+C content is 41.29% [range 41.11% – 41.42%]. The type strain of *Gardnerella vaginalis* subsp. *lacydonensis* remains Marseille-Q9181^T (=CSUR Q9181^T=CECT 31121^T).

Description of *Gardnerella vaginalis* subsp. *regina* subsp. nov.

Gardnerella vaginalis subsp. *regina* (re.gi'na. L. fem. n. *regina*, the queen, referring to its placement in the first validly published species of the *Gardnerella* genus).

We analyzed 17 curated genomes from cultivated strains of *Gardnerella vaginalis* subsp. *regina*,

isolated from the vagina of humans from Bangladesh, France, the United States, and Zambia

(Supplementary Table 2). The median genome length of the *Gardnerella vaginalis* subsp. *regina*

is 1.63 Mb [range 1.57 – 1.67], and the median G+C content is 41.41% [range 41.25% – 41.62%].

The isolate is a facultative anaerobe and has gram variable cells. Growth develops on Brucella agar

in anaerobic conditions at 37°C after one to two days. In a genome comparison of *Gardnerella*

vaginalis subsp. *regina* strain DNF00622A^T with validly published *Gardnerella* type strains and

the type strains proposed in this study, the highest ANI values correspond with its sister subspecies

Gardnerella vaginalis subsp. *vaginalis* ATCC14018^T (96.82%) and *Gardnerella vaginalis* subsp.

lacydonensis Marseille-Q9181^T (96.17%), followed by *Gardnerella piotii* subsp. *piotii* UGent

18.01^T (89.24%) (Figure 2). Within the genome, 83 CAZymes were identified, including α-

mannosidase (GH125), β-galactosidase (GH2), β-mannosidase (GH130), Class II α-mannosidase

(GH38), β-N-acetylglucosaminidase (GH20), α-L-fucosidase (GH29), and sialidase (GH33)[112].

KEGG analysis identified a total of 97 transporters, 13 secretion genes, and 365 enzymes[112].

Metabolism predictions using GapMind[103-105] include biosynthesis of cysteine, glutamine, glycine,

and proline and catabolism of ethanol, gluconate, and serine (Supplementary Table 6). The type

strain for *Gardnerella vaginalis* subsp. *regina* is DNF00622A^T, isolated from the vaginal fluid of

a woman positive for bacterial vaginosis in Seattle, Washington, USA. Accessions for the 16S

rRNA gene sequence and whole genome sequence for DNF00622A^T are PZ013896 and CP058395.

Strains used in previous classifying publications: 1-004-3, 819-1, and 819-3.

Description of *Gardnerella piotii* subsp. *piotii* (Vaneechoutte *et al.* 2019) Bouzek *et al.* comb.

nov.

Basonym: *Gardnerella piotii* Vaneechoutte *et al.* 2019[32]

Gardnerella piotii (pi.o'ti.i. N.L. gen. masc. n. *piotii*, of Piot, named after Peter Piot, a Belgian

microbiologist who first attempted to distinguish different groups within *G. vaginalis*).

The description of *Gardnerella piotii* subsp. *piotii* remains as given to *Gardnerella piotii* by

Vaneechoutte *et al.*[32] with the following additional information. There are 17 genomes of cultivated

Gardnerella piotii subsp. *piotii* in our curated collection, isolated from the vagina and endometrium

of humans from Bangladesh, Belgium, China, France, and the United States (Supplementary Table

2). The median genome length of these isolates is 1.54 Mb [range 1.50 – 1.59] and the median

G+C content is 42.43% [range 42.22% – 42.61%]. The *Gardnerella piotii* subsp. *piotii* type strain

remains UGent 18.01^T (= LMG 30818^T=CCUG 72427^T).

Description of *Gardnerella piotii* subsp. *pickettii* (Sousa *et al.* 2023) Bouzek *et al.* comb. nov.

Basonym: *Gardnerella pickettii* Sousa *et al.* 2023[48]

Gardnerella piotii subsp. *pickettii* (pic.ket'ti.i. N.L. gen. masc. n. *pickettii*, of Pickett, named after

M. John Pickett, an American microbiologist who proposed the transfer of *Haemophilus vaginalis*

to the new genus *Gardnerella*).

The description of *Gardnerella piotii* subsp. *pickettii* remains as given to *Gardnerella pickettii* by

Sousa *et al.*[48] with the following additional information. We included the curated genomes of 40

cultivated isolates of *Gardnerella piotii* subsp. *pickettii* in our analysis, isolated from the vagina,

endometrium, ectocervical mucosa, urine, bladder and rectum of humans from Bangladesh,

Canada, China, Kenya, Portugal, South Africa, the United States, and Zambia (Supplementary

Table 2). The median genome length of *Gardnerella piotii* subsp. *pickettii* is 1.56 Mb [range 1.51

– 1.68] and the median G+C content is 42.32% [range 42.00% - 42.74%]. The type strain of

Gardnerella piotii subsp. *pickettii* remains c17Ua_112^T (=DSM 113414^T=CCP 71^T).

Description of *Gardnerella piotii* subsp. *sialicivorans* subsp. nov.

Gardnerella piotii subsp. *sialicivorans* (si.a.li.ci.vo'rans. N.L. neut. n. *acidum sialicum*, sialic acid;

L. pres. part. *vorans*, devouring; N.L. fem. part. adj. *sialicivorans*, sialic acid devouring).

The subspecies *Gardnerella piotii* subsp. *sialicivorans* is named for the intact sialidase catabolism

operon and one or more sialidase genes present in the genomes of *Gardnerella piotii*. Our

collection of curated genomes includes three *Gardnerella piotii* subsp. *sialicivorans* cultivated

strains, all isolated from the vagina of humans from the United States (Supplementary Table 2).

The median genome length is 1.51 Mb [range 1.50 – 1.53], and the median G+C content is 42.48%

[range 42.39% - 42.48%]. This isolate is a gram-positive coccus and produces pinpoint colonies

(≤ 1 mm) on Brucella agar after three days of anaerobic incubation at 37˚C. Colonies are grey,

pleomorphic, and alpha-hemolytic. In a genome comparison of *Gardnerella piotii* subsp.

sialicivorans strain C0013B2^T with validly published *Gardnerella* type strains and the type strains

proposed in this study, the highest ANI values correspond with members of its sister subspecies

Gardnerella piotii subsp. *pickettii* c17Ua_112^T (95.68%) and *Gardnerella piotii* subsp. *piotii*

UGent 18.01^T (94.66%), followed by *Gardnerella rara* C0101A1^T (90.88%) (Figure 2). Within the

genome, 76 CAZymes were identified, including sialidase (GH33) and Class II α-mannosidase

(GH38)[112]. KEGG analysis identified a total of 79 transporters, 13 secretion genes, and 355

enzymes[112]. Metabolism predictions using GapMind[103-105] include biosynthesis of glutamine,

glycine, and proline, and catabolism of alanine, ethanol, and serine (Supplementary Table 6). The

Gardnerella piotii subsp. *sialicivorans* type strain is C0013B2^T, isolated from the vagina of an

asymptomatic menstruating woman in Birmingham, Alabama, USA. Accession for the 16S rRNA

gene sequence for C0013B2^T is PZ016854.

Strain used in previous classifying publications: 582.

Description of *Gardnerella rara* sp. nov.

Gardnerella rara (ra'ra. L. fem. adj. *rara*, rare, uncommon, referring to its rare genitourinary

colonization of humans).

Our curated collection includes six genomes of *Gardnerella rara* from isolates cultivated from the

vagina and endometrium of humans in Bangladesh, the United States, and Zambia (Supplementary

Table 2). The median genome length of *Gardnerella rara* is 1.52 Mb [range 1.47 – 1.58] and the

median G+C content is 43.23% [range 43.01% – 43.29%]. This isolate is a gram-positive bacillus

and produces small colonies on Brucella agar after two days of anaerobic incubation at 37˚C.

Colonies are grey, round, and alpha-hemolytic. In a genome comparison of *Gardnerella rara* strain

C0101A1^T with validly published *Gardnerella* type strains and the type strains proposed in this

study, the highest ANI values correspond with *Gardnerella piotii* subsp. *piotii* UGent 18.01^T at

91.38% (Figure 2). Within the genome, 71 CAZymes were identified, including β-mannosidase

(GH130), Class II α-mannosidase (GH38), and sialidase (GH33)[112]. KEGG analysis identified a

total of 78 transporters, 13 secretion genes, and 344 enzymes[112]. Metabolism predictions using

GapMind[103-105] include biosynthesis of glutamine, glycine, and proline, and catabolism of alanine

and ethanol (Supplementary Table 6). The type strain of *Gardnerella rara* type is C0101A1^T,

isolated from the vagina of a woman positive for *Chlamydia trachomatis* in Baltimore, Maryland,

USA. Accession for the 16S rRNA gene sequence for C0101A1^T is PZ016855.

Strains used in previous classifying publications: 1401E, GED7760B, and HUZ601048-2.

Description of *Gardnerella leopoldii* subsp. *leopoldii* (Vaneechoutte *et al*. 2019) Bouzek *et al*.

comb. nov.

Basonym: *Gardnerella leopoldii* Vaneechoutte *et al*. 2019[32]

Gardnerella leopoldii (le.o.pol'di.i. N.L. gen. masc. n. *leopoldii,* of Leopold, named after Sidney

Leopold, who described the first strains of the species).

The description of this subspecies remains as given to *Gardnerella leopoldii* by Vaneechoutte *et*

al.[32] with the following additional information. There are 24 genomes of cultivated isolates of

Gardnerella leopoldii subsp. *leopoldii* in our curated collection, isolated from the vagina and

catheter urine of humans from Belgium, Canada, France, Kenya, and the United States

(Supplementary Table 2). The median genome length is 1.53 Mb [range 1.37 – 1.67], and the

median G+C content is 42.15% [range 42.07% - 42.34%]. The type strain of *Gardnerella leopoldii*

subsp. *leopoldii* remains UGent 06.41^T (=LMG 30814^T=CCUG 72425^T).

Description of *Gardnerella leopoldii* subsp. *swidsinskii* (Vaneechoutte *et al*. 2019) Bouzek *et*

***al*. comb. nov.**

Basonym: *Gardnerella swidsinskii* Vaneechoutte *et al*. 2019[32]

Gardnerella swidsinskii (swid.sins'ki.i. N.L. gen. masc. n. *swidsinskii*, of Swidsinski, named after

Alexander Swidsinski, a German microbiologist, who recognized that clue cells in Gram stains of

vaginal swabs of women with BV are indicative for biofilm formation by *Gardnerella*).

The description of this subspecies remains as provided by Vaneechoutte *et al*. for *Gardnerella*

swidsinskii[32] with the following additional information. We included 46 genomes of cultivated

Gardnerella leopoldii subsp. *swidsinskii* in our curated collection, isolated from the vagina,

endometrium, rectum, and catheter urine of humans from China, France, Russia, and the United

States (Supplementary Table 2). The median genome length is 1.63 Mb [range 1.52 – 1.72], and

the median G+C content is 41.93% [range 41.64% - 42.29%]. The type strain of *Gardnerella*

leopoldii subsp. *swidsinskii* remains GS 9838-1^T (=LMG 30812^T=CCUG 72429^T).

Description of *Gardnerella leopoldii* subsp. *aggregata* subsp. nov.

Gardnerella leopoldii subsp. *aggregata* (ag.gre.ga'ta. L. fem. part. adj. *aggregata*, joined together,

referring to the presence of genes related to cell aggregation).

The genomes of *Gardnerella leopoldii* subsp. *aggregata*, as well as those of the overarching

species, differ from other *Gardnerell*a species in the presence of genes consistent with adherence

and biofilm formation. There are five genomes of cultivated *Gardnerella leopoldii* subsp.

aggregata in our curated collection, isolated from the vagina and ectocervical mucosa of humans

from Bangladesh, South Africa the United States, and Zambia (Supplementary Table 2). The

median genome length is 1.48 Mb [range 1.48 – 1.53], and the median G+C content is 42.19%

[range 42.11% - 42.50%]. This isolate is a gram variable coccobacillus and produces pinpoint

colonies (≤ 1 mm) on Brucella agar after three days of anaerobic incubation at 37˚C. Colonies are

glossy white, round, convex, and alpha-hemolytic. In a genome comparison of *Gardnerella*

leopoldii subsp. *aggregata* strain C0026F5[T] with validly published *Gardnerella* type strains and

the type strains proposed in this study, the highest ANI values correspond with its sister subspecies

Gardnerella leopoldii subsp. *phocaeensis* Marseille-Q9179[T] (96.09%), *Gardnerella leopoldii*

subsp. *leopoldii* UGent 06.41[T] (95.58%), and *Gardnerella leopoldii* subsp. *swidsinskii* GS 9838-

1[T] (94.70%), followed by *Gardnerella greenwoodii* subsp. *bretellae* Marseille-QA0894[T] (87.40%)

(Figure 2). Within the genome, 74 CAZymes were identified, including α-glucosidase (GH65),

endo-β-galactosidase (GH16), and β-glucosidase (GH3)[112]. KEGG analysis identified a total of 84

transporters, 14 secretion genes, and 353 enzymes[112]. Metabolism predictions using GapMind[103-

105] include biosynthesis of glutamine, glycine, methionine, proline, and threonine, and catabolism

of ethanol (Supplementary Table 6). The type strain of *Gardnerella leopoldii* subsp. *aggregata* is

C0026F5[T], isolated from the vagina of an asymptomatic menstruating woman from Birmingham,

Alabama, USA. Accession for the 16S rRNA gene sequence for C0026F5[T] is PZ016853.

Description of *Gardnerella leopoldii* subsp. *phocaeensis* (Allini Ntiguemassa *et al*. 2026)

Bouzek *et al*. comb. nov.

Basonym: *Gardnerella phocaeensis* Allini Ntiguemassa *et al*. 2026[49]

Gardnerella phocaeensis (pho.cae.en'sis. L. fem. adj. *phocaeensis*, referring to Phocaea, the name

of the Ionian Greek city where the founders of Marseille came from. The strain was isolated in

Marseille).

The description of *Gardnerella leopoldii* subsp. *phocaeensis* remains as given to *Gardnerella*

phocaeensis by Allini Ntiguemassa *et al.*[49] with the following additional information. There are

eleven genomes of cultivated *Gardnerella leopoldii* subsp. *phocaeensis* in our curated collection,

isolated from the vagina and endometrium of humans from Bangladesh, France, the United States,

and Zambia (Supplementary Table 2). The median genome length of *Gardnerella leopoldii* subsp.

phocaeensis is 1.54 Mb [range 1.52 – 1.66] and the median G+C content is 42.26% [range 42.03%

- 42.43%]. The type strain of *Gardnerella leopoldii* subsp. *phocaeensis* remains Marseille-Q9179[T]

(=CSUR Q9179[T]=CECT 31120[T]).

Description of *Gardnerella greenwoodii* subsp. *greenwoodii* (Sousa *et al.* 2023) Bouzek *et al.*

comb. nov.

Basonym: *Gardnerella greenwoodii* Sousa *et al.* 2023[48]

Gardnerella greenwoodii (green.woo'di.i. N.L. gen. masc. n. *greenwoodii*, of Greenwood, named

after James Greenwood, an American microbiologist who proposed the transfer of *Haemophilus*

vaginalis to the new genus *Gardnerella*).

The description of this subspecies remains as given to *Gardnerella greenwoodii* by Sousa *et al.*[48]

with the following additional information. We included 13 genomes of *Gardnerella greenwoodii*

subsp. *greenwoodii* in our curated collection, cultivated from the vagina, endometrium, and urine

of humans from France, Portugal, the United Kingdom, and the United States (Supplementary

Table 2). The median genome length of *Gardnerella greenwoodii* subsp. *greenwoodii* is 1.54 Mb

[range 1.49 – 1.65] and the median G+C content is 43.40% [range 42.90% - 43.44%]. The type

strain of *Gardnerella greenwoodii* subsp. *greenwoodii* remains c31Ua_26^T (=DSM 113415^T=CCP

72^T).

Description of *Gardnerella greenwoodii* subsp. *intermedia* subsp. nov.

Gardnerella greenwoodii subsp. *intermedia* (in.ter.me'di.a. L. fem. adj. *intermedia*, intermediate,

between, referencing the shared possession of operons with both *Gardnerella* subgenera).

Pangenome analysis revealed that *Gardnerella greenwoodii* subsp. *intermedia*, as well as other

members of the species, shares biologically relevant operons with the Set A subgenus (*Gardnerella*

vaginalis, *Gardnerella piotii* and *Gardnerella rara*) while also falling within the Set B subgenus,

suggesting that the species is an intermediate between Set A and Set B. Our curated set of genomes

contains two strains of *Gardnerella greenwoodii* subsp. *intermedia*, both cultivated from the vagina

of humans from the United States (Supplementary Table 2). The lengths of these genomes are 1.53

and 1.55 Mb, and the G+C contents are 43.04% and 43.05%. Cells are gram variable. Optimal

growth occurs within two to three days on Brucella agar in an anerobic atmosphere at 37°C but

can occur in aerobic environments. In a genome comparison of *Gardnerella greenwoodii* subsp.

intermedia strain KA00603^T with validly published *Gardnerella* type strains and the type strains

proposed in this study, the highest ANI values correspond with sister subspecies *Gardnerella*

greenwoodii subsp. *bretellae* Marseille-QA0894^T (95.76%) and *Gardnerella greenwoodii* subsp.

greenwoodii c31Ua_26^T (95.04%), followed by *Gardnerella kenyensis* NR047^T (88.32%) (Figure

2).

Within the genome, 66 CAZymes were identified, including β-glucosidase (GH3), and KEGG

analysis identified a total of 85 transporters, 13 secretion genes, and 347 enzymes[112]. Metabolism

predictions using GapMind[103-105] include biosynthesis of cysteine, glutamine, glycine, methionine,

proline, and threonine, and catabolism of ethanol and serine (Supplementary Table 6). The type

strain of *Gardnerella greenwoodii* subsp. *intermedia* is KA00603^T, isolated from the vaginal fluid

of a woman positive for bacterial vaginosis in Seattle, WA, USA. Accessions for the 16S rRNA

gene sequence and whole genome sequence for KA00603^T are PZ013893 and CP058371.

Strain used in previous classifying publications: 1500E.

Description of *Gardnerella greenwoodii* subsp. *bretellae* (Allini Ntiguemassa *et al*. 2026)

Bouzek *et al*. comb. nov.

Basonym: *Gardnerella bretellae* Allini Ntiguemassa *et al*. 2026

Gardnerella bretellae (bre.tel'lae. N.L. gen. fem. n. *bretellae*, honoring Florence Bretelle for her

contribution to the description of vaginal microbiota).

The description of *Gardnerella greenwoodii* subsp. *bretellae* remains as given to *Gardnerella*

bretellae by Allini Ntiguemassa *et al*.[49] with the following additional information. Our curated

collection of genomes contains eleven strains of *Gardnerella greenwoodii* subsp. *bretellae*,

cultivated from the vagina and ectocervical mucosa of humans from Canada, France, Kenya, South

Africa, the United States, and Zambia (Supplementary Table 2). The median genome length is 1.52

Mb [range 1.50 – 1.64], and the median G+C content is 43.22% [range 43.05% - 43.50%]. The

Gardnerella greenwoodii subsp. *bretellae* type strain remains Marseille-QA0894^T (=CSUR

QA0894^T=CECT 31122^T).

Description of *Gardnerella washingtoni* sp. nov.

Gardnerella washingtoni (wa.shing.to'ni. N.L. gen. n. *washingtoni*, named in honor of Washington

University in St. Louis, MO, USA, where the first two genomes of this species originated, and the

University of Washington in Seattle, WA, USA, which is associated with the first complete genome

of this species).

There are five genomes of *Gardnerella washingtoni* in our curated collection. These isolates have

been cultivated from the vagina and urine of humans in the United States (Supplementary Table

2). The median genome length of *Gardnerella washingtoni* is 1.60 Mb [range 1.56 – 1.65] and the

median G+C content is 42.96% [range 42.89% – 43.11%]. Cells are Gram variable. The isolate is

a facultative anaerobe that grows on Brucella agar in anaerobic conditions at 37°C in one to two

days. In a genome comparison of *Gardnerella washingtoni* strain DNF00536^T with validly

published *Gardnerella* type strains and the type strains proposed in this study, *Gardnerella*

greenwoodii subsp. *bretellae* Marseille-QA0894^T has the highest ANI value (87.40%) (Figure 2).

Within the genome, 69 CAZymes were identified, including endo-β-galactosidase (GH16), α-

glucosidase (GH65), and β-glucosidase (GH3)[112]. KEGG analysis identified a total of 83

transporters, 13 secretion genes, and 348 enzymes[112]. Metabolism predictions using GapMind[103-

105] include biosynthesis of cysteine, glutamine, glycine, methionine, and threonine, and catabolism

of ethanol (Supplementary Table 6). The type strain of *Gardnerella washingtoni* is DNF00536^T,

isolated from the vaginal fluid of a woman positive for bacterial vaginosis in Seattle, WA, USA.

Accessions for the 16S rRNA gene sequence and whole genome sequence for DNF00536^T are

PZ013897 and CP058398.

Strains used in previous classifying publications: JCP8481A, JCP8481B, and PSS7772B.

Description of *Gardnerella bivia* sp. nov.

Gardnerella bivia (bi'vi.a. L. fem. adj. *bivia*, having two ways, in reference to the genome

containing genes for two major pathways for mucin degradation).

Gardnerella bivia has genes associated with two major pathways for mucin degradation. There are

six isolates of this species in our curated collection, cultivated from the vagina of humans in

Bangladesh, the United States, and Zambia (Supplementary Table 2). The median genome length

of *Gardnerella bivia* is 1.59 Mb [range 1.55 – 1.64] and the median G+C content is 38.08% [range

37.90% – 38.31%]. The isolate is a gram variable obligate anaerobe. Growth occurs on Brucella

agar within one to two days in anaerobic conditions at 37°C. In a genome comparison of

Gardnerella bivia strain DNF00502[T] with validly published *Gardnerella* type strains and the type

strains proposed in this study, *Gardnerella hutchinsoni* KA00225[T] has the highest ANI value

(82.91%) (Figure 2). Within the genome, 75 CAZymes were identified, including β-galactoside

phosphorylase (GH112), α-L-fucosidase (GH29), and β-glucosidase (GH3)[112]. KEGG analysis

identified a total of 74 transporters, 13 secretion genes, and 349 enzymes[112]. Metabolism

predictions using GapMind[103-105] include biosynthesis of cysteine, glutamine, glycine, proline, and

threonine, and catabolism of acetate, ethanol, and serine (Supplementary Table 6). The type strain

of *Gardnerella bivia* is DNF00502[T], isolated from the vaginal fluid of a woman positive for

bacterial vaginosis in Seattle, WA, USA. Accessions for the 16S rRNA gene sequence and whole

genome sequence for DNF00502[T] are PZ013890 and CP058399.

Strain used in previous classifying publications: CMW7778B.

Description of *Gardnerella hutchinsoni* sp. nov.

Gardnerella hutchinsoni (hutch.in.so'ni. N.L. gen. n. *hutchinsoni*, named in honor of Fred

Hutchinson Cancer Center, where the isolate was first cultivated and the first complete genomes

of this subspecies were sequenced).

We included six genomes of *Gardnerella hutchinsoni* isolates in our curated collection, all

cultivated from the vagina of humans in the United States and Zambia (Supplementary Table 2).

The median genome length of *Gardnerella hutchinsoni* is 1.67 Mb [range 1.64 – 1.72] and the

median G+C content is 40.64% [range 40.53% – 40.76%]. Cells are gram variable. Optimal growth

occurs within one to two days on Brucella agar in an anerobic atmosphere at 37°C but can occur

in aerobic environments. In a genome comparison of *Gardnerella hutchinsoni* strain KA00225^T

with validly published *Gardnerella* type strains and the type strains proposed in this study,

Gardnerella leopoldii subsp. *aggregata* C0026F5^T has the highest ANI value (86.69%) (Figure 2).

Within the genome, 69 CAZymes were identified, including β-galactoside phosphorylase (GH112)

and β-glucosidase (GH3)[112]. KEGG analysis identified a total of 70 transporters, 13 secretion

genes, and 341 enzymes[112]. Metabolism predictions using GapMind[103-105] include biosynthesis of

asparagine, glutamine, glycine, methionine, and proline, and catabolism of ethanol (Supplementary

Table 6). The type strain of *Gardnerella hutchinsoni* is KA00225^T, isolated from the vaginal fluid

of a woman positive for bacterial vaginosis in Seattle, WA, USA. Accessions for the 16S rRNA

gene sequence and whole genome sequence for KA00225^T are PZ013891 and CP058376.

Strains used in previous classifying publications: 3-005-3V and HUZ601990-0.

Description of Candidatus *Gardnerella kenyensis* sp. nov.

Gardnerella kenyensis (ken.yen'sis. N.L. fem. adj. *kenyensis*, from Kenya, referring to the country

of isolation of the type strain).

All known strains of Candidatus *Gardnerella kenyensis* to date (NR002, NR021, and NR047) have

been isolated from the vaginal fluid of a human in Nairobi, Kenya (Supplementary Table 2). Two

of these are genomic duplicates (NR002 and NR047) and one (NR021) did not pass our genome

quality threshold metrics. As such, only NR047^T was included in our curated collection

(Supplementary Table 2). The genome length of this isolate is 1.66 Mb, and its G+C content is

43.62%. In a genome comparison of Candidatus *Gardnerella kenyensis* strain NR047T with validly

published *Gardnerella* type strains and the type strains proposed in this study, the highest ANI

values correspond with *Gardnerella insolita* subsp. *libera* C0085C4^T (90.59%) (Figure 2). Within

the genome, 66 CAZymes were identified, including sialidase (GH33), and KEGG analysis

identified a total of 84 transporters, 13 secretion genes, and 340 enzyme[112]. Metabolism predictions

using GapMind[103-105] include biosynthesis of asparagine, cysteine, glutamine, glycine, methionine,

proline, and threonine, and catabolism of ethanol (Supplementary Table 6). The type strain of

Candidatus *Gardnerella kenyensis* is NR047^T and the accession for the whole genome sequence is

NQOH00000000.2.

Description of *Gardnerella insolita* sp. nov.

Gardnerella insolita (in.so'li.ta. L. fem. adj. *insolita*, unusual, strange, uncommon, referring to the

low number of species cultivated).

To date, only three strains of *Gardnerella insolita* have been cultivated, all isolated from the vagina

of humans in the United States and Zambia (Supplementary Table 2). The median genome length

of *Gardnerella insolita* is 1.61 Mb [range 1.59 – 1.66] and the median G+C content is 43.60%

[range 41.27% – 43.80%]. The *Gardnerella insolita* type strain is C0396A3^T, isolated from the

vagina of a pregnant woman at 19 weeks of gestation and who delivered preterm in Zambia.

Accession for the 16S rRNA gene sequence C0396A3^T is PZ236487.

Description of *Gardnerella insolita* subsp. *insolita* comb. nov.

Basonym: *Gardnerella insolita* Bouzek *et al*. 2026

Gardnerella insolita (in.so'li.ta. L. fem. adj. *insolita*, unusual, strange, uncommon, referring to the

low number of species cultivated).

There are two strains of *Gardnerella insolita* subsp. *insolita* in our curated genome collection. The

genome lengths of these isolates are 1.66 and 1.61 Mb, and the G+C contents are 41.27% and

43.80%. This isolate is a gram-positive bacillus and produces pinpoint ($\leq$ 1 mm) colonies on

Brucella agar after two days of anaerobic incubation at 37˚C. Colonies are round, clear, and glossy.

In a genome comparison of *Gardnerella insolita* subsp. *insolita* strain C0396A3^T with validly

published *Gardnerella* type strains and the type strains proposed in this study, the highest ANI

values correspond with *Gardnerella insolita* subsp. *libera* C0085C4^T (95.24%), followed by

Gardnerella kenyensis NR047^T (89.89%) (Figure 2). Within the genome, 66 CAZymes were

identified, including β-glucosidase (GH3)[112]. KEGG analysis identified a total of 85 transporters,

13 secretion genes, and 347 enzymes[112]. Metabolism predictions using GapMind[103-105] include

biosynthesis of asparagine, glutamine, glycine, proline, and threonine, and catabolism of ethanol

(Supplementary Table 6).

Strain used in previous classifying publications: 1-004-1.

Description of *Gardnerella insolita* subsp. *libera* subsp. nov.

Gardnerella insolita subsp. *libera* (li'be.ra. L. fem. adj. *libera*, independent, free, referring to the rarity of the subspecies).

To date, only one strain of *Gardnerella insolita* subsp. *libera* has been cultivated, C0085C4^T, from the vagina of a woman positive for *Chlamydia trachomatis* in Baltimore, Maryland, USA. Its genome length is 1.59 Mb, and its G+C content is 43.60%. This isolate is a gram variable coccobacillus and produces pinpoint colonies (≤1 mm) on Brucella agar after two days of anaerobic growth at 37˚C. Colonies are grey, glistening, round, and β-hemolytic. In a genome comparison of *Gardnerella insolita* subsp. *libera* strain C0085C4^T with validly published Gardnerella type strains and the type strains proposed in this study, the highest ANI values correspond with *Gardnerella insolita* subsp. *insolita* C0396A3^T (95.30%), followed by Candidatus *Gardnerella kenyensis* NR047^T (90.82%) (Figure 2). Within the genome, 70 CAZymes were identified, including Class II α-mannosidase and α-glucosidase[112]. KEGG analysis identified a total of 84 transporters, 12 secretion genes, and 342 enzymes[112]. Metabolism predictions using GapMind[103-105] include biosynthesis of cysteine, glutamine, glycine, proline, and methionine, and catabolism of ethanol and serine (Supplementary Table 6). The type strain of *Gardnerella insolita* subsp. *libera* is C0085C4^T. Accession for the 16S rRNA gene sequence for C0085C4^T is PZ016852.

Acknowledgements

Research reported in this publication was supported in part by the National Human Genome Research Institute as part of the Human Microbiome Project Initiative focused on technology development under award number R01HG005816 (D.N.F.), the National Institute for Allergy and Infectious Diseases of the National Institutes of Health under award number U19AI084044 (J.R.), the National Institute of Dental and Craniofacial Research (NIDCR) of the National Institutes of Health Office of the Director (OD) Common Fund under award number R01DE027850 (C.D.J). and the Gates Foundation under award INV010470 to (J.R.). In addition, startup funds were provided by Fred Hutch Cancer Center and MD Anderson Cancer Center (C.D.J.).

Contributions

H.K.B., S.S., C.D.J., and D.N.F. conceived the study. E.F.M., D.S.J., S.M.S., T.F.L., E.M.L., and M.F. conducted experiments including strain isolation, DNA extraction, whole-genome sequencing, cloning, and testing of the shuttle vector. H.K.B. and M.Z.R. conducted all bioinformatics and curated data. S.S., C.D.J, J.R., and D.N.F. supervised and obtained funding for the project. H.K.B., M.Z.R., S.S., S.M.S., C.D.J and D.N.F. wrote the original draft. J.R. and M.K. provided critical edits to the manuscript and assisted in nomenclature. M.K. provided crucial nomenclature feedback and provided edits to the protologues and manuscript. All authors were involved in reviewing and editing this submission.

Data availability

NCBI Accession numbers of genomes used are available in Supplementary Table 1.

Declaration of interests

1004

1005   C.D.J is an inventor on intellectual property related to SyngenicDNA/SyMPL technology licensed

1006   by Fred Hutchinson Cancer Center to Azitra Inc., which received compensation related to this

1007   license, and C.D.J serves as a consultant to Azitra Inc. J.R. is the cofounder of LUCA Biologics.

1008   Azitra Inc. and LUCA Biologics had no role in this study. The remaining authors declare no

1009   competing interests.

References

1010

1   Fredricks, D. N., Fiedler, T. L. & Marrazzo, J. M. Molecular Identification of Bacteria
Associated with Bacterial Vaginosis. *New England Journal of Medicine* **353**, 1899-1911
(2005). https://doi.org/10.1056/nejmoa043802
2   Amsel, R. *et al.* Nonspecific vaginitis. Diagnostic criteria and microbial and
epidemiologic associations. *Am J Med* **74**, 14-22 (1983). https://doi.org/10.1016/0002-
9343(83)91112-9
3   Harwich, M. D. *et al.* Drawing the line between commensal and pathogenic Gardnerella
vaginalis through genome analysis and virulence studies. *BMC Genomics* **11**, 375 (2010).
https://doi.org/10.1186/1471-2164-11-375
4   Wiesenfeld, H. C., Hillier, S. L., Krohn, M. A., Landers, D. V. & Sweet, R. L. Bacterial
Vaginosis Is a Strong Predictor of<i>Neisseria
gonorrhoeae</i>and<i>Chlamydia</i><i>trachomatis</i>Infection. *Clinical Infectious*
Diseases **36**, 663-668 (2003). https://doi.org/10.1086/367658
5   Atashili, J., Poole, C., Ndumbe, P. M., Adimora, A. A. & Smith, J. S. Bacterial vaginosis
and HIV acquisition: a meta-analysis of published studies. *AIDS* **22** (2008).
6   Hillier, S. L. *et al.* Association between bacterial vaginosis and preterm delivery of a low-
birth-weight infant. *New England journal of medicine* **333**, 1737-1742 (1995).
7   Klebanoff, M. A. & Brotman, R. M. Treatment of bacterial vaginosis to prevent preterm
birth. *Lancet* **392**, 2141-2142 (2018). https://doi.org/10.1016/s0140-6736(18)32115-9
8   Berman, H. L., Goltsman, D. S. A., Anderson, M., Relman, D. A. & Callahan, B. J.
Gardnerella diversity and ecology in pregnancy and preterm birth. *mSystems* **9**, e0133923
(2024). https://doi.org/10.1128/msystems.01339-23
9   Gardner, H. L. & Dukes, C. D. Haemophilus vaginalis vaginitis: a newly defined specific
infection previously classified non-specific vaginitis. *Am J Obstet Gynecol* **69**, 962-976
(1955).
10   Schwebke, J. R., Muzny, C. A. & Josey, W. E. Role of Gardnerella vaginalis in the
Pathogenesis of Bacterial Vaginosis: A Conceptual Model. *The Journal of Infectious*
Diseases **210**, 338-343 (2014). https://doi.org/10.1093/infdis/jiu089

11   Muzny, C. A. *et al.* An Updated Conceptual Model on the Pathogenesis of Bacterial
Vaginosis. *The Journal of Infectious Diseases* **220**, 1399-1405 (2019).
https://doi.org/10.1093/infdis/jiz342
12   Gelber, S. E., Aguilar, J. L., Lewis, K. L. T. & Ratner, A. J. Functional and Phylogenetic
Characterization of Vaginolysin, the Human-Specific Cytolysin from<i>Gardnerella
vaginalis</i>. *Journal of Bacteriology* **190**, 3896-3903 (2008).
https://doi.org/10.1128/jb.01965-07
13   Patterson, J. L., Stull-Lane, A., Girerd, P. H. & Jefferson, K. K. Analysis of adherence,
biofilm formation and cytotoxicity suggests a greater virulence potential of Gardnerella
vaginalis relative to other bacterial-vaginosis-associated anaerobes. *Microbiology* **156**,
392-399 (2010). https://doi.org/10.1099/mic.0.034280-0
14   Yeoman, C. J. *et al.* Comparative Genomics of Gardnerella vaginalis Strains Reveals
Substantial Differences in Metabolic and Virulence Potential. *PLoS ONE* **5**, e12411
(2010). https://doi.org/10.1371/journal.pone.0012411
15   Alves, P., Castro, J., Sousa, C., Cereija, T. B. & Cerca, N. Gardnerella vaginalis
Outcompetes 29 Other Bacterial Species Isolated From Patients With Bacterial Vaginosis,
Using in an In Vitro Biofilm Formation Model. *The Journal of Infectious Diseases* **210**,
593-596 (2014). https://doi.org/10.1093/infdis/jiu131
16   Castro, J. *et al.* Using an in-vitro biofilm model to assess the virulence potential of
Bacterial Vaginosis or non-Bacterial Vaginosis Gardnerella vaginalis isolates. *Scientific
Reports* **5**, 11640 (2015). https://doi.org/10.1038/srep11640
17   Castro, J. *et al.* Comparative transcriptomic analysis of Gardnerella vaginalis biofilms vs.
planktonic cultures using RNA-seq. *NPJ Biofilms Microbiomes* **3**, 3 (2017).
https://doi.org/10.1038/s41522-017-0012-7
18   Janulaitiene, M. *et al.* Phenotypic characterization of Gardnerella vaginalis subgroups
suggests differences in their virulence potential. *PLoS One* **13**, e0200625 (2018).
https://doi.org/10.1371/journal.pone.0200625
19   Shvartsman, E., Hill, J. E., Sandstrom, P. & Macdonald, K. S. <i>Gardnerella</i>
Revisited: Species Heterogeneity, Virulence Factors, Mucosal Immune Responses, and
Contributions to Bacterial Vaginosis. *Infection and Immunity* **91** (2023).
https://doi.org/10.1128/iai.00390-22
20   Machado, A. & Cerca, N. Influence of Biofilm Formation by<i>Gardnerella
vaginalis</i>and Other Anaerobes on Bacterial Vaginosis. *Journal of Infectious Diseases*
212, 1856-1861 (2015). https://doi.org/10.1093/infdis/jiv338
21   Janulaitiene, M. *et al.* Prevalence and distribution of Gardnerella vaginalis subgroups in
women with and without bacterial vaginosis. *BMC Infectious Diseases* **17** (2017).
https://doi.org/10.1186/s12879-017-2501-y
22   Munch, M. M. *et al.* Gardnerella Species and Their Association With Bacterial Vaginosis.
J Infect Dis **230**, e171-e181 (2024). https://doi.org/10.1093/infdis/jiae026
23   Khan, S., Vancuren, S. J. & Hill, J. E. A Generalist Lifestyle Allows Rare Gardnerella
spp. to Persist at Low Levels in the Vaginal Microbiome. *Microbial Ecology* **82**, 1048-
1060 (2021). https://doi.org/10.1007/s00248-020-01643-1
24   Castro, J., Machado, D. & Cerca, N. Unveiling the role of Gardnerella vaginalis in
polymicrobial Bacterial Vaginosis biofilms: the impact of other vaginal pathogens living
as neighbors. *ISME J* **13**, 1306-1317 (2019). https://doi.org/10.1038/s41396-018-0337-0

25   Balashov, S. V., Mordechai, E., Adelson, M. E. & Gygax, S. E. Identification,
quantification and subtyping of Gardnerella vaginalis in noncultured clinical vaginal
samples by quantitative PCR. *Journal of Medical Microbiology* **63**, 162-175 (2014).
https://doi.org/https://doi.org/10.1099/jmm.0.066407-0

26   Piot, P. *et al.* Biotypes of Gardnerella vaginalis. *Journal of Clinical Microbiology* **20**,
677-679 (1984). https://doi.org/10.1128/jcm.20.4.677-679.1984

27   Benito, R., Vazquez, J. A., Berron, S., Fenoll, A. & Saez-Nieto, J. A. A Modified Scheme
for Biotyping Gardnerella Vaginalis. *Journal of Medical Microbiology* **21**, 357-359
(1986). https://doi.org/https://doi.org/10.1099/00222615-21-4-357

28   Nath, K., Devlin, D. & Beddoe, A. M. Heterogeneity in restriction patterns of Gardnerella
vaginalis isolates from individuals with bacterial vaginosis. *Research in Microbiology*
143, 199-209 (1992). https://doi.org/https://doi.org/10.1016/0923-2508(92)90009-D

29   Ingianni, A., Petruzzelli, S., Morandotti, G. & Pompei, R. Genotypic differentiation
of<i>Gardnerella vaginalis</i>by amplified ribosomal DNA restriction analysis
(ARDRA). *FEMS Immunology & Medical Microbiology* **18**, 61-66 (1997).
https://doi.org/10.1111/j.1574-695x.1997.tb01028.x

30   Ahmed, A. *et al.* Comparative Genomic Analyses of 17 Clinical Isolates of Gardnerella
vaginalis Provide Evidence of Multiple Genetically Isolated Clades Consistent with
Subspeciation into Genovars. *Journal of Bacteriology* **194**, 3922-3937 (2012).
https://doi.org/10.1128/jb.00056-12

31   Cornejo, O. E., Hickey, R. J., Suzuki, H. & Forney, L. J. Focusing the diversity of
<i>Gardnerella vaginalis</i> through the lens of ecotypes. *Evolutionary Applications* **11**,
312-324 (2018). https://doi.org/10.1111/eva.12555

32   Vaneechoutte, M. *et al.* Emended description of Gardnerella vaginalis and description of
Gardnerella leopoldii sp. nov., Gardnerella piotii sp. nov. and Gardnerella swidsinskii sp.
nov., with delineation of 13 genomic species within the genus Gardnerella. *Int J Syst Evol*
Microbiol **69**, 679-687 (2019). https://doi.org/10.1099/ijsem.0.003200

33   Potter, R. F., Burnham, C. D. & Dantas, G. In Silico Analysis of Gardnerella
Genomospecies Detected in the Setting of Bacterial Vaginosis. *Clin Chem* **65**, 1375-1387
(2019). https://doi.org/10.1373/clinchem.2019.305474

34   Innamorati, K. A. *et al.* Metronidazole response profiles of Gardnerella species are
congruent with phylogenetic and comparative genomic analyses. *Genome Medicine* **17**
(2025). https://doi.org/10.1186/s13073-025-01446-4

35   Tarracchini, C. *et al.* Assessing the Genomic Variability of Gardnerella vaginalis through
Comparative Genomic Analyses: Evolutionary and Ecological Implications. *Applied and*
Environmental Microbiology **87** (2020). https://doi.org/10.1128/aem.02188-20

36   Links, M. G., Dumonceaux, T. J., Hemmingsen, S. M. & Hill, J. E. The Chaperonin-60
Universal Target Is a Barcode for Bacteria That Enables De Novo Assembly of
Metagenomic Sequence Data. *PLoS ONE* **7**, e49755 (2012).
https://doi.org/10.1371/journal.pone.0049755

37   Jayaprakash, T. P., Schellenberg, J. J. & Hill, J. E. Resolution and Characterization of
Distinct cpn60-Based Subgroups of Gardnerella vaginalis in the Vaginal Microbiota.
PLoS ONE **7**, e43009 (2012). https://doi.org/10.1371/journal.pone.0043009

38   Zhang, G., Gao, B., Adeolu, M., Khadka, B. & Gupta, R. S. Phylogenomic Analyses and
Comparative Studies on Genomes of the Bifidobacteriales: Identification of Molecular

1129		Signatures Specific for the Order Bifidobacteriales and Its Different Subclades. *Frontiers*
1130		*in Microbiology* **7** (2016). https://doi.org/10.3389/fmicb.2016.00978
1131	39	Lugli, G. A. *et al.* Comparative genomic and phylogenomic analyses of the
1132		Bifidobacteriaceae family. *BMC Genomics* **18** (2017). https://doi.org/10.1186/s12864-
1133		017-3955-4
1134	40	Huys, G. *et al.* Alloscardovia omnicolens gen. nov., sp. nov., from human clinical
1135		samples. *International Journal of Systematic and Evolutionary Microbiology* **57**, 1442-
1136		1446 (2007). https://doi.org/https://doi.org/10.1099/ijs.0.64812-0
1137	41	García-Aljaro, C. *et al.* Neoscardovia arbecensis gen. nov., sp. nov., isolated from porcine
1138		slurries. *Systematic and Applied Microbiology* **35**, 374-379 (2012).
1139		https://doi.org/https://doi.org/10.1016/j.syapm.2012.06.007
1140	42	Simpson, P. J., Ross, R. P., Fitzgerald, G. F. & Stanton, C. Bifidobacterium
1141		psychraerophilum sp. nov. and Aeriscardovia aeriphila gen. nov., sp. nov., isolated from a
1142		porcine caecum. *International Journal of Systematic and Evolutionary Microbiology* **54**,
1143		401-406 (2004). https://doi.org/https://doi.org/10.1099/ijs.0.02667-0
1144	43	Jian, W. & Dong, X. Transfer of Bifidobacterium inopinatum and Bifidobacterium
1145		denticolens to Scardovia inopinata gen. nov., comb. nov., and Parascardovia denticolens
1146		gen. nov., comb. nov., respectively. *International Journal of Systematic and Evolutionary*
1147		*Microbiology* **52**, 809-812 (2002). https://doi.org/https://doi.org/10.1099/00207713-52-3-
1148		809
1149	44	Gerhardt, K. *et al.* FastAAI: efficient estimation of genome average amino acid identity
1150		and phylum-level relationships using tetramers of universal proteins. *Nucleic Acids*
1151		*Research* **53** (2025). https://doi.org/10.1093/nar/gkaf348
1152	45	Schellenberg, J. J. *et al.* Gardnerella vaginalis Subgroups Defined by cpn60 Sequencing
1153		and Sialidase Activity in Isolates from Canada, Belgium and Kenya. *PLOS ONE* **11**,
1154		e0146510 (2016). https://doi.org/10.1371/journal.pone.0146510
1155	46	Bohr, L. L., Mortimer, T. D. & Pepperell, C. S. Lateral Gene Transfer Shapes Diversity of
1156		Gardnerella spp. *Front Cell Infect Microbiol* **10**, 293 (2020).
1157		https://doi.org/10.3389/fcimb.2020.00293
1158	47	Hill, J. E. & Albert, A. Y. K. Resolution and Cooccurrence Patterns of Gardnerella
1159		leopoldii, G. swidsinskii, G. piotii, and G. vaginalis within the Vaginal Microbiome.
1160		*Infection and Immunity* **87** (2019). https://doi.org/10.1128/iai.00532-19
1161	48	Sousa, M. *et al.* Gardnerella pickettii sp. nov. (formerly Gardnerella genomic species 3)
1162		and Gardnerella greenwoodii sp. nov. (formerly Gardnerella genomic species 8) isolated
1163		from female urinary microbiome. *Int J Syst Evol Microbiol* **73** (2023).
1164		https://doi.org/10.1099/ijsem.0.006140
1165	49	Allini Ntiguemassa, P. A. *et al.* Further dissection of Gardnerella vaginalis: description of
1166		Gardnerella lacydonensis sp. nov. (formerly genomic species 2), Gardnerella bretellae sp.
1167		nov. (formerly genomic species 9), Gardnerella massiliensis sp. nov. (formerly genomic
1168		species 14) and Gardnerella phocaeensis sp. nov. *International Journal of Systematic and*
1169		*Evolutionary Microbiology* **76** (2026).
1170		https://doi.org/https://doi.org/10.1099/ijsem.0.007028
1171	50	Teixeira, P., Sousa, M., Ribeiro, T. G., Grosso, F. & Peixe, L. *Unraveling*
1172		*the<i>Gardnerella</i>Pangenome: Insights into Genetic Diversity, Virulence, and*
1173		*Antibiotic Resistance* (Cold Spring Harbor Laboratory, 2025).

51 Malki, K. *et al.* Genomes of Gardnerella Strains Reveal an Abundance of Prophages
within the Bladder Microbiome. *PLOS ONE* **11**, e0166757 (2016).
https://doi.org/10.1371/journal.pone.0166757
52 Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned
pangenome graph. *PLOS Computational Biology* **16**, e1007732 (2020).
https://doi.org/10.1371/journal.pcbi.1007732
53 Flores Ramos, S. *et al.* Genomic Stability and Genetic Defense Systems in
Dolosigranulum pigrum , a Candidate Beneficial Bacterium from the Human
Microbiome. *mSystems* **6** (2021). https://doi.org/10.1128/msystems.00425-21
54 Dillard, L. R., Glass, E. M., Lewis, A. L., Thomas-White, K. & Papin, J. A. Metabolic
Network Models of the *Gardnerella* Pangenome Identify Key Interactions with
the Vaginal Environment. *mSystems* **8** (2023). https://doi.org/10.1128/msystems.00689-22
55 Garcia, E. M., Kraskauskiene, V., Koblinski, J. E. & Jefferson, K. K. Interaction of
Gardnerella vaginalis and Vaginolysin with the Apical versus Basolateral Face of a Three-
Dimensional Model of Vaginal Epithelium. *Infection and Immunity* **87** (2019).
https://doi.org/10.1128/iai.00646-18
56 Cauci, S. *et al.* Pore-forming and haemolytic properties of the *Gardnerella
vaginalis* cytoiysin. *Molecular Microbiology* **9**, 1143-1155 (1993).
https://doi.org/10.1111/j.1365-2958.1993.tb01244.x
57 Robinson, L. S., Schwebke, J., Lewis, W. G. & Lewis, A. L. Identification and
characterization of NanH2 and NanH3, enzymes responsible for sialidase activity in the
vaginal bacterium Gardnerella vaginalis. *Journal of Biological Chemistry* **294**, 5230-5245
(2019). https://doi.org/10.1074/jbc.ra118.006221
58 Agarwal, K. & Lewis, A. L. Vaginal sialoglycan foraging by *Gardnerella
vaginalis*: mucus barriers as a meal for unwelcome guests? *Glycobiology* **31**, 667-680
(2021). https://doi.org/10.1093/glycob/cwab024
59 Pelayo, P. *et al.* *Prevotella* are major contributors of sialidases in the human
vaginal microbiome. *Proceedings of the National Academy of Sciences* **121** (2024).
https://doi.org/10.1073/pnas.2400341121
60 Perna, V. N., Barrett, K., Meyer, A. S. & Zeuner, B. Substrate specificity and
transglycosylation capacity of α-L-fucosidases across GH29 assessed by bioinformatics-
assisted selection of functional diversity. *Glycobiology* **33**, 396-410 (2023).
https://doi.org/10.1093/glycob/cwad029
61 Watanabe, S. Characterization of l-2-keto-3-deoxyfuconate aldolases in a
nonphosphorylating l-fucose metabolism pathway in anaerobic bacteria. *Journal of
Biological Chemistry* **295**, 1338-1349 (2020). https://doi.org/10.1016/s0021-
9258(17)49890-5
62 Watanabe, S. Characterization of a novel L-fuconate dehydratase involved in the non-
phosphorylated pathway of L-fucose metabolism from bacteria. *Bioscience,
Biotechnology, and Biochemistry* **88**, 177-180 (2024).
https://doi.org/10.1093/bbb/zbad161
63 Kurukulasuriya, S. P., Patterson, M. H. & Hill, J. E. Slipped-Strand Mispairing in the
Gene Encoding Sialidase NanH3 in *Gardnerella* spp. *Infection and Immunity* **89**
(2021). https://doi.org/10.1128/iai.00583-20

64   Bhandari, P., Tingley, J., Abbott, D. W. & Hill, J. E. Glycogen-Degrading Activities of
Catalytic Domains of α-Amylase and α-Amylase-Pullulanase Enzymes Conserved in
<i>Gardnerella</i> spp. from the Vaginal Microbiome. *Journal of Bacteriology* **205**
(2023). https://doi.org/10.1128/jb.00393-22
65   Bordigoni, A., Bouchard, S. & Desnues, C. Isolation and Characterization of
<i>Gardnerella</i> Phage vB_Gva_AB1, a Bacteriophage Infecting a Clinical Strain of
Gardnerella vaginalis. *Microbiology Resource Announcements* **10** (2021).
https://doi.org/10.1128/mra.00053-21
66   Meier-Kolthoff, J. P. & Göker, M. VICTOR: genome-based phylogeny and classification
of prokaryotic viruses. *Bioinformatics* **33**, 3396-3404 (2017).
https://doi.org/10.1093/bioinformatics/btx440
67   Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of bacteria.
Nature Reviews Microbiology **21**, 686-700 (2023). https://doi.org/10.1038/s41579-023-
00934-x
68   Oliveira, P. H., Touchon, M. & Rocha, E. P. C. The interplay of restriction-modification
systems with mobile genetic elements and their prokaryotic hosts. *Nucleic Acids Research*
42, 10618-10631 (2014). https://doi.org/10.1093/nar/gku734
69   Tock, M. R. & Dryden, D. T. F. The biology of restriction and anti-restriction. *Current*
Opinion in Microbiology **8**, 466-472 (2005).
https://doi.org/https://doi.org/10.1016/j.mib.2005.06.003
70   Tesson, F. *et al.* Systematic and quantitative view of the antiviral arsenal of prokaryotes.
Nature Communications **13** (2022). https://doi.org/10.1038/s41467-022-30269-9
71   Tal, N. *et al.* Bacteria deplete deoxynucleotides to defend against bacteriophage infection.
Nature Microbiology **7**, 1200-1209 (2022). https://doi.org/10.1038/s41564-022-01158-0
72   Garvey, P., Fitzgerald, G. F. & Hill, C. Cloning and DNA sequence analysis of two
abortive infection phage resistance determinants from the lactococcal plasmid pNP40.
Applied and Environmental Microbiology **61**, 4321-4328 (1995).
https://doi.org/10.1128/aem.61.12.4321-4328.1995
73   Deng, Y.-M., Harvey, M. L., Liu, C.-Q. & Dunn, N. W. A novel plasmid-encoded phage
abortive infection system from Lactococcus lactis biovar. diacetylactis. *FEMS*
Microbiology Letters **146**, 149-154 (1997). https://doi.org/10.1111/j.1574-
6968.1997.tb10185.x
74   Cohen, D. *et al.* Cyclic GMP–AMP signalling protects bacteria against viral infection.
Nature **574**, 691-695 (2019). https://doi.org/10.1038/s41586-019-1605-5
75   Millman, A. *et al.* An expanded arsenal of immune systems that protect bacteria from
phages. *Cell Host & Microbe* **30**, 1556-1569.e1555 (2022).
https://doi.org/10.1016/j.chom.2022.09.017
76   Mestre, M. R. *et al.* UG/Abi: a highly diverse family of prokaryotic reverse transcriptases
associated with defense functions. *Nucleic Acids Research* **50**, 6084-6101 (2022).
https://doi.org/10.1093/nar/gkac467
77   Gao, L. *et al.* Diverse enzymatic activities mediate antiviral immunity in prokaryotes.
Science **369**, 1077-1084 (2020). https://doi.org/10.1126/science.aba0372
78   Doron, S. *et al.* Systematic discovery of antiphage defense systems in the microbial
pangenome. *Science* **359**, eaar4120 (2018). https://doi.org/10.1126/science.aar4120

79   Chen, C. *et al.* Development of a Pre-Modification Strategy to Overcome Restriction–
Modification Barriers and Enhance Genetic Engineering in Lactococcus lactis for Nisin
Biosynthesis. *International Journal of Molecular Sciences* **26**, 2200 (2025).
https://doi.org/10.3390/ijms26052200

80   Hu, S. *et al.* Altering under-represented DNA sequences elevates bacterial transformation
efficiency. *mBio* **14** (2023). https://doi.org/10.1128/mbio.02105-23

81   Finn, M. B., Ramsey, K. M., Tolliver, H. J., Dove, S. L. & Wessels, M. R. Improved
transformation efficiency of group A Streptococcus by inactivation of a type I restriction
modification system. *PLOS ONE* **16**, e0248201 (2021).
https://doi.org/10.1371/journal.pone.0248201

82   Klimowicz, A. K., Garcia, E. M., Jefferson, K. K. & Dillard, J. P. Genetic transformation
of *Gardnerella* species and characterization of vaginolysin and sialidase mutants.
Infection and Immunity **93** (2025). https://doi.org/10.1128/iai.00299-25

83   Kularatne, B. M. D. N. & Hill, J. E. Transformation of *Gardnerella vaginalis* with
a *Bifidobacterium-Escherichia coli* shuttle vector plasmid. *Microbiology*
Spectrum **13** (2025). https://doi.org/10.1128/spectrum.00481-25

84   Zhu, M. *et al.* Vaginal *Lactobacillus* fatty acid response mechanisms reveal a
novel strategy for bacterial vaginosis treatment. *bioRxiv*, 2023.2012.2030.573720 (2023).
https://doi.org/10.1101/2023.12.30.573720

85   Kharsany, A. B., Hoosen, A. A. & Van Den Ende, J. Antimicrobial susceptibilities of
Gardnerella vaginalis. *Antimicrobial Agents and Chemotherapy* **37**, 2733-2735 (1993).
https://doi.org/10.1128/aac.37.12.2733

86   Johnston, C. D. *et al.* Systematic evasion of the restriction-modification barrier in
bacteria. *Proceedings of the National Academy of Sciences* **116**, 11454-11459 (2019).
https://doi.org/10.1073/pnas.1820256116

87   Srinivasan, S. *et al.* More Easily Cultivated Than Identified: Classical Isolation With
Molecular Identification of Vaginal Bacteria. *Journal of Infectious Diseases* **214**, S21-S28
(2016). https://doi.org/10.1093/infdis/jiw192

88   Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J. & Sayers, E. W. GenBank.
Nucleic Acids Research **44**, D67-D72 (2016). https://doi.org/10.1093/nar/gkv1276

89   Meier-Kolthoff, J. P. & Göker, M. TYGS is an automated high-throughput platform for
state-of-the-art genome-based taxonomy. *Nature Communications* **10** (2019).
https://doi.org/10.1038/s41467-019-10210-3

90   Alessandri, G., Van Sinderen, D. & Ventura, M. The genus Bifidobacterium: from
genomics to functionality of an important component of the mammalian gut microbiota.
Computational and Structural Biotechnology Journal **19**, 1472-1487 (2021).
https://doi.org/10.1016/j.csbj.2021.03.006

91   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Research* **25**, 1043-1055 (2015).
https://doi.org/10.1101/gr.186072.114

92   Aziz, R. K. *et al.* The RAST Server: Rapid Annotations using Subsystems Technology.
BMC Genomics **9**, 75 (2008). https://doi.org/10.1186/1471-2164-9-75

93 Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nature Communications **9** (2018). https://doi.org/10.1038/s41467-018-07641-9
94 Gu, Z. Complex heatmap visualization. *iMeta* **1** (2022). https://doi.org/10.1002/imt2.43
95 Auch, A. F., Von Jan, M., Klenk, H.-P. & Göker, M. Digital DNA-DNA hybridization for
microbial species delineation by means of genome-to-genome sequence comparison.
Standards in Genomic Sciences **2**, 117-134 (2010). https://doi.org/10.4056/sigs.531120
96 Meier-Kolthoff, J. P., Auch, A. F., Klenk, H.-P. & Göker, M. Genome sequence-based
species delimitation with confidence intervals and improved distance functions. *BMC*
Bioinformatics **14**, 60 (2013). https://doi.org/10.1186/1471-2105-14-60
97 Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic*
Acids Research **53**, D444-D456 (2025). https://doi.org/10.1093/nar/gkae1082
98 barrnap v. 0.9 (Australia, 2018).
99 Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**, 113 (2004). https://doi.org/10.1186/1471-2105-
5-113
100 Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592-593 (2011).
https://doi.org/10.1093/bioinformatics/btq706
101 Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the
phylogenetic tree display and annotation tool. *Nucleic Acids Research* **52**, W78-W82
(2024). https://doi.org/10.1093/nar/gkae268
102 Gardner, S. N., Slezak, T. & Hall, B. G. kSNP3.0: SNP detection and phylogenetic
analysis of genomes without genome alignment or reference genome. *Bioinformatics* **31**,
2877-2878 (2015). https://doi.org/10.1093/bioinformatics/btv271
103 Price, M. N., Deutschbauer, A. M. & Arkin, A. P. GapMind: Automated Annotation of
Amino Acid Biosynthesis. *mSystems* **5** (2020). https://doi.org/10.1128/msystems.00291-
20
104 Price, M. N., Deutschbauer, A. M. & Arkin, A. P. Filling gaps in bacterial catabolic
pathways with computation and high-throughput genetics. *PLOS Genetics* **18**, e1010156
(2022). https://doi.org/10.1371/journal.pgen.1010156
105 Price, M. N. *et al. Improving the annotation of amino acid biosynthesis pathways:*
GapMind 2024 (openRxiv, 2024).
106 Payne, L. J. *et al.* PADLOC: a web server for the identification of antiviral defence
systems in microbial genomes. *Nucleic Acids Research* **50**, W541-W550 (2022).
https://doi.org/10.1093/nar/gkac400
107 Payne, L. J. *et al.* Identification and classification of antiviral defence systems in bacteria
and archaea with PADLOC reveals new system types. *Nucleic Acids Research* **49**, 10868-
10878 (2021). https://doi.org/10.1093/nar/gkab883
108 Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and
curation of microbial viruses, and evaluation of viral community function from genomic
sequences. *Microbiome* **8** (2020). https://doi.org/10.1186/s40168-020-00867-0
109 Perfilyev, A. *et al.* Large-scale analysis of bacterial genomes reveals thousands of lytic
phages. *Nature Microbiology* **11**, 42-52 (2025). https://doi.org/10.1038/s41564-025-
02203-4

110   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
1350   Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530-1534 (2020).
1351   https://doi.org/10.1093/molbev/msaa015
1352   111   Greenwood, J. R. & Pickett, M. J. Transfer of Haemophilus vaginalis Gardner and Dukes
1353   to a New Genus, Gardnerella: G. vaginalis (Gardner and Dukes) comb. nov. *International*
1354   *Journal of Systematic Bacteriology* **30**, 170-178 (1980).
1355   https://doi.org/10.1099/00207713-30-1-170
1356   112   Hitch, T. C. A. *et al.* Automated analysis of genomic sequences facilitates high-
1357   throughput and comprehensive description of bacteria. *ISME Communications* **1** (2021).
1358   https://doi.org/10.1038/s43705-021-00017-z
1359